Evaluating Tumor Evolution via Genomic Profiling of Individual Tumor Spheroids in a Malignant Ascites from a Patient with Ovarian Cancer Using a Laser-aided Cell Isolation Technique

Sungsik Kim[1, 4, 11], Soochi Kim[3, 5, 11], Jinhyun Kim[1], Boyun Kim[6], Se Ik Kim[7], Min A Kim[8], Sunghoon Kwon[1-3, *], and Yong Sang Song[3, 5, 7, 9, 10 *]

[1] Department of Electrical and Computer Engineering, Seoul National University, Seoul 08826,

Republic of Korea

[2] Institutes of Entrepreneurial BioConvergence, Seoul National University, Seoul 08826,

Republic of Korea

[3] Seoul National University Hospital Biomedical Research Institute, Seoul National University Hospital, Seoul 03080, Republic of Korea

[4] Interdisciplinary Program for Bioengineering, Seoul National University, Seoul 08826, Republic of Korea

[5] Cancer Research Institute, Seoul National University College of Medicine, Seoul 03080, Republic of Korea

[6] Department of Anesthesiology, McGovern Medical School, The University of Texas Health Science Center at Houston, Houston, TX, USA

[7] Department of Obstetrics and Gynecology, Seoul National University College of Medicine,

Seoul 03080, Republic of Korea

[8] Department of Pathology, Seoul National University College of Medicine, Seoul 03080,

Republic of Korea

[9] Interdisciplinary Program in Cancer Biology, Seoul National University College of

Medicine, Seoul 03080, Republic of Korea

[10] Biomodulation Department of Agricultural Biotechnology, Seoul National University,

Seoul 03080, Republic of Korea

[11] These authors contributed equally to this work

[*] These authors jointly supervised this work. Correspondence should be addressed to SH.K.

(skwon@snu.ac.kr) or Y.S.S (yssong@snu.ac.kr)

Abstract

Background

Epithelial ovarian cancer (EOC) is a silent but mostly lethal gynecologic malignancy. Most

patients present with malignant ascites and peritoneal seeding at diagnosis. In the present

study, we used a laser-aided isolation technique to investigate the clonal relationship between

the primary tumor and tumor spheroids found in the malignant ascites of an EOC patient.

Somatic alteration profiles of ovarian cancer-related genes were determined for eight spatially

separated samples from primary ovarian tumor tissues and ten tumor spheroids from the

malignant ascites using next-generation sequencing.

Results

We observed high levels of intra-tumor heterogeneity (ITH) in copy number alterations

(CNAs) and single-nucleotide variants (SNVs) in the primary tumor and the tumor spheroids.

As a result, we discovered that tumor cells in the primary tissues and the ascites were

genetically different lineages. We categorized the CNAs and SNVs into clonal and subclonal

alterations according to their distribution among the samples. Also, we identified focal

amplifications and deletions in the analyzed samples. For SNVs, a total of 171 somatic

mutations were observed, among which 66 were clonal mutations present in both the primary

tumor and the ascites, and 61 and 44 of the SNVs were subclonal mutations present in only

the primary tumor or the ascites, respectively.

Conclusions

Based on the somatic alteration profiles, we constructed phylogenetic trees and inferred the

evolutionary history of tumor cells in the patient. The phylogenetic trees constructed using

the CNAs and SNVs showed that two branches of the tumor cells diverged early from an

69   ancestral tumor clone during an early metastasis step in the peritoneal cavity. Our data

70   support the monophyletic spread of tumor spheroids in malignant ascites.

71   **Keywords**: Tumor evolution, Intra-tumor heterogeneity, Malignant ascites, Next-generation

72   sequencing, Epithelial ovarian cancer, Phylogenetic analysis

Background

Epithelial ovarian cancer (EOC) is a silent but mostly lethal gynecologic malignancy. The most common histological EOC subtype is high-grade serous carcinoma, and the current treatment strategy involves a primary debulking surgery followed by chemotherapy to reduce the tumor burden [1, 2]. Recent advances in genomics have revealed the presence of extensive intra-tumor heterogeneity (ITH) in many cancers, including ovarian cancer [3-5]. The presence of extensive clonal diversity increases the capacity of a given tumor to survive upon an expected strike in the microenvironment and thus is thought to be responsible for a reduced response to current chemotherapy and to contribute to chemoresistance development [6-8].

Unlike other solid tumors, the primary route of metastasis in EOC patients is the transcoelomic metastasis route, which is a passive process and involves dissemination of tumor cells from the primary tumor tissue into the peritoneal cavity [9]. Thus, early disseminating clones may exist in the malignant ascites tumor microenvironment (TME) and may form an independent subclonal lineage and contribute to ITH. Both protumorigenic and antitumorigenic factors are known to be enriched in the malignant ascites TME [10]. However, genetic differences between tumor cells in the primary tissue and tumor cells surviving in the ascites TME are not yet fully understood. Multi-region sequencing of both the primary tumor and associated metastases in ovarian cancer has provided insights into spatial heterogeneity and has shown that metastatic tumors maintain the genetic alterations found in the primary tumor and arise with little accumulation of genetic alterations [5]. However, the extent of the genetic heterogeneity within and between the primary tumor and tumor cells found in ascites remains underestimated.

Here, to uncover the genetic heterogeneity of tumor cells in malignant ascites, we

introduced a genetic profiling method for individual tumor spheroids which are the common

form of tumor cells floating in malignant ascites [11]. Inspired by single-cell analysis, we

hypothesized that genetic profiling of individual tumor spheroids might uncover the

heterogeneity within and between the primary tumor and tumor cells in ascites. We isolated

individual tumor spheroids through a laser-aided isolation technique. Then, we performed

low-depth whole-genome sequencing (WGS) and high-depth whole-exome sequencing (WES)

for ten tumor spheroids and eight primary tumor samples from a high-grade serous (HGS)

EOC patient. We explored somatic copy number alterations (CNAs) and single-nucleotide

variants (SNVs) to determine the tumor evolution and ITH between the primary tissues and

the tumor spheroids from the malignant ascites. This study reports the feasibility of analyzing

tumor cells in malignant ascites to detect early disseminating EOC clones.

Results

Preparation and isolation of single tumor spheroids from the ascites of an ovarian

cancer patient

A malignant ascites was collected during a primary debulking surgery. The tumor

spheroids in the malignant ascites were purified, fixed and prepared on a discharging layer-

coated glass slide (Fig. 1A). Single tumor spheroids on the slide were isolated by an infrared

(IR) laser pulse as described in our previous publication [12]. Briefly, the discharging layer

consisted of indium tin oxide (ITO), which vaporizes when irradiated by an IR laser pulse.

The ITO vaporization generates pressure, by which cells in the irradiated area are discharged

from the slide. From the prepared sample on the slide, we isolated ten individual tumor

spheroids, which were tens of micrometers in diameter and contained hundreds of cells (Fig.

1B, C). Isolating and capturing each tumor spheroid took less than 1 second on average,

which means this technique is feasible for analyzing a large number of samples and could be

implemented in a routine procedure. The isolated single tumor spheroids were collected in

PCR tubes for further reactions.

Whole-genome amplification of the isolated individual tumor spheroids

The isolated single tumor spheroids were lysed by proteinase K. Then, the samples

underwent multiple displacement amplification (MDA, Fig. 2A). The amplification was

monitored via real-time PCR. The results showed that all the isolated samples yielded

successful amplification (10/10). Additionally, comparing the amplification plots between the

tumor spheroids and controls showed that there was no or a negligible amount of carry-over

contamination (Fig. 2B). Every reaction yielded over 2 µg of amplified DNA, which was

enough to conduct WGS and WES.

Next, we calculated and plotted the distributions of the normalized read depth (Fig.

2C) and variant allele frequency (VAF, Fig. 2D) based on the sequencing data to evaluate the

amplification uniformity of the MDA reaction. In the Fig. 2C and 2D, the distributions of the

MDA products from single cells were used for comparison. Normalized read depth indicates

the uniformity of the number of sequencing reads throughout the whole-genome. The DNA

from bulk tumor samples showed normal-like distributions with small variance, but whole-

genome amplified DNA from single cells presented a skewed distribution because of non-

uniform amplification. In contrast, the distributions of the tumor spheroids were similar to the

distributions of the tumor bulk samples, rather than the whole-genome amplified products

from the single cells. This result suggests that the effect of non-uniform amplification during

MDA was minimized because hundreds of cells were included in the individual tumor

spheroids. Similarly, the VAF distributions of the tumor spheroids were similar to those of the

bulk tumor samples but not to the distributions of the single cells. This result supports the

presumption that the MDA products of the tumor spheroids present a balanced allele

amplification without losing one of the two alleles.

Low-depth WGS reveals the somatic CNAs and genetic subclones

First, we assessed the somatic CNAs of the primary ovarian cancer tissues and the

tumor spheroids from the ascites (Additional file 1: Table S1). We carried out low-depth

WGS using the Illumina platform to produce 8.53 ± 0.879 ($\times 10^6$) sequenced reads for each

sample. As a result, we generated CNA profiles based on which we performed a hierarchical

clustering analysis (Fig. 3A). The clustering yielded three distinct genetic subgroups. The

primary ovarian cancer tissues (RO 1-7 and LO, named "Primary clone" and colored red)

were clustered together. In contrast, the tumor spheroids from the ascites were divided into

two clusters, one of which showed a primary-like CNA profile (AC 1-3 and 7-8, named

"Ascites clone 1" and colored yellow), but the other presented a normal-like profile (AC 4-6

and 9-10, named "Ascites clone 2", colored green).

Interestingly, the CNA profiles showed that deletion of FAT1 and amplification of

MYC, PARP10, and CYC1 were shared by most of the samples (Fig. 3B). These genes are

reported to be recurrently deleted (FAT1) or amplified (MYC, PARP10, and CYC1) in pan-

cancer data [13]. These facts suggest that the shared CNAs might be the driving alterations at

the first stage of cancer initiation. However, the primary clone had exclusive focal

amplifications of KDM5A and NOTCH3 (Fig. 3B), which are known as recurrently amplified

genes in ovarian cancer [13, 14]. These focal amplifications of KDM5A and NOTCH3 might

allow the primary clone to overwhelm the other subclones and finally dominate the left and

right ovaries. However, we did not find a critical focal amplification or a deep deletion

exclusive to Ascites clone 1. This implied that other types of alterations might drive Ascites

clone 1 to survive or propagate in the peritoneal fluid.

WES reveals somatic SNVs and genetic subclones

To identify the somatic SNVs, the samples underwent WES. For each sample, the sequencing run generated 134 ± 21.4 depth of data, covering the whole exome of the human genome. As a result, 171 somatic SNVs were identified by variant calling from all the samples (Additional file 2: Table S2). The results shown in Fig. 4A and 4B revealed that 38.6% of the SNVs were common to the primary tumor and tumor spheroids from the ascites, and 35.7% of the SNVs exclusively belonged to primary-only and 25.7% to ascites-only mutations. The exclusive mutations in the Ascites clone suggest that this clone evolved by accumulating mutations independent from the Primary clone. Interestingly, the Ascites clone had a nonsynonymous mutation in the KRAS gene at 12p12.1. A single nucleotide substitution (C>T) results in an activating KRAS mutation that is a well-known oncogenic mutation associated with the anchorage-independent growth of tumor cells through the acquisition of anoikis resistance in various malignancies [15, 16]. Therefore, the mutation in KRAS in the Ascites clone might provide an additional fitness gain for anchorage-independent survival in the ascites TME. However, both the Primary and Ascites clones shared somatic SNVs in TP53 and ARID1A, which are well-known driver mutations in ovarian cancer [17, 18]. At the initial stage of tumorigenesis, these mutated genes might be tumor-initiating SNVs in conjunction with the CNAs of FAT1, MYC, PARP10, and CYC1. In addition to these somatic variants, the patient had germline variants in BRCA1 (NM_007294.3:c.1511dupG) and TP53 (NM_001126118:c.C98G), which are well-known susceptibility genes of ovarian cancer and are likely to predispose individuals to ovarian cancer and promote carcinogenesis (Additional file 3: Table S3) [19,20].

Cellular composition of the tumor spheroids

Regarding the CNAs, Ascites clone 2 had no alteration except for amplification of the

8q24 region. Concerning the SNVs, Ascites clone 2 had fewer mutations than the other

clusters. Based on these facts, we examined the possibility that normal cells exist in a tumor

spheroid. We assumed that the VAF distribution of Ascites clones 1 and 2 would be similar if

the two subclones had a similar proportion of normal cells. However, the VAF of Ascites

clone 2 would be low if a single tumor spheroid from the clone included a high proportion of

normal cells. We tested this idea by plotting the VAF distribution of each sample (Fig. 5). The

results showed that most of the VAF distributions from the Primary clone and Ascites clone 1

were located at a higher range than those from Ascites clone 2. Therefore, we concluded that

the small number of CNAs and SNVs in Ascites clone 2 was not due to their true

characteristics but because the proportion of tumor cells in the tumor spheroid was small.

Consequently, we excluded Ascites clone 2 from the following phylogenetic analysis.

In addition to the presence of normal cells in the samples, we examined the possibility

of the presence of heterogeneous tumor cells in the samples. By comparing the allele

frequency distributions of the common and primary-only mutations for each sample, we

found that the allele frequencies of the common mutations were higher than those of the

primary-only mutations for the primary tissues (7 of 8 samples, $p < 0.01$). This result implies

that each of the primary tissues (except RO3) had two or more subclones sharing common

mutations but not subclonal mutations. In contrast, the allele frequencies of the common

mutations were similar to those of the ascites-only mutations for the tumor spheroids (8 of 10

samples). This result can be interpreted to indicate that, compared with the primary tissue

samples, each tumor spheroid was comprised of genetically homogeneous tumor cells.

Constructing phylogenetic trees based on the somatic CNAs and SNVs

The phylogenetic trees were constructed from the CNA and SNV data. We achieved a

CNA-based phylogeny analysis by identifying the common chromosomal breakpoints,

calculating a trinary event matrix, and constructing a maximum parsimony tree [28]. The

phylogenetic tree showed that an ancestral cancer clone accumulated CNAs and divided into

two clones, which gained additional exclusive CNAs (Fig. 6A). Notably, these two genetic

clones were composed of tumor spheroids from ascites and tumor tissues. Potentially,

physically separated and biologically distinct TMEs might drive cancer cells into different

alteration statuses.

Maximum parsimony tree generation using the CNA data has a couple of limitations.

First, this approach needs to set thresholds to define the amplified, neutral, and deleted status.

The resultant tree is significantly affected by thresholds, and there is no golden rule to set the

thresholds. Second, the proportion of normal cells in a sample has a substantial impact on a

tree because the CNA status might be incorrectly assigned according to the normal cell

portion. For example, the VAFs of RO6 (Fig. 5) show that the sample had a large number of

normal cells. In this case, the copy number value of RO6 was close to the normal value (Fig.

3A), although the overall pattern was not similar to that of the normal sample. Thus, the

thresholding led RO6 to be the same as the normal sample. For this reason, we excluded RO6

when constructing the maximum parsimony tree based on the CNA data.

Next, we constructed a phylogenetic tree from the SNV data. This approach does not

use manual thresholding, and a phylogenetic tree is less affected by a normal cell portion.

Therefore, we expected that, compared with the CNA-based approach, this approach would

provide a more accurate result. The results showed that the cancer cells accumulated

mutations as a single clone and divided into two independent clones (Fig. 6B). Moreover,

with the full advantage of the SNV information, the phylogenetic tree presented the

sequential creation of RO3, LO, and the rest of the Primary clones. Overall, the phylogenetic

tree based on the SNV data rather than the CNA data presented a more stable and biologically

explainable result.

Inferring the evolutionary trajectory of the primary ovarian cancer and the single

tumor spheroids in the ascites

This patient harbored a bilateral ovarian tumor at the time of the primary debulking

surgery. It is important to note whether these bilateral tumors arise independently or are the

result of metastasis. The clonal evolution of the tumorigenesis theory provides two

mechanisms of bilateral ovarian tumor development. If bilateral ovarian tumors arise from

independent ancestral clones, they would have distinct genomic profiles without sharing

somatic alterations. In contrast, bilateral tumors would have an identical set of somatic

variants if they resulted from metastasis [21]. The somatic CNAs and SNVs of the left and

right primary ovarian tumor in this study displayed comparable genomic profiles, strongly

indicating a monoclonal origin of the bilateral tumor in this patient. This was further

confirmed by calculating the clonality index (CI) based on previous reports [21, 22] revealing

that the bilateral ovarian tumors were clonally related (CI1 = 1.0).

Finally, the history of the ovarian cancer development and progression was

established based on the genomic profiles to understand the tumor evolution and its direction

in this patient. As noted earlier, ovarian cancer metastasis occurs through a passive process,

which initially involves physical shedding of tumor cells from the primary tumor into the

peritoneal cavity, and the accumulation of ascites facilitates distant seeding of tumor cells

along the peritoneal wall. Given a fixed chance of evolution, two scenarios are possible,

either a monoclonal or polyclonal seeding process. If only certain clones from the primary

tumor are fit to survive in the ascites TME, distinct clones, which may have diverged early,

may be selected and progress over time in the primary and ascites TMEs, showing a tendency

toward independent tumor evolution driven by different TMEs. In contrast, if tumor evolution

is entirely driven by clonal dominance and the physical shedding of tumor cells from the

primary tumor occurs by chance, then dominant clones expand in size and others may remain

unchanged or become extinct over time at the primary tumor site. As the tumor grows,

multiple clones may shed from the primary tumor into ascites. The ascites TME then acts as a

reservoir of clonal lineage, and tumor cells in the ascites would represent the entire

mutational landscape of a given tumor. For our case, we observed significant genetic

differences in the CNAs and SNVs among the primary tissue samples and tumor spheroids.

The dominant clones found in the right ovary were absent in the ascites TME, and we found

44 tumor spheroid–specific somatic SNVs (Additional file 2: Table S2). Furthermore, the

comparable allele frequencies between the common mutations and tumor spheroid–specific

mutations suggest that the tumor spheroids in the ascites TME are comprised of genetically

homogeneous tumor cells compared with the primary tissues. Therefore, we conclude that the

tumor spheroids were from a single subclonal lineage, supporting a mono- and early-seeding

origin of the tumor spheroids in this patient. Based on these perspectives, we drew a potential

evolutionary trajectory of the tumor from the patient (Fig. 7A). The tumor was initiated at the

right ovary to generate the ancestral clone. With further accumulation of mutations, the

ancestral clone evolved into two subclones, the first of which was found in the right ovary

and metastasized to the left ovary. The second subclone shed into the ascites TME and

became extinct or dominated by the first subclone in the right ovary (Additional file 4: Table

S4). Eventually, the Ascites subclone moved to the peritoneal cavity. In addition, the

summary of genome-wide somatic CNAs and SNVs indicated that the tumor cells in the

primary tissue and the ascites possessed exclusive alterations as well as common ones (Fig.

7B). This result shows that the tumor cells in the primary tissue and the ascites were two

subclonal lineages, which branched from one ancestral lineage.

Discussion

In this study, we attempted to determine the presence of genetic heterogeneity within

and between a primary tumor and the associated tumor spheroids in the ascites by performing

multi-region sequencing of the primary tumor and genetic profiling of the individual tumor

spheroids using the laser-aided cell isolation technique. We performed both WGS and WES

of the primary tumor and tumor spheroid samples. First, we discovered high ITH levels in

eight primary tissues and ten tumor spheroids. We also discovered that the CNA profiles in

the primary and associated tumor spheroids were separated into two distinct genetic clusters,

suggesting that the TME may be operative during tumor evolution. Second, we identified

somatic SNVs using WES. We discovered a total of 171 somatic SNVs from all the samples,

and 66 (38.6 %) of these SNVs were ubiquitous mutations that were common to the primary

tumor and tumor spheroids. The rest were either primary-only (61 SNVs, 35.7 %) or ascites-

only (44 SNVs, 25.7 %) mutations, highlighting the notion that the tumor spheroids might

have diverged early and accumulated additional mutations independently from the Primary

clone. Supporting this idea, both phylogenic analyses, using the CNAs and SNVs, showed

that the tumor spheroids might have diverged early from an ancestral tumor clone, evolved

further with distinctive genomic profiles, and formed an independent subclonal lineage,

thereby contributing to the ITH.

We also assessed the normal cell contamination in both the primary tumor and tumor

spheroids using the VAF distribution in each sample. Indeed, both the Primary clone and

Ascites clone 1 showed higher VAF distributions than Ascites clone 2, suggesting that the

normal-like CNA and SNV profiles in Ascites clone 2 were due to a high proportion of

normal cells. These findings are consistent with previous data from ovarian cancer patient-

derived tumor spheroids and mouse models that suggested the presence of tumor-associated

macrophages in the center of tumor spheroids [23].

Although we only studied a single high-grade EOC patient, our data support previous

studies demonstrating early divergence of the ascites sample from the primary tumor [24].

Further studies are needed to compare similarities and differences between the ascites

spheroids and distant metastasis samples. Our data suggest that the mutation set of ascites

spheroids does not represent the entire mutational landscape of a given EOC patient. This

disagrees with recent findings by Choi et al. [25] showing that ascites tumor cells represent

the entire mutational landscape of a given tumor, and no additional genetic aberrations were

detected. In contrast, our data showed the presence of genetic heterogeneity within and

between the primary tumor and the associated ascites spheroids. Moreover, the primary and

associated ascites spheroids diverged early in tumor development, and not all the Primary

clones disseminated into the ascites TME. However, our study is limited to a single ascites

TME and provided no insight into distant metastatic sites.

Additionally, our data demonstrated that, compared with the primary tissue samples,

each tumor spheroid was comprised of genetically homogeneous tumor cells (Fig. 5). This

can be interpreted in two ways. First, the tumor cells in an ascites may have low ITH. In this

case, the spheroids of the tumor cells would be genetically homogeneous. Second, the tumor

cells with a similar genetic profile may form individual tumor spheroids. In this case, the

tumor cells in each tumor spheroid might have the same genetic profile, but two different

tumor spheroids might be genetically different. For this case, isolating and analyzing the

individual tumor spheroids from ascites might be widely utilized to discover the ITH of

ovarian cancer.

Our data can partly be explained by the theory of Darwinian selection. For simplicity,

tumor evolution is described as a series of expansions of clones, where each expansion series

is driven by additional mutation acquisition, and clone fitness is tested by Darwinian

selection. This selective sweep is context-dependent, and thus, genetic variants that are

beneficial at a certain point may become extinct throughout the period of tumor progression.

As a consequence, these clones may be absent in a fully grown tumor [26]. The selective

pressures are further influenced by the dynamics of the TME, thereby increasing the

complexity of tumor evolution [27]. The presence of extensive ITH in tumor spheroids and

the early divergence of these subclones from the primary tumor suggests that we are currently

underestimating the tumor genomic landscape.

In addition to the importance of genetic differences between tumor cells in primary

tissue and those in ascites, knowledge regarding the genetic heterogeneity within the tumor

cells in ascites would be valuable. Although not thoroughly studied, the genetic diversity of

tumor cells in an ascites may have a large impact on tumor relapse and metastasis, given that

transcoelomic spread is the primary route of metastasis in ovarian cancer. However, there has

been no attempt to discover the genetic heterogeneity of individual tumor spheroids. In this

study, we evaluated 10 individual tumor spheroids, five of which contained sufficient tumor

cells for the analysis. Although we observed genetic heterogeneity of the individual ascites

spheroids, a follow-up study should analyze at least a few tens of individual tumor spheroids

per patient to find a clear signature of the genetic heterogeneity in an ascites.

Conclusion

In this study, we performed genome-wide sequence analysis of the primary tumor

and the associated tumor spheroids in the malignant ascites of an EOC patient. We analyzed

genetic heterogeneity in the primary tumor and tumor spheroids through multi-region

sequencing and the laser-aided cell isolation technique [12]. From the sequencing data, we

discovered clonal or subclonal somatic CNAs and SNVs, based on which we constructed

phylogenetic trees and inferred the evolutionary history of tumor cells in the patient. As a

result, we found that the tumor cells in the malignant ascites were an independent lineage

from the primary tumor. The phylogenetic analysis showed that the lineage branched before

the evolution of the cancer cells at the primary tissues, which suggests that analyzing

malignant ascites might be used to detect ovarian cancer or metastasis in the early stage. In

summary, the genetic plasticity and similarity between a primary tumor and associated tumor

spheroids are still not clear, and yet, the nature of the similarity may have profound

implications for both tumor progression and therapeutic outcomes in ovarian cancer.

Therefore, future prospective studies profiling the genomic information of primary ovarian

tumors, distant metastatic tumors, and tumor spheroids to determine the direction of tumor

evolution and metastasis of ovarian cancer are warranted.

Methods

Patient information and sample preparation

A 42 yr old female patient diagnosed with primary high-grade serous ovarian cancer (Grade 3,

stage IIIC) presented with malignant ascites and peritoneal seeding. Both primary tissues and

malignant ascites were collected during primary debulking surgery. Fresh primary tissues and

tumor cell clusters were mounted onto ITO-coated glass slides. Six samples were taken

randomly from the solid portions of right ovary and only one from left ovary. Blood was

collected to serve as the normal control. Ten tumor cell clusters were collected from the

malignant ascites and fixed in 10% (v/v) formaldehyde. This study was approved by the

Institutional Review Board (IRB) at Seoul National University Hospital (Registration number:

1305-546-487) and performed in compliance with the Helsinki Declaration. We obtained

informed consent from the patient prior to primary debulking surgery to be used in research.

Laser-aided isolation of tumor spheroids and their whole-genome amplification

Previously, we developed and published a laser-aided cell isolation technique [12], and

designed two different pieces of software written in Python scripts and available at Github

(https://github.com/BiNEL-SNU/PHLI-seq). Isolation of tumor spheroids was performed as

described in the prior publication. In brief, an infrared laser was applied to the target area,

vaporizing Indium Tin Oxide (ITO) layer and discharging the targeted tumor spheroid on the

region. We used glass slides with a 100-nm-thick ITO layer.

The 8-strip PCR tube caps for the retrieval of tumor spheroids were pre-exposed under O_2

plasma for 2 minutes. The tumor spheroids were lysed using proteinase K (cat no. P4850-

1ML, Sigma Aldrich) according to the manufacturer's directions after the PCR tubes were

centrifuged. For whole-genome amplification, we used GE's Illustra Genomiphi V2 DNA

amplification kit (cat no. 25-6600-30). We added 0.2 µl of SYBR green I (Life Technologies)

into the reaction solution for real-time monitoring of the amplification (Fig. 2B). All

amplified products were purified using Beckman Coulter's Agencourt AMPure XP kit (cat no.

A63880) immediately following the amplification. To prevent carry-over contamination, the

pipette tip, PCR tube, and cap for the reaction were stored in a clean bench equipped with UV

light and treated with O_2 plasma for 2 minutes before use. Additionally, we monitored the

real-time amplification of non-template controls to ensure that no contaminants were

transferred.

Sequencing library preparation, whole-genome, and whole-exome sequencing

The whole-genome amplified products or genomic DNA were fragmented using an EpiSonic

Multi-Functional Bioprocessor 1100 (Epigentek) to generate DNA fragments with 250-bp on

average. The fragmented products underwent Illumina library preparation using Celemics

NGS Library Preparation Kit (LI1096, Celemics, Seoul, Korea) for the whole-genome

sequencing library preparation, and SureSelectXT (Agilent, CA, US) for whole-exome

sequencing. DNA purification was performed by TOPQXSEP MagBead (XB6050, Celemics,

Seoul, Korea), and DNA libraries were amplified using the KAPA Library Amplification Kit

(KAPA Biosystems, KK2602). Finally, the products were quantified by TapeStation 2200

(Agilent, CA, US). We used HiSeq 2500 150 PE (Illumina) to generate 1 Gb/sample for

whole-genome sequencing and 5 Gb/sample for whole-exome sequencing, respectively.

Detecting copy number alterations

We used low-depth whole-genome sequencing data and the variable-size binning method [29]

to estimate the CNAs of the samples. Briefly, the whole genome was divided into 15,000

variable-sized bins (median genomic length of bin = 184 kbp), in which each bin had an

equal expected number of uniquely mapped reads. Then, each sequence read was assigned to

each bin followed by Lowess GC normalization to obtain the read depth of each bin. The

copy number was estimated by normalizing the read depth of each bin by the median read

depth of the reference DNA.

Detecting Single Nucleotide Variants

GATK (v3.5-0) IndelRealigner and BaseRecalibrator were used to locally realign reads

around the Indel and recalibrate the base quality score of BAM files [30]. Then, GATK

UnifedGenotyper, Varscan, and MuTect were used and combined the results to avoid false-

positive variant calls [31]. First, GATK UnifiedGenotyper was used with default parameters

followed by GATK VariantRecalibrator to obtain filtered variants [30]. Data of primary tissue

samples and ascites tumor spheroid samples were processed together to produce a single vcf

file. dbSNP build 137, HapMap 3.3, Omni 2.5, and 1000G phase1 were used as the training

data for variant recalibration. Also, annotation data including QD, MQ, FS,

ReadPosRankSum, and MQRankSum were used for the training. Variants detected in the

paired blood sample of the cancer patient were removed to produce the final list of GATK

called variants. Varscan2 [32] (ver 2.3.7) and Mutect [33] (ver 1.1.4) were used with default

parameters to produce the lists of Varscan and MuTect called variants, respectively. Here,

paired blood read data was also used to remove germline variants.

Among the variants from the three callers, variants called by at least two callers were

collected to obtain intra-sample double-called sites. We could reduce false-positive variant

caused by NGS errors by considering only double-called variants for subsequent analysis [31].

Among the intra-sample double called sites, variants found in at least two samples were

collected to remove WGA (whole genome amplification) errors, and the genomic loci with

the resultant variants were considered confident sites. Finally, a variant in the confident sites

was considered to be true if one of the three variant callers detected the variant at the locus

and the allele count of the variant was significantly larger than that of the other non-reference

bases (Fisher's exact test, $p < 10^{-3}$). The overall process is visually described in Additional

file5: Figure S1.

Abbreviations:

EOC: Epithelial ovarian cancer

ITH: Intra-tumor heterogeneity

CNA: Copy number alteration

SNV: Single-nucleotide variant

TME: Tumor microenvironment

WGS: Whole-Genome Sequencing

WES: Whole-exome sequencing

HGS: High-grade serous

ITO: Indium tin oxide

MDA: Multiple displacement amplification

VAF: Variant allele frequency

CI: Clonality index

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board (IRB) at Seoul National

University Hospital (Registration number: 1305-546-487) and performed in compliance with

the Helsinki Declaration. We obtained informed consent from the patient prior to primary

debulking surgery to be used in research.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request

Competing interests

The authors declare that they have no competing interests

Acknowledgements

Not applicable

References

1.   Reid BM, Permuth JB, Sellers TA. Epidemiology of ovarian cancer: a review. Cancer

Biol Med. 2017;14:9-32.

2.   Jessmon P, Boulanger T, Zhou W, Patwardhan P. Epidemiology and treatment patterns

of epithelial ovarian cancer. Expert Rev Anticancer Ther. 2017;17:427-37.

3.   Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, et al. Clonal

evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing.

Nature. 2012;481:506-10.

4.   Gerlinger M, Rowan AJ, Horswell S, Math M, Larkin J, Endesfelder D, et al.

Intratumor heterogeneity and branched evolution revealed by multiregion sequencing.

N Engl J Med. 2012;366:883-92.

5.   Lee JY, Yoon JK, Kim B, Kim S, Kim MA, Lim H, et al. Tumor evolution and

intratumor heterogeneity of an epithelial ovarian cancer investigated using next-

generation sequencing. BMC Cancer. 2015;15:85.

6.   Swanton C. Intratumor heterogeneity: evolution through space and time. Cancer Res.

537   2012;72:4875-82.

538   7.   Bilen MA, Hess KR, Campbell MT, Wang J, Broaddus RR, Karam JA, et al.

539   Intratumoral heterogeneity and chemoresistance in nonseminomatous germ cell tumor

540   of the testis. Oncotarget. 2016;7:86280-9.

541   8.   Greaves M. Evolutionary determinants of cancer. Cancer Discov. 2015;5:806-20.

542   9.   Tan DS, Agarwal R, Kaye SB. Mechanisms of transcoelomic metastasis in ovarian

543   cancer. Lancet Oncol. 2006;7:925-34.

544   10.   Kim S, Kim B, Song YS. Ascites modulates cancer cell behavior, contributing to

545   tumor heterogeneity in ovarian cancer. Cancer Sci. 2016;107:1173-8.

546   11.   Smolle E, Taucher V, Haybaeck J. Malignant ascites in ovarian cancer and the role of

547   targeted therapeutics. Anticancer Res. 2014;34:1553-61.

548   12.   Kim S, Lee AC, Lee H, Kim J, Jung Y, Ryu HS, et al. Constructing and visualizing

549   cancer genomic maps in 3D spatial context by Phenotype-based High-throughput

550   Laser-aided Isolation and Sequencing (PHLI-seq). bioRxiv 2018; doi:

551   10.1101/278010

552   13.   Zack TI, Schumacher SE, Carter SL, Cherniack AD, Saksena G, Tabak B, et al. Pan-

553   cancer patterns of somatic copy number alteration. Nat Genet. 2013;45:1134-40.

554   14.   Park JT, Li M, Nakayama K, Mao TL, Davidson B, Zhang Z, et al. Notch3 gene

555   amplification in ovarian cancer. Cancer Res. 2006;66:6312-8.

556   15.   Derouet M, Wu X, May L, Hoon Yoo B, Sasazuki T, Shirasawa S, et al. Acquisition of

557   anoikis resistance promotes the emergence of oncogenic K-ras mutations in colorectal

558   cancer cells and stimulates their tumorigenicity in vivo. Neoplasia. 2007;9:536-45.

559   16.   Rytomaa M, Martins LM, Downward J. Involvement of FADD and caspase-8

560   signalling in detachment-induced apoptosis. Curr Biol. 1999;9:1043-6.

17.   Ahmed AA, Etemadmoghadam D, Temple J, Lynch AG, Riad M, Sharma R, et al.

Driver mutations in TP53 are ubiquitous in high grade serous carcinoma of the ovary.

J Pathol. 2010;221:49-56.

18.   Wiegand KC, Shah SP, Al-Agha OM, Zhao Y, Tse K, Zeng T, et al. ARID1A

mutations in endometriosis-associated ovarian carcinomas. N Engl J Med.

2010;363:1532-43.

19.   Welcsh PL, King MC. BRCA1 and BRCA2 and the genetics of breast and ovarian

cancer. Hum Mol Genet. 2001;10:705-13.

20.   Xi Y, Liu C, Xin X. Association between a single nucleotide polymorphism in the

TP53 region and risk of ovarian cancer. Cell Biochem Biophys. 2014;70:1907-12.

21.   Yin X, Jing Y, Cai MC, Ma P, Zhang Y, Xu C, et al. Clonality, heterogeneity and

evolution of synchronous bilateral ovarian cancer. Cancer Res. 2017;77:6551-61.

22.   Schultheis AM, Ng CK, De Filippo MR, Piscuoglio S, Macedo GS, Gatius S, et al.

Massively parallel sequencing-based clonality analysis of synchronous

endometrioidendometrial and ovarian carcinomas. J Natl Cancer Inst.

2016;108:djv427.

23.   Yin M, Li X, Tan S, Zhou HJ, Ji W, Bellone S, et al. Tumor-associated macrophages

drive spheroid formation during early transcoelomic metastasis of ovarian cancer. J

Clin Invest. 2016;126:4157-73.

24.   Schwarz RF, Ng CK, Cooke SL, Newman S, Temple J, Piskorz AM, et al. Spatial and

temporal heterogeneity in high-grade serous ovarian cancer: a phylogenetic analysis.

PLoS Med. 2015;12:e1001789.

25.   Choi YJ, Rhee JK, Hur SY, Kim MS, Lee SH, Chung YJ, et al. Intraindividual

genomic heterogeneity of high-grade serous carcinoma of the ovary and clinical

utility of ascitic cancer cells for mutation profiling. J Pathol. 2017;241:57-66.

26.   Marusyk A, Polyak K. Tumor heterogeneity: causes and consequences. Biochim
Biophys Acta. 2010;1805:105-17.

27.   Polyak K, Haviv I, Campbell IG. Co-evolution of tumor cells and their
microenvironment. Trends Genet. 2009;25:30-8.

28.   Gao R, Davis A, McDonald TO, Sei E, Shi X, Wang Y, et al. Punctuated copy number
evolution and clonal stasis in triple-negative breast cancer. Nat Genet. 2016;48:1119-
30.

29.   Baslan, T. *et al.* Genome-wide copy number analysis of single cells. Nat. Protoc. **7,**
1024–1041 (2012).

30.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. Genome Res. **20,** 1297–1303 (2010).

31.   Lodato, M. A. *et al.* Somatic mutation in single human neurons tracks developmental
and transcriptional history. Science (80-.). **350,** 94–98 (2015).

32.   Koboldt, D. C. *et al.* VarScan 2□: Somatic mutation and copy number alteration
discovery in cancer by exome sequencing VarScan 2□: Somatic mutation and copy
number alteration discovery in cancer by exome sequencing. Genome Res. **22,** 568–
576 (2012).

33.   Cibulskis, K. *et al.* Sensitive detection of somatic point mutations in impure and
heterogeneous cancer samples. Nat. Biotechnol. **31,** 213–219 (2013).

607
608

<u>Figure 1: An overview of individual tumor spheroid isolation from malignant ascites.</u> (A)

A malignant ascites was collected during a primary debulking surgery. Tumor spheroids in

the malignant ascites were purified, fixed and prepared on a discharging layer (Indium Tin

Oxide (ITO), 100 nm in thickness)-coated glass slide. (B) The laser isolation technique was

used to isolate individual tumor spheroids. This technique utilizes an IR pulsed laser, which

vaporizes the discharging layer on the glass slide. Using this technique, ten individual tumor

spheroids were isolated from the slide. The isolated cells underwent WGA and sequencing.

(C) The images before and after isolation demonstrate that the targeted tumor spheroids in the

malignant ascites were specifically isolated without disturbing the neighboring cells. The

scale bars represent 100 μm.

<u>Figure 2: WGA of the isolated tumor spheroids and several quality metrics of the</u>

<u>amplified products.</u> (A) MDA was performed to amplify the DNA in each tumor spheroid.

MDA amplified tumor spheroid DNA 10^3- to 10^4-fold. (B) The amplification process was

monitored by observing the fluorescence signal in each reaction. A non-template control was

included in the reaction to testify carry-over contamination. The results showed that there was

no or a negligible amount of carry-over contamination. (C, D) The distributions of the

normalized read depth and VAF reflect the quality of the WGA products. Compared with the

distributions of the amplified products from single cells, the distributions of the tumor

spheroids were similar to those of the primary tissues. This indicated that the amplified

products from the tumor spheroids had a negligible amount of WGA artifacts.

Figure 3: CNA analysis based on the genetic subclones of the tumor cells identified via low-depth WGS. (A) A genome-wide CNA analysis was performed using the low-depth WGS data. Each row represents each sample, and the samples were reordered by the hierarchical clustering method. The clustering analysis generated three major clusters, which were named Primary clone (red), Ascites clone 1 (yellow), and Ascites clone 2 (green). The clear differentiation of the CNA profiles between the Primary clone and Ascites clones implied that the tumor spheroids in the Ascites clones were not derived from the tumor cells in the Primary clone but from another independent tumor lineage. (B) Representation of the CNA profiles in detail at several regions for RO1, AC1, and AC4. The three samples exhibited both shared and exclusive CNAs. For example, deletion of FAT1 (1^{st} column) and amplification of MYC, CYC1, and PARP10 (2^{nd} column) were shared in every sample. However, the amplification of KDM5A (3^{rd} column) and NOTCH3 (4^{th} column) was exclusive to the Primary clone. This might indicate that the FAT1, MYC, CYC1, or PARP10 alterations conferred a growth advantage to the common ancestor of the Primary clone and Ascites clones. In contrast, the KDM5A or NOTCH3 amplifications might cause branching from the common ancestor and proliferation of the Primary clone

Figure 4: SNV analysis based on the WES data. The WES data from the primary tissue samples and tumor spheroids were used to analyze the SNVs. The results showed that a significant portion of the SNVs was shared in the Primary clone and Ascites clone 1. At the same time, the Primary clone and Ascites clone 1 had unique mutations. This result suggests that the two clones might have branched from a common ancestor. Ascites clone 2 was excluded from the analysis because the tumor spheroids in Ascites clone 2 were presumed to contain a large number of normal cells in each tumor spheroid.

Figure 5: Analysis of the allele frequency to infer the cellular composition of each

sample. The VAF distribution was plotted for each sample from the (A) Primary site and (B,

C) Ascites. The mutations were categorized into common, primary-only or ascites-only

mutations. Common mutations were somatic SNVs, which were detected in both the Primary

clone and Ascites clones, and primary-only and ascites-only mutations, which were shared

somatic SNVs detected only in the Primary clone and Ascites clones, respectively. The results

showed that most of the VAF distributions from Ascites clone 2 were located at a much lower

range than those from the Primary clone and Ascites clone 1. This suggests that the tumor

spheroids in Ascites clone 2 had a large proportion of normal cells in each tumor spheroid.

Figure 6: Constructing phylogenetic trees. Phylogenetic trees were constructed using both

the (A) CNA profiles and (B) SNV profiles. The two trees presented similar topologies and

indicated that the Primary clone and Ascites clone 1 were derived from one ancestral clone at

the early stage of cancer development. In addition, the phylogenic trees indicated that the

analyzed tumor spheroids were not derived from the primary tumor cells that were present at

the time of sampling.

Figure 7: The inferred evolutionary history of the tumor and the Circos plot of the

major subclones. (A) Based on the sequencing data from the primary tissue samples and the

tumor spheroids from the ascites, the evolutionary trajectory was inferred. The tumor was

initiated at the right ovary to generate the ancestral clone. With the further accumulation of

mutations, the ancestral clone evolved into two subclones, the first of which was found in the

right ovary and metastasized to the left ovary. The second subclone shed into the ascites TME

and became extinct or dominated by the first subclone in the right ovary. Eventually, the

Ascites subclone moved to the peritoneal cavity. (B) The Circos plot presents the genome-

wide alterations in the ancestral, Primary, and Ascites subclones. For the SNVs, the black, red,

and yellow bars represent the ancestral, primary-only, and ascites-only mutations,

respectively. The tumor cells acquired the ancestral mutations before dividing into the

Primary and Ascites clones. After division, the Primary and Ascites clones acquired lineage-

specific SNVs. For the CNAs, the red and blue bars represent amplification and deletion,

respectively.

A

Single nucleotide variants

■ Mutated　□ Not mutated　■ Not confident

Primary tissue samples

RO 3
LO
RO 6
RO 4
RO 7
RO 1
RO 2
RO 5

Primary clone

Individual tumor spheroids

AS 1
AS 2
AS 3
AS 7
AS 8

Ascites clone 1

AS 10
AS 9
AS 4
AS 5
AS 6

Ascites clone 2

KRAS (C>T)　SEPT6 (T>A)　TP53 (T>C)　ARID1A (A>G)　PLCG2 (C>A)

B

Primary-only　Common　Ascites-only

61 mutations (35.7%)　66 mutations (38.6%)　44 mutations (25.7%)

TP53
ARID1A
PLCG2

KRAS
SEPT6

Figure 6

Figure 7