Genetic dissection of assortative mating behavior

Richard M. Merrill [1,2,3,*], Pasi Rastas[2], Maria C. Melo[3,4], Sarah Barker[2], John Davey[2,5], W. Owen McMillan[3] & Chris D. Jiggins[2]

1 Division of Evolutionary Biology, Ludwig-Maximilians-Universität, München, Germany **2** Department of Zoology, University of Cambridge, Cambridge, UK **3** Smithsonian Tropical Research Institute, Panama City, Panama **4** IST Austria, Klosterneuburg, Austria **5** Department of Biology, University of York, York, UK

 *merrill@bio.lmu.de

Abstract

The evolution of new species is made easier when traits under divergent ecological selection are also mating cues. Such ecological mating cues are now considered more common than previously thought, but we still know little about the genetic changes underlying their evolution, or more generally about the genetic basis for assortative mating behaviors. The warning patterns of *Heliconius melpomene* and *H. cydno* are under disruptive selection due to increased predation of non-mimetic hybrids, and are used during mate recognition. We carried out a genome-wide quantitative trait locus (QTL) analysis of preference behaviors between these species and showed that divergent male preference has a simple genetic basis. Three QTLs each explain a large proportion of the differences in preference behavior observed between the parental species. Two of these QTLs are on chromosomes with major color pattern genes, including one that is tightly associated with the gene *optix*. Different loci influence different aspects of attraction, suggesting that behavioral isolation in *Heliconius* involves the evolution of independently segregating modules, similar to those for the corresponding wing pattern cues. Hybridization and subsequent sharing of wing pattern loci has played an important role during adaptation and speciation in *Heliconius* butterflies. The existence of large effect preference loci could similarly assist the evolution of novel behavioral phenotypes through recombination and introgression, and should facilitate rapid speciation.

Introduction

During ecological speciation, reproductive isolation evolves as a result of divergent natural selection [1]. Although ecological barriers can reduce gene flow between divergent populations, speciation normally requires the evolution of assortative mating [1, 2]. This is made easier if traits under divergent ecological selection also contribute to assortative mating, as this couples ecological and mating traits [2-5]. Ecologically relevant mating cues (sometimes known as 'magic traits' [2, 4]) are now predicted to be widespread in nature [4, 6], and the last few years have seen considerable progress in our understanding of their genetic basis. For example, studies have explored the genetic basis of beak shape in Darwin's finches [7], body shape in sticklebacks [8], cuticular hydrocarbons in *Drosophila* [9], and wing patterns in *Heliconius* butterflies [10-12]. However, the extent to which these traits contribute to assortative mating depends on the evolution of corresponding preference behaviors. This process will be greatly influenced by the underlying genetics. Both the substitution of large effect preference alleles, and physical linkage will strengthen linkage disequilibrium between cue and preference, thereby reducing the effects of recombination that might otherwise impede the evolution of assortative mating. We still know little about the process by which ecological traits are co-opted as mating cues, and in particular how matching cues and preference behaviors are controlled genetically (but see [13]).

In central Panama, the butterfly *Heliconius melpomene rosina* is a precise Müllerian mimic of *H. erato* and normally occurs in forest-edge habitats, whereas the closely related species *H. cydno chioneus* mimics *H. sapho* and is more common in closed-forest habitats, although *H. melpomene* and *H. cydno* are often seen flying together (Fig. 1a & b). Hybrids are viable but occur at very low frequency in the wild

(estimated at ~0.001 [14]), consistent with strong assortative mating shown in

insectary experiments. Specifically, heterospecific mating was not observed in 50

choice and no-choice trials between Panamanian *H. melpomene* and *H. cydno* ([15,

16]; see also [17]).

The amenability of *Heliconius* color patterns to experimental manipulation has

led to the demonstration that color pattern is both under strong disruptive selection

due to predation [18], and also that males prefer live females and paper models with

the same color pattern as themselves [15]. These results led Servedio and

colleagues [4] to conclude that, unlike other putative examples, in *H. melpomene*

rosina and *H. cydno chioneus* both criteria for a magic trait have been confirmed with

manipulative experiments. Although female preferences undoubtedly contribute to

assortative mating [19-22], male preferences act first in these species such that

strong observed male discrimination against heterospecific females will have a

disproportionate absolute contribution to overall reproductive isolation [22].

Crossing experiments have shown that the shift in mimetic warning pattern

between *H. melpomene rosina* and *H. cydno chioneus* is largely controlled by three

major effect loci [23]. Candidate genes underlying these loci have now been

identified: the transcription factor *optix* controls red patterns [10], the *WntA* gene

controls forewing shape [11] and yellow patterns map to the gene *cortex* [12]. In

addition, a further locus, *K*, segregates in crosses between *H. melpomene rosina* and

H. cydno chioneus with more minor effect [23, 24]. Further modularity occurs within

these loci. For example, different regulatory elements of optix each result in distinct

red pattern elements [25]. The modular nature of individual color pattern loci and their

functionally sufficient enhancers means that they can be combined to produce

considerable phenotypic diversity [25, 26]. These loci are large-effect 'speciation

genes', in that they control traits that generate strong reproductive isolation. To complement our extensive knowledge of the genetics of their color pattern cues, here we explore the genetics of male preference behaviors between *H. melpomene* and *H. cydno*.

Results

We studied male mating preference among F1 and backcross hybrid families between *H. melpomene rosina* and *H. cydno chioneus*, in standardized choice trials [20]. We introduced individual males into an experimental cage and recorded courtship directed towards two virgin females, one of each species. In total, we collected data from 1347 behavioral trials, across 292 individuals. Multiple trials were performed for each backcross male, from which we determined the relative courtship time directed towards *H. melpomene* and *H. cydno* females, as well as the proportion of trials in which courtship was initiated towards each species. As reported previously [20], F1 males have a strong preference for the red *H. melpomene* females, and little segregation in mate preference is observed among the backcross to melpomene (no significant difference from pure *H. melpomene* males: $2\Delta\ln L = 1.33$, *d.f.* $= 1$, $P > 0.2$), implying that *melpomene* mate preference alleles are dominant. In contrast, courtship behavior segregates among *H. cydno* backcross males, permitting analysis of the genetic basis for this mating behavior (Fig. 1C).

Three loci contribute to species differences in preference behavior. We used a genome-wide quantitative trait locus (QTL) mapping approach to identify the genomic regions underlying divergence in mate attraction. Previous studies of *Heliconius* behavior have only considered individual chromosomes [24] or individual loci [20,

27], and may have missed important contributions made by other parts of the genome. Linkage maps were constructed from genotype data of 331 backcross-to-cydno individuals and their associated parents [28], including 146 individual males for which we had recorded attraction behaviors. We identified three unlinked QTLs on chromosomes 1, 17 and 18 associated with variation in the relative amount of time males spent courting red *H. melpomene* and white *H. cydno* females (Fig. 2A). Of these, one is tightly linked to the optix locus on chromosome 18, which controls the presence/absence of a red forewing band. Given the sample sizes feasible in *Heliconius*, our analysis lacks the power to resolve smaller effect QTLs. Nevertheless, modeling supports additive effects of all three detected loci (Table 1), and in our mapping population these three QTLs together explain ~60% of the difference between the parental species (Fig. 2B).

Preference QTL are of large effect. Individually, the measured effect of each QTL was large, explaining between 23 and 31% of the difference between males of two parental species (Fig. 2B). However, in studies with relatively small sample sizes such as ours (n = 139), estimated effects of QTL are routinely over-estimated (a phenomenon known as the "Beavis effect", after [29]). This is because effect sizes are determined only after significance has been determined, and QTL with artificially high effect sizes – due to variation in sampling – are more likely to achieve 'significance'. To determine the extent to which the effects of our QTL may be over-estimated, we simulated QTL across a range of effect sizes, and compared the distribution of measured effects for all simulations to those which would be significant in our analysis. Our simulations suggest that the reported effects of our QTL are not greatly over-estimated. We first considered what proportion of 'significant' simulations

would be smaller than our empirically measured effects (Fig. 3A). With the possible

exception of the QTL on chromosome 17, this suggests that the true effect sizes of

our QTL are likely to be large. A highly conservative threshold of 95% would suggest

that the QTL on chromosome 1 and 18 explain at least 10% and 20% of the

difference in behavior between the parental species, respectively. Adopting the

median values, our simulations would suggest true effects of 25%, 15% and 30%, or

greater, for the QTL on chromosomes 1, 17 and 18, respectively. Given simulated

effect sizes similar to those measured empirically, there was little bias among

simulation runs that achieved the genome-wide significance threshold (though there

is a slight deviation for the QTL on chromosome 17) (Fig. S1).

Although our simulations suggest the effects we have measured are

reasonable, ideally we would estimate effect sizes from a population of individuals

that were not used to determine significance. In evolutionary biology, follow-up

experiments such as this are uncommon; collecting phenotypic data across a large

number of hybrid individuals is often a considerable undertaking, and this is similarly

true for *Heliconius* behaviors. Nevertheless, we were able to follow-up our results for

the QTL on chromosome 18, using a sample of 35 individuals for which preference

behavior was measured, but for which we were unable to generate genotype data

(and so were not included in our initial QTL analysis). As reported above, the QTL on

chromosome 18 is in very tight linkage with the B/D color pattern locus, so that

segregation of the red forewing band can be used to determine genotype at this

locus. This analysis supports our previous result that the QTL on linkage group 18 is

of large effect: among these 35 hybrid males, the B/D locus explains 27% of the

difference in behavior between the parental species (c.f. 31% for the larger mapping

population, Fig. 3B).

The preference QTL in linkage with warning pattern genes influence initial attraction. The time *Heliconius* males spend courting a particular female might depend not only on male attraction, but also on the female's response (and in turn his response to her behavior) [19] . To test whether the QTLs we reported above control male approach behaviors (as opposed to other traits that may influence courtship), we repeated our genome-wide QTL analysis but this time considered the number of trials in which courtship was initiated towards cydno, and then separately, towards melpomene females. These analyses each revealed a significant QTL, one on chromosome 1 associated with initial attraction towards *H. cydno* females (LOD = 3.44, $P \sim 0.029$), and another on chromosome 18 for initial attraction towards *H. melpomene* females (LOD = 3.10, $P \sim 0.049$). These are coincident with QTLs on chromosome 1 and 18 for relative courtship time (Fig. S2).

Different preference QTL affect different aspects of behavior. Our analyses of courtship initiation confirmed that male attraction behavior is influenced by QTLs on chromosomes 1 and 18, but we were unable to detect a significant effect of chromosome 17. This is in direct contrast to our analysis of the relative courtship time, in which the influence of chromosome 17 was well supported. Although we found no evidence of pairwise interactions between individual QTLs in our model of relative courtship time, genome scans considering individuals with alternative genotypes at the QTL on chromosome 18 separately revealed a significant QTL on chromosome 17 (LOD = 4.16, $P < 0.05$) for heterozygous (i.e. LG18@0cM = cyd:melp), but not for homozygous (i.e. LG18@0cM = cyd:cyd) males (Fig. S3). Together, these results suggest that alleles on chromosome 17 may have little effect

on initial visual attraction but influence subsequent behaviors after courtship has

been initiated.

The male preference QTLs we have identified must influence differences in

male attraction towards red *H. melpomene* females, or white *H. cydno* females, or

towards both females types. To further explore the influence of segregating alleles at

these loci we considered the influence of all three QTLs on courtships directed

towards each female type separately. We have already robustly established a

significant effect of these loci on variation in the relative amount of time males spent

courting each female type (see Fig. 2A). Consequently, although we corrected for

multiple testing arising from considering three QTL across the two data sets, in these

analyses we did not account for multiple segregating loci across the entire genome

(in contrast to the results reported above). This greatly increases our power to detect

any influence of the QTLs on attraction towards the two species individually, but also

increases the likelihood of false positives. The QTL on chromosome 1 influenced the

number of courtships directed towards *H. cydno* females ($F_{1,145}$ = 11.88, P < 0.01),

but had no significant effect on how males behaved towards *H. melpomene* females

($F_{1,145}$ = 1.63, P > 0.1). In contrast, the QTL on chromosome 17 influenced the

degree of courtship directed towards *H. melpomene* ($F_{1,145}$ = 8.97, P = 0.018), but not

H. cydno females ($F_{1,145}$ = 0.93, P > 0.1). Similarly, the QTL on chromosome 18 had

a significant effect on courtships directed towards *H. melpomene* ($F_{1,145}$ = 10.16, P =

0.012) females (though we note that prior to Bonferroni correction there is also some

support for an effect on courtships directed towards *H. cydno* females: $F_{1,145}$ = 6.74,

P = 0.01).

Discussion

We have shown that the genetic architecture of mate preference behaviors can facilitate speciation in *Heliconius*. Specifically, we demonstrate that just three regions of the genome are largely responsible for an important component of behavioral isolation. For a behavior, presumably including a large environmental component, and measured in outdoor insectaries, these results are striking. Both our simulations and replication support the existence of large effect QTLs controlling this important interspecific difference in behavior. Even if we adopt an especially cautious approach, the QTLs on chromosomes 1 and 18 would explain *at least* 10% and 20% of the difference in male preference behavior, respectively. However, a follow-up analysis, exploiting individuals that were not used to determine significance (thereby evading the Beavis effect), suggests that these estimates are overly conservative; these data explicitly reinforce our initial estimate for the QTL on chromosome 18, which explains ~30% of the difference between parents. Additional smaller effect loci undoubtedly also contribute to variation in male preference, which we would be unlikely to detect without very large sample sizes (a caveat shared with many QTL studies of ecologically relevant behaviors e.g. [13, 29, 30]). Regardless, our results suggest that during speciation, divergence between populations in both mating cue and the corresponding preference behaviors can have a surprisingly simple genetic architecture. This architecture will facilitate the evolution of reproductive isolation in the face of gene flow by both evading the effects of drift and strengthening genetic associations (*i.e.* linkage disequilibrium) between cue and preference traits, even if the two are not physically linked [2, 6, 31].

Two of the behavioral QTL we have identified are situated on chromosomes with major color pattern loci (chromosome 1 includes the *K* locus, and chromosome

18 includes the *optix* locus). By further strengthening the genetic associations

between these key components of behavioral isolation, physical linkage between

cues and preferences is expected to facilitate speciation with gene flow [32]. Both

optix and the *K* locus have previously been associated with variation in *Heliconius*

courtship behaviors [20, 24, 27]. Nevertheless, we have not previously been able to

robustly estimate the position of QTLs along the chromosome. The QTL we identify

on chromosome 1 is not tightly linked to the *K* locus. It remains to be seen whether

this QTL underlies the association between male preference behavior and the *K*

locus phenotype (a shift between white and yellow color pattern elements) previously

observed in crosses between *H. cydno* and *H. pachinus* [24], and within a

polymorphic population of *H. cydno* [27]. In contrast, our results reveal that the QTL

for male attraction on chromosome 18 is tightly linked to the *optix* locus, which

controls presence/absence of a red forewing. The mechanistic basis for linkage of

trait and preference loci remains unclear. There is no evidence for an inversion at this

locus [28]; it also seems unlikely that the same mutations control both wing pattern

and the corresponding attraction behavior. However, *optix* is known to function during

eye and neural development in *Drosophila* [33], and is expressed in the optic lobe

and medulla of pupal *Heliconius* [34], so it is plausible – if not likely [35] – that the two

traits could be controlled by different regulatory elements of the same gene. Ours

joins a handful of examples where genetic linkage between cue and preference has

been reported [13, 20, 24, 27, 36, 37], which is expected to facilitate speciation with

gene flow [3, 32].

In addition to a very simple genetic architecture of male preference behavior,

different aspects of preference behavior are controlled by different QTL. Differences

associated with QTL1 and QTL17 in the relative amount of time spent courting each

female type are driven by differences in attraction to either *H. cydno* or *H. melpomene*, respectively, rather than both species. QTL18 also influences attraction to *H. melpomene* much more strongly than to *H. cydno* females. This genetic modularity can facilitate evolutionary change and innovation by providing a route for rapid evolution of novel phenotypes [29, 38]. In *Heliconius*, the observed modularity should allow different aspects of mating behavior to evolve independently. It might also allow novel composite behavioral preferences to arise through hybridization and recombination. Indeed, there is some evidence that this has occurred during hybrid speciation in *Heliconius*, where hybrids between *H. melpomene* and *H. cydno* have a novel preference similar to that of the natural hybrid species *Heliconius heurippa* [39].

Others factors, including habitat-use, female choice and intrinsic and extrinsic hybrid incompatibilities, contribute to reproductive isolation between *H. cydno* and *H. melpomene*. Each of these can impose a strong barrier to gene flow in their own right; however, because they act early in the life cycle, habitat choice and male preferences probably represent the most important contemporary barriers to gene flow between *H. cydno* and *H. melpomene* (with absolute contributions estimated at 0.66 and 0.32, respectively [22]). Of course, the current importance of isolating barriers may not reflect a barrier's historical importance, but in *Heliconius*, there are many examples of within-species divergence in color pattern and corresponding male preference (e.g. [27, 40, 41]), suggesting that behavioral isolation due to male preferences evolves early during divergence.

Divergence in male preferences is likely driven by initial divergence in female wing pattern. In *Heliconius*, female re-mating is a rare event [42], and males must compete to find virgin females within a visually complex environment [21]. Divergence in female (and male) wing patterns is driven primarily by strong selection

for mimicry, and is likely to impose divergent sexual selection on male preferences to improve their ability to find receptive females. Color pattern therefore effectively acts as a female 'preference' trait (sensu [43], p.17) in that it 'biases … fertilization success toward certain male phenotypes" (in this case particular behavioral phenotypes). This is important because strong ecological selection on female traits, which biases mating towards particular male phenotypes, will rapidly drive diversification of the corresponding male trait [5]. This can provide an effective route to speciation, as has been widely appreciated in examples of assortative mating driven by sensory drive, such as in cichlid fishes [44]. However, it is perhaps less well appreciated, that morphological traits under ecological selection (such as *Heliconius* wing patterns) might impose divergent sexual selection on male preferences in a similar fashion.

The male preferences observed here may have additionally been strengthened as result of reinforcement. Non-mimetic hybrids between *H. melpomene* and *H. cydno* suffer increased predation [18], as well as reduced mating success [16] and fertility [45]. Males make a considerable reproductive investment by donating a nutrient-rich spermatophore during mating [46, 47], so indirect selection against poorly adapted hybrids could strengthen divergent male preferences. Consistent with this, *H. melpomene* males from French Guiana, outside the range of *H. cydno*, are less choosy than males from Panama, where the species co-occur and are known to occasionally hybridise [15]; and similar patterns of reproductive character displacement have been observed elsewhere in the *melpomene-cydno* clade [48]. However, male preferences are also observed between *Heliconius* populations with few opportunities for hybridization e.g. [40]. It seems likely therefore

that both direct and indirect selection contribute to the evolution of male preferences

in *Heliconius*.

Under either selective regime, the genetic architecture we demonstrate here will

promote the evolution of behavioral isolation by strengthening genetic associations

between cue and preference. Dissasociation of alleles at loci that are physically close

on the chromosome is less likely compared to that between alleles at more distant

loci (due to reduced crossing over), or at loci on different chromosomes. Similarly,

the substitution of large effect alleles will also increase linkage disequilibrium

between cue and preference, even if they are not physically linked, because

preference alleles of larger effect will more often find themselves in the same

genome as alleles for the corresponding cue, compared to preference alleles with

smaller effects. We cannot currently distinguish whether preference QTL result from

single adaptive mutations, or represent multiple functional loci that have built up

during the course of speciation. Nevertheless, the genetic basis of *Heliconius* mate

preferences is remarkably similar to that for differences in the wing pattern cue.

Differences in individual color pattern elements probably do involve multiple,

sequential mutations (which target the same gene(s)), but 'ready-made' alleles of

large phenotypic effect can be brought together in new combinations through

adaptive introgression, a process we have termed 'enhancer shuffling'. This can

generate a multitude of novel wing pattern phenotypes. In this way, mimicry in

Heliconius can be achieved through hybridization and subsequent sharing of key loci

between closely related species [25, 49]. The existence of large effect preference

loci, potentially influencing different aspects of behavior, could similarly facilitate the

origin of novel phenotypes through recombination and introgression, and facilitate

rapid speciation.

Methods

Butterfly collection, rearing and crossing design. All butterfly rearing, genetic

crosses and behavioral experiments were conducted at the Smithsonian Tropical

Research Institute in Panama between August 2007 and August 2009.

We collected wild *Heliconius cydno chioneus* and *Heliconius melpomene rosina* from

Gamboa (9°7.4'N, 79°42.2' W, elevation 60 m) and the nearby Soberania National

Park, Panama. These were used to establish stocks maintained in insectaries in

Gamboa, which were further supplemented with wild individuals throughout the

experimental period. We established interspecific crosses by mating wild caught *H.

melpomene* males to *H. cydno* females from our stock population. In interspecific

crosses between *Heliconius cydno* females and *Heliconius melpomene* males, F1

hybrid females are sterile, restricting us to a backcrossing design. We generated

backcross broods to *H. cydno* and *H. melpomene* by mating F1 males to virgin

females from our stock populations. Brood mothers were kept individually in cages

(approx. 1 or 2 x 2x2m), and provided with ~10% sugar solution, a source of pollen

and Passiflora shoots for oviposition. Eggs were collected daily and caterpillars

raised individually in small pots until 4th or 5th instar, and then either in groups or

individually until pupation. Caterpillars were provided with fresh Passiflora leaves and

shoots daily.

Behavioral assays. We measured male attraction to *H. melpomene* and *H. cydno*

females in standardized choice trials [20]. Males were allowed to mature for at least 5

days after eclosion before testing. Males were introduced into outdoor experimental

cages (1x1x2m) with a virgin female of each species (0 – 10 days matched for age).

Fifteen-minute trials were divided into 1 min intervals, which were scored for

courtship (sustained hovering or chasing) directed towards each female as having

occurred or not occurred. Accordingly, if a male courted the same female twice within

a minute interval, it was recorded only once; if courtship continued into a second

minute, it was recorded twice. Where possible, trials were repeated for each male

(median = 5 trials). This resulted in three related datasets used in subsequent

analyses: i) the proportion of 'courtship minutes' directed towards *H. melpomene* as

opposed to *H. cydno* females (= "the relative amount of time males spent courting red

H. melpomene and white *H. cydno* females" = "relative courtship time"); ii) the

proportion of trials in which courtship was initiated towards *H. melpomene*; and iii) the

proportion of trials in which courtship was initiated towards *H. cydno*. In total we

conducted 1347 behavioral trials, and collected data from 28 *H. cydno*, 16 *H.*

melpomene, 23 F1 hybrid, 29 backcross-to-*melpomene* hybrid and 196 backcross-

to-*cydno* hybrid males (of which 11 performed no courtship behaviors).

Genotyping and linkage map construction. Genotyping and construction of

linkage maps has been described elsewhere [28]. In brief, backcross hybrids and

associated parents were preserved in 20% DMSO and 0.25M EDTA (pH 8.0) and

stored at -20°C. DNA was extracted with Qiagen DNeasy Blood & Tissue Kits

following the manufacture's protocol for animal tissue. Individuals were genotyped

using a RAD-sequencing approach [50] and sequenced by BGI using the Illumina

HiSeq 2500. Sequences were then aligned to version 2 of the *H. melpomene*

genome [51] using Stampy v1.0.23 [52] . Duplicates were removed with Piccard tools

v1.135 (http://broadinstitute.github.io/picard/), and genotype posteriors called using

SAMtools v1.2. Interspecific linkage maps were constructed using Lep-MAP2 [53]

and modules from Lep-MAP3 as described in [28]. To obtain the genotypic data for

QTL mapping, the parental-phased data was obtained using Lep-MAP3

option outputPhasedData=1. This option imputes data based on input genotypes and

the map order. These data were then compared to the subset of markers in which

grandparents could be used to phase the data for each family and chromosome

using custom scripts. Family and chromosome was inverted when required to obtain

matching phases. Finally, the non-informative markers between inferred

recombinations were masked (i.e. set to missing) to account for the fact the exact

recombination position was not known for these regions.

Data analysis. We were able to generate genotype data for 146 of the 196

backcross-to-*cydno* hybrid males for which we recorded behaviors in our choice

trials. The remaining 50 individuals include males from which we were unable to

extract sufficient DNA, were poorly sequenced, or were lost in the insectaries most

often due to ants or other predators. For each backcross individual, we calculated the

probabilities of the two alternative genotypes at every marker and centiMorgan (cM)

position along the chromosomes, conditional on the available marker data, using

R∕qtl package [54]. R∕qtl uses a hidden Markov model to calculate the probabilities of

the true underlying genotypes given the observed multipoint marker data. We then

tested for an association between phenotype and genotype at each position using

generalized linear mixed models (GLMMs) with binomial error structure and logit link

function (implemented with the R package lme4). We first considered the relative

time males courted *H. melpomene* as opposed to *H. cydno* females. For each

position along the genome we modeled the response vector of the number of

'courtship minutes' towards *H. melpomene* vs 'courtship minutes' towards *H. cydno*

with the genotype probability as the independent variable. LOD scores were obtained

by comparing this to a null model in which genotype probability was not included. An

individual level random factor was included in all models to account for over-

dispersion. This approach is analogous to the to the Haley-Knott regression

implemented in R/qtl [54, 55], but more appropriately accounts for the non-normal

structure of our data and for differences in total courtship data recorded for each

individual [56]. Seven individuals were excluded from these analyses for which,

although tested in multiple trials, no courtship towards either female type was

recorded. Using permutation [57], we determined the genome-wide significance

threshold for the association between marker genotype and phenotype (alpha = 0.05,

n = 1000 permutations) as LOD = 2.99. By using our GLMM approach we had more

power to detect QTL than would be permitted by adopting non-parametric methods.

Nevertheless, we repeated all genome-wide QTL analyses using non-parametric

interval mapping in R/qtl, using the 'scanone' and 'model = "np" ' commands. Results

of non-parametric analyses are reported in the supplementary materials (Table S1).

To consider all three QTL identified in our initial genome scans together, we again

modeled the number of 'courtship minutes' towards H. melpomene vs 'courtship

minutes' towards H. cydno but with the genotype at the max LOD score for each QTL

as explanatory variables. The fully saturated GLMM, including all three pairwise

interactions, was simplified in a stepwise manner using likelihood ratio tests, but

always retaining individual id as a random factor. To further test for effects of each

QTL we compared the penalized LOD scores of the full model (including all three

QTL as additive effects) to reduced models in which each QTL was eliminated in

turn. The penalized LOD score is calculated as: $pLODa(\gamma) = LOD(\gamma) - T|\gamma|$, where γ

denotes a model, $|\gamma|$ is the number of QTL in the model and T is a penalty

determined through permutation (i.e. the genome-wide significance threshold = 2.99)

[58] .

To identify QTL associated with courtship initiation towards each species we

again used GLMMs with binomial error structure and logit link function. This was

done in exactly the same way as our initial genome scan looking at relative time

males courted *H. melpomene* as opposed to *H. cydno* females except that the

response variables were i) trials in which courtship was initiated towards *H. cydno*

females vs trials in which courtship towards *cydno* females did not occur, and ii) trials

in which courtship was initiated towards *H. melpomene* females vs trials in which

courtship towards melpomene females did not occur. Here we determined the

genome-wide significance thresholds for the association between marker genotype

and phenotype (alpha = 0.05, n = 1000 permutations) as LOD = 3.08 (*cydno*) and

LOD = 3.02 (*melpomene*). We included all 146 backcross males for which we had

genotype data in this analysis.

Finally, to determine the contribution of each QTL to variation in courtship time

towards *H. cydno* and *H. melpomene* females separately, we considered the total

number of 'courtship minutes' directed to each female type, correcting for the number

of trials. We included all 146 backcross males for which we had genotype data in this

analysis. We square-root transformed courtship minutes/trial and then used the

makeqtl() and fitqtl() functions in R/qtl [54] to guide model simplification . Model

residuals were inspected visually for an approximate normal distribution, and we

tested for homogeneity of variance with Levene's tests (*H. cydno* females: $F_{3,142}$ =

0.25, P > 0.02; *H. melpomene* females: F3,142 = 0.29, P > 0.02). We corrected P

values to account for the 6 tests (i.e. 3 loci x 2 species) [59].

Simulations. We used simulations to estimate potential inflation of measured effect

sizes due to the Beavis effect. We generated 10000 simulated data-sets for each of a

range of 'true' effect sizes for each significant QTL (i.e. on chromosomes 1, 17 and

18), using the R package simr [60]. For each of these we determined the LOD score

and compared it to our genome-wide significance threshold (i.e. LOD = 2.99). This

allowed us to compare i) the entire range of simulated effects (where the mean is

expected to equal the 'true' effect size), with those that would be significant given our

sample size (n = 139) and linkage map (figure S1), and ii) the empirically measured

effects with simulated effects that would be significant (Fig. 3A).

Code and data availability. Sequence data used to make the linkage maps have

previously been submitted to the European Nucleotide Archive

(http://www.ebi.ac.uk/ena) [28], accession number ERP018627.

493   **Table 1. Individual and combined QTLs for differences in relative courtship time.**

Chromosome	Position (cM)	LOD score	ΔpLOD$_a$	2ΔlnL	P
1	4.2 (1.8-9.1)	4.54	-1.04	18.54	<0.001
17	24.4 (0-48.3)	3.50	-1.03	18.50	<0.001
18	0 (0-13.6)	6.83	-3.87	31.60	<0.001
1+17+18	**–**	**14.90**	**(5.93)**	**–**	**–**

Position in cM (1.5 LOD interval); LOD score, log odds ratio; ΔpLOD$_a$, penalized LOD score, change in penalized LOD score compared to the full (best supported) model incorporating all three putative QTLs (in bold); 2ΔlnL & P values compare the full model to reduced models in which individual QTLs were eliminated; *n* = 139.

Figure 1. Divergence in warning pattern cue and corresponding preference in sympatric

Heliconius **butterflies.** A, Wing pattern phenotypes of: top, *Heliconius cydno chioneus* (left), *H.*
melpomene rosina (right) their non-mimetic first generation hybrid (center), and bottom, their sympatric
co-mimics *H. sapho sapho* (left) and *H. erato demophoon* (right). B, Distribution of *H. cydno* (blue) and
H. melpomene (orange). Individuals were collected, and experiments performed in Panama (black
circle), where the two species co-occur in Central and northern South America. **C**, Proportion of
courtships directed towards *H. melpomene* (as opposed to *H. cydno*) females for *H. cydno* (CY), *H.*
melpomene (ME), their F1 and backcross hybrids to *H. cydno* (BC) and *H. melpomene* (BM). Values
in parentheses indicate total number of individuals with behavioral data. Solid colored boxes represent
expected average genome contribution of each generation. Note that a further 11 BC individuals were
tested but performed no courtship behaviors.

Figure 2. QTL analysis of variation in mate preference. A, QTLs for relative time males court *H.*

melpomene (as opposed to *H. cydno*) females on chromosomes 1, 17 and 18 (*n* = 139). Scale on right

axis depicts genome-wide significance (P_{perm}), determined through permutation. Dotted red line

represents log odds ratio (LOD) significance threshold (genome-wide alpha = 0.05, LOD = 2.99).

Dashes indicate position of genetic markers (SNPs) and red arrows indicate the position of the max

LOD score for each QTL (used in B). Horizontal blue lines represent the position of major color pattern

loci, and their phenotypic effects of major color pattern loci. Note that the *K* locus only has limited

phenotypic effects in crosses between *H. cydno chionius* and *H. melpomene rosina*, but is responsible

for the switch from yellow to white color pattern elements between other taxa within the *melpomene-*

cydno clade. B, Proportion of time males court *H. melpomene* (as opposed to *H. cydno*) females for

each of the two genotypes for respective QTLs (homozygous = cyd:cyd, and heterozygous = cyd:mel).

Error bars represent 95% confidence intervals. Lower dashed blue and upper orange bars represent

mean phenotypes measured in *H. cydno* and *H. melpomene,* respectively. Circle size depicts total

number of 'courtship minutes' for each male. Vertical black bars indicate the percentage of the

difference measured in the parental species explained.

Figure 3. QTL effects in consideration of the to the Beavis effect. A, Proportion of 'significant'

simulations that would be smaller than our empirically measured effects, for preference QTL on

chromosome 1 (blue), chromosome 17 (black), and chromomsome 18 (orange). 10000 simulations

were run for effect sizes corresponding to between 5 and 40% of the difference in male preference

behavior of the parental species difference between parental species behavioral. In each case the

distribution of sample effect sizes was determined for those simulations that reached the genome-wide

significance threshold determined through permutation (figure 2). B, Proportion of time males court *H.*

melpomene (as opposed to *H. cydno*) females for each of the two genotypes for red (homozygous =

cyd:cyd) and white (heterozygous = cyd:mel) hybrid males for which we were unable to generate

RADdata (and so which were not included in our initial QTL analysis). Error bars represent 95%

confidence intervals. Lower dashed blue and upper orange bars represent mean phenotypes

measured in *H. cydno* and *H. melpomene,* respectively. Circle size depicts total number of 'courtship

minutes' for each male. Vertical black bars indicate the percentage of the difference measured in the

parental species explained.

References

1. Nosil P Ecological speciation. Oxford University Press, Oxford 2012.

2. Gavrilets S Fitness landscapes and the origin of species. Princeton University Press, Princeton 2004.

3. Smadja CM, Butlin RK A framework for comparing processes of speciation in the presence of gene flow. Mol Ecol. 2011; 20:5123–5140.

4. Servedio MR, Doorn GSV, Kopp M, et al Magic traits in speciation: "magic" but not rare? Trends in Ecology & Evolution. 2011; 26:389–397.

5. Kopp M, Servedio MR, Mendelson TC, et al Mechanisms of Assortative Mating in Speciation with Gene Flow: Connecting Theory and Empirical Research. Am Nat. 2018; in press.

6. Thibert Plante X, Gavrilets S Evolution of mate choice and the so-called magic traits in ecological speciation. Ecol Lett. 2013; 16:1004–1013.

7. Lamichhaney S, Berglund J, Almén MS, et al Evolution of Darwin's finches and their beaks revealed by genome sequencing. Nature. 2015; 518:371–375.

8. Albert AYK, Sawaya S, Vines TH, et al The genetics of adaptive shape shift in stickleback: pleiotropy and effect size. Evolution. 2008; 62:76–85.

9. Chung H, Chung H, Loehlin DW, et al A single gene affects both ecological divergence and mate choice in Drosophila. Science. 2014; 343:1148–1151.

10. Reed RD, Papa R, Martin A, et al optix drives the repeated convergent evolution of butterfly wing pattern mimicry. Science. 2011; 333:1137–1141.

11. Martin A, Papa R, Nadeau NJ, et al Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. Proc Natl Acad Sci USA . 2012; 109:12632–12637.

12. Nadeau NJ, Pardo Diaz C, Whibley A, et al The gene cortex controls mimicry and crypsis in butterflies and moths. Nature. 2016; 534:106–110.

13. Bay RA, Arnegard ME, Conte GL, et al Genetic coupling of female mate choice with polygenic ecological divergence facilitates stickleback speciation. Curr Biol. 2017; 27:3344–3349.e4.

14. Bull V, Beltran M, Jiggins CD, et al Polyphyly and gene flow between non-sibling Heliconius species. BMC Biol. 2006; 4:11.

15. Jiggins CD, Naisbit RE, Coe RL, Mallet J Reproductive isolation caused by color pattern mimicry. Nature. 2001; 411:302–305.

16. Naisbit RE, Jiggins CD, Mallet J Disruptive sexual selection against hybrids contributes to speciation between Heliconius cydno and Heliconius melpomene. Proceedings of the Royal Society B: Biological Sciences. 2001; 268:1849–1854.

17. Maravez J, Salazar CA, Bermingham E, et al Speciation by hybridization in Heliconius butterflies. Nature. 2006; 441:868–871.

18. Merrill R, Wallbank RWR, Bull V, et al Disruptive ecological selection on a mating cue. Proceedings of the Royal Society B: Biological Sciences. 2012; 279:4907–4913.

19. Southcott L, Kronforst MR Female mate choice is a reproductive isolating barrier in Heliconius butterflies. 2017; bioRxiv 233270.

591   20.   Merrill R, Van Schooten B, Scott JA, Jiggins CD Pervasive genetic associations between traits
592   causing reproductive isolation in Heliconius butterflies. Proceedings of the Royal Society B:
593   Biological Sciences. 2011; 278:511–518.

594   21.   Merrill R, Dasmahapatra KK, Davey JW, et al The diversification of Heliconius butterflies: what
595   have we learned in 150 years? J Evol Biol. 2015; 28:1417–1438.

596   22.   Jiggins CD Ecological speciation in mimetic butterflies. BioScience. 2008; 58:541-548

597   23.   Naisbit RE, Jiggins CD, Mallet J Mimicry: developmental genes that contribute to speciation.
598   Evolution & Development. 2003; 5:269–280.

599   24.   Kronforst MR, Kapan DD, McNeely C, et al Linkage of butterfly mate preference and wing color
600   preference cue at the genomic location of wingless. Proceedings of the National Academy of
601   Sciences. 2006; 103:6575–6580.

602   25.   Wallbank RWR, Baxter SW, Pardo Diaz C, et al Evolutionary Novelty in a Butterfly Wing Pattern
603   through Enhancer Shuffling. PLoS Biol. 2016; 14:e1002353.

604   26.   Van Belleghem SM, Rastas P, Papanicolaou A, et al Complex modular architecture around a
605   simple toolkit of wing pattern genes. Nature Ecology & Evolution. 20171:s41559–016–0052.

606   27.   Chamberlain NL, Hill RI, Kapan DD, et al Polymorphic butterfly reveals the missing link in
607   ecological speciation. Science. 2009; 326:847–850.

608   28.   Davey JW, Barker SL, Rastas PM, et al No evidence for maintenance of a sympatric Heliconius
609   species barrier by chromosomal inversions. Evolution Letters. 2017; 1:138–154.

610   29.   Weber JN, Peterson BK, Hoekstra HE Discrete genetic modules are responsible for complex
611   burrow evolution in Peromyscus mice. Nature. 2013; 493:402–405.

612   30.   Diao W, Mousset M, Horsburgh GJ, et al Quantitative trait locus analysis of mating behavior and
613   male sex pheromones in Nasonia Wasps. G3. 2016; 6:1549–1562.

614   31.   Savolainen O, Lascoux M, Merilä J Ecological genomics of local adaptation. Nat Rev Genet. 2013;
615   14:807–820.

616   32.   Felsenstein J Skepticism towards santa rosalia, or why are there so few kinds of animals?
617   Evolution. 1981; 35:124–138.

618   33.   Seimiya M, Gehring WJ The Drosophila homeobox gene optix is capable of inducing ectopic eyes
619   by an eyeless-independent mechanism. Development. 2000; 127:1879–1886.

620   34.   Martin A, McCulloch KJ, Patel NH, et al Multiple recent co-options of optix associated with novel
621   traits in adaptive butterfly wing radiations. Evodevo. 2014; 5:7.

622   35.   Maynard Smith J Sympatric speciation. Am. Nat. 1966; 100:637–650.

623   36.   Shaw KL, Lesnick SC Genomic linkage of male song and female acoustic preference QTL
624   underlying a rapid species radiation. Proc Natl Acad Sci USA. 2009; 106:9737–9742.

625   37.   Wiley C, Ellison CK, Shaw Widespread genetic linkage of mating signals and preferences in the
626   Hawaiian cricket Laupala. Proceedings of the Royal Society B: Biological Sciences. 2012;
627   279:1203–1209.

628   38.   Greenwood AK, Wark AR, Yoshida K, Peichel CL Genetic and neural modularity underlie the
629   evolution of schooling behavior in threespine sticklebacks. Current Biology. 2013; 23:1884-1888.

630   39.   Melo MC, Salazar C, Jiggins CD, Linares M Assortative mating preferences among hybrids offers
631   a route to hybrid speciation. Evolution. 2009; 63:1660–1665.

40. Jiggins CD, Estrada C, Rodrigues A Mimicry and the evolution of premating isolation in Heliconius
melpomene Linnaeus. Journal Evolutionary Biology. 2004; 17:680–691.

41. Merrill R, Gompert Z, Dembeck LM, et al Mate preference across the speciation continuum in a
clade of mimetic butterflies. Evolution. 2011; 65:1489–1500.

42. Walters JR, Stafford C, Hardcastle TJ Evaluating female remating rates in light of spermatophore
degradation in Heliconius butterflies: pupal-mating monandry versus adult-mating polyandry.
Ecological Entomology. 2012; 37:257–268.

43. Rowe L, Arnqvist G Sexual conflict. Princeton University Press, Princeton 2005.

44. Seehausen O, Terai Y, Magalhaes IS, et al Speciation through sensory drive in cichlid fish. Nature.
2008; 455:620–626.

45. Naisbit RE, Jiggins CD, Linares M, et al Hybrid sterility, Haldane's rule and speciation in Heliconius
cydno and H. melpomene. Genetics. 2002; 161:1517–1526.

46. Boggs CL Selection pressures affecting male nutrient investment at mating in heliconiine
butterflies. Evolution. 1981; 35:931–940.

47. Boggs CL, Gilbert LE Male contribution to egg production in butterflies: evidence for transfer of
nutrients at mating. Science. 1979; 206:83–84.

48. Kronforst MR, Gilbert LE Reinforcement of mate preference among hybridizing Heliconius
butterflies. Journal Evolutionary Biology. 2007; 20:278–285.

49. Heliconius Genome Consortium Butterfly genome reveals promiscuous exchange of mimicry
adaptations among species. Nature. 2012; 487:94–98.

50. Baird NA, Etter PD, Atwood TS, et al Rapid SNP discovery and genetic mapping using sequenced
RAD markers. PloS ONE 3:e33754.   Davey JW, Chouteau M, Barker SL, et al Major
improvements to the Heliconius melpomene genome assembly used to confirm 10 chromosome
fusion events in 6 million years of butterfly evolution. G3. 2016; 6:695–708.

51. Davey JW, Chouteau M, Barker SL, et al Major improvements to the Heliconius melpomene
genome assembly used to confirm 10 chromosome fusion events in 6 million years of butterfly
evolution. G3. 2016; 6:695–708.

52. Lunter G, Goodson M Stampy: A statistical algorithm for sensitive and fast mapping of Illumina
sequence reads. Genome Res. 2010; 21:936–939.

53. Rastas P, Paulin L, Hanski I, et al Lep-MAP: fast and accurate linkage map construction for large
SNP datasets. Bioinformatics. 2013; 29:3128–3134.

54. Broman KW, Wu H, Sen Ś, Churchill GA R/qtl: QTL mapping in experimental crosses.
Bioinformatics. 2003; 19:889–890.

55. Haley CS, Knott SA A simple regression method for mapping quantitative trait loci in line crosses
using flanking markers. Heredity. 1992; 69:315–324.

56. Bolker BM, Brooks ME, Clark CJ, et al Generalized linear mixed models: a practical guide for
ecology and evolution. Trends in Ecology & Evolution. 2009; 24:127–135.

57. Churchill GA, Doerge RW Empirical threshold values for quantitative trait mapping. Genetics.
1994; 138:963–971.

58. Broman KW, Sen S A Guide to QTL Mapping with R/qtl. Springer, New York. 2009.

59. Forstmeier W, Wagenmakers E-J, Parker TH Detecting and avoiding likely false-positive
findings - a practical guide. Biol Rev Camb Philos Soc. 2017; 92:1941–1968.

60.   Green P, Evolution CMMIEA, SIMR: an R package for power analysis of generalized linear mixed
675   models by simulation. Methods Ecol Evo. 2016; 7:493-498

Acknowledgements We are grateful to Moisés Abanto, Josephine Dessmann, Janet Scott and Bas

van Schooten for help in the insectaries, to Justin Touchon for advice regarding statistics, and to

Stephen Montgomery, Markus Möst, Ricardo Pereira, Alexander Hausmann and Vera Warmuth for

comments on the manuscript. We thank the Smithsonian Tropical Research Institute for support and

the Ministerio del Ambiente for permission to collect butterflies in Panama. We are also grateful to

Edinburgh Genomics for sequencing support. RMM is currently funded by a DFG Emmy Noether

Fellowship, and during this work he was also funded by a Junior Research Fellowship at King's

College, Cambridge and by the ERC grant 339873 Speciation Genetics awarded to CJ. JD was

funded by a Herchel Smith Postdoctoral Fellowship.

Supporting Information

Figure S1. Simulations suggest QTL effect sizes are not greatly overestimated. For each simulated effect size, the distribution of all simulated effects (blue) and those which would be significant in our analysis (*i.e.* lod ≥ 2.99) (orange) are shown. In each case, 'recorded' refers to the empirically measure effect size.

Figure S2. QTL analysis of courtship initiation. QTL associated with the proportion of trials males

initiated courtship towards *H. cydno* females (blue) and *H. melpomene* females (orange). Dotted blue

and orange lines represents log odds ratio (LOD) significance threshold (genome-wide alpha = 0.05)

for *cydno* (LOD = 3.08) and *melpomene* (LOD = 3.02) females, respectively. Gray boxes indicate 1.5

LOD interval for QTL associated with proportion of time males court *H. melpomene* (as opposed to *H.*

cydno) females on chromsomes 1, 17 and 18 (see Fig. 1a). Horizontal bars represent 1.5 LOD interval

for significant QTL associated with the proportion of trials males initiated courtship towards *H. cydno*

females (chr. 1) and *H. melpomene* females (chr. 18).

Figure S3. QTL analysis of variation of mate preference for individuals with alternative

genotypes at LG18@0cM. QTL associated with the proportion of time males court *H. melpomene* (as

opposed to *H. cydno*) females on chromosomes 17 for individuals homozygous (i.e. white, cyd:cyd =

blue line) and heterozygous (i.e. red, cyd:mel = orange line) at LG18@0cM. Dashed line represents

log odds ratio (LOD) significance threshold (genome-wide alpha = 0.05, LOD =) for heterozygous

(i.e.red, cyd:mel) individuals. Horizontal black bars represent 1.5 LOD interval of significant the QTL.

Table S1. Summary of genome-wide QTL analyses using binomial GLMM methods (reported in

main text) and non-parametric methods implemented in R/qtl.

a) Courtship (*n* = 139)

	Binomial GLMM			Non-parametric		
Chromosome	Position (cM)	LOD	*P*	Position (cM)	LOD	*P*
1	4.23	4.54	<0.001	4.23	3.6	~0.009
17	24.47	3.5	~0.013	24.47	2.8935341	~0.049
18	0	6.83	<0.001	0	5.3432172	<0.001

731 b) Courtship initiation (proportion of trial in which courtship occurred) (*n* = 146)

		Binomial GLMM			Non-parametric		
Chromosome	Female	Position (cM)	LOD	*P*	Position (cM)	LOD	*P*
1	CP	4.23	3.44	~0.029	4.23	3.86	~0.005
	MP	4.23	1.66	~0.682	3.02	1.30	~0.871
18	CP	0	2.06	~0.371	0	2.19	~0.243
	MP	6.65	3.10	~0.049	6.65	2.98	~0.038

733
734 CP = *H. cydno chionius*; MP = *H. melpomene rosina*; *P* values are determined through permutation as
735 described in the methods. QTL are reported only for chromosomes where the genome-wide
736 significance threshold was achieved. For comparison, for (b) courtship initiation, non-significant peaks
737 are reported for attraction towards MP and CP females for chromosomes 17 and 18, respectively.
738
739