Contrasting patterns of coding and flanking region evolution in mammalian keratin associated protein-1 genes

Huitong Zhou[1,2,*,†], Tina Visnovska[3,†], Hua Gong[1,2], Sebastian Schmeier[3], Jon Hickford[1,2], and Austen R.D. Ganley[4,*]

1   State Key Laboratory of Sheep Genetic Improvement and Heathy Production, Shihezi 832000, China

2   Faculty of Agricultural and Life Sciences, Lincoln University, Lincoln 7647, New Zealand

3   Institute of Natural and Mathematical Sciences, Massey University Auckland, Auckland 0632, New Zealand

4   School of Biological Sciences, University of Auckland, Auckland 1142, New Zealand

†   Huitong Zhou and Tina Visnovska should be considered joint first author

Author contributions:

ARDG and HZ conceived of the study. HZ, HG, and JH collected the data. TV, HZ, and ARDG

performed analyses. HZ, TV, HG, SS, JH, and ARDG interpreted the data and wrote the

manuscript.

To whom correspondence should be addressed:

Austen Ganley: School of Biological Sciences, University of Auckland, 3A Symonds St,

Building 110N, Auckland 1142, New Zealand; +64 9 923 2906; a.ganley@auckland.ac.nz

Huitong Zhou: Faculty of Agriculture and Life Sciences, Lincoln University,

34   Cnr Springs Road & Ellesmere Junction Road, Lincoln 7647, New Zealand; +64 3 423 0684;

35   zhouh@lincoln.ac.nz

37   **Acknowledgements:**

38   This work was supported by a Marsden Fund award (14-MAU-053) to ARDG, an AGMARDT

39   Postdoctoral Fellowship to HG, and a Vernon Willey Trust Fellowship to HZ.

Abstract

DNA repeats are common in eukaryotic genomes, and recombination between copies can occur. This recombination can result in concerted evolution, where within-genome repeats are more similar to each other than to orthologous repeats in related species. We investigated the tandemly-repeated keratin-associated protein (KAP) gene family, *KRTAP1*, which encodes proteins that are important components of hair and wool in mammals. Comparison of *KRTAP1* gene repeats across the mammalian phylogeny shows strongly contrasting evolutionary patterns between the coding regions, that have a concerted evolution pattern, and the flanking regions, that have a normal, radiating pattern of evolution. This dichotomy transitions abruptly at the start and stop codons, and is not the result of purifying selection, codon adaptation, or reverse transcription of *KRTAP1-n* mRNA. Instead, our results suggest that short-tract gene conversion events coupled with selection for these events in the coding region drives the contrasting *KRTAP1* repeat evolutionary patterns. Our work shows the power that repeat recombination has to complement selection and shape the evolution of repetitive genes, and this interplay may be a more common mechanism than currently appreciated for achieving adaptive outcomes in eukaryotic multi-gene families. Thus, our work argues for greater emphasis on exploring the evolution of these families.

Introduction

Repetitive DNA is widespread in most eukaryote genomes (Britten and Kohne 1968; Richard, et al. 2008; Lopez-Flores and Garrido-Ramos 2012). There are two basic repeat DNA types: tandem repeats that are typically arranged in head-to-tail arrays; and dispersed repeats, and these can occur in either coding or non-coding DNA. Repeats are thought to arise from recombination-based duplication/amplification events (Stephan 1989). Sequence identity between duplicates will then decay through the diversifying force of mutation, unless counteracting processes operate (Brown, et al. 1972; Dover 1982). The balance between duplication, diversification, selection, and counteracting forces thus dictate the evolutionary dynamics of repeats. Two main paradigms have been proposed to account for the long-term maintenance of repeat identity: concerted evolution and birth-and-death evolution. Concerted evolution describes a pattern of evolution where the repeats within a genome show greater sequence identity to each other than to orthologous repeats in related genomes (Elder and Turner 1995). The pattern of concerted evolution is proposed to result from recombination-based processes, such as gene conversion and unequal cross-over events, that replace the DNA sequence from one repeat with that from another repeat (Liao 1999). In so doing, these recombination processes maintain sequence identity between repeat copies in the face of mutation, and thus homogenize the repeats (Dover 1982). 'Birth-and-death' evolution involves purifying selection maintaining sequence identity between repeats that are generated by occasional duplication events (i.e. birth), as well as death, which results from repeat loss or pseudogenization (Nei, et al. 1997; Nei, et al. 2000). While there has been debate as to which of these processes best describes the evolutionary dynamics of repetitive DNA (Nei and Rooney 2005; Rooney and Ward 2005; Eirin-Lopez, et al. 2012), a basic characterization of the evolutionary dynamics of most repeat families is lacking.

The keratin-associated proteins (KAPs) are a diverse group of proteins, and are rich in either

sulphur, or glycine and tyrosine. They are important structural components of hair and wool

fibres, and form a matrix that cross-links the keratin intermediate filaments. The genes encoding

the KAPs are called *KRTAP*s (Gong, et al. 2012), and can be classified into 27 families, with

each family comprising 1-12 members that are usually tandemly arranged (Rogers and

Schweizer 2005; Rogers, et al. 2006; Gong, et al. 2016). The *KRTAP*s are single exon (intron-

less) genes, with small coding sequences (less than 1 kb) (Rogers and Schweizer 2005), and they

have low numbers of pseudogenes. For example, in humans the pseudogene:gene *KRTAP* ratio is

approximately 1:5 (Gong, et al. 2016), while across all human genes the ratio is close to 1:1

(Torrents, et al. 2003; Stein 2004). In addition, the *KRTAP*s show high levels of population

variation, with all known *KRTAP* genes being polymorphic in sheep (Gong, Zhou, McKenzie, et

al. 2010; Gong, et al. 2016; Zhou, et al. 2016), where they are well studied because of their roles

in determining wool phenotypes (Zhou, et al. 2015; Li, Zhou, Gong, Zhao, Hu, et al. 2017; Li,

Zhou, Gong, Zhao, Wang, Liu, et al. 2017; Li, Zhou, Gong, Zhao, Wang, Luo, et al. 2017; Tao,

Zhou, Gong, et al. 2017; Tao, Zhou, Yang, et al. 2017). Despite this variation, it has been

reported that at least some *KRTAP* genes show a pattern of concerted evolution between the

paraglogous gene copies (Rogers, et al. 1994; Wu, et al. 2008; Khan, et al. 2014).

The KAP1 proteins form the best characterised KAP family, and they show a high degree of

sequence heterogeneity compared to other KAP families. These KAP1 proteins appear to be

restricted in expression to the middle to upper cortex region of the hair and wool follicle, and are

absent in the cuticle (Powell and Rogers 1997; Shimomura, et al. 2002). Their precise role in hair

and wool function, has yet to be determined. The genes encoding the KAP1 proteins (*KRTAP1-*

n) have been characterized in a number of mammalian species, where they are usually arranged

as four tandem copies (**Figure 1**) (Khan, et al. 2014). The coding regions of the *KRTAP1-n* genes

vary in length within species, predominantly as a consequence of variation in the number of

imperfect tandem decapeptide repeat units (Gong, et al. 2016) (**Figure 1**).

Here we analyse the *KRTAP1* genes from a number of mammalian species, including four

species for which the *KRTAP1-n* loci have not been described. Together with the existing

KRTAP1-n sequences, we reveal that the *KRTAP1-n* coding regions display a pattern of

concerted evolution. In stark contrast to the coding region though, we find that the repeat

flanking regions display no evidence of concerted evolution, and instead appear to be evolving

by normal vertical or radiating evolution. Surprisingly, we find that this pattern of coding region

restricted concerted evolution is not the result of purifying selection, nor does it result from

codon adaptation or reverse transcription/reintegration of *KRTAP1-n* mRNA sequences. Instead,

the results are best explained by a combination of on-going short-tract gene conversion events

between the *KRTAP1-n* copies, and negative selection. We argue that these gene conversion

events act as an unusual mechanism of purifying selection to prevent excessive intra-genomic

divergence between the four gene copies, while also allowing inter-species diversity. This

unusual mode of evolution may apply to other multicopy genes that encode products subject to

diversifying selection.

Materials and Methods

Sequence Resources and Gene Identification: All genome sequences were sourced from the

NCBI GenBank. Previously identified *KRTAP1-n* sequences (Itenge-Mweza, et al. 2007; Wu, et

al. 2008; Gong, Zhou and Hickford 2010; Gong, et al. 2011) were used to search the genomes of

cattle, horses, rabbits and African elephants using BLAST with default parameters, and the genes

retrieved were identified by sequence identity within both the coding and flanking regions

(**Table S1**).

Sequence Alignments: *KRTAP1* nucleotide sequences (**Table S1**) for all four paralogs from the

ten species (sheep, cattle, dog, elephant, horse, human, macaque, mouse, rat and rabbit) were

separated into 5' flanking regions, coding sequences, and 3' flanking regions. The multiple

sequence alignment tool *mafft* (v7.123b) (Katoh and Standley 2013) was used to separately align

the 5' and 3' flanking regions as nucleotide sequences, using the arguments '--nuc --localpair --

maxiterate 1000'. To align the coding sequences at the predicted amino acid level, *mafft* with the

arguments '--amino --localpair --maxiterate 1000' was run.

The coding sequence alignment was subsequently reverse translated using *revTrans* (v1.4)

(Wernersson and Pedersen 2003) with two input files: the sequences of all the coding regions,

and the amino acid sequence alignments. The sequences in the two files were paired by name

using the '-match name' parameter, and default values were used for all other parameters. A

number of regions align poorly and have many indels, therefore we used the longest continuous

coding sequence block (198 nucleotides; covers on average around 40% of the coding region)

where none of the 40 sequences had indels. For the flanking region alignments, we used *Gblocks*

(v0.91b) (Talavera and Castresana 2007) to select blocks that cover approximately 40% of the

flanking regions having the best alignment. We also used *Gblocks* with less stringent criteria to

create multiple sequence alignments of the coding and flanking regions that included more

poorly aligning regions.

Phylogenetic Trees: *PhyML* (v3.1) (Guindon, et al. 2010) was used to construct phylogenies

based on the coding and flanking region sequences. The number of resampled bootstrap data sets

was set to 1000 (parameter '-b 1000'), and the additional arguments '-q -s BEST -o tlr' were

employed. The Bioconductor package *ggtree* (v1.9.4) (Yu, et al. 2017) was used to plot the

phylogenies.

Codon Adaptation Index: The CAIcal server (http://genomes.urv.es/CAIcal(Puigbo, et al.

2008) was used to calculate CAI values for the *KRTAP1*s, as well as expected CAI values from

permutated sequences using default parameters and published codon usage data (Nakamura, et

al. 2000).

Motifs in the Coding Sequences: We used MEME motif finder (v4.12.0) (Bailey, et al. 2006) to

explore repetitive elements in the coding sequences. The repetitive structure of the coding

regions reported in the Results was obtained with parameters '-dna -oc . -nostatus -time 18000 -

maxsize 60000 -mod anr -nmotifs 6 -minw 6 -maxw 30 -minsites 20 -maxsites 600 –revcomp'

and all the other parameters set to the default values.

KRTAP1-n **Polymorphism in Sheep:** Intra-specific variation was assessed using three

sequences for *KRTAP1-1* (Itenge-Mweza, et al. 2007), eleven sequences for *KRTAP1-2* (Gong, et

al. 2011; Gong, et al. 2015), nine sequences for *KRTAP1-3* (Itenge-Mweza, et al. 2007), and nine

sequences for *KRTAP1-4* (Gong, Zhou and Hickford 2010). These were aligned using

DNAMAN (v5.2.10; Lynnon BioSoft, Canada) with default parameters, and polymorphic sites

were identified manually.

Data Availability: Sequence data are available at GenBank and the accession numbers and

positions are listed in the **Materials and Methods** (sheep polymorphism data) and Table S1

(*KRTAP1* sequences).

Results

Mammalian *KRTAP1-n* repeats show a concerted evolution pattern in the coding but not

the flanking regions

To better understand the genetic architecture of the mammalian *KRTAP1* cluster, we selected the

KRTAP1 genomic region from key members of the mammalian phylogeny for analysis. The

Basic Local Alignment Search Tool (BLAST) was used to search GenBank with known

KRTAP1-n sequences to identify and retrieve the *KRTAP1* clusters from the genomes of four

species (cattle, horses, rabbits and African elephants) for whom *KRTAP1-n* sequence information

has not been reported (**Figure S1**). We then combined these with previously-identified *KRTAP1-*

n sequences from other mammalian species to obtain sampling across the mammalian phylogeny

(**Figure 2**).

Previously, the *KRTAP1* genes of sheep were shown to contain a variable number of occurrences

of a QTSCCQPXXX decapeptide tandem repeat in the N-terminal region of the protein (Rogers,

et al. 1994; Gong, et al. 2011; Gong, et al. 2016). We used a motif finding tool (MEME; (Bailey,

et al. 2006) to search for repetitive motifs in the coding regions of all the mammalian *KRTAP1-n*

sequences. This revealed that the decapeptide repeat is present at the N-terminus in all

mammalian *KRTAP1-n* genes we obtained (**Figure S2)**, albeit with less amino acid conservation

than that observed in sheep. MEME also identified nucleotide level tandem copies of this repeat

at the C-terminus of the protein. Furthermore, both the N- and C-terminal repeats vary in copy

number, within and between genomes. This copy number variation is responsible for much of the

length variation between *KRTAP1-n* sequences.

To determine the genetic relationships between of the mammalian *KRTAP1-n* genes, we

generated a *KRTAP1* phylogenetic tree from an alignment of our mammalian *KRTAP1-n* coding

region sequences. This revealed that, in most cases, the *KRTAP1* genes are more related to each

other within a species than to their orthologs in other species, thus exhibiting a concerted

evolution pattern. This manifests as clades that group by species, rather than by repeat, in the

phylogenetic tree (**Figure 3**). This concerted evolution pattern breaks down between the most

closely-related species pairs (cattle/sheep, rat/mouse, human/macaque), presumably because the

signal is confounded by these species having more recent shared ancestry. Nevertheless, for most

species there is a clear pattern of concerted evolution.

For concertedly evolving tandem repeat sequences such as the ribosomal RNA gene repeats,

homogenization occurs for the complete repeat unit, including the non-coding regions (Ganley

and Kobayashi 2007). To test whether the *KRTAP1* clusters display a 'whole-unit' pattern of

concerted evolution, we generated *KRTAP1* phylogenetic trees from multiple alignments of the

5' and 3' flanking sequences of the mammalian *KRTAP1* genes. Surprisingly, the phylogenies

derived from these flanking sequences did not show any pattern of concerted evolution, and in

contrast to the coding region phylogeny, the clades in these phylogenetic trees were group by

KRTAP1 repeat number, not by species (**Figure 3**). We note that bootstrap support is not strong

for all the clades in these phylogenetic trees, but the contrast between the coding region

concerted versus flanking region radiating evolutionary patterns is unmistakable. Furthermore,

the topology within many of the *KRTAP1* flanking region clades is consistent with the reported

mammalian phylogeny (refer to **Figures 2** and **3**). These phylogenies were generated from

multiple sequence alignments that encompass the regions that align well, but phylogenies

derived from sequence alignments that include poorly aligned regions give qualitatively similar

results (**Figure S3**). Overall, in stark contrast to the coding region, the flanking regions show a

phylogenetic pattern expected for normal radiating evolution, and exhibit no evidence of

concerted evolution.

What is responsible for the different evolutionary patterns of the *KRTAP1* coding and

flanking regions?

The difference in evolutionary pattern between the coding and flanking regions is striking, hence

we sought to identify the mechanism(s) responsible.

Purifying selection: Previous studies have shown that multi-gene loci undergoing birth-and-

death evolution can show high levels of identity within the coding region due to strong purifying

selection (Nei, et al. 2000; Piontkivska, et al. 2002). It is possible that purifying selection

maintains sequence identity between *KRTAP1-n* copies within a species, whilst diversifying

selection results in differences between species. If so, we would predict that while the non-

synonymous sites would show a concerted evolution pattern, the synonymous sites would instead

show a normal radiating pattern of evolution (resembling the flanking regions).

To investigate this, we looked at the pattern of evolution of the synonymous sites in the coding

sequences compared to the non-synonymous sites. The number of KAP1 amino acid changes

present within and between species makes it difficult to consistently call sites as synonymous or

non-synonymous, so third codon positions were used as a proxy for synonymous sites, and first

and second codon positions were used as a proxy for non-synonymous sites. We generated

phylogenetic trees from multiple sequence alignments of the first-second (which we refer to as

"non-synonymous"), and third (which we refer to as "synonymous") codon sites of the *KRTAP1-*

n coding regions to test for different evolutionary patterns. Surprisingly, while the non-

synonymous sites displayed a pattern of concerted evolution as was expected (**Figure 4A**), the

synonymous sites also revealed the same pattern of concerted evolution (**Figure 4B**). The

concerted evolution pattern for the synonymous sites seems to be stronger than that of the non-

synonymous sites, as they separate sheep and cattle into separate clades, and also resolve dog,

elephant, and rat/mouse into separate clades (**Figure 4**).

Codon adaptation: We considered whether this pattern of concerted evolution amongst the

synonymous sites might result from codon adaptation (Lin, et al. 2006), as a result of

synonymous mutations being selected to follow changes in the favoured codons between species.

The *KRTAP1-n* genes display strong evidence for codon adaptation (the degree to which the

favoured codons for that species are used in a gene). For example, the human *KRTAP1-n* genes

collectively show a codon adaptation index (CAI) of 0.91 (out of a maximum of 1), higher than

the CAI of randomly permuted human *KRTAP1* sequences (CAI=0.78). Using the *KRTAP1*

coding sequence alignment used for the phylogenies presented in **Figure 3**, we identified nine

synonymous differences between human and mouse that exhibit a concerted evolution pattern

(similarity within species versus difference between species). If codon adaptation can explain

this pattern, these synonymous mutations should change in a manner consistent with a change in

codon usage preference for that amino acid. Five of these mutations show the pattern expected,

given the change in codon usage between human and mouse (synonymous change creates the

more favoured codon in the species it is found in). However, four of these mutations show the

opposite pattern, and most of the codon usage preference changes between human and mouse are

small (**Table S2**). These results provide no evidence for adaptation to different codon usage

preferences driving the pattern of *KRTAP1* concerted evolution.

Reverse transcription of *KRTAP1* mRNA: Another potential explanation for the incongruence

in evolutionary pattern between the *KRTAP1* coding and flanking regions is reverse transcription

of *KRTAP1-n* mRNAs, followed by homologous recombination-mediated replacement of a

genomic *KRTAP1-n* with the reverse transcribed copy (Coulombe-Huntington and Majewski

2007). This is feasible given that *KRTAP1-n* are single-exon genes. If reverse transcription

events occur, the 5' and particularly 3' flanking regions should show a concerted evolution

pattern that is similar to the coding region. Inspection of the 5' and 3' flanking regions revealed

that sequence similarity between *KRTAP1-n* sequences within a genome tends to decay

immediately upstream of the ATG codon and downstream of the stop codon (**Figure 5**). This

suggests that reverse transcription/integration of *KRTAP1-n* mRNA is unlikely to explain the

pattern of *KRTAP1* concerted evolution, as the transcribed flanking regions of the gene would be

expected to 'hitch-hike' with the coding regions through such a mechanism.

We also considered whether the *KRTAP1-n* sequences might have arisen through a pure birth-

and-death process by independent gene duplication events. However, we think this is improbable

as it would require the same number of duplications to occur in at least seven of the species, and,

independently, that each of these duplications would not involve any flanking sequence

(including promoter and terminator sequences) and have inserted into the same site in each

species.

Gene conversion: Finally, we considered whether gene conversion could explain the pattern of

KRTAP1 repeat evolution. Gene conversion events within a genome that convert a section of one

repeat to the sequence of another can create homogeneity (Chen, et al. 2007), and the degree of

homogeneity depends on the relative rates of gene conversion and mutation (Teshima and Innan

2004; Harpak, et al. 2017). Our results imply that if gene conversion does occur, it is somehow

restricted to the coding region. This pattern could occur if there is selective pressure to maintain

a degree of intra-genome homogeneity between the repeat copies. If so, under the assumption

that gene conversion occurs in both the coding and flanking regions, those events occurring in

the flanking region will not have a selective advantage, while those occurring in the coding

region will. Therefore, the probability of gene conversion events becoming fixed in the

population will be greater for events that involve the coding region. There is considerable intra-

genomic variation between *KRTAP1* repeats (**Figure 3**), but this incomplete level of

homogenization can be explained by relatively infrequent gene conversion events and/or relative

infrequent fixation of these events. Therefore, the sequence features of the *KRTAP1* repeats that

we document here can all be accounted for by gene conversion coupled with selection.

Evidence for gene conversion events in the *KRTAP1-n* repeats

Inspection of the *KRTAP1* coding region multiple sequence alignment provides evidence for

tracts of gene conversion. Specifically, sites where there are mutations that are shared between

copies within a species, but that differ between species, are frequently clustered together rather

than scattered throughout the gene (**Figure 6**). Such patches of homogeneity are expected if there

has been occasional, short-tract gene conversion events. The patches we observe are small, but

are within the expected range for mammalian gene conversion events (Chen, et al. 2007). In

addition, we collected population polymorphism data for *KRTAP1-n* sequences in sheep, as

comprehensive sequence variation data are scarce in other species. For many of the sites that are

polymorphic, the polymorphism is shared across some, or all, of the *KRTAP1-n* sequences

(**Figure 7**). While we cannot rule out independent mutation events in each *KRTAP1* copy, we

think that gene conversion is a more parsimonious explanation for this observation, particularly

for the polymorphisms at synonymous sites. Gene conversion has also previously been suggested

as an explanation for the pattern of polymorphism in the ovine *KRTAP1* genes (Rogers, et al.

1994). Collectively, our results suggest that the unusual evolutionary pattern of the *KRTAP1*

repeats, where the coding region evolutionary dynamics are uncoupled from those of the flanking

region, is the result of occasional short-tract gene conversion events that are selected for in the

coding region but not the flanking regions, and that drive partial homogenization.

Discussion

Here we have shown that *KRTAP1-n* genes are conserved as a block of four tandem repeats in

mammalian species, and this suggests they derive from a relatively ancient gene-amplification

event or events that probably pre-date mammalian speciation. The four tandem copies display a

strong pattern of concerted evolution in the coding regions, yet the regions flanking show a

normal radiating pattern of evolution. We suggest that this dichotomous pattern of evolution is

not the result of purifying selection acting to retard changes to the amino acid sequence, but

instead results from short gene conversion tracts that periodically homogenize sequences

between the four *KRTAP1* genes within a genome.

The role of gene conversion is supported by two key pieces of evidence: 1) unique amino acid

tracts that are shared by KAP1 copies within a species, but are unique to that species/group of

related species; and 2) the possession of shared nucleotide variants between *KRTAP1* gene

copies in sheep populations. These results extend previous reports of homogenization via

ongoing short-tract gene conversion events in other protein coding genes (Noonan, et al. 2004;

Lamping, et al. 2017).

We propose that gene conversion is being utilized as an unusual form of purifying selection that

prevents accumulation of too much divergence between *KRTAP1* gene copies. We speculate that

homogeneity of the *KRTAP1* coding sequences is beneficial as it enables the production of more

homogenous components of the hair and wool fibre matrix, and thus potentially facilitates better

associations with the keratin intermediate filaments. We cannot, however, rule out the possibility

that individual *KRTAP1* repeats might have functional differences, the signal of which is

overwhelmed by the concerted evolution signal from the majority of the gene. However, we note

that, particularly in dogs, some of the *KRTAP1-n* genes are very similar in sequence. Therefore,

we favour the explanation that *KRTAP1* concerted evolution results from ongoing, stochastic

gene conversion events coupled with selection within the coding region against inter-repeat

heterogeneity.

Purifying selection is evident in the *KRTAP1-n* coding regions, as the rate of synonymous

change is about twice that of the non-synonymous rate (**Figure 4**). While this may seem to

contradict the similarity in the synonymous and non-synonymous concerted evolution tree

topologies, it can be simply explained by purifying selection acting on residues that are

conserved between species, and thus not contributing to the synapomorphies that influence the

tree topologies. Any gene conversion events that homogenize unfavourable amino acids will be

selected against, thereby preventing deleterious mutations from spreading between copies.

However, this same process also allows tolerable and advantageous amino acid changes to sweep

through the copies (Dover 1982). The *KRTAP1-n* sequences from closely related species (i.e.

human and macaque, rat and mouse, sheep and cattle) were not separated into different clades for

most of the phylogenetic trees we generated (**Figures 3 and 4**). This suggests that the rate of

homogenization is relatively slow, and insufficient to drive substantial homogeneity over the

evolutionary time frames separating these species pairs. In this context, the shared

polymorphisms that we observe in sheep (that are evidence for gene conversion events) are likely

intermediate stages in the accumulation of homogenized *KRTAP1-n* sequences.

The sharp border between a concerted evolution pattern in the coding region and a radiating

evolution pattern in the immediate flanking regions is striking. This can partially be explained by

the selection for gene conversion events within the coding region, as we have proposed.

However, it is intriguing to speculate that this may also be a consequence of differential

expression between the *KRTAP1* genes that is mediated by copy-specific differences in the

regulatory regions. Although not direct, some evidence for differential regulation of *KRTAP1-n*

gene expression was found in two transcriptome studies looking for differentially expressed

genes (Fan, et al. 2013; Chang, et al. 2014). If the *KRTAP1-n* genes do have functionally distinct

roles, gene conversion events in the *KRTAP1* regulatory regions that perturb their differential

regulation may be maladaptive and therefore selected against. Thus, selective pressure for coding

region homogeneity versus regulatory region diversity, coupled with ongoing gene conversion,

may be a powerful way to achieve the dichotomy in evolutionary patterns we observe. Clearly, a

better understanding of the transcriptional regulation of the *KRTAP1* genes is required to address

this hypothesis.

Gene conversion is frequently viewed through the lens of impeding sub-functionalization of gene

duplicates. This view is consistent with the well characterized case of the opsin gene duplicates

in primates, where there is a much stronger signal of gene conversion/concerted evolution in the

introns, than in the exons (Shyue, et al. 1994; Hiwatashi, et al. 2011). The interpretation is that

selection has largely rejected gene conversion events that include the coding (exon) regions,

whilst allowing those occurring in the non-coding (intron) regions to spread in the population

(Shyue, et al. 1994). This is the opposite of what we observe, and illustrates how gene

conversion and selection can intersect to produce a constellation of evolutionary patterns:

homogenization of the non-coding but not the coding regions in the opsin paralogs (Shyue, et al.

1994); homogenization of the coding but not the non-coding regions in the *KRTAP1* genes (this

study); and homogenisation of both coding and non-coding regions equally in the ribosomal

RNA gene repeats (Ganley and Kobayashi 2007).

The extent to which gene conversion acts to homogenize gene duplicates remains controversial

(Gao and Innan 2004; Casola, et al. 2012; Harpak, et al. 2017). Furthermore, even in examples

where recurrent gene conversion events can be detected, they are often not sufficient to produce

a strong concerted evolution pattern (Petronella and Drouin 2011, 2014). There are two potential

explanations for why such a strong pattern of concerted evolution is observed in the case the

KRTAP1 genes, despite the relatively high levels of divergence between copies. First, unlike

many of the examples that have aroused controversy (Gao and Innan 2004; Casola, et al. 2012;

392 Harpak, et al. 2017), the *KRTAP1-n* repeats are tandemly-arranged. Proximity effects as a

393 consequence of tandem arrangement may increase the chances of unequal alignment of the

394 repeats during DNA repair-based homologous recombination compared to dispersed repeats, and

395 thus may increase the chances of inter-repeat gene conversion events. However, this does not

396 explain examples where tandemly repeated paralogs do not show a strong concerted evolution

397 pattern (Nei, et al. 2000; Perina, et al. 2011). A second explanation relates to the imperfect

398 decapeptide tandem repeat motif found in the coding region. Variation in the copy number of

399 decapeptide repeats between *KRTAP1* genes is possibly the result of unequal recombination

400 (Liao and Weiner 1995; Ganley and Scott 1998; Morrill, et al. 2016). If so, the *KRTAP1* genes

401 may harbour a recombination hotspot that drives both decapeptide repeat copy number variation

402 and gene conversion at higher than average levels.

403 Repeats are ubiquitous denizens of eukaryote genomes, where they exist in different forms

404 (coding, non-coding) and organizations (tandem, dispersed). Our results add to the growing list

405 of examples that illustrate how different molecular and evolutionary processes can impinge on

406 repeats to structure their sequences and create distinctive patterns of evolution (Shyue, et al.

407 1994; Noonan, et al. 2004; Ganley and Kobayashi 2007; Storz, et al. 2007; Hiwatashi, et al.

408 2011; Lamping, et al. 2017). However, it is unclear how widespread these sorts of evolutionary

409 dynamics are for eukaryotic gene repeats, largely because the patterns of evolution have not been

410 investigated for the vast majority of multi-gene families. The increasing availability of high

411 quality genome sequences for a wide range of eukaryotes puts us in an excellent position to

412 determine, on a much more systematic and wide-ranging basis, the patterns of repeat sequence

413 dynamics and evolution. This will, in turn, make it clear whether the impact of recombination on

the *KRTAP1*s is unusual, or highlights a common mechanism to finely scale patterns of

homogeneity and divergence between repeat copies over time.

References

Bailey TL, Williams N, Misleh C, Li WW. 2006. MEME: discovering and analyzing DNA and

protein sequence motifs. Nucleic Acids Res. 34:W369-373.

Britten RJ, Kohne DE. 1968. Repeated sequences in DNA. Science 161:529-540.

Brown DD, Wensink PC, Jordan E. 1972. A comparison of the ribosomal DNA's of *Xenopus*

laevis and *Xenopus mulleri*: the evolution of tandem genes. J. Mol. Biol. 63:57-73.

Casola C, Conant GC, Hahn MW. 2012. Very low rate of gene conversion in the yeast genome.

Mol. Biol. Evol. 29:3817-3826.

Chang TH, Huang HD, Ong WK, Fu YJ, Lee OK, Chien S, Ho JH. 2014. The effects of actin

cytoskeleton perturbation on keratin intermediate filament formation in mesenchymal

stem/stromal cells. Biomaterials 35:3934-3944.

Chen JM, Cooper DN, Chuzhanova N, Ferec C, Patrinos GP. 2007. Gene conversion:

mechanisms, evolution and human disease. Nature Reviews Genetics 8:762-775.

Coulombe-Huntington J, Majewski J. 2007. Characterization of intron loss events in mammals.

Genome Res. 17:23-32.

Dover GA. 1982. Molecular drive: a cohesive mode of species evolution. Nature 299:111-117.

Eirin-Lopez JM, Rebordinos L, Rooney AP, Rozas J. 2012. The birth-and-death evolution of

multigene families revisited. Genome Dynamics 7:170-196.

Elder JF, Jr., Turner BJ. 1995. Concerted evolution of repetitive DNA sequences in eukaryotes.

Q. Rev. Biol. 70:297-320.

Fan R, Xie J, Bai J, Wang H, Tian X, Bai R, Jia X, Yang L, Song Y, Herrid M, et al. 2013. Skin

transcriptome profiles associated with coat color in sheep. BMC Genomics 14:389.

Ganley ARD, Kobayashi T. 2007. Highly efficient concerted evolution in the ribosomal DNA

repeats: total rDNA repeat variation revealed by whole-genome shotgun sequence data. Genome

Res. 17:184-191.

Ganley ARD, Scott B. 1998. Extraordinary ribosomal spacer length heterogeneity in a

Neotyphodium endophyte hybrid: implications for concerted evolution. Genetics 150:1625-1637.

Gao L-Z, Innan H. 2004. Very low gene duplication rate in the yeast genome. Science 306:1367-

1370.

Gong H, Zhou H, Forrest RHJ, Li S, Wang J, Dyer JM, Luo Y, Hickford JGH. 2016. Wool

keratin-associated protein genes in sheep—a review. Genes 7:24.

Gong H, Zhou H, Hickford JGH. 2010. Polymorphism of the ovine keratin-associated protein 1-

4 gene (KRTAP1-4). Mol. Biol. Rep. 37:3377-3380.

Gong H, Zhou H, Hodge S, Dyer JM, Hickford JGH. 2015. Association of wool traits with

variation in the ovine KAP1-2 gene in Merino cross lambs. Small Rumin. Res. 124:24-29.

Gong H, Zhou H, McKenzie GW, Hickford JG, Yu Z, Clerens S, Dyer JM, Plowman JE. 2010.

Emerging issues with the current keratin-associated protein nomenclature. International Journal

of Trichology 2:104-105.

Gong H, Zhou H, McKenzie GW, Yu Z, Clerens S, Dyer JM, Plowman JE, Wright MW, Arora

R, Bawden CS. 2012. An updated nomenclature for keratin-associated proteins (KAPs). Int. J.

Biol. Sci. 8:258-264.

Gong H, Zhou H, Yu Z, Dyer J, Plowman JE, Hickford J. 2011. Identification of the ovine

keratin-associated protein KAP1-2 gene (KRTAP1-2). Exp. Dermatol. 20:815-819.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms

and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML

3.0. Syst. Biol. 59:307-321.

Harpak A, Lan X, Gao Z, Pritchard JK. 2017. Frequent nonallelic gene conversion on the human

lineage and its effect on the divergence of gene duplicates. PNAS 114:12779-12784.

Hiwatashi T, Mikami A, Katsumura T, Suryobroto B, Perwitasari-Farajallah D, Malaivijitnond

S, Siriaroonrat B, Oota H, Goto S, Kawamura S. 2011. Gene conversion and purifying selection

shape nucleotide variation in gibbon L/M opsin genes. BMC Evol. Biol. 11:312.

Itenge-Mweza TO, Forrest RH, McKenzie GW, Hogan A, Abbott J, Amoafo O, Hickford JG.

2007. Polymorphism of the KAP1.1, KAP1.3 and K33 genes in Merino sheep. Mol. Cell. Probes

21:338-342.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30:772-780.

Khan I, Maldonado E, Vasconcelos V, Stephen JO, Johnson WE, Antunes A. 2014. Mammalian

keratin associated proteins (KRTAPs) subgenomes: disentangling hair diversity and adaptation to

terrestrial and aquatic environments. BMC Genomics 15:779.

Lamping E, Zhu JY, Niimi M, Cannon RD. 2017. Role of ectopic gene conversion in the

evolution of a *Candida krusei* pleiotropic drug resistance transporter family. Genetics 205:1619-

1639.

Li S, Zhou H, Gong H, Zhao F, Hu J, Luo Y, Hickford JGH. 2017. Identification of the ovine

keratin-associated protein 26-1 gene and its association with variation in wool traits. Genes

8:225.

Li S, Zhou H, Gong H, Zhao F, Wang J, Liu X, Luo Y, Hickford JGH. 2017. Identification of the

ovine keratin-associated protein 22-1 (KAP22-1) gene and its effect on wool traits. Genes 8:27.

Li S, Zhou H, Gong H, Zhao F, Wang J, Luo Y, Hickford JGH. 2017. Variation in the ovine

KAP6-3 gene (KRTAP6-3) is associated with variation in mean fibre diameter-associated wool

traits. Genes 8:204.

Liao D. 1999. Concerted evolution: molecular mechanism and biological implications. Am. J.

Hum. Genet. 64:24-30.

Liao D, Weiner AM. 1995. Concerted evolution of the tandemly repeated genes encoding

primate U2 small nuclear RNA (the RNU2 locus) does not prevent rapid diversification of the

(CT)n.(GA)n microsatellite embedded within the U2 repeat unit. Genomics 30:583-593.

Lin Y-S, Byrnes JK, Hwang J-K, Li W-H. 2006. Codon-usage bias versus gene conversions in

the evolution of yeast duplicate genes. PNAS 103:14412-14416.

Lopez-Flores I, Garrido-Ramos MA. 2012. The repetitive DNA content of eukaryotic genomes.

Genome Dynamics 7:1-28.

Morrill SA, Exner AE, Babokhov M, Reinfeld BI, Fuchs SM. 2016. DNA instability maintains

the repeat length of the yeast RNA polymerase II C-terminal domain. J. Biol. Chem. 291:11540-

11550.

Nakamura Y, Gojobori T, Ikemura T. 2000. Codon usage tabulated from international DNA

sequence databases: status for the year 2000. Nucleic Acids Res. 28:292.

Nei M, Gu X, Sitnikova T. 1997. Evolution by the birth-and-death process in multigene families

of the vertebrate immune system. PNAS 94:7799-7806.

Nei M, Rogozin IB, Piontkivska H. 2000. Purifying selection and birth-and-death evolution in

the ubiquitin gene family. PNAS 97:10866-10871.

Nei M, Rooney AP. 2005. Concerted and birth-and-death evolution of multigene families. Annu.

Rev. Genet. 39:121-152.

Noonan JP, Grimwood J, Schmutz J, Dickson M, Myers RM. 2004. Gene conversion and the

evolution of protocadherin gene cluster diversity. Genome Res. 14:354-366.

Perina A, Seoane D, Gonzalez-Tizon AM, Rodriguez-Farina F, Martinez-Lage A. 2011.

Molecular organization and phylogenetic analysis of 5S rDNA in crustaceans of the genus

Pollicipes reveal birth-and-death evolution and strong purifying selection. BMC Evol. Biol.

11:304.

Petronella N, Drouin G. 2011. Gene conversions in the growth hormone gene family of primates:

stronger homogenizing effects in the Hominidae lineage. Genomics 98:173-181.

Petronella N, Drouin G. 2014. Purifying selection against gene conversions in the folate receptor

genes of primates. Genomics 103:40-47.

Piontkivska H, Rooney AP, Nei M. 2002. Purifying selection and birth-and-death evolution in

the histone H4 gene family. Mol. Biol. Evol. 19:689-697.

Powell BC, Rogers GE. 1997. The role of keratin proteins and their genes in the growth,

structure and properties of hair. In. Formation and structure of human hair: Birkhäuser Verlag. p.

59-148.

Puigbo P, Bravo IG, Garcia-Vallve S. 2008. CAIcal: a combined set of tools to assess codon

usage adaptation. Biol. Direct 3:38.

Richard GF, Kerrest A, Dujon B. 2008. Comparative genomics and molecular dynamics of DNA

repeats in eukaryotes. Microbiol. Mol. Biol. Rev. 72:686-727.

Rogers GR, Hickford JGH, Bickerstaffe R. 1994. Polymorphism in two genes for B2 high sulfur

proteins of wool. Anim. Genet. 25:407-415.

Rogers MA, Langbein L, Praetzel-Wunder S, Winter H, Schweizer J. 2006. Human hair keratin-

associated proteins (KAPs). Int. Rev. Cytol. 251:209-263.

Rogers MA, Schweizer J. 2005. Human KAP genes, only the half of it? Extensive size

polymorphisms in hair keratin-associated protein genes. J. Invest. Dermatol. 124:vii-ix.

Rooney AP, Ward TJ. 2005. Evolution of a large ribosomal RNA multigene family in

filamentous fungi: birth and death of a concerted evolution paradigm. PNAS 102:5084-5089.

Shimomura Y, Aoki N, Schweizer J, Langbein L, Rogers MA, Winter H, Ito M. 2002.

Polymorphisms in the human high sulfur hair keratin-associated protein 1, KAP1, gene family. J.

Biol. Chem. 277:45493.

Shyue SK, Li L, Chang BH, Li W-H. 1994. Intronic gene conversion in the evolution of human

X-linked color vision genes. Mol. Biol. Evol. 11:548-551.

Stein LD. 2004. End of the beginning. Nature 431:915-916.

Stephan W. 1989. Tandem-repetitive noncoding DNA: Forms and forces. Mol. Biol. Evol.

6:198-212.

Storz JF, Baze M, Waite JL, Hoffmann FG, Opazo JC, Hayes JP. 2007. Complex signatures of

selection and gene conversion in the duplicated globin genes of house mice. Genetics 177:481–

500.

Talavera G, Castresana J. 2007. Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. Syst. Biol. 56:564-577.

Tao J, Zhou H, Gong H, Yang Z, Ma Q, Cheng L, Ding W, Li Y, Hickford JGH. 2017. Variation

in the KAP6-1 gene in Chinese Tan sheep and associations with variation in wool traits. Small

Rumin. Res. 154:129-132.

Tao J, Zhou H, Yang Z, Gong H, Ma Q, Ding W, Li Y, Hickford JGH. 2017. Variation in the

KAP8-2 gene affects wool crimp and growth in Chinese Tan sheep. Small Rumin. Res. 149:77-

80.

Teshima KM, Innan H. 2004. The effect of gene conversion on the divergence between

duplicated genes. Genetics 166:1553-1560.

Torrents D, Suyama M, Zdobnov E, Bork P. 2003. A genome-wide survey of human

pseudogenes. Genome Res. 13:2559-2567.

Wernersson R, Pedersen AG. 2003. RevTrans: Multiple alignment of coding DNA from aligned

amino acid sequences. Nucleic Acids Res. 31:3537-3539.

Wu DD, Irwin D, Zhang YP. 2008. Molecular evolution of the keratin associated protein gene

family in mammals, role in the evolution of mammalian hair. BMC Evol. Biol. 8:241.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. GGTREE: an R package for visualization

and annotation of phylogenetic trees with their covariates and other associated data. Methods

Ecol. Evol. 8:28–36.

Zhou H, Gong H, Li S, Luo Y, Hickford J. 2015. A 57-bp deletion in the ovine KAP6-1 gene

affects wool fibre diameter. Journal of Animal Breeding and Genetics.

Zhou H, Gong H, Wang J, Dyer JM, Luo Y, Hickford JGH. 2016. Identification of four new

gene members of the KAP6 gene family in sheep. Sci. Rep. 6:24074.

Figure 1. Tandem repeat organization of the keratin associated protein-1 (*KRTAP1*) genes
The organization of mammalian *KRTAP1* genes is illustrated by the arrangement found in sheep. The four *KRTAP1-n* paralogs are represented by arrows that indicate the direction of transcription. Diagram is drawn to scale, with *KRTAP1-n* lengths bracketed below the genes. These repeats are numbered *KRTAP1-1, 3, 4*, and *5* in human.

Figure 2. Mammalian *KRTAP1-n* gene phylogenetic relationships
Representative phylogenetic tree illustrating the relationships between
the *KRTAP1-n* genes in the species used in this study. Branch lengths
are not to scale. The phylogeny is adapted from that presented in
McCormack et al. (2012).

Figure 3. Phylogenetic trees of *KRTAP1-n* coding and flanking region sequences

Phylogenetic trees were constructed for the mammalian *KRTAP1-n* 5' flanking region, coding region, and 3' flanking region using PhyML. The species are indicated by three-letter abbreviations. The number following this for the coding regions indicates the *KRTAP1-n* gene name. The major clades within the trees are indicated by coloured boxes. The 5' and 3' flanking region phylogenies group by repeat number, while the coding region phylogeny tends to group by species. Numbers on nodes indicate bootstrap supports over 50%, and substitution rates are indicated at the top left. Human *KRTAP1-n* gene names have been altered for consistency with other species.

Figure 4. The *KRTAP1-n* concerted evolution pattern is not explained by purifying selection
Phylogenetic trees were constructed for the 1st and 2nd codon positions ("non-synonymous"; **A**), and the 3rd codon position ("synonymous"; **B**), as per **Figure 3**. The major clades in both phylogenies tend to group by species, with this concerted evolution pattern being stronger for the synonymous phylogeny. Numbers on nodes indicate bootstrap supports with values over 50%, and substitution rates are indicated at the top left.

Figure 5. The switch between concerted and radiating evolution patterns is located close to the start/stop sites

A) Alignment of the region flanking the *KRTAP1-1* gene start site. The boundary between the 5' flanking and coding regions is marked by a vertical line (followed by the ATG). Underneath is an alignment of the same region for all four *KRTAP1-n* sequences from sheep. Mismatches have a white background, conservation is indicated graphically above each alignment, and consensus sequences are shown at the top. **B**) As in **(A)**, except the region flanking the stop site is shown, with the vertical line marking the boundary between the coding and 3' flanking regions (preceded by the stop codon).

Figure 6. Evidence for short gene conversion tracts between *KRTAP1-n* sequences within species
Alignment of KAP1 amino acid sequences from the ten mammalian species. Amino acid tracts boxed in green represent sequences unique to a species or related species pairs. The grey vertical box represents the conserved decapeptide repeat sequences (which have been removed). Dots represent gaps in the alignment.

Figure 7. Shared polymorphisms between *KRTAP1-n* sequences in sheep

Alignment of the four sheep *KRTAP1-n* coding region sequences. Dashes represent nucleotides identical to the top sequence, and dots represent gaps. The 30 bp repeats are not shown, as the insertion/deletion positions cannot be precisely determined. Shared nucleotide substitutions between repeat copies are highlighted in red.