1   **QTL mapping of natural variation reveals that the developmental regulator**

2   ***bruno* reduces tolerance to *P*-element transposition in the *Drosophila* female**

3   **germline**

4   Erin S. Kelleher[*][†], Jaweria Jaweria[*], Uchechukwu Akoma[*][‡], Lily Ortega[*] and Wenpei

5   Tang[*].

[*] Department of Biology and Biochemistry, University of Houston, Houston, TX
[†] To whom correspondence should be addressed: eskelleher@uh.edu
[‡] Department of Molecular Physiology and Biophysics, Baylor College of Medicine,
Houston, TX, US 77030

ABSTRACT

Transposable elements (TEs) are obligate genetic parasites that propagate in host genomes by replicating in germline nuclei, thereby ensuring transmission to offspring. This selfish replication not only produces deleterious mutations---in extreme cases, TE mobilization induces genotoxic stress that prohibits the production of viable gametes. Host genomes could reduce these fitness effects in two ways: resistance and tolerance. Resistance to TE propagation is enacted by germline specific small-RNA-mediated silencing pathways, such as the piRNA pathway, and is studied extensively. However, it remains entirely unknown whether host genomes may also evolve tolerance, by desensitizing gametogenesis to the harmful effects of TEs. In part, the absence of research on tolerance reflects a lack of opportunity, as small-RNA-mediated silencing evolves rapidly after a new TE invades, thereby masking existing variation in tolerance.

We have exploited the recent the historical invasion of the *Drosophila melanogaster* genome by *P*-element DNA transposons in order to study tolerance of TE activity. In the absence of piRNA-mediated silencing, the genotoxic stress imposed by *P*-elements disrupts oogenesis, and in extreme cases leads to atrophied ovaries that completely lack germline cells. By performing QTL-mapping on a panel of recombinant inbred lines (RILs) that lack piRNA-mediated silencing of *P*-elements, we uncovered multiple QTL that are associated with differences in tolerance of oogenesis to *P*-element transposition. We localized the most significant QTL to a small 230 Kb euchromatic region, with the LOD peak occurring in the *bruno*

locus, which codes for a critical and well-studied developmental regulator of

oogenesis. We further demonstrate that multiple *bruno* loss-of-function alleles are

strong dominant suppressors of ovarian atrophy, allowing for the development of

mature egg-chambers in the face of *P*-element activity. Genetic and cytological

analyses suggest that *bruno* tolerance is explained by enhanced retention of

germline stem cells in dysgenic ovaries, which are typically lost due to DNA damage.

Our observations reveal segregating variation in TE tolerance for the first time, and

implicate gametogenic regulators as a source of tolerant variants in natural

populations.

INTRODUCTION

Transposable elements (TEs) are omnipresent and abundant constituents of

eukaryotic genomes, comprising up to 80% of genomic DNA in some lineages

(reviewed in [1]). The evolutionary footprint of TEs is extensive, including dramatic

genome size expansions [2,3], acquisition of new regulatory networks [4,5],

structural mutations [6], novel genes [7–9], and adaptive insertions [10–12].

However, the charismatic and occasionally beneficial impact of TEs over

evolutionary time masks their fundamental identity as intragenomic parasites and

mutagens. In addition to causing deleterious mutations [13,14], TEs can exert lethal,

genotoxic effects on host cells by producing abundant double-stranded breaks

(DSBs) during insertion and excision [15,16]. TEs are therefore intragenomic

parasites.

Host developmental and evolutionary responses to parasites, pathogens, and

herbivores are broadly delineated into two categories: resistance and tolerance

(reviewed in [17,18]). Mechanisms of resistance prevent—or limit the spread of—

infection or herbivory. By contrast, mechanisms of tolerance do not effect the

propagation, but rather limit the fitness costs to the host. With respect to TEs,

resistance by eukaryotic genomes is enacted by small RNA mediated silencing

pathways [19], and Kruppel-associated box zinc-finger proteins (KRAB-ZFPs)[20],

which regulate the transcription and subsequent transposition of endogenous TEs.

However, it remains unknown whether genomes can also evolve tolerance of TEs,

by altering how host cells are affected by TE activity. Tolerance therefore represents

a wholly unexplored arena of the evolutionary dynamics between TEs and their hosts. Germline tolerance of TEs is predicted to be of particular importance, due to the significance of this cell lineage in ensuring the vertical transmission of the parasite, and the reproductive fitness of the host.

The absence of research on tolerance is at least partially due the primacy of resistance: endogenous TEs are overwhelmingly repressed by host factors in both germline and somatic tissues [19,21,22]. However, the invasion of the host genome by a novel TE family, which happens recurrently over evolutionary time scales (reviewed in [23]), provides a window of opportunity through which tolerance could be viewed, both empirically and by natural selection. The absence of evolved resistance in the host against a new invader could reveal differential responses of germline cells to unrestricted transposition. A classic example of genome invasion by a novel TE is provided by *P*-elements, DNA transposons that have recently colonized two *Drosophila* species. *P*-elements first appeared in genomes of *D. melanogaster* around 1950 (reviewed in [24]), and later colonized it's sister species *D. simulans* around 2006 [25,26]. Particularly for *D. melanogaster*, a large number of naïve strains collected prior to *P*-element invasion are preserved in stock centers and laboratories, providing a record of ancestral genetic variation in tolerance [27,28]. Furthermore, 15 of these naïve strains were recently used to develop the *Drosophila* Synthetic Population Resource (DSPR), a panel of highly Recombinant Inbred Lines (RILs) that serve as a powerful toolkit for discovering the natural genetic variants influencing quantitative traits [29–31].

Here, we harness the mapping power of the DSPR to screen for phenotypic

and genetic variation in the tolerance of the *D. melanogaster* female germline to

unrestricted *P*-element activity. We developed a novel screen for phenotypic

variation in host tolerance by taking advantage of the classic genetic phenomenon of

hybrid dysgenesis, in which TE families that are inherited exclusively paternally can

induce a sterility syndrome in offspring, due to an absence of complementary

maternally-transmitted regulatory small RNAs (piRNAs, reviewed in [24,32,33]).

The dysgenesis syndrome induced by *P*-elements in the female germline is

particularly severe, and can be associated with a complete loss of germline cells

[15,34,35]. *P*-element hybrid dysgenesis is directly related to genotoxic stress, as

apoptosis is observed in early oogenesis in dysgenic females [15], and the DNA

damage response factors *checkpoint kinase 2* and *p53* act as genetic modifiers of

germline loss [35]. Variation in the sensitivity of the DNA damage response to

double-stranded breaks therefore represents one potential cellular mechanism for

tolerance.

By phenotyping the germline development of >32,000 dysgenic female

offspring of RIL mothers, we uncovered substantial heritable variation in female

germline tolerance of *P*-element activity. We furthermore mapped this variation to a

small 230 Kb quantitative trait locus (QTL) on the second chromosome, containing

only 18 genes. Only one locus, *bruno*, is a strong candidate for the source of

causative variation. Surprisingly, *bruno* encodes a well-studied developmental

regulator of oogenesis, with no known function in TE repression or DNA damage

response [36–39]. We further demonstrate that *bruno* loss-of function alleles act as

dominant suppressors of germline loss, and relate these effects to the retention of

germline stem cells (GSCs) in dysgenic females. Our findings represent the first

demonstration of natural variation in TE tolerance in any organism. They further

implicate regulators of gametogenesis such as *bruno*, as a source of tolerant variants

that could be beneficial when new TEs invade the host.

RESULTS

Maternal genotype, but not zygotic age, is a strong predictor of dysgenic F1

atrophy.

We first sought to uncover phenotypic variation in germline tolerance of *P*-

element activity among a panel of highly recombinant inbred lines (RILs) derived

from eight founder genomes [29]. To quantify tolerance, we performed dysgenic

crosses between RIL females and males from the *P*-element containing strain

Harwich (Figure 1A). Harwich is a strong paternal inducer of hybrid dysgenesis,

producing F1 females with 100% atrophied ovaries in crosses with naïve females at

the restrictive temperature of 29°C [28]. We therefore performed our crosses at

25°C, a partially permissive temperature at which intermediate levels of ovarian

atrophy are observed [40,41]. Because *P*-element dysgenic females may recover

their fertility as they age, through zygotic production of *P*-element derived piRNAs

[42], we assayed both 3-day old and 21-day old F1 female offspring from each RIL.

In total, we documented the incidence of atrophied ovaries among 17,150 3-day-old

and 15,039 21-day-old F1 female offspring, and estimated the proportion of F1

atrophy within broods of $\geq$ 20 3-day old and 21-day old offspring from 592 and 492

RILs, respectively (Figure 1B, 1C).

We observed continuous variation in the proportion of F1 atrophy among

both 3-day old and 21-day old offspring of different RIL genotypes (Figure 1B, 1C).

After accounting for the effects of experimenter and experimental block, the

incidence of ovarian atrophy is strongly correlated among 427 RILs for which we

sampled broods of both 3-day old and 21-day old F1 females (Pearson's $R = 0.56$, $p >$

10^{-15}, Figure 1D). Because broods of different age classes were sampled from

separate crosses and experimental blocks, this correlation strongly implies that

phenotypic differences are explained by the maternal genotype. Indeed, based on

the F1 atrophy proportions measured in broods of different ages, we estimate that

the broad-sense heritability of F1 ovarian atrophy among the RIL offspring was

40.35% (see Materials and Methods). Furthermore, we saw greater reproducibility

across a small sample of 14 RILs for which we phenotyped two independent 21-day

broods (Pearson's $R = 0.97$, $p > 10^{-8}$), suggesting even higher heritability among

offspring of the same age class.

Despite the previous observation of developmental recovery from hybrid

dysgenesis [42], the relationship between age and the proportion of F1 atrophy is

only marginally significant in our data (*Likelihood Ratio Test* = 3.60, *p* = 0.058).

Furthermore, 21-day old dysgenic females exhibited only a 0.63% decrease in the

proportion of F1 atrophy when compared to 3-day old females, indicating that the

overall effect of age in our crosses was very modest. Finally, we did not observe a

group of RILs in which ovarian atrophy is much more common among 3-day-old as

compared to 21-day-old F1 females (Figure 1D), as would be predicted if there were

genetic variation for development recovery across the RIL panel. The absence of

developmental recovery of in our experiments could reflect differences in

developmental temperature between our two studies (22°C in [42] and 25°C here).

Alternatively, the causative variant that allows for developmental recovery could be

absent from the founder RILs.

QTL Mapping of Dysgenic F1 Atrophy.

To identify the genomic regions that harbor causative genetic variation in

tolerance, we performed a QTL analysis using the published RIL genotypes [29]. In

these data, the founder allele (A1-A8) carried by each RIL is inferred

probabilistically for 10 Kb windows along the euchromatic regions of the major

autosomes 2 and 3, and the X-chromosome [29]. The fourth chromosome is ignored

because the absence of recombination makes it uninformative for QTL mapping

(reviewed in [43]).

Consistent with the strongly correlated phenotypes of 3-day-old and 21-day-

old offspring (Figure 1D), we identified a single major effect QTL associated with

phenotypic variation at both developmental time points (Figure 2A). Additionally,

the Δ2-LOD drop confidence intervals (Δ2-LOD CI) of the LOD peak from each

analysis are both narrow (<300 Kb) and highly overlapping (Table 1). The peak

explains 14.2% and 14.8% of variation in ovarian atrophy among 3-day-old and 21-

day-old F1 females, indicating it is a major determinant of heritable variation.

Multiple minor peaks close to the centromere of 3L may represent another source of

heritable variation in 3-day-old females.

To further narrow the location of genetic variation in tolerance, we took

advantage of the striking concordance in QTL mapping for the 3-day old and 21-day

old datasets, and performed a combined analysis including all 660 RILs whose F1

offspring were sampled at either developmental time point. F1 female age was

included as a covariate (see methods). From this analysis we obtained a final Δ2-

LOD CI for the major QTL peak, which corresponds to a 230 Kb genomic region

containing 18 transcribed genes, 15 protein-coding and 3 non-coding (Figure 2B).

Simulation testing of the statistical properties of the DSPR indicate that a causative

variant explaining 10% of phenotypic variation lies within the Δ2-LOD drop CI 96%

of time for sample sizes of 600 RILs [30]. Therefore, given our increased sample

(660 RILs) and effect (~14%) sizes, the Δ2-LOD CI we infer for our major peak

should be conservative. Interestingly, of the genes within the QTL peak, only *bruno*

and *crooked* are highly expressed in the *D. melanogster* ovary [44]*. While *bruno* is

translational repressor whose major essential role is in oogenesis [37,45,46],

crooked is a more broadly expressed component of septate junctions, which is

essential for viability [47]. The LOD peak resides within the 138 Kb *bruno* locus

which, in addition to its function, make it the strongest candidate for the source of

causative variation.

Two classes of tolerance alleles.

We next sought to partition founder alleles at the QTL peak into phenotypic

classes, in order to better understand the complexity of causative genetic variation.

First we identified all sampled RILs whose genotype was probabilistically assigned

($p > 0.95$) to a single founder (A1-A8) at the LOD peak, and estimated the

phenotypic effect associated with each founder allele (Figure 3). We then used

stepwise regression (see Materials and Methods) to identify the minimum number

of allelic classes required to explain phenotypic variation associated with the

founder alleles. For both age classes, we found strong evidence for 2-allelic classes,

one sensitive and one tolerant, were sufficient explain phenotypic variation in

tolerance. This implies that the major QTL peak could correspond to a single

segregating genetic variant. Furthermore, with the exception of founder A8, founder

alleles were assigned to the same allelic class for both age cohorts, revealing allelic

behavior is highly biologically reproducible.

In light of the simple biallelic behavior of our phenotype, we sought to

identify polymorphisms within the founder strains whose genotypic differences

matched their phenotypic classifications (*i.e.* "in-phase" polymorphisms [29,31],

Figure 3C). We excluded A8 from these analyses due to the ambiguity of its allelic

class. In total we identified 36 in-phase single nucleotide polymorphisms (SNPs),

which potentially affect the function of only seven transcribed genes in the QTL

interval ([48] Supplementary Table 3, Figure 2B, Figure 3C). We did not identify any

in phase, segregating TE insertions, although a recent re-assembly of the founder A4

genome based on long single-molecule real-time sequencing reads suggests many

TE insertions remain un-annotated [49]. 22 of the in-phase SNPs are within the

bruno locus or immediately upstream, while none are found in the gene-body or

upstream of *crooked*. None of the 36 in-phase SNPs are non-synonymous, implying a

regulatory difference between the tolerant and sensitive alleles.

bruno is a strong dominant suppressor of P-element induced ovarian atrophy.

8   We next examined whether *bruno* loss of function alleles affected the atrophy

phenotype. For comparison, we also considered available alleles from three other

ovary-expressed genes that are located within the Δ2-LOD CI of the 21-day old or 3-

day old female analyses, but not in the combined analysis: *ced-12*, *Rab6* and *ThrRS*.

We reasoned that since the causative variant is almost certainly not recessive, being

found only in the maternal RIL genotype, mutant alleles might also exhibit non-

recessive effects. We therefore used balanced heterozygous females as mothers in

dysgenic crosses with Harwich males, and compared the incidence of F1 ovarian

atrophy among their 3-7 day-old F1 females (*mutant/+ vs. balancer/+,* Figure 4).

Strikingly, while *ced-12, Rab6* and *ThrRS* alleles had no effect on the incidence of

ovarian atrophy, two different *bruno* alleles ($bruno^{RM}$ and $bruno^{QB}$) acted as

dominant suppressors. In contrast to their balancer control siblings, who exhibited

64-75% ovarian atrophy, $bruno^{RM}/+$ and $bruno^{QB}/+$ offspring exhibited only 13%

and 20% atrophy, respectively, a dramatic reduction in the atrophy phenotype

(Figure 4). We further observed that an independently-derived *bruno* deficiency

[50,51] suppresses ovarian atrophy to a similar degree (Figure 5), indicating these

effects cannot be attributed to a shared linked variant on the *bruno^{RM}* and *bruno^{QB}*

chromosomes [45]. Importantly, these non-recessive, fertility enhancing effects of

bruno loss-of-function alleles in dysgenic females also contrast their effects on the

fertility of non-dysgenic females, where they act as recessive female steriles [45].

***bruno's* effects on ovarian atrophy are associated with germline stem cell**

retention.

bruno is a translational regulator with three known functions in *Drosophila*

melanogaster oognesis. At the start of oogenesis in the ovarian substructure called

the germaria, *bruno* is required to promote the differentiation of cystoblasts (CBs)

the immediate daughters of germline stem cells (GSCs) [37,52]. In mid-oogenesis,

Bruno protein blocks vitellogenesis if not properly sequestered by its mRNA target

oskar [38,39]. Finally, in late oogenesis, Bruno repression of *oskar* translation is

required to establish dorsoventral (DV) patterning in the subsequent embryo [36] .

This final role of *bruno* affects only the morphology of egg chambers, and but not

their production, suggesting that it can not account for *bruno's* effects in dysgenic

germlines. We therefore focused on *bruno's* earlier roles in GSC differentiation and

vitellogenesis, which are distinguishable by their dependency on *oskar* mRNA. While

bruno's functions in GSC differentiation are independent of *oskar* mRNA [36,37,53],

bruno and *oskar*'s impact on vitellogenesis are interdependent, due to the

requirement for *oskar* mRNA to sequester Bruno protein [39]*.*

To determine if the effect of *bruno* alleles on hybrid dysgenesis are

independent of *oskar* mRNA, we examined whether two *oskar* mRNA null alleles,

osk⁰ and *osk^{A87}*, as well as an *oskar* deficiency, affected the incidence of ovarian

atrophy when compared to a balancer control. We observed that *osk⁰* and *Df(3R)osk*

exhibited no effect on the atrophy phenotype, while *osk^{A87}* was associated with only

a marginal increase in ovarian atrophy ($p = 0.07$). If *bruno* suppression of ovarian

atrophy reflects reduced sequestration by *oskar* mRNA in the dysgenic female

germline, atrophy should be enhanced by *oskar* mutants.

Comparison of single and double heterozygotes of *bruno* and *oskar*, also

suggest that *bruno* suppression occurs independently of *oskar* mRNA. *Df(2L)bru/+;*

osk^{A87}/+; double heterozygotes did not differ from their single

Df(2L)bru/+;balancer/+ heterozygous siblings with respect to ovarian atrophy,

suggesting that *bruno* suppression is unrelated to *oskar* dosage. Although a second

double-heterozygote *Df(2L)bru/+;Df(3R)osk/+;* was associated with significantly

increased atrophy when compared to the single heterozygote (*Df(2L)bru/+;*

balancer/+), we suspect this is a consequence of the double-deficiency, because

Df(3R)osk alone had no effect (Figure 5). We also did not detect any Bruno

mislocalization in the developing egg chambers of wild-type dysgenic females

(Supplementary Figure 1), nor did we see any evidence of an arrest in mid-stage

oogenesis in wild-type dysgenic ovaries, as occurs when Bruno is not sequestered

by *oskar* mRNA [38,39]. Taken together, these experiments strongly suggest that

bruno's effects on oogenesis under *P*-element dysgenesis are independent of *oskar*.

To evaluate whether *bruno* suppression of ovarian atrophy could be

explained by its *oskar*-independent role in GSC differentiation [52,54], we directly

examined and compared GSCs between *bruno^QB/+* and *CyO/+* dysgenic ovaries.

While did not observe any direct evidence of delayed GSC differentiation in

bruno^QB/+ germaria, we did observe that ovarioles containing developing egg

chambers overwhelmingly retained all oogenic stages, including GSCs (Figure 6B).

In contrast, atrophied ovaries lacked any developing egg chambers including GSCs

(Figure 6A). These observations are consistent with recent evidence that GSC

retention is a key determinant of *P*-element induced ovarian atrophy [35]. They

further suggest that reduced *bruno* signaling for differentiation may stabilize GSCs

in their niche, allowing them to be retained despite the genotoxic effect of *P*-

elements.

DISCUSSION:

While the evolution of TE resistance through small-RNA-mediated silencing

is a topic of immense research interest, the existence and evolution of tolerance

factors that may reduce the fitness costs of TEs on the host remain undocumented.

By opportunistically employing a panel of genotyped recombinant inbred lines,

which uniformly lack small-RNA-mediated silencing of the recently invaded *P*-

element, we have here uncovered the first example of segregating variation in host-

TE tolerance in any organism. The natural variation in tolerance that we uncovered

is unlikely to be exceptional. A complementary QTL analysis of tolerance in DSPR

mapping population B has revealed a distinct peak of major effect in a different

genomic location (Lama and Kelleher, *unpublished*). Furthermore, major QTL peak

that we identified here in population A explains only about 35% of the heritable

variation in tolerance. Therefore, other segregating variants in the population A

RILs must also affect female germline response to *P*-element activity. Our inability to

map these variants likely reflects the fact that they are rare, exhibit small effects, or

both [30].

While differences in tolerance may be masked by resistance after small-RNA-

mediated silencing evolves, our study reveals that in the absence of silencing,

tolerance can be a major determinant of fitness. While the major peak we identified

is modest in it's functional effects, explaining ∼14% of variation in ovarian atrophy,

variation in fertility of this scale would be dramatic in the eyes of natural selection.

Segregating tolerance alleles, such as the one we have detected here, could therefore

be subject to strong positive selection during genome invasion by a novel TE.

Tolerance may therefore be an important feature of the host evolutionary response

to invading TEs. Indeed, the correlation between *P*-element dosage and the severity

of hybrid dysgenesis is poor at best, causing many to suggest that other genetic

factors, such as tolerance alleles, may also be important determinants of the

dysgenic phenotype [55–57]. Furthermore, the hybrid dysgenesis induced by recent

collections of *D. melanogster* tends to be mild when compared to collections from

the 1970's and 1980's, providing circumstantial evidence that tolerance factors may

have increased in frequency over time [57]. Once the causative variant responsible

for the tolerance phenotype we uncovered here is identified, we will be poised to

ask whether its increase in frequency has enhanced tolerance in extant populations.

Based on its high expression in the *Drosophila* female ovary [44], the

presence of 22 SNPs that are in-phase with founder QTL alleles, and the dominant

suppressive effect of classical loss-of-function alleles on dysgenic ovarian atrophy,

bruno is a very strong candidate for the source of causative variation in *P*-element

tolerance that we mapped on chromosome 2L. Identifying the causative variant(s)

within the very large (138 Kb) *bruno* locus, and understanding how its altered

function relates to hybrid dysgenesis presents an exciting challenge for future work.

On the surface, it is not obvious how *bruno* function could be related to *P*-element

activity, since *bruno* is neither a TE regulator nor a component of the DNA damage

response that responds to TE-induced genotoxicity. However, the connection is

plausibly explained by *bruno*'s function in promoting GSC differentiation [37,52]. We

recently have discovered that the DNA damage response factor p53 is ectopically

activated in the GSCs and CBs of dysgenic germlines, indicating that *P*-element

induced genotoxic stress begins at the earliest stages of oogenesis [35]. GSCs are

destabilized and lost in the presence of DNA damage [58], which explains why GSCs

are frequently absent from dysgenic ovaries (Figure 6, [15,34,35]). Because GSC

maintenance also requires the repression of differentiation factors including *bruno*

[54,59–61], we propose that *bruno* loss-of-function alleles could stabilize damaged

GSCs in their niche in dysgenic germaria. Indeed, loss-of-function mutations in two

other GSC differentiation factors, *bam* and *bgcn*, have been associated with

enhanced retention of GSCs in the niche of non-dysgenic germaria [62]. This model

is fully consistent with our observation that *bruno* suppression of ovarian atrophy is

accompanied by a rescue of oogenesis at all stages, including enhanced maintenance

of GSCs (Figure 6). It is additionally supported by the fact the *bruno* repression of *P*-

element ovarian atrophy is independent of its mRNA target oskar, precluding

bruno's function in later stages of oogenesis from explaining its repressive effects

(Figure 5).

Our observations with *bruno* reveal unexpected and novel role for

developmental regulators of gametogenesis as determinants of germline tolerance

of transposition. Interestingly, multiple regulators of female GSC maintenance and

differentiation in *Drosophila melanogaster* exhibit recent or ongoing signatures of

positive selection [63–65]. Tolerance to the selfish bacterial endosymbiont

Wolbacchia has already been implicated in driving some of this adaptive evolution

[66]. The fact that *bruno* alleles act as strong repressors of *P*-element hybrid

dysgenesis suggests that another class of parasites, transposable elements, may also

contribute to adaptive evolution of stem cell determinants.

MATERIALS and METHODS:

***Drosophila* Strains and Husbandry.** Recombinant inbred lines (RILs) from

Population A were generously provided by Stuart Macdonald. Harwich (#4264),

Ced-12^{c06760}/CyO (#17781), *Rab6^{GE13031}/CyO* (#26898), *Rab6^{D23D}/CyO* (#5821), and

ThrRS^{K04203}/CyO (#10539), were obtained from the Bloomington *Drosophila* stock

center. Harwich was sib-mated for one generation to increase homozygosity. *Bruno*

and *oskar* mutants and deficiencies, in single and double heterozygous

combinations, were generously provided by Paul MacDonald. *Canton-S* was obtained

from Richard Meisel. All flies were maintained in standard cornmeal media.

Phenotyping. Virgin RIL females were crossed to Harwich males, maintained at

25°C, and flipped onto fresh food every 3-5 days. Resulting F1 offspring were

maintained at 25°C for 3 days or 21 days at which their ovaries were examined

using a squash prep [56]. 21 day old females were transferred onto new food every

5 days as they aged to avoid bacterial growth. For the squash prep, individual

females were squashed in a food-dye solution allowed to incubate for $\geq$ 5 minutes.

After incubation, the slide was examined for the presence of stage 14, chorionated

egg chambers, which preferentially absorb the dye. Females who produced $\geq$ 1 egg

chambers were scored as having non-atrophied ovaries, and females producing 0

egg chambers were scored as having atrophied ovaries. A phenotyping schematic is

provided in Figure 1A.

Crosses and phenotyping were performed for 656 RILs across 24

experimental blocks for 3 day-old F1 females, and 606 RILs across 21 experimental

blocks for 21 day-old F1 females. If fewer than 21 F1 offspring were phenotyped for

a given cross, it was discarded and repeated if possible. In total, we phenotyped >20

3-day old and 21 days-old F1 female offspring for 592 RILs and 492 RILs,

respectively, and 660 RILs were assayed for at least one of the age groups.

QTL mapping. For age-class specific QTL mapping (3-day and 21-day old), the

arcsine transformed proportion of F1 females with atrophied ovaries produced by

each RIL was used as the response variable in a random effects multiple regression

model that included experimental block and undergraduate experimenter.

$$\boldsymbol{asin}\sqrt{atrophy\ proportion} = Block\ + Experimenter$$

7   For the combined analysis of both age classes, we used the full set of arcsine

8   transformed proportions, and accounted for female age as an additional fixed effect

9   in the regression model.

$$\boldsymbol{asin}\sqrt{atrophy\ proportion} = Block\ + Experimenter + Age$$

All models were fit using the lmer function from the lme4 package[67].

The raw residuals of the regression models above were used as the

phenotypic response for QTL analysis (Supplementary Tables 2-4), implemented

with the DSPRqtl package[29] in R 3.02 [68]. The output yields a logarithm of the

odds (LOD) score for the observed association between phenotype and genotype at

11,768 10 Kb intervals along the euchromatic arms of the X, 2 and 3rd chromosomes.

The LOD significance threshold was determined from 1,000 permutations of the

observed data, and the confidence interval around each LOD peak was identified by

a difference of -2 from the LOD peak position (Δ2-LOD).

Broad sense heritability. Maternal genotype was added as a random effect to the

models above, in order to determine the genetic variance in the phenotype (V^G). V^G

was obtained by extracting the variance component for maternal genotype using the

VarCorr() function from the nlme package [69]. Broad sense heritability (H^2) was

then estimated proportion of overall variance (V^G/V^P).

Estimation of Founder Phenotypes and QTL phasing. To estimate the phenotypic

effect of each founder at the QTL peak, we considered the residual phenotype for

each used in QTL mapping and then determined the founder allele carried by the

RIL at the LOD peak position [29]. RILs whose genotype at the LOD peak could not

be assigned to a single founder with >95% probability were discarded.

Founder alleles were phased into phenotypic classes by identifying the

minimal number of groups of alleles that describes phenotypic variation associated

with the QTL peak [29]. Briefly, founder alleles were sorted based on their average

estimated phenotypic effect, which was provided by the sampled RILs. Linear

models containing all possible sequential partitions of founder alleles were then fit,

and compared to a null model in which all founder alleles are in a single partition

using an extra sum-of-squares *F*-test. The two-partition model with the highest *F*-

statistic was retained and fixed only it if provided a significantly better fit *(P < 10⁻⁴)*

than the null model. The two partitions of founder haplotypes were then fixed, and

all possible three-partition models were explored. This process was continued until

the model fit could not be improved.

Identification of in-phase polymorphisms and TEs. Founders were assigned a

"hard" genotype for all annotated TEs [70] and SNPs [29] in the QTL window if their

genotype probability for a given allele was greater than 0.95 [29]. We then looked

for alternate alleles that were in-phase with our inferred allelic classes [29,31]: the

sensitive class (A3 and A5) and the tolerant class (A1, A2, A4, A6, and A7). A8 was

excluded because its assignment to the sensitive or tolerant class differed between

the data sets from 3-day old and 21-day old females.

Mutational analysis of candidate genes. Single and double heterozygote mutant

virgin females (*mutant/balancer*) were crossed to Harwich males at 25°C. Because

the vast majority of *Drosophila* lab stocks are *P*-element free, with the exception of

any *P*-element derived transgenes, these crosses are dysgenic. Resulting F1 dysgenic

female offspring were collected and aged at 25°C for 3-7 days, when their ovaries

were assayed using the squash prep described above [56]. The incidence of ovarian

atrophy was then compared between *mutant/+* and *balancer/+* siblings from the

same cross.

Immunolabelling. Ovaries from 3-7 day old females offspring of dysgenic crosses

were dissected and immediately fixed with 4% EM-grade methanol-free

paraformaldehyde (Thermo Scientific). Ovaries were washed with 0.1% Triton X-

100 in PBS, and blocked with 5% goat serum albumin (Sigma-Aldrich). Primary

antibody concentrations were as follows: anti-Hts 1B1 1:4 (DSHB [71]), anti-Vasa

1:40 (DSHB), anti-Bruno 1:1000 (Provided by Paul MacDonald [46]), and anti-Orb

4H8 and 6H4 1:20 (DSHB [46]). Secondary antibody concentrations were 1:500.

Microscopy. Ovaries were visualized with a SP8 Upright Confocal DM6000 CFS

(Leica) Microscope, outfitted with a 60X oil immersion lens. Images were collected

using an EM-CCD camera (Hamamatsu) and LAS-AF software (Leica).

Acknowledgements. We are grateful to Stuart Macdonald, Paul MacDonald, and

Richard Meisel for *Drosophila* strains, and Paul MacDonald for anti-Bruno

antibodies. Ryan Castelluci, Ashka Shah and Wesley Adalume assisted in

phenotyping atrophy among F1 offspring. Stuart Macdonald and Daniel Barbash

provided helpful discussion and feedback on this manuscript. This research was

funded by NSF DEB #1457800 to Erin Kelleher.

REFERENCES:

1. Chénais B, Caruso A, Hiard S, Casse N. The impact of transposable elements on

    eukaryotic genomes: from genome size increase to genetic adaptation to

    stressful environments. Gene. 2012;509: 7–15.

    doi:10.1016/j.gene.2012.07.042

2. Vitte C, Panaud O. LTR retrotransposons and flowering plant genome size:

    emergence of the increase/decrease model. Cytogenet Genome Res. Karger

    Publishers; 2005;110: 91–107. doi:10.1159/000084941

3.   Sun C, Shepard DB, Chong RA, López Arriaza J, Hall K, Castoe TA, et al. LTR

retrotransposons contribute to genomic gigantism in plethodontid

salamanders. Genome Biol Evol. 2012;4: 168–83. doi:10.1093/gbe/evr139

4.   Kunarso G, Chia N-Y, Jeyakani J, Hwang C, Lu X, Chan Y-S, et al. Transposable

elements have rewired the core regulatory network of human embryonic

stem cells. Nat Genet. 2010;42: 631–4. doi:10.1038/ng.600

5.   Lynch VJ, Leclerc RD, May G, Wagner GP. Transposon-mediated rewiring of

gene regulatory networks contributed to the evolution of pregnancy in

mammals. Nat Genet. 2011;43: 1154–9. doi:10.1038/ng.917

6.   Startek M, Szafranski P, Gambin T, Campbell IM, Hixson P, Shaw CA, et al.

Genome-wide analyses of LINE-LINE-mediated nonallelic homologous

recombination. Nucleic Acids Res. 2015; doi:10.1093/nar/gku1394

7.   Rajaei N, Chiruvella KK, Lin F, Astrom SU. Domesticated transposase Kat1 and

its fossil imprints induce sexual differentiation in yeast. Proc Natl Acad Sci.

2014;111: 15491–15496. doi:10.1073/pnas.1406027111

8.   Kapitonov V V, Jurka J. RAG1 Core and V(D)J Recombination Signal Sequences

Were Derived from Transib Transposons. Nemazee D, editor. PLoS Biol. Public

Library of Science; 2005;3: e181. doi:10.1371/journal.pbio.0030181

9.   Silva-Sousa R, López-Panadès E, Casacuberta E. Drosophila telomeres: an

example of co-evolution with transposable elements. Genome Dyn. 2012;7:

46–67. doi:10.1159/000337127

10.   Aminetzach YT, Macpherson JM, Petrov DA. Pesticide resistance via

transposition-mediated adaptive gene truncation in Drosophila. Science.

2005;309: 764–7. doi:10.1126/science.1112699

11. Mateo L, Ullastres A, González J. A Transposable Element Insertion Confers Xenobiotic Resistance in Drosophila. Feschotte C, editor. PLoS Genet. 2014;10: e1004560. doi:10.1371/journal.pgen.1004560

12. Hof AE van't, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The industrial melanism mutation in British peppered moths is a transposable element. Nature. Nature Research; 2016;534: 102–105. doi:10.1038/nature17951

13. Dupuy AJ, Fritz S, Largaespada DA. Transposition and gene disruption in the male germline of the mouse. Genesis. 2001;30: 82–8. Available: http://www.ncbi.nlm.nih.gov/pubmed/11416868

14. Spradling AC, Stern D, Beaton A, Rhem EJ, Laverty T, Mozden N, et al. The Berkeley Drosophila Genome Project gene disruption project: Single P-element insertions mutating 25% of vital Drosophila genes. Genetics. 1999;153: 135–77. Available: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1460730&tool=pmcentrez&rendertype=abstract

15. Dorogova N V., Bolobolova EU, Zakharenko LP. Cellular aspects of gonadal atrophy in Drosophila P-M hybrid dysgenesis. Dev Biol. 2017;424: 105–112. doi:10.1016/j.ydbio.2017.02.020

16. Noutsopoulos D, Markopoulos G, Vartholomatos G, Kolettas E, Kolaitis N, Tzavaras T. VL30 retrotransposition signals activation of a caspase-independent and p53-dependent death pathway associated with

mitochondrial and lysosomal damage. Cell Res. 2010;20: 553–562.

doi:10.1038/cr.2010.48

17.   Roy BA, Kirchner JW. Evolutionary dynamics of pathogen resistance and

tolerance. Evolution. 2000;54: 51–63. Available:

http://www.ncbi.nlm.nih.gov/pubmed/10937183

18.   Mauricio R. Natural selection and the joint evolution of toleranceand

resistance as plant defenses. Evol Ecol. Kluwer Academic Publishers; 2000;14:

491–507. doi:10.1023/A:1010909829269

19.   Blumenstiel JP. Evolutionary dynamics of transposable elements in a small

RNA world. Trends Genet. Elsevier Ltd; 2011;27: 23–31.

doi:10.1016/j.tig.2010.10.003

20.   Yang P, Wang Y, Macfarlan TS. The Role of KRAB-ZFPs in Transposable

Element Repression and Mammalian Evolution. Trends Genet. 2017;

doi:10.1016/j.tig.2017.08.006

21.   Senti K-A, Brennecke J. The piRNA pathway: a fly's perspective on the

guardian of the genome. Trends Genet. 2010;26: 499–509.

doi:10.1016/j.tig.2010.08.007

22.   Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an

adaptive defense in the transposon arms race. Science. 2007;318: 761–4.

doi:10.1126/science.1146484

23.   Schaack S, Gilbert C, Feschotte C. Promiscuous DNA: Horizontal transfer of

transposable elements and why it matters for eukaryotic evolution. Trends

Ecol Evol. 2010;25: 537–546. doi:10.1016/j.tree.2010.06.001

24.   Kelleher ES. Reexamining the P-Element Invasion of Drosophila melanogaster

Through the Lens of piRNA Silencing. Genetics. 2016;203: 1513–31.

doi:10.1534/genetics.115.184119

25.   Kofler R, Hill T, Nolte V, Betancourt AJ, Schlötterer C. The recent invasion of

natural Drosophila simulans populations by the P-element. Proc Natl Acad Sci.

2015;112: 201500758. doi:10.1073/pnas.1500758112

26.   Hill T, Schlötterer C, Betancourt AJ. Hybrid Dysgenesis in Drosophila simulans

Associated with a Rapid Invasion of the P-Element. PLoS Genet. 2016;12:

e1005920. doi:10.1371/journal.pgen.1005920

27.   Kidwell MG. Evolution of hybrid dysgenesis determinants in Drosophila

melanogaster. Proc Natl Acad Sci U S A. 1983;80: 1655–9. Available:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=393661&tool=p

mcentrez&rendertype=abstract

28.   Kidwell MG, Frydrk T, Novy J. The hybrid dysgenesis potential of Drosophila

melanogaster from diverse temporal and geographic origins. Drosoph Inf

Serv. 1983;59: 63–69.

29.   King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, et al. Genetic

dissection of a model complex trait using the Drosophila Synthetic Population

Resource. Genome Res. 2012;22: 1558–66. doi:10.1101/gr.134031.111

30.   King EG, Macdonald SJ, Long AD. Properties and power of the Drosophila

Synthetic Population Resource for the routine dissection of complex traits.

Genetics. 2012;191: 935–49. doi:10.1534/genetics.112.138537

31.   Long AD, Macdonald SJ, King EG. Dissecting complex traits using the

Drosophila Synthetic Population Resource. Trends Genet. 2014;30: 488–495.

doi:10.1016/j.tig.2014.07.009

32. Bucheton A. I transposable elements and I-R hybrid dysgenesis in Drosophila.

Trends Genet. 1990;6: 16–21. Available:

http://www.ncbi.nlm.nih.gov/pubmed/2158161

33. Malone CD, Lehmann R, Teixeira FK. The cellular basis of hybrid dysgenesis

and Stellate regulation in Drosophila. Curr Opin Genet Dev. 2015;34: 88–94.

doi:10.1016/j.gde.2015.09.003

34. Schaefer RE, Kidwell MG, Fausto-Sterling A. Hybrid Dysgenesis in Drosophila

melanogaster: Morphological and Cytological Studies of Ovarian Dysgenesis.

Genetics. 1979;92: 1141–52. Available:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1214061&tool=

pmcentrez&rendertype=abstract

35. Tasnim S, Kelleher ES. p53 is required for female germline stem cell

maintenance in P- element hybrid dysgenesis. Dev Biol. 2018;434: 215–220.

doi:10.1016/j.ydbio.2017.12.021

36. Kim-Ha J, Kerr K, Macdonald PM. Translational regulation of oskar mRNA by

bruno, an ovarian RNA-binding protein, is essential. Cell. 1995;81: 403–12.

Available: http://www.ncbi.nlm.nih.gov/pubmed/7736592

37. Wang Z, Lin H. Sex-lethal is a target of Bruno-mediated translational

repression in promoting the differentiation of stem cell progeny during

Drosophila oogenesis. Dev Biol. 2007;302: 160–168.

doi:10.1016/j.ydbio.2006.09.016

38. Jenny A, Hachet O, Závorszky P, Cyrklaff A, Weston MDJ, Johnston DS, et al. A

translation-independent role of oskar RNA in early Drosophila oogenesis.

Development. 2006;133. Available:

http://dev.biologists.org/content/133/15/2827

39. Kanke M, Jambor H, Reich J, Marches B, Gstir R, Ryu YH, et al. oskar RNA plays

multiple noncoding roles to support oogenesis and maintain integrity of the

germline/soma distinction. RNA. 2015;21: 1096–109.

doi:10.1261/rna.048298.114

40. Kidwell MG, Kidwell JF, Sved JA. Hybrid Dysgenesis in Drosophila

melanogaster: A Syndrome of Aberrant Traits Including Mutation, Sterility

and Male Recombination. Genetics. 1977;86: 813–33. Available:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1213713&tool=

pmcentrez&rendertype=abstract

41. Engels WR, Preston CR. Hybrid dysgenesis in Drosophila melanogaster: the

biology of female and male sterility. Genetics. 1979;92: 161–74. Available:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1213938&tool=

pmcentrez&rendertype=abstract

42. Khurana JS, Wang J, Xu J, Koppetsch BS, Thomson TC, Nowosielska A, et al.

Adaptation to P element transposon invasion in Drosophila melanogaster.

Cell. 2011;147: 1551–63. doi:10.1016/j.cell.2011.11.042

43. Hartmann MA, Sekelsky J. The absence of crossovers on chromosome *4* in

Drosophila melanogaster : Imperfection or interesting exception? Fly (Austin).

Taylor & Francis; 2017;11: 253–259. doi:10.1080/19336934.2017.1321181

44.   Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. Diversity

and dynamics of the Drosophila transcriptome. Nature. 2014;512: 393–399.

doi:10.1038/nature12962

45.   Schüpbach T, Wieschaus E. Female sterile mutations on the second

chromosome of Drosophila melanogaster. II. Mutations blocking oogenesis or

altering egg morphology. Genetics. 1991;129: 1119–36. Available:

http://www.ncbi.nlm.nih.gov/pubmed/1783295

46.   Filardo P, Ephrussi A. Bruno regulates gurken during Drosophila oogenesis.

Mech Dev. 2003;120: 289–97. Available:

http://www.ncbi.nlm.nih.gov/pubmed/12591598

47.   Nilton A, Oshima K, Zare F, Byri S, Nannmark U, Nyberg KG, et al. Crooked,

Coiled and Crimpled are three Ly6-like proteins required for proper

localization of septate junction components. Development. 2010;137: 2427–

2437. doi:10.1242/dev.052605

48.   Casper J, Zweig AS, Villarreal C, Tyner C, Speir ML, Rosenbloom KR, et al. OUP

accepted manuscript. Nucleic Acids Res. 2017;46: D762–D769.

doi:10.1093/nar/gkx1020

49.   Chakraborty M, VanKuren NW, Zhao R, Zhang X, Kalsow S, Emerson JJ. Hidden

genetic variation shapes the structure of functional elements in Drosophila.

Nat Genet. Nature Publishing Group; 2018;50: 20–25. doi:10.1038/s41588-

017-0010-y

50.   Thibault ST, Singer MA, Miyazaki WY, Milash B, Dompe NA, Singh CM, et al. A

complementary transposon tool kit for Drosophila melanogaster using P and

piggyBac. Nat Genet. 2004;36: 283–287. doi:10.1038/ng1314

51. Parks AL, Cook KR, Belvin M, Dompe NA, Fawcett R, Huppert K, et al. Systematic generation of high-resolution deletion coverage of the Drosophila melanogaster genome. Nat Genet. 2004;36: 288–92. doi:10.1038/ng1312

52. Parisi MJ, Deng W, Wang Z, Lin H. The arrest gene is required for germline cyst formation during Drosophila oogenesis. Genesis. 2001;29: 196–209. Available: http://www.ncbi.nlm.nih.gov/pubmed/11309853

53. Kim-Ha J, Smith JL, Macdonald PM. oskar mRNA is localized to the posterior pole of the Drosophila oocyte. Cell. 1991;66: 23–35. Available: http://www.ncbi.nlm.nih.gov/pubmed/2070416

54. Xin T, Xuan T, Tan J, Li M, Zhao G, Li M. The Drosophila putative histone acetyltransferase Enok maintains female germline stem cells through regulating Bruno and the niche. Dev Biol. 2013;384: 1–12. doi:10.1016/j.ydbio.2013.10.001

55. Bergman CM, Han S, Nelson MG, Bondarenko V, Kozeretska I. Genomic analysis of *P* elements in natural populations of *Drosophila melanogaster*. PeerJ. PeerJ Inc.; 2017;5: e3824. doi:10.7717/peerj.3824

56. Srivastav SP, Kelleher ES. Paternal Induction of Hybrid Dysgenesis in *Drosophila melanogaster* Is Weakly Correlated with Both *P* -Element and *hobo* Element Dosage. G3: Genes|Genomes|Genetics. 2017; g3.117.040634. doi:10.1534/g3.117.040634

57. Ignatenko OM, Zakharenko LP, Dorogova N V, Fedorova SA. P elements and the determinants of hybrid dysgenesis have different dynamics of

propagation in Drosophila melanogaster populations. Genetica. 2015;143: 751–9. doi:10.1007/s10709-015-9872-z

58. Ma X, Han Y, Song X, Do T, Yang Z, Ni J, et al. DNA damage-induced Lok/CHK2 activation compromises germline stem cell self-renewal and lineage differentiation. Development. 2016;143: 4312–4323. doi:10.1242/dev.141069

59. Li Y, Minor NT, Park JK, McKearin DM, Maines JZ. Bam and Bgcn antagonize Nanos-dependent germ-line stem cell maintenance. Proc Natl Acad Sci. 2009;106: 9304–9309. doi:10.1073/pnas.0901452106

60. Ma X, Zhu X, Han Y, Story B, Do T, Song X, et al. Aubergine Controls Germline Stem Cell Self-Renewal and Progeny Differentiation via Distinct Mechanisms. Dev Cell. 2017;41: 157–169.e5. doi:10.1016/j.devcel.2017.03.023

61. Chen D, McKearin DM. A discrete transcriptional silencer in the bam gene determines asymmetric division of the Drosophila germline stem cell. Development. 2003;130: 1159–70. Available: http://www.ncbi.nlm.nih.gov/pubmed/12571107

62. Jin Z, Kirilly D, Weng C, Kawase E, Song X, Smith S, et al. Differentiation-Defective Stem Cells Outcompete Normal Stem Cells for Niche Occupancy in the Drosophila Ovary. Cell Stem Cell. 2008;2: 39–49. doi:10.1016/j.stem.2007.10.021

63. Civetta A, Rajakumar SA, Brouwers B, Bacik JP. Rapid evolution and gene-specific patterns of selection for three genes of spermatogenesis in Drosophila. Mol Biol Evol. 2006;23: 655–62. doi:10.1093/molbev/msj074

1   64.   Bauer DuMont VL, Flores HA, Wright MH, Aquadro CF. Recurrent positive

2   selection at bgcn, a key determinant of germ line differentiation, does not

3   appear to be driven by simple coevolution with its partner protein bam. Mol

4   Biol Evol. 2007;24: 182–91. doi:10.1093/molbev/msl141

5   65.   Flores HA, DuMont VLB, Fatoo A, Hubbard D, Hijji M, Barbash DA, et al.

6   Adaptive Evolution of Genes Involved in the Regulation of Germline Stem Cells

7   in *Drosophila melanogaster* and *D. simulans*. G3:

8   Genes|Genomes|Genetics. 2015;5: 583–592. doi:10.1534/g3.114.015875

66.   Flores HA, Bubnell JE, Aquadro CF, Barbash DA. The Drosophila bag of

marbles Gene Interacts Genetically with Wolbachia and Shows Female-

Specific Effects of Divergence. Malik HS, editor. PLOS Genet. 2015;11:

e1005453. doi:10.1371/journal.pgen.1005453

67.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models

Using lme4. J Stat Softw. 2015;67: 1–48. doi:10.18637/jss.v067.i01

68.   R Development Core Team. R: A Language and Environment for Statistical

Computing [Internet]. Vienna, Austria; 2008. Available: http://www.r-

project.org

69.   Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. {nlme}: Linear and

Nonlinear Mixed Effects Models [Internet]. 2016. Available: http://cran.r-

project.org/package=nlme

70.   Cridland JM, Macdonald SJ, Long AD, Thornton KR. Abundance and

distribution of transposable elements in two Drosophila QTL mapping

resources. Mol Biol Evol. 2013;30: 2311–27. doi:10.1093/molbev/mst129

71.   Zaccai M, Lipshitz HD. Role ofAdducin-like (hu-li tai shao) mRNA and protein

localization in regulating cytoskeletal structure and function

duringDrosophila oogenesis and early embryogenesis. Dev Genet. 1996;19:

249–257. doi:10.1002/(SICI)1520-6408(1996)19:3<249::AID-

DVG8>3.0.CO;2-9

72.   Hoskins RA, Carlson JW, Wan KH, Park S, Mendez I, Galle SE, et al. The Release

6 reference sequence of the Drosophila melanogaster genome. Genome Res.

2015;25: 445–58. doi:10.1101/gr.185579.114

Data Set	LOD Score	Peak Position	Δ2-LOD CI	% variation
3-days	19.59	2L:12260000	2L:12080000-12380000	14.2
21-days	17.22	2L:12250000	2L:12110000-12390000	14.8
combined	34.90	2L:12260000	2L:12120000-12350000	13.8

Table 1. QTL peak positions. The position of the major QTL peak for 3-day old F1

females, 21-day old F1 females, and the combined data set are shown. For each

analysis, the peak position an Δ2-LOD drop confidence interval in *Drosophila*

melanogaster genome release 6 (dm6 [72]) are provided. The percent of phenotypic

variation explained by the QTL peak is based on the genotype of each sampled RIL at

the LOD peak position.

Figure 1. Heritable phenotypic variation in tolerance of *P*-element activity in the female germline. A) Crossing scheme for documenting variation in tolerance of *P*-element activity among RIL offspring. Representative images of atrophied and normal ovaries were published previously [35]. B and C) phenotypic variation in the proportion of 3-day old (B) and 21-day old (C) F1 female offspring of different RILs. RILs are sorted according the proportion of F1 atrophy observed in their offspring.

D) Scatterplot of the arsine-transformed proportion of F1 atrophy observed among

3-day old and 21-day old offspring of the same line, after accounting for the effects

of experimenter and experimental block. Specifically, adjusted F1 atrophy values are

the residuals of a random effects linear model including block and student

experimenter (see methods).

Figure 2. QTL mapping of variation in *P*-element tolerance. A) Logarithm of the

odds ratio (LOD) of the observed association between maternal RIL genotype [29],

and the adjusted proportion of F1 atrophy phenotype. Adjusted proportions of F1

atrophy are the residuals of a random effects linear regression that included

experimenter and experimental block. Higher LOD scores correspond to stronger

evidence of linkage, and significant LOD scores are above the threshold (dotted

line), which was obtained from 1,000 permutations of the observed data. B) 2 LOD-

drop confidence interval of the QTL peak based on a combined QTL analysis

1   including both 3-day old and 21-day old F1 females. Genes indicated by red letters

2   are potentially effected by polymorphisms in the RIL founders that are in phase the

3   inferred allelic classes (Figure 3 [48]). Gene boxes indicated red are highly

4   expressed in the *D. melanogaster* ovary [44].

A)

B)

C)

A1	**TTCCACAAGGA**
A2	**TTCCACAAGGA**
A3	**TTCCAGAAGGA**
A4	**TTCCACAAGGA**
A5	**TTCCAGAAGGA**
A6	**TTCCACAAGGA**
A7	**TTCCACAAGGA**
A8	**TTCCACAAGGA**

Figure 3. Phenotypic classes of founder alleles at the major QTL peak. The adjusted F1 atrophy among 3-day old females A) and 21-day old females B) is shown for RILs carrying each of the 8 founder alleles. QTL phasing (see methods) detects two allelic classes for both the 3-day old and 21-day old phenotypes, a sensitive allele that increases the odds F1 ovarian atrophy (pink) and a tolerant allele that decreases the odds of F1 ovarian atrophy 1 (red). The assignment of founder alleles to phenotypic classes across ages is consistent for all founders except A8. C) An example in-phase SNP in the population A RIL founder genomes at 2L:12225285 (*Drosophila melanogaster* reference genome dm6 [72]).

Figure 4. Mutational analysis of candidate genes. Loss of function heterozygotes for four candidate genes are compared to their siblings inheriting the balancer chromosome *CyO*, in order to detect zygotic effects on the incidence of ovarian atrophy among 3-7 day-old dysgenic F1 offspring. Only *bruno* alleles are significantly associated with altered odds of ovarian atrophy, acting as dominant suppressors of the atrophy phenotype.

Figure 5. Mutational analysis of *bruno* and *oskar* single and double mutants.

Single and double loss of function and deficiency heterozygotes for *bruno* and *oskar*

are compared to their siblings inheriting the balancer chromosomes *CyO* (*bruno*)

and *TM2, TM3* or *MKRS* (*oskar*), in order to detect zygotic effects on the incidence of

atrophy among 3-7 day-old dysgenic F1 offspring. *Bruno* crosses are the same as in

Figure 4, and are provided as a reference. While *bruno* alleles act as consistent

dominant suppressors in both single and double mutants, *oskar* deficiencies and

mRNA null mutants exhibit only stochastic effects on the atrophy phenotype,

1 suggesting that the mechanism of *bruno* suppression of F1 atrophy is independent

2 of *oskar* mRNA function.

A)

Hts-1B1   Vasa   DNA

B)

Figure 6. Reduced atrophy in *bruno^{QB}/+* ovaries is associated with enhanced

retention of GSCs. A) Atrophied ovaries, which are less frequent in *bruno^{QB}/+*

dysgenic females than their *CyO* siblings (Figure 4), lack any vasa-positive germline

cells in the ovarioles, including GSCs. Nuclear Vasa external to Hts-1B1 corresponds

to somatic ovarian sheath [35]. B) Non-atrophied (*i.e.* morphologically normal)

ovaries, which are more common among *bruno^{QB}/+* dysgenic females than their *CyO*

siblings (Figure 4), exhibit the full range of developing oocytes, including GSCs

(white arrow). Arrowheads correspond to cystoblasts (CBs), the undifferentiated

daughters of GSCs. Cytological markers are hu li tai shao (Hts-1B1), which labels

somatic follicle cell membranes the circular fusomes of GSCs and CBs, and the

spectrosomes that connect cells in the developing cyst, Vasa, which labels cytoplasm

of germline cells and nuclei of the ovarian sheath, and DAPI staining of nuclear DNA.