The complex architecture and epigenomic impact of plant T-DNA insertions

Florian Jupe[1,3,*], Angeline C. Rivkin[1*], Todd P. Michael[4*], Mark Zander[1,2,6*], S. Timothy Motley[4],

Justin P. Sandoval[1], R. Keith Slotkin[5], Huaming Chen[1], Rosa Castanon[1], Joseph R. Nery[1], and

Joseph R. Ecker[1,2,6,$]

[1]Genomic Analysis Laboratory, Salk Institute for Biological Studies, La Jolla, CA 92037 USA

[2]Plant Biology Laboratory, Salk Institute for Biological Studies, La Jolla, CA, 92037, USA

[3]Current address: Bayer Crop Science, Chesterfield, MO 63017 USA

[4]J. Craig Venter Institute, La Jolla, CA 92037 USA

[5]Donald Danforth Plant Science Center, Creve Coeur, MO 63132 USA

[6]Howard Hughes Medical Institute, The Salk Institute for Biological Studies, La Jolla, CA 92037

USA

[*]These authors contributed equally to this work

[$]Correspondence: ecker@salk.edu

Keywords: T-DNA, Agrobacterium, *Arabidopsis thaliana*, nanopore sequencing, optical

mapping, siRNA, DNA methylation, chromatin, transgenic, silencing

BNG: Bionano Genomics Irys optical genome mapping

CaMV 35S: cauliflower mosaic virus 35S

Col: *Arabidopsis thaliana* Columbia ecotype, wild type control

LB: T-DNA left border

NOSp: nopaline synthase promoter

NOSt: nopaline synthase terminator

ONT: Oxford Nanopore Technologies MinION long-read nanopore sequencer

PRC2: polycomb-repressive complex-2

RdDM: RNA-directed DNA methylation

RNAi: RNA interference

siRNA: small interfering RNA

sRNA: small RNA

T-DNA: transfer DNA

Ti: Tumor-inducing

Abstract

The bacterium *Agrobacterium tumefaciens* has been the workhorse in plant genome engineering. Customized replacement of native tumor-inducing (Ti) plasmid elements enabled insertion of a sequence of interest called Transfer-DNA (T-DNA) into any plant genome. Although these transfer mechanisms are well understood, detailed understanding of structure and epigenomic status of insertion events was limited by current technologies. Here we applied two single-molecule technologies and analyzed *Arabidopsis thaliana* lines from three widely used T-DNA insertion collections (SALK, SAIL and WISC). Optical maps for four randomly selected T-DNA lines revealed between one and seven insertions/rearrangements, and the length of individual insertions from 27 to 236 kilobases. *De novo* nanopore sequencing-based assemblies for two segregating lines partially resolved T-DNA structures and revealed multiple translocations and exchange of chromosome arm ends. For the current TAIR10 reference genome, nanopore contigs corrected 83% of non-centromeric misassemblies. The unprecedented contiguous nucleotide-level resolution enabled an in-depth study of the epigenome at T-DNA insertion sites. SALK_059379 line T-DNA insertions were enriched for 24nt small interfering RNAs (siRNA) and dense cytosine DNA methylation, resulting in transgene silencing via the RNA-directed DNA methylation pathway. In contrast, SAIL_232 line T-DNA insertions are predominantly targeted by 21/22nt siRNAs, with DNA methylation and silencing limited to a reporter, but not the resistance gene. Additionally, we profiled the H3K4me3, H3K27me3 and H2A.Z chromatin environments around T-DNA insertions using ChIP-seq in SALK_059379, SAIL_232 and five additional T-DNA lines. We discovered various effect s ranging from complete loss of chromatin marks to the *de novo* incorporation of H2A.Z and trimethylation of H3K4 and H3K27 around the T-DNA integration sites. This study provides new insights into the structural impact of inserting foreign fragments into plant genomes and demonstrates the utility of state-of-the-art long-range sequencing technologies to rapidly identify unanticipated genomic changes.

Introduction

Plant genome engineering using the soil microorganism *Agrobacterium tumefaciens* has

revolutionized plant science and agriculture by enabling identification and testing of gene

functions and providing a mechanism to equip plants with superior traits [1, 2, 3]. Transfer DNA

(T-DNA) insertional mutant projects have been conducted in important dicot and monocot

models, and over 700,000 lines with gene affecting insertions have been generated in

Arabidopsis thaliana (*Arabidopsis* henceforth) alone (reviewed in O'Malley [4]). Targeted T-DNA

sequencing approaches were conducted on approximately 325,000 of these lines to identify the

disruptive transgene insertions and to link genotype with phenotype. This wealth of sequence

information, much of which has been made available prior to publication, is available at:

http://signal.salk.edu/Source/AtTOME_Data_Source.html, has been iteratively updated since

2001, and accessed by the community over 10 million times by 2017.

The *Agrobacterium* strains used in research projects are no longer harmful to the plant

because the oncogenic elements of the tumor-inducing (Ti) plasmid have been replaced by a

customizable cassette that includes a diverse set of *in planta* regulatory elements.

Agrobacterium-mediated transgene integration occurs through excision of the T-DNA strand

between two imperfect terminal repeat sequences [5], the left border (LB) and right border (RB)

[6], and translocation into the host genome (reviewed in Nester [7]). Hijacking the plant

molecular machinery, the T-DNA is integrated at naturally occurring double strand breaks

through annealing and repair at sites of microhomology [8, 9]. While the exact mechanisms

behind this error prone integration are poorly understood, it is known that insertion events

generally occur at multiple locations throughout the genome [5, 10]. T-DNA insertions also

frequently contain the vector backbone and occur as direct or inverted repeats of the T-DNA,

resulting in large intra- and inter-chromosomal rearrangements [6, 11, 12, 13, 14, 15, 16, 17, 18,

19, 20, 21]. The phenomenon of long T-DNA concatemers has previously been attributed to the

replicative T-strand amplification specific to the floral dip method, and which is less often

observed after tissue explant transformation of roots or leave discs [22].

Knowledge of structural variations induced by transgene insertions, including insertion

site, copy number and potential backbone insertions, as well as evidence for epigenetic

changes to the host genome is crucial from scientific as well as regulatory perspectives. These

aspects are routinely assessed using laborious Southern blotting, Thermal Asymmetric

Interlaced (TAIL) PCR, targeted short-read sequencing, or recently digital droplet PCR [4, 23].

One of the few attempts to gain deeper insight into an engineered genome was for transgenic

Papaya, using a Sanger sequencing approach [14]. This work identified three insertion events,

each less than 10 kilobases (kb) in length, however large repeat structures with high sequence

identity are generally impossible to assemble using short-read sequences [24].

Current knowledge of structural genome changes and epigenetic stability in transgenic

plants is limited. In this study we report on the genome structures of four *Arabidopsis* T-DNA

floral-dip transformed plants, and for the first time we report the lengths of T-DNA insertions up

to 236 kilobases along with long-molecule evidence for genome structural rearrangements

including chromosomal translocations and induced epigenomic variation. To study such large

insertions and rearrangements at the sequence level, we *de novo* assembled the genomes of

two multi-insert lines (SALK_059379 and SAIL_232) and the reference accession Columbia-0

(Col-0) using Oxford Nanopore Technologies MinION (ONT) reads to very high contiguity. We

present polished contigs that span chromosome arms and reveal the scrambled nature of T-

DNA and vector backbone insertions and rearrangements in high detail. We subsequently

tested transgene expression and functionality and show differential epigenetic effects of the

insertions on the transgenes between the two tested vector backbones. Small interfering RNA

(siRNA) species induced transgene silencing through the RNA-directed DNA methylation

(RdDM) pathway of the entire T-DNA strand in the SALK-vector background, and in contrast the

transgene remained active in the SAIL-line. Moreover, by profiling the occupancy of H3K4me3,

H3K27me3 and H2A.Z in SALK_059379, SAIL_232 and five additional T-DNA lines, we

uncovered various effects of T-DNA insertions on the adjacent chromatin landscape. In

summary, new technological advances have enabled us to assemble and analyze the genomes

and epigenomes of T-DNA insertion lines with unprecedented detail, revealing novel insights

into the impact of these events on plant genome/epigenome integrity.

Results

BNG optical genome maps reveal size and structure of T-DNA insertions

To study both the location and size of T-DNA insertions into the genome, we randomly

selected four transgenic *Arabidopsis* lines and assembled optical genome maps from nick-

labelled high-molecular weight DNA molecules using the Bionano Genomics Irys system (BNG;

San Diego, CA) from pooled leaf tissue. These Columbia-based (Col) plant lines have

previously been transformed by *Agrobacterium* using different vector constructs: SALK_059379

and SALK_075892 with pROK2 (T-DNA strand 4.4 kb; [25]), SAIL_232 with pCSA110 (T-DNA

strand 7.1 kb; [26, 27]), and WiscDsLox_449D11 with pDSLox (T-DNA strand 9.1 kb; [20]). The

segregating plant lines resemble their Columbia parent plants showing no obvious visual T-DNA

insertion-induced phenotypes. The fluorescently-labeled DNA molecules used for BNG mapping

had an average length of up to 288 kb, which enables conservation of long-range information

across transgene insertion sites (Table 1 and Fig. 1, 2; Supplementary Table 1). We compared

assembled BNG genome maps for each of the four lines to the *Arabidopsis* TAIR10 reference

genome sequence and observed up to four structural variations, due to T-DNA insertions, with

sizes ranging from 27 kb to 236 kb (Fig. 1, 2, Table 1, and Supplementary Table 2). Although

insertion sizes larger than the actual T-DNA cassette have been reported [6, 11, 12, 17, 20, 21],

the measured length for these insertions exceeded expectations based on vector size by ~20-

60 fold. Moreover, BNG genome maps of the SAIL_232 line identified a total of seven genomic

changes including three insertion events, one inversion involving ~500 kb on chromosome 1, an

inverted translocation on chromosome 3 that involves the exchange of two adjacent regions

between 2.6-3.4Mb (847 kb) with 8.9-10.1 Mb (1,193 kb), as well as a swap between

chromosome arm ends (chromosomes 3 and 5; Fig. 2). Previous short-read sequencing

projects to identify insertion numbers (T-DNA seq; http://signal.salk.edu/cgi-bin/tdnaexpress)

provided evidence for two (WiscDsLox_449D11), three (SALK_075892), four (SALK_059379)

and five (SAIL_232) insertion sites, thus less than what we have observed through the BNG

mapping approach (Supplementary Table S5).

Assembly of highly contiguous genomes from ONT MinION data

The number of insertions in SALK_059379 and the type of rearrangements observed in

SAIL_232 sparked our interest in analyzing these genomes at greater (nucleotide) resolution.

We sequenced these engineered genomes, alongside the parent reference Col-0 plant (ABRC

accession CS70000) using the Oxford Nanopore Technologies (ONT) MinION device. We

performed ONT sequencing on each line using a single R9.4 flow cell (Table 1 and

Supplementary Table 1) and assembled each genome using minimap/miniasm followed by

three rounds of racon [28] and one round of Pilon [29]. We assembled the three lines into 40

contigs (Col-0; longest 16,115,063 bp), 59 contigs (SAIL_232; longest 16,070,966 bp) and 139

contigs (SALK_059379; longest 8,784,268 bp) (Supplementary Table 1). Individual whole

genome alignments to the TAIR10 reference show over 98% coverage with 39 and 57 contigs

for SAIL_232 and SALK_059379, respectively (Table 1; Supplementary Fig. 1; Supplementary

Table 2).

The remaining short contigs (< 50 kb) encode only highly repetitive sequences such as

ribosomal DNA and centromeric repeats that cannot be placed onto the reference.

Chromosome arms are generally contained within one or two contigs, and contiguity declines

with repeat content towards the centromere (Supplementary Fig. 1). Chromosome arm-

spanning contigs covered telomere repeats and at least the first centromeric repeat, thus

capturing 100% of the genic content. When we aligned the contigs of all assembled genomes

back to the TAIR10 reference, we consistently did not cover ~3.9Mb of the reference. The Col-0

contigs covered over 99% of the TAIR10 reference, and the only discrepancies occur at the

centromeres (Supplementary Fig. 1). The high contiguity and quality of this genome assembly

allowed correction of previously identified misassemblies (38/46 'N'-regions) in the TAIR10

reference genome (Fig. 1a, Supplementary Table 3). Our contigs were not able to span the

remaining eight. BNG alignments confirmed that all ONT contigs were chimera-free, while only

eleven (Col-0 and SALK_059379) or three contigs (SAIL_232) contained misassembled non-T-

DNA repeats (Supplementary Table 4).

One aim of creating near complete genome assemblies was to enable the structural

resolution of transgene insertions at nucleotide level, rather than with genome scaffolding alone.

We next assessed contiguity at sites of T-DNA insertion. After alignment these assemblies to

BNG maps we concluded that the shorter insertions, SALK:chr2_18Mb (28 kb), SAIL:chr3_9Mb

(11 kb), SAIL:chr3_21Mb (25 kb; Fig. 2c) and SAIL:chr5_22Mb (11 kb) were completely

assembled. Because of extensive repeats, much larger T-DNA insertions collapsed upon

themselves, although contigs reaching up to 39 kb into the insertions from the flanking genomic

sequences could be assembled.

T-DNA independent chromosomal inversion in the SAIL Col-3 background

Chromosome 1 in the SAIL_232 line was assembled into a single contig

(SAIL_contig_20), spanning an entire chromosome arm from telomere to the first centromeric

repeat arrays (Supplementary Fig. 1). Compared with the Col-0 reference genome, we found a

512 kb inversion in the upper arm (SAIL_chr1:11,703,634-12,215,749). Because we could not

find a signature of T-DNA insertion at the inversion edges, we posited that this event may have

resulted from a preexisting structural variation of the particular Columbia strain used for the

SAIL project (Col-3; ABRC accession CS873942). To test this hypothesis, we genotyped

SAIL_232 alongside two randomly selected lines of the same collection (SAIL_59 and

SAIL_107) using primers specific to the reference Col-0 CS70000 genome and the SAIL-

inverted state (see Methods). PCR analysis confirmed that this "inversion" was common to all

three independent SAIL-lines tested, and absent from Col-0 CS70000 (Supplementary Fig. 2).

Thus the event was not due to the T-DNA mutagenesis, rather is an example of the genomic

"drift" occurring during the propagation Columbia "reference" accession within individual

laboratories.

SALK_059379 T-DNA insertions are conglomerates of T-strand and vector backbone

To annotate T-DNA insertion sites within the assembled genomes, we searched for

pROK2 and pCSA110 plasmid vector sequence fragments within the assembled contigs

(Supplementary Table 5). The assembled SALK_059379 genome contained three of the four

BNG map identified T-DNA insertions: SALK:chr1_5Mb, chr2_15Mb and chr2_18Mb (Table 1,

Fig. 1 a,c,e). Specifically, SALK:chr2_18Mb, the shortest identified insertion with 28,356 bp, was

completely assembled (contig_7:3,690,254-3,719,373) and included a genomic deletion of

5,497 bp (chr2:18,864,678-18,870,175). Annotation of the insertion revealed two independent

insertions; a T-DNA/backbone-concatemer (11,838 bp) from the centromere proximal end, and

a T-DNA/backbone/T-DNA-concatemer (16,463 bp) from the centromere distal end, both linked

by a guanine-rich (26 G bases) segment of 55 bp (Fig. 1e). Two independent insertion events,

5,497 bp apart, potentially created a double-hairpin through sequence homology, that was

eventually excised and removed the intermediate chromosomal stretch (Supplementary Fig. 3).

The second and third insertions, SALK:chr1_5Mb (131 kb) and SALK:chr2_15Mb (207

kb), were partially assembled into contigs. SALK:chr1_5Mb contig_10 and contig_5 contain

25,132 bp and 33,736 bp T-DNA segments. Similarly, the extremely long chr2_15Mb insertion

was partially contained within contig_4 and contig_7 (Fig. 1c), leaving an unassembled gap of

131 kb. The recovered structure of this insertion is noteworthy as it represents a conglomerate

of intact T-DNA/backbone concatemers, as well as various breakpoints that introduced partial

vector fragments with frequent changes of the insertion direction (Fig. 1c). Finally, the fourth

insertion SALK_059379 (SALK:chr4_10Mb) was absent from the assembly. However, we

observed a single ONT read (length 10,118 bp) supporting the presence of an insertion at this

location. We also recovered a further ONT read (length 15,758 bp) that anchors at position

chr3:20,141,394 and extends 14,103 bp into a previously unidentified T-DNA insertion

(Supplementary Fig. 4). PCR amplification from DNA samples using the genomic/T-DNA

junctions sequences from segregant and homozygous seeds, in fact, confirmed the presence of

all five insertions, revealing heterozygosity within the ABRC-sourced seed material.

While BNG maps were successful in placing long "T-DNA only" sequence contigs into

the large gaps (e.g. SALK:chr1_5Mb), the four "short" T-DNA-only sequence contigs of ~50 kb

or less did not contain sufficiently unique nicking pattern to confidently facilitate contig

placement.

Large-scale rearrangements reshape the SAIL_232 genome

We next searched the SAIL_232 ONT contigs for pCSA110 vector fragments, and were

able to confirm all BNG-observed genome insertions (Table 1). This search additionally

identified a further T-DNA insertion at chr5:20,476,509 (Supplementary Table 5) that was not

assembled in the BNG maps.

We found that chromosome 3 harbored two major rearrangements (Fig. 2). The first was

a translocation of a 1.19 Mb fragment (chr3:8,902,305-10,095,395), which split at an internal T-

DNA insertion at reference position 9,343,053 bp (Fig. 2a). The resulting two fragments were

independently inverted prior to integration just before position chr3:2,586,494 (Fig. 2 a,b). The

second major change was a swap between the distal arms of chromosomes 3 and 5, which is

supported by two SAIL_232 BNG genome maps as well as two ONT contigs (Fig. 2a-d). Here,

chromosome 3 broke at 21,094,402 nt, and chromosome 5 at 18,959,379 nt and 20,476,664 nt,

and the larger chromosomal fragments swapped places. Specifically, this translocation was

captured in SAIL_contig_31, showing the fusion of chr5:20,476,664-end to chromosome 3 after

position 21,094,402 nt. The reciprocal event joined (SAIL_contig_2) chromosome 3 fragment

(21,094,407-end), almost seamlessly to chromosome 5 after reference position 18,959,379 nt

(Fig. 2c,d). The genomic location of an excised fragment of chromosome 5 (chr5:18,959,380 -

20,476,663 found within SAIL_contig_11), was not determined (Fig. 2d).

Finally, the 81-kb insertion at SAIL:chr1_19Mb was identified from a BNG contig that

aligned together with a contig that did not harbor the insertion (Fig. 2e). This apparent phasing

of this heterozygous region was the only occasion where we observed this. The insertion

consists of four tandem T-DNA copies (~30 kb) followed by ~20 kb of breakpoint interspersed T-

DNA and vector backbone was partially assembled (50,676 bp) at the 5' end of contig_5 (Fig.

2e). Although not assembled as part of the flanking SAIL_contig_47, we recovered single ONT

read that contain T-DNA as well as genomic DNA sequences. In summary, the BNG maps

perfectly aligned with the ONT assembly of the T-DNA insertion haplotype (Fig. 2e).

T-DNA Integration occurs independently from both double-strand break ends

While both sequenced lines share similar numbers of T-DNA insertion events, the

genome of the SAIL_232 plant line underwent more significant changes to its architecture. All

genome insertion sites began and ended with the LB of the T-DNA strand, providing evidence

for independent transgene integration at both ends of the DNA double-strand break. We did not

recover any LB sequences at the chromosome/T-DNA junction, in line with literature reports that

usually 73 - 113 bp are missing from the LB sequence inwards [15, 30]. Internal T-DNA

sequence deletions were also seen at breakpoints within the insertion (Fig. 1c, Fig. 2c). As

observed for the SALK:chr2_18Mb chromosomal deletion (Fig. 1e and Supplementary Fig. 3),

we cannot exclude that long homologous stretches between the independently inserted T-

DNA/vector backbone concatemers represent inverted repeats.

Transgenes are functional in SAIL-lines but are silenced in SALK-lines

We next wanted to assess the effects of T-DNA insertions on the epigenomic landscape.

The pROK2 T-DNA strand contains the kanamycin antibiotic-resistance gene *nptII* under control

of the bacterial nopaline synthase promoter (NOSp) and terminator (NOSt), and an empty

multiple cloning site under control of the widely used Cauliflower mosaic virus 35S (CaMV 35S)

promoter and NOSt. The CaMV 35S constitutive overexpression promoter has previously been

described to cause homology-dependent transcriptional gene silencing (TGS) in crosses with

other mutant plants that already contain a CaMV 35S promoter driven transgene [31, 32]. In

germination assays, we confirmed that the kanamycin selective marker is not functional in SALK

lines propagated for more than a few generations [4]. While ~75% of SALK_059379 seed

initially germinated on kanamycin-containing plates, these seedlings stopped growth after

primary root and cotyledons emergeance (Fig. 3a). The SAIL_232 pCSA110 T-DNA segment

encodes the herbicide resistance gene *bar* (phosphinothricin acetyltransferase), under control of

a mannopine synthase promoter [26]. In contrast to SALK_059379, we confirmed proper

transgene function by applying herbicide to soil-germinated plants [33, 34] (Fig. 3b). We

corroborated these differential phenotypes by mapping RNA-seq reads to the corresponding

transformation plasmid and found that the SAIL_232 *bar* gene was expressed, while in

SALK_059379 the *nptII* gene was not expressed, most likely due to epigenetic silencing (Fig.

3c,d).

Differential siRNA species define transgene silencing

We hypothesized that the observed plasmid dependent transgene expression is dictated

by gene silencing via RdDM pathways [35]. To test for the presence of small RNA (sRNA) and

cytosine DNA methylation within T-DNA loci, we sequenced sRNA populations, as well as

bisulfite-converted whole genome DNA libraries for both lines. Our analyses identified abundant,

yet divergent sRNA species (15-20nt, 21nt, 22nt, 23nt, 24nt) that mapped to genomic insertions

of the *nptII* (pROK2) and *bar* (pCSA110) vector sequences in the SALK and SAIL lines,

respectively (Fig. 3c,d). The *nptII* gene was highly targeted by 21/22nt short interfering

(si)RNAs, as well as RdDM promoting 24nt siRNAs. The NOSp and duplicated NOSt were

equally targeted by 22nt and 24nt siRNAs, although at lower numbers than the *nptII* gene itself

(Fig. 3e). In contrast, the SAIL_232 *bar* gene and its promoter were targeted by high numbers of

21nt siRNAs, and lacking 24-mers (Fig. 3f). Interestingly, the pCSA110 encoded pollen specific

promoter pLAT52, promoting the reporter gene *GUS*, was the only element highly targeted by

24nt siRNAs as well as high cytosine methylation levels in all sequence contexts (CG, CHG and

CHH methylation, where H is A, C or T) in the SAIL background (Fig. 3d, f; Supplementary

Table 6). In contrast, the entire pROK2 T-DNA region showed high cytosine methylation in all

three contexts. We found all elements to be fully CG and CHG methylated, and between 30%

(*nptII*) and 71% (NOSt) of CHH's were methylated (Supplementary Table 7). In summary, the

GUS reporter in pCSA110 and all pROK2 transgenic elements are targeted by RdDM gene

silencing pathway 24nt siRNAs. In contrast, the SAIL_232 pCSA110 *bar* gene was only targeted

by 21/22nt siRNAs without any apparent effects on expression or the chemical resistance

phenotype.

We were curious whether the 24nt siRNA and DNA methylation of the *GUS* gene

observed in leaf tissue suppresses expression in pollen, where it is driven by the strong pLAT52

promoter. Indeed staining of mature pollen identified GUS activity in the tested SAIL line, thus

suggesting that these epigenetic effects can be partially overcome by when transgene

expression is driven by a strong promoter (Fig. 3g-i).

While these observations are limited to the transformation vectors, we specifically looked

at the individual junctions between genome and T-DNA to examine for epigenetic effects on the

flanking genomic DNA sequences/genes. We only found few siRNA reads at two SALK_059379

(of eight) and six SAIL_232 (of 11), and thus were not able to draw any conclusions. Similarly

for bisulfite-converted DNA reads, where we were able to identify DNA methylation signatures at

only one junction in each transgenic line. With the caveat that only a few cases were examined,

we were unable to observe any signatures for siRNA or DNA methylation spreading outside of

the T-DNA borders.

T-DNA-insertions shape the local chromatin environment

Besides DNA methylation, modifications of histone tails as well as histone variants are

also key epigenome features playing crucial roles in instructing gene expression dynamics [36].

To study the role of T-DNA insertions in shaping the adjacent chromatin in SALK_059379 and

SAIL_232 seedlings, we employed ChIP-seq assays to profile the occupancy of two histone

modifications   (H3K4me3,   H3K27me3)   and   the   histone   variant   H2A.Z

(http://neomorph.salk.edu/T-DNA_ChIP.php). H3K4me3 marks active or poised genes [37],

whereas H3K27me3 is involved in Polycomb Repressive Complex 2 (PRC2)-mediated gene

silencing [38]. The histone variant H2A.Z is known to be involved in the gene responsiveness to

environmental stimuli in *Arabidopsis* [39]. SALK_059379 carries five T-DNA-insertions, two of

which are homozygous and could thus be analyzed and visualized in our ChIP-seq data (Fig. 4a

and Supplementary Fig. 5a-c). The SALK:chr2_18Mb T-DNA insertion led to a 5.5 kb deletion

ranging from the promoter of *At2g45820* to the gene body of *At2g45840* (Fig. 1d, 4a). We

observed that the H3K4me3 domain at *At2g45820* (domain 1) is affected by the deletion,

displaying lower levels of H3K4me3 (Fig. 4a and Supplementary Fig. 5a). In Col-0, this deleted

fragment shows a large H3K27me3 domain which extends into the gene body of *At2g45840*.

Levels of H3K27me3 are also reduced at that region (domain 2) in SALK_059379 (Fig. 4a and

Supplementary Fig. 5a). The SALK:chr2_15Mb insertion occurred in the first intron of

At2g36290, and leads to a strong reduction of H3K4me3 and H2A.Z compared to Col-0

(Supplementary Fig. 5b, c).

For SAIL_232, we began analysis of the chromatin environment at the Col-3 specific

large inversion on chromosome 1. The most profound impact at the flanking sites was found on

the level of H3K27me3, where a large H3K27me3 domain, that spans *At1g32420* and

At1g32430 in Col-0 seedlings, was split and yet still maintained at the new inverted

chromosomal positions in Col-3 (Supplementary Fig. 5d, e). As a result, we discovered a new

H3K27me3 domain (domain 2) adjacent to *At1g33700* that is likely caused by spreading of

H3K27me3 from *At1g32420* into the inverted genomic region (Supplementary Fig. 5d, e). In

contrast, the H3K27me3 domain at the gene body of *At1g32430* was detached from the larger

H3K27me3 domain and decreased in size compared to Col-0, possibly due to the loss of

H3K27me3 reinforcement (Supplementary Fig. 5d, e). The T-DNA insertion in SAIL:chr5_15Mb

occurred in the first exon of *At5g38850,* and changed the H3K4me3 pattern in this genomic area

(Supplementary Fig. 5f, g). Levels of H3K4me3 were clearly lower at *At5g38850* and started to

spread into the adjacent *At5g38840* gene (Supplementary Fig. 5f, g).

To further our understanding of how the insertion of T-DNA affects the adjacent

chromatin environment, we expanded our analysis to include eight additional T-DNA integration

events (http://neomorph.salk.edu/T-DNA_ChIP.php), previously discovered in five T-DNA lines

(SALK_061267, SALK_017723, SALK_069537, 50B_HR80 and SALK_117411)

(Supplementary Table 8). Initially, we focused on two single exon genes: *At1g32640*

(SALK_061267) and *At5g11930* (SALK_017723). In Col-0 WT *At1g32640* carries H3K4me3 and

H2A.Z over the entire gene body whereas levels of H3K27me3 were comparably low

(Supplementary Fig. 5h, i). The T-DNA insertion at *At1g32640* disrupts the H3K4me3 domain,

leading to an overall reduction, and also disrupts the H2A.Z domain resulting in a clear increase,

where especially the first domain shows a 3-fold increase (Supplementary Fig. 5h, i). At an

additional T-DNA insertion site, 865 bp outside of the gene body of *At1g23910*, we observed a

decrease in all three gene-body localized marks (Supplementary Fig. 5j, k). We discovered an

even stronger impact at *At5g11930* in SALK_017723 where the gene body T-DNA insertion

caused a complete loss all three profiled histone modifications/variants (Supplementary Fig. 6a,

b).

Next, we analyzed T-DNA insertions in multi-exon genes. Contrary to *At1g32640* and

At5g11930 where H3K4me3 and H2A.Z can be detected over the entire gene body in Col-0

seedlings (Supplementary Fig. 5h, 6a), chromatin marks are restricted to the +1 and +2

nucleosome region at *At5g23110* (SALK_069537) (Fig. 4b). Strikingly, SALK_069537 seedlings

display a new H3K4me3 and H2A.Z domain downstream of the T-DNA insertion in *At5g23110*

which is not present in Col-0 seedlings (Fig. 4b, c). This is the first evidence of a T-DNA-induced

de novo tri-methylation of H3K4 and incorporation of H2A.Z in the host genome.

In order to examine whether this pattern was specific to *At5g23110*, we inspected

another T-DNA line with a T-DNA insertion in *At3g57300*, another large multi-exon gene

(Supplementary Fig. 6c, d). This 50B_HR80 T-DNA line is derived from a T-DNA mutant-based

screen for defects in homologous recombination [40]. Similar to SALK_061267, 50B_HR80 also

shows the *de novo* H3K4 trimethylation and H2A.Z incorporation adjacent to the T-DNA

insertion site (Supplementary Fig. 6c, d), suggesting that a more general mechanism may takes

place at T-DNA insertions within larger genes.

In addition, we also found evidence of *de novo* trimethylation of H3K27 next to a T-DNA

insertion site in the gene body of *At3g53680* in SALK_117411 seedlings (Fig. 4d, e). This new

H3K27me3 domain expands into the promoter region of *At3g53680* and affects H3K4me3 and

H2A.Z only slightly (Fig. 4d, e). SALK_117411 carries another T-DNA in the 3'UTR of

At1g01700 which has a very selective impact on the chromatin environment at *At1g01700*

reducing only H3K27me3 whereas H3K4me3 and H2A.Z remained unaffected (Supplementary

Fig. 6e, f). Another 3'UTR-located T-DNA insertion in *At3g46650* (Supplementary Fig. 6g) also

impacted the adjacent chromatin in SALK_069537 but this time H3K4me3 and H2A.Z and not

H3K27me3 were profoundly affected (Supplementary Fig. 6g, h). Taken together, our ChIP-seq

assays clearly demonstrate that T-DNA integration affects the local chromatin environment at

the site of insertion.

Discussion

During the process of Agrobacterium transformation, T-DNA can insert itself into the

plant genome at random or induced (e.g. through nuclease activity) sites of chromosomal

double-strand breaks, utilizing host DNA repair mechanisms. The ideal transgene delivery

system would result in the gene of interest being integrated as a single functional copy.

Unfortunately, the majority of T-DNA transformed plants deviate from this ideal (reviewed by

Gelvin [5]), containing only partially integrated or concatemers of T-DNA that often are not

expressed (silenced) (e.g. Gelvin [10], Peach and Velten [41]). The genome structure at sites of

T-DNA insertion or the structure of the T-DNA insertion itself can additionally induce epigenetic

alterations with detrimental effects on transgene function. To better understand these effects,

short-read sequencing technologies have been employed [42]. Unfortunately, the repetitiveness

of concatenated T-DNAs (carrying transgene sequences) and vector backbone insertions [13]

has hampered efforts to fully appreciate the complexity of these events and their impact on the

plant genome and epigenome.

Using a combination of long-read sequencing/mapping tools, we identified and analyzed

perturbations to the genome structure of four randomly selected transgenic, floral dip

transformed *Arabidopsis thaliana* (Columbia accession) T-DNA insertion lines from three of the

most widely utilized plant mutant collections (SALK, SAIL, WISC). Optical genome physical

maps (BNG) created from single DNA molecules with an average size of up to 288 kb were

critical to unveil both the size and structure of genomic transgene insertions which ranged in

size up to 236 kb in length. Using nanopore long read DNA sequencing technology we were

subsequently able to assemble contigs up to chromosome arm length, which closed 83% of

non-centromeric (26% of all) misassemblies within the gold-standard reference genome

(TAIR10) [43]. Sequence contigs for the two transgenic lines (SAIL_232 and SALK_059379)

captured up to 39 kb of assembled T-DNA insertion sequence and revealed the complexity of

Agrobacterium-mediated transgene insertions. Less repetitive rearrangements, like the SAIL-

specific 512 kb inversion on chromosome 1, were perfectly captured within single chromosome

arm spanning contigs. Although ONT long read sequences were not sufficient to provide

complete contiguity of the highly repetitive T-DNA insertions, BNG maps preserved contiguity

over all insertions and rearrangements, and thus provided absolute proof for these events - in

contrast to e.g. split-read mapping of short Illumina reads [42].

We hypothesize that the final T-DNA insertion derives from two independent T-DNA

insertions at each end of the genomic breakpoint, subsequent connection into the observed long

T-DNA concatemers, and possible interaction among the many resulting homologous regions.

Previously these long repetitive segments were attributed to the floral dip transformation method

Arabidopsis [22]. Long homologous stretches between the independently inserted T-DNA/vector

backbone concatemers represent inverted repeats which can lead to hairpin formation, which

stalls the recombination fork and excision during DNA repair [45]. We identified a single

chimeric vector/vector insertion (SALK:chr2_15Mb) that most likely occurred through homology-

dependent recombination of the two identical NOSt sequences in the T-DNA vector

(pROK2:6778-7038 bp and 8607-8867 bp). While internal breakpoints and the scrambled

insertion pattern that we observed are in support of this, we have no data to confirm whether

this happened before, during, or after integration (Fig. 1, 2, Supplementary Fig. 3). Further

experiments are necessary to characterize the mechanism of T-DNA strand concatenation, the

role of floral dip and whether internal rearrangements and excisions occur during the insertion

process or at later meiotic stages.

The relationship between *Agrobacterium*-mediated T-DNA integration and the host

epigenome are largely uncharted territory. Reports and anecdotal experiences suggested

silencing of the kanamycin antibiotic resistance *nptII* marker gene in a large number of SALK

lines [4]. In contrast silencing of the SAIL line inherent herbicide tolerance *bar* gene has not

been reported. Our analyses revealed distinct epigenomic features for the two phenotypically

indistinct transgenic lines; both transgenes are targeted for sRNA-level silencing. For pROK2,

the sRNAs have progressed to the RdDM phase, and are targeting the entire transgene for

methylation and functional shut-down of the resistance gene. It appears that the antibiotic

resistance marker *nptII* is undergoing a mixture of expression-dependent silencing and

conventional Pol IV-RdDM. It lacks heavy 24mers at the promoter and makes 21-22mers within

the coding region. These even levels of 21-22 and 24nt siRNAs further suggest that the *nptII*

gene is transcribed at least at some level but degraded very efficiently and targeted for RdDM.

This transgene is a heavy target of some type of RdDM (equal levels of CHH and CG) and

produces no steady state mRNA, confirming the antibiotic sensitive phenotype.

For pCSA110, besides the Lat52 promoter, the *bar* transcript is targeted by RNAi, likely

RDR6-dependent since 21mers are derived from both strands. The few observed 24nt siRNA

traces are potentially degradation products from the highly expressed *bar* gene. However, the

siRNAs are not successfully targeting RdDM, most likely due to absence of Pol V expression, so

there is likely a reduced level of steady state mRNA, but enough to make a protein and provide

the observed herbicide tolerance. It is not surprising that the pollen LAT52 promoter is heavily

targeted by Pol IV-RdDM. This tomato-derived promoter is made up of retrotransposon

sequences that show sufficient sequence conservation among plant species, that this promoter

is always recognized and targeted by Pol IV-RdDM. We were able to show that this strong

promoter can override the silencing machinery to drive *GUS* expression in mature pollen.

Our ChIP-seq-based analyses of H3K4me3, H3K27me3 and H2A.Z occupancy

surrounding ten T-DNA insertion sites and one deletion site revealed a strong impact of T-DNA

insertion on the local chromatin environment. The most profound effects were found at T-DNA

insertions in large multi-exonic genes which gave rise to completely new H3K4me3/H2A.Z

domains. From other systems such as yeast or humans it is known that the histone variant

H2A.Z as well as the histone modification H3K4me3 are involved in the DNA damage response,

both rapidly accumulating at double strand breaks [51, 52]. Interestingly the T-DNA integration

in SALK_069537 and 50B_80HR T-DNA lines results in only one new domain at the 3' side of

the gene, clearly arguing against a DNA damage response scenario, suggesting that the

direction of transcription spatially determines the T-DNA insertion-facilitated *de novo* histone

domain formation. Moreover, we also discovered a T-DNA insertion-induced *de novo*

trimethylation of H3K27 spanning from the site of insertion into the promoter. The asymmetric

shape of the domain with its gradual increase towards the promoter suggests that it is caused

by the *de novo* recruitment of PRC2 and/or the blocking of H3K27me3 demethylases such as

REF6 [50] in that region and not by PRC2-mediated spreading of H3K27me3 from the T-DNA.

Our chromatin study identified a highly diverse and complex impact of T-DNA insertions on the

adjacent chromatin environment. Future studies will need to be conducted to analyze how T-

DNA insertions shape the local host epigenome.

Its small genome size and the widespread utilization of the T-DNA mutant collections

made *Arabidopsis thaliana* an ideal organism to study the structural and transgenerational

effects of transgene insertions. Our findings pave the way for structural genomic studies of

transgenic crop plants and provide insights into the effects of transgene insertions on the

epigenetic landscape.

Material and Methods

Plant Material

Seeds were ordered from ABRC (https://abrc.osu.edu): SALK_059379 (segregant and homozygous lines), SALK_075892, SAIL_232 (seg.), SAIL_59 (seg.), SAIL_107 (seg.), WiscDsLox_449D11 (seg.), SALK_061267, SALK_017723, SALK_069537 and SALK_117411. 50B_HR80 seeds were kindly provided from Susan M. Gasser. Plants were grown in a 20□ growth room with 13h light/11h dark cycles in peat-based soil supplemented with nutrients, and collected approximately three weeks after bolting. Growth conditions for plant material that was used for ChIP-seq is listed in Supplementary Table 8.

Testing Selection Markers

Kanamycin resistant lines- SALK and Wisc

MS/MES media was prepared (1L NP H2O, 2.2g MS, 0.5g MES, pH5.7) and autoclave sterilized. Kanamycin (50ug/mL) was added to half of the media, and plates with and without the antibiotic were poured. Seeds from the lines SALK_059379 (homozygous), SAIL_232, WiscDsLox_449D11, and Col-0 CS70000 control were sterilized and spotted on the plates, which were placed in the same growth conditions as the plant material in soil. Survival rates were measured 5 days after seeds were spotted and growth was observed for further 14 days.

Finale™ *resistant lines- SAIL*

One flat each of Col-0 (control) and SAIL_232 were grown in soil as outlined in *Plant Material*. Plants were sprayed with Finale™ at three different concentrations (5mg/L, 10mg/L, 20mg/L) at 5, 14, and 21 days after germination [1, 2].

GUS reporter staining- SALK and SAIL

Leaf and flower tissue was submerged in 1.5mL staining buffer (0.5mM ferrocyanide,

0.5mM ferricyanide, 0.5% Triton, 1mg/mL X-Gluc, 100mM Sodium Phosphate Buffer pH 7) and

incubated at 37°C overnight. A light microscope was used to visualize the staining at 10x

magnification for the leaves and 4x for the flowers.

ONT Library Prep and Sequencing

5g of flash-frozen leaf tissue, pooled from the segregant seed stocks, was ground in

liquid nitrogen and extracted with 20mL CTAB/Carlson lysis buffer (100mM Tris-HCl, 2% CTAB,

1.4M NaCl, 20mM EDTA, pH 8.0) containing 20µg/mL proteinase K for 20 minutes at 55□. To

purify the DNA, 0.5x volume chloroform was added, mixed by inversion, and centrifuged for 30

minutes at 3000 RCF. Purification was followed by a 1x volume 1:1 phenol: [24:1

chloroform:isoamyl alcohol] extraction. The DNA was further purified by ethanol precipitation

(1/10 volume 3M sodium acetate pH 5.3, 2.5 volumes 100% ethanol) for 30 minutes on ice. The

resulting pellet was washed with freshly-prepared ice-cold 70% ethanol, dried, and resuspended

in 350µL 1x TE buffer (10mM Tris-HCl, 1mM EDTA, pH 8.0) with 5µL RNase A (Qiagen, Hilden)

at 37□ for 30 minutes, then at 4□ overnight. The RNase A was removed by double extraction

with 24:1 chloroform:isoamyl alcohol, centrifuging at 22,600xg for 20 minutes at 4□ each time to

pellet. An ethanol precipitation was performed as before for 3 hours at 4□, washed, and

resuspended overnight in 350µL 1x TE buffer. The genomic DNA samples were purified with the

Zymogen Genomic DNA Clean and Concentrator-10 column (Zymo Research, Irvine). The

purified DNA was prepared for sequencing with the Ligation Sequencing Kit 1D (SQK-LSK108,

ONT, Oxford, UK) sequencing kit protocol. Briefly, approximately 2 µg of purified DNA was

repaired with NEBNext FFPE Repair Mix for 60 min at 20^{0}C. The DNA was purified with 0.5X

Ampure XP beads (Beckman Coulter, Brea). The repaired DNA was End Prepped with

NEBNExt Ultra II End-repair/dA tail module and purified with 0.5X Ampure XP beads. Adapter

mix (ONT, Oxford, UK) was added to the purified DNA along with Blunt/TA Ligase Master Mix

(NEB) and incubated at 20^0C for 30 min followed by 10 min at 65^0C. Ampure XP beads and

ABB wash buffer (ONT, Oxford, UK) were used to purify the library molecules, which were

recovered in Elution buffer (ONT, Oxford, UK). The purified library was combined with RBF

(ONT, Oxford, UK) and Library Loading Beads (ONT, Oxford, UK) and loaded onto a primed

R9.4 Spot-On Flow cell. Sequencing was performed with an ONT MinION Mk1B sequencer

running for 48 hrs. Resulting FAST5 (HDF5) files were base-called using the ONT Albacore

software (0.8.4) for the SQK-LSK108 library type.

Sequence extraction, assembly, consensus and correction

Raw ONT reads (fastq) were extracted from base-called FAST5 files using poretools [3].

Overlaps were generated using minimap [4] with the recommended parameters (-Sw5 -L100 -

m0). Genome assembly graphs (GFA) were generated using miniasm [4]. Unitig sequences

were extracted from GFA files. Three rounds of consensus correction was performed using

Racon [5] based on minimap overlaps, and the resulting assembly was polished using Illumina

PCR-free 2x250 bp reads mapped with bwa [6] and pilon [7]. Genome statistics were generated

using QUAST [8].

The ONT reads were also assembled with the CANU assembler [9]. However, the

minimap/miniasm assemblies were of higher contiguity and resolved longer stretches of the T-

DNA insertions. In addition, the minimap/miniasm assemblies were computed within 4-6 hours,

while the CANU assemblies took between one (Col-0) and four weeks (SAIL). We hypothesize

that the SAIL lines inherently complex repeat structure caused the long compute time, while the

reference Col-0 required only the expected time of one week. Also, we hypothesize that

minimap/miniasm resolved the T-DNA structures more fully due to the fact it does not have a

read correction step, which could lead to the collapsing of highly repetitive yet distinct T-DNA

insertions.

Identification of individual T-DNA matching reads and annotation

Reads obtained via ONT sequencing were imported into Geneious R10 [10], along with

the pROK2, and pCSA110 plasmid sequences. BLAST databases, were created for the

complete set of SALK_059379 and SAIL_232 ONT reads, and their respective plasmid

sequences were BLASTn searched against these databases. Reads with hits on the plasmid

sequences were extracted, and BLASTn searched against the TAIR10 reference. NCBI

megaBLAST [11] and Geneious 'map to reference' functions were used to annotate regions on

each read corresponding to either TAIR10 or the plasmid sequence. The chromosomal regions

were compared to the regions identified by T-DNA seq and BNG mapping to verify and refine T-

DNA insertion start and stop coordinates.

BNG Optical Genome Mapping

High molecular weight (HMW) DNA for BNG mapping and ONT sequencing was

extracted as outlined in Kawakatsu [12]. Briefly, up to 5g of fresh mixed leaf and flower tissue

(excluding chlorotic leaves and stems) pooled from the segregant seed stocks were

homogenized in 50mL nuclei isolation buffer. Nuclei were separated from debris using a Percoll

layer. Extracted nuclei were subsequently embedded in low melting agarose plugs and exposed

to lysis buffer overnight. DNA was released by digesting the agarose with Agarase enzyme

(Thermo Fisher Scientific).

HMW DNA was nicked with the enzyme Nt.BspQI (New England Biolabs, Ipswich, MA),

fluorescently labeled, repaired and stained overnight following the Bionano Genomics nick-

labeling protocol and accompanying reagents (Bionano Genomics, San Diego, CA) [13]. Each

Arabidopsis T-DNA insertion line was run for up to 120 cycles on a single flow cell on the Irys

platform (Bionano Genomics, San Diego, CA). Collected data was filtered (SNR = 2.75; min

length 100kb) using IrysView 2.5.1 software, and assembled using default "small" parameters.

Average molecule length for assembly varied between 201 kb and 288 kb, and resulted in BNG

map N50 0.97 to 1.03 Mb. Derived assembled maps were anchored to converted TAIR10

chromosomes using the RefAligner tool and standardized parameters (-maxthreads 32 -output-

veto-filter _intervals.txt$ -res 2.9 -FP 0.6 -FN 0.06 -sf 0.20 -sd 0.0 -sr 0.01 -extend 1 -outlier

0.0001 -endoutlier 0.01 -PVendoutlier -deltaX 12 -deltaY 12 -xmapchim 12 -hashgen 5 7 2.4 1.5

0.05 5.0 1 1 1 -hash -hashdelta 50 -mres 1e-3 -hashMultiMatch 100 -insertThreads 4 -nosplit 2 -

biaswt 0 -T 1e-10 -S -1000 -indel -PVres 2 -rres 0.9 -MaxSE 0.5 -HSDrange 1.0 -outlierBC -

xmapUnique 12 -AlignRes 2 -outlierExtend 12 24 -Kmax 12 -f -maxmem 128 -stdout -stderr).

We utilized the resulting *.xmap, *_q.cmap and *_r.cmap files in the structomeIndel.py script

(https://github.com/RyanONeil/structome) to identify T-DNA insertion locations and sizes. We

used this script to further determine misassemblies in the ONT genome assemblies. Known Col-

0 mis-assemblies [12] were subtracted from the list of derived locations.

Aligning contigs to TAIR10

We aligned ONT contigs to the TAIR10 reference using the BioNano Genomics

RefAligner as above (Supplementary Table 3). ONT contigs were in silico digested with

Nt.BspQI and used as template in RefAligner. The alignment output was manually derived from

the .xmap output file.

Genotyping of structural genome variations

Primer3 was used to create oligos (Supplementary Table 5) for each insertion identified

in the SALK_059379 and SAIL_232 lines. The forward primers correspond to the chromosome

sequence ~500bp before the insertion start site, and the reverse to the plasmid sequence

~500bp after the insertion start site. A second set of reverse primers corresponding to the

original WT chromosome ~1kb after the original forward primers. Individuals and pooled tissue

from SALK_059379, SAIL_232, SAIL_59, SAIL_107 and Col_0 CS7000 DNA was extracted

using Qiagen DNeasy Plant Mini kit (following the protocol) and genotyped using these primers,

with Col-0 as the negative control.

Bisulfite Sequencing

DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Hilden, D) according to the

manufacturer's protocol from segregant plant pools, and quantified using the Qubit dsDNA BR

assay kit. Illumina sequencing library preparation and bisulfite conversion was conducted as

described in Kawakatsu [12]. Briefly, DNA was End Repaired using the End-It kit (Epicentre,

Madison, WI), A-tailed using dA-Tailing buffer and 3µL Klenow (3' to 5' exo minus) (NEB,

Ipswich, MA) and Truseq Indexed Adapters (Illumina, San Diego, CA) were ligated overnight.

Bead purified DNA was quantified using the Qubit dsDNA BR assay kit, and stored at -20□. At

least 450ng of adapter ligated DNA was taken into bisulfite conversion which was performed

according to the protocol provided with the MethylCode Bisulfite Conversion kit (Thermo Fisher,

Waltham, MA). Cleaned, converted single stranded DNA was amplified by PCR using the KAPA

U+ 2x Readymix (Roche Holding AG, Basel, CH): 2 min at 95□, 30 sec at 98□ [15 sec at 98□,

30 sec at 60□, and 4 min at 72□] x 9, 10 min at 72□, hold at 4□. After amplification, the DNA

was bead purified, the concentration was assessed using the Qubit dsDNA BR assay kit, and

the samples were stored at -20□. WGBS library was sequenced as part of a large multiplexed

pool paired-end 150 bp on an Illumina HiSeq 4000.

RNA Libraries

RNA extractions performed using the RNeasy Plant Mini Kit (Qiagen, Hilden, D) from

segregant plant pools. RNA-seq libraries were prepared manually (SAIL_232) as described in

Kawakatsu or using the Illumina NeoPrep Library Prep System (SALK_059379) (Illumina, San

Diego, CA), following the NeoPrep control software protocol. The completed libraries were

quantified using the Qubit dsDNA HS assay kit and stored at -20□. SALK_059379 was

sequenced in duplicate as single end 150 bp in a multiplexed pool on two lanes of an Illumina

HiSeq 2500. SAIL_232 was sequenced as part of a large multiplexed pool paired-end 150 bp on

an Illumina HiSeq 4000.

small RNA Libraries

We have extracted sRNA according to the protocol described in Vandivier [14] with

modifications to the RNA extraction and size selection steps. In brief, 300mg flash-frozen leaf

tissue from segregant plant pools was ground in liquid nitrogen and extracted with 700µL QIAzol

lysis reagent (Qiagen, Hilden, D). The RNA was separated from the lysate using QIAshredder

columns (Qiagen, Hilden, D) and purified with a □ volume 24:1 chloroform:isoamyl alcohol

(Sigma-Aldrich, St. Louis, MO) extraction followed by a 1mL wash of the aqueous phase with

100% ethanol. The RNA was further purified using miRNeasy columns (Qiagen, Hilden, D) with

two 82µL DEPC-treated water washes. 37µL DNase was added to the flow-through and

incubated at RT for 25 minutes. The DNase-treated RNA was precipitated with 20µL 3M sodium

acetate (pH 5.5) and 750µL 100% ethanol overnight at -80□. The pellet was washed with 750µL

ice-cold 80% ethanol and resuspended in a 12:1 DEPC-treated water:RNase OUT Recombinant

Ribonuclease Inhibitor (Thermo Fisher, Waltham, MA) mixture on ice for 30 minutes and

quantified (Qubit RNA HS assay kit).

Size selection and library prep were conducted exactly as described in Vandivier [14].

The RNA from 15-35bp was cut out of the gel and purified. sRNA libraries were sequenced in

duplicate as single end 150 bp in a multiplexed pool on two lanes of an Illumina HiSeq 2500.

Short read analysis

RNA-seq and sRNA Illumina reads were adapter trimmed and mapped against the

junction sequences as well as the corresponding plasmid sequences, using Bowtie2 and

Geneious aligner (Geneious R10.2.3; custom sensitivity, iterate up to 5 times, zero mismatches

or gaps per read, word length 14) at highest stringency. RNA-seq reads were mapped against

AtACT1 as control, and found this gene expressed. RNA-seq reads were mapped against Pol V

AT2G40030 as control, and found this gene not expressed. Resulting reads were extracted and

re-mapped against the TAIR10 reference genome to ensure that no off-target reads mapped to

the plasmid sequences. Because we analyzed various genomic insertions against a single

reference, we did not normalize read mapping.

The MethylC-Seq data (paired-end) reads in FastQ format and the second pair reads

were converted to their reverse complementary strand nucleotides. Then reads were aligned to

the *Arabidopsis thaliana* reference genome (araport 11) and pCSA110/pROK2 genome. The

Chloroplast genome was used for quality control (< 0.35% non-conversion rate). After mapping,

overlapping bases in the paired-end reads were trimmed. The base calls per reference position

on each strand were used to identify methylated cytosines at 1% FDR.

ChIP-seq assays and analysis

ChIP-seq experiments were performed as previously described [15] using antibodies against

H3K4me3 (04-745, EMD Millipore), H2A.Z (39647, Active Motif) and H3K27me3 (39156, Active

Motif). Incubation with mouse IgG (015-000-003, Jackson ImmunoResearch) served as our

negative control. ChIP DNA was used to generate sequencing libraries following the

manufacturer's instructions (Illumina). Libraries were sequenced on the Illumina Genome

Analyzer II. Sequencing reads were aligned to the TAIR10 genome assembly using Bowtie2

[16]. Histone domains next to the T-DNA insertion events were identified with SICER [17]. The

bigWigAverageOverBed tool executable from the UCSC genome browser [18] was used to

quantify the occupancy of H3K4me3, H2A.Z and H3K27me3 next to T-DNA integration sites.

The TAIR10 reference domain coordinates can be found in Supplementary Table 9.

Data Availability

Raw ONT sequencing data was deposited in the European Nucleotide Archive (ENA)

under project PRJEB23977 (ERP105765). Final polished assemblies were deposited in the

ENA Genome Assembly Database PRJEB23977. Raw BNG molecules and assembled maps

are deposited under BioProjects PRJNA387199, PRJNA387199, PRJNA387199 and

PRJNA387199. Short-read datasets are deposited under GEO accession GSE108401.

Author contributions

FJ, TPM, JRE conceptualized the study; AR, FJ, STM, MZ, RC, JN and JS created

sequencing libraries and/or conducted sequencing; FJ, AR, HC, MZ and TPM analyzed

sequencing data; AR, JS and FJ conducted plant studies; MZ conceptualized and conducted

ChIP-seq experiments; FJ, AR, TPM, RKS, MZ and JRE wrote the manuscript.

Acknowledgments

We would like to thank Cesar Barragan and Dr. Ronan O'Malley for insights into the T-

DNA project, and Bruce Jow and Christopher Santos for excellent greenhouse support. We

thank Detlef Weigel and Christa Lanz, both Max Planck Institute for Developmental Biology

(Tuebingen, Germany) for performing and providing Illumina short read sequencing for Col-0

CS70000. FJ was supported through a Human Frontier Science Program Organization long-

term fellowship. M.Z. was supported by the Salk Pioneer Postdoctoral Endowment Fund as well

as by a Deutsche Forschungsgemeinschaft (DFG) research fellowship (Za-730/1-1). JRE is an

Investigator of the Howard Hughes Medical Institute.

Conflict of Interests

FJ is an employee of Bayer Crop Science. JRE serves on the scientific advisory boards of

Cibus, Zymo Research and Pathway Genomics Inc.

References

[1] Baulcombe DC, Saunders GR, Bevan MW, Mayo MA, Harrison BD. Expression of

biologically active viral satellite RNA from the nuclear genome of transformed plants. Nature.

1986 May 22;321(6068):446–9.

[2] Caplan A, Herrera-Estrella L, Inzé D, Van Haute E, Van Montagu M, Schell J, et al.

Introduction of genetic material into plant cells. Science. 1983 Nov 18;222(4625):815–21.

[3] Fraley RT, Rogers SG, Horsch RB, Sanders PR, Flick JS, Adams SP, et al. Expression of

bacterial genes in plant cells. Proc Natl Acad Sci USA. 1983 Aug;80(15):4803–7.

[4] O'Malley RC, Ecker JR. Linking genotype to phenotype using the Arabidopsis unimutant

collection. Plant J. 2010 Mar;61(6):928–40.

[5] Gelvin SB. Agrobacterium-mediated plant transformation: the biology behind the "gene-

jockeying" tool. Microbiol Mol Biol Rev. 2003 Mar;67(1):16–37, table of contents.

[6] Zambryski P, Depicker A, Kruger K, Goodman HM. Tumor induction by Agrobacterium

tumefaciens: analysis of the boundaries of T-DNA. J Mol Appl Genet. 1982;1(4):361–70.

[7] Nester EW. Agrobacterium: nature's genetic engineer. Front Plant Sci. 2014;5:730.

[8] Van Kregten M, de Pater S, Romeijn R, van Schendel R, Hooykaas PJJ, Tijsterman M. T-

DNA integration in plants results from polymerase-θ-mediated DNA repair. Nat Plants. 2016 Oct

31;2(11):16164.

[9] Zelensky AN, Schimmel J, Kool H, Kanaar R, Tijsterman M. Inactivation of Pol θ and C-

NHEJ eliminates off-target integration of exogenous DNA. Nat Commun. 2017 Jul 7;8(1):66.

[10] Gelvin SB. Integration of Agrobacterium T-DNA into the Plant Genome. Annu Rev Genet.

2017 Nov 27;51:195–217.

[11] Clark KA, Krysan PJ. Chromosomal translocations are a common phenomenon in

Arabidopsis thaliana T-DNA insertion lines. Plant J. 2010 Dec;64(6):990–1001.

[12] Feldmann KA. T-DNA insertion mutagenesis in Arabidopsis: mutational spectrum. Plant J.

1991 Jul;1(1):71–82.

[13] Jorgensen R, Snyder C, Jones JDG. T-DNA is organized predominantly in inverted repeat

structures in plants transformed with Agrobacterium tumefaciens C58 derivatives. Mol Gen

Genet. 1987 May;207(2-3):471–7.

[14] Ming R, Hou S, Feng Y, Yu Q, Dionne-Laporte A, Saw JH, et al. The draft genome of the

transgenic tropical fruit tree papaya (Carica papaya Linnaeus). Nature. 2008 Apr

24;452(7190):991–6.

[15] Nacry P, Camilleri C, Courtial B, Caboche M, Bouchez D. Major chromosomal

rearrangements induced by T-DNA transformation in Arabidopsis. Genetics. 1998

Jun;149(2):641–50.

[16] Ooms G, Bakker A, Molendijk L, Wullems GJ, Gordon MP, Nester EW, et al. T-DNA

organization in homogeneous and heterogeneous octopine-type crown gall tissues of Nicotiana

tabacum. Cell. 1982 Sep;30(2):589–97.

[17] Thomashow MF, Nutter R, Montoya AL, Gordon MP, Nester EW. Integration and

organization of Ti plasmid sequences in crown gall tumors. Cell. 1980 Mar;19(3):729–39.

[18] Ulker B, Peiter E, Dixon DP, Moffat C, Capper R, Bouché N, et al. Getting the most out of

publicly available T-DNA insertion lines. Plant J. 2008 Nov;56(4):665–77.

[19] Windels P, De Buck S, Van Bockstaele E, De Loose M, Depicker A. T-DNA integration in

Arabidopsis chromosomes. Presence and origin of filler DNA sequences. Plant Physiol. 2003

Dec;133(4):2061–8.

[20] Woody ST, Austin-Phillips S, Amasino RM, Krysan PJ. The WiscDsLox T-DNA collection:

an arabidopsis community resource generated by using an improved high-throughput T-DNA

sequencing pipeline. J Plant Res. 2007 Jan;120(1):157–65.

[21] Zhu Q-H, Ramm K, Eamens AL, Dennis ES, Upadhyaya NM. Transgene structures suggest

that multiple mechanisms are involved in T-DNA integration in plants. Plant Sci. 2006

Sep;171(3):308–22.

[22] De Buck S, Podevin N, Nolf J, Jacobs A, Depicker A. The T-DNA integration pattern in

Arabidopsis transformants is highly determined by the transformed target cell. Plant J. 2009

Oct;60(1):134–45.

[23] Collier R, Dasgupta K, Xing Y-P, Hernandez BT, Shao M, Rohozinski D, et al. Accurate

measurement of transgene copy number in crop plants using droplet digital PCR. Plant J. 2017

Jun;90(5):1014–25.

[24] Michael TP, Jupe F, Bemm F, Motley ST, Sandoval JP, Lanz C, et al. High contiguity

Arabidopsis thaliana genome assembly with a single nanopore flow cell. Nat Commun. 2018

Feb 7;9(1):541.

[25] Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, et al. Genome-wide

insertional mutagenesis of *Arabidopsis thaliana*. Science. 2003 Aug 1;301(5633):653–7.

[26] McElver J, Tzafrir I, Aux G, Rogers R, Ashby C, Smith K, et al. Insertional mutagenesis of

genes required for seed development in Arabidopsis thaliana. Genetics. 2001 Dec;159(4):1751–

63.

[27] Sessions A, Burke E, Presting G, Aux G, McElver J, Patton D, et al. A high-throughput

Arabidopsis reverse genetics system. Plant Cell. 2002 Dec;14(12):2985–94.

[28] Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from

long uncorrected reads. Genome Res. 2017 Jan 18;27(5):737–46.

[29] Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. PLoS ONE. 2014 Nov 19;9(11):e112963.

[30] Kleinboelting N, Huep G, Appelhagen I, Viehoever P, Li Y, Weisshaar B. The Structural

Features of Thousands of T-DNA Insertion Sites Are Consistent with a Double-Strand Break

Repair-Based Insertion Mechanism. Mol Plant. 2015 Nov 2;8(11):1651–64.

[31] Daxinger L, Hunter B, Sheikh M, Jauvion V, Gasciolli V, Vaucheret H, et al. Unexpected

silencing effects from T-DNA tags in Arabidopsis. Trends Plant Sci. 2008 Jan;13(1):4–6.

[32] Gao Y, Zhao Y. Epigenetic suppression of T-DNA insertion mutants in Arabidopsis. Mol

Plant. 2013 Mar;6(2):539–45.

[33] LeClere S, Bartel B. A library of Arabidopsis 35S-cDNA lines for identifying novel mutants.

Plant Mol Biol. 2001 Aug;46(6):695–703.

[34] Nakamura S, Mano S, Tanaka Y, Ohnishi M, Nakamori C, Araki M, et al. Gateway binary

vectors with the bialaphos resistance gene, bar, as a selection marker for plant transformation.

Biosci Biotechnol Biochem. 2010 Jun 7;74(6):1315–9.

[35] Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in

plants and animals. Nat Rev Genet. 2010 Mar;11(3):204–20.

[36] Fritsch O, Benvenuto G, Bowler C, Molinier J, Hohn B. The INO80 protein controls

homologous recombination in Arabidopsis thaliana. Mol Cell. 2004 Nov 5;16(3):479–85.

[37] Coleman-Derr D, Zilberman D. Deposition of histone variant H2A.Z within gene bodies

regulates responsive genes. PLoS Genet. 2012 Oct 11;8(10):e1002988.

[38] Peach C, Velten J. Transgene expression variability (position effect) of CAT and GUS

reporter genes driven by linked divergent T-DNA promoters. Plant Mol Biol. 1991 Jul;17(1):49–

60.

[39] Schouten HJ, Vande Geest H, Papadimitriou S, Bemer M, Schaart JG, Smulders MJM, et

al. Re-sequencing transgenic plants revealed rearrangements at T-DNA inserts, and integration

of a short T-DNA fragment, but no increase of small mutations elsewhere. Plant Cell Rep. 2017

Mar;36(3):493–504.

[40] Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant

Arabidopsis thaliana. Nature. 2000 Dec 14;408(6814):796–815.

[41] Dürrenberger F, Crameri A, Hohn B, Koukolíková-Nicola Z. Covalently bound VirD2 protein

of Agrobacterium tumefaciens protects the T-DNA from exonucleolytic degradation. Proc Natl

Acad Sci USA. 1989 Dec;86(23):9154–8.

[42] Schuermann D, Fritsch O, Lucht JM, Hohn B. Replication stress leads to genome

instabilities in Arabidopsis DNA polymerase delta mutants. Plant Cell. 2009 Sep 29;21(9):2700–

14.

[43] Simpson JT, Workman RE, Zuzarte PC, David M, Dursi LJ, Timp W. Detecting DNA

cytosine methylation using nanopore sequencing. Nat Methods. 2017 Apr;14(4):407–10.

[44] Sidorenko LV, Lee T-F, Woosley A, Moskal WA, Bevan SA, Merlo PAO, et al. GC-rich

coding sequences reduce transposon-like, small RNA-mediated transgene silencing. Nat Plants.

2017 Nov;3(11):875–84.

[45] Shilo S, Tripathi P, Melamed-Bessudo C, Tzfadia O, Muth TR, Levy AA. T-DNA-genome

junctions form early after infection and are influenced by the chromatin state of the host

genome. PLoS Genet. 2017 Jul 24;13(7):e1006875.

[46] O'Malley RC, Barragan CC, Ecker JR. A user's guide to the Arabidopsis T-DNA insertion

mutant collections. Methods Mol Biol. 2015;1284:323–42.

<u>Methods References</u>

[1] LeClere S, Bartel B. A library of Arabidopsis 35S-cDNA lines for identifying novel mutants.

Plant Mol Biol. 2001 Aug;46(6):695–703.

[2] Nakamura S, Mano S, Tanaka Y, Ohnishi M, Nakamori C, Araki M, et al. Gateway binary

vectors with the bialaphos resistance gene, bar, as a selection marker for plant transformation.

Biosci Biotechnol Biochem. 2010 Jun 7;74(6):1315–9.

[3] Loman NJ, Quinlan AR. Poretools: a toolkit for analyzing nanopore sequence data.

Bioinformatics. 2014 Dec 1;30(23):3399–401.

[4] Li H. Minimap and miniasm: fast mapping and de novo assembly for noisy long sequences.

Bioinformatics. 2016 Jul 15;32(14):2103–10.

[5] Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from

long uncorrected reads. Genome Res. 2017 Jan 18;27(5):737–46.

[6] Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009 Jul 15;25(14):1754–60.

[7] Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an integrated

tool for comprehensive microbial variant detection and genome assembly improvement. PLoS

ONE. 2014 Nov 19;9(11):e112963.

[8] Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome

assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

[9] Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and

accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res.

2017 Mar 15;27(5):722–36.

[10] Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious

Basic: an integrated and extendable desktop software platform for the organization and analysis

of sequence data. Bioinformatics. 2012 Jun 15;28(12):1647–9.

[11] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J

Mol Biol. 1990 Oct 5;215(3):403–10.

[12] Kawakatsu T, Huang S-SC, Jupe F, Sasaki E, Schmitz RJ, Urich MA, et al. Epigenomic

Diversity in a Global Collection of Arabidopsis thaliana Accessions. Cell. 2016 Jul

14;166(2):492–505.

[13] Lam HYK, Clark MJ, Chen R, Chen R, Natsoulis G, O'Huallachain M, et al. Performance

comparison of whole-genome sequencing platforms. Nat Biotechnol. 2011 Dec 18;30(1):78–82.

[14] Vandivier LE, Li F, Gregory BD. High-throughput nuclease-mediated probing of RNA

secondary structure in plant transcriptomes. Methods Mol Biol. 2015;1284:41–70.

Table 1: BNG maps and ONT assemblies identified T-DNA insertions from segregant

samples. BNG maps and ONT assemblies were aligned against the TAIR10 reference genome

to detect coverage, number of insertions, and maximum insertion site. Individual ONT reads

identified further insertions or rearrangements that were absent from the segregant assemblies.

Fig. 1: SALK_059379 T-DNA insertions are T-strand and backbone conglomerations.

Schematic representation of *Arabidopsis* mutant line SALK_059379 chromosome 2 T-DNA

insertions as identified from (a, c, e) ONT sequencing *de novo* genome assemblies and (b, d)

BNG optical genome maps. (a) Graphical alignment of Col-0 and SALK_059379 ONT contigs to

TAIR10 chromosome 2; two of the three TAIR10 misassemblies (red) are resolved within a

single Col-0 ONT contig. Contig joints are represented as vertical black bars. Blue boxes

indicate a broken (chr2_15Mb) and an assembled T-DNA insertion (chr2_18Mb). (b, d)

represent the BNG maps that identify T-DNA insertions including an alignment of individual

BNG molecules. (c) ONT contigs 4 and 7 have the chr2_15Mb insertion site partially assembled

and identify a mix of direct T-DNA (blue) and pROK2 backbone (red) concatemers or scrambled

stretches with breakpoints as red bars. (e) The chr2_18Mb insertion is assembled and identifies

a 5,497 bp chromosomal deletion.

Fig. 2: T-DNA integration induced large scale rearrangements in the SAIL_232 genome.

Two T-DNA induced translocations occurred on chromosomes 3 (light blue) and 5 (lime green).

(a) Inverted translocation of a 1.1 Mb segment split at a T-DNA insertion site (blue bar) was

moved to a distal part of the same chromosome arm. The other chromosome arm was swapped

with chromosome 5 (d). (b) The ONT assembly identified the underlying T-DNA inserts (boxed

arrows) including a breakpoint (red line). (c) The chromosome swap joint site contains four

pCSA110 vector fragments (boxed arrows) interspersed with breakpoints. These insertions

change the CG signature in the respective region, and are confirmed by BNG alignment. (e)

BNG maps (blue), aligned to *Arabidopsis* reference TAIR10 (green), are able to phase the WT

(top) and T-DNA haplotypes (T-DNA; bottom) for this particular insertion site. Black lines

indicate Nt.BspQI nicking sites.

1011

Fig. 3: SALK and SAIL T-strand sequences show divergent epigenomic signatures.

Transgene activity was tested by exposing plants to the respective selective agent: (a)

SALK_059379 grown on media containing the antibiotic kanamycin (50ug/uL) (or empty control)

and (b) the SAIL_232 sprayed with the herbicide Finale™ at two different concentrations

through the middle of the tray. All phenotypes are compared to the WT Col-0 (a,b). Analysis of

expression and epigenetic signatures on the corresponding T-DNA sequence is captured in

browser shots for SALK_059379 plasmid pROK2 (c) and SAIL_232 plasmid pCSA110 (d):

Illumina read mapping of bisulfite sequencing, RNA-seq and different small RNA (sRNA)

species. Quantification of individual siRNA read length against individual parts of the two

plasmid sequences are reported in (e = pROK2) and (f = pCSA110). GUS staining of leaves and

flowers of SAIL_232 (g), with Col-0 (h) and SALK_059379 (i) as control.

1023

Fig. 4: T-DNA insertions alter the chromatin landscape

(a) AnnoJ genome browser visualization of ChIP-seq reads derived from H3K4me3, H2A.Z and

H3K27me3 occupancy in Col-0 and SALK_059379 seedlings around the T-DNA-induced

deletion on the chromosome 2. Histone domains that are adjacent to the T-DNA deletion are

indicated as domain 1 and 2. (b) Annoj genome browser visualization of H3K4me3, H2A.Z and

H3K27me3 occupancy in Col-0 and SALK_069537 seedlings around the T-DNA insertion at

At5g23110. Schematic illustration shows the gene structure of *At5g23110* and the approximate

location of the corresponding T-DNA insertion. (c) Quantification of H3K4me3, H2A.Z and

H3K27me3 around the T-DNA insertion site at *At5g23110* (SALK_069537). (d) Visualization of

H3K4me3, H2A.Z and H3K27me3 occupancy in Col-0 and SALK_117411 seedlings indicates

de novo trimethylation of H3K27 around the T-DNA insertion in *At3g53680*. Scheme of gene

structure of *At3g53680* shows the approximate location of the T-DNA insertion in genome of

SALK_117411 plants. (b) Levels of H3K4me3, H2A.Z and H3K27me3 around the T-DNA

insertion at *At3g53680* in SALK_117411 seedlings are shown. A red arrow marks the

approximate location of the T-DNA integration site in the Annoj genome browser screenshots.

The respective occupancies were identified with ChIP-seq and all shown AnnoJ genome

browser tracks were normalized to the respective sequencing depth. The Col-0 IgG track serves

as a control. The occupancy of the respective domains was calculated as the ratio between the

respective ChIP-seq sample and the Col-0 IgG control.

1043

1044

1045 **Supplementary Table 1: Bionano Genomics and Oxford Nanopore Technologies**

1046 **sequencing and assembly statistics.**

1047

1048 **Supplementary Table 2: Genome Alignment of ONT assembled contigs against TAIR10**

1049 **using the Bionano Genomics RefAligner algorithm.**

1050

1051 **Supplementary Table 3: Identification of 'N' regions in the TAIR10 reference genome and**

1052 **the corresponding Col-0 ONT contigs.**

1053

1054 **Supplementary Table 4: BNG maps identify misassembled regions in ONT contigs as**

1055 **'False Duplications' or 'False Deletions'.**

1056

1057 **Supplementary Table 5: List and analysis of genomic T-DNA insertion sites.**

1058

1059 **Supplementary Table 6: Analysis of methylated cytosines on the pCSA110 plasmid for**

1060 **SAIL_232.**

1061

1062 **Supplementary Table 7: Analysis of methylated cytosines on the pROK2 plasmid for**

1063 **SALK_059379.**

1064

1065 **Supplementary Table 8: Experimental information for conducted ChIP experiments in T-**

1066 **DNA and WT Arabidopsis lines.**

1067

1068 **Supplementary Table 9: TAIR10 reference histone domain coordinates.**

1069

1070 **Supplementary Table 10: PCR oligo sequences used for genotyping T-DNA inserts**

1071

Supplementary Fig. 1: Alignment of *de novo* assembled ONT contigs for Col-0 (CS70000) and

the transgenic lines SALK_059379 and SAIL_232 relative to the reference TAIR10.

Uninterrupted colored blocks indicate contig length, black bars indicate the start and end of

contigs, black boxes indicate centromere gaps. Arrowheads represent T-DNA insertion sites;

orange lines and boxes indicate sites of translocations. Drawn to scale for each chromosome

individually.

1078

Supplementary Fig. 2: Background inversion in SAIL lines derived from Col-3. Two

additional SAIL lines were chosen at random and genotyped along with SAIL_232. The primers

were designed to amplify 1000bp spanning the junction between the genomic and inverted DNA

sequence (based on *de novo* reads from ONT sequencing). The inversion on chromosome 1

was seen in all SAIL lines tested, unlike the other chromosomal architecture disruptions found.

Thus, it can be concluded that this inversion is part of the Col-3 background from which SAIL

lines are derived. The bands at 1500bp in the SAIL_107 and SAIL_59 lines are off-target

amplification products.

1087

Supplementary Fig. 3: Model for microhomology dependent excision of T-DNA/genome

fragments. (a) A double strand chromosome break leads to (b) two multi-copy T-DNA strand

insertions in opposing directions. The exact structure and length of the inserted sequence is

unknown, as indicated by question marks. (c) SALK_chr2:18Mb insertion features two individual

double strand breaks, around 5 kb apart. High homology between the T-DNA strands as well as

the hairpin forming original DNA piece created a secondary structure (d), that was potentially

excised (e) and resulted in the deletion of the ~5kb chromosomal fragment, as shown in main

text Fig. 2. Arrowheads on the red T-DNA strand show direction. The blue line represents

double stranded DNA.

1097

Supplementary Fig. 4: ONT reads identify insertions with lower allele frequency that are

not part of assembled contigs. We applied blastn searches of all unassembled ONT reads

against the utilized vector sequence, and subsequently the TAIR10 reference genome. This

identified reads such as the depicted, that confirm insertion events (like SALK_059379 on Chr 4

at 10.4 Mb) not present in the assembly or BNG maps. Our blastn strategy identified

chromosomal sequence (yellow), and individual alignments with the pROK plasmid sequence

revealed T-strand (blue) and vector backbone (green) sequence. The pink plasmid sequence

within the vector backbone shows an internal breakpoint, which was likely caused by multiple

independent insertion events within the same region. Percent identity of the raw read stretch to

the reference sequence are listed within the annotations.

1108

1109

Supplementary Fig. 5: Variable effects of T-DNA insertions on the local chromatin environment

(a) Quantification of H3K4me3, H2A.Z and H3K27me3 at the two domains next to the T-DNA deletion in SALK_059379 is shown. (b), (c) Impact of the T-DNA integration on the local chromatin environment at *At2g36280* in SALK_059379 seedlings. Visualization (b) and quantification (c) of H3K4me3, H2A.Z and H3K27me3 occupancy around the T-DNA insertion site in Col-0 and SALK_059379 seedlings is shown as well as the schematic illustration of the gene structure of *At2g36280* (b) including the approximate localization of T-DNA insertion. (d), (e) Visualization (d) and quantification (e) of H3K4me3, H2A.Z and H3K27me3 occupancy at flanking regions of the SAIL specific WT inversion on chromosome 1. The corresponding flanking regions in Col-0 were indicated as domain 1 and 2. (f), (g), Impact of the T-DNA integration on the local chromatin environment at *At5g38850* in SAIL_232 seedlings. Visualization (f) and quantification (g) of H3K4me3, H2A.Z and H3K27me3 occupancy around the T-DNA insertion site in Col-0 and SAIL_232 seedlings is shown. Schematic illustration of the gene structure of *At5G38850* (f) indicates the approximate localization of the T-DNA insertion. (h), (i) Impact of the T-DNA integration on the local chromatin environment at *At1g32640* in SALK_061267 seedlings. Visualization (h) and quantification (i) of H3K4me3, H2A.Z and H3K27me3 occupancy around the T-DNA insertion site in Col-0 and SALK_061267 seedlings is shown as well as the schematic illustration of the gene structure of *At1g32640* (h) including the approximate localization of T-DNA insertion. (j), (k) Impact of the T-DNA integration on the local chromatin environment at *At1g23910* in SALK_061267 seedlings. Visualization (j) and quantification (k) of H3K4me3, H2A.Z and H3K27me3 occupancy around the T-DNA insertion site in Col-0 and SALK_061267 seedlings is shown. Schematic illustration of the gene structure of *At1g23910* (j) indicates the approximate localization of the T-DNA insertion. A red arrow marks the approximate location of the T-DNA integration site in the Annoj genome browser screenshots. H3K4me3, H2A.Z and H3K27me3 occupancy was profiled with ChIP-seq whereby

1136 the Col-0 IgG sample serves as a negative control. All shown AnnoJ genome browser tracks

1137 were normalized to the respective sequencing depth. The occupancy of the respective domains

1138 was calculated as the ratio between the respective ChIP-seq sample and the Col-0 IgG control.

1139

Supplementary Fig. 6: T-DNAs affect the local chromatin in a highly diverse manner

(a), (b) Impact of the T-DNA integration on the local chromatin environment at *At5g11930* in

SALK_017723 seedlings. Visualization (a) and quantification (b) of H3K4me3, H2A.Z and

H3K27me3 occupancy around the T-DNA insertion site in Col-0 and SALK_017723 seedlings is

shown as well as the schematic illustration of the gene structure of *At5g11930* (a) including the

approximate localization of T-DNA insertion. (c) Genome browser visualizes the T-DNA-induced

de novo trimethylation of H3K4 and incorporation of H2A.Z at in 50B_HR80 seedlings. The new

T-DNA-induced domain within the gene body of *At3g57300* is indicated as well as a schematic

illustration of the gene structure of *At3g57300* including the approximate localization of the

corresponding T-DNA insertion. (d) Quantification of the new H3K4me3 and H2A.Z domain

within the gene body of *At3g57300* is shown. (e), (f) Impact of the T-DNA integration on the

local chromatin environment at *At1g01700* in SALK_117411 seedlings. Visualization (e) and

quantification (f) of H3K4me3, H2A.Z and H3K27me3 occupancy around the T-DNA insertion

site in Col-0 and SALK_117411 seedlings is shown as well as the schematic illustration of the

gene structure of *At1g01700* (e) including the approximate localization of T-DNA insertion. (g),

(h) T-DNA integration impacts the local chromatin environment at *At3g46650* in SALK_069537

seedlings. Visualization (g) and quantification (h) of H3K4me3, H2A.Z and H3K27me3

occupancy around the T-DNA insertion site in Col-0 and SALK_069537 seedlings is shown.

Schematic illustration of the gene structure of *At3g46650* (g) indicates the approximate

localization of the T-DNA insertion. All ChIP-seq data was visualized with the AnnoJ genome

browser. A red arrow marks the approximate location of the T-DNA integration site in the Annoj

genome browser screenshots. H3K4me3, H2A.Z and H3K27me3 occupancy was profiled with

ChIP-seq whereby the Col-0 IgG sample serves as a negative control. All shown AnnoJ genome

browser tracks were normalized to the respective sequencing depth. The occupancy of the

respective domains was calculated as the ratio between the respective ChIP-seq sample and

the Col-0 IgG control.

1166

	BNG Maps			ONT Assemblies			ONT Individual Reads	
	TAIR Coverage	# Insert./ Rearrange.	Max. Insert kb	TAIR Coverage	# Insert./ Rearrange.	Max. Resolved Insert kb	# Reads	# Insert./ Rearrange.
SALK_ 059379	98%	4	206	98%	3	39	503,388	2
SALK_ 075892	96%	3	96	-	-	-	-	-
SAIL_ 232	97%	7	236	98%	9	50	1,297,359	3
WiscDs Lox_ 449D11	94%	1	164	-	-	-	-	-

Figure 1

a

3.501 Mb   5.202 Mb   7.493 Mb

TAIR10 chr2

Col-0 ONT

SALK_059379 ONT

15.213 Mb   18.866 Mb

ctg_4   ctg_7

| TAIR10 N's | TAIR10 centromere |
| ONT contig joints | T-DNA sequence |

b

15.213 Mb

TAIR10 chr2

SALK_059379 BNG

SALK_059379 BNG
molecules

d

18.866 Mb

TAIR10 chr2

SALK_059379 BNG

SALK_059379 BNG
molecules

27,546 bp

e

SALK_059379
ONT contig_7

5,497 bp

206,818 bp

c

36,841 bp   130,828 bp   39,149 bp

SALK_059379 ONT contig_4   SALK_059379 ONT contig_7

| T-DNA | pROK2 |
| breakpoint | Deletion |

Figure 2

Figure 4

a SALK_059379

Col-0 IgG
Col-0 αH3K4me3
T-DNA αH3K4me3
Col-0 αH2A.Z
T-DNA αH2A.Z
Col-0 αH3K27me3
T-DNA αH3K27me3

deletion

domain 1 2

b SALK_069537
At5g23110
ATG   TAA

Col-0 IgG
Col-0 αH3K4me3
T-DNA αH3K4me3
Col-0 αH2A.Z
T-DNA αH2A.Z
Col-0 αH3K27me3
T-DNA αH3K27me3

c SALK_069537

$\log_2(\text{ChIP/IgG})$

H3K4me3
H2A.Z
H3K27me3

Col-0 T-DNA   Col-0 T-DNA   Col-0 T-DNA

d SALK_117411
At3g53680
ATG   TGA

Col-0 IgG
Col-0 αH3K4me3
T-DNA αH3K4me3
Col-0 αH2A.Z
T-DNA αH2A.Z
Col-0 αH3K27me3
T-DNA αH3K27me3

e SALK_117411

$\log_2(\text{ChIP/IgG})$

H3K4me3
H2A.Z
H3K27me3

Col-0 T-DNA   Col-0 T-DNA   Col-0 T-DNA