Information-dense transcription factor binding site clusters identify target genes with

similar tissue-wide expression profiles and serve as a buffer against mutations

Ruipeng Lu[1], Peter K. Rogan[1,2,3*]

[1] Department of Computer Science, Western University, London, Ontario, Canada,

rlu45@uwo.ca

[2] Department of Biochemistry, Western University, London, Ontario, Canada

[3] Cytognomix Inc., London, Ontario, Canada

* Corresponding author (progan@uwo.ca)

Running title: Binding site clusters identify transcription targets

ABSTRACT

Background: The distribution and composition of *cis*-regulatory modules (e.g. transcription factor binding site (TFBS) clusters) in promoters substantially determine gene expression patterns and TF targets, whose expression levels are significantly regulated by TF binding. TF knockdown experiments have revealed correlations between TF binding profiles and gene expression levels. We present a general framework capable of predicting genes with similar tissue-wide expression patterns from activated or repressed TF targets using machine learning to combine TF binding and epigenetic features.

Methods: Genes with correlated expression patterns across 53 tissues were identified according to their Bray-Curtis similarity. DNase I-accessible promoter intervals of direct TF target genes were scanned with previously derived information theory-based position weight matrices (iPWMs) of 82 TFs. Features from information density-based TFBS clusters were used to predict target genes with machine learning classifiers. The accuracy, specificity and sensitivity of the classifiers were determined for different feature sets. Mutations in TFBSs were also introduced to *in silico* examine their impact on cluster densities and the regulatory states of target genes.

Results: We initially chose the glucocorticoid receptor gene (*NR3C1*), whose regulation has been extensively studied, to test this approach. *SLC25A32* and *TANK* were found to exhibit the most similar expression patterns to this gene across 53 tissues. A Decision Tree classifier exhibited the largest area (0.9987 and 0.9956 respectively before and after eliminating inaccessible promoter intervals based on DNase I HyperSensitive Regions (DHSs)) under the Receiver Operating Characteristic (ROC) curve in detecting such coordinately regulated genes. Target gene prediction was confirmed using siRNA knockdown data of TFs, which was more accurate than CRISPR inactivation. *In-silico* mutation analyses of TFBSs also revealed that one

or more information-dense TFBS clusters in promoters are required for accurate target gene

prediction.

Conclusions: Machine learning based on TFBS information density, organization, and

chromatin accessibility accurately identifies gene targets with comparable tissue-wide

expression patterns. Multiple, information-dense TFBS clusters in promoters appear to protect

promoters from the effects of deleterious binding site mutations in a single TFBS that would

effectively alter the expression state of these genes.

INTRODUCTION

The distinctive organization and combination of transcription factor binding sites (TFBSs) and

regulatory modules in promoters dictates specific expression patterns within a set of genes [1].

Clustering of multiple adjacent binding sites for the same TF (homotypic clusters) and for

different TFs (heterotypic clusters) defines *cis*-regulatory modules (CRMs) in human gene

promoters and can amplify the influence of individual TFBSs on gene expression through

increasing binding affinities, facilitated diffusion mechanisms and funnel effects [2]. Because

tissue-specific TF-TF interactions in TFBS clusters are prevalent, these features can assist in

identifying correct target genes by altering binding specificities of individual TFs [3]. Previously,

we derived iPWMs from ChIP-seq data that can accurately detect TFBSs and quantify their

strengths by computing associated R_i values (Rate of Shannon information transmission for an

individual sequence [4]), with $R_{sequence}$ being the average of R_i values of all binding site

sequences and representing the average binding strength of the TF [3]. Furthermore,

information density-based clustering (IDBC) can effectively identify functional TF clusters by

taking into account both the spatial organization (i.e. intersite distances) and information density

(i.e. R_i values) of TFBSs [5].

TF binding profiles, either derived from *in vivo* ChIP-seq peaks [6–8] or computationally

detected binding sites and CRMs [9], have been shown to be predictive of absolute gene

expression levels using a variety of tissue-specific machine learning classifiers and regression

models. Because signal strengths of ChIP-seq peaks are not strictly proportional to TFBS

strengths [3], representing TF binding strengths by ChIP-seq signals may not be appropriate;

nevertheless, both achieved similar accuracy [10]. CRMs have been formed by combining two

or three adjacent TFBSs [9], which is inflexible, as it arbitrarily limits the number of binding sites

contained in a module, and does not consider differences between information densities of

different CRMs. Chromatin structure (e.g. histone modification (HM) and DNase I

hypersensitivity) were also found to be statistically redundant with TF binding in explaining

tissue-specific mRNA transcript abundance at a genome-wide level [7,8,11,12], which was

attributed to the heterogeneous distribution of HMs across chromatin domains [8]. Combining

these two types of data explained the largest fraction of variance in gene expression levels in

multiple cell lines [7,8], suggesting that either contributes unique information to gene expression

that cannot be compensated for by the other.

The number of genes directly bound by a TF significantly exceeds the number of differentially

expressed (DE) genes whose expression levels significantly change upon knockdown of the TF.

Only a small subset of direct target genes whose promoters overlap ChIP-seq peaks were DE

after individually knocking 59 TFs down using small interfering RNAs (siRNAs) in the GM19238

cell line [13]. Using these knockdown data on 8,872 genes as the gold standard, correlation

between TFBS counts and gene expression levels across 10 different cell lines was more

predictive of DE targets than setting a minimum threshold on TFBS counts [14]. Their TFBS

counts were defined as the number of ChIP-seq peaks overlapping the promoter, though it was

unknown how many binding sites were present in these peaks; positives might not be direct

targets in the TF regulatory cascade, as the promoters of these targets were not intersected

with ChIP-seq peaks. By perturbing gene expression with CAS9-directed clustered regularly

interspaced short palindromic repeats (CRISPR) of 10 different TF genes in K562 cells, the

regulatory effects of each TF on 22,046 genes were dissected by single cell RNA sequencing

with a regularized linear computational model [15]; this accurately revealed DE targets and new

functions of individual TFs, some of which were likely regulated through direct interactions at

TFBSs in their corresponding promoters. Machine learning classifiers have also been applied in

a small number of gene instances to predict targets of a single TF using features extracted from

n-grams derived from consensus binding sequences [16], or from TFBSs and homotypic binding

site clusters [5].

To investigate whether the distribution and composition of CRMs in promoters substantially

determines gene expression profiles of direct TF targets, we developed a general machine

learning framework that predicts which genes have similar expression profiles to a given gene

and predicts DE direct TF targets by combining information theory-based TF binding profiles

with DHSs. Upon filtering for accessible promoter intervals with DHSs, features designed to

capture the spatial distribution and information composition of CRMs were extracted from

clusters identified by the IDBC algorithm from iPWM-detected TFBSs. Though not all direct

targets regulated by multiple TFs share a common tissue-wide expression profile, this

framework provides insight into the transcriptional program of genes with similar profiles by

dissecting their *cis*-regulatory element organization and strengths. We identify genes with

comparable tissue-wide expression profiles by application of Bray-Curtis similarity [17]. Using

transcriptome data generated by CRISPR- [15] and siRNA-based [13] TF knockdowns, we

predicted DE TF target genes that are simultaneously direct targets whose promoters overlap

tissue-specific ChIP-seq peaks.

MATERIALS AND METHODS

To identify genes with similar tissue-wide expression patterns, we formally define gene

expression profiles and pairwise similarity measures between profiles of different genes. A

general machine learning framework relates features extracted from the organization of TFBSs

in these genes to their tissue-wide expression patterns. Since protein-coding (PC) sequences

represent the most widely studied and best understood component of the human genome [18],

positives and negatives for predicting DE direct TF target genes that encode proteins (TF

targets for short below) were obtained from CRISPR- and siRNA-generated knockdown data

(see below).

Similarity between gene expression profiles

For each of 56,238 genes, the Genotype-Tissue Expression (GTEx) project measured its

expression levels in 53 tissues in a number of individuals (N=5-564), and provides the median

expression value (in RPKM (Reads Per Kilobase of transcript per Million mapped reads) in the

GTEx Analysis v6p release) of each gene in each tissue [19]. To capture the tissue-wide overall

expression pattern of a gene instead of within a single tissue, the expression profile of a gene

was defined as its median RPKM across the 53 tissues. In Equation 1, the expression profile of

a gene is described by a vector of size 53; note that different isoforms whose expression

patterns may significantly differ from each other cannot distinguished by this approach.

134
$$EP^A = \left[MEV_1^A, MEV_2^A, \ldots, MEV_{53}^A \right] \ (in \ RPKM) \quad (1)$$

where EP^A is the expression profile of Gene A, MEV_1^A is the median expression value of Gene A

in Tissue 1, MEV_2^A is the median expression value of Gene A in Tissue 2, etc.

To discover other genes whose expression profiles are similar to a given gene, we computed

the Bray-Curtis Similarity (Equation 2) between the expression profiles of gene pairs. Compared

to other similarity metrics (Table 1, Example 1, Additional file 1), the application of this function

is justified by desirable properties, including: 1) maintaining bounds between 0 and 1, 2)

achieving the maximal similarity value 1 if and only if two vectors are identical, and 3) larger

values having a larger impact on the resultant similarity value.

$$sim_{Bray-Curtis}(EP^A, EP^B) = \begin{cases} 1, & if \ \sum_{i=1}^{53} MEV_i^A = \sum_{i=1}^{53} MEV_i^B = 0 \\ 1 - \frac{\sum_{i=1}^{53} |MEV_i^A - MEV_i^B|}{\sum_{i=1}^{53} (MEV_i^A + MEV_i^B)}, & otherwise \end{cases}$$ (2)

Example 1. Assume that Genes *A,B,C,D,E,F* respectively have the following expression profiles

across two tissues: $EP^A = [1,1]$, $EP^B = [2,2]$, $EP^C = [3,3]$, $EP^D = [1,2]$, $EP^E = [1,99]$, $EP^F =$

$[1,100]$. The ground-truth similarity relationships that we can intuitively infer include

$sim(EP^C, EP^A) < sim(EP^C, EP^B) < 1$, and $sim(EP^A, EP^D) < sim(EP^E, EP^F) < 1$. Only the results

computed by the Bray-Curtis Similarity are completely concordant with these ground-truth

relationships (Table 2).

Prediction of genes with similar expression profiles

The framework for identifying genes whose expression profiles most resemble a particular

gene is shown in Figure 1A and B. The genomic locations of all DHSs in 95 cell types generated

by the ENCODE project [18; hg38 assembly] were selected for known promoters [21], then 94

iPWMs exhibiting primary binding motifs for 82 TFs [3] were used to detect TFBSs within the

overlapping intervals. When detecting heterotypic TFBS clusters with the IDBC algorithm, a

minimum threshold $0.1 * R_{sequence}$ was specified for the R_i values of TFBSs in order to remove

weak binding sites that were more likely to be false positive TFBSs [3].

The information density-related features derived from each TFBS cluster included: 1) The

distance between this cluster and the transcription start site (TSS); 2) The length of this cluster;

3) The information content of this cluster (i.e. the sum of R_i values of all TFBSs in this cluster); 4)

The number of binding sites of each TF within this cluster; 5) The number of strong binding sites

($R_i > R_{sequence}$) of each TF within this cluster; 6) The sum of R_i values of binding sites of each TF

within this cluster; 7) The sum of R_i values of strong binding sites ($R_i > R_{sequence}$) of each TF

within this cluster.

For a gene, each of Features 1-3 was defined as a vector whose size equals the number of

clusters in the promoter; thus, the entire vector could be input into a classifier. If two genes

contained different numbers of clusters, the maximum number of clusters among all instances

was determined, and null clusters were added at the 5' end of promoters with fewer clusters,

enabling all instances to have the same cluster count. Machine learning classifiers with default

parameter values in MATLAB were used to generate ROC curves (Additional file 1).

Prediction of TF targets

Using gene expression in the CRISPR-based perturbations

Dixit et al. performed CRISPR-based perturbation experiments using multiple guide RNAs for

each of ten TFs in K562 cells, resulting in a regulatory matrix of coefficients that indicate the

effect of each guide RNA on each of 22,046 genes [15]. The coefficient of a guide RNA on a

gene is defined as the $\log_{10}$(fold change in gene expression level) [15]. Among these ten TFs,

we have previously derived iPWMs exhibiting primary binding motifs for seven (EGR1, ELF1,

ELK1, ETS1, GABPA, IRF1, YY1) [3]. Therefore, the framework for predicting TF targets in the

K562 cell line (Figure 1A and 1C) was applied to these TFs. The criteria for defining a positive

(i.e. a target gene), of a TF were:

1) The fold change in the expression level of this gene for each guide RNA of the TF was >

(or <) 1, consistent with the possibility that the gene was regulated by the TF, and

2) The average fold change in the expression level of this gene for all guide RNAs of the TF

was > threshold ε (or $< 1/\varepsilon$), and

3) The promoter interval (10 kb) upstream of a TSS of this gene overlaps a ChIP-seq peak of

the TF in the K562 cell line.

If the coefficients of all guide RNAs of the TF for a gene are zero, the gene was defined as a

negative. As the threshold ε increases, the number of positives strictly decreases; as ε

decreases, we have increasingly lower confidence in the fact that the positives were indeed

differentially expressed because of the TF perturbation. To achieve a balance between

sensitivity and specificity, we evaluated three different values (i.e. 1.01, 1.05 and 1.1) for ε. For

each TF, all ENCODE ChIP-seq peak datasets from the K562 cell line were merged to

determine positives. To make the numbers of negatives and positives equal to avoid

imbalanced datasets that significantly compromise the classifier performance [22], the Bray-

Curtis function was applied to compute the similarity values in the expression profile between all

negatives and the positive with the largest average coefficient, then the TNs with the smallest

values were selected (Figure 1C).

The DHSs in the K562 cell line were intersected with known promoters. Because TFs may

exhibit tissue-specific sequence preferences due to different sets of target genes and binding

sites in different tissues [3], the iPWMs of EGR1, ELK1, ELF1, GABPA, IRF1, YY1 from the

K562 cell line were used to most accurately detect binding sites; for ETS1, we used the only

available iPWM from the GM12878 cell line [3]. Six features were derived from each homotypic

cluster (i.e. Features 3 and 6 converged to the same value, because only binding sites from a

single TF were used).

Using gene expression in the siRNA-based knockdown

In the GM19238 cell line, 59 TFs were individually knocked down using siRNAs, and

significant changes in the expression levels of 8,872 genes were indicated according to their

corresponding P-values [13]. In these cases, the P-value of a gene for a TF is the probability of

observing the change in the expression level of this gene under the null hypothesis of no

differential expression after TF knockdown; thus the larger the change in the expression level,

the smaller the P-value and the more likely this gene is differentially expressed. They also

indicated whether the promoters of these genes display evidence of binding to TFs by

intersecting with ChIP-seq peaks in the GM12838 cell line. Among these 59 TFs, we have

previously derived accurate iPWMs exhibiting primary binding motifs for 11 (BATF, JUND,

NFE2L1, PAX5, POU2F2, RELA, RXRA, SP1, TCF12, USF1, YY1) [3]. Therefore, the

framework for predicting TF targets in the GM19238 cell line (Figure 1A and 1D) was applied to

these 11 TFs.

We defined a positive (i.e. a target gene) for a TF, if the P-value of this gene for the TF was ≤

0.01, and the promoter interval (10kb) upstream of a TSS of this gene overlapped a ChIP-seq

peak of the TF in the GM12878 cell line. A negative for a TF exhibited the following property: a

P-value > 0.01 for the TF.

The DHSs in the GM19238 cell line mapped from the hg19 genome assembly were first

remapped to the hg38 assembly using liftOver (available at genome.ucsc.edu) prior to being

intersected with known promoters [23]. Aside from RELA, RXRA and NFE2L1, the iPWMs of

TFs from the GM12878 cell line were used to detect binding sites. For RELA, the iPWM from

the GM19099 cell line was used; for RXRA and NFE2L1, the only available iPWMs were

respectively derived from HepG2 and K562 cells and were applied. Although the knockdown

was performed in GM19238, GM12878 and GM19099 are also lymphoblastic cell lines, with

GM19099 and GM19238 both being derived from Yorubans. For this analysis, the iPWMs

derived in GM12878 and GM19099 were more appropriate sources of accessible TFBSs than

those from HepG2 and K562, since GM12878 and GM19099 are of the same tissue type and

are thus more likely comparable to GM19238 than HepG2 and K562.

Mutation analyses on promoters of TF targets

To better understand the significance of individual binding sites for information-dense

clusters and the regulatory state of direct targets, we evaluated the effects of sequence changes

that altered the R_i values of these sites on cluster formation and whether a gene was predicted

to be a TF target. Mutations were sequentially introduced into the strongest binding sites in

TFBS clusters of the EGR1 target gene, *MCM7*, to determine the threshold for cluster formation

after disappearing clusters disabled induction of *MCM7* expression. For one target gene of each

TF from the CRISPR-generated perturbation data, effects of naturally occurring TFBS variants

present in dbSNP (https://www.ncbi.nlm.nih.gov/projects/SNP/) [24] were also evaluated to

explore aspects of TFBS organization that enabled both clusters and promoter activity to be

resilient to binding site mutations. This was done by analyzing whether the occurrence of

individual or multiple single nucleotide polymorphisms (SNPs) lead to the loss of binding sites

and the clusters that contain them, and result in changes in the predictions of these targets.

RESULTS

Similarity between gene expression profiles

To confirm that the Bray-Curtis Similarity can indeed effectively measure how akin the

expression profiles of two genes are to one another, Equation 2 was applied to compute the

similarity values between the expression profiles of the glucocorticoid receptor (*GR* or *NR3C1*)

gene and all other 18,812 PC genes. *NR3C1* is an extensively characterized TF with many

known direct target genes [25]. As a constitutively expressed TF activated by glucocorticoid

ligands, it can mediate the up-regulation of anti-inflammatory genes by binding of homodimers

to glucocorticoid response elements and down-regulation of proinflammatory genes by

complexing with other activating TFs (e.g. NFKB and AP1) and eliminating their ability to bind

targets [25]. NR3C1 can bind its own promoter forming an auto-regulatory loop, which also

contains functional binding sites of 11 other TFs (e.g. SP1, YY1, IRF1, NFKB) whose iPWMs

have been developed and/or mutual interactions have been described previously [3,25].

However, as the expression profile of *NR3C1* integrates all different splicing and translational

isoforms (e.g. *GRα-A* to *GRα-D*, *GRβ*, *GRγ*, *GRδ*), the tissue-specific expression patterns of

these isoforms cannot be distinguished (e.g. levels of the *GRα-C* isoforms are significantly

higher in the pancreas and colon, whereas levels of *GRα-D* are highest in spleen and lungs)

[25]. *SLC25A32* and *TANK* have the greatest similarity in expression to *NR3C1* (0.880 and

0.877 respectively), which is evident based on their overall similar expression patterns across

the 53 tissues (Figure 2).

Prediction of genes with similar expression profiles

In prediction of genes with similar expression profiles to *NR3C1*, we generated ROC curves

to compare the performance of different classifiers (Naïve Bayes, Decision Tree, Random

Forest and Support Vector Machines (SVM) with four different kernels), under two scenarios

depending on whether promoter sequences were first intersected with DHSs (Figure 3).

Decision Tree (DT) exhibited the largest AUC (area under the curve) under both scenarios, and

was one of two most stable classifiers (i.e. ΔAUC < 0.01), with the other being the SVM with

RBF kernel. Inclusion of DHS information significantly improved other classifiers' AUC except

for Naïve Bayes, and generally all TFBSs in a DHS formed a binding site cluster.

Prediction of TF targets

The best-performing DT classifier in distinguishing genes with *NR3C1*-like expression profiles

from others was used to predict TF targets respectively based on the CRISPR- [15] and siRNA-

generated [13] perturbation data. To validate that using all six machine learning features more

comprehensively capture the distribution and composition of CRMs in the promoter, feature

removal was performed by only using TFBS counts. The classifier performance decreased,

except for CRISPR-perturbed GABPA, IRF1 and YY1 after inclusion of DHS information

(Additional file 5).

On the CRISPR-generated knockdown data, after eliminating TFBSs in inaccessible

promoter intervals, i.e. those excluded from tissue-specific DHSs, the DT classifier predicted TF

targets with greater sensitivity and specificity (Table 3). Specifically, predictions for TFs: EGR1,

ELK1, ELF1, ETS1, GABPA, and IRF1 were more accurate than for YY1, which itself represses

or activates a wide range of promoters by binding to sites overlapping the TSS (Table 3).

Accordingly, the perturbation data indicated that YY1 has ~4-22 times more PC targets in the

K562 cell line than the other TFs ($\varepsilon = 1.05$), and its binding has a more significant impact on the

expression levels of target genes (for YY1, the ratio of the PC target counts at $\varepsilon = 1.1$ vs $\varepsilon =$

1.01 was 0.334, which significantly exceeded those of the other TFs (0.017-0.082); Additional

file 3). This is concordant with our previous finding that YY1 extensively interacts with 11

cofactors (e.g. DNA-binding IRF9 and TEAD2; non-DNA-binding DDX20 and PYGO2) in K562

cells, consistent with a central role in specifying erythroid-specific lineage development [3].

Despite a high accuracy of target recognition, sensitivity did not exceed specificity except for

IRF1 (Table 3), due to a relatively large number of false negative genes. Promoters of most TF

targets contain accessible, functional binding sites that significantly change gene expression

levels upon binding. By contrast, promoters of non-targets contain either no accessible binding

sites at all, or accessible, but non-functional sites. The fact that these false negatives were

erroneously predicted to non-targets was attributable to the indistinguishability between

functional binding sites in their promoters and non-functional ones in non-targets in the classifier.

In vivo co-regulation mediated by interacting cofactors, which was excluded by the classifier,

assisted in distinguishing these non-functional sites that do not significantly affect gene

expression [13].

As the threshold ε increased, the accuracy of the classifier for all the TFs monotonically

increased as expected (Figure 4). For a gene to be defined as a DE target of a TF, the average

fold change in its expression level for all guide RNAs that downregulated the TF were required

to reach the minimum threshold ε. Upon TF knockdown, ε is inversely correlated with the

number of target genes, but positively correlated with fold changes in their corresponding

expression levels. In general, more significantly DE genes have been associated with a higher

number of TFBSs in their promoters [13]. Thus, at greater ε, there are larger differences in the

values of machine learning features derived from TFBS clusters between direct targets and non-

targets.

With the siRNA-generated knockdown data, the performance of the DT classifier was

compared to the approach inferring DE targets by correlating TF binding with gene expression

levels across ten cell types [14]. In this correlation-based approach, three measures (i.e. the

absolute Pearson correlation coefficient (PCC), the absolute Spearman correlation coefficient

(SCC), and the absolute combined angle ratio statistic (CARS)), whose performance was

evaluated with precision-recall curves, were alternatively used to compute a correlation score

between the number of ChIP-seq peaks overlapping the promoter and gene expression values.

Genes predicted to be DE targets had scores above the threshold resulting in a 1.5-fold

increase compared to the background precision (i.e. the DE target count / 8,872). For example,

in the case of YY1, which was the only TF analyzed by both approaches, the performance of the

DT classifier was 0.98 (precision) and 0.55 (recall) after including DHS information (Table 4).

This classifier outperformed all three correlation measures (PCC: 0.467 and 0.003; SCC: 0.467

and 0.006; CARS: 0.467 and 0.003 directly obtained from [14]), even though the correlation-

based approach used a less stringent P-value threshold (0.05) for defining differential

expression of likely non-direct targets, and intersected ChIP-seq peaks over shorter 5kb

promoter intervals upstream of the TSS. Three reasons explain why the correlation-based

approach exhibited lower recall, including that 1) it did not use machine learning classifiers, 2)

its larger P-value threshold (0.05) generated a larger number of positives and 3) its positives

also include DE targets that cannot be directly bound.

Intersection of genes with similar expression profiles and TF targets

To determine how many TF targets have similar tissue-wide expression profiles, we

intersected the set of targets with the set of 500 PC genes with the most similar expression

profiles for each TF (Table 5, Additional file 6). The TFs PAX5 and POU2F2 are primarily

expressed in B cells, and their respective targets *IL21R* and *CD86* are also B cell-specific,

which accounts for the high similarity in the expression profile between them. There are

respectively 21 and 7 nuclear mitochondrial genes (e.g. *MRPL9* and *MRPS10*, which are

subunits of mitochondrial ribosomes) in the intersections for YY1 in the K562 and GM19238 cell

lines [26]. Previous studies reported that YY1 upregulates a large number of mitochondrial

genes by complexing with PGC-1α in C2C12 cells [27], and genes involved in the mitochondrial

respiratory chain in K562 cells [15], which is consistent with the idea that YY1 may broadly

regulate mitochondrial function (within all 53 tissues in addition to the erythrocyte, lymphocyte

and skeletal muscle cell lines).

Between 0.4%-25% of genes with similar expression profiles to the TFs are actually their

targets (Table 5); the majority are non-targets whose promoters contain non-functional binding

sites that are distinguished from targets by their lack of coregulation by corresponding cofactors.

For YY1 and EGR1, we validated this hypothesis by contrasting the flanking cofactor binding

site distributions and strengths in the promoters of the most similarly expressed target genes

(YY1: *MRPL9*, *BAZ1B*; EGR1: *CANX*, *NPM1*) and non-target genes (YY1: *ADNP*, *RNF25*;

EGR1: *GUCY2F*, *AWAT1*). Strong and intermediate recognition sites for TFs: SP1, KLF1,

CEBPB formed heterotypic clusters with adjacent YY1 sites; as well TFBSs of SP1, KLF1, and

NFY were frequently present adjacent to EGR1 binding sites. These patterns contrasted with

the enrichment of CTCF and ETV6 binding sites in gene promoters of YY1 and EGR1 non-

targets (Additional file 7). Previous studies have reported that KLF1 is essential for terminal

erythroid differentiation and maturation [26], direct physical interactions between YY1 and the

constitutive activator SP1 synergistically induce transcription [29], the activating CEBPB

promotes differentiation and suppresses proliferation of K562 cells by binding the promoter of

the *G-CSFR* gene encoding a hematopoietin receptor [30], EGR1 and SP1 synergistically

cooperate at adjacent non-overlapping sites on *EGR1* promoter but compete binding at

overlapping sites [31]; whereas CTCF binding sites function as an insulator blocking the effects

of *cis*-acting elements and preventing gene activation by mediating long-range DNA loops to

alter topological chromatin structure [32,33], and ETV6, a member of the ETS family, is a

transcriptional repressor required for bone marrow hematopoiesis and associated with leukemia

development [34].

Mutation analyses on promoters of direct targets

Because the promoters of most direct targets contain multiple binding site clusters, we

anticipate that this enables these genes' expression to be naturally robust against binding site

mutations; in other words, the other clusters can compensate for the loss of a cluster destroyed

by mutations in binding sites, so that the mutated promoters are still capable of effectively

inducing gene transcription upon TF binding. First, we validated this hypothesis *in silico* by

examining whether introducing artificial variants into binding sites in the promoter of the target

gene *MCM7* of EGR1 changes the classifier output (Figure 5). Specifically, in the K562 cell line,

MCM7 is upregulated by EGR1. Knockdown of *MCM7* has an anti-proliferative and pro-

apoptotic effect on K562 cells [35] and the loss of EGR1 increases leukemia initiating cells [36],

which suggests that EGR1 may act as a tumor suppressor in K562 cells through the *MCM7*

pathway.

First, the strongest binding site (at position chr7:100103347 [hg38], - strand, R_i = 12.0 bits) in

the promoter was eliminated by a G->A mutation, resulting in the disappearance of Cluster 1,

which consists of two sites (the other site at chr7:100103339, -, 4.3 bits). EGR1 was still

predicted to compensate for this mutation, due to the presence of the other two clusters

comprising weaker binding sites of intermediate strength (chr7:100102252, +, 7.0 bits;

chr7:100102244, +, 1.3 bits; chr7:100101980, +, 8.9 bits; chr7:100101977, +, 2.2 bits;

chr7:100101984, +, 1.9 bits), enabling the promoter to maintain capability of inducing *MCM7*

expression (Figure 5). These adjacent clustered sites, which may not be strong enough to bind

TFs and individually activate transcription, can stabilize each other's binding [2]. The weaker

sites flanking a strong binding site in a cluster can direct the TF molecule to the strong site and

extend the period of the molecule physically associating with the strong site, which is termed,

the funnel effect [2]. Further, Clusters 2 and Cluster 3 were respectively removed by G->T and

C->G mutations abolishing the strongest site in either cluster, which altered the prediction, that

is, EGR1 lost the capability to induce *MCM7* transcription (Figure 5). The remaining four sparse

weak sites do not form a cluster and cannot completely supplant the disrupted strong sites.

Further, we examined the impacts of known natural SNPs on binding site strengths, clusters

and the regulatory state of the promoter for a direct target of each of the seven TFs from the

CRISPR-generated perturbation data (Table 6). Often a single SNP (e.g. rs996639427 of EGR1)

can affect the strengths of multiple binding sites (Table 6). Apart from SNPs that are predicted

to abolish binding (Figure 5), leaky variants that merely weaken TF binding are common (Table

6). Binding stabilization between adjacent sites and the funnel effect enable the CRMs

comprised of information-dense clusters to be robust to mutations in individual binding sites. In

this way, neither mutations that abolish TFBSs nor leaky SNPs in flanking weak sites can

destroy functional clusters (e.g. rs1030185383 and rs5874306 of IRF1), whereas SNPs with

large reductions in R_i values of central strong sites are more likely to abolish clusters (e.g.

rs865922947, rs946037930, rs917218063 and rs928017336 of YY1) (Table 6). More generally,

the presence of multiple clusters enables promoters to be effectively resilient to the effects

binding site mutations; only the complete abolishment of all clusters resulting from the

simultaneous occurrence of multiple SNPs can transform the promoter to be unresponsive to TF

binding to residual weak sites (e.g. rs997328042, rs1020720126 and rs185306857 of GABPA)

(Table 6). Furthermore, a relatively small number of SNPs that strengthen TF binding and

eventually amplify the regulatory effect of the TF on the gene expression level are also present

(e.g. rs887888062 of EGR1 and rs751263172 of ELF1) (Table 6), suggesting that, in addition to

deleterious mutations, benign variants may also be found in promoters, consistent with the

expectations of neutral theory [37].

DISCUSSION

In this study, the Bray-Curtis similarity function was initially shown (for the *NR3C1* gene) to

measure the relatedness of overall expression patterns between genes across a diverse set of

tissues. The resulting machine learning framework distinguished Bray-Curtis function-defined

similar from dissimilar genes based on the distribution, strengths and compositions of TFBS

clusters in accessible promoters, which can substantially account for the corresponding gene

expression patterns. Using knockdown data as the gold standard, the combinatorial use of TF

binding profiles and chromatin accessibility was also demonstrated to be predictive of TF targets.

A binding site comparison confirmed that coregulatory cofactors can be used to responsible for

distinguishing between functional sites in targets and non-functional ones in non-targets.

Furthermore, mutation analyses on binding sites of targets demonstrated that the existence of

both multiple TFBSs in a cluster and multiple information-dense clusters in a promoter enables

both the cluster and the promoter to be resilient to binding site mutations.

The DT classifier improved after intersecting promoters with DHSs in prediction of TF targets

with the CRISPR-generated knockdown data (Table 3). This intersection eliminated noisy

binding sites that are inaccessible to TF proteins in promoters; specifically, it widened

discrepancies in feature vectors between positives and negatives. If the 10kb promoter of a

gene instance does not overlap DHSs, its feature vector will only consist of 0; the percentages

of negatives whose promoters do not overlap DHSs considerably exceeded those of positives

(Additional file 8), which led to an excess of negatives with feature vectors containing only 0

after intersection. This explains why these negatives are not DE targets of the TFs in the K562

and GM19238 cell lines, because their entire promoters are not open to TF molecules; other

regulatory regions besides the proximal promoters (e.g. intronic enhancers [38]) still enable the

TFs to effectively control the expression of the positives with inaccessible promoters. The

relatively poor performance of the classifier on YY1 (Table 3) is attributable to its smaller

percentage of negatives with inaccessible promoters (Additional file 8) and the larger number of

functional binding sites in the K562 cell line.

Our *in-silico* mutation analyses revealed that some deleterious TFBS mutations could be

compensated for by other information-dense clusters in a promoter; thus predicting the effects

of mutations in individual binding sites would not be sufficient to interpretation of downstream

effects. Though compensatory clusters may maintain gene expression, the promoter will provide

lower levels of activity than the wild-type promoter could, which is a recipe for achieving natural

phenotypic diversity. Few published studies in molecular diagnostics have specifically examined

the effects of naturally occurring variants within clustered TFBSs; thus IDBC-based machine

learning provided an alternative computational approach to predict deleterious mutations that

actually impact (i.e. repress or abolish) transcription of target genes and result in abnormal

phenotypes, and to simultaneously minimize false positive calls of TFBS mutations that

individually have little or no impact.

Apart from these TFs, the Bray-Curtis Similarity can be directly applied to identify the ground-

truth genes with overall similar tissue-wide expression patterns to any other gene whose

expression profile is known. Further studies could investigate the biological significance

underlying the phenomenon that all these genes share a common expression pattern, including

the similarity between other regulatory regions besides proximal promoters in terms of TFBSs

and epigenetic markers. This machine learning framework can also be applied to predict target

genes for other TFs and in other cell lines, depending on the availability of corresponding

knockdown data.

There are a number of limitations of our approach. The Bray-Curtis function seems unable to

accurately measure the similarity between the expression profiles of a gene (e.g. *MIR23A*)

without any detectable mRNA in any of the 53 tissues analyzed and genes (e.g. the ubiquitously

expressed *NR3C1* and stomach-specific *PGA3*) that are expressed in at least one tissue.

Intuitively, in terms of expression patterns *PGA3* is more similar to *MIR23A* than *NR3C1*;

however, the Bray-Curtis similarity values indicate that both *PGA3* and *NR3C1* bear no similarity

to *MIR23A* (i.e. $sim_{Bray-Curtis}(NR3C1, MIR23A) = sim_{Bray-Curtis}(PGA3, MIR23A) = 0$). Another

possible limitation in classifier performance in the prediction of genes with similar tissue-wide

expression profiles is that only binding sites of 82 TFs were analyzed due to a lack of available

iPWMs for other TFs, given that 2000-3000 sequence-specific DNA-binding TFs are estimated

to be encoded in the human genome [39]. For example, four TFs (CREB, MYB, NF1, GRF1)

were previously reported to bind the promoter of the *NR3C1* gene to activate or repress its

expression, however their iPWMs exhibiting known primary motifs could not be successfully

derived from ChIP-seq data [3,25]. Regarding the CRISPR-generated knockdown data used

here, positives were inferred to be direct targets by intersecting their promoters with

corresponding ChIP-seq peaks, which may not be completely accurate, due to the presence of

noise peaks that do not contain true TFBSs [3,40]. Small fold changes in the expression levels

of DE targets could arise from compromised efficiency of knockdowns as a result of suboptimal

guide RNAs or the limitations of perturbing only a single allele of the TF. Finally, the framework

developed here only takes into account the 10kb interval proximal to the TSS, and would not

therefore capture long range enhancer effects beyond this distance; by contrast, correlation

based approaches have successfully incorporated multiple definitions of promoter length [14].

CONCLUSIONS

The Bray-Curtis function is able to effectively quantify the similarity between tissue-wide gene

expression profiles. By analysis of information theory-based TF binding profiles that captured

the spatial distribution and information contents of TFBS clusters, ChIP-seq and chromatin

accessibility data, we described a machine learning framework that distinguished tissue-wide

expression profiles of similar vs dissimilar genes and identified TF target genes. Functional

binding sites in target genes that significantly alter expression levels upon direct binding are at

least partially distinguished by TF-cofactor coregulation from non-functional sites in non-targets.

Finally, *in-silico* mutation analyses demonstrated that the presence of multiple information-

dense clusters in the promoter, as a protective mechanism, reduces deleterious mutations that

can significantly alter the regulatory state and expression level of the gene.

LIST OF ABBREVIATIONS

TF: transcription factor, TFBS: transcription factor binding site, CRM: *cis*-regulatory modules,

iPWM: information theory-based position weight matrix, IDBC: information density-based

clustering, R_i: Rate of Shannon information transmission for an individual sequence, ChIP-seq:

chromatin immunoprecipitation with massively parallel DNA sequencing, HM: histone

modification, mRNA: messenger RNA, siRNA: small interfering RNA, CRISPR: clustered

regularly interspaced short palindromic repeats, DE: differentially expressed, DHS:

deoxyribonuclease I hypersensitive region, RPKM: reads per kilobase of transcript per million

mapped reads, GR: glucocorticoid receptor, GTEx: genotype-tissue expression, ENCODE:

encyclopedia of DNA elements, TSS: transcription start site, SVM: support vector machine, RBF:

radial basis function, DT: decision tree, ROC: receiver operating characteristic, AUC: area under

the curve, PC: protein-coding, PCC: absolute Pearson correlation coefficient, SCC: absolute

Spearman correlation coefficient, CARS: the absolute combined angle ratio statistic, SNP:

single nucleotide polymorphism.

ADDITIONAL FILES

Additional file 1: The mathematical definitions of the four other similarity metrics, the workflow

of the IDBC algorithm, the mathematical definitions of five statistical variables to measure

classifier performance, and the default parameter values of classifiers in MATLAB

Format: .docx

Additional file 2: The lists of positives and negatives in the machine learning classifiers to

predict genes with similar tissue-wide expression profiles

Format: .xlsx

Additional file 3: The lists of positives and negatives in the DT classifier to predict TF targets

based on the CRISPR-generated knockdown data

Format: .xlsx

Additional file 4: The lists of positives and negatives in the DT classifier to predict DE direct

targets based on the siRNA-generated knockdown data

Format: .xlsx

Additional file 5: The performance of the DT classifier using only TFBS counts

Format: .xlsx

Additional file 6: The list of the most similar 500 PC genes to each TF in terms of expression

profiles, and the intersection of these 500 genes and target genes of the TF

Format: .xlsx

Additional file 7: Cofactor binding sites adjacent to YY1 and EGR1 sites in the promoters of

their targets and non-targets

Format: .docx

Additional file 8: The percentages of positives and negatives whose promoters do not overlap

DHSs for the CRISPR-perturbed TFs

Format: .xlsx

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The median RPKM, TSS coordinate, DNase I hypersensitivity and ChIP-seq data are

respectively available from the GTEx Analysis V6p release (www.gtexportal.org), Ensembl

Biomart (www.ensembl.org) and ENCODE (www.encodeproject.org). The CRISPR- and siRNA-

generated knockdown data are available from the supplementary information files of Dixit et al.

[15] and Cusanovich et al. [13]. The code implementing this machine learning framework and

the datasets generated and/or analysed by this framework are available in Zenodo

(https://doi.org/10.5281/zenodo.1249635). All other data generated or analysed during this

study are included in this published article and its supplementary information files.

Competing interests

PKR is the inventor of US Patent 5,867,402 and other patents pending, which apply iPWMs

to the prediction and validation of mutations. He cofounded Cytognomix, Inc., which is

developing software based on this technology for complete genome or exome mutation analysis.

Funding

Natural Sciences and Engineering Research Council of Canada Discovery Grant [RGPIN-

2015-06290]; Canada Foundation for Innovation; Canada Research Chairs; Cytognomix Inc.

Funding for open access charge: University of Western Ontario and the Natural Sciences and

Engineering Research Council.

Authors' contributions

PKR defined the objectives and directed the study. RL and PKR devised the general

machine learning framework. RL implemented this framework and collected the results. Both RL

and PKR interpreted the results and wrote the manuscript.

Acknowledgements

We are grateful to Ben Shirley and Eliseos Mucaki for constructive comments on the paper.

REFERENCES

1. Hosseinpour B, Bakhtiarizadeh MR, Khosravi P, Ebrahimie E. Predicting distinct organization

of transcription factor binding sites on the promoter regions: a new genome-based approach to

expand human embryonic stem cell regulatory network. Gene. 2013;531:212–9.

2. Ezer D, Zabet NR, Adryan B. Homotypic clusters of transcription factor binding sites: A model

system for understanding the physical mechanics of gene expression. Comput Struct Biotechnol

J. 2014;10:63–9.

3. Lu R, Mucaki EJ, Rogan PK. Discovery and validation of information theory-based

transcription factor and cofactor binding site motifs. Nucleic Acids Res. 2017;45:e27.

4. Schneider TD. Information content of individual genetic sequences. J Theor Biol.

1997;189:427–41.

5. Dinakarpandian D, Raheja V, Mehta S, Schuetz EG, Rogan PK. Tandem machine learning

for the identification of genes regulated by transcription factors. BMC Bioinformatics. 2005;6:204.

6. Ouyang Z, Zhou Q, Wong WH. ChIP-Seq of transcription factors predicts absolute and

differential gene expression in embryonic stem cells. Proc Natl Acad Sci U S A.

2009;106:21521–6.

7. Cheng C, Alexander R, Min R, Leng J, Yip KY, Rozowsky J, et al. Understanding

transcriptional regulation by integrative analysis of transcription factor binding data. Genome

Res. 2012;22:1658–67.

8. Budden DM, Hurley DG, Cursons J, Markham JF, Davis MJ, Crampin EJ. Predicting

expression: the complementary power of histone modification and transcription factor binding

data. Epigenetics Chromatin. 2014;7:36.

9. Smith AD, Sumazin P, Xuan Z, Zhang MQ. DNA motifs in human and mouse proximal

promoters predict tissue-specific expression. Proc Natl Acad Sci U S A. 2006;103:6275–80.

10. McLeay RC, Lesluyes T, Cuellar Partida G, Bailey TL. Genome-wide in silico prediction of

gene expression. Bioinforma Oxf Engl. 2012;28:2789–96.

11. Karlić R, Chung H-R, Lasserre J, Vlahovicek K, Vingron M. Histone modification levels are

predictive for gene expression. Proc Natl Acad Sci U S A. 2010;107:2926–31.

12. Dong X, Greven MC, Kundaje A, Djebali S, Brown JB, Cheng C, et al. Modeling gene

expression using chromatin features in various cellular contexts. Genome Biol. 2012;13:R53.

13. Cusanovich DA, Pavlovic B, Pritchard JK, Gilad Y. The functional consequences of variation

in transcription factor binding. PLoS Genet. 2014;10:e1004226.

14. Banks CJ, Joshi A, Michoel T. Functional transcription factor target discovery via compendia

of binding and expression profiles. Sci Rep. 2016;6:20649.

15. Dixit A, Parnas O, Li B, Chen J, Fulco CP, Jerby-Arnon L, et al. Perturb-Seq: Dissecting

Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. Cell.

2016;167:1853-1866.e17.

16. Cui S, Youn E, Lee J, Maas SJ. An improved systematic approach to predicting transcription

factor target genes using support vector machine. PloS One. 2014;9:e94519.

17. Bray JR, Curtis JT. An Ordination of the Upland Forest Communities of Southern Wisconsin.

Ecol Monogr. 1957;27:325–349.

18. Consortium IHGS. Finishing the euchromatic sequence of the human genome. Nature.

2004;431:931–45.

19. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. Nat Genet.

2013;45:580–5.

20. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human

genome. Nature. 2012;489:57–74.

21. Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, et al. The

accessible chromatin landscape of the human genome. Nature. 2012;489:75–82.

22. He H, Garcia EA. Learning from Imbalanced Data. IEEE Trans Knowl Data Eng.

2009;21:1263–1284.

23. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The human

genome browser at UCSC. Genome Res. 2002;12:996–1006.

24. Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, Smigielski EM, et al. dbSNP: the NCBI

database of genetic variation. Nucleic Acids Res. 2001;29:308–11.

25. Vandevyver S, Dejager L, Libert C. Comprehensive overview of the structure and regulation

of the glucocorticoid receptor. Endocr Rev. 2014;35:671–93.

26. Calvo SE, Clauser KR, Mootha VK. MitoCarta2.0: an updated inventory of mammalian

mitochondrial proteins. Nucleic Acids Res. 2016;44:D1251-1257.

27. Cunningham JT, Rodgers JT, Arlow DH, Vazquez F, Mootha VK, Puigserver P. mTOR

controls mitochondrial oxidative function through a YY1-PGC-1alpha transcriptional complex.

Nature. 2007;450:736–40.

28. Tallack MR, Perkins AC. KLF1 directly coordinates almost all aspects of terminal erythroid

differentiation. IUBMB Life. 2010;62:886–90.

29. Seto E, Lewis B, Shenk T. Interaction between transcription factors Sp1 and YY1. Nature.

1993;365:462–4.

30. Ferrari-Amorotti G, Mariani SA, Novi C, Cattelani S, Pecorari L, Corradini F, et al. The

biological effects of C/EBPalpha in K562 cells depend on the potency of the N-terminal

regulatory region, not on specificity of the DNA binding domain. J Biol Chem. 2010;285:30837–

50.

31. Huang RP, Fan Y, Ni Z, Mercola D, Adamson ED. Reciprocal modulation between Sp1 and

Egr-1. J Cell Biochem. 1997;66:489–99.

32. Bell AC, West AG, Felsenfeld G. The protein CTCF is required for the enhancer blocking

activity of vertebrate insulators. Cell. 1999;98:387–96.

33. Hou C, Zhao H, Tanimoto K, Dean A. CTCF-dependent enhancer-blocking by alternative

chromatin loop formation. Proc Natl Acad Sci U S A. 2008;105:20398–403.

34. Wang LC, Swat W, Fujiwara Y, Davidson L, Visvader J, Kuo F, et al. The TEL/ETV6 gene is

required specifically for hematopoiesis in the bone marrow. Genes Dev. 1998;12:2392–402.

35. Tian L, Liu J, Xia G-H, Chen B-A. RNAi-mediated knockdown of MCM7 gene on CML cells

and its therapeutic potential for leukemia. Med Oncol Northwood Lond Engl. 2017;34:21.

36. Maifrede S, Liebermann D, Hoffman B. Egr-1, a Stress Response Transcription Factor and

Myeloid Differentiation Primary Response Gene, Behaves As Tumor Suppressor in CML. Blood.

2014;124:2211.

37. Kimura M. The neutral theory of molecular evolution. Sci Am. 1979;241:98–100, 102, 108

passim.

38. Hural JA, Kwan M, Henkel G, Hock MB, Brown MA. An intron transcriptional enhancer

element regulates IL-4 gene locus accessibility in mast cells. J Immunol Baltim Md 1950.

2000;165:3239–49.

39. Vaquerizas JM, Kummerfeld SK, Teichmann SA, Luscombe NM. A census of human

transcription factors: function, expression and evolution. Nat Rev Genet. 2009;10:252–63.

40. Kidder BL, Hu G, Zhao K. ChIP-Seq: technical considerations for obtaining high-quality data.

Nat Immunol. 2011;12:918–22.

41. Pearson K. Note on Regression and Inheritance in the Case of Two Parents. Proc R Soc

Lond. 1895;58:240–2.

42. Spearman C. The Proof and Measurement of Association between Two Things. Am J

Psychol. 1904;15:72–101.

 FIGURE LEGENDS

 Figure 1. The general framework for predicting genes with similar tissue-wide expression

 profiles and TF targets

     A) An overview of the machine learning framework. The steps enclosed in the dashed

 rectangle vary across prediction of genes with similar expression profiles and TF targets. The

 step with a dash-dotted border that intersects promoters with DHSs is a variant of the primary

 approach. In the IDBC algorithm (Additional file 1), the parameter l is the minimum threshold on

 the total information contents of TFBS clusters. In prediction of genes with similar expression

 profiles, the minimum value was 939, which was the sum of mean information contents ($R_{sequence}$

 values) of all 94 iPWMs; in prediction of direct targets, this value was the $R_{sequence}$ value of the

 single iPWM used to detect TFBSs in each promoter. The parameter d is the radius of initial

 clusters in base pairs, whose value, 25, was determined empirically. The performance of seven

 different classifiers were evaluated with statistics (accuracy, sensitivity and specificity)

 (Additional file 1). **B)** Obtaining of the positives and negatives for identifying genes with similar

 expression profiles to a given gene(Additional file 2). **C)** Obtaining of the positives and negatives

 for predicting target genes of seven TFs using the CRISPR-generated perturbation data in K562

 cells (Additional file 3). **D)** Obtaining of the positives and negatives for predicting target genes of

 11 TFs using the siRNA-generated knockdown data in GM19238 cells (Additional file 4).

 Figure 2. Expression profiles of NR3C1, SLC25A32 and TANK

     Visualization of the expression values (in RPKM) of these genes across 53 tissues from

 GTEx. For each gene, the colored rectangle belonging to each tissue indicates the valid RPKM

 of all samples in the tissue, the black horizontal bar in the rectangle indicates the median RPKM,

 the hollow circles indicate the RPKM of the samples considered as outliers, and the grey vertical

 bar indicates the sampling error. By comparing the pictures, the overall expression patterns of

the three genes across the 53 tissues resemble each other (e.g. all three genes exhibit the

highest expression levels in lymphocytes and the lowest levels in brain tissues).

Figure 3. Comparison between the performance of different classifiers in prediction of

genes with similar expression profiles to *NR3C1*.

A) ROC curves and AUC of seven classifiers without intersecting promoters with DHSs. **B)**

ROC curves and AUC of seven classifiers after intersecting promoters with DHSs. The Decision

Tree classifier exhibited the largest AUC under both scenarios, and inclusion of DHS

information significantly improved other classifiers' AUC except for Naïve Bayes.

Figure 4. Accuracy of the Decision Tree classifier when using three different values for ε

Each accuracy value was averaged from 10 rounds of 10-fold cross validation, when the

minimum threshold ε on the average fold change in gene expression levels under all guide

RNAs of the TF took three different values 1.01, 1.05 and 1.1. As ε increased, accuracy for all

seven TFs monotonically increased.

Figure 5. Mutation analyses on the target MCM7 of EGR1

This figure depicts the effect of a mutation in each EGR1 binding site cluster of the MCM7

promoter on the expression level of MCM7, which is a target of the TF EGR1. The strongest

binding site in each cluster were abolished by a single nucleotide variant. Upon loss of all three

clusters, only weak binding sites remained and EGR1 was predicted to no longer be able to

effectively regulate MCM7 expression. Multiple clusters in the promoters of TF targets confer

robustness against mutations within individual binding sites that define these clusters.

TABLES

Table 1. Comparison between metrics in measurement of similarity between gene
expression profiles

Similarity metric	Property 1[†,‡]	Property 2[†]	Property 3[†]
Bray-Curtis	√; [0,1]	√	√
Euclidean	√; (0,1]	√	×
Cosine	√; [0,1]	×	√
Pearson correlation [41]	×; [-1,1]	×	×
Spearman correlation [42]	×; [-1,1]	×	×

704 [†] √ and × respectively indicate that the similarity metric satisfies and does not satisfy the
705 property.

706 [‡] The interval in each cell indicates the range in which the result computed by the similarity
707 metric lies.

709 **Table 2. Similarity values computed by different metrics**

Similarity metric	$sim(EP^C, EP^B)$	$sim(EP^C, EP^A)$	$sim(EP^E, EP^F)$	$sim(EP^A, EP^D)$
Bray-Curtis	0.8	0.5	≈ 0.995	0.8
Euclidean	≈ 0.41	≈ 0.26	0.5	0.5
Cosine	1	1	≈ 0.999999995	≈ 0.949
Pearson correlation	Undefined	Undefined	1	1
Spearman correlation	1	1	1	1

711 **Table 3**. **The Decision Tree classifier performance for predicting TF targets using the**

712 **CRISPR-generated knockdown data**

TF	Excluding DHS information[†]			Including DHS information[†]		
	Sensitivity	Specificity	Accuracy	Sensitivity	Specificity	Accuracy
EGR1	0.58	0.62	0.60	0.78	0.81	0.80
ELF1	0.59	0.65	0.62	0.83	0.87	0.85
ELK1	0.59	0.59	0.59	0.80	0.81	0.81
ETS1	0.59	0.6	0.59	0.81	0.81	0.81
GABPA	0.55	0.57	0.56	0.72	0.75	0.74
IRF1	0.54	0.55	0.54	0.76	0.64	0.70
YY1	0.50	0.51	0.51	0.45	0.69	0.57

713 [†]The average performance of 10 rounds of 10-fold cross validation when setting ε to 1.05 is

714 indicated. The CRISPR-generated knockdown data were obtained from Dixit et al [15].

716   **Table 4**. **The Decision Tree classifier performance for predicting TF targets using the**

717   **siRNA-generated knockdown data**

TF	Excluding DHS information[†]			Including DHS information[†]		
	Sensitivity	Specificity	Accuracy	Sensitivity	Specificity	Accuracy
BATF	0.96	0.97	0.96	0.85	1	0.93
JUND	0.86	0.90	0.88	0.80	1	0.90
NFE2L1	0.92	0.95	0.94	0.71	0.93	0.82
PAX5	0.96	0.97	0.96	0.88	0.98	0.93
POU2F2	0.97	0.97	0.97	0.89	0.99	0.94
RELA	0.95	0.96	0.96	0.83	0.97	0.90
RXRA	0.93	0.91	0.92	0.84	0.95	0.89
SP1	0.98	0.98	0.98	0.89	0.99	0.94
TCF12	0.98	0.98	0.98	0.86	0.99	0.93
USF1	0.97	0.98	0.97	0.83	0.98	0.90
YY1	1	1	1	0.55	0.99	0.77

718   [†] The average performance of 10 rounds of 10-fold cross validation is indicated. The siRNA-

719   generated knockdown data were obtained from Cusanovich et al [13].

Table 5. Intersection of TF targets and 500 protein-coding genes with the most similar expression profiles

TF	Cell line	Number of targets	Size of intersection	Targets among the most similar 10 genes[§]
EGR1	K562	169	12	None
ELF1		78	5	None
ELK1		112	4	GNL1(8^{th})
ETS1		267	15	None
GABPA		513	25	TAF1(1^{st})
IRF1		457	10	None
YY1		1752	127	MRPL9(2^{nd}), BAZ1B(6^{th}), ENY2(7^{th}), NUB1(8^{th}), USP1(9^{th}), HNRNPR(10^{th})
	GM19238	1040	61	MED4(1^{st}), SURF6(3^{rd}), BAZ1B(6^{th})
BATF		186	21	None
JUND		44	2	None
NFE2L1		58	4	None
RELA		247	13	HMG20B(9^{th})
RXRA		181	3	None
SP1		1595	81	None
TCF12		655	20	None
USF1		301	21	None
PAX5		918	86	IL21R(8^{th})
POU2F2		532	26	CD86(3^{rd})

[§] The rank of each target in the list of similar genes in the descending order of Bray-Curtis similarity values is shown in the brackets immediately following the target.

 Table 6. Mutation analyses on promoters of TF targets

TF	Target	Normal cluster	Normal allele§	SNP ID§	Variant allele§	Variant cluster‡	Classifier output: Variant†	Wild-type
EGR1 (R_sequence = 12.2899 bits)	EID2B	Cluster 1 of 2	GAGGGGGCATC (chr19:39540286, -, 7.22 bits)	rs538610162 (chr19:39540296C>G)	CAGGGGGCATC (chr19:39540286, -, 4.84 bits)	Abolished	√	√
				rs759233998 (chr19:39540294C>T)	GAAGGGGCATC (chr19:39540286, -, 0.06 bit)	Abolished	√	
				rs974735901 (chr19:39540288T>A)	GAGGGGGCTTC (chr19:39540286, -, 6.90 bits)	Cluster 1 of 2	√	
				rs978230260 (chr19:39540287A>T)	GAGGGGGCAAC (chr19:39540286, -, 5.31 bits)	Abolished	√	
		Cluster 2 of 2	GCGTGCGTGGG (chr19:39540162, +, 1.59 bits)	rs764734511 (chr19:39540162G>A)(chr19:39540162G>C)	ACGTGCGTGGG (chr19:39540162, +, -0.72 bit)	Cluster 2 of 2	√	×
					CCGTGCGTGGG (chr19:39540162, +, -0.79 bit)	Cluster 2 of 2	√	
				rs996639427 (chr19:39540170G>C)	GCGTGCGTCGG (chr19:39540162, +, -5.21 bits)	Abolished	√	
					GCGTCGGCGCT (chr19:39540165, +, -0.85 bit)	Abolished	√	
			GCGTGGGCGCT (chr19:39540166, +, 9.72 bits)	rs1027751538 (chr19:39540174G>A)	GCGTGGGCACT (chr19:39540166, +, 5.16 bits)	Abolished	√	
				rs887888062 (chr19:39540176T>A)	GCGTGGGCGCA (chr19:39540166, +, 10.94 bits)	Cluster 2 of 2	√	
ELF1 (R_sequence = 11.2057 bits)	HIST1H4H	Cluster 1 of 2	GCGGAAGCGTG (chr6:26286540, +, 9.92 bits)	rs760968937 (chr6:26286547C>T)(chr6:26286547C>A)	GCGGAAGTGTG (chr6:26286540, +, 10.71 bits)	Cluster 1 of 2	√	√
					GCGGAAGAGTG (chr6:26286540, +, 8.84 bits)	Cluster 1 of 2	√	
				rs1000196206 (chr6:26286542G>C)	GCCGAAGCGTG (chr6:26286540, +, -6.26 bits)	Abolished	√	
				rs144759258 (chr6:26286543G>A)	GCGAAAGCGTG (chr6:26286540, +, -3.60 bits)	Abolished	√	×
				rs966435996 (chr6:26286544A>G)	GCGGGAGCGTG (chr6:26286540, +, 5.28 bits)	Abolished	√	
				rs950986427 (chr6:26286548G>A)	GCGGAAGCATG (chr6:26286540,	Cluster 1 of 2	√	

TF	Gene	Cluster	Binding site	rs ID	Variant	Result			
					+, 8.28 bits)				
		Cluster 2 of 2	C*AGGAG*ATGC*G* (chr6:26286473, -, 6.98 bits)	rs373649904 (chr6:26286483G>A)	*T*AGGAGATGCG (chr6:26286473, -, 0.61 bit)	Abolished	√		
				rs926919149 (chr6:26286480C>T)	CAG*A*AGATGCG (chr6:26286473, -, -6.53 bits)	Abolished	√		
				rs751263172 (chr6:26286479T>G)	CAGG*C*GATGCG (chr6:26286473, -, 1.24 bits)	Abolished	√		
				rs369076253 (chr6:26286473C>G)	CAGGAGATGC*C* (chr6:26286473, -, 6.92 bits)	Cluster 2 of 2	√		
				<u>rs751263172 (chr6:1044474314C>T)</u>	<u>CAGGA*A*ATGCG (chr6:26286473, -, 11.43 bits)</u>	Cluster 2 of 2	√	√	
ELK1 (R_sequence = 11.9041 bits)	G0S2	Cluster 1 of 2	C*AG*GGAAG*A*CC (chr1:209667959, -, 1.92 bits)	<u>rs146048477 (chr1:209667961T>A)</u>	<u>CAGGGAAG*T*CC (chr1:209667959, -, 2.24 bits)</u>	Cluster 1 of 2	√	√	
				rs887606802 (chr1:209667968T>C)	C*G*GGGAAGACC (chr1:209667959, -, -3.35 bits)	Cluster 1 of 2	√		
				rs1021034916 (chr1:209667967C>T)	CA*A*GGAAGACC (chr1:209667959, -, -3.57 bits)	Cluster 1 of 2	√		
			GAGGA*A*ATGAG (chr1:209667969, +, 8.14 bits)	rs941962117 (chr1:209667974A>G)	GAGGA*G*ATGAG (chr1:209667969, +, 4.11 bits)	Abolished	√		
		Cluster 2 of 2	*C*TGGAAGA*GC*A (chr1:209673544, -, 5.91 bits)	rs896117033 (chr1:209673545G>A)	CTGGAAGAG*T*A (chr1:209673544, -, 3.95 bits)	Cluster 2 of 2	√	×	√
				rs971962577 (chr1:209673546C>T)	CTGGAAGA*A*CA (chr1:209673544, -, 3.49 bits)	Cluster 2 of 2	√		
				rs1011969709 (chr1:209673554G>C)	*G*TGGAAGAGCA (chr1:209673544, -, 3.92 bits)	Abolished	√		
			CCA*G*AAGTCA*A* (chr1:209673551, +, 7.44 bits)		CCA*C*AAGTCAA (chr1:209673551, +, -5.50 bits)	Abolished	√		
				<u>rs1023312090 (chr1:209673561A>G)</u>	<u>CCAGAAGTCA*G* (chr1:209673551, +, 8.40 bits)</u>	Cluster 2 of 2	√	√	
ETS1 (R_sequence = 10.0788 bits)	TTC19	Cluster 1 of 1	GCA*G*GGAA*A*GG (chr17:16022293, +, 7.92 bits)	rs1022234223 (chr17:16022296G>C)	GCA*C*GGAAAGG (chr17:16022293, +, -4.98 bits)	Abolished	×	×	√
				<u>rs968299415 (chr17:16022301A>T)</u>	<u>GCAGGGAA*T*GG (chr17:16022293, +, 10.01 bits)</u>	Cluster 1 of 1	√	√	
GABPA (R_sequence =	PLEKHB2	Cluster 1 of 1	A*CAG*GAAAGGG (chr2:131112770,	rs997328042 (chr2:131112771C>T)	A*T*AGGAAAGGG (chr2:131112770,	Abolished	×	×	√

10.8567 bits)			+, 10.36 bits)		+, -3.68 bits)				
				rs1020720126 (chr2:131112773G>C)	ACACGAAAGGG (chr2:131112770, +, -4.16 bits)	Abolished	×		
			TCTGGAAACTA (chr2:131112760, +, 1.53 bits)	rs185306857 (chr2:131112761C>A)	TATGGAAACTA (chr2:131112760, +, -2.86 bits)	Cluster 1 of 1	√		
				rs772728699 (chr2:131112762T>A)	TCAGGAAACTA (chr2:131112760, +, 5.23 bits)	Cluster 1 of 1	√		
				rs965753671 (chr2:131112769T>C)	TCTGGAAACCA (chr2:131112760, +, 2.13 bits)	Cluster 1 of 1	√		
IRF1 ($R_{sequence}$ = 13.5544 bits)	SMIM13	Cluster 1 of 1	GAGAATGAAAGCA (chr6:11093663, +, 12.56 bits)	rs950528541 (chr6:11093663G>C)	CAGAATGAAAGCA (chr6:11093663, +, 8.97 bits)	Cluster 1 of 1	√		√
				rs886259573 (chr6:11093664A>G)	GGGAATGAAAGCA (chr6:11093663, +, 9.65 bits)	Cluster 1 of 1	√		
				rs982931728 (chr6:11093666A>G)	GAGGATGAAAGCA (chr6:11093663, +, 8.09 bits)	Cluster 1 of 1	√		
				rs1020218811 (chr6:11093668T>G)	GAGAAGGAAAGCA (chr6:11093663, +, 9.36 bits)	Cluster 1 of 1	√		
				rs570723026 (chr6:11093672A>G)	GAGAATGAAGGCA (chr6:11093663, +, 8.01 bits)	Cluster 1 of 1	√	×	
				rs1004825794 (chr6:11093675A>C) (chr6:11093675A>T)	GAGAATGAAAGCC (chr6:11093663, +, 10.47 bits)	Cluster 1 of 1	√		
					GAGAATGAAAGCA (chr6:11093663, +, 10.42 bits)	Cluster 1 of 1	√		
			AAGACCAAAGGCA (chr6:11093641, +, 2.43 bits)	rs1030185383 (chr6:11093649A>C)	AAGACCAACGGCA (chr6:11093641, +, -3.39 bits)	Cluster 1 of 1	√		
				rs5874306 (chr6:11093650delG)	AAGACCAAAGCAG (chr6:11093641, +, 0.90 bit)	Cluster 1 of 1	√		
				rs558896490 (chr6:11093643G>A)	AAAACCAAAGGCA (chr6:11093641, +, 7.06 bits)	Cluster 1 of 1	√	√	
YY1 ($R_{sequence}$ = 12.8554 bits)	CKLF	Cluster 1 of 1	GCGGCCATCGGC (chr16:66549785, -, 10.06 bits)	rs865922947 (chr16:66549791G>A)	CCGGCCATCGGC (chr16:66549785, -, 6.80 bits)	Cluster 1	√		√
				rs946037930 (chr16:66549794C>A)	GCTGCCATCGGC (chr16:66549785, -, 8.02 bits)	Cluster 1	√	×	
				rs917218063 (chr16:66549793C>T)	GCGACCATCGGC (chr16:66549785, -, 5.41 bits)	Abolished	×		

				rs928017336 (chr16:66549791G>A)	GCGGC*T*ATCGGC (chr16:66549785, -, -3.62 bits)	Abolished	×		
			GCCGCCCCCGTC (chr16:66549792, +, 1.34 bits)						

§ All coordinates are based on the hg38 genome assembly. A bold italic letter in a binding site

sequence indicates the base where a SNP occurs. The SNPs strengthening binding sites and

corresponding variant binding site sequences are underlined.

‡ The impact on whether the occurrence of a single SNP resulted in the disappearance of the

cluster containing it is shown.

† After a single SNP occurred or multiple SNPs simultaneously occurred, the classifier produced

a new prediction on whether the TF is still capable of significantly affecting gene expression via

the variant promoter.

A

For a specific gene A / each TF

```
Obtain gene expression data    Details in Panels
                               B, C and D
        ↓
Obtain positives and negatives
```

For each gene instance

```
Obtain the union of 10kb promoters upstream of all TSSs
        ↓
Intersect these promoters with DNase I hypersensitive sites
        ↓
Scan the (overlapping) promoter intervals for TFBSs
using 94 iPWMs of 82 TFs / one iPWM of this TF
        ↓
Detect TFBS clusters using the IDBC algorithm ($d=25$, $l=939$)
        ↓
Extract features from the clusters in the 10kb promoter of the longest transcript
```

```
Machine learning classifiers (e.g. SVM, Decision tree, Naïve Bayes)
        ↓
Performance evaluation
```

B Obtain expression profiles of 56,238 genes from GTEx

For a specific gene A

```
Compute the Bray-Curtis similarity value between
$A$'s profile and every other gene's profile
```

```
Select the 500 PC genes          Select the 500 PC genes
with the largest similarity      with the smallest similarity
values as positives              values as negatives
```

C Obtain the CRISPR-generated regulatory matrix

For each TF

```
Select the PC genes meeting      Select the PC genes meeting
three criteria as positives      one criterion as negatives
```

```
Find the TF $B$ with the largest average correlation
        ↓
Compute the Bray-Curtis similarity value between
$B$ and each negative based on expression profiles
        ↓
Select a subset of negatives with the smallest similarity
values whose size is the number of positives
```

D Obtain the siRNA-generated P-value matrix

For each TF

```
Select the PC genes meeting      Select the same number of
two criteria as positives        PC genes meeting one
                                 criterion as negatives
```

The threshold c on the average fold change

Legend: ⊔ : A binding site of EGR1   ⬦ : An EGR1 molecule   ✔/✘ : Is / is not a DE target of EGR1

Cluster 1 — 12.0 bits, 4.3 bits; Cluster 2 — 7.0 bits, 1.3 bits; Cluster 3 — 1.9 bits, 8.9 bits, 2.2 bits ✔

GAGGG**G**GAGTG → GAGGG**A**GAGTG

Cluster 2 — 7.0 bits, 1.3 bits; Cluster 3 — 1.9 bits, 8.9 bits, 2.2 bits; 4.3 bits ✔

GT**G**GTGGCGGG → GT**T**GTGGCGGG   CGGGAGG**C**GGG → CGGGAGG**G**GGG

4.3 bits, 1.3 bits, 1.9 bits, 2.2 bits ✘