Genome-wide fitness assessment during diurnal growth reveals an expanded role of the cyanobacterial circadian clock protein KaiA

David G. Welkie[1], Benjamin E. Rubin[2], Yong-Gang Chang[3], Spencer Diamond[2,4], Scott A. Rifkin[2], Andy LiWang[3], Susan S. Golden[1,2]

[1]Center for Circadian Biology, University of California, San Diego, La Jolla, CA; [2]Division of Biological Sciences, University of California San Diego, La Jolla CA; [3]School of Natural Sciences, University of California, Merced, CA; [4]Department of Earth and Planetary Science, University of California, Berkeley, CA.

Abstract

The recurrent pattern of light and darkness generated by Earth's axial rotation has profoundly influenced the evolution of organisms, selecting for both biological mechanisms that respond acutely to environmental changes and circadian clocks that program physiology in anticipation of daily variations. The necessity to integrate environmental responsiveness and circadian programming is exemplified in photosynthetic organisms such as cyanobacteria, which depend on light-driven photochemical processes. The cyanobacterium *Synechococcus elongatus* PCC 7942 is an excellent model system for dissecting these entwined mechanisms. Its core circadian oscillator, consisting of three proteins KaiA, KaiB, and KaiC, transmits time-of-day signals to clock-output proteins, which reciprocally regulate global transcription. Research performed under constant light facilitates analysis of intrinsic cycles separately from direct environmental responses, but does not provide insight into how these regulatory systems are integrated during light-dark cycles. Thus, we sought to identify genes that are specifically necessary in a day-night environment. We screened a dense bar-coded transposon library in both continuous light and daily cycling conditions and compared the fitness consequences of loss of each nonessential gene in the genome. Although the clock itself is not essential for viability in light-dark cycles, the most detrimental mutations revealed by the screen were those that disrupt KaiA. The screen broadened our understanding of light-dark survival in photosynthetic organisms, identified unforeseen clock-protein interaction dynamics, and reinforced the role of the clock as a negative regulator of a night-time metabolic program that is essential for *S. elongatus* to survive in the dark.

Significance

Understanding how photosynthetic bacteria respond to and anticipate natural light–dark cycles is necessary for predictive modeling, bioengineering, and elucidating metabolic strategies for diurnal growth. Here, we identify the genetic components that are important specifically under light-dark cycling conditions and determine how a properly functioning circadian clock prepares metabolism for darkness, a starvation period for photoautotrophs. This study establishes that the core circadian clock protein KaiA is necessary to enable rhythmic de-repression of a night-time circadian program.

Specific Contributions

DGW BR SD and SSG conceived and designed the project. DGW BR and YGC performed the experiments and analyzed the data. SAR wrote the R scripts for RB-TnSeq conditional fitness analysis. YGC and AL interpreted fluorescence anisotropy data. DGW BR SSG YGC and AL wrote the manuscript.

\body

Introduction

Photosynthetic organisms experience a dramatic change in physiology and metabolism each day when the sun sets and the daytime processes related to photosynthesis become inoperative. Like many diverse organisms throughout nature, prokaryotic cyanobacteria have evolved circadian clocks that aid in the temporal orchestration of activities that are day- or night-appropriate. The circadian clock's contribution towards fitness in changing environments was first demonstrated in 1998, when it was shown that strains of the cyanobacterium *Synechococcus elongatus* PCC 7942 with intrinsic circadian periods that match that of external light-dark cycles (LDC) outcompete strains that have different periods (1, 2). More recent experiments showed the value of preparing for night: when cells synchronized to different phases of the circadian cycle are mixed and then exposed to a pulse of darkness, those whose internal clock corresponds to a dawn or day-time phase have a higher occurrence of arrested growth upon re-illumination than those synchronized to anticipate darkness at the time of the pulse (3). These data support the hypothesis that the clock acts to prepare the cell for conditions that are predictable and recurrent, such as night following day.

A fitness advantage of appropriate circadian timing signals is not exclusive to cyanobacteria, as is observed in the model plant *Arabidopsis* (4, 5) and in humans, where circadian disruption can result in myriad adverse health effects including cardiovascular disease, cancer, and sleep disorders (6–8). Although the fundamental properties of circadian rhythms are shared across the domains of life, the molecular networks that generate them vary (9). In *S. elongatus* a three-protein oscillator comprising the proteins KaiA, KaiB, and KaiC sets up a circadian cycle of KaiC phosphorylation. This oscillator even functions *in vitro* when mixed at appropriate ratios with ATP (10), which has provided a powerful tool for exploring the mechanism that generates circadian rhythms. Genetic, biochemical, metabolic, and structural inquiries have revealed that the *S. elongatus* oscillator actively turns off a night-time metabolic program prior to dawn by triggering the dephosphorylation of the master transcription factor Regulator of Phycobilisome Association A (RpaA). It accomplishes this task by recruiting Circadian Input Kinase A (CikA) to a ring of KaiB that forms on KaiC during the night-time portion of the cycle, thus activating CikA phosphatase activity against RpaA (11–13). Among the targets of the RpaA regulon are the genes that facilitate catabolism of glycogen and generation of NADPH via the oxidative pentose phosphate pathway (14). These enzymatic reactions are essential for the cyanobacterium to survive the night when photosynthetic generation of reductant is disabled (15, 16).

While the molecular mechanisms of the circadian clock are well established in constant light for *S. elongatus* (17), its role in response to LDC, and the mechanisms by which the cell integrates circadian and environmental data, are still not understood. There have been few publications on the topic, and this study is the first to report on a genome-wide screen for LDC-sensitive mutants. Studies that report conditional LDC effects have been limited to a small number of genes. For instance, mutants defective for the circadian clock output pathway genes, *rpaA* and *sasA* (15, 16, 18, 19), and mutants unable to mobilize carbon through the oxidative pentose phosphate pathway (OPPP) and glycogen breakdown (16, 20, 21), or unable to synthesize the alarmone nucleotide ppGpp (22, 23), have conditional defects specific to growth in LDC. To identify the genetic contributors to fitness under LDC and acquire a more comprehensive picture of the regulatory network, we used an unbiased population-based screen that tracks the fitness contributions of individual *S. elongatus* genes to reproduction in diel cycles. The method, random bar code transposon-site sequencing (RB-TnSeq), quantitatively compares the changes in abundances of individuals in a pooled mutant population over time. This assay allowed the calculation of a fitness contribution to LDC survival for each nonessential gene in the genome by leveraging multiple inactivating mutations spread across a genomic locus. This library was previously used to identify the essential gene set of *S. elongatus* under standard laboratory

continuous light conditions (CLC) (24) and to generate an improved the metabolic model for
generating phenotype predictions (25).

The results described here provide a comprehensive picture of genes that contribute to fitness
under diurnal growth conditions. In addition to identifying the genetic components previously
known to be critical for growth in LDC we found ~100 new genes whose mutation confers weak
to strong fitness disadvantages. Of these genes, we focused on the top-scoring elements for
LDC-specific defects which, unexpectedly, lie within the *kaiA* gene of the core circadian
oscillator. As no previously known roles for KaiA predicted this outcome, we characterized the
mechanism that underlies the LDC sensitivity of a *kaiA*-null mutant, and revealed interactions
within the oscillator complex that regulate the activation of CikA, an RpaA phosphatase.

Results

Growth of library mutants to assess genetic fitness contributions under light-dark cycles.

The RB-TnSeq mutant library includes a unique identifier sequence, or bar code, at each
insertion site to track each transposon-insertion mutant. Next-generation sequencing is used to
quantify these bar codes within the total DNA of the population, such that the survival and
relative fitness of the approximately 150,000 mutants in the library can be tracked under control
and experimental conditions (26). In this way, the *S. elongatus* RB-TnSeq library can be used
for pooled, quantitative, whole-genome mutant screens.

First, the library was thawed, placed into 250-mL flasks, and sampled during incubation in CLC
under only the selective pressure for the transposon antibiotic-resistance marker. Genomic DNA
was extracted, and the bar codes of the mutant population were sequenced to establish a
baseline abundance (**Fig. 1**). The culture was then divided into photobioreactors set to maintain
constant optical density under either CLC, or LDC conditions that mimic day and night. After a
treatment period allowing for approximately 6 generations, the library was again sampled and
bar codes were quantified in a similar fashion. By comparing the frequencies of the bar codes
after growth in LDC to those in CLC (**Dataset S1**), we estimated a quantitative measure of the
fitness contribution for each gene specifically under LDC (**Dataset S2**).

After filtering the data as described in the Materials and Methods section we were able to
analyze insertions in 1,872 of the 2,005 non-essential genes (24) in the genome. Each fitness
score that passed our statistical threshold for confidence (FDR <1%) was classified as having a
"strong" (>1.0 or <-1.0), "moderate" (between 1.0 and 0.5 or -1.0 and -0.5) , or "negligible /
minor" (between 0 and ±0.5) effect on fitness in LDC. This analysis revealed 90 loci whose
disruption caused strong changes to LDC fitness (41 negatively and 49 positively) and an
additional 130 loci that had moderate effects (**Fig. 2A**). A volcano plot showing the data with
selected genes highlighted in this study is displayed in **Fig. 2B**. Because our initial goal was to
understand the suite of genes required to survive the night, we focused first on those mutants
whose loss specifically debilitates growth in LDC.

Metabolic pathways important for LDC growth

The functional composition of genes found to be either detrimental or favorable to fitness is
indicative of the metabolic processes that are important for growth in LDC (**Fig. 2C**). Genes
identified as necessary in LDC encode enzymes that participate in carbohydrate and central
carbon metabolism, protein turnover and chaperones, CO_2 fixation, and membrane biogenesis.
Strikingly, all genes of the OPPP, *zwf*, *opcA*, *pgl*, and *gnd,* have strong fitness effects (**Fig. 3A**).
In addition, *tal*, the only non-essential gene of the reductive phase, is also essential for LDC
growth. It is known that *tal* mRNA peaks at dusk and that its product acts in a reaction that is
shared between the oxidative and reductive phases of the PPP, linking it to glycolysis; the *tal*
mutant's sensitivity to LDC supports the hypothesis that Tal is a less recognized, but important,

component of the OPPP reactions. Furthermore, all OPPP genes have peak expression at dusk
(27).

Disruptions in genes that encode enzymes responsible for glycogen metabolism previously
have been shown to cause LDC sensitivity (21). Our screen clearly identified the genes for
catabolic glycogen metabolism *glgX* and *glgP* as important LDC fitness factors (**Fig. 2B**).
However, the importance of glycogen metabolism in LDC fitness is complicated by evidence
that the anabolic reactions of *glgA* and *glgC* contribute to fitness in CLC as well, due to the
utilization of glycogen as a sink for excess photosynthetically derived electrons (28, 29). Thus,
for genes critical for glycogen anabolism, the differential LDC fitness score is not as striking as
might be expected from phenotypic assays of those mutants. The *glgA* mutants were classified
as having a minor LDC defect, and insufficient reads for *glgC* removed this locus from our
analysis pipeline, leading to a classification of "not determined". Despite this complication,
mutations in the gene *glgB,* whose product catalyzes the last non-essential step in glycogen
synthesis, caused a strong decrease in LDC fitness, supporting the conclusion that both
glycogen synthesis and breakdown are critical aspects of metabolism in LDC conditions.
Together, the reactions facilitated by these enzymes provide a major source of reductant
(NADPH) in the dark and facilitate the flow of glucose compounds via the breakdown of stored
glycogen into downstream metabolites.

Genes involved in DNA repair (such as the transcription-repair coupling factor *mdf*) and
chaperone systems (*clpB1*) are also important for light-dark growth. ClpB1 is known to
contribute to thermotolerance stress and is induced upon dark-light transitions in cyanobacteria
(30, 31). The importance of these genes for surviving light-dark cycles suggests a role in
managing light-induced stress that the transition from dark to light may impose. Moreover,
mutation of any of the three genes that encode enzymes of the glycine cleavage system confer
increased LDC fitness. The glycine decarboxylase complex (GDC) is linked to photorespiratory
mechanisms that respond to high-light stress (32, 33). This result further supports our finding
that variations in the fitness of metabolism genes between CLC and LDC are linked to pathways
that mitigate light stress.

In addition to underscoring the importance of the OPPP, glycogen metabolism, and repair
mechanisms in LDC, we also found a strong link between these metabolic pathways and
circadian clock management. Many of the genes essential for growth in LDC are regulated by
the clock output protein RpaA. For instance, the circadian clock gene *kaiC* is itself regulated by
RpaA and its expression is reduced by over 50% in an *rpaA*-null background (12, 14). This
attribute is also shared by the core glycogen metabolism and PPP genes *glgP*, *zwf*, *opcA*, *gnd*,
and *tal,* and genes involved with metal ion homeostasis (*smtA*) and cobalamin biosynthesis
(*cobL*), all of which show significantly reduced transcripts (<-1.0 log2(*ΔrpaA*/WT, wild type) in
the absence of the RpaA (14). These core glycogen metabolism and PPP genes , with the
exception of *tal* and *gnd,* are also direct targets of the transcription factor. This finding is a
common trend in mutants that show strong fitness decreases in LDC, as ~20% (9/41) have
dramatic reductions in mRNA levels in an *rpaA*-null strain. This trend is not the case for any of
the genes identified as mutants that confer increased fitness (**Fig 3B**). These findings support a
growing body of evidence that the ability of *S. elongatus* to initiate a night-time transcriptional
program, and particularly to activate the OPPP pathway genes, is critical for survival through
darkness (15, 16).

Fitness contributions of circadian clock constituents

The mechanism of circadian timekeeping and regulation in cyanobacteria is well understood at
the molecular level (17, 34). The three core clock proteins, KaiA, KaiB, and KaiC orchestrate
circadian oscillation in *S. elongatus* (35)*,* whereby KaiC undergoes a daily rhythm of

phosphorylation and dephosphorylation that provides a time indicator to the cell (36, 37). During
the day period, KaiA stimulates KaiC autophosphorylation, and, during the night, KaiB opposes
KaiA's stimulatory activity by sequestering an inactive form of KaiA (38), leading to KaiC
autodephosphorylation. Two histidine kinase proteins, SasA and CikA, engage with the
oscillator at different phases of the cycle, and this interaction stimulates SasA kinase activity to
phosphorylate RpaA and later stimulates CikA phosphatase activity to dephosphorylate RpaA.
Thus, phosphorylated RpaA accumulates during the day, peaking at the day-night transition,
and activates the night-time circadian program.

Mutations that caused the most serious defect specifically in LDC were mapped to the gene that
encodes the clock oscillator component KaiA (**Fig. 2B**). This outcome was unexpected because
the known roles of KaiA are to stimulate phosphorylation of KaiC and to enable normal levels of
KaiC and KaiB expression (38, 39). Thus, a KaiA-less strain, in which KaiC levels are low and
hypophosphorylated (39), was expected to behave like a *kaiC* null. A *kaiC* null mutant has no
notable defect in LDC when grown in pure culture (1), and even a *kaiABC* triple deletion,
missing KaiA as well as KaiB and KaiC, grows well in LDC (18).

Nonetheless, a *kaiC* null mutant is known to be outcompeted by WT under LDC conditions in
mixed culture (1, 2), which is the situation in this population-based screen. Mutations in *kaiC*
returned negative fitness scores in the current screen, as did bar codes associated with *rpaA*
and *sasA*, two clock-related genes that are known to be important for growth in LDC (18, 20).
RpaA is important for redox balancing, metabolic stability, and carbon partitioning during the
night, and an *rpaA* mutant is severely impaired in LDC (15, 16, 20). While it is identified in the
current screen as LDC sensitive, inactivation of *rpaA* also confers a moderate fitness decrease
under CLC (~-1.0), thus diminishing the differential LDC-specific fitness calculation.

Disruptions to other genes related to circadian clock output scored as beneficial to growth in
LDC, increasing the fitness of the strains against the members of the population that carry an
unmutated copy. For instance, mutations within the gene that encodes CikA showed improved
fitness, along with disruptions to the sigma factor encoded by *rpoD2* (40). CikA has several
roles that affect the clock, including synchronization with the environment and maintenance of
WT period; additionally, CikA is stimulated to dephosphorylate RpaA by association with the
oscillator at night (13, 41). Thus, a phenotype for *cikA* mutants that is opposite to that of *sasA*
mutants forms a consistent pattern.

From the published biochemical data we would predict that, in a *kaiA* null, SasA kinase would
still be activated by the low levels of unphosphorylated KaiC but that CikA phosphatase would
not be: CikA binds not to KaiC, but to a ring of KaiB that engages hyperphosophorylated KaiC, a
state that is not achieved in the absence of KaiA. Therefore, we would expect that RpaA would
be constitutively phosphorylated in a *kaiA* null mutant and that the cell would be fixed in the
night-time program, where it would be insensitive to LDC. We further investigated the *kaiA*-
mutant phenotype under LDC to test the validity of the RB-TnSeq results.

Growth attenuation and rescue of a kaiA mutant. The strong negative scores of bar codes
for *kaiA* insertion mutants pointed to a severe growth defect in LDC. In order to vet this finding
we tested the phenotype of an insertional knockout of the *kaiA* gene (*kaiA*[ins]), similar to the
strains in the RB-TnSeq library. Assays in CLC vs LDC verified the LDC-specific defect in the
mutant (**Fig. 4A**). Because *kaiABC* deletion mutants and *kaiC* deletion mutants both do not
have an LDC defect in pure culture, the LDC defect seemed to be specific for situations in which
KaiC and KaiB are present without KaiA. The *kaiA* coding region has a known negative element
located within its C-terminal coding region that negatively regulates expression of the
downstream *kaiBC* operon (42, 43) (*kaiA*[ins]), keeping KaiBC levels low. We reasoned that the
loss of KaiA would be exacerbated if expression of KaiBC increased. To test this hypothesis, we

measured the growth characteristics of a *kaiA* mutant in which the coding region of *kaiA*,
including the negative element, has been removed and replaced with a drug marker (*kaiA^del^*)
(39). This strain, which is known to have normal or slightly elevated levels of KaiB and KaiC,
exhibited even more dramatic sensitivity to LDC and supported the hypothesis that an
unbalanced clock output signal underlies the LDC sensitivity in the *kaiA* mutant.

Knowing that *rpaA*-null mutants, which cannot activate the OPPP genes, have similarly severe
LDC phenotypes, we hypothesized that the *kaiA*-less strains have insufficient phosphorylation of
RpaA to turn on those genes. We reasoned that additionally interrupting the *cikA* gene, whose
product dephosphorylates RpaA in a clock-dependent manner, would help to conserve RpaA
phosphorylation in the *kaiA^del^* strain and increase LDC fitness. Indeed, mutations that disrupt
cikA were identified from the population screen as enhancing fitness (**Fig. 2B),** a finding we
confirmed in a head-to-head competitive growth assay in a mixed culture with WT (**Fig. 4B**).
Complementation of this double mutant with a *cikA^WT^* gene, or the *kaiA^del^* strain with *kaiA^WT^*
expressed from a neutral site in the genome, resulted in a reversal of the respective phenotypes
(**Fig. S1**). The LDC growth defects in *kaiA^del^* and improvement of the *kaiA^del^cikA* strain were
reproducible when grown in liquid cultures in the same photobioreactors used to grow the library
for the LDC screen (**Fig. 4C**). These results are consistent with a model in which, in the
absence of KaiA, CikA is constitutively activated as a phosphatase to dephosphorylate RpaA.

Arrhythmic and low* kaiBC *expression correlates with LDC phenotype. Hallmarks of RpaA
deficiency or dephosphorylation are low and arrhythmic expression from the *kaiBC* locus under
CLC, as visualized using a luciferase-based reporter (P*~kaiBC~::luc*) (12). We used this assay to
determine whether the severity of defect in LDC growth in mutants correlates with level of
expression from this promoter as a proxy for RpaA phosphorylation levels. Consistent with the
hypothesis, the *kaiA^ins^* mutant has low constitutive levels of luciferase activity, tracking near the
troughs of WT circadian expression, the *kaiA* deletion mutant has almost undetectable
expression, and the constitutive expression from the *kaiABC^del^* strain tracks near peak WT
levels (**Fig. 5**). Moreover, the *kaiA^del^cikA* strain shows expression levels much higher than the
*kaiA^del^* strain, approaching those observed in the *kaiABC^del^* mutant and around the midline of
WT oscillation. Intermediate levels of growth in LDC between the *kaiA^del^* and *kaiA^del^cikA* strains
were observed when a *cikA* variant (C644R) that is predicted to have reduced KaiB binding, and
thus less phosphatase activation, was expressed in *trans* in the *kaiA^del^cikA* strain (11). Growth
of this strain also displayed intermediate LDC sensitivity (**Fig. 4A**). These data support a
correlation between expression from the P*~kaiBC~::luc* reporter strain, likely reflecting RpaA
phosphorylation levels, and growth in LDC.

***Low RpaA phosphorylation in* kaiA*-null cells as the cause of decreased fitness*.** We next
directly assayed the RpaA phosphorylation levels in this group of mutants by using immunoblots
and Phos-tag reagent to separate phosphorylated (P-RpaA) from unphosphorylated RpaA. As
predicted, P-RpaA was low in the *kaiA^ins^* mutant and almost undetectable in the *kaiA^del^* strain
(**Fig. 6A**). Adding an ectopic WT *kaiA* allele, or disrupting *cikA*, in the *kaiA^del^* mutant restored
RpaA phosphorylation. Thus, even though adequate KaiC is present to engage SasA to
phosphorylate RpaA (**Fig. 5B**), the absence of KaiA is sufficient to suppress RpaA
phosphorylation. These results support the conclusion that the rescue of the *kaiA* growth defect
by disrupting *cikA* is due to the elimination of CikA phosphatase activity.

Because the model suggests that turning on RpaA-dependent genes at dusk is critical for
survival in LDC, we measured P-RpaA specifically at light-to-dark and dark-to-light transitions in
*kaiA^del^* and *kaiA^del^cikA* over three diel cycles. WT has a marked pattern of high P-RpaA at dusk
and low levels at dawn (**Fig. 6C**). The *kaiA^del^* strain had almost undetectable levels of P-RpaA,
visible only when the overall RpaA signal was very high. The P-RpaA phosphorylation status in

the *kaiA^del^cikA* was restored but did not show diel variations, consistent with the arrhythmicity of
kaiA mutants.

Other physiological similarities between the *kaiA*-null and *rpaA*-null strains exist. An *rpaA*
mutant accumulates excessive reactive oxygen species (ROS) during the day that it is unable to
alleviate during the night (15). Metabolomic analysis points towards a deficiency in reductant,
needed to power detoxification reactions for photosynthetically-generated ROS, due to lack of
NADPH production in the dark when OPPP genes are not activated. Likewise, levels of ROS in
*kaiA^del^* were significantly higher at night than in WT and *kaiA^del^cikA* (**Fig. 6D**).

Molecular basis of CikA engagement in the absence of KaiA. The proposal that the LDC
fitness defects in *kaiA* mutants result from excessive CikA phosphatase activity was derived
from genetic data related to known LDC-defective mutants. However, previous studies
regarding KaiA activities had not predicted this outcome. Very recent structural data, enabled
through the assembly of KaiABC and KaiBC-CikA complexes, pointed to a likely mechanism:
loss of competition for a binding site on KaiB that KaiA and CikA share (11). Although this
hypothesis was appealing, the current model of progression of the Kai oscillator cycle could not
predict this outcome. KaiA and CikA bind to a ring of KaiB that forms after KaiC has become
phosphorylated at Serine 431 and Threonine 432 (44). Maturation to this state induces
intramolecular changes in KaiC that expose the KaiB binding site. In the absence of KaiA, KaiC
autophosphorylation is extremely abated as measured both *in vivo* (45) and *in vitro* (1-10%, **Fig.
S5C**), so how can KaiB bind?

We used fluorescence anisotropy to directly determine whether KaiB can engage with KaiC in
the absence of KaiA, when KaiC is mainly unphosphorylated (**Fig. 7** and **Fig. S5**). If KaiC-KaiB
binding can occur, it could support binding of CikA – expected to be a necessary event for CikA
phosphatase activation. This method measures the magnitude of polarization of fluorescence
from KaiB tagged with a fluorophore (6-iodoacetamidofluorescein), which increases when KaiB
becomes part of a larger complex. KaiC phosphomimetic mutants were used to quantify
interaction of KaiB with each of the phosphostates of KaiC. Initially, we hypothesized that the
KaiB binding site might become exposed on KaiC in the absence of phosphorylation when cells
enter the dark and the ATP/ADP ratio is altered (46). For this reason, experiments were
performed at different ATP/ADP ratios. As a control, we confirmed that no significant association
of KaiB with CikA occurs in the absence of KaiC (**Fig. 7A**). However, KaiB is able to associate
with hypophosphorylated WT KaiC approximately as well as with the phosphomimetic, KaiC-EA,
which mimics a form of KaiC (KaiC-pST) that binds KaiB, which then forms a new ring on the N-
terminal face of KaiC that captures KaiA and CikA. This binding was similar in both 1 mM ATP
and 0.5 mM ATP+0.5 mM ADP conditions (**Fig. 7B**), suggesting that darkness (associated with
decreased ATP/ADP *in vivo*) is not required to enable KaiB binding to KaiC. Moreover, CikA
associated with KaiBC under both conditions. Association of KaiB was minimal with KaiC-AE, a
mimic of the daytime state (KaiC-SpT) when KaiA is usually well-associated with the A-loops of
KaiC in an interaction that does not require KaiB. However, the presence of CikA enhanced the
binding of KaiB to KaiC-AE, presumably due to formation of a KaiC-KaiB-CikA complex. For the
WT KaiC experiments in **Fig. 7**, KaiB (0.05 µM) is most likely binding to the small percentage (1-
10%) of the KaiC population that remains phosphorylated (0.04-0.4 µM). *In vivo*, it is likely that
in the absence of KaiA, KaiB also binds selectively to a phosphorylated fraction of the
hypophosphorylated KaiC pool.

These data generally match the patterns expected for the current oscillator model; however,
they also demonstrate that the KaiC-KaiB-CikA complex can form in the absence of KaiA, when
the KaiC pool is hypophosphorylated, and that the ATP ratio (mimicking day and night) has little
effect on this association. Furthermore, they support the competitive binding of KaiA and CikA to
the KaiB ring. These observations are consistent with the proposal that the low survival rate of

kaiA-null mutants is due to over-stimulation of CikA phosphatase activity, stripping the cell of P-
RpaA and disabling expression of the critical night-time reductant-producing OPPP.

Discussion

Recent research and interest in growth of cyanobacteria in LDC conditions (15, 16, 20, 22, 47–
49) provides a rich dataset to inform engineering strategies, to model metabolic flux predictions,
and to elucidate the intersection of the clock and cell physiology. While previous studies have
identified a handful of genes important for growth in LDC, the conditional defect of these
mutants was determined from targeted studies that left large unknowns in the pathways
important for LDC. This study used an unbiased quantitative method to comprehensively query
the *S. elongatus* genome for the full set of genes that are specifically necessary to survive the
night. This approach eliminated the requirement to test individual mutants for a scorable
phenotype under diel conditions and allowed for the discovery of genes that would not be
predicted to have such a phenotype. In addition to identifying genes of pathways previously
unknown to be involved in LDC, this screen also strongly reinforced two emerging stories from
recent research: the breakdown of glycogen and flux through the OPPP, the major source of
reductant under non-photosynthetic conditions, is essential (15, 16); and, the role of the clock is
repressive, rather than enabling (12), as its complete elimination has little consequence in LDC,
but its constitutive output signaling is harmful under these conditions because of its negative
effect on the OPPP genes.

A boon of the Rb-TnSeq data is in how it reveals non-intuitive insights into the regulatory
measures cells need for growth in specific conditions. One such discovery is that a *kaiA* mutant
is severely LDC sensitive. Investigation of the mechanism that underlies this phenotype led to
increased understanding of the function of the Kai oscillator, a nanomachine whose structural
interactions have been revealed with an unusual degree of clarity (11, 50). KaiA is
conventionally thought to have a specific daytime role in stimulating KaiC autophosphorylation –
an activity that promotes SasA stimulation and brings about the conformational change in KaiC
that enables KaiB engagement (**Fig. 8***A*). This latter step ushers in the KaiC dephosphorylation
phase, during which KaiA is a passive, inactive player. Results from this study provide a new
perspective of KaiA in its KaiB-bound form as regulating the night-time clock output by its
competition with CikA for KaiB-binding (**Fig. 8***B*). A cell without KaiA leaves the major
transcription factor RpaA hypophosphorylated for two reasons. First, without KaiC undergoing
its daytime phosphorylation cycle, SasA engagement with the oscillator is diminished, reducing
its kinase activity acting on RpaA. Secondly, without KaiA present as a competitor, the
opportunities for CikA binding on KaiB-KaiC complexes increase, leading to hyperactive
phosphatase activity and ultimately eliminating any chance for phosphorylated RpaA to
accumulate. The result is a severe decrease in fitness in LDC. However, cells without KaiA and
CikA (and even without KaiABC) have ample P-RpaA, indicating that other pathways contribute
to the phosphorylation of RpaA. This fact emphasizes the counterintuitive point that activation of
CikA phosphatase in the night-time complex, rather than activation of SasA kinase, is the key
signaling state of the cyanobacterial clock (12).

The screen revealed expected mutants as well. Although a *kaiC* mutant does not have a notable
LDC defect in pure culture, it is known to do poorly in LDC when grown in a population with
other cells that have a functioning circadian rhythm, as is the case in the pooled growth scenario
used here, illustrating soft selection for circadian rhythms in cyanobacteria (1). As expected,
kaiC scored as deficient in LDC, but not as severely as many other mutants. One of the most
severely LDC-defective mutants we know of is *rpaA*, which dies within a few hours of dark
exposure and requires special handling in the lab. Strains with disruptions in *rpaA* did score as
negative, but only to a moderate degree (-0.7). This discrepancy between severe known
phenotype and moderate score can be attributed to the fact that *rpaA* mutants were also less fit

than the population in CLC, decreasing its magnitude of relative fitness in LDC, similar to the
situation for mutants defective in glycogen anabolic processes.

Importantly, the screen also identified an equivalent list of genes whose loss improves growth in
LDC. Although not the target of the current study, these genes have great potential for providing
new insights into *S. elongatus* diurnal physiology and perhaps improved genetic backgrounds
for metabolic engineering. In contrast to many of the mutants that negatively affect LDC growth,
the list of top positive mutants includes many genes with less transparent roles in LDC
conditions, including some that carry no functional annotation. The phenotype may be
attributable to a specific pathway or may result from the additive effects of many unrelated
genes whose expression is altered in a regulon indirectly. Some mutants may outcompete
others in the population because they have alleviated photochemical intermediate buildup or
lessened the uptake of harmful metabolites in the medium. The scant information available
regarding these pathways makes it difficult to predict why their value would be different in CLC
vs. LDC.

In summary, this study has improved our understanding of the mechanism of core oscillator
protein interactions and generated a comprehensive map of genetic pathways needed for
survival in LDC in *S. elongatus*. We have shown that KaiA, originally thought to only regulate
KaiC autophosphorylation, also modulates CikA-mediated clock output. More broadly, the use of
RB-TnSeq on a dense transposon library in this study has identified the genes in the genome
necessary for growth in alternating light-dark conditions, established a firm connection between
circadian clock regulation and the metabolic pathways critical for fitness in more natural
environments, and contributed to a more complete picture of the finely balanced mechanisms
that underlie circadian clock output.

Materials and Methods

Bacterial strains and culture conditions. All cultures were constructed using WT *S. elongatus*
PCC 7942 stored in our laboratory as AMC06. Cultures (**Table S1**) were grown at 30 °C using
either BG-11 liquid or solid medium with antibiotics as needed for selection at standard
concentrations: 20 µg/mL for kanamycin (Km) and gentamicin (Gm), 22 µg/mL for
chloramphenicol (Cm), and 10 µg/mL each for spectinomycin and streptomycin (SpSm) (51).
Liquid cultures were either cultivated in 100 mL volumes in 250 mL flasks shaken at ~150 rpm
on an orbital shaker or in 400 mL volume in top-lit bioreactors (Phenometrics Inc. ePBR
photobioreactors version 1.1) mixed via filtered ambient-air bubble agitation with a flow rate of
0.1 mL per min at 30 °C. Viable cell plating for LDC-sensitivity testing and quantification of
cellular ROS were performed as previously described (15). Spot-plate growth was quantified by
converting the plate image to 8-bit and performing densitometric analyses using National
Institutes of Health ImageJ software (52). Transformation was performed following standard
protocols (53). Strains were checked periodically for contamination using BG-11 Omni plates
(BG-11 supplemented with 0.04% glucose and 5% LB) (53). Verification of the disruption of the
kaiA gene in mutants was performed by PCR and is shown in **Fig. S1**. For immunoblotting
experiments and ROS measurements cultures were grown in flasks. Samples for protein were
taken after 3 light-dark cycles at specified time points immediately before the lights turned on
and turned off.

One-on-one competition experiments. The relative fitness of *cikA* null and *cikA+* strains was
assessed in growth competition experiments that leveraged different antibiotic resistance
markers in the strains. The *cikA* mutant carries a Gm-resistance marker inserted in the *cikA*
locus. AMC06 was transformed with pAM1579 to generate a Km-resistant strain that carries the

drug cassette in neutral site II (designated as "WT-Km^R" for this experiment). The axenic
cultures were grown to OD$_{750}$ ~ 0.4, washed three times with BG-11 via centrifugation at 4,696
G for 10 min to remove antibiotics, and diluted with BG-11 to an OD$_{750}$ of 0.015. Cultures were
then mixed in a 1:1 ratio and grown without antibiotics for 8 days under ~100 µmol photos· m^2
· s^{-1} light. To assay survival of each genotype at 0, 3, and 8 days, mixed cultures were diluted
using a 1:10 dilution series and plated in 20 µL spots in duplicate onto ~2 mL of BG-11 agar
with either Gm or Km poured in the wells of a 24-well microplate. CFU's of each strain were
calculated from the dilution series across the two antibiotic regimes and were used to determine
the percentage of each strain in the mixture population over time. To ensure that the expression
of the antibiotic markers did not influence fitness, WT cultures that carried each drug-resistance
marker (Km in neutral site II via plasmid pAM1579 or Gm in neutral site III via plasmid
pAM5328) were grown together and assayed, and displayed no significant differences (**Fig. S2).**

Rb-Tnseq Assay. Samples of the library archived at -80 C were quickly thawed in a 37 C water
bath for 2 min and then divided into three flasks of 100 mL BG-11 with Km, and incubated at 30
°C in 30 µmol photos· m^2· s^{-1} light for 1 day without shaking. The library culture was then
moved to 70 µmol photos· m^2· s^{-1} light on an orbital shaker till OD$_{750}$ reached ~0.3. The
cultures were then combined, diluted to starting density of OD$_{750}$ 0.025, and 4 replicates of 15
mL were spun down at 4,696 g and frozen at -80 °C as Time 0 samples to determine population
baseline. After transfering to bioreactors, CLC bioreactors were set for continuous light at 500
µmol photos· m^2· s^{-1} and LDC bioreactors were set to a square-wave cycle consisting of 12 h
light - 12 h dark. Bioreactors were set to run in turbidostat mode maintaining the density at
OD$_{750}$ = 0.1. Bioreactors were sampled after approximately 6 generations: the CLC reactors
after 3 days (three 24-h light periods) and LDC reactors after 4 days (four 12-h light-12-h dark
light dark periods). Two of the three biological replicates for each condition were performed
concurrently (using 4 bioreactors) while a third was performed later using the same procedure
and conditions.

Bioluminescence monitoring. Flask-grown batch cultures that reached a density of OD$_{750}$ =
0.3-0.4 were diluted as needed to OD$_{750}$ of 0.3 and 20 µL of each culture was placed on a pad
of BG-11 agar containing 10 µL of 100 mM firefly luciferin, in a well of a 96-well dish. Plates
were covered with clear tape to prevent drying and holes were poked using a sterile needle to
allow air transfer. Cultures were synchronized by incubating the plate under permissive LDC
conditions (0 and 30 µmol photos· m^2· s^{-1} light) for 3 days, and then returned to constant light
(LL) conditions for bioluminescence sampling by a Packard TopCount luminometer
(PerkinElmer Life Sciences). Bioluminescence of P$_{kaiBC}$::*luc* firefly luciferase fusion reporter was
monitored at 30 °C under CLC as described previously (54). Data were analyzed with the
Biological Rhythms Analysis Software System (http://millar.bio.ed.ac.uk/pebrown/
brass/brasspage.htm) import and analysis program using Microsoft Excel. Results shown are
the average of four biological replicate wells located in the innermost section of the plate, where
drying is minimal.

Protein sample preparation and gel electrophoresis, and Phos-tag acrylamide gels.
Protein was isolated from cells and diluted to final loading concentration of 5 µg/well for RpaA
analysis and 10 µg/well for KaiC analysis (55). SDS/PAGE was performed according to
standard methods with the following exceptions. Phosphorylation of RpaA was detected using
10% SDS-polyacrylamide gels supplemented with Phos-tag ligand (Wako Chemicals USA) at a
final concentration of 25 µM and manganese chloride at a final concentration of 50 µM. Gels

were incubated once for 10 min in transfer buffer supplemented with 100 mM EDTA, followed by
a 10-min incubation in transfer buffer without EDTA before standard wet transfer. Protein
extracts and the electrophoretic apparatus were chilled to minimize hydrolysis of heat-labile
phospho-Asp. Protein extracts for use in Phos-tag gels were prepared in Tris-buffered saline,
and extracts for standard SDS/PAGE were prepared in PBS. RpaA antiserum (a gift from E.
O'Shea, Harvard University, Cambridge, MA) was used at a dilution of 1:2,000 and secondary
antibody (goat anti-rabbit IgG, Calbiochem Cat # 401315) at 1:5000. KaiC immunoblotting was
performed similarly using KaiC antiserum diluted to 1:2000 and secondary antibody diluted to
1:10,000 (Horseradish Peroxidase-labeled goat anti-chicken IgY, Aves Labs, Cat # H-1004)
(56). Densitometric analyses were performed using National Institutes of Health ImageJ
software (52) (**Fig. S3, Dataset S3**). Samples were immediately stored at -80 °C, thawed once,
and always kept on ice.

Fitness calculations

Of the 2,723 genes comprising the genome of *S. elongatus* 718 are essential, and mutants in
the essential genes are not present in the library (24). To estimate the fitness effects of gene
disruptions in LDC relative to CLC, we developed an analysis pipeline of curating the data from
the 2,005 nonessential genes, normalizing it, and then analyzing it using linear models (**Dataset
S1**). We first counted the number of reads for each sample to use as a normalizing factor
between samples. Bar codes are dispersed across the genome, and we removed any bar code
falling outside of a gene (24,868 bar codes out of 154,949 total bar codes) or within a gene but
not within the middle 80% (27,763/154,949). Based on the bar codes remaining, we removed
any gene not represented by at least three bar codes in different positions (114 genes out of
2,075 total). This filtering left us with 102,136 bar codes distributed across 1,961 genes.

For each bar code in each sample we added a pseudocount of one to the number of reads,
divided by the total number of reads for the sample as calculated before, and took the log-2
transformation of this sample-normalized number of reads. The experiment involved two
different starting pools of strains (called T0), each of which was divided into CLC and LDC
samples. To account for different starting percentages of each bar code within the T0 pools, we
averaged the log-2 transformed values for a bar code across the four replicate T0 samples for
each pool then subtracted these average starting bar-code values from the CLC and LDC
values in the respective pools. We also removed any gene without at least 15 T0 reads (across
the four replicates and before adding the pseudocount) in each pool (89/1,961), leaving 1,872
genes and 101,258 bar codes.

For each gene, we used maximum likelihood to fit a pair of nested linear mixed effects models
to the sample- and read-normalized log-2 transformed counts:

(1) $y_{i,j,k} = \mu_g + C_j + B_i + \varepsilon_{i,j,k}$; $B_i \sim iid\ N(0,\varsigma_g^2)$; $\varepsilon_{i,j,k} \sim iid\ N(0,\sigma_g^2)$

(2) $y_{i,j,k} = \mu_g + B_i + \varepsilon_{i,j,k}$; $B_i \sim iid\ N(0,\varsigma_g^2)$; $\varepsilon_{i,j,k} \sim iid\ N(0,\sigma_g^2)$

where $y_{i,j,k}$ is the normalized log-2 value for bar code *i* in gene *g* in condition *j* for sample *k*, μ_g is
the average value for the gene, C_j is the fixed effect of condition *j*, B_i is a random effect for bar
code *i*, and $\varepsilon_{i,j,k}$ is the residual. We identified genes with significant fitness differences between
conditions by comparing the difference in the -2*log likelihoods of the models to a chi-square
distribution with one degree of freedom, estimating a p-value, accounting for multiple testing by
the false-discovery rate method of Benjamini and Hochberg (57), and selecting those gene with

adjusted p-values less than 0.01. We took the contrast $C_{LDC} - C_{CLC}$ to be the estimated LDC-
specific fitness effect of knocking out the gene (**Dataset S2**).

Results obtained across biological replicates and across similar screens performed in either
batch cultures grown in 250 mL Erlenmeyer shake flasks and on solid agar on Petri plates were
consistent (**Fig. S4, Dataset S4**). Mutants that had estimated fitness scores below a confidence
threshold (associated FDR adjusted p-value <0.01) (**Dataset S2**) were not considered. Mutants
with gene disruptions that had an LDC fitness score ≥ 1.0 or ≤ -1.0 with a false discovery rate ≤
1% were classified having strong fitness phenotype specific to growth in LDC, and those with a
fitness score of < 1.0 but ≥ 0.5 and > -1.0 but ≤ -0.5 were classified as causing a moderate
fitness phenotype. Fitness scores <0.5 and >-0.5 were categorized as negligent/minor, i.e., little
change in abundance over the course of the experiment in LDC vs. CLC.

Fluorescence Spectroscopy. To monitor the formation of the CikA activating complex,
fluorescence anisotropy measurements were performed at 30 °C on an ISS PC1
spectrofluorometer equipped with a three-cuvette sample compartment, at excitation and
emission wavelengths of 492 nm and 530 nm, respectively, for samples including 6-
iodoacetamidofluorescein (6-IAF) labeled KaiB-FLAG-K251C alone (0.05 µM), and its mixtures
with KaiC (4 µM) or FLAG-CikA (4 µM) or KaiC (4 µM) + FLAG-CikA (4 µM). Each sample (400
µL) was either in an ATP buffer [20 mM Tris, 150 mM NaCl, 5 mM MgCl$_2$, 1 mM ATP, 0.5 mM
EDTA, 0.25 mM TCEP (tris(2-carboxyethyl)phosphine), pH 8.0] or in ATP/ADP buffer [20 mM
Tris, 150 mM NaCl, 5 mM MgCl$_2$, 0.5 mM ATP, 0.5 mM ADP, 0.5 mM EDTA, 0.25 mM TCEP,
pH 8.0]. TCEP was introduced from the FLAG-CikA stock [20 mM Tris, 150 mM NaCl, 5 mM
TCEP, pH 8.0]. KaiB stock was in the buffer [20 mM Tris, 150 mM NaCl, pH 8.0] and KaiC stock
in the buffer [20 mM Tris, 150 mM NaCl, 5 mM MgCl$_2$, 1 mM ATP, 0.5 mM EDTA, pH 8.0].

The KaiC samples included freshly dephosphorylated WT KaiC, and KaiC phosphomimetics
KaiC-AE and KaiC-EA. To prepare the freshly dephosphorylated KaiC, WT KaiC at 20 µM in the
buffer [20 mM Tris, 150 mM NaCl, 5 mM MgCl$_2$, 1 mM ATP, 0.5 mM EDTA, pH 8.0] was
incubated at 30 °C for 24 hours before the fluorescence binding assays. SDS-PAGE analysis
was performed to determine KaiC and CikA stability and KaiC phosphorylation levels (**Fig. S5**,
Dataset S5).

Acknowledgments

552 This work was supported by research grants (R35GM118290 to SSG; R01GM107521 to AL) and
553 training grants (T32GM007240 to BER and SD) from the National Institutes of Health and the
554 National Science Foundation (MCB1517482 to SAR).

References

1. Ouyang Y, Andersson CR, Kondo T, Golden SS, Johnson CH (1998) Resonating circadian clocks enhance fitness in cyanobacteria. *Proc Natl Acad Sci U S A* 95(15):8660–8664.

2. Woelfle MA, Ouyang Y, Phanvijhitsiri K, Johnson CH (2004) The adaptive value of circadian clocks: an experimental assessment in cyanobacteria. *Curr Biol* 14(16):1481–1486.

3. Lambert G, Chew J, Rust MJ (2016) Costs of Clock-Environment Misalignment in Individual Cyanobacterial Cells. *Biophys J* 111(4):883–891.

4. Yerushalmi S, Yakir E, Green RM (2011) Circadian clocks and adaptation in *Arabidopsis*. *Mol Ecol* 20(6):1155–1165.

5. Dodd AN, et al. (2005) Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. *Science* 309(5734):630–633.

6. Morris CJ, Yang JN, Scheer FAJL (2012) The impact of the circadian timing system on cardiovascular and metabolic function. *Prog Brain Res* 199:337–358.

7. Kelleher FC, Rao A, Maguire A (2014) Circadian molecular clocks and cancer. *Cancer Lett* 342(1):9–18.

8. Kim JH, Duffy JF (2018) Circadian Rhythm Sleep-Wake Disorders in Older Adults. *Sleep Med Clin* 13(1):39–50.

9. Chaix A, Zarrinpar A, Panda S (2016) The circadian coordination of cell biology. *J Cell Biol* 215(1):15–25.

10. Nakajima M, et al. (2005) Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation in vitro. *Science* 308(5720):414–415.

11. Tseng R, et al. (2017) Structural basis of the day-night transition in a bacterial circadian clock. *Science* 355(6330):1174–1180.

12. Paddock ML, Boyd JS, Adin DM, Golden SS (2013) Active output state of the *Synechococcus* Kai circadian oscillator. *Proc Natl Acad Sci U S A* 110(40):E3849–57.

13. Gutu A, O'Shea EK (2013) Two antagonistic clock-regulated histidine kinases time the activation of circadian gene expression. *Mol Cell* 50(2):288–294.

14. Markson JS, Piechura JR, Puszynska AM, O'Shea EK (2013) Circadian control of global gene expression by the cyanobacterial master regulator RpaA. *Cell* 155(6):1396–1408.

15. Diamond S, et al. (2017) Redox crisis underlies conditional light-dark lethality in cyanobacterial mutants that lack the circadian regulator, RpaA. *Proc Natl Acad Sci U S A* 114(4):E580–E589.

16. Puszynska AM, O'Shea EK (2017) Switching of metabolic programs in response to light availability is an essential function of the cyanobacterial circadian output pathway. *Elife* 6. doi:10.7554/eLife.23210.

17. Swan JA, Golden S, LiWang A, Partch CL (2018) Structure, function, and mechanism of the
core circadian clock in cyanobacteria. *J Biol Chem*. doi:10.1074/jbc.TM117.001433.

18. Iwasaki H, et al. (2000) A *kaiC*-interacting sensory histidine kinase, SasA, necessary to
sustain robust circadian oscillation in cyanobacteria. *Cell* 101(2):223–233.

19. Boyd JS, Bordowitz JR, Bree AC, Golden SS (2013) An allele of the crm gene blocks
cyanobacterial circadian rhythms. *Proc Natl Acad Sci U S A* 110(34):13950–13955.

20. Diamond S, Jun D, Rubin BE, Golden SS (2015) The circadian oscillator in *Synechococcus*
elongatus controls metabolite partitioning during diurnal growth. *Proc Natl Acad Sci U S A*
112(15):E1916–25.

21. Gründel M, Scheunemann R, Lockau W, Zilliges Y (2012) Impaired glycogen synthesis
causes metabolic overflow reactions and affects stress responses in the cyanobacterium
Synechocystis sp. PCC 6803. *Microbiology* 158(Pt 12):3032–3043.

22. Puszynska AM, O'Shea EK (2017) ppGpp Controls Global Gene Expression in Light and in
Darkness in *S. elongatus*. *Cell Rep* 21(11):3155–3165.

23. Hood RD, Higgins SA, Flamholz A, Nichols RJ, Savage DF (2016) The stringent response
regulates adaptation to darkness in the cyanobacterium *Synechococcus elongatus*. *Proc*
Natl Acad Sci U S A 113(33):E4867–76.

24. Rubin BE, et al. (2015) The essential gene set of a photosynthetic organism. *Proc Natl*
Acad Sci U S A 112(48):E6634–43.

25. Broddrick JT, et al. (2016) Unique attributes of cyanobacterial metabolism revealed by
improved genome-scale metabolic modeling and essential gene analysis. *Proc Natl Acad*
Sci U S A 113(51):E8344–E8353.

26. Wetmore KM, et al. (2015) Rapid quantification of mutant fitness in diverse bacteria by
sequencing randomly bar-coded transposons. *MBio* 6(3):e00306–15.

27. Vijayan V, Zuzow R, O'Shea EK (2009) Oscillations in supercoiling drive circadian gene
expression in cyanobacteria. *Proc Natl Acad Sci U S A* 106(52):22564–22568.

28. Miao X, Wu Q, Wu G, Zhao N (2003) Changes in photosynthesis and pigmentation in an
agp deletion mutant of the cyanobacterium *Synechocystis* sp. *Biotechnol Lett* 25(5):391–
396.

29. Li X, Shen CR, Liao JC (2014) Isobutanol production as an alternative metabolic sink to
rescue the growth deficiency of the glycogen mutant of *Synechococcus elongatus* PCC
7942. *Photosynth Res* 120(3):301–310.

30. Porankiewicz J, Clarke AK (1997) Induction of the heat shock protein ClpB affects cold
acclimation in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *J Bacteriol*
179(16):5111–5117.

31. Gill RT, et al. (2002) Genome-wide dynamic transcriptional profiling of the light-to-dark
transition in *Synechocystis* sp. strain PCC 6803. *J Bacteriol* 184(13):3671–3681.

32. Hagemann M, Vinnemeier J, Oberpichler I, Boldt R, Bauwe H (2005) The glycine

decarboxylase complex is not essential for the cyanobacterium *Synechocystis* sp. strain
PCC 6803. *Plant Biol* 7(1):15–22.

33. Hackenberg C, et al. (2009) Photorespiratory 2-phosphoglycolate metabolism and
photoreduction of O2 cooperate in high-light acclimation of *Synechocystis* sp. strain PCC
6803. *Planta* 230(4):625–637.

34. Cohen SE, Golden SS (2015) Circadian Rhythms in Cyanobacteria. *Microbiol Mol Biol Rev*
79(4):373–385.

35. Ishiura M, et al. (1998) Expression of a gene cluster *kaiABC* as a circadian feedback
process in cyanobacteria. *Science* 281(5382):1519–1523.

36. Nishiwaki T, et al. (2004) Role of KaiC phosphorylation in the circadian clock system of
Synechococcus elongatus PCC 7942. *Proc Natl Acad Sci U S A* 101(38):13927–13932.

37. Xu Y, et al. (2004) Identification of key phosphorylation sites in the circadian clock protein
KaiC by crystallographic and mutagenetic analyses. *Proc Natl Acad Sci U S A*
101(38):13933–13938.

38. Williams SB, Vakonakis I, Golden SS, LiWang AC (2002) Structure and function from the
circadian clock protein KaiA of *Synechococcus* elongatus: a potential clock input
mechanism. *Proc Natl Acad Sci U S A* 99(24):15357–15362.

39. Ditty JL, Canales SR, Anderson BE, Williams SB, Golden SS (2005) Stability of the
Synechococcus elongatus PCC 7942 circadian clock under directed anti-phase expression
of the kai genes. *Microbiology* 151(Pt 8):2605–2613.

40. Nair U, Ditty JL, Min H, Golden SS (2002) Roles for sigma factors in global circadian
regulation of the cyanobacterial genome. *J Bacteriol* 184(13):3530–3538.

41. Schmitz O, Katayama M, Williams SB, Kondo T, Golden SS (2000) CikA, a
bacteriophytochrome that resets the cyanobacterial circadian clock. *Science*
289(5480):765–768.

42. Chen Y, et al. (2009) A novel allele of *kaiA* shortens the circadian period and strengthens
interaction of oscillator components in the cyanobacterium *Synechococcus elongatus* PCC
7942. *J Bacteriol* 191(13):4392–4400.

43. Kutsuna S, Nakahira Y, Katayama M, Ishiura M, Kondo T (2005) Transcriptional regulation
of the circadian clock operon kaiBC by upstream regions in cyanobacteria. *Mol Microbiol*
57(5):1474–1484.

44. Chang Y-G, et al. (2015) Circadian rhythms. A protein fold switch joins the circadian
oscillator to clock output in cyanobacteria. *Science* 349(6245):324–328.

45. Kitayama Y, Iwasaki H, Nishiwaki T, Kondo T (2003) KaiB functions as an attenuator of
KaiC phosphorylation in the cyanobacterial circadian clock system. *EMBO J* 22(9):2127–
2134.

46. Rust MJ, Golden SS, O'Shea EK (2011) Light-driven changes in energy metabolism directly
entrain the cyanobacterial circadian oscillator. *Science* 331(6014):220–223.

47. Ito H, et al. (2009) Cyanobacterial daily life with Kai-based circadian and diurnal genome-
wide transcriptional control in *Synechococcus elongatus*. *Proc Natl Acad Sci U S A*
106(33):14168–14173.

48. Matson MM, Atsumi S (2017) Photomixotrophic chemical production in cyanobacteria. *Curr*
Opin Biotechnol 50:65–71.

49. McEwen JT, Kanno M, Atsumi S (2016) 2,3 Butanediol production in an obligate
photoautotrophic cyanobacterium in dark conditions via diverse sugar consumption. *Metab*
Eng 36:28–36.

50. Snijder J, et al. (2017) Structures of the cyanobacterial circadian oscillator frozen in a fully
assembled state. *Science* 355(6330):1181–1184.

51. Taton A, et al. (2014) Broad-host-range vector system for synthetic biology and
biotechnology in cyanobacteria. *Nucleic Acids Res* 42(17):e136.

52. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* 9(7):671–675.

53. Clerico EM, Ditty JL, Golden SS (2007) Specialized techniques for site-directed
mutagenesis in cyanobacteria. *Methods Mol Biol* 362:155–171.

54. Mackey SR, Golden SS, Ditty JL (2011) The itty-bitty time machine genetics of the
cyanobacterial circadian clock. *Adv Genet* 74:13–53.

55. Ivleva NB, Golden SS (2007) Protein extraction, fractionation, and purification from
cyanobacteria. *Methods Mol Biol* 362:365–373.

56. Dong G, et al. (2010) Elevated ATPase activity of KaiC applies a circadian checkpoint on
cell division in *Synechococcus elongatus*. *Cell* 140(4):529–539.

57. Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J R Stat Soc Series B Stat Methodol* 57(1):289–300.

58. Ditty JL, Williams SB, Golden SS (2003) A cyanobacterial circadian timing mechanism.
Annu Rev Genet 37:513–543.

Figure Legends

Fig 1. Using RB-TnSeq to assess genetic contributions to fitness in LDC. A dense pooled mutant library containing unique known bar-code identification sequences linked to each individual gene insertion was used to estimate the fitness of each loss-of-function mutant grown under alternating light-dark conditions. The library was grown under continuous light and sampled to determine the baseline abundance of each strain in the population prior to transfer to photobioreactors which were then either exposed to continuous light or alternating 12-h light - 12-h dark regimes. Bar-code quantification of the library population in both conditions was normalized to the baseline and compared against each other to estimate the fitness consequences of loss of function of each gene specific to growth in LDC.

Fig 2. LDC Rb-TnSeq screen results breakdown. *(A)* Of the 2723 genes in the genome, 1872 genes were analyzed in the library population. Of those, data from mutants with loss-of-function of 1420 genes fell below our false discovery threshold. Of the remaining 452 genes, 220 had moderate to strong fitness effects. *(B)* A volcano plot highlighting genes of interest. Red-shaded region indicates an estimated fitness decrease in LDC and the green-shaded region indicates a fitness increase. *(C)* Functional category composition of genes that gave strong fitness scores. Categories based on COG classification or GO ID. Green corresponds to strong fitness increases; Red corresponds to strong fitness decreases.

Fig. 3. Connection between essential night-time metabolism and RpaA regulation. *(A)* A metabolic map showing the reactions (red) that are controlled by genes that are essential in LDC. Reactions that are non-essential (solid arrows) or essential (dashed arrows) in CLC are are also indicated. Stars mark NADPH-generating reactions. * gene was not revealed in the screen but is known to be required in LDC; # minor LDC decrease measured, likely due to fitness contribution in CLC which affects calculation **(B)** LDC-sensitive mutant population enriched for genes known to be positively regulated by RpaA.

Fig. 4. *kaiA* mutant growth and *cikA*-based intervention in LDC. *(A)* Dilution spot-plate growth of strains in CLC and LDC. A representation of the genotype of each strain is portrayed next to each dilution series. Shaded grey bar at end of *kaiA* = negative regulatory element. *(B)* *cikA* mutant fitness increase in competition with WT in LDC and CLC. *(C)* Growth of liquid cultures in replicate photobioreactors in LDC. Vertical shaded areas represent dark conditions. Shaded areas around each line = standard deviation of biological replicates.

Fig. 5. Bioluminescence levels from *kaiBC* reporter in mutant backgrounds. The intensity of P$_{kaiBC}$::*luc* output signal correlates with LDC sensitivity. Rhythms were measured in CLC conditions driven by the *kaiBC* promoter after exposure to 72 h of entraining LDC that includes a low-light intensity that is permissive for the mutants. Shaded areas around each line = standard deviation of biological replicates.

Fig. 6. RpaA status correlates with growth success in LDC. *(A)* RpaA phosphorylation at dusk (after 12 h light) showing average ± SD values. Letters indicate groupings from ANOVA analysis. *(B)* Levels of KaiC from cells taken at dusk. **C.** RpaA phosphorylation across the light-dark transitions beginning with dawn. *(D)* H$_2$DCFDA fluorescence over a 12-h dark period during an LDC, indicating total cellular ROS in WT, *kaiA^{del}*, and *kaiA^{del}cikA*. The shaded area indicates the period of darkness following a 12-h light period. P-RpaA denotes phosphorylated protein.

Fig. 7. CikA-activating complex formation. *(A)* Fluorescence anisotropy of the binding kinetics of 6-IAF–labeled KaiB (black) to WT KaiC, KaiA-AE, or KaiC-EA (red in each case) with and without CikA (blue), and in the presence of 0.5 mM ATP + 0.5 mM ADP. *(B)* Binding of KaiB (black) to WT KaiC or KaiC-AE (red in each case) with and without CikA (blue) in the presence of 1.0 mM ATP.

Fig. 8. Model for LDC sensitivity due to circadian clock disfunction. *(A)* Conventionally, the role of KaiA is primarily to bind to the A-loops of the CII domain of KaiC, stimulating its autophosphorylation and consequently increasing the activity of the kinase SasA; this activity in turn results in accumulation of phosphorylated RpaA throughout the day. P-RpaA initiates the expression of circadian-controlled night-time class 1 genes at dusk. *(B)* Without KaiA, KaiC phosphorylation is suppressed and, thus, SasA kinase activity is diminished, reducing the pools of SasA-mediated P-RpaA. Upon KaiC-KaiB complex formation, CikA has access to binding sites usually occupied by KaiA, and becomes hyperactive; excessive CikA phosphatase activity extinguishes intracellular pools of P-RpaA and eliminates the cell's ability to express essential genes needed for survival in LDC. Figure modified from (11). The dotted line in thepanel B graph represents the WT levels of P-RpaA from A.

Supplemental Material

Dataset Files

Dataset S1: This zipped file contains the annotated R script for determining genetic fitness estimates under LDC and the data files processed by the script: 1) "lightdarkbioreactors_all.poolcount.csv.", the location and reads of bar-coded transposon mutants for each sample 2) "lightdarkbioreactors_exp.csv", designation of the samples as T0, control genotype, and experimental genotype 3) "genes.tab", the gene coordinate file used to map bar-coded transposon mutants to genes.

Dataset S2: This Excel file has the following tabs: 1) Fitness estimates determined by processing the files in Dataset S1. 2) Search utility for displaying information easily for a desired gene locus id.

Dataset S3: This Excel table contains two sheets: 1) The data calculated from the Phos-tag™ immunoblots for RpaA phosphorylation and the corresponding blots they were derived from in Fig. S3; and 2) Data in a format for entering into R and using the script provided in the text box to perform the statistical analysis that generated the groupings in Fig. 6A.

Dataset S4: RB-TnSeq data comparing photobioreactor biological replicates, and the estimates from analysis of the library grown in triplicate in photobioreactors with the library grown on Petri plates and in 100 mL shake flasks under similar LDC conditions.

Dataset S5: This Excel file contains the data from the anisotropy measurements of *in vitro* KaiB, KaiC, and CikA.

Supplemental Table

Table S1

Strain	Genetic background	Reporter	Antibiotic resistance	Reference
AMC541	WT *S.elongatus*	P_{kaiBC} : : *luc*;	Cm	(42)
AMC06	WT *S.elongatus*	none	none	Lab collection
AMC1161	*kaiA* insertion; pAM2969 in AMC541	P_{kaiBC} : : *luc*;	Cm Km	(58)
AMC702	*kaiA* deletion; in-frame markerless constructed in AMC541	P_{kaiBC} : : *luc*;	Cm	(58)
AMC1936	*kaiABC* knockout; pAM4252 in AMC541	P_{kaiBC} : : *luc*;	Cm Km	(12)
AMC1679	*kaiA cikA* double knockout; pAM2152 in AMC702	P_{kaiBC} : : *luc*;	Cm Gm	(56)
AMC2479	Complemented *kaiA cikA* double knockout; pAM3389 expressing *cikA* in AMC1679	P_{kaiBC} : : *luc*;	Cm SpSm	This study
AMC1100	Complemented *kaiA* deletion; pAM2246 in AMC702 expressing WT *kaiA* in NS1.	P_{kaiBC} : : *luc*;	Cm SpSm	(39)
AMC2480	Complemented *kaiA cikA* double knockout AMC1679 with pAM5294 expressing *cikA*^{C644R} variant.	P_{kaiBC} : : *luc*;	Cm SpSm	This study
AMC2481	*kaiA* deletion; pAM5389 in AMC541	P_{kaiBC} : : *luc*;	Cm SpSm	This study
AMC2482	Remade *kaiA cikA* double knockout; pAM2152 in AMC2481	P_{kaiBC} : : *luc*;	Cm SpSm Gm	This study
AMC2483	Remade *kaiA cikA* double knockout; pAM2152 in AMC702	P_{kaiBC} : : *luc*;	Cm Gm	This study

Fig. S1. Mutant strain complementation reverses phenotype. *(A)* Dilution spot plate showing growth of various *kaiA* mutant strains under CLC or LDC conditions. Reintroduction of *kaiA* greatly improves the growth of the *kaiA* deletion mutant (compare rows 6 and 3 in LDC); reintroduction of *cikA* into the *kaiA*^{del}*cikA* double mutant reverses the alleviating effects of *cikA* disruption (compare rows 7 and 4) *(B)* Genetic map showing a representation of the strain backgrounds. **C.** DNA gel electrophoresis of PCR amplification confirming mutations in *kaiA* gene in *kaiA* mutant strains.

Fig. S2. Effect of antibiotic marker expression during head-to-head growth competition. WT cultures with antibiotic resistance cassettes for either Gm or Km in a neutral site of the genome were mixed in a 1:1 ratio and competed against each other in a head-to-head growth assay without antibiotics under CLC and LDC conditions.

Fig. S3. Immunoblots for determining RpaA phosphorylation and KaiC levels. For blots A-F Phos-tag^{TM} was added to cell extracts to distinguish phosphorylated from non-phosphorylated proteins and antisera were used to identify the labeled protein in each panel. Panels A-B and G show proteins taken from cells sampled at the end of the light period after 3 consecutive 12-h light:12-h dark cycles. Panels E-F show samples taken every 12 h over the course of a 12-h light: 12-h dark cycle; even lanes are samples at the end of the dark period and odd lanes are samples taken at the end of the light period. Genotypes of strains are listed in Table S1.

(A) lanes: 1, AMC541; 2, AMC702; 3, AMC2481; 4, AMC2481; 5, AMC2482; 6, AMC2482; 7, AMC1679; 8, AMC2480; 9, AMC2480; 10, AMC2479; 11, AMC2479; 12, AMC2483; 13, AMC1161; 14, AMC1936;

(B) lanes: 1, AMC541; 2, AMC1161; 3, AMC702; 4, AMC1936; 5, AMC1679; 6, AMC2479.

(C) lanes: 1, AMC541; 2, AMC1161; 3, AMC702; 4, AMC1679; 5, AMC1100; 6, AMC2479; 7, empty; 8, AMC541; 9, AMC2481; 10, AMC702; 11, empty; 12, AMC1936; 13, empty.

(D) lanes: 1-6, AMC541; 7-12, AMC702.

(E) lanes: 1-6, AMC541; 7-12, AMC702.

(F) lanes 1-6, AMC541; 7-12, AMC1679.

(G) lanes: 1, AMC541; 2, AMC1161; 3, AMC2481; 4, AMC2482; 5, empty; 6, AMC2482; 7, empty; 8, AMC541; 9, AMC1161; 10, AMC702; 11, AMC1679; 12, AMC1936.

Fig. S4. Comparison of LDC Rb-TnSeq results across replicates and growth conditions. *(A)* Strain fitness for each photobioreactor replicate was estimated as previously described in (24)**.** Fitness scores with t-statistics ≥ |4| were compared against each other showing agreement among biological replicates. *(B)* Comparison of LDC fitness results screening the library in different growth formats including photobioreactors, Erlenmeyer flasks, and Petri plates. Because there was only 1 measurement each for the plates and flask screens, the stringency for sensitivity scores was relaxed. *(C)* Summary of the 13 mutant strains that showed strong LDC defects across all growth media situations.

Fig. S5. SDS-PAGE analysis of KaiC and CikA stability and KaiC dephosphorylation. *(A)* 17% SDS-PAGE analysis of KaiC-EA or KaiC-AE and CikA before and after KaiB binding assays. *(B)* 17% SDS-PAGE analysis of KaiC and CikA stability before and after KaiB binding assays. *(C)* 6.5% SDS-PAGE analysis showing that KaiC retains a hypophosphorylated state during KaiB-binding assays. Note that all samples other than the controls contain 6-IAF-labeled KaiB-FLAG-K25C. Due to the low concentration of KaiB (0.05 µM), SDS-PAGE runs were optimized only for visualization of KaiC and CikA. The stable anisotropy readings of free KaiB, shown in Fig. 7, suggest its stability for the duration of the binding assays.

828 **Figure 1**

830   **Figure 2**

Figure 3

Figure 4

A

B

837 **Figure 5**

840
841

842   **Figure 7**

844 **Figure 8**

A

KaiA stimulates KaiC
phosphorylation; SasA
phosphorylates **RpaA**

KaiB binds to KaiC
and displaces SasA

KaiA sequestered on KaiB;
Activated CikA
dephosphorylates **RpaA**

B

KaiC autophosphorylation
ceases without KaiA;
SasA kinase activity stifled

KaiB binds to KaiC

CikA binds to KaiB with no
KaiA competition, becomes
hyperactivated to
dephosphorylate **RpaA**

Figure S1

A

B

C

848   **Figure S2**

Figure S3

 Figure S4

A

B

Fitness Decrease in LDC (PBR)
Fitness score <-1.0, FDR <.01

Fitness Decrease in LDC
on Petri Plates Score <-0.5

1
2%

27
66%

13
32%

Fitness Decrease in LDC in
Erlenmeyer Flasks Score <-0.5

C

locusId	Description	Flask	Plate	PBR
0039	6-phosphogluconate dehydrogenase	-3.4	-3.2	-1.4
2334	glucose-6-phosphate 1-dehydrogenase	-3.6	-3.2	-2.7
1218	circadian clock protein KaiA	-3.5	-2.5	-3.0
2114	sensory kinase SasA	-2.4	-2.4	-1.5
0518	hypothetical protein	-1.1	-2.1	-1.4
2297	transaldolase	-2.7	-1.4	-2.4
0347	hypothetical protein	-2.3	-1.4	-1.3
1216	circadian clock protein KaiC	-1.9	-1.1	-1.0
0156	phosphoglucomutase	-0.9	-3.8	-1.1
1417	hypothetical protein	-0.8	-1.2	-1.3
0529	6-phosphogluconolactonase	-0.8	-1.0	-2.1
0244	glycogen phosphorylase	-3.7	-0.8	-2.8
2076	ribosome-associated GTPase	-0.9	-0.6	-1.1

Figure S5

SDS-PAGE analysis of KaiC and CikA stability and KaiC dephosphorylation:
(A) 17% SDS-PAGE analysis of KaiC-EA or KaiC-AE and CikA; (B) 17% SDS-PAGE analysis of KaiC and CikA;
(C): 6.5 % SDS-PAGE gels showing that KaiC remained hypophosphorylated during KaiB binding assays