Voltage- and Branch-specific Climbing Fiber Responses in Purkinje Cells

Yunliang Zang[1], Stéphane Dieudonné[2] & Erik De Schutter[1*]

[1] Computational Neuroscience Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa 904-0495, Japan
[2] Inhibitory Transmission IBENS, CNRS UMR UMR8197, INSERM U1024, Ecole Normale Supérieure, 75005 Paris, France
[*] Correspondence: erik@oist.jp

Summary

Climbing fibers (CFs) provide instructive signals driving cerebellar learning. However, conflicting experimental studies have been reported about the reliability of CF mediated Ca^{2+} influx in Purkinje cell (PC) distal dendrites. Mechanisms causing the wide variation in duration and spikelet numbers of complex spikes (CSs) have not been explored systematically. Using a new experimentally validated PC model, we describe the full range of modifiability of CF responses to explain the experimental data and make new predictions. We find voltage state gates the initiation and propagation of dendritic spikes. PC dendrites exhibit inhomogeneous excitability with individual branches as computational units for CF input. Somatic CSs are regulated by voltage state, CF activation phase and instantaneous CF firing rate. Concurrent synaptic inputs can affect CSs by modulating dendritic responses in a spatially precise way. These voltage- and branch-specific CF responses will increase dendritic computational capacity and give PCs an active role in integrating CF signals.

Introduction

According to the Marr-Albus-Ito theory (Albus, 1971; Ito, 1972; Marr, 1969), climbing fiber (CF) inputs to Purkinje cells (PCs) carry movement error information and evoke cerebellar learning by depressing the strength of parallel fiber (PF) synaptic inputs. However, how PCs process error information conveyed by CF input remains elusive. Recently, the traditional description of 'all-or-none' CF responses (Eccles, 1977) has been questioned and lead to increased attention for the modulation of CF-evoked dendritic responses as global signals (Davie et al., 2008; Kitamura and Hausser, 2011; Najafi et al., 2014). Nevertheless, conflicting observations have been reported regarding the reliability of Ca^{2+} influx in distal spiny dendrites (Ohtsuki et al., 2012; Otsu et al., 2014; Zagha et al., 2010). Whether CF-evoked dendritic responses are 'global' or 'local' determines the functional role of CF input, because the spatial range of Ca^{2+} influx controls the dendritic sites undergoing synaptic plasticity and even the direction of cerebellar learning (Coesmans et al., 2004). Intrinsically, the confusion about the range of Ca^{2+} influx stems from a lack of understanding about mechanisms governing dendritic spike generation and propagation. Furthermore, whether individual branches of PC dendritic trees are homogeneously excitable has never been investigated due to technical limitations. The effect of clustered PF synaptic inputs onto PC dendrites (Wilms and Hausser, 2015) on CF-evoked dendritic responses is also unknown. The answers to these questions will determine whether the computational unit of the CF responses is entire dendritic tree or individual branches.

Somatic Complex spikes (CSs) were assumed to be a result of synaptic input current and somatic ionic currents (Schmolesky et al., 2002). Observations that dendritic spike variation plays a minimal role in evoking an extra somatic spikelet seemed to confirm this assumption (Davie et al., 2008), however, other experiments reported contrary observations (Ohtsuki et al., 2012; Otsu et al., 2014). Additionally, the somatic CS is quite variable *in vivo* (Burroughs et al., 2017; Warnaar et al., 2015) and its duration has been linked with the degree of trial-over-trail learning (Yang and Lisberger, 2014), but the mechanism causing CS variability remains unresolved. Spikelet numbers per CS correlate with preceding simple spike (SS) firing rate (SSFR) in some studies (Burroughs et al., 2017; Gilbert, 1976), but not in others (Mano, 1970; Warnaar et al., 2015). Recently, SSFR was reported to reliably reflect the proximal dendrite voltage state in mice during self-paced locomotion (Jelitai et al., 2016). Together, these correlations motivate us to ask whether an increase or decrease of SSFR (manifestation of bi-directional voltage variation) can modulate the CF-evoked somatic and dendritic responses. Similarly, contradictory correlations between CS spikelet numbers and the instantaneous CF firing rate (Burroughs et al., 2017; Maruta et al., 2007; Warnaar et al., 2015) require further examination.

To address these questions, a systematic exploration of both somatic and dendritic responses in the whole PC is required. Given the limited simultaneous recording sites by patch-clamp (Larkum et al., 2001) and the lack of direct spike information from Ca^{2+} imaging (Deneux et al., 2016), computer models can play an indispensable role. Here we have built a 'canonical' PC model that reproduces most available experimental observations instead of fitting it to a specific cell recording. We use this model to systematically explore factors that affect CF-evoked somatic and dendritic responses.

Results

Electrophysiological properties of SSs
Before using the model to make predictions, we first compare its basic properties with experimental data. Example cell recordings are shown in Fig. 1A-D to help compare SS properties between the model and experimental data. Our PC model spontaneously fires at 40 Hz in the absence of synaptic inputs or current injections (27 - 140 Hz in experiments). SSs initiate first at the axon initial segment (AIS) with an axosomatic delay of 0.12 ms (experimentally reported ~ 0.1 ms by Palmer et al. (2010)). The SS peak is -1 mV and its amplitude is 68 mV (-11 - -3 mV and 56 - 68 mV respectively in experiments). SSs have a half-amplitude duration of 0.22 ms (0.22 - 0.28 ms in experiments). The peak *dv/dt* of SSs is 530 mV/ms (440 - 540 mV/ms in experiments). The d^2v/dt^2 of SSs shows a biphasic increase, with the first component reflecting the contribution of axial current from the AIS and the second one reflecting the contribution of somatic Na^+ current. The peak value of d^2v/dt^2 is 6600 mV/ms^2 (5600 - 8200 mV/ms^2 in experiments).

Due to the absence of local Na^+ channels, SSs fail to invade the dendritic tree. The distance-dependent passive decay of SS amplitudes in the model matches experimental data by Ohtsuki et al. (2012)(Fig. 1E). The model also performs well in response to somatic current injections. The F-I curve falls completely within the experimentally measured range (Fig. 1F). When a large enough negative current is injected, the cell is silenced and shows a typical 'sag' response (illustrated in the inset of Fig. 1G). Both the peak and steady state of the 'sag' response are within the range

Figure 1. Model properties compare well with experimental data.
A. Spontaneously firing somatic SSs and dendritic membrane potentials (103 μm from soma) from the model (upper panel) and example experimental recording (lower panel, dendritic site is ~ 100 μm from soma). **B.** Single somatic SS waveforms from the model (upper panel) and experiment (lower panel) are shown for comparison. The spike at the AIS is shown with somatic SS to measure the axosomatic delay in the model. The *dv/dt* and *d²v/dt²* of the somatic spike in the model (upper panel) are compared in shape and amplitude with experimental recordings (lower panel) in **C** and **D** separately. **E**, Distance-dependent decay of somatic SS amplitudes in the model and experimental recordings (data from Ohtsuki et al. (2012)). **F**, The F-I curve of the model compared with experimental recordings. Red circles represent the F-I curve of the example cell used in A-D. **G**, The peak values (the strongest hyperpolarization) and steady state values of 'sag' responses caused by negative current injections are compared between model and experimental recordings. Red symbols represent the data from the example cell used in A-D. **H**, Distinct spike (blue traces) peaks and principal ionic current profiles in intact and dissociated PCs. In the dissociated PC model, only the dendritic root section before the first bifurcation is preserved.

of experimental data (Fig. 1G) (All the preceding experimental data except Fig. 1E were shared by Ede Rancz and Michael Häusser from UCL). This good overall performance in response to current injections suggests that the model is reliable in predicting excitatory and inhibitory synaptic responses in PCs.
Dissociated PCs are often used to investigate the role of ionic currents in PC electrophysiological properties (Carter and Bean, 2009; Khaliq et al., 2003; Swensen and Bean, 2003). However, through the model, we find that the profiles of Na current (principal depolarization current (Carter and Bean, 2009)) and of Kv3 current (principal repolarization current (Martina et al., 2007)) in an intact PC differ from in a dissociated PC owing to their distinct spike properties (Fig. 1H). The SS peak in the intact PC (-1 mV) is much lower than the dissociated PC (+26 mV in the model and +32 mV in experiments by Carter and Bean (2011)). Consequently, due to their high

activation threshold (Martina et al., 2007), a much smaller fraction of Kv3 channels
activate in an intact PC compared to a dissociated PC. In the intact PC, the passive
dendrite repolarizes the somatic spike together with the 'less' activated Kv3 current.
This large axial current (flow into dendrite) keeps pace with the somatic spike
depolarization to constrain the spike peak to -1 mV (a smaller axial current exists in
the dissociated PC due to the remaining dendritic stump). At the peak of the spike, the
'dip' in the Na^+ current observed in dissociated PCs (Carter and Bean, 2009) is absent
in the intact PC model. The Na^+ entry ratio (defined as the ratio of total Na^+ influx to
the Na^+ influx during the rising phase (Carter and Bean, 2009)) decreases from 3.1 in
the dissociated PC model to 2.2 in the intact PC model. The decreased Na^+ entry ratio
in the intact PC suggests that dendrites lower the metabolic cost of somatic spikes.

Generation of CF responses in the PC
140

Figure 2. The generation of CF responses in the PC.
A. CF responses recorded at different sites of the PC model. Two sites on each main branch are
selected as indicated in the figure. Point A (smooth dendrite) and point B (spiny dendrite) on branch 1;
Point C and point D on branch 2; Point E and point F on branch 3. CF responses at dendritic root
section, soma and AIS are labeled by the corresponding arrows. **B.** The peak *dv/dt* of the first spikelet
in the CS increases compared with a SS in both the model (upper panel) and experiment (lower panel,
the data is from Warnaar et al. (2015). **C.** The axosomatic delay of the first spikelet in the CS decreases
compared with that in the SS. **D.** The composition of the somatic CS. Currents at the initial
depolarization phase are enlarged in the inset. Dendritic currents at the smooth dendrite (point A) and
spiny dendrite (point B) are shown in **E** and **F** in sequence. In D-F. Ionic currents are computed by
multiplying the density of currents in the central segment of the corresponding section with their

surface areas (Carnevale and Hines, 2006). Surface areas of the segments chosen at the soma, smooth
dendrite and spiny dendrite are 204 μm², 64 μm², and 23 μm², respectively. CF EPSC is absent in the
spiny dendrite. Notice that different scales are used in D-F, due to different segment surface areas.
156
When PCs receive synaptic inputs from CFs, stereotypical somatic CSs occur at their
somas. CF responses at different sites of a spontaneously firing PC model (no holding
currents or background synaptic inputs) are shown in Fig. 2A. The peak dv/dt of the
first spikelet in the CS is ~ 270 mV/ms larger than that of the SS, as confirmed by
experimental data (Fig. 2B). The increased depolarization speed can facilitate the
propagation of the first spikelet down to the cerebellar nuclei (CN). In the model,
similar to experimental observations (Palmer et al., 2010), the CS still initiates first at
the AIS. The first spikelet has an axosomatic delay of 0.06 ms, which is much shorter
than that of the SS (Fig. 2C). This decreased axosomatic delay is caused by the
depolarizing CF EPSC during the initial depolarization phase (Fig. 2D). The model
also replicates the varied CF responses due to bursting CF input (Mathy et al., 2009)
(Fig. S1).
In agreement with previous experimental findings (Swensen and Bean, 2003), the
SK2 current terminates the CS (Fig. 2D). Each spikelet of the somatic CS is sharp
(Fig. 2A), suggesting they are generated by Na^+ channels (Fig. 2D). Two factors
determine the absence of Ca^{2+} spikes at the soma. First, the P-type Ca^{2+} current is
much smaller than the Na^+ current; therefore, the Na^+ current shunts it. Second, K^+
currents are always larger than the P-type Ca^{2+} current (Fig. 2D). Magnitudes of the
CF EPSC and net axial current are comparable to ionic currents at the soma (Fig. 2D),
suggesting an important role in generating somatic CSs. Consequently, variation of
either the CF EPSC or dendritic axial current has the potential to alter somatic CSs, as
will be shown in the following sections.
The dendritic root section (the section between the soma and the first dendritic
bifurcation) is electrically compact with the soma and shows mainly a passive
propagation of a somatic CS. With distance from the soma, the clamping effect of a
somatic CS on the dendrite weakens. The P-type Ca^{2+} current gradually dominates
and cause Ca^{2+}-driven dendritic responses in the PC dendrite (Fig. 2A, E-F).

Voltage state regulates CF-evoked dendritic responses
Motivated by the variable voltage states of PCs observed *in vivo* (Jelitai et al., 2016),
we systematically explore how voltage state regulates CF-evoked dendritic responses
(Fig. 3). Voltage states of the PC model are manipulated by varying the somatic
holding currents. Depolarization facilitates higher peaks of dendritic voltage
responses with further propagation into distal spiny dendrites (Fig. 3A-C). In distal
spiny dendrites, axial currents from more proximal smooth dendrites are the only
current sources that can depolarize to approach the Ca^{2+} spike threshold and trigger a
local dendritic spike (Fig. 3J-L). When the soma is held at -76 mV (Fig. 3A), the A-
type Kv4 current is always larger than the P-type Ca^{2+} current in distal spiny dendrites
(Fig. 3J). Consequently, axial currents cause only small passive depolarizations in
spiny dendrites (Fig. 3D,J), and P-type Ca^{2+} channels are hardly activated (Fig. 3J). In
proximal smooth dendrites, the CF provides powerful synaptic input to depolarize
(Fig. 3D,M). However, this CF-evoked depolarization is still passive because the local
Kv4 current and axial currents (flow into distal parts) stop the P-type Ca^{2+} current
from dominating depolarization (Fig. 3M). Accordingly, the dendritic voltage
responses decrease with distance from the soma in both proximal smooth dendrites
and distal spiny dendrites (Fig. 3G). Distance-dependent propagation of dendritic
202

Figure 3. Voltage states regulate CF-evoked dendritic spike generation and propagation.
A-C. Color-coded representation of dendritic voltage responses at different holding potentials (-76 mV, -70 mV and -61 mV respectively). **D-F.** Example somatic CSs (black), dendritic voltage responses on the smooth dendrite (red) and spiny dendrite (blue) at different voltage states. In **E**, an extra example dendritic voltage response (green trace, the position is labeled in B) on the spiny dendrite is shown. **G-I.** Distance-dependent propagation of dendritic responses at different voltage states. Experimental data (squares) shown in G are from Ohtsuki et al. (2012). **J-L.** Ionic currents in the distal spiny dendritic site at different voltage states. **M-O.** Ionic currents in the proximal smooth dendritic site at different

voltage states. The surface areas of the segments chosen at the smooth dendrite and spiny dendrite are
64 μm^2 and 23 μm^2 respectively. **P-R.** Color-coded peak Ca^{2+} concentrations at corresponding
membrane potentials.

voltage responses matches experimental recordings under the same conditions
(Ohtsuki et al., 2012) (Fig. 3G).
As the soma is held at -70 mV (Fig. 3B), dendritic spikes occur in part of the dendritic
tree (the left half, proximal to the soma), where the voltage responses now increase
with distance from the soma. However, the responses still decay with distance from
the soma in other parts of the dendritic tree (Fig. 3B,E,H). With depolarization of the
soma to -61 mV (Fig. 3C), the Kv4 current gradually inactivates and becomes smaller
than the P-type Ca^{2+} current in both smooth and spiny dendrites during the initial
depolarization phase (Fig. 3L,O). As a result, the P-type Ca^{2+} current depolarizes the
dendrites and activates more Ca^{2+} channels in a positive feedback loop until the Kv3
current is highly activated to repolarize the spikes. Because dendritic spikes now
occur globally, its peak increases with distance from the soma (Fig. 3F,I). In line with
the modifiable voltage responses, the CF-evoked dendritic Ca^{2+} influx is also
facilitated by the depolarization, causing larger and more widespread elevations in
dendritic Ca^{2+} concentration (Fig. 3P-R). Additional CF-evoked voltage and Ca^{2+}
responses at other voltage states are shown in Fig. S2.
Taken together, these results suggest that the CF-evoked Ca^{2+} influx can be gated as
either a local or global signal by the voltage state of PCs. Previous seemingly
conflicting observations regarding the spatial range of Ca^{2+} influx fit well with our
new mechanism (Kitamura and Hausser, 2011; Najafi et al., 2014; Ohtsuki et al.,
2012; Otsu et al., 2014; Zagha et al., 2010). Similarly, the discrepant dendritic spike
waveforms recorded by patch-clamp from different groups can also be reconciled by
distinct voltage conditions (Davie et al., 2008; Ohtsuki et al., 2012) in Fig. S3.

Inhomogeneous excitability of the dendritic tree
Dendritic spikes can occur in one part of the dendritic tree, without occurring in other
parts (Fig. 3B), suggesting an asynchronous occurrence of the first spikelet at
different sites of the dendrite. Similar to the first dendritic spikelet, the secondary
dendritic spikelet occurs asynchronously in different branches around its threshold
(Fig. 4). The secondary dendritic spikelet initially occurs at branch 1 (points A and B
in Fig. 2A); then it gradually occurs at branch 2 (points C and D in Fig. 2A) and
finally occurs at branch 3 (points E and F in Fig. 2A) with slight increasing
depolarization. Even when secondary dendritic spikelets occur globally, their time to
peak still differs between branches (Fig. 4C). Secondary dendritic spikelets peak at
8.36 ms, 10.74 ms and 13.64 ms at point A, point C and Point E respectively (relative
to CF activation time). With further depolarization, secondary dendritic spikelets peak
more synchronously (with 0.89 nA current injection, the respective values become
6.06 ms, 6.88 ms and 7.32 ms). The asynchronous occurrence of individual dendritic
spikelets reflects the inhomogeneous excitability of each dendritic branch. In the
model, the density of ionic currents is homogeneous in the spiny dendritic tree.
Therefore, we investigate whether morphological differences between individual
branches can account for their heterogeneity in excitability. The excitability of each
branch is determined by the ratio of spiny dendrite area to smooth dendrite length in
that branch (essentially the ratio of spiny dendrite capacitance load to CF synaptic
input, 'load/input' ratio, see **Methods**). The 'load/input' ratios are 69, 96 and 130 at
branches 1, 2 and 3 respectively, explaining their heterogeneous excitability.
Moreover, within each 'big' branch, 'small' child branchlets also show heterogeneity

Figure 4. Branch-specific occurrence of secondary dendritic spikelets.
In all simulations, the holding current is -0.17 nA. CF input is activated at three different dendritic voltage states (different phases of somatic interspike intervals) as shown in **A**, **B**, and **C**. Points A-F are defined in Fig. 2.

in excitability, which are caused by other morphological factors such as the electrotonic distance to smooth dendrites (Fig. 3A-C).

To test the role of Kv4 current in heterogeneous dendritic excitability, we simulate the CF-evoked dendritic responses after globally decreasing its activation (left shift of the steady state availability by 11 mV). As expected, after reducing the Kv4 current, both the first and secondary dendritic spikelets occur at more hyperpolarized voltage states (Fig. S4). The whole dendrite also gets close to homogeneous at a more hyperpolarized level. Essentially the uneven distributed 'load/input' ratios underlie the heterogeneous dendritic excitability, but dendritic Kv4 channels enhance this heterogeneity.

Voltage- and phase-dependent somatic CSs
CF-evoked dendritic responses can be strongly regulated by PC voltage states. Similarly, somatic CSs also vary significantly with the depolarization of PC (Fig. 5A-C). The number of spikelets in CSs increases with initial depolarization, but decreases with further depolarization. A somatic CS results from the interaction of intrinsic ionic currents, axial current, and CF EPSC (Fig. 2D). At low voltage, although Na^+ channel availability decreases slightly (Fig. 5C), increased axial current from larger dendritic voltage responses (Fig. 5A) dominates and triggers more somatic spikelets with initial depolarization. Interestingly, the largest number of somatic spikelets coincides with the longest occurrence latency of secondary dendritic spikelets (Fig. 5A). With further depolarization, Na^+ channel availability decreases at a higher rate

Figure 5. Voltage- and phase-dependent somatic CSs.
A. Voltage states affect somatic CSs and corresponding dendritic responses. The definition of dendritic sites is the same as in Fig. 2. The required holding currents and the firing rates at each voltage state are listed on the left. **B.** Voltage states modulate CS durations, represented by the timing of the last spikelets. **C.** In parallel with Na$^+$ channels availability (blue circles), amplitudes of the first three spikelets in the CS (black circles) decrease with depolarization. **D.** Amplitudes of the first and second spikelets are phase-dependent in both the model (upper left panel) and experimental recordings (lower panel). Phase-dependent recovery of Na$^+$ channel availability is shown in upper right panel. Experimental data are from Warnaar et al. (2015).

(Fig. 5C). Also, secondary dendritic spikelets peak much faster and closer to a preceding somatic spikelet (Fig. 5A). The shortened interval between secondary dendritic spikelets and their preceding somatic spikelet means fewer Na$^+$ channels are recovered to evoke an extra somatic spikelet (right axis of Fig. 5D). The critical role

of secondary dendritic spikelet timing in evoking an extra somatic spikelet agrees
with previous experimental observations (Davie et al., 2008). Both factors determine
the decreased somatic spikelets with further depolarization. Concurrent with spikelet
numbers, CS durations are regulated by voltage states (or SSFRs), as represented by
the timing of the last spikelets (Fig. 5B).
Voltage states also affect individual spikelet amplitudes in CSs. We analyzed the first
three spikelets, which exist in all investigated voltage conditions. Spikelet amplitudes
decrease with depolarization due to reduced availability of Na^+ channels (Fig. 5C).
Individual spikelet amplitude is critical because it determines the probability of
propagation down to the CN (Khaliq and Raman, 2005).
Finally, we observed phase-dependency of somatic CSs. Amplitudes of the first two
spikelets in CSs decrease with shortening of the interval between the CF activation
and its preceding SS (Fig. 5D). This phase-dependency reflects the time course of Na^+
channel recovery from inactivation (Fig. 5D). Subsequently, we confirmed this phase-
dependency by reanalyzing *in vitro* experimental recordings (Warnaar et al., 2015),
Fig. 5D).
So far, voltage states of the PC model were manipulated by varying the holding
currents at the soma. We implemented further simulations, in which the net balance of
excitatory PF and inhibitory synaptic inputs determines the dendritic voltage state
changes, reflecting the situation *in vivo* (Jelitai et al., 2016). We find that the effect of
voltage state on CF-evoked somatic and dendritic responses found *in vitro* also holds
in vivo (Fig. S5).

Spatially constrained modulation by clustered PF/stellate cell synaptic input
Sensory stimuli can activate clustered PF synaptic inputs *in vivo* (Wilms and Hausser,
2015). Here we examine how simultaneous clustered PF or stellate cell synaptic
inputs regulate CF responses. We find that clustered synaptic inputs preferentially
regulate dendritic responses in a spatially constrained manner. A secondary dendritic
spikelet only occurs at branch 1 with CF input in isolation (Fig. 6A). With 5 PF
synapses simultaneously activated at the indicated branchlet of branch 2, the
secondary dendritic spikelet also occurs at branch 2. Concurrently, an extra spikelet is
evoked in the somatic CS. Nonetheless, clustered PF synaptic inputs within branch 2
do not significantly affect the dendritic responses at branches 1 and 3. Seemingly
paradoxically, the extra somatic spikelet is eliminated if the number of simultaneously
activated PF synapses increases to 20. As explained in the previous section, Na^+
channels need time to recover from inactivation after each somatic spikelet.
Increasing the number of PF synaptic inputs accelerate the occurrence of secondary
dendritic spikelets, making them occur closer to the preceding somatic spikelet, and
reducing the likelihood of triggering an extra spikelet due to fewer recovered Na^+
channels. Similarly, simultaneous activation of 10 stellate cell synaptic inputs can
evoke an extra somatic spikelet by delaying the occurrence of the secondary dendritic
spikelet at branch 2 (Fig. 6B). Again, dendritic responses at other branches are
minimally affected. The secondary dendritic spikelet still occurs in the proximal
smooth dendrite even when it is eliminated in the distal part by activating 20 stellate
cell synaptic inputs.
We also show that compartment-specific dendritic excitability (Ohtsuki et al., 2012),
simulated here by regional Kv4 current down-regulation, modulates dendritic
responses in a spatially precise way (Fig. S6).

Figure 6. Simultaneous clustered PF or stellate cell synaptic input regulates the CF-evoked somatic CSs by locally modulating the dendritic responses.
A. Somatic and dendritic responses with only CF input, CF input + 5 PF synaptic inputs and CF input + 20 PF synaptic inputs are shown from top to bottom. PF synapses are randomly distributed on the child branchlet of the indicated dendritic section (green dots). **B.** Somatic and dendritic responses with only CF input, CF input + 10 stellate cell synaptic inputs and CF input + 20 stellate cell synaptic inputs are shown from top to bottom. Similar placement of stellate cell synapses (orange dots). The definition of recorded dendritic sites is the same as in Fig. 2.

Paired-pulse depression regulates CF responses in the PC

The CF activation rate can transiently reach ~ 5 - 8 Hz after sensory stimuli *in vivo* (Najafi et al., 2014; Warnaar et al., 2015). Different intervals between two consecutive CF signals (instantaneous CF firing rate) can correlate with varied spikelet numbers in CSs. However, inconsistent observations have been reported (Burroughs et al., 2017; Maruta et al., 2007; Warnaar et al., 2015) and the mechanism

Figure 7. PPD regulates somatic and dendritic responses.
Examples of CF responses (both somatic and dendritic) with unchanged (case I), decreased (case II) and increased (case III) spikelet numbers per CS at decreased CF EPSC are shown in **A**, **B** and **C** in sequence. The effect of decreased CF EPSC on the spikelet number and duration of the CS is shown in **D**.

underpinning the correlation remains unclear. Paired-pulse depression (PPD) has been demonstrated for CF synaptic inputs *in vitro* (Andjus et al., 2005) and may be a candidate cause. To investigate the effect of PPD on CF responses, we systematically explore CF responses with baseline CF EPSC or 80% of baseline CF EPSC (~ 4 Hz of instantaneous CF firing rate in PPD experiment by Andjus et al. (2005)). We find a smaller CF EPSC consistently reduces the dendritic responses. At low voltage, the peak of the dendritic response decreases (Fig. 7A); at higher voltage, the appearance of a secondary dendritic spikelet is prevented (Fig. 7B) or delayed (Fig. 7C). The effect of PPD on the somatic CSs is more complicated. The spikelet numbers in CSs show no changes, decreases or increases with a smaller CF EPSC at different voltage states (Fig. 7A-C). A smaller CF EPSC always delays occurrence of individual somatic spikelets by providing less depolarization current. Therefore, PPD increases CS duration if the number of spikelets either does not change or increases (case I and case III in Fig. 7D). When a smaller CF EPSC evokes fewer somatic spikelets, CS

duration decreases (case II in Fig. 7D). Voltage state ranges for the occurrence of
cases I-III depend on the degree of PPD. For example, when comparing with 90% of
baseline CF EPSC, case I can occur at both low and high voltages (data not shown).
Therefore, PPD can bridge the gap between instantaneous CF firing rates and CS
variations. This new finding also reconciles the unchanged (Warnaar et al., 2015),
decreased (Maruta et al., 2007), and increased (Burroughs et al., 2017) number of
somatic spikelets at higher instantaneous CF firing rates (smaller CF EPSC) observed
in experiments.

Discussion

We built a new PC model based on extensive experimental data to systematically
investigate CF-evoked dendritic and somatic responses. Simulation results help to
reveal the nature of different forms of spikes in PCs, reconcile conflicting
experimental data and make new predictions. We demonstrate that CF-evoked
dendritic responses are voltage-dependent and individual branches behave as
computational units. Branch-specific computation can be strengthened by concurrent
synaptic inputs or dendritic plasticity. We further correlate CS properties with voltage
states (or SSFRs) and identify PPD as the cause of SSFR-related CS variations. Our
findings offer a new perspective on how PCs dynamically process information from
CF inputs.

Model accuracy
It has been demonstrated that a data set recorded from two locations in the same
neuron is insufficient to accurately constrain a compartment model (Keren et al.,
2005). We constrained model parameters by hand tuning and tried to replicate as
many experimental observations as possible. We started by constraining somatic ionic
currents based on ionic current measurements taken from dissociated PCs. These data
allowed us to uniquely constrain principal somatic currents. Given the larger surface
area of the soma compared to the AIS, somatic ionic currents are extremely important
for PC electrophysiological properties. Subsequently, the ionic currents in the AIS
and dendrites were constrained to obtain the spike properties summarized in Fig. 1. It
was more difficult to constrain dendritic ionic currents due to a lack of sufficient data.
The total Ca^{2+} influx during one dendritic spikelet was constrained by the
experimentally estimated value (see **Supplementary Information**). In addition, the
replicated CF-evoked dendritic responses under different conditions (Figs. 3,S2,S3)
suggest that the dendrite model produces realistic results. Nonetheless, there is still
room for improvement in the future. Due to the lack of specific data, we did not try to
reproduce the role of SK2 current (Ohtsuki et al., 2012) and Kv1 current (Khavandgar
et al., 2005) in regulating dendritic excitability, but such changes are not expected to
affect our model predictions qualitatively.
The detailed PC model presented here follows the tradition of the biophysical
modeling by De Schutter and Bower (1994a, 1994b), which has successfully
predicted experimental findings (Steuber et al., 2007). Although the 1994 model is
clearly outdated by recent experimental findings on PC channels, there was no
suitable replacement until now. A popular PC model by Khaliq et al. (2003) has no
dendrite and can therefore not be used to simulate synaptic responses. A more recent
PC model (Masoli et al., 2015) has not been validated extensively against
experimental data. For example, it fails to produce the stereotypical 'sag' responses to
hyperpolarization.

Physiological predictions and implications

The model replicates spiking properties of both intact and dissociated PCs in detail. This allows us to critically evaluate predictions about the metabolic cost of spiking in PCs. Generation of spikes is energetically expensive. In a cortical pyramidal neuron, $\sim 400 \times 10^6$ ATP molecules are consumed to restore the Na^+ and K^+ ion gradients after each spike (Attwell and Laughlin, 2001). Carter and Bean (2009) reported that spiking is energetically less efficient in PCs compared with cortical pyramidal neurons. However, this study was performed in dissociated PCs. In our model, the intact PC has a smaller Na^+ entry ratio compared with the dissociated PC (Fig. 1H), suggesting that dendrites increase the metabolic efficiency of somatic spikes. Given the energetic cost of spikes, it will be interesting to explore whether this dendritic role also applies to other neurons in future work.

This model allows us to systematically explore the origin of spikes observed at different parts of a PC and how they can be modulated. Dendritic spikes and somatic CSs are generated by different channels intrinsically, but they communicate via the axial currents through the dendritic root section (Fig. 2). Due to technical challenges, dendritic patch-clamp recording is usually performed on smooth dendrites of large diameter. Though we could reproduce the experiment by Davie et al. (2008) in the model (Fig. S7), we show that a single-site dendritic recording cannot reliably reflect spike occurrence and propagation in the rest of the dendritic tree. Both the peak and occurrence latency of secondary dendritic spikelets can be modulated. The peak of secondary dendritic spikelets decreases in the somatopetal direction due to unevenly distributed impedances (Roth and Hausser, 2001) and K^+ channels (Martina et al., 2003). When a secondary dendritic spikelet is 'weak', with small amplitude and long latency, it may fail to trigger a regenerative dendritic spike in the proximal dendrites (branch 1 in Fig. S7). By single-site dendritic patch-clamp, a dendritic response showing one spikelet may not correspond to a one-spikelet spike in the rest of the dendrite and therefore the role of dendritic spikes in evoking extra somatic spikelets may have been underestimated.

Our results suggest a new mechanism for PCs to process the information from CF inputs: the voltage states of PCs gate the initiation and propagation of CF-evoked dendritic responses (Fig. 3,S2,S5). This new mechanism not only reconciles the conflicting observations regarding the spatial range of CF-evoked Ca^{2+} influx (Kitamura and Hausser, 2011; Najafi et al., 2014; Ohtsuki et al., 2012; Otsu et al., 2014; Zagha et al., 2010), but also can play a functional role in regulating CF efficacy to induce plasticity in cerebellum learning. The Marr-Albus-Ito theory posits that error signals evoke CF inputs, which trigger plasticity at PF synapses (Albus, 1971; Ito, 1972; Marr, 1969). The voltage-modulated spatial range and amplitude of dendritic responses indicate that PCs may play active roles in response to error signals from CF inputs rather than passively accepting instruction from their CF 'teachers'. PF synapses can undergo not only LTD, but also long-term potentiation (LTP) (Lev-Ram et al., 2002). PF-LTD and -LTP are both Ca^{2+}-dependent, with a higher Ca^{2+} induction threshold for LTD (Coesmans et al., 2004; Piochon et al., 2016). The higher induction threshold of LTD means CF may trigger LTP instead of LTD, if the CF-evoked Ca^{2+} influx does not reach the induction threshold (Carey and Regehr, 2009; Coesmans et al., 2004). When PCs are relatively hyperpolarized, i.e. at low SSFRs *in vivo* (Jelitai et al., 2016), CF activation evokes very limited Ca^{2+} influx in the spiny dendrites and PF synapses may undergo LTP or no change instead of LTD; when PCs undergo slight depolarization, CF activation evokes dendritic spikes in only part of the distal dendrites and LTD will probably just occur in that part; when PCs are

highly depolarized (high SSFR), Ca^{2+} influx can be modulated by the number of spikelets in the dendritic spikes to affect the degree of LTD. Such effects will be further modified by stochastic gating of Ca^{2+}-activated K^+ channels (Anwar et al., 2013), a phenomenon that was not simulated in this study.

The PC dendritic tree exhibits inhomogeneous excitability in individual branches, suggesting a computational unit at the level of individual branches, or even smaller branchlets (Fig. 4). Building upon the modulated voltage states of PCs, this new concept of branch-specific computation can undoubtedly increase the capacity of information processing in PC dendrites, in contrast to the traditionally view of the whole dendrite as the computational unit. Notably, branch-specific plasticity has already been demonstrated in other neurons such as L5 pyramidal neurons *in vitro* (Golding et al., 2002) and *in vivo* (Cichon and Gan, 2015). When PCs receive concurrent clustered synaptic input (Wilms and Hausser, 2015) or undergo localized dendritic excitability plasticity (Ohtsuki et al., 2012), CF-evoked dendritic responses can be modulated more precisely in space and magnitude to affect the plasticity induction (Fig. 6, Fig. S6).

Recently, CS durations have been linked to the degree of trial-over-trial learning in monkeys (Yang and Lisberger, 2014), with the mechanism underlying the variable CSs unresolved. The authors assumed that the variation of CS durations is due to the bursting of CF inputs, based on Mathy et al. (2009). We correlate the spikelet numbers and durations of CSs with voltage states or SSFRs (Fig. 5A-B). Besides reconciling previous contradictory correlations between spikelet numbers and SSFRs (Burroughs et al., 2017; Gilbert, 1976; Mano, 1970; Warnaar et al., 2015), our results provide an alternative interpretation of the work by Yang and Lisberger (2014). Our interpretation highlights the importance of all synaptic inputs to the PCs and resulting SSFR, not just CF, to control learning. Additionally, the CS spikelet amplitudes can be modulated by voltage states (Fig. 5C,). The timing of CF activation (phase-dependency) and concurrent synaptic input also affect somatic CSs (Fig. 5D, 6). All these factors change CS waveforms to vary the PC output and affect inhibition of the CN.

In reality, the CF input itself is not 'all-or-none' either. The variable number of spikes per CF burst can affect both somatic and dendritic responses (Mathy et al., 2009) (Fig. S1). Furthermore, CF synaptic inputs can be regulated by PPD (Andjus et al., 2005) and by norepinephrine (Carey and Regehr, 2009). Both PPD and norepinephrine can reduce the CF EPSC to modulate somatic and dendritic responses (Fig.7). In particular, PPD bridges the gap between instantaneous CF firing rates and varied CS spikelet numbers (Burroughs et al., 2017; Maruta et al., 2007; Warnaar et al., 2015) and play an important role in regulating CF responses. It is reasonable to infer that both CF inputs and PC voltage states are modulated in concert to optimize the transmission of error information down to PCs.

Conclusion

To the best of our knowledge, this study is the first to systematically explore CF responses in the PC. We believe that multiple mechanisms working synergistically to modulate CF responses will have important consequences for cerebellar learning an for signals transmitted to the CN. We hope this study will inspire experiments to directly validate modulation of CF responses *in vivo*. Special attention should be paid to PCs with low SSFRs.

Methods

All simulations were performed with NEURON, version 7.4 (Hines and Carnevale, 1997). We use the morphology of a 21-day-old Wistar rat PC (Roth and Hausser, 2001). The model consists of an axon initial segment (AIS), a soma, smooth dendrites, and spiny dendrites. The first 17 μm of the reconstructed axon are preserved as the AIS. When exploring the spike initiation site, AIS indicates the end of AIS next to the myelinated axon. The definition of other morphological parts is the same as in the original work (Roth and Hausser, 2001). The model has the following passive parameters: $R_m = 120.2$ kΩ cm^2, $R_i = 120$ Ω cm, $C_m = 0.64$ μF/cm^2 (Roth and Hausser, 2001). To compensate for the absence of spines in the reconstructed morphology, the conductance of passive current and C_m are scaled by a factor of 5.3 in the spiny dendrite and 1.2 in the smooth dendrite. The AIS contains Na$^+$ current, T-type Ca^{2+} current, P-type Ca^{2+} current, H-current, high-threshold-activated K$^+$ current (Kv3 current), large-conductance Ca^{2+}-activated K$^+$ current (BK current), and a small-conductance Ca^{2+}-activated K$^+$ current (SK2 current). Current distribution in the soma is similar to the AIS without the T-type Ca^{2+} current. T-type Ca^{2+} current, P-type Ca^{2+} current, H-current, Kv3 current, BK current, SK2 current, Kv1 current, and Kv4 current are distributed in both the smooth dendrite and the spiny dendrite. The conductance densities of ionic currents in corresponding parts of the model were found by hand tuning. Detailed information about ionic current equations and the process of parameter tuning can be found in the **Supplementary Information**. The source code of this model is available at ModelDB (future link).

The synaptic conductance activated by CF input is approximated as a bi-exponential waveform, with 0.3 ms and 3 ms as the rise and decay time constants (Davie et al., 2008). Unless otherwise specified, baseline peak conductance is 1.2 nS. 500 synaptic inputs are distributed on the soma and smooth dendrite at a constant density per length to simulate the CF response. The conductance varies in the range of 80% - 100% to simulate how the EPSC changes by paired-pulse depression (PPD). Due to the elaborate structure, the dendrite is not isopotential with the soma when holding current is injected at the soma. The somatic membrane potential is selected to represent the voltage state of the whole PC except in Fig. 4, in which dendritic membrane potential is used. When measuring the distance-dependence of the dendritic response peak, we selected the first dendritic spikelet if there were two spikelets. To measure excitability of individual branches, we define a new measure as the ratio of the spiny dendrite capacitance load to CF synaptic input, 'load/input' ratio, which can be transformed into 'the area of spiny dendrite'/'length of smooth dendrite' within each branch in the model. To exclude the effect of phase-dependence on CSs in Fig. 5, all the CF signals are activated with 3 ms after the preceding SS under different membrane potentials. To explore how simultaneous clustered synaptic inputs regulate dendritic response and somatic output, PF or stellate cell synaptic inputs are activated 5 ms after the CF input (Fig. 6). To simply mimic the PC voltage states *in vivo*, 1105 PF synapses and 1105 Inhibitory synapses are uniformly distributed on the spiny dendrites. Different voltage states of the PC are achieved by randomly activating excitatory and inhibitory synapses at different average firing rates.

Data availability. All the simulation data and the model code are available from the authors.

Acknowledgements
We thank Ede Rancz, Michael Häusser, Gen Ohtsuski, Christian Hansel, João Couto
(the data shared by above-mentioned people were used in the manuscript for
validation of the model results), Bruce Bean, Indira Raman (the data shared by above-
mentioned people were used to constrain ionic current properties), Mathew Nolan,
Edward Zagha and Bernardo Rudy for sharing related experimental data.
588
589 **Author Contributions**
590 Y.L.Z., S.D., and E.D.S. conceived this study; Y.L.Z. performed all the simulations;
591 Y.L.Z. and E.D.S. analyzed the simulation data; Y.L.Z. and E.D.S. wrote the
592 manuscript.
593
References
Albus, J.S. (1971). A theory of cerebellar function. Mathematical Biosciences *10*,
25-61.
Andjus, P.R., Bajic, A., Zhu, L., Schachner, M., and Strata, P. (2005). Short-term
facilitation and depression in the cerebellum: some observations on wild-type
and mutant rodents deficient in the extracellular matrix molecule tenascin C.
Annals of the New York Academy of Sciences *1048*, 185-197.
Anwar, H., Hepburn, I., Nedelescu, H., Chen, W., and De Schutter, E. (2013).
Stochastic calcium mechanisms cause dendritic calcium spike variability. The
Journal of neuroscience : the official journal of the Society for Neuroscience *33*,
15848-15867.
Attwell, D., and Laughlin, S.B. (2001). An energy budget for signaling in the grey
matter of the brain. Journal of cerebral blood flow and metabolism : official
journal of the International Society of Cerebral Blood Flow and Metabolism *21*,
1133-1145.
Burroughs, A., Wise, A.K., Xiao, J.Q., Houghton, C., Tang, T.Y., Suh, C.Y., Lang, E.J.,
Apps, R., and Cerminara, N.L. (2017). The dynamic relationship between
cerebellar Purkinje cell simple spikes and the spikelet number of complex spikes.
J Physiol-London *595*, 283-299.
Carey, M.R., and Regehr, W.G. (2009). Noradrenergic control of associative
synaptic plasticity by selective modulation of instructive signals. Neuron *62*, 112-
122.
Carnevale, N.T., and Hines, M.L. (2006). The NEURON book (Cambridge, UK ; New
York: Cambridge University Press).
Carter, B.C., and Bean, B.P. (2009). Sodium entry during action potentials of
mammalian neurons: incomplete inactivation and reduced metabolic efficiency
in fast-spiking neurons. Neuron *64*, 898-909.
Carter, B.C., and Bean, B.P. (2011). Incomplete inactivation and rapid recovery of
voltage-dependent sodium channels during high-frequency firing in cerebellar
Purkinje neurons. Journal of neurophysiology *105*, 860-871.
Cichon, J., and Gan, W.B. (2015). Branch-specific dendritic Ca(2+) spikes cause
persistent synaptic plasticity. Nature *520*, 180-185.
Coesmans, M., Weber, J.T., De Zeeuw, C.I., and Hansel, C. (2004). Bidirectional
parallel fiber plasticity in the cerebellum under climbing fiber control. Neuron
44, 691-700.

Davie, J.T., Clark, B.A., and Hausser, M. (2008). The origin of the complex spike in
cerebellar Purkinje cells. The Journal of neuroscience : the official journal of the
Society for Neuroscience *28*, 7599-7609.
De Schutter, E., and Bower, J.M. (1994a). An active membrane model of the
cerebellar Purkinje cell II. Simulation of synaptic responses. Journal of
neurophysiology *71*, 401-419.
De Schutter, E., and Bower, J.M. (1994b). An active membrane model of the
cerebellar Purkinje cell. I. Simulation of current clamps in slice. Journal of
neurophysiology *71*, 375-400.
Deneux, T., Kaszas, A., Szalay, G., Katona, G., Lakner, T., Grinvald, A., Rozsa, B., and
Vanzetta, I. (2016). Accurate spike estimation from noisy calcium signals for
ultrafast three-dimensional imaging of large neuronal populations in vivo.
Nature communications *7*, 12190.
Eccles, J.C. (1977). An instruction-selection theory of learning in the cerebellar
cortex. Brain research *127*, 327-352.
Gilbert, P.F. (1976). Simple spike frequency and the number of secondary spikes
in the complex spike of the cerebellar Purkinje cell. Brain research *114*, 334-338.
Golding, N.L., Staff, N.P., and Spruston, N. (2002). Dendritic spikes as a
mechanism for cooperative long-term potentiation. Nature *418*, 326-331.
Hines, M.L., and Carnevale, N.T. (1997). The NEURON simulation environment.
Neural computation *9*, 1179-1209.
Ito, M. (1972). Neural design of the cerebellar motor control system. Brain
research *40*, 81-84.
Jelitai, M., Puggioni, P., Ishikawa, T., Rinaldi, A., and Duguid, I. (2016). Dendritic
excitation-inhibition balance shapes cerebellar output during motor behaviour.
Nature communications *7*, 13722.
Keren, N., Peled, N., and Korngreen, A. (2005). Constraining compartmental
models using multiple voltage recordings and genetic algorithms. Journal of
neurophysiology *94*, 3730-3742.
Khaliq, Z.M., Gouwens, N.W., and Raman, I.M. (2003). The contribution of
resurgent sodium current to high-frequency firing in Purkinje neurons: an
experimental and modeling study. The Journal of neuroscience : the official
journal of the Society for Neuroscience *23*, 4899-4912.
Khaliq, Z.M., and Raman, I.M. (2005). Axonal propagation of simple and complex
spikes in cerebellar Purkinje neurons. The Journal of neuroscience : the official
journal of the Society for Neuroscience *25*, 454-463.
Khavandgar, S., Walter, J.T., Sageser, K., and Khodakhah, K. (2005). Kv1 channels
selectively prevent dendritic hyperexcitability in rat Purkinje cells. The Journal
of physiology *569*, 545-557.
Kitamura, K., and Hausser, M. (2011). Dendritic calcium signaling triggered by
spontaneous and sensory-evoked climbing fiber input to cerebellar Purkinje cells
in vivo. The Journal of neuroscience : the official journal of the Society for
Neuroscience *31*, 10847-10858.
Larkum, M.E., Zhu, J.J., and Sakmann, B. (2001). Dendritic mechanisms
underlying the coupling of the dendritic with the axonal action potential
initiation zone of adult rat layer 5 pyramidal neurons. The Journal of physiology
533, 447-466.
Lev-Ram, V., Wong, S.T., Storm, D.R., and Tsien, R.Y. (2002). A new form of
cerebellar long-term potentiation is postsynaptic and depends on nitric oxide

678 but not cAMP. Proceedings of the National Academy of Sciences of the United
679 States of America *99*, 8389-8393.

680 Mano, N. (1970). Changes of simple and complex spike activity of cerebellar
681 purkinje cells with sleep and waking. Science *170*, 1325-1327.

682 Marr, D. (1969). A theory of cerebellar cortex. The Journal of physiology *202*,
683 437-470.

684 Martina, M., Metz, A.E., and Bean, B.P. (2007). Voltage-dependent potassium
685 currents during fast spikes of rat cerebellar Purkinje neurons: inhibition by BDS-
686 I toxin. Journal of neurophysiology *97*, 563-571.

687 Martina, M., Yao, G.L., and Bean, B.P. (2003). Properties and functional role of
688 voltage-dependent potassium channels in dendrites of rat cerebellar Purkinje
689 neurons. The Journal of neuroscience : the official journal of the Society for
690 Neuroscience *23*, 5698-5707.

691 Maruta, J., Hensbroek, R.A., and Simpson, J.I. (2007). Intraburst and interburst
692 signaling by climbing fibers. The Journal of neuroscience : the official journal of
693 the Society for Neuroscience *27*, 11263-11270.

694 Masoli, S., Solinas, S., and D'Angelo, E. (2015). Action potential processing in a
695 detailed Purkinje cell model reveals a critical role for axonal
696 compartmentalization. Front Cell Neurosci *9*, 47.

697 Mathy, A., Ho, S.S., Davie, J.T., Duguid, I.C., Clark, B.A., and Hausser, M. (2009).
698 Encoding of oscillations by axonal bursts in inferior olive neurons. Neuron *62*,
699 388-399.

700 Najafi, F., Giovannucci, A., Wang, S.S., and Medina, J.F. (2014). Coding of stimulus
701 strength via analog calcium signals in Purkinje cell dendrites of awake mice.
702 eLife *3*, e03663.

703 Ohtsuki, G., Piochon, C., Adelman, J.P., and Hansel, C. (2012). SK2 channel
704 modulation contributes to compartment-specific dendritic plasticity in
705 cerebellar Purkinje cells. Neuron *75*, 108-120.

706 Otsu, Y., Marcaggi, P., Feltz, A., Isope, P., Kollo, M., Nusser, Z., Mathieu, B., Kano,
707 M., Tsujita, M., Sakimura, K._, et al._ (2014). Activity-dependent gating of calcium
708 spikes by A-type K+ channels controls climbing fiber signaling in Purkinje cell
709 dendrites. Neuron *84*, 137-151.

710 Palmer, L.M., Clark, B.A., Grundemann, J., Roth, A., Stuart, G.J., and Hausser, M.
711 (2010). Initiation of simple and complex spikes in cerebellar Purkinje cells. The
712 Journal of physiology *588*, 1709-1717.

713 Piochon, C., Titley, H.K., Simmons, D.H., Grasselli, G., Elgersma, Y., and Hansel, C.
714 (2016). Calcium threshold shift enables frequency-independent control of
715 plasticity by an instructive signal. Proceedings of the National Academy of
716 Sciences of the United States of America *113*, 13221-13226.

717 Roth, A., and Hausser, M. (2001). Compartmental models of rat cerebellar
718 Purkinje cells based on simultaneous somatic and dendritic patch-clamp
719 recordings. The Journal of physiology *535*, 445-472.

720 Schmolesky, M.T., Weber, J.T., De Zeeuw, C.I., and Hansel, C. (2002). The making
721 of a complex spike: ionic composition and plasticity. Annals of the New York
722 Academy of Sciences *978*, 359-390.

723 Steuber, V., Mittmann, W., Hoebeek, F.E., Silver, R.A., De Zeeuw, C.I., Hausser, M.,
724 and De Schutter, E. (2007). Cerebellar LTD and pattern recognition by Purkinje
725 cells. Neuron *54*, 121-136.

Swensen, A.M., and Bean, B.P. (2003). Ionic mechanisms of burst firing in
dissociated Purkinje neurons. The Journal of neuroscience : the official journal of
the Society for Neuroscience *23*, 9650-9663.
Warnaar, P., Couto, J., Negrello, M., Junker, M., Smilgin, A., Ignashchenkova, A.,
Giugliano, M., Thier, P., and De Schutter, E. (2015). Duration of Purkinje cell
complex spikes increases with their firing frequency. Front Cell Neurosci *9*.
Wilms, C.D., and Hausser, M. (2015). Reading out a spatiotemporal population
code by imaging neighbouring parallel fibre axons in vivo. Nature
communications *6*, 6464.
Yang, Y., and Lisberger, S.G. (2014). Purkinje-cell plasticity and cerebellar motor
learning are graded by complex-spike duration. Nature *510*, 529-532.
Zagha, E., Manita, S., Ross, W.N., and Rudy, B. (2010). Dendritic Kv3.3 potassium
channels in cerebellar purkinje cells regulate generation and spatial dynamics of
dendritic Ca2+ spikes. Journal of neurophysiology *103*, 3516-3525.
740
741

Supplementary Information

Supplementary Figures

Figure S1. PC responses to CF bursts.
We reproduced the experiment by Mathy et al. (2009). Example somatic and dendritic responses to CF burst with 1 (up), 3 (middle) and 6 spikes (bottom) in the CF burst. The simulated instantaneous CF bursting frequency is 250 Hz close to the experimentally reported ~ 273 Hz.

751
Figure S2. Voltage states regulate CF-evoked dendritic spike generation and propagation.
A-C. Color-coded representation of dendritic voltage responses at different holding potentials (-84 mV,
-80 mV and -60 mV respectively). **D-F.** Example somatic CSs (black), dendritic voltage responses on
the smooth dendrite (red) and spiny dendrite (blue) at different voltage states. **G-I.** Distance-dependent
propagation of dendritic responses at different voltage states. **J-L.** Ionic currents in the distal spiny
dendritic site at different voltage states. **M-O.** Ionic currents in the proximal smooth dendritic site at
different voltage states. The surface areas of the segments chosen at the smooth dendrite and spiny
dendrite are 64 μm^2 and 23 μm^2 respectively. **P-R.** Color-coded peak Ca^{2+} concentrations at
corresponding membrane potentials.

761
762
Figure S3. CF-evoked dendritic voltage response waveforms with different current injections
(holding potentials).
A. CF-evoked dendritic spike with two spikelets in the experiment (left, reproduced from Fig.5A of
Davie et al. (2008)) and in the model (right). **B.** CF-evoked dendritic voltage response in the
experiment (left, data from Ohtsuki et al. (2012)) and in the model (right). The 'deflection' caused by a
somatic Na$^+$ spikelet can be found on top of the dendritic response under this condition in both
experimental recording and in the model (circled in the figure).
770

771
Figure S4. Global inactivation of the Kv4 current can facilitate the occurrence of dendritic spikes
and make the dendritic excitability more homogenous.
The Kv4 inactivation was simulated by left shifting its availability by 11 mV. Dendritic responses to
holding currents of -0.89 nA, -0.71 nA and -0.53 nA are shown from left to right. Color-coded
representation of dendritic response peaks, distance-dependent propagation of dendritic response peaks,
and dendritic responses (EPSP or spike) at different branches are shown from top to bottom.
778

779
Figure S5. CF responses when the voltage states of the PC are manipulated by combined
excitatory and inhibitory synaptic inputs.
A-C show color-coded representation of dendritic voltage response peaks with the increase of SSFRs.
D-F show the corresponding example somatic and dendritic responses. CF responses at a higher time
scale are shown in the bottom panel. **G** shows the effect of somatic voltage states on the CS spikelet
numbers and durations. **H.** Effect of somatic voltage states on the amplitudes of the first 3 somatic
spikelets. **I-K** show color-coded CF-evoked peak Ca^{2+} concentrations with the corresponding SSFR.
787

788
Figure S6. Local down-regulation of Kv4 current mdulates dendritic responses in a spatially
precise manner.
In the child branchlet of the labeled dendritic section, the availability of Kv4 current is left shifted by
11 mV (bottom panel). Only the dendritic responses located within the labeled branchlet (For example,
Point F) are facilitated by the local Kv4 down-regulation. In other parts of the dendrite, dendritic
responses are not affected.

Figure S7. The dendritic spike measured on the proximal smooth dendrite can't reflect dendritic responses in the rest of the dendrite.
Here we reproduce the experiment by Davie et al. (2008). The holding current is -0.2 nA. The CF is activated at different phases of the somatic interspike interval to achieve a slight change of dendritic voltage state. In case I (blue traces), a secondary dendritic spikelet occurs at point B of the distal dendrite. However, at point A and P of the same branch closer to the soma, no secondary dendritic spikelets are triggered. With a slight depolarization of the dendrite in case II (black traces), secondary dendritic spikelets occur at both proximal sites (point A and point P) and at the distal site (point B).

Supplementary Experimental Procedures

Current equations and calcium handling

1) Na$^+$ current in the PC.

Na$^+$ current in the model is distributed in the AIS and soma. Na$^+$ current in the PC is
composed of 'transient', 'persistent' and 'resurgent' components. We started from the
previous model (Khaliq et al., 2003) and made slight changes (Narsg). In the soma,
we adjusted the reaction rates between the I1 - I5 states by right shifting 5 mV (Yan et
al., 2014): $\alpha = 150\ e^{(V-5)/20}$, $\beta = 3\ e^{(V-5)/-20}$. Other parameters are the same as
the original model. This adjustment mainly right-shifts the inactivation curve of this
current by $\sim$ 2.5 mV. The inactivation curve of the modified model lies within
measurements (Raman and Bean, 2001; Yan et al., 2014). In the AIS, beside the
above-mentioned adjustment, we also adjusted deactivation rates between C1 - C5 to
lower the spike threshold in the AIS, as done by Colbert and Pan (2002). Therefore,
$\beta = 3\ e^{(V+15)/-15}$ for reaction rates between C1 - C5. First, by this adjustment, the
contribution of AIS spikes to somatic spikes is closer to measured data (represented
by d^2V/dt^2 in Fig. 1D). Second, we removed the frequently failed recovery of simple
spikes after a complex spike when there is no obvious after-hyperpolarization
following the complex spike (although this can be observed in some experiments, data
shared by Paul Mathews from UCLA).

It is impossible to approximate Na$^+$ currents using a single model, given their multiple
components. In the original model (Khaliq et al., 2003), the persistent component is
tiny. Therefore, we incorporated an additional persistent Na$^+$ current model to
compensate (NaP). The steady state activation curve is from Carter et al. (2012),
$m_{ss} = 1/(1 + e^{-(V+66)/5})$. The time constant is from Baker (2005), $\tau_m = 1/(\alpha_m +$
$\beta_m)$, $\alpha_m = 1/(\ 17.235/(1+e^{(V+27.58)/-11.47}\ ))$, $\beta_m\ = 1/(\ 17.235/(1+e^{(V+86.2)/19.8}\ ))$.
This persistent component helps to maintain spike firing during negative current
injections.

2) H-current in the PC
H-current (Ih) is distributed in the whole PC (Angelo et al., 2007). In our model, the
formulations are the same as the model formulations from Angelo et al. (2007),
except we shifted the activation curve to a more hyperpolarized direction by 6 mV to
make 'sag' responses with different holding currents more comparable to
experimental recordings (Fig. 1G).

3) T-type Ca^{2+} current in the PC
T-type Ca^{2+} current is distributed in the AIS (Bender and Trussell, 2009), the smooth
dendrite and the spiny dendrite (Hildebrand et al., 2009; Otsu et al., 2014). The
formulation of this current is according to data from Otsu et al. (2014). $m_{ss} = 1/(1 +$
$e^{-(V-35.606)/4.7})$, $\tau_m = 1/(1.2757 + \alpha_m + \beta_m)$,
$\alpha_m = -2.3199/(1 + e^{(V+41.448)/-30.655})$, $\beta_m = 2.5712/(1 + e^{(V+21.7866)/-9.6306})$,
$h_{ss} = 1/(1 + e^{(V+81.718)/6.4635})$, $\tau_h = 1/(0.0076 + \alpha_h + \beta_h)$, $\alpha_h = 0.17746/(1 +$
$e^{(V+41.448)/-30.655})$, $\beta_h = 0.13402/(1 + e^{(V+94.836)/-5.5845})$. In the model, both BK
and SK2 don't sense Ca^{2+} influx from T-type Ca^{2+} channels.

4) P-type Ca^{2+} current in the PC

P-type Ca^{2+} current is distributed throughout the entire PC (Indriati et al., 2013). The
formulation of this current is according to the data fromBenton and Raman (2009).

$m_{ss} = 1/(1 + e^{-(V-25.5)/4.113})$, $\tau_m = 0.2 + 0.7031 e^{-((V+25)/14)^2}$.

5) Ca^{2+} handling mechanism
In the dendrite, we used the formulation from Anwar et al. (2012) with 4 shells. In
both soma and AIS, we made BK channels sense Ca^{2+} signals closer to the membrane
surface (subspace) or Ca^{2+} channels by increasing the number of shells (Indriati et al.,
2013). In the AIS, we still used the formulation as in Anwar et al. (2012), but we
increased the number of shells to 10. In the soma, we used the fixed depth scheme for
the outermost shell and 20 shells in total to reduce the computational load (Anwar et
al., 2014). We also incorporated a sarcoplasmic endoplasmic reticulum Ca^{2+}ATPase
(SERCA) pump to reduce the spike-over-spike Ca^{2+} summation in AIS and soma.
$J_{SERCA} = 0.1 * vol(i) * Ca[i]/(Ca[i] + 0.0058(mM))$, $vol(i)$ means the volume of
the i_{th} shell. $Ca[i]$ means the free Ca^{2+} concentration of the i_{th} shell.

6) BK current in the PC
We used the same formulation from Anwar et al. (2012) as the fast (or IBTX-sensitive)
component of the BK current (Benton et al., 2013). We also incorporated the slow (or
IBTX-insensitive) component of BK current (Benton et al., 2013) from Jaffe et al.
(2011). The formulation of their type II model was used. The fast component is
distributed in the whole PC and the slow component is only distributed in the soma
and the AIS.

7) SK2 current in the PC
SK2 current is distributed in the whole PC (Belmeguenai et al., 2010; Swensen and
Bean, 2003). We used the formulation from Solinas et al. (2007) according to the data
by Hirschberg et al. (1998).

8) Kv3 current in the PC
KV3 current is distributed in the whole PC (Martina et al., 2003). The formulation is
from Akemann et al. (2009) according to the data from Martina et al. (2007). In our
model, the steady state activation curve is left shifted by 4 mV with $V_{0.5} \approx 2$ mV,
which is still within the experimental range (Martina et al., 2007; Martina et al., 2003;
Zagha et al., 2010).

9) Kv4 current in the PC
In our model, Kv4 current has both fast and slow inactivation components and is
distributed in the smooth and the spiny dendrites (Otsu et al., 2014). We took the
steady state activation curve from Gunay et al. (2008). The inactivation curve and
time constants are according to experimental data of Stéphane Dieudonné. $m_{ss} =$
$1/(1 + e^{(V+49)/-12.5})$, $\tau_m = 1/(\alpha_m + \beta_m)$, $\alpha_m = 0.1342/(1 + e^{(V+60)/32.19976})$,
$\beta_m = 0.15743/(1 + e^{(V+57)/37.51346})$, $h_{ss} = 1/(1 + e^{(V+75.30348)/6.06329})$,
$\tau_{hf} = 1/(\alpha_{hf} + \beta_{hf})$, $\alpha_{hf} = 0.01342/(1 + e^{(V+60)/7.86476})$, $\beta_{hf} = 0.04477/(1 +$
$e^{(V+54)/-11.6315})$, $\tau_{hs} = 100$.

10) Kv1 current in the PC
In our model, Kv1 is distributed in the dendrite (Khavandgar et al., 2005). The
formulation is from Akemann and Knopfel (2006). We added an inactivation gate,

according to data from Otsu et al. (2014). $h_{ss} = 1/(1 + e^{(V+66.16)/6.1881})$. Due to the
slow inactivation process, the inactivation time constant was set to be 1000 ms.
906

Parameter tuning
The model was hand-tuned because extensive preceding efforts using automatic
parameter searching did not achieve good results (Achard and De Schutter, 2006; Van
Geit et al., 2007). Due to technical limitations, we can only get simultaneous somatic
and dendritic patch-clamp recordings (single dendritic site) to constrain the model in
most cases. However, it has been demonstrated that data recorded from just two
locations in the same L5 pyramidal neuron are not sufficient to accurately constrain a
compartmental model (Keren et al., 2005). For the PC, we were able to get good F-I
curves and 'sag' responses with automated search methods, but the dendrite model
could not generate physiologic responses. This is because we don't have sufficient
data to constrain the active properties of the dendrite and the interaction between
soma and dendrite by automatic parameter searching. Therefore, we decided to use
hand-tuning instead, and we tried to incorporate as many experimental observations
as possible into the model. After repeated modification over a period of more than
two years, we achieved the present model.
922

1) Currents in the soma
The Na$^+$ current profile nearly coincides with the total net ionic current during the
depolarization phase of dissociated PC AP-clamp recordings (Carter and Bean, 2009).
The total net ionic current is calculated by –C*dV/dt, where C is the cell capacitance
and dV/dt is the time derivative of the voltage. The results of Carter and Bean suggest
that Na$^+$ current is nearly the sole current supporting the depolarization and K$^+$
currents are not activated during the depolarization phase (as illustrated in Fig. 2 of
their paper). This also agrees with the high activation threshold of K$^+$ currents in the
PC (Martina et al., 2007). This finding provided an opportunity to firmly constrain the
somatic ionic currents. The measured peak Na$^+$ current amplitude is 12.5 nA in
dissociated mouse PCs with a capacitance of 12 pF, but the ratio of
current/capacitance in the soma may be underestimated due to the possible remaining
stump of the dendritic tree. In our model, we have a peak somatic Na$^+$ current of 17
nA relative to a capacitance of 10.5 pF in the soma. Then we fixed the conductance
density of somatic Kv3 current by constraining the spike half-amplitude duration
(0.15 ms) close to experimental measurements (0.17 ms, we also gave consideration
to the spike duration in the intact PC as shown in Fig. 1B). The high activation
threshold of the Kv3 current makes it active mainly during the repolarization phase
(Carter and Bean, 2009) and constrains the spike duration (Martina et al., 2007).
Subsequently, we constrained the conductance density of P-type Ca^{2+} current and BK
current to have an AHP (-77 mV) comparable to the recorded (-82 mV, still we also
considered the AHP in the intact PC ~ -70 mV). The conductance density of P-type
Ca^{2+} is constrained to be the same as that in the smooth dendrite. Then only SK2 and
Ih remained unconstrained for the somatic ionic currents. We initially assigned
tentative conductance density values for these two currents to develop a firing somata
model (like the previous Kaliq-Raman model (Khaliq et al., 2003)). To reflect the
negligible role of Ih in dissociated PC firing (Raman and Bean, 1999), the
conductance density of Ih is half of that in smooth dendrite. In the intact PC, the
conductance density of somatic SK2 current is adjusted to get a F-I curve close to the
experimental measurements (Fig. 1G). The persistent component of Na$^+$ current is

constrained so that the intact PC still fires with -0.17 nA somatic holding current, but
it stops firing when the current amplitude changes to -0.36 nA.
955
2) Currents in the AIS
We constrained ionic currents in the AIS to make the model behave like experimental
data in two aspects: the axosomatic delay of the simple spike (Fig. 1B) and the second
derivative of the simple spike (Fig. 1D). The first reflects how the spike in the AIS
precedes the somatic spike and the second reflects the contribution of the spike in the
AIS to the somatic spike. We assumed the conductance densities of Ih in AIS are the
same as in soma.
963
3) Currents in the dendrite
Due to the limited quantitative data, currents in the dendrite are more difficult to
constrain compared with somatic currents. We first constrained the P-type Ca^{2+}
current in the dendrite. In the spiny dendrite, Ca^{2+} influx during one dendritic spikelet
is 1.3e-14 $C/\mu m^3$ in our model, which is close to the experimentally estimated 1.1e-14
$C/\mu m^3$ (Otsu et al., 2014). The CF-evoked Ca^{2+} influx increases with distance from
soma when a Ca^{2+}-supported dendritic spike occurs (Kitamura and Hausser, 2011;
Otsu et al., 2014). Therefore, in the smooth dendrite, the current density was
constrained to have a calcium influx ratio (smooth dendrite relative to the spiny
dendrite at same membrane surface area) of 0.36. The conductance densities of P-type
Ca^{2+} current are further supported by the time to peak of the first dendritic spikelet
being ~ 1.5 ms in the smooth dendrite (Davie et al., 2008; Kitamura and Hausser,
2011). Conductance densities of Kv4 current are assumed to be homogeneously
distributed and then repeatedly adjusted to make this current able to 'brake' the
dendritic spikes at low voltage ranges. We distribute more T-type Ca^{2+} channels
(Hildebrand et al., 2009) in the spiny dendrite given they are preferentially distributed
on spines. This current helps to offset the window current of Kv4 current to maintain
the dendritic interspike interval membrane potential. We distribute larger conductance
densities of BK current on the smooth dendrite compared to on spiny dendrite, given
the BK channel clusters are mainly distributed on the soma and smooth dendrite
(Kaufmann et al., 2010). BK current contributes to reasonable AHP after complex
spikes as found in Davie et al. (2008). Ih is constrained to get a 'sag' response close to
the experimental recordings (Fig. 1G). Kv1 and SK2 help to regulate dendritic
excitability to stop the occurrence of Na^+-Ca^{2+} bursting at low current injections. In
our model, as with most cell recordings in rat PCs, Na^+-Ca^{2+} bursts begin to occur
with 1.25 nA somatic current injection. In addition, this dendrite model can replicate
the distance-dependent decay of the simple spike amplitude (Fig. 1E), the regulatory
role of Kv4 current in voltage-dependent dendritic spike generation and propagation
(Fig. 3,S2,S5), dendritic voltage response waveforms at different conditions (Fig. S3),
and peak dendritic voltage response propagation with distance from the soma at
specific condition (Fig. 3G).
995
996
997 **Current conductance densities in different part of PC**
998

Currents (nS/cm²)	AIS	soma	smooth dendrite	spiny dendrite
Narsg	0.56	0.0317	0	0
NaP	0.0023	1.4e-4	0	0

Ih	1.08e-4	1.08e-4	2.16e-4	3.24e-4
T- type Ca^{2+}	1.28e-4	0	2.7e-5	1.08e-4
P-type Ca^{2+}	1.05	1.9e-4	1.9e-4	7.6e-4
BKfast	6	0.8736	0.21	0.0448
BKslow	1.05	0.3	0	0
SK2	0.0278	0.0075	2.16e-4	3.6e-4
Kv3	115	1.8	0.1512	0.2268
Kv4	0	0	0.015	0.015
Kv1	0	0	0.002	0.001

999
1000
1001

Supplementary References

Achard, P., and De Schutter, E. (2006). Complex parameter landscape for a
complex neuron model. PLoS computational biology *2*, e94.
Akemann, W., and Knopfel, T. (2006). Interaction of Kv3 potassium channels and
resurgent sodium current influences the rate of spontaneous firing of Purkinje
neurons. The Journal of neuroscience : the official journal of the Society for
Neuroscience *26*, 4602-4612.
Akemann, W., Lundby, A., Mutoh, H., and Knopfel, T. (2009). Effect of voltage
sensitive fluorescent proteins on neuronal excitability. Biophysical journal *96*,
3959-3976.
Angelo, K., London, M., Christensen, S.R., and Hausser, M. (2007). Local and global
effects of I(h) distribution in dendrites of mammalian neurons. The Journal of
neuroscience : the official journal of the Society for Neuroscience *27*, 8643-8653.
Anwar, H., Hong, S., and De Schutter, E. (2012). Controlling Ca2+-activated K+
channels with models of Ca2+ buffering in Purkinje cells. Cerebellum *11*, 681-
693.
Anwar, H., Roome, C.J., Nedelescu, H., Chen, W., Kuhn, B., and De Schutter, E.
(2014). Dendritic diameters affect the spatial variability of intracellular calcium
dynamics in computer models. Front Cell Neurosci *8*, 168.
Baker, M.D. (2005). Protein kinase C mediates up-regulation of tetrodotoxin-
resistant, persistent Na+ current in rat and mouse sensory neurones. The Journal
of physiology *567*, 851-867.
Belmeguenai, A., Hosy, E., Bengtsson, F., Pedroarena, C.M., Piochon, C., Teuling, E.,
He, Q., Ohtsuki, G., De Jeu, M.T., Elgersma, Y*., et al.* (2010). Intrinsic plasticity
complements long-term potentiation in parallel fiber input gain control in
cerebellar Purkinje cells. The Journal of neuroscience : the official journal of the
Society for Neuroscience *30*, 13630-13643.
Bender, K.J., and Trussell, L.O. (2009). Axon initial segment Ca2+ channels
influence action potential generation and timing. Neuron *61*, 259-271.
Benton, M.D., Lewis, A.H., Bant, J.S., and Raman, I.M. (2013). Iberiotoxin-sensitive
and -insensitive BK currents in Purkinje neuron somata. Journal of
neurophysiology *109*, 2528-2541.
Benton, M.D., and Raman, I.M. (2009). Stabilization of Ca current in Purkinje
neurons during high-frequency firing by a balance of Ca-dependent facilitation
and inactivation. Channels *3*, 393-401.

Carter, B.C., and Bean, B.P. (2009). Sodium entry during action potentials of mammalian neurons: incomplete inactivation and reduced metabolic efficiency in fast-spiking neurons. Neuron *64*, 898-909.

Carter, B.C., Giessel, A.J., Sabatini, B.L., and Bean, B.P. (2012). Transient Sodium Current at Subthreshold Voltages: Activation by EPSP Waveforms. Neuron *75*, 1081-1093.

Colbert, C.M., and Pan, E. (2002). Ion channel properties underlying axonal action potential initiation in pyramidal neurons. Nature neuroscience *5*, 533-538.

Davie, J.T., Clark, B.A., and Hausser, M. (2008). The origin of the complex spike in cerebellar Purkinje cells. The Journal of neuroscience : the official journal of the Society for Neuroscience *28*, 7599-7609.

Gunay, C., Edgerton, J.R., and Jaeger, D. (2008). Channel density distributions explain spiking variability in the globus pallidus: a combined physiology and computer simulation database approach. The Journal of neuroscience : the official journal of the Society for Neuroscience *28*, 7476-7491.

Hildebrand, M.E., Isope, P., Miyazaki, T., Nakaya, T., Garcia, E., Feltz, A., Schneider, T., Hescheler, J., Kano, M., Sakimura, K.*, et al.* (2009). Functional Coupling between mGluR1 and Ca(v)3.1 T-Type Calcium Channels Contributes to Parallel Fiber-Induced Fast Calcium Signaling within Purkinje Cell Dendritic Spines. Journal of Neuroscience *29*, 9668-9682.

Hirschberg, B., Maylie, J., Adelman, J.P., and Marrion, N.V. (1998). Gating of recombinant small-conductance Ca-activated K+ channels by calcium. The Journal of general physiology *111*, 565-581.

Indriati, D.W., Kamasawa, N., Matsui, K., Meredith, A.L., Watanabe, M., and Shigemoto, R. (2013). Quantitative localization of Cav2.1 (P/Q-type) voltage-dependent calcium channels in Purkinje cells: somatodendritic gradient and distinct somatic coclustering with calcium-activated potassium channels. The Journal of neuroscience : the official journal of the Society for Neuroscience *33*, 3668-3678.

Jaffe, D.B., Wang, B., and Brenner, R. (2011). Shaping of action potentials by type I and type II large-conductance Ca(2)+-activated K+ channels. Neuroscience *192*, 205-218.

Kaufmann, W.A., Kasugai, Y., Ferraguti, F., and Storm, J.F. (2010). Two distinct pools of large-conductance calcium-activated potassium channels in the somatic plasma membrane of central principal neurons. Neuroscience *169*, 974-986.

Keren, N., Peled, N., and Korngreen, A. (2005). Constraining compartmental models using multiple voltage recordings and genetic algorithms. Journal of neurophysiology *94*, 3730-3742.

Khaliq, Z.M., Gouwens, N.W., and Raman, I.M. (2003). The contribution of resurgent sodium current to high-frequency firing in Purkinje neurons: an experimental and modeling study. The Journal of neuroscience : the official journal of the Society for Neuroscience *23*, 4899-4912.

Khavandgar, S., Walter, J.T., Sageser, K., and Khodakhah, K. (2005). Kv1 channels selectively prevent dendritic hyperexcitability in rat Purkinje cells. The Journal of physiology *569*, 545-557.

Kitamura, K., and Hausser, M. (2011). Dendritic calcium signaling triggered by spontaneous and sensory-evoked climbing fiber input to cerebellar Purkinje cells in vivo. The Journal of neuroscience : the official journal of the Society for Neuroscience *31*, 10847-10858.

1087   Martina, M., Metz, A.E., and Bean, B.P. (2007). Voltage-dependent potassium currents during fast spikes of rat cerebellar Purkinje neurons: inhibition by BDS-I toxin. Journal of neurophysiology *97*, 563-571.

Martina, M., Yao, G.L., and Bean, B.P. (2003). Properties and functional role of voltage-dependent potassium channels in dendrites of rat cerebellar Purkinje neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *23*, 5698-5707.

Otsu, Y., Marcaggi, P., Feltz, A., Isope, P., Kollo, M., Nusser, Z., Mathieu, B., Kano, M., Tsujita, M., Sakimura, K*., et al.* (2014). Activity-dependent gating of calcium spikes by A-type K+ channels controls climbing fiber signaling in Purkinje cell dendrites. Neuron *84*, 137-151.

Raman, I.M., and Bean, B.P. (1999). Ionic currents underlying spontaneous action potentials in isolated cerebellar Purkinje neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *19*, 1663-1674.

Raman, I.M., and Bean, B.P. (2001). Inactivation and recovery of sodium currents in cerebellar Purkinje neurons: evidence for two mechanisms. Biophysical journal *80*, 729-737.

Solinas, S., Forti, L., Cesana, E., Mapelli, J., De Schutter, E., and D'Angelo, E. (2007). Computational reconstruction of pacemaking and intrinsic electroresponsiveness in cerebellar Golgi cells. Front Cell Neurosci *1*, 2.

Swensen, A.M., and Bean, B.P. (2003). Ionic mechanisms of burst firing in dissociated Purkinje neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *23*, 9650-9663.

Van Geit, W., Achard, P., and De Schutter, E. (2007). Neurofitter: a parameter tuning package for a wide range of electrophysiological neuron models. Frontiers in neuroinformatics *1*, 1.

Yan, H., Pablo, J.L., Wang, C., and Pitt, G.S. (2014). FGF14 modulates resurgent sodium current in mouse cerebellar Purkinje neurons. eLife *3*, e04193.

Zagha, E., Manita, S., Ross, W.N., and Rudy, B. (2010). Dendritic Kv3.3 potassium channels in cerebellar purkinje cells regulate generation and spatial dynamics of dendritic Ca2+ spikes. Journal of neurophysiology *103*, 3516-3525.