6mer seed toxicity in tumor suppressive microRNAs

Quan Q. Gao[1,6], William E. Putzbach[1,6], Andrea E. Murmann[1], Siquan Chen[2], Aishe A. Sarshad[3], Johannes M. Peter[4], Elizabeth T. Bartom[5], Markus Hafner[3], and Marcus E. Peter[1,5,*]

[1]Department of Medicine/Division Hematology/Oncology and [5]Department of Biochemistry and Molecular Genetics, Northwestern University, Chicago, IL 60611, USA, [2]Cellular Screening Center, Institute for Genomics & Systems Biology, The University of Chicago, Chicago, IL 60637, USA, [3]Laboratory of Muscle Stem Cells and Gene Regulation, NIAMS, NIH, Bethesda, MD 20892, USA, [4]DigiPen Institute of Technology, Redmond, WA 98052, USA,

[6] Authors share first authorship

Corresponding author: Marcus E. Peter, E-mail: m-peter@northwestern.edu, phone: 312-503-1291; FAX: 312-503-0189.

Keywords: RNAi, miRNAs, strand selection, toxicity, evolution

SUMMARY

Many small interfering (si)RNAs are toxic to cancer cells through a 6mer seed sequence (position 2-7 of the

guide strand). A siRNA screen with all 4096 possible 6mer seed sequences in a neutral siRNA backbone

revealed a preference for guanine in positions 1 and 2 and a high overall G or C content in the seed of the

most toxic siRNAs for four tested human and mouse cell lines. Toxicity of these siRNAs stems from

targeting survival genes with C-rich 3'UTRs. The master tumor suppressor miRNA miR-34a-5p is toxic

through such a G-rich 6mer seed and is upregulated in cells subjected to genotoxic stress. In the absence of

most canonical miRNAs, a modified highly toxic form of miR-320a-3p is induced. An analysis of all mature

miRNAs suggests that during evolution most miRNAs evolved to avoid guanine at the 5' end of the 6mer

seed sequence of the guide strand. In contrast, for certain tumor suppressive miRNAs the guide strand

contains a G-rich toxic 6mer seed, presumably to eliminate cancer cells.

INTRODUCTION

RNA interference (RNAi) is a form of post-transcriptional regulation exerted by 19-21 nt long double

stranded RNAs that negatively regulate gene expression at the mRNA level. RNAi-active guide RNAs can

come from endogenous siRNAs and micro(mi)RNAs. For a miRNA, the RNAi pathway begins in the

nucleus with transcription of a primary miRNA precursor (pri-miRNA) [1]. Pri-miRNAs are first processed by

the Drosha/DGCR8 microprocessor complex into pre-microRNAs [2], which are then exported from the

nucleus to the cytoplasm by Exportin 5 [3]. Once in the cytoplasm, Dicer processes them further [4,5] and these

mature dsRNA duplexes are then loaded into Argonaute (Ago) proteins to form the RNA-induced silencing

complex (RISC) [6]. The sense/passenger strand is ejected/degraded, while the guide strand remains associated

with the RISC [7]. Depending on the degree of complementarity between the guide strand and its target, the

outcome of RNAi can either be target degradation - most often achieved by siRNAs with full

complementarity to its target mRNA [8] - or miRNA-like cleavage-independent silencing, mediated by

deadenylation/degradation or translational repression [9]. The latter mechanism can be initiated with as little as

six nucleotide base-pairing between a guide RNA's so-called seed sequence (positions 2 to 7) and fully

complementary seed matches in the target RNA [10,11]. This seed-based targeting most often occurs in the

3'UTR of a target mRNA [12,13].

A number of miRNAs function either as tumor suppressors or as oncogenes [14]. Their cancer specific

activities are usually explained by their identified targets, being oncogenes tumor or suppressors,

respectively [14]. Examples of targets of tumor-suppressive miRNAs are the oncogenes Bcl-2 for miR-15/16 [15]

and c-Myc for miR-34a [16]. While many miRNAs have been reported to have both tumor suppressive and

oncogenic activities depending on the cancer context, examples for widely established tumor promoting

miRNAs are miR-221/222, miR-21, miR-155, and members of the miR-17~92 cluster, or its paralogues

miR-106b~25 and miR-106a~363 [17, 18]. In contrast, two of the major tumor suppressive miRNA families are

miR-15/16 and the p53 regulated miR-34a/c and miR-34b [19].

We recently discovered that many si- and shRNAs can kill all tested cancer cell lines through RNAi by

targeting the 3'UTRs of critical survival genes [20]. Cancer cells have difficulty developing resistance to this

mechanism both *in vitro* and when treated *in vivo* [21]. We reported that a 6mer seed sequence in the toxic

siRNAs is sufficient for effective killing [20]. We have now performed a strand specific siRNA screen with a

library of individual siRNAs representing all 4096 possible 6mer seed sequences in a neutral RNA duplex.

This screen, while based on siRNA biochemistry was not designed to identify targets that are degraded

through siRNA mediated slicing activity but to identify toxicity caused by moderately targeting hundreds of

genes required for cell survival in a mechanism similar to miRNA-induced silencing.

We report that the most toxic 6mer seeds are G-rich with a G enrichment towards the 5' end targeting

survival genes with a high C content in their 3'UTR in a miRNA-like manner. Many tumor suppressive

miRNAs such as miR-34a-5p but none of the established oncogenic miRNAs contain G-rich 6mer seeds and

most of miR-34a-5p's toxicity comes from its 6mer seed sequence. Mature miRNAs from older and more

conserved miRNAs contain less toxic seeds. We demonstrate that for most miRNAs the more abundant

mature form corresponds to the arm that contains the less toxic seed. In contrast, for major tumor suppressive

miRNAs, the mature miRNA is derived from the arm that harbors the more toxic seed. Our data allow us to

conclude that while most miRNAs have evolved to avoid targeting survival and housekeeping genes, certain

tumor suppressive miRNAs function to kill cancer cells through a toxic G-rich 6mer seed targeting the

3'UTR of survival genes.

RESULTS

Identifying the most toxic 6mer seeds

To test how certain 6mer seeds present in the guide strand of a siRNA affect cancer cell survival, we recently

designed a neutral 19mer oligonucleotide scaffold with two nucleotide 3' overhangs, and we demonstrated

that modifying a siRNA strand at positions 1 and 2 by 2'-O-methylation (OMe) completely blocks its loading

into the RISC [22]. Different 6mer sequences can be inserted at positions 2-7 of the guide strand with the

designated passenger strand modified by OMe (two red Xs in **Fig. 1a**). Transfection efficiency and

conditions were optimized for each cell line used. To determine the general rules of seed-based toxicity, we

individually transfected 4096 siRNAs with all possible 6mer seed sequences in this 19mer scaffold into two

human, HeyA8 (ovarian cancer) and H460 (lung cancer), and two mouse cell lines M565 (liver cancer) and

3LL (lung cancer). This allowed us to rank all 4096 6mer seeds according to their toxicity (**Fig. 1b**,

Supplementary Table 1, and **6merdb.org**). The congruence between the results of the two human cell lines

and the two human and two mouse cell lines was quite high (r=0.68 and 0.65, respectively, **Fig. 1c**)

suggesting that many siRNAs were toxic through a mechanism independent of cancer origin and species.

Toxicity was caused by the different 6mer seeds in the *guide* strand. A siRNA duplex highly toxic to all cell

lines (#2733, HeyA8 cell viability = 1.4%) strongly inhibited cell growth and reduced cell viability of

HeyA8 cells only when the passenger strand was modified by the OMe modification. Toxicity was

completely blocked when the guide strand was modified (**Fig. S1a**). The toxicity was due to RNAi as

knockdown of AGO2 abolished the toxicity of two of the most toxic siRNAs (**Fig. S1b**).

We previously reported that the CD95 ligand (CD95L) coding region (CDS) is enriched in sequences that

when converted into si- or shRNAs are toxic to cancer cells [20] and that the CD95L mRNA itself is toxic to

cells [23]. We now report a substantial correlation between the most toxic CD95L-derived shRNAs and the

toxicity of their predicted 6mer seed (**Fig. 1d**) suggesting the CD95L-derived si/shRNAs kill cancer cells

through 6mer seed toxicity. Consistent with this assumption we found that the 6mer seeds of 4 previously

tested siRNAs derived from CD95L [24] in this screen were about as toxic as the full length siRNAs, with

siL3^{Seed} being the most toxic followed by siL2^{Seed} and less or no toxicity associated with siL4^{Seed} and siL1^{Seed}

(**Fig. 1b, Fig. 1c**). Our recent analysis suggested that the toxic si/shRNAs act like artificial miRNAs by

targeting the 3'UTR of mRNAs [20].

6mer seeds enriched in G at the 5'end are most toxic to human and mouse cell lines

We noticed that the 6mer seeds in siL3 and siL2 have a higher G content than the ones in siL4 and siL1 (**Fig.**

1b). By analyzing the screen results of all four cell lines (**Fig. S2**), we found that a high G content of the seed

correlated better with toxicity than a high C content. Almost no toxicity was found with seeds with a high A

content. To test the effect of nucleotide content on toxicity directly, we retested the 19 seed duplexes with

the highest content (>80%) for each of the four nucleotides in the four cell lines (**Fig. 2**). The reanalysis also

allowed us to determine the reproducibility of the results obtained in the large screens (which for technical

reasons had to be performed in three sets). All data on the three cell lines were highly reproducible especially

for the most toxic seeds (**Fig. S3a**). When the data on the four cell lines were compared, it became apparent

that in all cell lines, the G-rich seeds were by far the most toxic followed by the C-rich, U-rich, and A-rich

seeds (**Fig. 2a**). This indicates it is mostly the G content that determines toxicity.

Most genome-wide siRNA libraries designed to study functions of individual genes are highly

underrepresented in G and C to increase RNAi specificity [25] (**Fig. S3b**, left panel). In contrast, our complete

set of 6mer seed duplexes exhibits no nucleotide composition bias, allowing us to test the contributions of all

four nucleotides in each of the 6 seed positions (**Fig. S3b**, right panel). To determine the nucleotide content

of the most toxic seeds, we averaged the nucleotide content at each of the 6 positions of either the 200 most

or 200 least toxic seed duplexes for each of the two human and the two mouse cells (**Fig. 2b**,

Supplementary Table 1). In all four cell lines, we found that a high G content towards the 5' end of the seed

and a C in position 6 was most toxic (**Fig. 2b** and **Fig. S3c**). In contrast, non-toxic seeds were A and U-rich

especially when positioned at the 5' end. While the rules of toxicity that emerged are almost identical

between human and mouse, we noticed a subtle difference. C-rich 6mer seeds are slightly more toxic to

mouse than to human cells. This can be seen in the nucleotide composition of the most toxic seeds (**Fig. 2b**),

a slightly higher toxicity of the most C rich seeds in the siRNA retest (**Fig. 2a**), and this can also be explored

at 6merdb.org.

siRNAs with toxic seeds target housekeeping genes enriched in C-rich sequences in their 3'UTR

We previously showed that si- and shRNAs are toxic through 6mer seed toxicity preferentially targeting

hundreds of genes critical for cell survival [20]. We had developed a Toxicity Index (TI), a simple tool to

predict the most toxic seeds based on the ratio of putative seed match occurrences in the 3'UTR of set of

survival genes (SGs) versus a set of genes not involved in cell survival (nonSGs). We now compared the TI

with our experimentally determined 6mer seed toxicity in the three cell lines screened (**Fig. S4**) and found a

significant correlation between these two types of analyses, further supporting the mechanism of toxicity.

Knowing that seed sequences rich in G are most toxic suggested the targeted genes carry C-rich seed

matches in their 3'UTR. To be most stringent, we used a list of the 20 6mer seed containing siRNAs that

were most toxic to both HeyA8 and H460 cells (**Supplementary Table 2**). The G richness towards the 5'

end of the 6mers in these toxic seeds and the 5' A/U richness of the nontoxic seeds was even more

pronounced than in the top/bottom 200 most toxic seeds (**Fig. 3a**). We scored for the occurrence of seed

matches to the 20 seeds in each group in the 3'UTR, the CDS and the 5'UTR of a set of 938 critical SGs

similar to one recently described [20] and an expression-matched set of 938 nonSGs. We found a significantly

higher count ratio of toxic versus nontoxic seed matches in the 3'UTR of SGs when compared to nonSGs

(**Fig. 3b**, right panel). Consistent with a miRNA-like function no such enrichment was found when the CDS

was analyzed (**Fig. 3b**, center panel). An inverse ratio of sequences complementary to the 6mer seeds of

unknown significance was found in the 5'UTR (**Fig. 3b**, left panel). This result was very similar when we

scored for seed matches to the 100 seeds most toxic and least toxic to both human cell lines (**Fig. S5a**). An

enrichment of the exact seed matches in 3'UTRs was consistent with the overall higher C content of 3'UTRs

of SGs when compared to nonSGs (different peak maxima in **Fig. 3c**). A metaplot analysis of the 500 bases

upstream and downstream of the translational start and stop site of all human genes showed that as expected

the 3'UTR was enriched in A and U (**Fig. 3d**, top). Interestingly, SGs had a lower A/U content in a region

~150-500 bases into the 3'UTR than expression matched nonSGs (**Fig. 3d**, green horizontal bar, center and

bottom). To determine where survival genes are being targeted by toxic seed containing siRNAs we again

performed a metaplot analysis this time plotting the locations of seed matches to the 20 6mer seeds that were

most and least toxic to both human cell lines (**Fig. 3e**, an analysis with the 100 most/least toxic seeds is

provided in **Fig. S5b**). When analyzing all human coding genes we found the reverse complements of the

most toxic seeds to be highly enriched at the beginning of the 3'UTR whereas the reverse complements of the

least toxic seeds were underrepresented in this region (**Fig. 3e**, top). This effect was not due to a much higher

G/C or lower A/U content in this region (**Fig. 3d**, top). A comparison of the location of these seed matches in

the SGs and in expression matched nonSGs confirmed this general trend, however, two differences between

SGs and nonSGs became apparent: 1) nonSGs have more nontoxic seed matches ~150-500 bases into the

3'UTR (**Fig. 3e**, bottom, green horizontal bar) maybe due to the higher A/U content of this region (**Fig. 3d**,

center and bottom). 2) SGs contain a small stretch at positions 42-65 into the 3'UTR (**Fig. 3e** and **Fig. S5b**,

center, red horizontal bar) that is enriched in seed matches for the most toxic seeds. This region in SGs seems

to be a preferential target site for siRNAs carrying toxic G-rich seed sequences.

Tumor suppressive miR-34a-5p kills cancer cells through its toxic 6mer seed

The toxic siRNAs kill cancer cells through 6mer seed toxicity by a mechanism reminiscent of the function of

miRNAs. To test whether actual miRNAs could kill cancer cells with the help of toxic 6mer seeds, we

analyzed the seed toxicity determined in our screen for all known ~2600 mature miRNAs expressed as either

the 3p or 5p arm (**6merdb.org**). While none of the 6mer seeds present in the predominant arm (guide strand)

of the most oncogenic miRNAs (miR-221/222, miR-21, miR-155, the miR-17~92, miR-106b~25, and miR-

106a~36 clusters) were toxic (reduced viability >50%, stippled line in **Fig. 4a**), two of the major tumor

suppressive miRNA families, miR-15/16 and p53 regulated miR-34a/c and miR-34b contained toxic seeds in

the guide strand (**Fig. 4a**). This suggested these two families were killing cancer cells through toxic 6mer

seeds. Interestingly, two other major tumor suppressive families, let-7 and miR-200, did not contain toxic G-

rich seeds in their guide strand, suggesting they may be tumor suppressive through other mechanisms, such

as inducing and maintaining cell differentiation [26].

When transfecting the pre-miRs of miR-34a-5p, miR-15a-5p and let-7a-5p into HeyA8 cells, the potency

of these three miRNAs to reduce cell growth mimicked the toxicity of their 6mer seed containing siRNAs

(**Fig. 4b** and **4a**). This suggested that a large part of their toxicity comes from the composition of the seed

position 2-7. The most toxic seed in a major tumor suppressive miRNA was present in miR-34a5p/34c-5p, a

master regulator of tumor suppression [27]. We directly compared the toxicity of pre-miR-34a-5p and its toxic

seed in the neutral scaffold with blocked passenger strand (si34a-5p^{Seed}) in the same assays (**Fig. 4c**).

Strikingly, the toxicity evoked by these two RNA species (assessed by growth inhibition and DNA

fragmentation) was similar. Cells showed the typical morphology we found in cells dying from toxic siRNAs

(**Fig. 4d**) [20, 24, 28]. To determine the contribution of the 6mer seed sequence of miR-34a-5p to its toxicity and

the mode of cell death, we performed a RNA-Seq analysis on HeyA8 cells transfected with either miR-34a-

5p or si34a-5p^{Seed} (**Fig. 4e**, top) (**Supplementary Table 3**). The vast majority of genes were significantly up-

and downregulated by both RNA duplexes (**Fig. 4e**, bottom). While miR-34a-5p targeted a subset of genes

not affected by miR-34a-5p^{Seed}, the majority of genes (>78%) were downregulated >1.5 fold by both the

premiR and the 6mer seed duplex (**Fig. 4f**, left). A Sylamer analysis is a unbiased approach allowing to

identify which seed matches are enriched in the 3'UTRs of downregulated genes from RNAseq data [29]. In

this analysis both duplexes caused similar and highly effective downregulation of the mRNAs that carry a

6mer seed match (**Fig. 4g**). When the Sylamer analysis was performed with either 7mer or 8mer seeds,

enrichment of seed matches was much less significant (**Fig. S6a**) suggesting that most of the targeting by

both RNAs only required a 6mer seed.

Consistent with this activity, targeting by both RNA duplexes resulted in a very similar reduction of

survival genes (**Fig. 4h**). The genes downregulated by both the premiR and the 6mer seed construct were

highly enriched in genes involved in regulation of cell cycle, cell division, DNA repair, and nucleosome

assembly (**Fig. 4f**, right). These GO terms were very similar to the ones we found enriched in downregulated

genes in cells dying after transfection with CD95R/L-derived si/shRNAs containing toxic 6mer seeds [20]. In

contrast, no such GO terms were found enriched when the same analysis was performed with the upregulated

genes as control (**Fig. S6b**). While both miR-34-5p and si34a-5p^{Seed} caused the most significant

downregulation of genes carrying 8mers in their 3'UTR (**Fig. S6c**), only the most highly downregulated

genes that carry the shared 6mer seed match were grouped in a number of GO terms that are consistent with

6mer seed toxicity as previously reported [20] and barely any GO terms were shared among genes that

contained 7 or 8mer seed matches (**Fig. S6d**). All these data suggest that miR-34a-5p kills cancer cells using

its toxic 6mer seed. While optimal miRNA targeting requires at least a 7mer seed interaction and also

involves nucleotides at positions 13-16 of the miRNA [30], the cell death inducing activity of this tumor

suppressive miRNA may only require the 6mer seed.

Toxic 6mer seed toxicity is shaping the miRNA repertoire

Toxic 6mer seeds may be a driving force in miRNA evolution, whereby toxic seed sequences are either

selected against - because they contribute to cell toxicity - or are preserved to operate as tumor suppressors.

Based on the composition of toxic 6mer seeds and the enrichment of corresponding seed matches in survival

genes, we could now ask whether and when miRNAs that contain toxic G-rich sequences in positions 2-7 of

their seeds evolved. When comparing all mature miRNA arms annotated in TargetScan Human 7, we noticed

that miRNAs in highly conserved miRNA seed families contained 6mer seed sequences that were much less

toxic in our screen than seeds in poorly conserved miRNAs (**Fig. 5a,** left panel and **Fig. S7a**). Weakly

conserved miRNA seed families would be expected to be younger in evolutionary age than highly conserved

ones. Consistent with this assumption we found that the 6mer seeds of younger miRNAs (<10 million years

old) were more likely to be toxic to cells than the ones of older miRNAs (>800 million years old) [31] (**Fig. 5a,**

right panel and **Fig. S7b**). Most importantly, when comparing miRNAs of different ages, it became
apparent that seeds of miRNAs over the last 800 million years were gradually depleted of G beginning at the
5' end and eventually also affecting positions 3-5 until the oldest ones, where A and U had replaced G as the
most abundant nucleotide in all six positions (**Fig. 5b**). These analyses indicated that most highly conserved
miRNAs avoid G in potentially toxic seed positions. Interestingly, the most toxic seed sequences were found
in miRtrons (**Fig. 5c** and **Fig. S7c**), miRNAs that are derived by splicing short introns [32].

miRNAs are expressed as pre-miRs and usually only one major species of mature miRNA (either the 5p
or the 3p arm) is significantly expressed in cells produced from one of the two strands of the premiR stem [33].
Consistent with the assumption that cells cannot tolerate toxic 6mer seeds, we now tested across 780
miRNAs which have been shown to give rise to both a 3p and a 5p arm whether the more highly expressed
arm contains a seed of lower toxicity than the lesser expressed arm (**Fig. 5d**). We ranked the miRNAs
according to the ratio of the 6mer seed toxicity associated with the guide arm to the lesser-expressed arm
(**Supplementary Table 4**). When we labeled the major tumor suppressive and oncogenic miRNAs, we
noticed the highly expressed arm of most of the oncogenic miRNAs contained a 6mer seed that was not toxic
in our screen (**Fig. 5d**, green dots). In contrast, for most of the tumor suppressive miRNAs, the dominant arm
contained a seed much more toxic than the lesser arm (**Fig. 5d**, red dots). The overall difference in ratio
between the two groups of miRNAs was highly significant. A more detailed analysis of these data revealed
that the three oncogenic miRNAs with the highest ratio in toxicity between their arms, miR-363, miR-92a-2,
and miR-25, were almost exclusively expressed as the non-toxic 3p form (**Fig. 5e**, top). In contrast, the
dominant arm of the three tumor suppressive miRNAs, miR-34a, miR-34c, and miR-449b contained the
most toxic seed sequence (**Fig. 5e**, bottom). Interestingly, miR-449b has the same seed sequence as miR-34a
and has been suggested to act as a backup miRNA for miR-34a [34]. These data are consistent with most tumor
suppressive miRNAs using 6mer seed toxicity to kill cancer cells and suggest that this mechanism developed
over hundreds of millions of years.

Genotoxic drugs upregulate toxic 6mer seed containing miRNAs
Our data showing that miR-34a-5p contains a toxic 6mer seed, along with miR-34a being upregulated after
genotoxic stress [19], led us to wonder whether miR-34a-5p would contribute to cell death induced by
genotoxic drugs and whether this type of cell death had similarities to the death observed in cells dying from
toxic 6mer seed containing si/shRNAs. This would be consistent with the observation that many genotoxic
drugs induce multiple cell death pathways [35, 36, 37, 38, 39]. To compare cell death induced by different genotoxic
agents with that of toxic si/shRNAs, we treated the p53 wild-type ovarian cancer cell line HeyA8 with
doxorubicin (Doxo), carboplatin (Carbo), or etoposide (Eto) and performed a RNA-Seq analysis. Drug
concentrations were chosen so that after 80 hrs, treatment would slow down cell growth and induce signs of

stress without major cell death occurring to capture changes that could be causing cell death rather than

being the result of it (**Fig. S8a** and data not shown). The morphological changes in the cells treated with the

drugs were very similar to the ones seen in cells treated with si/shRNAs (**Fig. S8b**), and similar to reported

morphologies of cells treated with genotoxic drugs [40, 41].

The ranked lists of downregulated RNAs isolated from HeyA8 cells treated with the three drugs were

subjected to a gene set enrichment analysis (GSEA) to determine whether survival genes were enriched in

the downregulated genes (**Fig. S9a**). There was strong enrichment of downregulated survival genes towards

the top of the ranked list. 102 of the survival genes were downregulated in cells treated with any of the three

drugs (**Fig. S9b**). In a DAVID gene ontology analysis, these genes were strongly enriched in clusters

involved in chromosome segregation, DNA replication, cell cycle regulation, and mitosis, typical for 6mer

seed toxicity induced cell death (**Supplementary Table 5**). We quantified 30 of the 102 survival genes in

HeyA8 cells treated with Doxo at different time points using an arrayed quantitative PCR (**Fig. S9c**). 24 of

the 30 genes' mRNAs were significantly downregulated as early as 7 hours after treatment with no further

reduction beyond 15 hrs after treatment, suggesting that their repression was the cause of cell death rather

than a consequence. A Metascape analysis of all RNA-Seq data of downregulated RNAs in response to the

toxic siL3, si34a-5p[Seed], miR-34a-5p, and the three genotoxic drugs suggested a common mode of action

(**Fig. S9d**). The GO clusters that were most significantly downregulated in all data sets were again related to

DNA repair, cell cycle, and mitosis as described before [24].

To test whether treatment of cells with genotoxic drugs results in loading the RISC with toxic miRNAs,

HeyA8 cells were treated with Doxo for 0, 20, 40 and 80 hrs and all 4 Ago proteins were pulled down using

a GW182 peptide [42]. Interestingly, while the amount of AGO2 pulled down was the same at all time points,

the amount of bound miRNA-sized RNAs substantially increased with longer treatment times (**Fig. 6a**). This

was most likely the result of an overall increase in total small RNAs in the treated cells (**Fig. 6b**).

Alternatively, this could also be a result of cells dividing more slowly and a stable RISC. After Doxo

treatment, >99% of the Ago-bound small RNAs were miRNAs (data not shown). miR-34a/b/c-5p bound to

Ago proteins were upregulated at all time points (**Fig. S10a**). To determine the contribution of miR-34a-5p

and other miRNAs to the toxicity seen in cells exposed to the genotoxic drugs, we treated Drosha k.o. cells -

devoid of most canonical miRNAs [43] - with the three genotoxic drugs (**Fig. S11**). These cells were

hypersensitive to the toxicity induced by any of the three drugs. We attributed this response to the absence of

most canonical miRNAs that protect cells from toxic RNAi active sequences [20]. This result also suggested

involvement of small RNAs that do not require Drosha for processing. As expected, the composition of small

RNAs bound to Ago proteins dramatically varied between wild-type and Drosha k.o. cells (**Fig. 6c**). In the

absence of most canonical miRNAs, miR-320a-3p, which was previously shown not to require Drosha for its

biogenesis [43], represented more than 86% of all Ago-bound miRNAs. Similar to HeyA8 cells (see **Fig. S10a**),

Ago-bound miR-34a-5p was upregulated in wild-type but not in Drosha k.o. HCT116 cells upon Doxo

treatment (**Fig. S10b**, right). Interestingly, the average 6mer seed toxicity of all Ago-bound miRNAs >1.5

fold upregulated in HCT116 wt cells was significantly higher than the ones >1.5 fold downregulated in cells

treated with Doxo (**Fig. 6d**, left). While in the Drosha k.o. cells, a number of nontoxic miRNAs were

downregulated, the only miRNA that was upregulated in the RISC (1.49 fold) was miR-320a-3p (**Fig. 6d**,

right). However, upon closer inspection it became clear that this form of miR-320a-3p was shortened by two

nucleotides at the 5' end. This resulted in the shift of the 6mer seed into a G-rich sequence (**Fig. 6d**, right),

converting a moderately toxic miRNA (average viability = 49.2%) into a highly toxic one (average viability

= 9.3%, **Fig. 1b**). To test this predicted increase in toxicity experimentally, we transfected HeyA8 cells with

either the authentic pre-miR-320a-3p or a miR-320a-3p^{Seed} duplex that corresponded to the shifted Ago-

bound miR-320a-3p sequence (miR-320a-3p^{Ago}) (**Fig. 6e**). While pre-miR-320a-3p was not toxic, miR-320a-

3p^{Ago} completely blocked the growth of the cells. Toxicity of miR-320a-3p^{Ago} was established in four human

and mouse cell lines (**Fig. 6f**). These data suggested that in the absence of other miRNAs that could kill cells

through 6mer seed toxicity, miR-320a-3p (and possibly other small RNAs) may represent an alternative

mechanism that ensures that genotoxic stressors can kill cells with defective miRNA processing often

observed in cancer [44, 45]. In summary, our data suggest that certain tumor suppressive miRNAs, such as miR-

34a-5p and miR-320a-3p exert their tumor suppressive activities by carrying toxic 6mer seed sequences that

can kill cancer cells by targeting survival genes in C-rich regions close to the start of their 3'UTR. This

activity may contribute to the cell death induced by genotoxic drugs.

DISCUSSION

We previously discovered a fundamental cell type- and species-independent form of toxicity that is evoked

by the 6mer seed sequence in si-/shRNAs that function similar to miRNAs [20]. We have now performed an

siRNA screen that effectively tested the miRNA activities of all 4096 different 6mer seed sequences.

Performing the screen in four cell lines (two human and two mouse) ensured that the results were relatively

independent of species or cell type specific transcriptomes. The screen has discovered the rules underlying

this seed toxicity and allows prediction of the 6mer seed toxicity for any siRNA, shRNA, miRNA with a

known 6mer seed (https://6merdb.org).

Based on this screen, the toxicity of a number of tumor suppressive miRNAs could be predicted solely

on the basis of their 6mer seed sequences. The enrichment of G in the first 2-3 positions of the most toxic

seeds is consistent with the way Ago proteins scan mRNAs as targets. This involves mainly the first few

nucleotides (positions 1-3) of the seed [46]. miR-34a-5p contains two Gs in positions 1 and 2 of its 6mer seed.

While miR-34a-5p is considered a master tumor suppressive miRNA, no single target has been identified to

be responsible for this activity. Over 700 targets implicated in cancer cell proliferation, survival, and

resistance to therapy have been described [16]. Our data now suggest that miR-34a-5p uses 6mer seed toxicity

to target hundreds of housekeeping genes. Our data provide the means to rationally design new artificial

miRNAs as anti-cancer reagents that attack networks of survival genes. In humans, miR-34a is highly

expressed in many tissues. Consistent with our data that delivering siRNAs with toxic 6mer seeds to mice are

not toxic to normal cells [21] miR-34a exhibits low toxicity to normal cells *in vitro* and *in vivo* [47]. miR-34a

(MRX34) became the first miRNA to be tested in a phase I clinical trial of unresectable primary liver cancer

[27, 48]. The study was recently terminated and reported immune-related adverse effects in several individuals. It

was suggested that these adverse effects may have been caused by either a reaction to the liposome-based

carrier or the use of double-stranded RNA [16]. In addition, they may be due to an undesired gene modulation

by miR-34a itself defined by sequences outside the 6mer seed [16]. Our data suggest that miR-34a exerts

toxicity mostly through the 6mer seed of its 5p arm and that its 700 known targets may be part of the

network of survival genes that are targeted. The comparison of the RNA-Seq data of cells treated with either

miR-34a-5p or si34-5p^{Seed} now allows to determine whether these two activities can be separated.

Our data provide evidence that genotoxic drugs kill cancer cells, at least in part, by triggering the toxic

6mer seed mechanism. Exposure of cancer cells to such drugs resulted in upregulation of tumor suppressive

miRNAs, most prominently of the p53 regulated miR-34 family [19]. While one report demonstrated that

inhibiting miR-34a rendered cancer cells more resistant to cell death induced by genotoxic stress [49], another

one found no effect of knocking out miR-34a on the sensitivity of HCT116 or MCF-7 cells to Doxo [50]. This

mechanism may be highly redundant and may involve many miRNAs. Our analysis of Ago-bound miRNAs

in Drosha k.o. cells suggest that in the absence of miR-34a, the noncanonical miR-320a-3p which was

recently also found to be p53 regulated [51] may act as a backup miRNA that can still respond to genotoxic

stress in case the amounts of other miRNAs are reduced, for instance in cases of mutations in miRNA

biogenesis associated genes frequently found in human cancers [44]. In addition, recent data suggest that other

toxic small RNAs can also be taken up by the RISC and negatively regulate cell growth through their toxic

6mer seed (data not shown).

It was shown before that miRNAs overall avoid seed sequences that target the 3'UTR of

survival/housekeeping genes [52, 53]. Survival genes therefore are depleted in seed matches for the most

abundant miRNAs in a cell. That also means 3'UTRs of survival genes must be enriched in sequences not

targeted by the seeds present in most miRNAs. Our combined data now suggest it is these sequences that

toxic siRNAs and tumor suppressive miRNAs with toxic 6mer seeds are targeting. Our analyses also suggest

that most miRNAs have evolved over the last 800 million years by gradually depleting G in their seeds

beginning at the 5' end. In addition, the most abundant miRNAs have evolved to use the arm with the lower

6mer seed toxicity as the active guide strand, presumably to avoid killing cells. Only in a minority of tumor-

suppressive miRNAs does the dominant guide strand contain a toxic seed. By ranking miRNAs according

to whether they express the arm with the seed of higher toxicity, it is now possible to identify novel tumor

suppressive miRNAs (see 6merdb.org).

In summary, we have determined the rules of RNAi targeting by toxic 6mer seeds. These rules allowed

us to predict with some confidence which si/shRNAs or miRNAs have the potential to kill cells through their

toxic 6mer seed. Toxic miRNAs seem to be involved in killing cancer cells in response to genotoxic drugs.

Toxic 6mer seeds are present in a number of tumor-suppressive miRNAs that can kill cancer cells. Our data

allow new insights into the evolution of miRNAs and provide evidence that 6mer seed toxicity is shaping the

miRNA repertoire.

METHODS

Reagents, cell lines and antibodies

HeyA8 (RRID:CVCL_8878) and H460 (ATCC HTB-177) cells were cultured in RPMI1640 medium

(Cellgro Cat#10-040) supplemented with 10% FBS (Sigma Cat#14009C) and 1% L-Glutamine (Corning

Cat#25-005). 3LL cells (ATCC CRL-1642) were cultured in DMEM medium (Gibco Cat#12430054)

supplemented with 10% FBS and 1% L-Glutamine. Mouse hepatocellular carcinoma cells M565 were

previously described [54] and cultured in DMEM/F12 (Gibco Cat#11330) supplemented with 10% FBS, 1% L-

Glutamine and ITS (Corning #25-800-CR). HCT116 parental (Cat#HC19023, RRID:CVCL_0291) and the

Drosha k.o. clone (clone #40, Cat#HC19020) were purchased from Korean Collection for Type Cultures

(KCTC). Both HCT116 cell lines were cultured in McCoy's 5A medium (ATCC, Cat#30-2007)

supplemented with 10% FBS and 1% L-Glutamine. All cell lines were authenticated by STR profiling. Anti-

Argonaute-2 antibody (cat#ab186733) was purchased from Abcam, anti-β-actin antibody from Santa Cruz

(#sc-47778), and secondary antibody for Western blot was Goat anti-rabbit, IgG-HRP from Southern

Biotech (#SB-4030-05). Etoposide (Cat#BML-GR307-0100) was purchased from Enzo Life Sciences,

propidium iodide (#P4864) doxorubicin (Cat#D1515) and carboplatin (Cat#C2538) were from Sigma-

Aldrich.

siRNA screens and cell viability assay

The non-toxic siRNA backbone used in the 4096 screen was designed as previously described [22]. Briefly, the

siNT2 sequence was used as a starting point and four positions in the center of siNT2 were replaced with the

complementary nucleotides in order to remove any identity between the backbone siRNA and the toxic siL3

while retaining the same GC content. Two 2'-O-methylation groups were added to positions 1 and 2 of the

passenger strand to prevent loading into the RISC. The 6mer seed region (position 2-7 on the guide strand)

was then replaced with one of the 4096 possible seeds. Transfection efficiency was optimized for each of the

four cell lines individually. For HeyA8 and M565 cells, the reverse transfection and the quantification of

cellular ATP content in the 4096 seed duplexes screen was done as previously described [22]. For 3LL and

H460 cells screens were done in a very similar manner except for the amount of RNAiMAX and the number

of cells plated. For 3LL cells, 150 cells per well were added in a 384 well plate and 9.3 μl of Lipofectamine

RNAiMax was mixed with 990.7 μl of Opti-MEM. For H460 cells, 420 cells per well in a 384 well plate was

added and 7.3 μl of Lipofectamine RNAiMax was mixed with 993.7 μl of Opti-MEM.

Transfection with short oligonucleotides

For IncuCyte experiments, HeyA8 cells were plated in 50 μl antibiotic free medium in a 96 well plate at

1000 cells/well, and 50 μl transfection mix with 0.1 μl RNAiMAX and siRNAs or miRNA precursors were

added during the plating. For the AGO2 knockdown experiment, 100,000 cells/well HeyA8 cells were

reverse-transfected in six-well plates with either non-targeting (Dharmacon, cat#D-001810-10-05) or an

AGO2 targeting siRNA SMARTpool (Dharmacon, cat#L004639-00- 005) at 25 nM. 1 μl RNAiMAX per

well was used for HeyA8 cells. Twenty-four hours after transfection with the SMARTpools, cells were

reversed-transfected in a 96-well plate with siNT2, si2733, or si2733 (see **Supplementary Table 1**) at 10

nM and monitored in the IncuCyte Zoom. 48 hours after transfection, cells were lysed in RIPA buffer for

Western blot analysis as previously described [22].

All custom siRNA oligonucleotides were ordered from integrated DNA technologies (IDT) and annealed

according to the manufacturer's instructions. In addition to the 4096 siRNAs of the screen the following

siRNA sequences were used:

siNT1 sense: rUrGrGrUrUrUrArCrArUrGrUrCrGrArCrUrArATT;

siNT1 antisense: rUrUrArGrUrCrGrArCrArUrGrUrArArArCrCrAAA;

siNT2 sense: rUrGrGrUrUrUrArCrArUrGrUrUrGrUrGrUrGrATT;

siNT2 antisense: rUrCrArCrArCrArArCrArUrGrUrArArArCrCrAAA;

siL3 sense: rGrCrCrCrUrUrCrArArUrUrArCrCrCrArUrArUTT;

siL3 antisense: rArUrArUrGrGrGrUrArArUrUrGrArArGrGrGrCAA;

si-miR-34a-5p^{Seed} sense: mUmGrGrUrUrUrArCrArUrGrUrArCrUrGrCrCrATT;

si-miR-34a-5p^{Seed} antisense: rUrGrGrCrArGrUrArCrArUrGrUrArArArCrCrAAA;

miR-320a-3p^{Ago} sense: mCmGrCrCrCrUrCrUrCrArArCrCrCrArGrCrUrUTT

miR-320a-3p^{Ago} antisense: rArArGrCrUrGrGrGrUrUrGrArGrArGrGrGrCrGAA.

The following miRNA precursors and negative controls were used: hsa-miR-34a-5p (Ambion, Cat. No#

PM11030), hsa-let-7a-5p (Ambion, Cat. No# PM10050), hsa-miR-320a-3p (Ambion, Cat. No# PM11621),

hsa-miR-15a-5p (Ambion, Cat. No# PM10235), and miRNA precursor negative control #1 (Ambion, Cat.

No# AM17110).

Monitoring growth over time by IncuCyte and cell death assays

Cells were seeded between 1000 and 3,000 per well in a 96-well plate in triplicates. The plate was then

scanned using the IncuCyte ZOOM live-cell imaging system (Essen BioScience). Images were captured

every six hours using a 10× objective. Cell confluence was calculated using the IncuCyte ZOOM software

(version 2015A). For treatment with genotoxic drugs HeyA8 cells were seeded at 750 cells/well and

HCT116 cells were seeded at 3000 cells/well in 96-well plate and treated with one of the three genotoxic

drugs (carboplatin, doxorubicin, or etoposide) at various concentrations at the time of plating. Solvent treated

(0.025% DMSO in medium) cells were used as control for etoposide. Medium treated cells were used as

control for carboplatin and doxorubicin. To assess cell viability, treated cells were subjected to a

quantification of nuclear fragmentation or ATP content both as previously described [20, 22].

RNA-Seq analysis

For RNA-Seq data in **Fig. 4e**, 50,000 cells/well HeyA8 cells were reversed transfected in 6-well plates with

10 nM of either pre-miR-34a-5p or si-miR-34a-5p^{Seed} with their respective controls. The transfection mix was

replaced 24 hours after transfection. Cells were lysed after 48 hours after transfection using Qiazol. For the

RNA-Seq data in **Fig. S9**, HeyA8 cells were seeded at 50,000 cells per well in a 6-well plate and treated with

three genotoxic drugs in duplicate: carboplatin (25 μg/ml), doxorubicin (50 ng/ml), and etoposide (500 nM).

Medium treated cells were used as control for carboplatin and doxorubicin treated cells. Solvent control

treated cells (0.025% DMSO in medium) were used as control for etoposide. Cells were lysed after 80 hours

drug incubation using Qiazol. Total RNA was isolated using the miRNeasy Mini Kit (Qiagen, Cat.No#

74004) following the manufacturer's instructions. An on-column digestion step using the RNAse-free

DNAse Set (Qiagen, Cat.No# 79254) was included for all RNA-Seq samples. The RNA libraries were

prepared and sequenced as previously described [20] at the Genomics Core facility at the University of

Chicago. Reads were aligned to the hg38 version of the human genome, using either Tophat v2.1.0 (RNA-

Seq data in **Fig. S9**) or STAR v2.5.2 (RNA-Seq data in **Fig. 4e**). In either case, aligned reads were

associated with genes using HTSeq v0.6.1, and the UCSC hg38 transcriptome annotation from

iGenomes. Differentially expressed genes were identified using the edgeR R package.

Arrayed real-time PCR

The top 30 most downregulated survival genes shared among HeyA8 cells treated with carboplatin,

doxorubicin, and etoposide based on the RNA-Seq analysis were selected for a kinetics analysis using real-

time PCR. To prepare the RNAs for the kinetics analysis, 75,000/plate HeyA8 cells were plated in 15 cm

plate. Twenty-four hours after plating, one plate of HeyA8 cells were lysed in QIAzol as the control sample.

The rest of the plates were treated with 50 ng/ml doxorubicin for 7 hrs, 14.5 hrs, and 21 hrs respectively

before being lysed in QIAzol. To perform the arrayed real-time PCR, 200 ng total RNA per sample was used

as the input to make cDNA using the high-capacity cDNA reverse Transcription Kit (Applied Biosystems

#4368814). For TaqMan Low Density Array (TLDA) profiling, custom-designed 384-well TLDA cards

(Applied Biosystems, Cat. No#4346799) were selected and used according to the manufacturer's protocols.

For each sample, 20 μl cDNA was mixed with 80 μl water and 100 μl TaqMan Universal PCR Master Mix

(Applied Biosystems, Cat. No#4304437). A total volume of 100μl of each sample is loaded into the 8

loading ports on the TLDA card (2 ports for each sample, 4 samples total on one card). The qPCR assays

used to detect the 30 genes on the TLDA card are as follows: HIST1H2AI (Hs00361878_s1), CENPA

(Hs00156455_m1), HJURP (Hs00251144_m1), FAM72D (Hs00416746_m1), CCNA2 (Hs00996788_m1),

KIF20A (Hs00993573_m1), PRC1 (Hs01597839_m1), KIF15 (Hs01085295_m1), BUB1B

(Hs01084828_m1), SCD (Hs01682761_m1), AURKA (Hs01582072_m1), NUF2 (Hs00230097_m1),

NCAPH (Hs01010752_m1), SPC24 (Hs00699347_m1), KIF11 (Hs00189698_m1), TTK (Hs01009870_m1),

PLK4 (Hs00179514_m1), AURKB (Hs00945858_g1), CEP55 (Hs01070181_m1), HMGCS1

(Hs00940429_m1), TOP2A (Hs01032137_m1), KIF23 (Hs00370852_m1), INCENP (Hs00934447_m1),

CDK1 (Hs00938777_m1), HIST2H2BE (Hs00269023_s1), KNL1 (Hs00538241_m1), NCAPD2

(Hs00274505_m1), RACGAP1 (Hs01100049_mH), SPAG5 (Hs00197708_m1), KNTC1 (Hs00938554_m1).

GAPDH (Hs99999905_m1) was used as the endogenous control. qPCR assay for individual gene was done

in technical triplicates on each TLDA cards. Statistically analysis was performed using Student's t test.

Ago affinity peptide purification

The FLAG-GST-T6B WT and mutant peptides was expressed and purified as described [42]. Briefly,

constructs were expressed in BL21-Gold(DE3)pLysS competent cells (Agilent). Bacteria, induced with 1

mM Isopropyl β-D-1-thiogalactopyranoside (IPTG), were grown in 1 L overnight at 18ºC to OD 0.6. The

bacteria were sedimented at 4000*g* for 15 min and resuspended in 25 ml GST-A buffer (1 mM 4-(2-

aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF), 1 mM DTT in PBS) supplemented with 1

mg/mL lysozyme (Sigma). Samples were sonicated three times for 3 min at 100% amplitude (Sonics,

VCX130) and cleared by centrifugation at 20,000*g* for 20 min. The lysate was loaded onto a column

containing 2 ml of bead volume glutathione Sepharose beads (Sigma) and washed two times with GST-A

buffer. The GST-tagged protein was eluted in 10 ml of GST-B buffer (20 mM Tris, pH 8.0, and 10 mM

glutathione in PBS). The peptide was concentrated using Amicon Ultra-15 Centrifugal Filter Unit (Millipore)

and desalted using Zeba Spin Desalting Columns (ThermoFisher).

Ago pull down and small RNA-seq

HeyA8 (5-7 x 10^6), HCT116 wild-type (1.2-1.6 x 10^8) or Drosha k.o. (4.8-6.3 x 10^7) cells treated with doxorubicin were lysed in NP40 lysis buffer (20 mM Tris, pH 7.5, 150 mM NaCl, 2 mM EDTA, 1% (v/v) NP40, supplemented with phosphatase inhibitors) on ice for 15 minutes. The lysate was sonicated 3 times for 30 s at 60% amplitude (Sonics, VCX130) and cleared by centrifugation at 12,000g for 20 minutes. AGO1-4 were pulled down by using 500 µg of Flag-GST-T6B peptide [42] and with 60 µl anti-Flag M2 magnetic beads (Sigma-Aldrich) for 2 hrs at 4°C. The pull-down was washed 3 times in NP40 lysis buffer. During the last wash, 10% of beads were removed and incubated at 95°C for 5 minutes in 2x SDS-PAGE sample buffer. Samples were run on a 4-12% SDS-PAGE and transferred to nitrocellulose membrane. The pull-down efficiency was determined by immunoblotting against AGO2 (Abcam 32381). To the remaining beads 500 µl TRIzol reagent were added and the RNA extracted according to the manufacturer's instructions. The RNA pellet was diluted in 20 µl of water. The sample was split and half of the sample was dephosphorylated with 0.5 U/µl of CIP alkaline phosphatase at 37°C for 15 min and subsequently radiolabeled with 0.5 µCi γ-^{32}P-ATP and 1 U/µl of T4 PNK kinase for 20 min at 37°C. The AGO1-4 interacting RNAs were visualized on a 15% urea-PAGE. The remaining RNA was taken through a small RNA library preparation as previously described [55]. Briefly, RNA was ligated with 3' adenylated adapters and separated on a 15% denaturing urea-PAGE. The RNA corresponding to insert size of 19-35 nt was eluted from the gel, ethanol precipitated followed by 5' adapter ligation. The samples were separated on a 12% Urea-PAGE and extracted from the gel. Reverse transcription was performed using Superscript III reverse transcriptase and the cDNA amplified by PCR. The cDNA was sequenced on Illumina HiSeq 3000.

Adapter sequences:

Adapter 1 – NNTGACTGTGGAATTCTCGGGTGCCAAGG;

Adapter 2 – NNACACTCTGGAATTCTCGGGTGCCAAGG,

Adapter 3 – NNACAGAGTGGAATTCTCGGGTGCCAAGG,

Adapter 4 – NNGCGATATGGAATTCTCGGGTGCCAAGG,

Adapter 47 – NNTCTGTGTGGAATTCTCGGGTGCCAAGG,

Adapter 48 – NNCAGCATTGGAATTCTCGGGTGCCAAGG,

Adapter 49 – NNATAGTATGGAATTCTCGGGTGCCAAGG,

Adapter 50 – NNTCATAGTGGAATTCTCGGGTGCCAAGG.

RT primer sequence: GCCTTGGCACCCGAGAATTCCA;

PCR primer sequences:

CAAGCAGAAGACGGCATACGAGATCGTGATGTGACTGGAGTTCCTTGGCACCCGAGAATTCCA.

Data analyses

GSEA was performed using the GSEA software version 3.0 from the Broad Institute downloaded from https://software.broadinstitute.org/gsea/. A ranked list was generated by sorting genes according the Log_{10}(fold downregulation). The Pre-ranked function was used to perform GSEA using the ranked list. 1000 permutations were used. Default settings were used. The ~1800 survival genes and ~420 non-survival genes defined previously [20] were used as custom gene sets. Default settings were used.

The list of survival genes and expression matched non-survival genes were generated by taking the survival and expression matched non-survival genes used previously (*20*) and retaining only the 938 genes in each group of expression matched survival and non-survival genes with an average expression across all RNA seq datasets above 1000 RPMs (see **Supplementary Table 6**).

Sylamer analysis [29] was used to find enrichment of small word motifs in the 3'UTRs of genes enriched in those that are most downregulated. A list of 3'UTRs was generated by eliminating repetitive regions as described previously [20] to use for the Sylamer analysis. Default settings were used. Bonferroni adjusted p-values were calculated by multiplying the unadjusted p-values by the number of permutations for each length of word searched for.

The GO enrichment analyses shown in **Fig. 4f** and **Fig. S6b** were performed using the GOrilla GO analysis tool at http://cbl-gorilla.cs.technion.ac.il using default setting using different p-value cut-offs for each analysis. GO analysis in **Supplementary Table S5** was done using DAVID 6.8 (https://david.ncifcrf.gov) using default settings. GO analyses across multiple data sets were performed using the software available on www.Metascape.org and default running parameters.

Density plots showing the contribution of the four nucleotides G, C, A and U at each of the 6mer seed positions were generated using the Weblogo tool at http://weblogo.berkeley.edu/logo.cgi using the frequency plot setting.

Venn diagrams were generated using http://bioinformatics.psb.ugent.be/webtools/Venn/ using default settings.

The scatter plot in **Fig. 4e** was generated using R package ggplot2. 10875 genes with RPM > 1 (average RPM of the 8 RNAseq samples) and adjusted p-value < 0.05 were included. 3696 genes were significantly upregulated in both mir-34a-5p and si34a-5p^{Seed} treated samples. 4207 genes were significantly downregulated in both mir-34a-5p and si34a-5p^{Seed} treated samples. 713 genes were only downregulated and 792 genes were only upregulated in si34a-5p^{Seed} treated samples. 730 genes were only downregulated and 737 genes were only upregulated in mir-34a-5p treated samples. 193 genes out of the total 10875 genes were omitted in the graph as the range for X and Y axes were set as -3 to 3.

Identification the 6mer seeds most and least toxic to human cancer cells.

To identify the 20 and 100 most and least toxic seeds to both human cell lines all 4096 seeds were ranked

for each cell line from highest to lowest toxicity. The 20 seeds with the highest toxicity to both HeyA8 and

H460 cells were found in the top 46 most toxic seeds to in both cells and the 20 seeds shared to be least toxic

were found in the bottom 149 seeds in each ranked group. The 100 most and least toxic seeds for both cell

lines were identified in the same way and all groups of seeds are shown in **Supplementary Table 2**.

Metaplots of 6-mer seed match locations

3pUTR sequences were downloaded from Ensembl Biomart. In order to reduce redundancy in the sequences,

a single longest 3pUTR (and associated transcript) was chosen to represent each gene. A custom perl script

(makeSeedBed.pl) was written to identify exact matches to all seeds (reverse complement) in all sequences,

and to output the coordinates of those matches in bed file format. A custom R script (plotBedMetaPlot.R)

was written that uses the GenomicRanges (https://doi.org/10.1371/journal.pcbi.1003118) and Sushi

(https://dx.doi.org/10.1093%2Fbioinformatics%2Fbtu379) R packages to calculate the coverage of seeds

across all sequences in a given set, and to create a plot of that coverage. The custom scripts and the input

data are available in the cloud-based computational reproducibility platform Code Ocean at

https://codeocean.com/capsule/9a3eb292-6e89-44f0-b9b0-bdd779f97516/code .

eCDF plots

A custom perl script (annotateWithSeeds.pl) was written to identify exact seed matches (reverse

complement) to all seeds in all sequences, and to output the total counts of the different types of seeds

(generally toxic vs. nontoxic) in the sequences. To compare the presence of toxic and nontoxic seed matches

in expression matched survival and non survival genes, a custom R script (makeECDFplot.180615.R) was

written that takes as input two different sets of genes (SGs and nonSGs) and the list of the counts of toxic

and nontoxic seeds (reverse complement) in all genes, and plots the cumulative distribution function for the

count statistics in each gene set. In **Fig. 3b** the ratio of the seed match counts to the 20 most and least toxic

seeds in the 5'UTR, CDS, first 1000bp of 3'UTR, and full 3'UTR (not shown) were compared between pairs

of 938 expression matched survival and non-survival genes. In **Fig. S5a**, this analysis was repeated with the

100 most and least toxic seeds to both human cell lines. The custom scripts and the input data are available in

the cloud-based computational reproducibility platform Code Ocean at

https://codeocean.com/capsule/b755ec2b-00d8-4281-9fa1-2a484fd7521b/code. To determine the dependence

of mRNAs regulation on miR-34a-5p seed presence in their 3'UTR, a custom R script

(makeECDFplot.cetoData.R) was written that takes as input a list of gene sets and a table of logFC

expression for those genes upon miR-34a-5p or si34a-5p^{Seed} over-expression. This Rscript then plots the

cumulative distribution function for the logFC expression data in each gene set. The custom scripts and the

input data are available in the Code Ocean at https://codeocean.com/capsule/31ec8deb-8282-4a90-98e6-

b80a0ba881cb/code.

Relation between miRNA seed conservation, age and toxicity

Information on miRNA seed family conservation and seed sequence were downloaded at

http://www.targetscan.org/vert_71/ from TargetScan Human 7.1. The toxicity of each mature human miRNA

arm sequence in the TargetScan dataset was assigned according to the average toxicity induced by the

siRNA in HeyA8 and H460 siRNA screens harboring the identical 6mer seed sequence. A list of miRNA

ages corresponding to ~1025 miRNA loci was acquired from [31] and was calculated using a modified version

of ProteinHistorian [31]. This list was used to assign ages to roughly ~1400 mature miRNA arms found in the

TargetScan dataset.

TargetScan 7.1 partitions the seed family conservation into four groups: highly conserved (group #2),

conserved (group #1), low conservation but still annotated as a miRNA (group #0), and low conservation

with the possibility of misannotation (group #-1). Probability density and eCDF plots for the assigned seed-

dependent toxicity were generated for each seed family conservation group using ggplot2 in R. Probability

density and eCDF plots were also generated to show how young (<10 million years) and old (>800 million

years) miRNAs compare in terms of the seed-dependent toxicity. All differences between groups in terms of

seed-dependent toxicity (always the average of the toxicity determined in HeyA8 and H460 cells) were

analyzed using a two-sample two-tailed Kolmogorov-Smirnov test in R.

Assessment of dominant arm seed toxicity

The expression (RPM) of miRNA 5p and 3p arms across 135 tissue samples were collected from MIRMINE

[56]. A miRNA was considered expressed if the sum of the normalized reads for both arms was above 5 for

each sample. A value of 0 was replaced with 0.01 to avoid a division by 0 error. A miRNA arm was

considered the dominant species if its expression was at least 25% greater than the other arm per sample. The

dominant arm for each miRNA across all samples was calculated by determining which arm was dominantly

expressed in more samples (>50% of samples where the miRNA was considered expressed). The miRNAs

that have only one annotated arm in miRBase were considered to have only one dominant arm. The seed

toxicity values for each arm were extracted from the 4096 siRNA screen data (% average viability for the

human HeyA8 and H460 cells) and used to calculate the ratio between the dominant arm's toxicity and the

lesser arm's toxicity; miRNAs that only had one expressed arm were not considered in the analyses shown in

Fig. 5d but are all included on the website: 6merdb.org.

To compare the 6mer seed toxicity between up regulated and down regulated Ago bound miRNA

populations in HCT116 cells after Doxo treatment we analyzed the reads obtained from RNA Seq analysis of

Ago bound RNAs. After removing the reads that were either shorter than 19 nt or longer than 26 nt in
length, the reads were blasted against a miRNA database consisting of all human miRNA mature sequence
information obtained from miRBase. A threshold of 100% identity for an at least 16 nt-long stretch without
any gaps was set for the BLAST analysis. After discarding sequences with no significant BLAST result, the
remaining sequences were trimmed from the 3'end so that all reads were now 19nt in length. This was done
to determine for each miRNA the relevant 5' end to obtain the 6mer seed sequence (position 2-7). All the
reads that shared the same 5' sequence and miRNA names were collapsed while adding up the number of
reads of each sequence in each condition. To compare the 6mer seed toxicity between up and down regulated
miRNAs, we calculated the average 6mer seed toxicity for miRNA sequences that were either 1.5 fold up or
1.5 fold down regulated in Doxo treated wt samples compared to medium treated control samples (after
removing sequences that had less than 100 collapsed reads in Doxo treated wt samples). In each group
miRNAs were ranked according to highest base mean expression and groups were compared. Statistically
significance was determined using the Mann-Whitney U test. The comparison was repeated for Drosha k.o.
cells where the 5' shifted form of miR-320a-3p was the only miRNA found to be upregulated (1.5 fold).
656

Analysis of miRtron and non-miRtron groups
MiRtrons and non-miRtrons were recently reported [57] and consisted of miRNAs that are listed in miRBase
v21 as expressing both arms. Comparing 6mer toxicity of miRtrons and non-miRtrons from this list was
done as described above for young/old non-conserved/conserved miRNAs. Both arms were considered.
Probability density (**Fig. 5c**) and eCDF plots (**Fig. S7c**) were generated to show how miRtrons and non
miRtrons compare in terms of the seed-dependent toxicity.
663

Comparing the seed viability of shRNAs derived from the CD95L sequence
An RNAi lethality screen composed of every shRNA sequence that can be derived from the CD95L CDS
was conducted previously (*20*). In this screen, toxicity of each shRNA was assessed in two ways: 1) fold
underrepresentation of the shRNA after infection with the shRNA-expressing lentivirus compared to its
representation in the plasmid pool and 2) fold representation of the shRNA after infection and treatment with
doxycycline compared to cells that were infected but did not receive doxycycline. The first analysis allowed
us to quantify toxicity associated with "leaky" shRNA expression. The second analysis quantified toxicity
associated with strong shRNA expression following treatment with doxycycline.
For each shRNA, the average fold downregulation was calculated from both of these toxicity assessments.
Then, the seed sequence of each shRNA was extracted and assigned an average viability score, which was a
composite of the % viabilities determined in the 4096 siRNA arrayed screen for both HeyA8 and H460 cells.

Pearson's correlation was determined for each CD95L-derived shRNA between its associated fold
downregulation in the screen (*20*) and the average seed sequence viability determined in the 4096 siRNA
screen.

In addition, the CD95L shRNAs were split into two groups: 1) 137 shRNAs with an average fold
downregulation (as determined in the shRNA lethality screen) above 5 and 2) a control group with a
matching number of shRNAs whose fold deregulation had an absolute value closest to 0. The average seed
viability (as determined from the siRNA screen) was extracted for the shRNAs in these two groups and
compared using the two-sample, two-tailed Rank Sum test.

Statistical analyses

Continuous data were summarized as means and standard deviations (except for all IncuCyte experiments
where standard errors are shown) and dichotomous data as proportions. Continuous data were compared
using t-tests for two independent groups and one-way ANOVA for 3 or more groups. For evaluation of
continuous outcomes over time, two-way ANOVA was performed using the Stata 14 software with one
factor for the treatment conditions of primary interest and a second factor for time treated as a categorical
variable to allow for non-linearity. Comparisons of single proportions to hypothesized null values were
evaluated using binomial tests. Statistical tests of two independent proportions were used to compare
dichotomous observations across groups. Pearson correlation coefficients (r) and p - values as well as Mann
Whitney U test were calculated using StatPlus (v. 6.3.0.5). Kolmogorov–Smirnov two-sample two-sided test
was used to compare different probability distributions shown in all density plots and eCDF plots. Wilcoxon
rank sum test was used to test for statistical significance in the analysis of the toxicity all miRNA arms in
Fig. 5d.

The effects of treatment vs. control over time were compared for Drosha k.o. and wild-type cells by
fitting regression models that included linear and quadratic terms for value over time, main effects for
treatment and cell type, and two- and three-way interactions for treatment, cell-type and time. The three-way
interaction on the polynomial terms with treatment and cell type was evaluated for statistical significance
since this represents the difference in treatment effects over the course of the experiment for the varying cell
types.

Data availability

RNA sequencing data generated for this study is available in the GEO repository: GSE111379
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE111379, reviewer access token: srileyyktjctduz)
and GSE111363 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE111363, reviewer access token:

udedukauvjexbmp). All 6mer seed toxicity data of the 4096 siRNA screen in HeyA8, H460, M565, and

3LL cells are available in searchable form at https://6merdb.org. All source data are available upon request.

ACKNOWLEDGEMENTS

We are indebted to Dr. Leon Platanias for his generous support of the seed screen and to Denise Scholtens

for help with biostatistics. M.H. was supported by the Intramural Research Program of NIAMS and A.A.S.

by the Swedish Research Council postdoctoral fellowship. This work was funded by training grant

T32CA009560 (to W.P.), R35CA197450 (to M.E.P.), R50CA221848 (to E.T.B.), and RO1 211916 (to EL).

AUTHOR CONTRIBUTIONS

M.E.P., Q.Q.G., W.E.P. and A.E.M. conceived the study. Q.Q.G., W.E.P. performed the majority of the

experiments and data analysis. S.C. performed the 6mer seed screens. A.A.S. performed the Ago pull-down

experiments. E.T.B. performed data analyses. J.M.P designed and implemented the website. M.H. provided

critical conceptual input. M.E.P. designed the study, guided the interpretation of the results and drafted the

manuscript. All authors discussed the results, edited and approved the draft and final versions of the

manuscript.

DECLARATION OF INTEREST

The authors declare no competing interests.

Figures

Fig. 1 A comprehensive screen identifies the most toxic 6mer seeds. **a** Schematic of the siRNA backbone

used in the 4096 seed duplexes toxicity screen. 2'-O-methylation modifications at position 1 and 2 of the

passenger strand were added to prevent passenger strand loading (marked by red Xs) [22]. The nucleotides that

remained constant in all the duplexes are in blue. The variable 6mer seed sequence is in red (Y and

complementary Z) and boxed in blue. **b** *Top:* Results of the 4096 6mer seed duplex screen in human HeyA8

and H460 cells. *Bottom:* Results of the 4096 6mer seed duplex screen in mouse M565 and 3LL cells. Cells

were reverse transfected in triplicates in 384 well plates with 10 nM of individual siRNAs. The cell viability

of each 6mer seed duplex was determined by quantifying cellular ATP content 96 hours after transfection.

All 4096 6mer seeds are ranked by the average effect on cell viability of the two cell lines from the most

toxic (left) to the least toxic (right). Rankings of the 6mer seeds of four previously characterized CD95L

derived siRNAs (siL1, siL2, siL3, and siL4) are highlighted in green. We consider an siRNA highly toxic if

it reduces cell viability 90% or more and moderately toxic if it reduces cell viability 50% or more (black

stippled line). **c** Regression analysis showing the correlation between the 6mer seed toxicity observed in

the human lung cancer cell line H460 (y axis) and the matching 6mer toxicity observed in the human ovarian

cancer cell line HeyA8 (x axis) (left) and average toxicity in the two human cell lines (y axis) and two mouse

cell lines (x axis) (right). p-values were calculated using Pearson correlation analysis. Toxicity of siRNAs

and miRNAs in all four cell lines can be interrogated at https://.6merdb.org. **d** *Left:* Correlation between the

$\log_{10}$ (fold down underrepresentation) of all possible shRNAs that can be derived from the mRNA sequence

of CD95L following their expression from a DOX-inducible lentiviral vector [20] and the toxicity of their seed

sequences as determined in a 4096 arrayed siRNA screen (average of both human cell lines) representing

every seed 6mer permutation [20]. *Right*: Difference in average seed toxicity between the 137 CD95L-derived

shRNAs downregulated at least 5 fold (= toxic) in this screen compared to a size matched group of 137

shRNAs that were the least altered in their abundance in that screen. Pearson correlation coefficient is given

as well as p-value (left) and p-value in analysis on the right was calculated using unpaired two arm ttest.

Fig. 2 The most toxic seeds are G rich. **a** Cell viability of the 19 seed duplexes with the highest content

(>80%) in the 6mer seed region for each nucleotide in two human and two mouse cell lines. p-values

between groups of duplexes were calculated using Student's t-test. **b** Nucleotide composition at each of the 6

seed positions in the top 200 most toxic (left) or the top 200 least toxic (right) seed duplexes in the four cell

lines. siRNAs are considered to be toxic when viability is inhibited >50% (grey stippled line).

Fig. 3 The most toxic G-rich seed containing duplexes preferentially target housekeeping genes enriched

in Cs close to the 3'UTR start site. **a** Nucleotide composition of 20 seeds that are most and least toxic in both

human cell lines (see **Fig. 1b**). **b** eCDF comparing the ratio of occurrences of the 20 most and least toxic

6mer seed matches in the mRNA elements of two sets of expression matched survival genes and nonsurvival

genes. Significance was calculated using a two-sample two-sided Kolmogorov–Smirnov (K-S) test.

Significance was higher in the analysis shown with the first 1000 bases of each 3'UTR than with the entire

3'UTR (data not shown). **c** Probability density plots comparing the nucleotide content between the groups of

expression matched SGs and nonSGs. p-values were calculated using a two-sample two-sided K-S test

comparing the density distribution of SGs and nonSGs. Relevant peak maxima are given. **d** Single nucleotide

frequency distribution in human mRNAs around the boundaries of the 5'UTR and the beginning of the CDS

and the end of the CDS and the beginning of the 3'UTR (shown are 500 bases in each direction). Data are

shown for either all human coding genes (top), a set of 938 SGs or an expression matched set of 938 nonSGs.

Green horizontal bars, area of reduced A/U content in SGs. **e** Distribution of the seed matches to the 20 most

and least toxic 6mer seeds to human cells in human mRNAs around the boundaries of the 5'UTR and the

beginning of the CDS and the end of the CDS and the beginning of the 3'UTR (shown are 500 bases in each

direction). Data are shown for either all genes (top) or the expression matched SGs and nonSGs (center and

bottom). Red horizontal bar, area of enriched toxic seed matches in SGs compared to nonSGs. Green

horizontal bar, area of fewer toxic seed matches in SGs.

Fig. 4 Tumor suppressive miRNAs inhibit cancer cell growth via toxic 6mer seeds. **a** All 4096 6mer

seeds ranked from the lowest average viability (highest toxicity) to the highest viability (lowest toxicity) in

HeyA8 and H460 cells. Locations of 6mer seeds present in major tumor suppressive (red) or tumor

promoting (green) miRNAs are highlighted as individual bars. miRNAs are considered to be toxic when

viability is inhibited >50% (blue stippled line). **b** Percent cell confluence over time of HeyA8 cells

transfected with 5 nM of either the indicated tumor suppressive miRNA precursors or a miRNA precursor

nontargeting control. Data are representative of two independent experiments. Each data point represents

mean ± SE of four replicates. *Two-way ANOVA p-value between cells treated with pre-miR-(NC) and pre-

let-7a-5p is 0.0000. **c** *Left*: Percent cell confluence over time of HeyA8 parental cells transfected with either

pre-miR-34a-5p or si34a-5p^{Seed} and compared to their respective controls (pre-miR (NC) for pre-miR-34a

and siNT2 for si34a^{Seed}) at 10 nM. Data are representative of two independent experiments. Each data point

represents mean ± SE of four replicates. *Right*: % cell death of the same cells harvested 4 days after

transfection. Data are representative of two experiments. Each data point represents mean ± SD of three

replicates. **d** Morphology of HeyA8 cells transfected with either pre-miR-34a-5p or si34a-5p^{Seed} compared to

their respective controls at 10 nM three days after transfection. **e** *Top*: Alignment of the sequences of miR-

34a-5p and miR-34a-5p^{Seed} with the 6mer highlighted. *Bottom:* Comparison of deregulated mRNAs (adjusted

p<0.05, RPM > 1) in HeyA8 cells 48 hrs after transfection with either miR-34a-5p or si34a-5p^{Seed}. Pearson

correlation p-value is given. **f** Overlap of RNAs detected by RNA-Seq downregulated in HeyA8 cells (>1.5

fold) 48 hrs after transfection with either si34a-5p^{Seed} or miR-34a-5p when compared to either siNT2 or a

nontargeting pre-miR, respectively. *Right*: Results of a GOrilla gene ontology analyses of the genes

downregulated in both cells transfected with miR-34a-5p or si34a-5p^{Seed} (top, significance of enrichment <10^{-}

^{11}), or only in cells transfected with miR-34a-5p (bottom, significance of enrichment <10^{-4}). **g** Sylamer plots

for the list of 3'UTRs of mRNAs in cells treated with either miR-34a-5p (top) or si34-5p^{Seed} (bottom) sorted

from down-regulated to up-regulated. The most highly enriched sequence is shown which, in each case, is

the 6mer seed match of the introduced 6mer seed. Bonferroni-adjusted p-values are shown. **h** Gene set

enrichment analysis for a group of 1846 survival genes (top 4 panels) and 416 non-survival genes (bottom 2

panels) [20] after transfecting HeyA8 cells with either miR-34a-5p or si34a-5p^{Seed}. siNT1 and a nontargeting

premiR served as controls, respectively. p-values indicate the significance of enrichment.

Fig. 5 Toxic 6mer seeds and the evolution of cancer regulating miRNAs. **a** Probability density plot of

cell viability of the 6mer seeds of either highly conserved (from humans to zebrafish) or poorly conserved

miRNA seed families (left panel, total number of mature miRNAs = 2164) or of very old (>800 Million

years) miRNAs or very young (<10 Million years) miRNAs (right panel, total number of miRNAs = 299).

For the analysis on the right, miRNA arms with identical sequences (gene duplications) were collapsed and

counted as one arm. Two-sample two-sided K-S test was used to calculate p-values. **b** Change in nucleotide

composition in the 6mer seeds of miRNAs of different ages. **c** Probability density plot of cell viability of the

6mer seeds of mature miRtrons or non-miRtrons. miRNAs with identical sequences (gene duplications) were

collapsed and counted as one seed. Two-sample two-sided K-S test was used to calculate p-value. **d** 780

miRNAs (**Supplementary Table 4**) ranked according to the ratio of viability of the seed (as determined in

the seed screen) of the guide strand and the lesser-expressed arm. Established oncogenic miRNAs are shown

in green, tumor-suppressive miRNAs are shown in red. The guide strand is given for each miRNA (in

parenthesis). p-value of the distribution of oncogenic versus tumor suppressive miRNAs was calculated

using two-sample two-sided K-S test. **e** Cumulative read numbers from the 5p or the 3p arm (according to

miRBase.org) of three oncogenic and three tumor-suppressive miRNAs with the highest (top three) or lowest

(bottom three) ratio of the viability of the guide strand versus the lesser arm. The viability numbers of the

matching 6mer seeds according to the siRNA 6mer seed screen are given. The sequences of the mature 5p or

3p arms are boxed in blue and black, respectively. Toxic seeds are shown in red, non-toxic ones in green.

Fig. 6 Genotoxic drugs cause upregulation of tumor suppressive miRNAs. **a** *Top*: Plot of radiolabeled RNAs pulled down with the Ago proteins from HeyA8 cells treated with doxorubicin (Doxo) for 20, 40, or 80 hours. *Bottom*: Western blot for the pulled down AGO2 of the same samples shown above. The images are representative of two biological duplicates. **b** Fold change of the total reads of Ago bound small RNAs after 20, 40, or 80 hours of Doxo treatment compared to the control sample from Ago-IP sequencing data (Ago bound). Fold change of the total reads of cytosolic small RNAs in HeyA8 cells treated with Doxo for 80 hours compared to the control sample from small RNA-Seq data is given (Total). Data are the

combination of biological duplicates. **c** Pie charts showing the composition of miRNAs bound to Ago proteins after 50 hrs Doxo treatment in HCT116 wild-type (left) or Drosha k.o. cells (right). **d** *Left*, 6mer seed viability (average between HeyA8 and H460 cells) of the Ago-bound miRNAs most up and downregulated in wt or Drosha k.o. cells after Doxo treatment. *Right*, Comparison of the predicted (and most abundant) sequence of miR-320a-3p and Ago-bound sequence of miR-320a-3p and their average viability found most upregulated in Drosha k.o. cells after Doxo treatment. Shown is variance of two biological replicates. Mann-Whiney U test was used to calculate p-value. **e** Percent cell confluence over time of HeyA8 cells transfected with 5 nM of controls, pre-miR-320a or an siRNA duplex that corresponds to the shifted form of miR-320a-3p (si-miR-320a-3p^Ago) that was found to be upregulated and bound to Ago proteins upon Doxo treatment. Data are representative of two independent experiments. Each data point represents mean ± SE of four replicates. **f** Viability changes (ATP content) in four cell lines 48 hrs after transfection with Lipid only, 10 nM of siNT1, siL3, a nontargeting pre-miR, or miR-320a-3p^Ago - the only shared upregulated miRNA in HeyA8 cells, HCT116 wild-type, and HCT116 Drosha k.o. cells - after Doxo treatment. p-values were determined using Student's t-test. *** p<0.0001. Samples were performed in triplicate (siNT1, siL3), 6 repeats (miR-320a-3p^Ago) and 8 repeats (Lipid).

REFERENCES

1. Lee Y, *et al*. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* **23**, 4051-4060 (2004).

2. Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN. The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* **18**, 3016-3027 (2004).

3. Yi R, Qin Y, Macara IG, Cullen BR. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* **17**, 3011-3016 (2003).

4. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**, 363-366 (2001).

5. Hutvagner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD. A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* **293**, 834-838 (2001).

6. Wang Y, Sheng G, Juranek S, Tuschl T, Patel DJ. Structure of the guide-strand-containing argonaute silencing complex. *Nature* **456**, 209-213 (2008).

7. Leuschner PJ, Ameres SL, Kueng S, Martinez J. Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep* **7**, 314-320 (2006).

8. Schirle NT, MacRae IJ. The crystal structure of human Argonaute2. *Science* **336**, 1037-1040 (2012).

884 9.   Eulalio A, Huntzinger E, Izaurralde E. GW182 interaction with Argonaute is essential for miRNA-
885   mediated translational repression and mRNA decay. *Nat Struct Mol Biol* **15**, 346-353 (2008).
886

887 10. Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB. Prediction of mammalian microRNA
888   targets. *Cell* **115**, 787-798 (2003).
889

890 11. Lai EC. Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-
891   transcriptional regulation. *Nat Genet* **30**, 363-364 (2002).
892

893 12. Selbach M, Schwanhausser B, Thierfelder N, Fang Z, Khanin R, Rajewsky N. Widespread changes in
894   protein synthesis induced by microRNAs. *Nature* **455**, 58-63 (2008).
895

896 13. Baek D, Villen J, Shin C, Camargo FD, Gygi SP, Bartel DP. The impact of microRNAs on protein
897   output. *Nature* **455**, 64-71 (2008).
898

899 14. Esquela-Kerscher A, Slack FJ. Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer* **6**, 259-
900   269 (2006).
901

902 15. Balatti V, Pekarky Y, Rizzotto L, Croce CM. miR deregulation in CLL. *Adv Exp Med Biol* **792**, 309-
903   325 (2013).
904

905 16. Slabakova E, Culig Z, Remsik J, Soucek K. Alternative mechanisms of miR-34a regulation in cancer.
906   *Cell Death Dis* **8**, e3100 (2017).
907

908 17. Hua YJ, Larsen N, Kalyana-Sundaram S, Kjems J, Chinnaiyan AM, Peter ME. miRConnect 2.0:
909   Identification of antagonistic, oncogenic miRNA families in three human cancers. *BMC Genomics* **14**,
910   179 (2013).
911

912 18. Concepcion CP, Bonetti C, Ventura A. The microRNA-17-92 family of microRNA clusters in
913   development and disease. *Cancer journal* **18**, 262-267 (2012).
914

915 19. He X, He L, Hannon GJ. The guardian's little helper: microRNAs in the p53 tumor suppressor
916   network. *Cancer Res* **67**, 11099-11101 (2007).
917

918 20. Putzbach W, *et al*. Many si/shRNAs can kill cancer cells by targeting multiple survival genes through
919   an off-target mechanism. *eLife* **6**, e29702 (2017).
920

921 21. Murmann AE, *et al*. Induction of DISE in ovarian cancer cells in vivo. *Oncotarget* **8**, 84643-84658
922   (2017).
923

924 22. Murmann AE, *et al*. Small interfering RNAs based on huntingtin trinucleotide repeats are highly
925   toxic to cancer cells. *EMBO Rep* **19**, e45336 (2018).
926

927 23. Putzbach W, *et al*. CD95L mRNA is toxic to cells *bioRxive*, https://doi.org/10.1101/330324 (2018).
928

929 24. Hadji A, *et al*. Death induced by CD95 or CD95 ligand elimination. *Cell Reports* **10**, 208-222 (2014).
930

931 25. Bramsen JB, *et al*. A large-scale chemical modification screen identifies design rules to generate
932   siRNAs with high activity, high stability and low toxicity. *Nucleic Acids Res* **37**, 2867-2881 (2009).
933

934   26.   Peter ME. Let-7 and miR-200 microRNAs: guardians against pluripotency and cancer progression.
935   *Cell Cycle* **8**, 843-852 (2009).
936
937   27.   Agostini M, Knight RA. miR-34: from bench to bedside. *Oncotarget* **5**, 872-881 (2014).
938
939   28.   Patel M, Peter ME. Identification of DISE-inducing shRNAs by monitoring cellular responses. *Cell*
940   *Cycle* **17**, 506-514 (2018).
941
942   29.   van Dongen S, Abreu-Goodger C, Enright AJ. Detecting microRNA binding and siRNA off-target
943   effects from expression data. *Nat Methods* **5**, 1023-1025 (2008).
944
945   30.   Grimson A, Farh KK, Johnston WK, Garrett-Engele P, Lim LP, Bartel DP. MicroRNA targeting
946   specificity in mammals: determinants beyond seed pairing. *Mol Cell* **27**, 91-105 (2007).
947
948   31.   Patel VD, Capra JA. Ancient human miRNAs are more likely to have broad functions and disease
949   associations than young miRNAs. *BMC Genomics* **18**, 672 (2017).
950
951   32.   Berezikov E, Chung WJ, Willis J, Cuppen E, Lai EC. Mammalian mirtron genes. *Mol Cell* **28**, 328-
952   336 (2007).
953
954   33.   Meijer HA, Smith EM, Bushell M. Regulation of miRNA strand selection: follow the leader?
955   *Biochem Soc Trans* **42**, 1135-1140 (2014).
956
957   34.   Concepcion CP, *et al*. Intact p53-dependent responses in miR-34-deficient mice. *PLoS Genet* **8**,
958   e1002797 (2012).
959
960   35.   Meredith AM, Dass CR. Increasing role of the cancer chemotherapeutic doxorubicin in cellular
961   metabolism. *J Pharm Pharmacol* **68**, 729-741 (2016).
962
963   36.   Yeung TK, Germond C, Chen X, Wang Z. The mode of action of taxol: apoptosis at low
964   concentration and necrosis at high concentration. *Biochem Biophys Res Commun* **263**, 398-404
965   (1999).
966
967   37.   Yoo SH, *et al*. Etoposide induces a mixed type of programmed cell death and overcomes the
968   resistance conferred by Bcl-2 in Hep3B hepatoma cells. *Int J Oncol* **41**, 1443-1454 (2012).
969
970   38.   Kwon HK, *et al*. Etoposide Induces Necrosis Through p53-Mediated Antiapoptosis in Human Kidney
971   Proximal Tubule Cells. *Toxicol Sci* **148**, 204-219 (2015).
972
973   39.   Wang D, Lippard SJ. Cellular processing of platinum anticancer drugs. *Nat Rev Drug Discov* **4**, 307-
974   320 (2005).
975
976   40.   Chang BD, *et al*. A senescence-like phenotype distinguishes tumor cells that undergo terminal
977   proliferation arrest after exposure to anticancer agents. *Cancer Res* **59**, 3761-3767 (1999).
978
979   41.   Eom YW, *et al*. Two distinct modes of cell death induced by doxorubicin: apoptosis and cell death
980   through mitotic catastrophe accompanied by senescence-like phenotype. *Oncogene* **24**, 4765-4777
981   (2005).
982

983   42.   Hauptmann J, *et al*. Biochemical isolation of Argonaute protein complexes by Ago-APP. *Proc*
984   *Natl Acad Sci U S A* **112**, 11841-11845 (2015).
985

986   43.   Kim YK, Kim B, Kim VN. Re-evaluation of the roles of DROSHA, Export in 5, and DICER in
987   microRNA biogenesis. *Proc Natl Acad Sci U S A* **113**, E1881-1889 (2016).
988

989   44.   Walz AL*, et al*. Recurrent DGCR8, DROSHA, and SIX homeodomain mutations in favorable
990   histology Wilms tumors. *Cancer Cell* **27**, 286-297 (2015).
991

992   45.   Kumar MS, Lu J, Mercer KL, Golub TR, Jacks T. Impaired microRNA processing enhances cellular
993   transformation and tumorigenesis. *Nat Genet* **39**, 673-677 (2007).
994

995   46.   Chandradoss SD, Schirle NT, Szczepaniak M, MacRae IJ, Joo C. A Dynamic Search Process
996   Underlies MicroRNA Targeting. *Cell* **162**, 96-107 (2015).
997

998   47.   Di Martino MT, *et al*. Synthetic miR-34a mimics as a novel therapeutic agent for multiple myeloma:
999   in vitro and in vivo evidence. *Clin Cancer Res* **18**, 6260-6270 (2012).
1000

1001   48.   Beg MS, *et al*. Phase I study of MRX34, a liposomal miR-34a mimic, administered twice weekly in
1002   patients with advanced solid tumors. *Invest New Drugs* **35**, 180-188 (2017).
1003

1004   49.   Cao W, *et al*. miR-34a regulates cisplatin-induce gastric cancer cell death by modulating
1005   PI3K/AKT/survivin pathway. *Tumour Biol* **35**, 1287-1295 (2014).
1006

1007   50.   Navarro F, Lieberman J. miR-34 and p53: New Insights into a Complex Functional Relationship.
1008   *PLoS One* **10**, e0132767 (2015).
1009

1010   51.   Alfano L, Costa C, Caporaso A, Antonini D, Giordano A, Pentimalli F. HUR protects NONO from
1011   degradation by mir320, which is induced by p53 upon UV irradiation. *Oncotarget* **7**, 78127-78139
1012   (2016).
1013

1014   52.   Stark A, Brennecke J, Bushati N, Russell RB, Cohen SM. Animal MicroRNAs confer robustness to
1015   gene expression and have a significant impact on 3'UTR evolution. *Cell* **123**, 1133-1146 (2005).
1016

1017   53.   Zare H, Khodursky A, Sartorelli V. An evolutionarily biased distribution of miRNA sites toward
1018   regulatory genes with high promoter-driven intrinsic transcriptional noise. *BMC Evol Biol* **14**, 74
1019   (2014).
1020

1021   54.   Ceppi P*, et al*. CD95 and CD95L promote and protect cancer stem cells. *Nature Commun* **5**, 5238
1022   (2014).
1023

1024   55.   Hafner M, Renwick N, Farazi TA, Mihailovic A, Pena JT, Tuschl T. Barcoded cDNA library
1025   preparation for small RNA profiling by next-generation sequencing. *Methods* **58**, 164-170 (2012).
1026

1027   56.   Panwar B, Omenn GS, Guan Y. miRmine: a database of human miRNA expression profiles.
1028   *Bioinformatics* **33**, 1554-1560 (2017).
1029

1030   57.   Rorbach G, Unold O, Konopka BM. Distinguishing mirtrons from canonical miRNAs with data
1031   exploration and machine learning methods. *Sci Rep* **8**, 7560 (2018).
1032

1033
1034