**Auxin is not asymmetrically distributed in initiating Arabidopsis leaves**

**Neha Bhatia[1] and Marcus G. Heisler[1*]**

**Affiliations**

[1]School of Life and Environmental Sciences, University of Sydney, NSW, Australia

*Author for correspondence: Marcus G Heisler (marcus.heisler@sydney.edu.au)

**Abstract**

It has been proposed that asymmetric auxin levels within initiating leaves help

establish leaf polarity, based in part on observations of the DII auxin sensor. Here we

show that the mDII control sensor also exhibits an asymmetry and that according to

the ratio-metric auxin sensor R2D2, no obvious asymmetry in auxin exists. Together

with other recent findings, our results argue against the importance of auxin

asymmetry in establishing leaf polarity.

**Results and Discussion**

The leaves of seed plants are usually flat with distinct cell types making up

their dorsal (upper) and ventral (lower) tissues. A fundamental question in plant

development is how this dorsal-ventral polarity is first specified. Studies based on

wounding experiments have suggested the presence of an inductive signal originating

from the meristem that promotes dorsal identity in the adaxial (adjacent to the

meristem) tissues of the leaf primordium[1,2]. However more recently, it was found that

exogenous application of the plant hormone auxin to tomato leaf primordia resulted in

the formation of radialized leaves that appeared ventrilized. Hence, relatively high

levels of auxin are proposed to promote ventral or inhibit dorsal leaf cell fate[3].

Extending these conclusions, it was also proposed that an asymmetry in the auxin

distribution across the leaf adaxial-abaxial axis at leaf initiation acts to help specify

leaf polarity during regular development[3]. A critical piece of evidence supporting this

latter proposal is that an auxin sensor, the DII[4,5] indicates low levels of auxin in

adaxial leaf tissues compared to abaxial tissues at leaf initiation[3]. Hence asymmetries

in auxin concentrations between the adaxial and abaxial leaf tissues, as a result of

PIN1 mediated auxin transport, are proposed to help establish leaf polarity[3]. Building

on this conclusion, a more recent study proposed that low levels of auxin in adaxial

tissues are necessary to restrict the expression of the WOX1 and PRS genes to the

middle domain, since auxin promotes their expression[6]. Finally, the reported

asymmetry in auxin has also been linked to asymmetries in the mechanical properties

of leaf tissues and their morphogenesis[7].

Despite these studies, reason to reassess the role of any asymmetry in auxin in

specifying leaf polarity has arisen due to the recent finding that the expression

patterns of genes involved in leaf dorsoventrality are already patterned in meristem

tissues where leaves originate, with little change occurring to their expression during

organ initiation. Hence leaf polarity appears pre-patterned [8]. Furthermore, rather than

compromise dorsal cell fate, auxin was found to promote dorsal cell fate by

maintaining Class III HD-ZIP expression and repressing KAN1 expression in the

adaxial cells of organ primordia[8], which is the opposite of what would be expected

according to the auxin asymmetry hypothesis. In addition, Caggiano et al. (2017) also

found that exogenous auxin application to young leaf primordia did not influence the

spatial pattern of WOX1 and PRS, only the intensity of expression[8]. This also

conflicts with the proposal that low levels of auxin are required in adaxial leaf tissues

to prevent ectopic WOX1 and PRS expression as previously proposed[6].

Given the concerns listed above, we decided to examine the distribution of auxin

during leaf initiation using not only by the DII sensor assessed in previous studies but

also, by looking more closely at the expression pattern of the non-auxin degradable

mDII auxin sensor control[4] as well as the ratiometric R2D2 auxin sensor[9]. We

initially focused on the first two leaves three and four days after stratification (DAS),

when these leaves are initiating as well as a day later as they begin to elongate and

included the PIN1-GFP marker in our analysis to correlate auxin levels with PIN1

expression and polarity. At 3DAS, the expression patterns of REV and KAN1 are

already polar within such primordia, although the FIL expression domain at this stage

of development is still being refined[8]. Also at this stage, PIN1 is polarized towards the

distal tip of leaf primordia but has reversed polarity away from the primordia, towards

the meristem, in cells adjacent to the primordia on the adaxial side. According to the

ratio-metric auxin sensor R2D2, auxin concentrations were relatively low in adaxial

cells of the primordia but also low in abaxial and lateral regions proximally (Fig1-a

and d; Fig.S1). High levels of auxin were only found in more distal regions towards

the tip of the primordia, matching the overall pattern of signal from PIN1-GFP. No

obvious asymmetry in signal between the adaxial and abaxial sides of the primordia

was observed. This same overall pattern of signal was found in 16 out of 18 leaves

that were examined. At 4DAS, the auxin distribution according to the R2D2 sensor

was more uniform although higher levels of auxin appeared associated with the

vasculature, again correlating with PIN1 expression (n=12/12 leaves) (Fig. 1 g and j).

As our results using the R2D2 auxin sensor indicate a different auxin distribution

compared to that reported previously using the DII marker [3], we next re-examined the

pattern of DII auxin sensor expression at the same developmental stages. In contrast

to the R2D2 pattern, the DII pattern showed an asymmetry of expression in leaf

primordia at 3 DAS, indicating relatively low auxin levels in adaxial primordium

cells, as found previously. DII signal appeared strongest in the adaxial epidermis but

was also stronger in the adaxial sub-epidermal cell layer compared to abaxial

epidermal and sub-epidermal cell layers (Fig.1- b, e, h and Fig. S2) (n=18/18 leaves).

One day later, although DII signal was still higher in the adaxial cells of the

primordia, it started to show a relatively increased expression in the abaxial sub-

epidermal layers as well compared to earlier stage (n=13/14 leaves) (Fig1-k). Overall

our results using the DII marker are similar to those obtained previously and

consistent with the proposal that there are low levels of auxin in the adaxial regions of

leaf primordia, in contrast to our results using the R2D2 sensor. Given this

discrepancy, we next examined the expression of the mDII sensor which is driven by

the same 35S promoter as the DII sensor but is not auxin sensitive. Surprisingly we

found that, like the DII results, expression of the mDII marker was also higher in the

adaxial cells of the primordia (n= 16/16 leaves) (Fig.1-c,f,i and Fig.S3). The pattern

appeared almost identical to the pattern found using the DII marker except that the

mDII marker also showed high levels of expression in the shoot meristem whereas the

DII sensor did not (compare Fig1-b,c; Fig S2 a-h and Fig S3 a-f ). The similarity of

expression between DII and mDII was also apparent at 4DAS when the leaves had

started to elongate (Fig.1-l) (n=14/14 leaves). To verify the auxin sensitivity of the

sensors used we imaged seedlings before and after treatment with 5mM NAA and

found a strong decrease in DII expression compared to mDII and an increase in the

ratio of VENUS compared to tdTomato signal for the R2D2 sensor, consistent with an

increase in auxin levels (Figure S4).

All together these results indicate that the asymmetry in expression found previously

for the DII auxin sensor in very young leaf primordia[3] is not due to an asymmetry in

auxin levels but rather, likely due to differences in transcription driven by the 35S

promoter used to drive both DII and mDII in adaxial compared to abaxial leaf tissues.

We note that although a single section showing control expression of mDII in older

leaves was cited by Guan et al., (2017)[10], our results show that this information was

not adequate for assessing similarities and differences between DII and mDII at early

developmental stages. To check whether adaxial expression in leaf primordia is a

characteristic common to other reporters driven by the 35S promoter, we also

examined the expression of 35S::H2B-mRFP1 and 35S::EGFP-LTI6b which had

previously been combined into the same plant line by crossing[11]. Surprisingly the

expression patterns for these markers were not only different to the mDII marker but

also different to each other with the H2B marker showing stronger expression in the

adaxial and abaxial epidermis and the LTI6b marker showing a more uniform pattern

(Fig. S5). These results reveal that even when the same promoter is utilized to drive

FP (fluorescent protein) expression, distinct differences in intensity patterns can occur

*in planta*. This may be due to differences in the position of T-DNA insertion in the

genome or differences in the surrounding DNA of the vector used. All together then,

our results highlight the importance of using ratio-metric sensors for *in vivo*

measurements since otherwise it is difficult to adequately control for differences in

signal intensity due to promoter activity or other confounding factors. Finally, while

the R2D2 marker utilizes a single transgene to express two distinct FPs from two

identical promoters[9], a potentially improved approach to assess auxin levels *in vivo*

may be to utilize a single promoter to drive the auxin-dependent degradation domain

II from Aux/IAA proteins linked to a tandem fusion of fast and slowly folding FP

variants, such as VENUS and mCherry. Such an approach can enable a more direct

readout of protein degradation rates[12,13] and therefore potentially improve

measurements of relative auxin concentration.

While our findings are inconsistent with the proposal that asymmetries in the auxin

distribution influence leaf polarity[3], pattern WOX/PRS gene expression [6] or regulate

tissue mechanics[7], they also leave other observations unexplained. In particular, to

understand the apparent ventrilization of leaf primordia in tomato in response to

exogenous auxin[3] will require further work in assessing how auxin distribution

patterns change in response to exogenous treatments and how auxin is distributed

during regular development in tomato, preferentially using a ratio-metric auxin sensor

such as R2D2.

**Methods**

**Plant material and growth conditions.**

Seeds of the plants expressing *p35S::DII-VENUS* and *p35S::mDII-VENUS*

transgenes (*Columbia* ecotype) were obtained from Dr. Teva Vernoux [4]. An

independent batch of seeds expressing *p35S::mDII-VENUS* transgenes (*Columbia*

ecotype) was also obtained from NASC (Arabidopsis stock center, NASC ID:

N799174) for analysis. R2D2 reporter line carrying *pPIN1::PIN1GFP* transgene

(*Landsberg* ecotype) has been described previously[9,14]. Seeds were germinated and

grown on GM medium (pH-7 with 1M KOH) containing 1% sucrose, 1X Murashige

and Skoog salts (Sigma M5524), MES 2-(MN-morpholino)- ethane sulfonic acid

(Sigma M2933), 0.8 % Bacto Agar (Difco), 1% MS vitamins (Sigma M3900) in long

day conditions.

**Confocal imaging and data analysis**

Seedlings aged 3DAS (days after stratification) and 4DAS were dissected as

described previously[14]. Dissected seedlings were then oriented appropriately to obtain

a view of the young leaves either from above or from the side. Seedlings were imaged

live, on a Leica TCS-SP5 upright confocal laser scanning microscope with hybrid

detectors (HyDs) using a 25X water objective (N.A 0.95). VENUS was imaged using

argon laser (excitation wavelength 514nm) while tdTomato was imaged using a white

light laser (excitation wavelength 561 nm). Z-stacks were acquired in a 512x512 pixel

format with section spacing of 1μm and line averaging 2.

Ratio-metric calculations for R2D2 auxin sensor were performed using ImageJ (FIJI,

https://fiji.sc) as described previously[14]. All the images were processed using IMARIS

9.0.0 (bit-plane). Optical sections (transverse or longitudinal) were reconstructed

using orthogonal slicer in IMARIS.

**Auxin treatment**

Seedlings aged 3DAS were dissected imaged and treated with approximately 10μL of 5mM NAA (1-Napthaleneacetic acid) solution in water (0.5M stock in 1M KOH) for 60 minutes and imaged again with same settings as prior to treatment.

**Acknowledgements**

We thank Associate Professor Mary Byrne for critical feedback on the manuscript. We thank Dr. Teva Vernoux for kindly providing the seeds of plants expressing *p35S::DII-VENUS* and *p35S::mDII-VENUS* transgenes. We also thank Professor Jim Haseloff for kindly providing the seeds of plants expressing *p35S::H2B-mRFP1* and *p35S::EGFP-LTI6b* transgenes.

**References**

Sussex, I. M. Morphogenesis in Solanum tuberosum I: experimental investigation of leaf dorsiventrality and orientation in the juvenile shoot. *Phytomorphology* **5**, 286-300 (1955).

Reinhardt, D., Frenz, M., Mandel, T. & Kuhlemeier, C. Microsurgical and laser ablation analysis of leaf positioning and dorsoventral patterning in tomato. *Development* **132**, 15-26 (2005).

Qi, J. *et al.* Auxin depletion from leaf primordia contributes to organ patterning. *Proc Natl Acad Sci U S A* **111**, 18769-18774, doi:10.1073/pnas.1421878112 (2014).

Brunoud, G. *et al.* A novel sensor to map auxin response and distribution at high spatio-temporal resolution. *Nature* **482**, 103-106, doi:10.1038/nature10791 (2012).

Vernoux, T. *et al.* The auxin signalling network translates dynamic input into robust patterning at the shoot apex. *Mol Syst Biol* **7**, 508, doi:10.1038/msb.2011.39 (2011).

Guan, C. *et al.* Spatial Auxin Signaling Controls Leaf Flattening in Arabidopsis. *Curr Biol* **27**, 2940-2950 e2944, doi:10.1016/j.cub.2017.08.042 (2017).

Qi, J. *et al.* Mechanical regulation of organ asymmetry in leaves. *Nat Plants* **3**, 724-733, doi:10.1038/s41477-017-0008-6 (2017).

Caggiano, M. P. *et al.* Cell type boundaries organize plant development. *eLife* **6**, doi:10.7554/eLife.27421 (2017).

Liao, C. Y. *et al.* Reporters for sensitive and quantitative measurement of auxin response. *Nat Methods* **12**, 207-210, 202 p following 210, doi:10.1038/nmeth.3279 (2015).

Wang, Y. *et al.* The Stem Cell Niche in Leaf Axils Is Established by Auxin and Cytokinin in Arabidopsis. *Plant Cell* **26**, 2055-2067, doi:10.1105/tpc.114.123083 (2014).

11   Federici, F., Dupuy, L., Laplaze, L., Heisler, M. & Haseloff, J. Integrated
genetic and computation methods for in planta cytometry. *Nature*
*Methods* **9**, 483-U104, doi:10.1038/nmeth.1940 (2012).
12   Khmelinskii, A. *et al.* Tandem fluorescent protein timers for in vivo
analysis of protein dynamics. *Nat Biotechnol* **30**, 708-714,
doi:10.1038/nbt.2281 (2012).
13   Barry, J. D., Dona, E., Gilmour, D. & Huber, W. TimerQuant: a modelling
approach to tandem fluorescent timer design and data interpretation for
measuring protein turnover in embryos. *Development* **143**, 174-179,
doi:10.1242/dev.125971 (2016).
14   Bhatia, N. *et al.* Auxin Acts through MONOPTEROS to Regulate Plant Cell
Polarity   and   Pattern   Phyllotaxis.   *Curr   Biol*   **26**,   3202-3208,
doi:10.1016/j.cub.2016.09.044 (2016).

**Figure 1 Predicted auxin distribution in young leaves as indicated by different auxin sensors.**

**(a-c)** Confocal projections of *Arabidopsis* seedlings aged 3DAS (days after stratification) showing predicted auxin distribution based on the ratio-metric auxin sensor R2D2 (magenta) along with PIN1-GFP (green) (a); expression pattern of *p35S::DII-VENUS* (magenta) (b) and *p35S::mutatedDII-VENUS* (mDII, magenta) (c). **(d-f)** Longitudinal reconstructed optical sections of (a-c), respectively, along the dashed lines. **(g-i)** Representative examples of transverse reconstructed optical sections of 3DAS Arabidopsis seedlings showing predicted auxin distribution as indicated by the R2D2 sensor along with PIN1-GFP expression (g), DII-VENUS expression (h) and mDII-VENUS expression (i). R2D2 auxin sensor indicates low but symmetric auxin levels on adaxial and abaxial sides and relatively high auxin levels at the distal tip and in the provasculature where PIN1 expression is also high (d, g). DII-

VENUS is more strongly expressed adaxially indicating low auxin levels on the

adaxial side of the leaves relative to the abaxial side. However, mDII-VENUS also

shows high expression on the adaxial side of the leaf (compare e, h with f, i) and in

the shoot meristem. **(j-l)** Representative examples of longitudinal reconstructed

optical sections of 4DAS Arabidopsis seedlings showing predicted auxin distribution

by R2D2 sensor along with PIN1-GFP expression (green) (i), DII-VENUS expression

(j) and mDII-VENUS expression (k). At this stage, the R2D2 sensor indicates a more

uniform auxin distribution in the leaf but higher auxin levels in the vasculature where

PIN1-GFP expression is also high (j). DII-VENUS shows slightly high expression in

adaxial cells and an absence of expression in the vasculature (k). mDII-VENUS

shows a similar pattern to DII but is also expressed in the vasculature (l). Scale bars

15μm (a-i) and 20μm (j-l).

FigureS1

**FigureS1 Additional examples of R2D2, ratio-based predictions of auxin**
**distribution in 3DAS old Arabidopsis seedlings. (a-g)** Confocal projections of
*Arabidopsis* seedlings aged 3DAS (days after stratification) showing predicted auxin
distribution based on the ratio-metric auxin sensor R2D2 (magenta) along with PIN1-
GFP (green). **(h-n)** Median transverse optical sections of (a-g). **(o-u)** Median
longitudinal optical sections of (a-g). Scale bars 20μm (a-u).

Figure S2

**FigureS1 Additional examples *p35S::DII-3xVENUS-N7* expression in 3DAS old
Arabidopsis seedlings. (a-g)** Confocal projections of *Arabidopsis* seedlings aged
3DAS (days after stratification) showing expression pattern of p35s::DII-3xVENUS-

270 N7 sensor (magenta). **(h-n)** Median transverse optical sections of (a-g). **(o-u)** Median

271 longitudinal optical sections of (a-g). Scale bars 20μm (a-u).

**FigureS1 Additional examples of *p35S::mDII-3xVENUS-N7* expression in 3DAS old Arabidopsis seedlings. (a-f)** Confocal projections of *Arabidopsis* seedlings aged 3DAS (days after stratification) showing expression pattern of p35s::mDII-3xVENUS-N7 sensor (magenta). Note that plants obtained from two different sources

278  showed similar expression pattern.  **(h-l)** Median transverse optical sections of (a-f).

279  **(o-r)** Median longitudinal optical sections of (a-g). Scale bars 20μm (a-r).

FigureS4

**Figure S4 Response of different auxin sensors to external auxin application**

**(a-f)** Confocal projections (a-c) and transverse optical reconstructions (d-f) of
*Arabidopsis* seedlings, 3DAS, showing predicted auxin distribution based on R2D2
ratio-metric sensor (magenta) along with PIN1-GFP expression in green (a,d), DII-
VENUS sensor (magenta) (b,e) and mDII-VENUS sensor (magenta) (c, f) before
auxin application. **(g-l)** Confocal projections (g-i) and transverse optical
reconstructions (j-l) of *Arabidopsis* seedlings, 3DAS, showing predicted auxin
distribution based on R2D2 ratio-metric sensor (magenta) along with PIN1-GFP
expression in green (a,d), DII-VENUS sensor (magenta) (b,e) and mDII-VENUS
sensor (magenta) (c, f) 1hour after the application of 5mM NAA. Note, R2D2 sensor
indicates an increased and broadening of auxin levels after NAA application (compare
d and j) (n=3/3). Consistently, DII-VENUS shows an attenuated expression within 1
hour of auxin application (compare b,e with h,j) (n=4/4). mDII-VENUS levels do not
decrease upon auxin application (compare c,f with i,l) (n=4/4). Scale bars 20μm (a-l).

Figure S5

**Figure S5 Different 35S promoter-driven reporter genes show different**
**expression patterns in young leaves.**

**(a-c)** Confocal projections of *Arabidopsis* seedlings aged 3DAS  showing expression
patterns of a *p35S:eGFP-Lti6b* (membrane marker, green) and *p35S::H2B-*
*mRFP1*(nuclear marker red) together (a), *p35S::H2B-mRFP1* only (magenta) (b) and
*p35S:eGFP-Lti6b* only (magenta) (c). **(d-f)** Transverse optical reconstructions of (a-c)
respectively; merged (d), *p35S::H2B-mRFP1* only (magenta) (e) and *p35S:eGFP-*
*Lti6b* only (magenta) (f). **(g-i)** Longitudinal optical reconstruction of  (a-c)
respectively; merged (g), *p35S::H2B-mRFP1* only (magenta) (h) and *p35S:eGFP-*
*Lti6b* only (magenta) (i). Note *p35S::H2B-mRFP1* shows similarly high levels of
expression in both adaxial (ad) as well as abaxial (ab) epidermal cell layers and
weaker expression in the middle regions (e and h). However, *p35S:eGFP-Lti6b* shows
no consistent asymmetry in expression throughout the leaf (f and i). Scale bars 20μm
(a-i), (n=10 leaves).