Synthetic standards combined with error and bias correction improves the accuracy and quantitative resolution of antibody repertoire sequencing in human naïve and memory B cells

Simon Friedensohn*[1], John M. Lindner*[2], Vanessa Cornacchione[2], Mariavittoria Iazeolla[2], Enkelejda Miho[1,3], Andreas Zingg[1], Simon Meng[1], Elisabetta Traggiai[2], and Sai T. Reddy[1]

[1]*Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland*
[2]*Novartis Institutes for BioMedical Research, Basel, Switzerland*
[3]*aiNET GmbH, c/o ETH Zürich, Basel, Switzerland*

*equal contribution

Correspondence: elisabetta.traggiai@novartis.com and sai.reddy@ethz.ch

ABSTRACT

High-throughput sequencing of immunoglobulin repertoires (Ig-seq) is a powerful method for quantitatively interrogating B cell receptor sequence diversity. When applied to human repertoires, Ig-seq provides insight into fundamental immunological questions, and can be implemented in diagnostic and drug discovery projects. However, a major challenge in Ig-seq is ensuring accuracy, as library preparation protocols and sequencing platforms can introduce substantial errors and bias that compromise immunological interpretation. Here, we have established an approach for performing highly accurate human Ig-seq by combining synthetic standards with a comprehensive error and bias correction pipeline. First, we designed a set of 85 synthetic antibody heavy chain standards (*in vitro* transcribed RNA) to assess correction workflow fidelity. Next, we adapted a library preparation protocol that incorporates unique molecular identifiers (UIDs) for error and bias correction which, when applied to the synthetic standards, resulted in highly accurate data. Finally, we performed Ig-seq on purified human circulating B cell subsets (naïve and memory), combined with a cellular replicate sampling strategy. This strategy enabled robust and reliable estimation of key repertoire features such as clonotype diversity, germline segment and isotype subclass usage, and somatic hypermutation (SHM). We anticipate that our standards and error and bias correction pipeline will become a valuable tool for researchers to validate and improve accuracy in human Ig-seq studies, thus leading to potentially new insights and applications in human antibody repertoire profiling.

INTRODUCTION

Adaptive immune responses are governed by cooperative interactions between B and T lymphocytes upon antigen recognition. A hallmark of these cells is the somatic generation of clonally unique antigen receptors during primary lymphocyte differentiation. In particular, B cell antigen receptors (BCRs, and their analogous secreted form, antibodies) result from rearrangement of the germline-encoded variable (V), diversity (D, heavy chain only), and joining (J) gene segments. V(D)J recombination in B cells creates a highly complex receptor population (generally interchangeably referred to as BCR, antibody, or immunoglobulin (Ig) repertoires), which matures upon antigen experience to produce the more targeted, high-affinity memory BCR network. In-

depth and accurate characterization of these repertoires provides valuable insight into the generation and maintenance of immunocompetency, which can be used to monitor changes in immune status, and to identify potentially reactive clones for therapeutic or other uses. Due to rapid technological advances, high-throughput sequencing of Ig genes (Ig-seq) has become a major approach to catalog the diversity of antibody repertoires (1-3). Ig-seq applied to human B cells has potential in a variety of applications (4), particularly in antibody drug discovery (5-7), diagnostic profiling for vaccines (8, 9), and biomarker-based disease detection (10, 11). Additionally, Ig-seq is enabling a more comprehensive understanding of basic human immunobiology, such as B cell clonal distribution across physiological compartments in health and disease (12, 13).

A major challenge in Ig-seq is the requirement of accurate and high-quality datasets. Several current library preparation protocols are based on target enrichment from genomic DNA or mRNA (14). For example, the conversion of mRNA (more commonly used due to transcript abundance and isotype splicing) into antibody sequencing libraries relies on a number of molecular reagents and amplification steps (e.g. reverse transcriptase, multiplex primer sets, PCR), which potentially introduce errors and bias. Due to the highly polymorphic nature of repertoires especially from affinity-matured memory B cells and plasma cells, it becomes essential to determine if such technical noise occurs at non-negligible rates, as this could alter quantitation of critical repertoire features such as clonal frequencies, germline gene usage, and somatic hypermutation (SHM) (14, 15). One way to address this is through the use of synthetic control standards, for which the sequence and abundance is known prior to sequencing, thus providing a means to assess quality and accuracy (16). Several examples of standards have been presented for Ig-seq; Shugay et al. sequenced libraries prepared from a small polyclonal pool of B and T lymphocyte cell lines, and observed nearly 5% erroneous reads, resulting in approximately 100 false-positive variants per clone (17). Recently, Khan et al. developed a set of synthetic RNA (*in vitro* transcribed) spike-in standards based on mouse antibody sequences, which were used to show that a substantial amount of errors and bias are introduced during multiplex-PCR library preparation and sequencing (18).

Various experimental and computational workflows exist to mitigate the effects of errors and bias in Ig-seq. One of the most advanced and powerful strategies is to prepare libraries with the incorporation of random and unique molecular identifiers (UIDs, also commonly referred to as UMIs or molecular barcodes). Following sequencing, error correction can be performed by clustering and consensus building of reads that share the same UID; reads sharing the same UID are assumed to be derived from the same original mRNA/cDNA molecule (19). Furthermore, bias correction for cDNA abundance can be performed by counting the number of UIDs (instead of total reads) (20, 21). Several iterations of UID-tagging have been developed for Ig-seq, such as UID labeling during first- and second-strand cDNA synthesis (22), UID addition during RT template switching (23), and so-called "tagmentation" of UID-labelled amplicons (24). Recently, we developed an innovative strategy to add UIDs both during first-strand cDNA synthesis as well as multiplex-PCR amplification; this protocol, known as molecular amplification fingerprinting (MAF), results in comprehensive error and bias correction of mouse antibody repertoires (18).

Here, we describe an experimental-computational approach to generate highly accurate human Ig-seq data. We first designed a comprehensive set of synthetic standards based on human antibody heavy chain variable (IGHV) sequences: a total of 85 *in vitro* transcribed RNA standards, each with a unique complementarity determining region 3 (CDR3) sequence and covering nearly the entire set

of productive human Ig heavy chain (IgH) germline (IGHV) gene segments. We used these synthetic standards to quantify the impact of errors and bias introduced during multiplex-PCR library preparation, and the robustness with which our previously developed method for UID addition by MAF could correct these artifact sequences. Finally, we implemented MAF-based error and bias correction on human B cell subsets (naïve and memory), which enabled us to make accurate clonal diversity estimates and quantify divergent repertoire features across B cell compartments.

RESULTS

Design of a comprehensive set of human synthetic standards

Our previously established set of murine synthetic antibody standards contained 16 unique clones (CDR3s) covering 7 IGHV gene segments (out of more than 140 annotated murine IGHV gene segments) (18). For our human standards, we developed a more comprehensive set consisting of 85 clones encompassing nearly the entire germline IGHV repertoire. The most commonly used repository for human germline segments is the International ImMunoGenetTics Database (IMGT), which has annotated 61 IGHV alleles as functional or having an open reading frame (25). After filtering out paralogs and selecting only gene segments that have been found in productive rearrangements (2), we chose 48 IGHV gene segments as the basis for our standards (**Table S1**). Each standard contained the following elements (5' to 3'): (i) a conserved non-coding region, (ii) ATG start codon and a leader peptide sequence spliced to its respective IGHV gene segment, (iii) a synthetic CDR3 sequence, (iv) a germline IgH J (IGHJ) gene segment, (v) a non-coding synthetic sequence identifier (for the separation of standards from biological sequences), (vi) a partial segment of the constant region from isotypes IgM, IgG, and IgA (**Figure 1A**). This design allows amplification of our synthetic controls with a variety of PCR primer sets. Notably, for control singleplex-PCR experiments, all standards can be amplified by a single forward primer (targeting the conserved 5' non-coding region) and a single reverse primer (targeting one of the isotype constant regions). Since IGHV gene segment usage has been reported to be non-uniform (2, 26), we selected the most abundant segments for use in multiple standards (**Figure 1B, Table S1**).

All standards carry a unique CDR3 sequence, which visually aids the analysis of sequencing results. Furthermore, all clones were designed to be resilient against sequencing and PCR errors: at least 9 specific nucleotide (nt) deletions, insertions, and/or mutations are needed in order to turn one CDR3 nt sequence into another (**Figure 1B**). For our experiments, synthetic standard genes were *in vitro* transcribed to RNA and subsequently reverse transcribed to cDNA. We measured individual cDNA molecules by digital droplet PCR (ddPCR) and capillary electrophoresis. Standards were then pooled in a non-uniform concentration distribution and maintained as a master stock (**Table S1**).

Human Ig-seq library preparation using the MAF protocol

We adapted our previously described library preparation protocol for murine antibody repertoires to be compatible for human Ig-seq (18). This protocol is based on targeted amplification via RT of RNA to first-strand cDNA, followed by two PCR amplification steps (27, 28), the first of which uses a forward multiplex primer set targeting the IGHV framework region 1 (FR1). Each step also

incorporates fragments of Illumina sequencing adapters (IA), such that the final product of the workflow is already compatible with the Illumina sequencing platform (**Figure 2A**).

Importantly, our library preparation protocol used primers incorporating random-nucleotide UIDs, thus enabling MAF-based error and bias correction. A reverse-UID (RID) with theoretical diversity up to $2\text{x}10^7$ unique sequences is present in the RT primer (between the Ig constant region-specific and partial IA regions), and a forward-UID (FID with additional diversity of approximately $7\text{x}10^5$ unique sequences) is present on the forward multiplex primer set (between the FR1-specific and partial IA regions) used in the first PCR reaction. Such high diversity among RIDs is necessary in order to prevent tagging of different cDNA molecules with the same barcode. Our multiplex forward primer set was designed to target all IMGT-annotated IGHV gene segments (**Figure 2B**). To compromise between maintaining similar amplicon length across gene segment families and creating thermodynamically equivalent oligonucleotides, we placed the primers at or near the beginning of each FR1, resulting in a melting temperature range of 57°C to 63°C (**Figure 2C**). This range and the accompanying (unavoidable) variability in GC content have been shown to potentially cause differences in amplification efficiencies, which in turn leads to a biased representation of segment usage frequencies in Ig-seq data. Our workflow aimed to solve this problem in two ways: first, since the RID labels cDNA at the single molecule level, we are able to resolve the number of molecules by counting the number of RIDs instead of raw reads. Second, by using the FIDs on our forward primers, we are able to further normalize our molecular count, since Ig genes preferentially amplified by our primer set should show a higher ratio of FIDs to RIDs. Additionally, the RIDs can be used to correct for errors introduced during PCR and the sequencing process itself by grouping sequencing reads based on their RID, then correcting diverging nt positions by generating a consensus sequence (majority voting scheme). This is especially useful in Ig-seq when attempting to distinguish true SHM variants from erroneous sequences.

Combining standards with MAF to correct errors and bias in Ig-seq

To evaluate the extent of errors and bias present in human Ig-seq data, standards were mixed (spiked-in) with cDNA prepared from circulating purified human B cells. In total, we sequenced 28 independently prepared libraries and annotated them with a custom aligner (18, 29). Prior to alignment, reads were either kept as uncorrected (raw) reads or were corrected using our MAF pipeline that takes into account RID and FID information. In this way, we could directly compare the number of erroneous sequences produced in uncorrected vs. MAF-corrected datasets. Clonal assignment of uncorrected reads produced many erroneous CDR3 amino acid (a.a.) variants (sequences with at least 1 a.a. difference from the nearest standard control sequence); for example, in a dataset with 100,000 aligned reads, there was a median value of 23 errors per clone (**Figure 3A**). The number of erroneous variants produced showed a clear correlation with the individual abundance of each clone within the master stock ($r = 0.89$). When taking the entire VDJ nt sequence into account, an even greater number of erroneous variants were observed (≥ 1 nt difference from the standard sequence) (**Figure 3B**). We observed a median value of 118 erroneous nt variants per standard (per 100,000 aligned sequences). Again, the number of erroneous variants exhibited a clear correlation with clone abundance ($r = 0.90$). However, we did not observe any significant trend linking IGHV family to the error rate (F-Test on full and reduced linear model, p = 0.083).

After performing error correction with our MAF pipeline, there was a dramatic reduction of CDR3 and VDJ errors: we observed a median value of 0 and 1 error per clone, respectively. Across all datasets, we remove an average of 94.2% CDR3 a.a. and 97.4% VDJ nt erroneous variants (**Figure 3A-B**). For example, prior to error correction the standard "CARGINGERALEW" (from dataset Donor 1, IgG aliquot 1, see **Table S3**) displayed 39 additional CDR3 a.a. variants and 217 additional VDJ nt variants (**Figure 3D**). After error correction, we retain only the correct CDR3 a.a. sequence and only one additional (erroneous) nt variant. With our current filtering criteria (see Methods), we observed 16 instances in which we removed a standard control sequence that was present in the raw data. However, in the vast majority of cases, we kept the standard when it was observed in the raw data (2,321 instances). In only 43 instances, a standard sequence was either too low in abundance or too frequently mutated to be annotated in either the raw or error-corrected datasets.

In order to assess potential biases introduced by library preparation we prepared control libraries containing only the pool of synthetic standards (from the master stock). The libraries were generated in the same manner as the described MAF protocol, with the exception that in the first PCR step, instead of using a multiplex forward primer set, a single forward primer targeting the conserved 5' non-coding region (singleplex-PCR) was used. Ig-seq on these samples allowed us to establish a baseline for pipetting accuracy by comparing the obtained standard frequencies from the singleplex-PCR against the expected frequencies based on our pooling scheme: this yielded an R^2 of 0.88 and average mean squared error (MSE) of 0.29 ± 0.02% (**Figure S1A**). These values indicate that only small systematic deviations occurred, most likely due to minor pipetting error. Next, we compared standard frequencies (expected relative concentration) with frequencies generated in our previous multiplex-PCR libraries, both with and without MAF correction (**Figure 3C**). On uncorrected data, the multiplex-PCR libraries achieved an R^2 of 0.84 with an average MSE of 0.34 ± 0.06%, which is significantly worse than the value obtained by singleplex-PCR (Student's t-test p = 0.008). After error and bias correction on these same datasets, the correlation improved to an R^2 of 0.89 and an average MSE of 0.28 ± 0.08%. While MAF-corrected MSE values show no significant difference to the singleplex-PCR libraries (Student's t-test p = 0.49), they do highlight a significant improvement over the uncorrected data (paired Student's t-test, p = 0.0007).

Impact of MAF error correction on human B cell repertoires

Next, we analyzed the impact of MAF on the BCR repertoires of B cells isolated from the peripheral blood of three healthy donors. We used flow cytometry sorting and a gating strategy to select for CD27$^-$ IgM$^+$ (naïve) and CD27$^+$ IgG$^+$ (memory) B cell populations (**Figure 4A**). Across all donors, the fraction of CD19$^+$ peripheral blood B cells was 16-29% for CD27$^-$ IgM$^+$ and 5-9% for CD27$^+$ IgG$^+$. Importantly, each donor population was split into 4-5 separate aliquots containing 200,000 cells each (cellular replicates) prior to cell lysis. Total RNA was extracted and RT for cDNA synthesis was performed independently in order to prevent the mixing of transcripts across cellular replicates. The cDNA of synthetic standards (from the master stock) was then mixed with the B cell cDNA, and corresponding molecular quantities were measured by ddPCR (**Figure 4B, Table S3**). All cDNA libraries were then processed into libraries using the MAF protocol (**Figure 2A**) and subjected to Ig-seq.

A simple global analysis of Ig-seq data revealed that diversity measurements were dramatically exaggerated, as the number of unique antibody sequence variants obtained from the raw,

uncorrected data often exceeded both the number of cells and the number of total cDNA transcripts
in a given aliquot. Following error correction by MAF, the variant count returned to ranges that are
physically possible, thereby highlighting the importance of proper error correction and quality
control when globally determining repertoire diversity (**Figure 4B**). We further examined the
influence of erroneous variants on CDR3 clonotype analysis. In order to identify clonotypes, we
used hierarchical clustering (30) based on sequences sharing the following features: identical IGHV
and IGHJ gene segment usage, identical CDR3 length, and a CDR3 a.a. similarity of at least 80% to
one other sequence in the given clonotype. When performing such an analysis on uncorrected data,
clonotypes contained an artificially high number of distinct clones (**Figure 4C**, left tree). Here, the
IgG-derived clonotype with the consensus CDR3 of 'CARAAGSQYYMDVW' (from the same
sample and IGHV gene segment used by the standard shown in **Figure 3D**) contains 249 unique nt
variants and 70 unique a.a. sequences. After MAF-based error correction, only 15 nt variants and 6
distinct CDR3 a.a. sequences remained. It is worthy to note that although both the standard
sequence (**Figure 3D**) and the biological clonotype (**Figure 4C**) had a large number of CDR3 a.a.
variants in uncorrected data, after error correction only the biological clonotype retained multiple
a.a. variants, suggesting these may be true variants generated *in vivo* by SHM. This general trend of
each IgG memory B cell-derived clonotype to contain more variants relative to antigen
inexperienced IgM-expressing B cells was clear across our biological data sets (**Figure 5A**).

Clonal diversity measurements of human B cell repertoires after error and bias correction

After establishing the value of performing MAF error correction on biological repertoires, we next
focused on determining the clonotype diversity present in each B cell sample. First, we determined
the overlap of clonotypes present in each cellular replicate (**Figure 6A**). Notably, we observed an
overlap of several clonotypes in the $CD27^-IgM^+$ subset; this overlap was unexpected given that this
subset should be highly enriched for naïve B cells, which by definition are not antigen experienced
or clonally expanded, and should therefore be mostly unique (not present in multiple replicates).
For each donor in the $CD27^-IgM^+$ subset, 1- 2% of all clonotypes were present in at least one other
cellular replicate. In the $CD27^+IgG^+$ subset, clonotypes shared between at least two cellular
replicates were nearly tenfold more frequent (12-15%), which was expected given that this
population is comprised of antigen experienced, clonally expanded memory B cells. Another
observation discordant with the expected naïve B cell properties of the $CD27^-IgM^+$ subset was that
overlapping clonotypes (in donors 1 and 3) were significantly more likely to have acquired
mutations (**Figure 6B**), which are not a typical feature of naïve B cells. In comparison, over 90% of
all $CD27^+IgG^+$ (overlapping and non-overlapping) clonotypes possessed at least one SHM, an
expected observation in a memory subset.

The high amount of overlap within the $CD27^+IgG^+$ B cell replicates of each donor allowed us to use
established population diversity estimation techniques to calculate clonal diversity (31). Rarefaction
curves were generated and estimates were extrapolated as a function of real and predicted cellular
replicates (**Figure 6C**). The asymptote was determined by the standard form of the Chao2 estimator
and yielded the following values for clonotype numbers: donor 1 = 164,268 ± 2,365, donor 2 =
38,034 ± 1,302, and donor 3 = 76,904 ± 1,409. Since the 95% confidence intervals for the three
donors did not overlap, we could also infer that the size of each donor's repertoire at the collection
time point was significantly different. This analysis indicates that we would need to sample at least
tenfold more cellular replicates in order to observe > 90% of all clonotypes; however the first five

samples analyzed here were sufficient to observe > 25% of the clonotypic memory repertoire. We
also generated rarefaction and extrapolation curves rescaled to the RID count (**Figures S2A and**
S2B). In the case of the CD27$^-$IgM$^+$ repertoire data, while asymptotic curves could be generated, a
diversity estimation is impractical. This is because plotting the observed numbers of newly
discovered clonotypes for each additional RID and donor shows that the number of newly
discovered clonotypes in the CD27$^-$IgM$^+$ dataset continues to grow over the observed range,
whereas the number of new clonotypes starts to converge at approximately 20,000 RIDs for
CD27$^+$IgG$^+$ repertoires (**Figure S2C**).

Divergent features of CD27$^-$IgM$^+$ and CD27$^+$IgG$^+$ repertoires

After pooling all clonotypes (expanded and unique to a single cellular replicate) for each donor, we
globally characterized sequences of the naïve CD27$^-$IgM$^+$ and memory CD27$^+$IgG$^+$ subsets. First,
we determined the SHM count (nt) of each clone with respect to its nearest germline IGHV and
IGHJ gene segment sequence. The median values of SHM for the CD27$^-$IgM$^+$ repertoires were
zero, which was to be expected for a naïve B cell subset. In contrast, the median values for
CD27$^+$IgG$^+$ repertoires were 20-24 mutations per clone (**Figure 5B**). It is widely appreciated that
human heavy chain CDR3 sequences are much longer than their murine counterparts, which we
also observed here, with a slight (but consistent across donors) variation between naïve and memory
B cells (**Figure 5C**). Interestingly, the IGHV gene segment family usage correlated with B cell
subset. The CD27$^+$IgG$^+$ repertoires across all donors were relatively enriched for IGHV1 and
IGHV3 gene segment family members, whereas the relative share of the IGHV4 gene segment
family was larger in the CD27$^-$IgM$^+$ repertoires (**Figure 7A**). We validated these observations
quantitatively using linear discriminant analysis (LDA) fitted on the centered log ratio (CLR)-
transformed frequencies of each cellular replicate (**Figure S3**). The LDA classifier was fit on
different splits of the data (based on two of the donors) and used to predict a holdout set (based on
the remaining donor). This showed that the fitted classifier in each instance is highly predictive of
the remaining aliquots and that prediction is robustly driven by the relative abundance of IGHV4
segment family usage in the CD27$^-$IgM$^+$ repertoires and the IGHV1, 2, and 3 families in the
CD27$^+$IgG$^+$ repertoires (**Figure S3A-C**). Next, we utilized LDA to perform dimensionality
reduction of all data points to into a single one-dimensional axis; again, the most important
components were the relative abundance of IGHV3 and IGHV4 gene segment families (**Figure**
S3D).

Next, we leveraged the ability of our reverse primer to distinguish among IgG subclasses (**Figure**
7B). The majority of sequences mapped either to IgG1 (40-66%), IgG2 (23%-36%), or IgG3 (23%-
36%), whereas IgG4 sequences were extremely rare, observed solely in donor 3 (0.3%). Finally, we
compared the CD27$^-$IgM$^+$ and CD27$^+$IgG$^+$ repertoires of each donor to determine the clonotype
overlap of each B cell subset and isotype. Strikingly, the observed overlap was very small, with
only 269 shared clonotypes for donor 1, 30 shared clonotypes for donor 2, and 215 clonotypes for
donor 3 (**Figure 7C**). In each donor, these represented less than 0.5% of identified clonotypes.
Closer examination revealed that clonotypes shared between the CD27$^-$IgM$^+$ and CD27$^+$IgG$^+$
subsets were also significantly enriched for intraclonal variants (SHM in CDR3) in one of the two
populations (**Figure 7D and Figure S4**). Furthermore, we could see that clonotypes with multiple
IgM variants were also shared specifically among IgM cellular replicates (**Figure 7D**, contingency
tables). This intraclonal variant bias was not limited to heavy chain isotype: IgG clonotypes with ≥

5 intraclonal variants also exhibited subclass composition skewing toward the IgG1/2 or IgG1/3
axis, but rarely at proportions similar to the overall IgG subclass distribution (**Figure 7E**).

DISCUSSION

Ig-seq is becoming an essential tool for the quantitative analysis of antibody repertoire diversity and
distribution. However, similar to other areas of high-throughput sequencing, Ig-seq also suffers
from technical errors and bias; thus, standardized experimental and analytical methods that increase
the validity of immunological interpretations must be developed. Here, we establish a
comprehensive set of synthetic standards which, when combined with UID-labeling and MAF-
based error and bias correction, results in highly accurate antibody repertoire data. By applying this
approach to human B cell subsets, we gain unique insights into repertoire features such as clonal
diversity, germline gene usage, SHM, and clonal history.

The synthetic standards developed here allowed quantitative interrogation of several accuracy-
related features in Ig-seq. One major observation was that raw uncorrected data has a high number
of erroneous variants, found both within the clonotype-defining CDR3 and across the entire VDJ
region (**Figure 3A, B**). The number of false-positive variants correlated with the abundance of each
standard; this is of particular concern because high frequency, clonally expanded B cells are often
correlated with antigen specificity and thus important for biological interpretations (32). However,
when applying our MAF-error correction protocol, we were able to remove nearly all erroneous
CDR3 and VDJ variants (94-97%); this correction was robust even for high-frequency standards
where the number of erroneous variants was especially high. Errors not removed by MAF could
potentially be addressed with more stringent filtering criteria (e.g. read number cutoffs); however,
this may come at the cost of reducing overall dataset size and removing legitimate intraclonal
variants in biological samples.

Another aspect we quantified with our standard pool was the impact of multiplex primer sets, which
have been shown to introduce substantial bias during library preparation (18, 33). By designing our
standards with a 5' conserved singleplex region (**Figure 1A**), we were able to directly compare Ig-
seq data from libraries (on the same master stock) prepared by singleplex-PCR vs. multiplex-PCR.
Our newly designed FR1-targeting multiplex primer set (**Figure 2B, C**) demonstrated a relatively
strong correlation with singleplex-PCR data ($R^2 = 0.84$) (**Figure 3C**). However, by performing
multiple MAF error and bias correction steps, the correlation was improved by an additional 7%
(**Figure 3D**). The remaining variability does not appear to be restricted to a particular IGHV-gene
family, indicating there is little systematic bias with respect to homologous sequences within the
standard pool. MAF therefore represents an essential step in eliminating technical artifacts from
human Ig-seq workflows, as it is able to generate data that closely mirrors that of the original
sample. In future applications, these synthetic standards could be a critical asset for evaluating
newly designed primer sets, library preparation protocols, or implementing new error and bias
correction pipelines.

Having established a comprehensive set of synthetic standards and a validated error and bias
correction pipeline, we were able to perform several analyses on human B cell repertoires with
greater confidence in the accuracy and quantitative resolution of the Ig-seq data. A simple approach

to estimating repertoire diversity is to calculate the number of unique antibody sequences as a
fraction of total transcript (cDNA) input. However, performing this analysis on our samples
suggests that the $CD27^+IgG^+$ memory B cell compartment is significantly more diverse than the
naïve $CD27^-IgM^+$ B cells (82% vs. 35% unique nt variants, respectively, averaged across donors
and cellular aliquots, Student's t-test $p < 10^{-4}$). This is potentially due to sample size variability with
respect to the number of transcripts and our ability to oversample smaller libraries. Critically, when
using bulk-sorted cells with UID labeling, it is not possible to discriminate between transcript
copies that are identical because they came from the same lysed cell, and those which are identical
because they represent two distinct, clonally related B cells. Thus, by biological subsampling
through cellular replicates, we ensured that clonotypes observed in multiple samples must come
from distinct, clonally related B cells, thereby providing an effective solution for estimating clonal
diversity.

Applying computational approaches from ecology (31) to our biological subsampling strategy, we
attempted to estimate the number of unique clonotypes in a given antibody repertoire. The $CD27^-$
IgM^+ B-cell subset did not show substantial clonotype overlap among cellular aliquots (**Figure 6A**).
As it is commonly assumed (and typically the case as shown in **Figure 5A**) that each newly
generated B cell is a unique clone, the size of the naïve repertoire in the human peripheral blood
would be equal to the total number of naïve B cells, in the range of 10 to 30 million. While it is
improbable to sample this subset in its entirety, and its diversity is also too high to estimate based
on the cell numbers obtained here, our observations are consistent with this model, since each
additional cellular replicate produced overwhelmingly unique sequences. One donor did show an
unexpected presence of overlapping sequences (shared clonotypes) across IgM cellular replicates
(**Figure 6A**, donor 1); these clonotype sequences were significantly enriched for SHM (**Figure 6B**),
suggesting the possible presence of an antigen-experienced B cell subset within $CD27^-IgM^+$
population, and highlighting the need for improved characterization of the heterogeneity within
circulating human B cell subsets. In the $CD27^+IgG^+$ B cell subset, we observed substantially more
overlap across cellular replicates, which was expected given that memory-enriched B cells would
have experienced antigen and undergone clonal expansion. By extrapolating the numbers of
additional uniquely observed clonotypes with each subsequent cellular aliquot, we were able to
predict the clonotype size of the peripheral $CD27^+IgG^+$ B-cell repertoire to be on the order of 10^5
(**Figure 6C**). Indeed, rough estimates of the number of antigen-specific clonotypes generated by a
single immune response ($\approx$100, a number in line with what has been described regarding serum
antibody clonotypes (34)) and the number of structurally distinct pathogens against which an
individual has mounted a response ($\approx$1000, a generous estimate given work showing that
worldwide, individuals have on average a serological history against less than 100 viral species
(35)) suggest that a memory repertoire of this size could reasonably protect against latent infection
and/or subsequent antigen encounter.

We observed a clear shift in IGHV segment family usage from the naïve to the memory BCR
repertoire (**Figure 7A**). Consistently observing this reshaping in three independent healthy donors,
and comparing to our standard controls to exclude the possibility of biased amplification, we can
conclude that it is a genuine phenomenon. Relatively more abundant IGHV1 and less abundant
IGHV4 segment usage in IgG memory B cells has been previously observed in one three-donor
cohort (1) but not in another which pooled sequences from both class-switched and IgM-memory

cells (36), underscoring the importance of experimental design and accurate bias correction in antibody repertoire analysis.

Our Ig-seq workflow also allowed us to unambiguously assign IgG antibody sequences to their appropriate subclass, offering further insight into patterns of class-switch recombination present in memory-enriched B cells. While plasma cell-secreted IgG proteins in human serum are present at ratios of approximately 14:8:1:1 (for IgG1:2:3:4, respectively (37)), CD27$^+$IgG$^+$ B cells showed a distribution of approximately 5:3:1 for IgG1:2:3, with a nearly complete absence of IgG4 (**Figure 7B**). Cole et al. similarly observed a lack of IgG4 heavy chains in a single donor but described an enrichment of IgG2 relative to IgG1 and IgG3 (24). The abundance of IgG3$^+$ B cells relative to its presence in the serum seen here indicates IgG3 may play a more important role in maintaining the reactive memory response compared to the protective memory response provided by serum IgG. Notably, the IgG3 locus is the most proximal, and thereby the most plastic of the human IgG subclasses; that is, an IgG3$^+$ B cell still retains the capacity to class-switch to any of the remaining three IgG subclasses, whereas IgG1, IgG2, and IgG4 cannot return to any of the previous states. Similar to these findings, a flow cytometry-based investigation has also found healthy human donors to have low frequencies of IgG4-expressing circulating memory B cells (38).

With new daily production and relatively rapid turnover of naïve B cells, it was not unexpected to see little overlap of clonotypes between the intradonor CD27$^-$IgM$^+$ and CD27$^+$IgG$^+$ populations (**Figure 7C**). An interesting finding was that for clonotypes present in both B cell subsets, intraclonal variation was largely restricted to one of the two isotypes. Assuming that clonotype overlap among CD27$^-$IgM$^+$ cellular replicates represents the presence of antigen-experienced, clonally expanded B cells, this suggests that the antigen specificity of antibody variable domains may to some extent be influenced by the downstream constant regions, which has been observed functionally for small cohorts of human and murine IgG and IgA antibodies (39). Notably, we also observe similar clonal restriction within IgG clonotypes with respect to heavy chain subclass (**Figure 7E**). This may be driven by the type of antigen and the nature of the elicited immune response, or governed by physical constraints as we suggest for the differences between the IgM and IgG repertoires. A larger scale functional study, including IgM sequences, could provide crucial support for this model, which would shed new light on the role of Ig isotypes and subclasses on B cells in the post-antigen encounter setting.

FIGURE LEGENDS

Figure 1. A comprehensive set of human synthetic spike-in standards for Ig-seq. **(A)** Schematic showing the prototypical spike-in with the following regions (5' to 3'): a conserved non-coding region, ATG start codon, IGHV region with FR1 specific for multiplex-PCR, IGHJ regions, non-coding synthetic sequence identifier (specific for ddPCR probes), and downstream heavy chain constant domain sequences (IGHG, IGHM, IGHA) containing primer binding sites used for cDNA synthesis. Each spike-in contains a complete VDJ open reading frame, including nucleotides upstream of the ATG start codon, and downstream constant domain sequences containing primer binding sites used for sample cDNA synthesis. **(B)** Pairwise comparisons based on a.a. Levenshtein edit distance of all 85 standard CDR3 sequences. The germline IGHV and IGHJ segment family usage and IgG subclasses are denoted. 39 spike-ins contain rationally designed nt SHM (black circles) across the IGHV regions.

Figure 2. Library preparation of immunoglobulin (Ig) heavy chain genes for high-throughput sequencing (Ig-seq) using molecular amplification fingerprinting (MAF). **(A)** In step 1, reverse transcription (RT) is performed to generate first-strand cDNA with a gene-specific (IgM or IgG) primer which includes a unique reverse molecular identifier (RID) and partial Illumina adapter (IA) region. This results in single-molecule labeling of each cDNA with an RID. In step 2, several cycles of multiplex-PCR are performed using a forward primer set with gene-specific regions targeting heavy chain variable (V_H) framework region 1 (FR1), with overhang regions comprised of a forward unique molecular identifier (FID) and partial IA. In step 3, singleplex-PCR is used to extend the partial IAs. The result (Step 4) is the generation of antibody amplicons with FID, RIDs, and full IA ready for Ig-seq and subsequent MAF-based error and bias correction. **(B)** List of oligonucleotides sequences annealing to the V_H FR1 used in multiplex-PCR (Step 2) of the MAF library preparation protocol. The nearest germline IGHV segment(s) likely to be amplified by the respective primer are listed in the rightmost column. **(C)** The estimated melting temperature distribution of the V_H FR1 forward primer set.

Figure 3. Synthetic standards used to validate performance of error and bias correction of Ig-seq data by MAF. **(A)** The number of erroneous CDR3 sequences (at least one a.a. difference from the correct CDR3) per 100,000 reads is plotted against the relative concentration of each standard (from a master stock). Color-coded diamonds correspond to germline IGHV segment family of the respective standard and show the number of erroneous variants in uncorrected (raw) data; gray diamonds indicate the number of variants remaining after MAF error correction. **(B)** The number of erroneous VDJ variants derived from each standard was calculated by finding all variants that carried the correct CDR3 a.a. sequence, but differed by at least one nt across the entire VDJ region. Colored diamonds represent uncorrected data; gray diamonds indicate variants remaining after MAF error correction. **(C)** Sequencing bias introduced by multiplex-PCR using the FR1 primer set was assessed by plotting the measured frequencies of each standard against its relative concentration (from a master stock). Dashed line represents a bias-free ideal scenario ($R^2 = 1$). The left and right plots show observed frequencies before and after MAF bias correction, respectively. **(D)** Phylogenetic trees visualizing the CDR3 a.a. variants present for a selected standard with the CDR3 a.a. sequence *CARGINGERALEW* and IGHV1-8 and IGHJ1 segment usage. Prior to error correction, 39 erroneous CDR3 a.a. variants (branches) and 218 VDJ nt variants (black circles) were observed. Following MAF error correction, only the original correct CDR3 a.a. and two VDJ nt variants remain. The Ig-seq data sets used in A-C consisted of ~300,000 preprocessed full-length antibody reads from each of the synthetic spike-in only samples. IgG1_D1 dataset was used for panel **(D)** (see **Table S3**).

Figure 4. Ig-seq analysis of human naïve (CD27⁻IgM⁺) and memory (CD27⁺IgG⁺) B cells. **(A)** The flow cytometric workflow for isolating CD27⁻IgM⁺ and CD27⁺IgG⁺ B cells from peripheral blood. Boxed-in values indicate the frequency of each sorted subset as a percentage of the total B cell (CD19⁺) population from each of three donors (1-3 from left to right). **(B)** Experimental and Ig-seq based quantitation of antibody diversity; points represent cDNA molecule counts (using ddPCR) or unique reads (before and after MAF error correction) from cellular replicates (with mean and standard deviation shown) isolated from each donor and B cell subset. Unique read counts were based on the VDJ nt seqeunce. Dashed line represents the number of B cells isolated per cellular replicate (2×10^5 cells). **(C)** Phylogenetic trees illustrating CDR3 a.a. and nt variants present for the

selected clonotype with the consensus CDR3 sequence *CARAAGSQYYYMDVW* and IGHV1-8 to
IGHJ1 recombination. Prior to error correction, 70 erroneous CDR3 a.a. variants and 249 VDJ nt
variants (black circles) were observed. Following MAF error correction, only 6 CDR3 a.a. and 15
VDJ nt variants remain. The Ig-seq data sets used in **(B)** are described in **Table S3**; IgG1_D1 was
used for the tree in panel **(C)**.

Figure 5. Ig-seq analysis of molecular features highlight global differences between naïve (CD27$^-$
IgM$^+$) and memory (CD27$^+$IgG$^+$) B cells. **(A)** Clonotype size (calculated as the total number of
variants within a clonotype) for naïve IgM (blue lines) and memory IgG (red lines) B cells. Each
pair of lines represents a single donor. **(B)** Graph showing the distribution of average SHM
frequencies for VDJ nt variants per clonotype. The CD27$^-$IgM$^+$ B cell repertoires (blue lines) have a
median SHM value of zero, whereas only a small fraction of clonotypes (approximately 8%)
contain an average of one or more mutations. CD27$^+$IgG$^+$ repertoires (red lines) have a median of
20 to 24 SHM per nt variant within each clonotype. **(C)** CDR3 a.a. length distribution across
clonotypes from naïve (blue lines) and memory (red lines) B cell subsets.

Figure 6. Clonotype diversity analysis across cellular replicates of naïve (CD27$^-$IgM$^+$) and memory
(CD27$^+$IgG$^+$) B cells. **(A)** Venn diagrams show the presence of clonotypes (80% CDR3 a.a.
similarity to least one clone in the cluster, same CDR3 a.a. length, same IGHV and IGHJ gene
segment usage) across cellular replicates (2 x 10^5 cells each) from each donor B cell subset. **(B)** Bar
graph showing the fraction of clonotypes containing at least one variant with at least one nt SHM.
Blue and red bars indicate clonotypes identified either in only one aliquot (unique) or in several
aliquots (shared), respectively. The p-values represent significance using Fisher's exact test. **(C)**
Species accumulation curves for CD27$^+$IgG$^+$ B cells: the number of newly discovered clonotypes
from each additional cellular replicate (black circles) is plotted. Extrapolating the observed overlap
provides an estimate for the total number of distinct clonotypes (Chao2 estimator: D1 = 164,268 ±
2,365; D2 = 38,034 ± 1,302; D3 = 76,904 ± 1,409) and the approximate amount of cellular
replicates needed to discover all clonotypes present in the peripheral blood CD27$^+$ IgG$^+$ population.

Figure 7. Genetic features of naïve and IgG memory BCR repertoires. **(A)** Block map shows IGHV
gene segment usage sorted by family (color-coded) across donors and B cell subset; blocks are
normalized to the total number of clonotypes within each group. **(B)** IgG subclass usage in
CD27$^+$IgG$^+$ donor repertoires. IgG$_4$ sequences (dark blue bars) were virtually absent among three
donors; a small fraction of sequences could not be unambiguously mapped based on the sequencing
read (gray bars). **(C)** Venn diagrams showing the overlap of clonotypes (80% CDR-H3 amino acid
similarity to least one clone in the cluster, same CDR3 a.a. length, same IGHV and IGHJ gene
segment usage) between the naïve (CD27$^-$IgM$^+$) and memory (CD27$^+$IgG$^+$) BCR heavy chain
repertoire in each of three donors. **(D)** Each plot shows the clonal composition of each shared
clonotype (from panel **(C)**) in terms of its IgG and IgM intraclonal variants. Red triangles indicate
clonotypes found in multiple IgM cellular aliquots; blue triangles show clonotypes which could
only be found in one IgM cellular aliquot. The total number of clonotypes found are depicted in the
corresponding contingency table. Fisher's exact test was used to quantitatively analyze enrichment
of expanded IgM clonotypes in the shared IgM/IgG subset. **(E)** Ternary plots comprised of three
axes representing the IgG1, IgG2, and IgG3 isotype subclasses. The relative subclass composition
of intraclonal variants per IgG clonotype (each represented by a circle colored according to the
number of variants belonging to that clonotype) is depicted by the position of the circle within the

triagonal space. Red circles represent the average subclass composition for all IgG variants of each
donor (cf. **Figure 6C**).
525
526
527
528
529 **METHODS**
530
531 **Preparation of spike-in master stocks**
532
The spike-in standards were ordered from GeneArt (Invitrogen) in the form of plasmids. Each
spike-in sequence contained a T7 promoter for *in vitro* transcription. Approximately 1.5 μg of each
plasmid was digested with 10 U of EcoRV-HF (New England BioLabs) and purified with DNA-
binding magnetic beads (SPRI select, Beckman Coulter). Approximately 1 μg of the digested
plasmid was then used for *in vitro* transcription (MEGAscript T7 Transcription Kit, ThermoFisher
Scientific). RNA was purified by lithium-chloride precipitation, eluted (TE with 1U/μl RiboLock)
and aliquoted. The final concentration was then determined with the TapeStation (Agilent
Technologies).
The spike-in RNA obtained this way was reverse transcribed with the Maxima Reverse
Transcriptase kit (Thermofisher Scientific). 500 ng mRNA was mixed with 20 pmol of IgM reverse
primer and 3 μl dNTP-mix (10 mM each) and was then filled up to 14.5 μl with water. The
reaction-mix was incubated at 65 °C for 5 minutes. 4 μl of 5x RT-buffer, 0.5 μl (20 U) RiboLock
and 1 μl Maxima reverse transcriptase (200 U) were then added. The resulting reaction mix was
incubated for 35 minutes at 55 °C, followed by a termination step at 85 °C for 5 minutes. 2.5 μl of
RNase A (Thermofisher Scientific) was added and the mix was again incubated at 60 °C for 30
minutes. The resulting cDNA was purified with SPRI Select magnetic beads and eluted in nuclease-
free water. The concentration of each cDNA reaction was determined afterwards with the Fragment
Analyzer and pooled according to Table S1. The exact concentration of the pooled spike-ins was
determined by ddPCR with dilutions of the pool ranging from 10^{-3} to 10^{-6}. The measured spike-in
pool was afterwards diluted to a final storage concentration of 250,000 transcripts per μl.
553
554 **Transcript quantitation by ddPCR**
555
Quantifiying cDNA and PCR products by ddPCR was conducted for all measurements in the
following way: A dilution series with 3 or 4 points was prepared. Droplets were generated with
BioRad's droplet generator using 12.25 μl of ddPCR Supermix (BioRad) combined with 10 μl of
the diluted sample, 25 pmol of the biological ddPCR probe (SF_21), 25 pmol of the spike-in
specific ddPCR probe (TAK_499), 22.5 pmol of the forward (SF_63) and reverse ddPCR primer
(TAK_522) and 55 μl of droplet generation oil (BioRad). Droplets were then transferred to a 96-
well reaction plate, which was heat sealed with easy pierce foil (VWR International). Then a PCR
reaction was performed using the following conditions: 95°C for 10 min; 45 cycles of 94 °C for
30s, 53 °C for 30s, 64 °C for 1 min; 98°C for 10 min; and holding at 4 °C. After the PCR step,
every 96-well plate was read using BioRad's droplet reader.
566
567 **B-cell isolation, sorting, and lysis**
568

Peripheral blood leukocyte-enriched fractions ('buffy coats') were received from the Bern
(Switzerland) blood donation center after obtaining the proper informed consent from healthy
human donors. Blood samples were diluted 1:3 with sterile PBS and overlaid on Ficoll-Paque
PLUS (GE Healthcare) using LeukoSep conical centrifuge tubes (Greiner Bio-One). Peripheral
blood mononuclear cells (PBMCs) were harvested after separation for 30 minutes at 400 x *g*
without braking. Successive centrifugation steps were performed to wash the mononuclear cell
fraction and remove residual neutrophils and granulocytes. Total B cells were isolated from PBMCs
by negative selection with the EasySep Human B Cell Enrichment Kit (STEMCELL Technologies)
according to the manufacturer's instructions. The following fluorescently labeled antibodies were
used to stain the enriched B-cell fraction prior to sorting by flow cytometry: anti-CD3-APC/Cy7
(clone HIT3a BioLegend # 300318), anti-CD14-APC/Cy7 (clone HCD14 BioLegend #325620),
anti-CD16-APC/Cy7 (clone 3G8 BioLegend #302018), anti-CD19-BV785 (clone HIB19,
BioLegend #302240), anti-CD20-BV650 (clone 2H7, BioLegend #302336), anti-CD27-V450
(clone M-T271, BD Horizon #560448), anti-CD24-BV510 (clone ML5, BioLegend #311126), anti-
CD38-PC5 (clone LS198-4-3, Beckman Coulter #A07780), anti-IgD-PEcy7 (clone IA6-2,
BioLegend #348210), anti-IgG-Alexa Fluor 647 (Jackson ImmunoResearch #109-606-003) anti-
IgA-Alexa Fluor 488 (Jackson ImmunoResearch #109-549-011), anti-IgM-PE (clone SA-DA4,
eBioscience #12-9998). Cell sorting was performed on a BD FACS Aria III following the gating
strategy depicted in Figure 4A. After sorting, isolated fractions were centrifuged 5 minutes at 300 x
g, the supernatants were aspirated, and the cell pellets were re-suspended in 1 ml PBS. Recovered
cells were hand-counted using a Neubauer hemocytometer, and aliquots containing equal numbers
of cells were prepared from the cellular suspension. These aliquots were centrifuged, and the
supernatant aspirated. Cell pellets were lysed directly in 200 μl TRI Reagent (Sigma), allowed to
dissociate for 5 minutes at room temperature, then frozen on dry ice prior to storage at -80°C.
593

594   **RNA isolation from sorted B-cell populations**
595

Immediately prior to use, Phase Lock Gel (PLG) tubes were pelleted at 12000 - 16000 x g in a
microcentrifuge for 20 to 30 seconds. Each TRIzol aliquot was then thawed on ice. After thawing
and an incubation time of 5 minutes at room temperature, 1 mL of the TRIzol homogenate was
transferred to the phase lock tube. 0.2 mL chloroform was added and the tube was shaken
vigorously by hand (~15 seconds). After an incubation time of 3 minutes, the phase lock tube was
centrifuged at 12'000 x g during 15 minutes at 4 °C. The resulting upper aqueous phase was
transferred to a fresh Eppendorf tube, an equal volume of 70% ethanol was added and the solution
was purified on a PureLink RNA column according to the manufacturer's instructions (Life
Technologies). Finally, RNA was eluted in 25 μl nuclease-free water.
605

606   **Library preparation and NGS on Illumina's MiSeq**
607

First-strand cDNA synthesis was carried out using Maxima reverse transcriptase (Life
Technologies). The protocol for one reaction is as follows: A 29 μl reaction mix was prepared using
up to 2 μg of RNA together with 40 pmol of the respective gene-specific reverse primer (for IgG
sequences, 5'-RID-GTTCTGGGAAGTAGTCCTTGACCAG-3' (IgH_10r); for IgM, 5'-RID-
ACGAGGGGGAAAAGGGTTGG-3' (CH1_1r)), 2 μl dNTP (10 mM each) and the required
amount of nuclease-free water. This mix is incubated for 5 minutes at 65 °C. A master mix of 8 μl
5x RT buffer, 1 μl of (20 U) RiboLock and 2 μl of Maxima reverse transcriptase is then prepared

and added to the reaction mix. Finally, the mix is incubated for 30 minutes at 50°C and the reaction
is terminated by incubating at 85°C for 5 minutes.
The obtained cDNA is then cleaned with a left-sided SPRI-Select bead clean up (0.8x) according to
the manufacturer's instructions (Beckman coulter) and subsequently measured by ddPCR.
Up to 135,000 cDNA transcripts were then pooled together with 12,500 spike-in transcripts.
Multiplex-PCR was performed using an equimolar pool of the forward primer mix (Figure 2B) and
the reverse primer (TAK_423) targeting the overhang introduced during cDNA synthesis. Due to
low cDNA yields, the first PCR was carried out for 20 cycles and the following cycling protocol: 2
min at 95 °C; 20 cycles of 98°C for 20s; 60°C for 50; 72°C for 1min; 72°C and then holding at 4°C.
PCR reactions were prepared using 15 μl of Kapa HiFi HotStart ReadyMix (KAPA Biosystems), 50
pmol of the forward mix and the reverse primer and 9 μl of the cDNA mix. After the first PCR, we
again performed a left-sided bead clean-up (0.8x) and measured the PCR product concentration
using ddPCR. We use 800,000 transcripts from PCR 1 as input into the adapter extension PCR. For
this PCR 25 μl of Kapa HiFI Hotstart ReadyMix was combined with 25 pmol of the forward primer
(TAK_424) and 25 pmol of the index primer (TAK_531) as well as the diluted PCR product.
Finally, the reaction volume was adjusted to 50 μl by the addition of nuclease-free water.
Thermocycling was performed as follows: 95°C for 5 min; 23 cycles of 98°C for 20 s, 65°C for 15
s, 72°C for 15 s; 72°C for 5 min; and 4°C indefinitely. Following second-step adapter extension
PCR, reactions were cleaned using a double-sided SPRIselect bead cleanup process (0.5x to 0.8x),
with an additional ethanol wash and elution in TE buffer.
Libraries were then quantified by capillary electrophoresis (Fragment analyzer, Agilent). After
quantitation, libraries were pooled accordingly and sequenced on a MiSeq System (Illumina) with
the paired-end 2x300bp kit.
638
639 **Bioinformatic pipeline**
640
Paired-end fastq files were merged using PandaSeq (40). Afterwards, sequences were filtered for
quality and length using the FASTX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). After the
quality trim, sequences were processed with a custom Python script that performed error correction
by consensus building on our sequences and RIDs. In order to utilize as many sequencing reads as
possible, we required UIDs to have at least 3 reads, but did not remove sequences that only had one
UID group mapping to them. VDJ annotation and frequency calculation was then performed by our
in-house aligner (18) which was updated with the human reference database downloaded from
IMGT. The complete error-correction and alignment pipeline is available under
https://gitlab.ethz.ch/reddy/MAF.
650
651 **Statistical analysis**
652
653 All statistical and computational analyses following the alignment step were performed in R.
654 Details about specific tests that were used can be found in the results section and in the figure
655 legends. Scripts are available upon request.
656
657 **Data availability**
658
659 In adherence to the data sharing recommendations of the AIRR community our data is publically
660 available in the following repositories: BioProject, BioSample, SRA and GenBank and can be
661 accessed with the accession number PRJNA430091 (BioProject). The exact data processing steps,

662 including software tools and version numbers can be found on zonodo.org under the following doi:
663 10.5281/zenodo.1201416.
664
665 Likewise, the designed spike-in sequences are also stored on GenBank (Accession number
666 MG785894-MG785978).
667

Author contributions

J.M.L., S.F., E.T., and S.T.R. designed experiments. V.C. performed B-cell enrichment, sorting,
and mRNA extraction. M.I. and S.F. prepared IgH libraries. J.M.L. designed primer sequences.
J.M.L. and A.Z. designed antibody spike-ins. A.Z., S.M., and M.I. conducted preliminary
experiments. E.M. analysed preliminary data. S.F. was responsible for the bioinformatics pipeline.
J.M.L. and S.F. analysed the final data and prepared figures. J.M.L., S.F., E.T., and S.T.R. wrote the
paper. All authors provided scientific guidance.

Acknowledgments

678 We would like to acknowledge the Genomics Facility Basel of ETH Zurich for Illumina sequencing
679 support, in particular E. Burcklen, K. Eschbach and C. Beisel. We also want to thank H.
680 Ruscheweyh for bioinformatic code support.

REFERENCES

1.   Wu Y-C, Kipling D, Leong HS, Martin V, Ademokun AA, Dunn-Walters DK. High-throughput immunoglobulin
repertoire analysis distinguishes between human IgM memory and switched memory B-cell populations. *Blood* (2010)
116(7):1070-8. doi: 10.1182/blood-2010-03-275859. PubMed PMID: 20457872; PubMed Central PMCID:
PMCPMC2938129.
2.   Arnaout R, Lee W, Cahill P, Honan T, Sparrow T, Weiand M, et al. High-resolution description of antibody
heavy-chain repertoires in humans. *PLoS ONE* (2011) 6(8):e22365. doi: 10.1371/journal.pone.0022365. PubMed PMID:
21829618; PubMed Central PMCID: PMCPMC3150326.
3.   DeKosky BJ, Kojima T, Rodin A, Charab W, Ippolito GC, Ellington AD, et al. In-depth determination and
analysis of the human paired heavy- and light-chain antibody repertoire. *Nature medicine* (2015) 21(1):86-91. doi:
10.1038/nm.3743. PubMed PMID: 25501908.
4.   Robinson WH. Sequencing the functional antibody repertoire--diagnostic and therapeutic discovery. *Nat Rev*
Rheumatol (2015) 11(3):171-82. doi: 10.1038/nrrheum.2014.220. PubMed PMID: 25536486; PubMed Central PMCID:
PMCPMC4382308.
5.   Williams LD, Ofek G, Schätzle S, McDaniel JR, Lu X, Nicely NI, et al. Potent and broad HIV-neutralizing
antibodies in memory B cells and plasma. *Science immunology* (2017) 2(7):eaal2200. doi: 10.1126/sciimmunol.aal2200.
PubMed PMID: 28783671.
6.   Doria-Rose NA, Schramm CA, Gorman J, Moore PL, Bhiman JN, DeKosky BJ, et al. Developmental pathway
for potent V1V2-directed HIV-neutralizing antibodies. *Nature* (2014) 509(7498):55-62. doi: 10.1038/nature13036.
PubMed PMID: 24590074; PubMed Central PMCID: PMCPMC4395007.
7.   Zhu J, Ofek G, Yang Y, Zhang B, Louder MK, Lu G, et al. Mining the antibodyome for HIV-1-neutralizing
antibodies with next-generation sequencing and phylogenetic pairing of heavy/light chains. *Proceedings of the National*
Academy of Sciences (2013) 110(16):6470-5. doi: 10.1073/pnas.1219320110. PubMed PMID: 23536288; PubMed
Central PMCID: PMCPMC3631616.
8.   Lavinder JJ, Wine Y, Giesecke C, Ippolito GC, Horton AP, Lungu OI, et al. Identification and characterization of
the constituent human serum antibodies elicited by vaccination. *Proceedings of the National Academy of Sciences*
(2014) 111(6):2259-64. doi: 10.1073/pnas.1317793111. PubMed PMID: 24469811; PubMed Central PMCID:
PMCPMC3926051.
9.   Jiang N, He J, Weinstein JA, Penland L, Sasaki S, He X-S, et al. Lineage structure of the human antibody
repertoire in response to influenza vaccination. *Science translational medicine* (2013) 5(171):171ra19-ra19. doi:
10.1126/scitranslmed.3004794. PubMed PMID: 23390249; PubMed Central PMCID: PMCPMC3699344.

10.   Roskin KM, Simchoni N, Liu Y, Lee J-Y, Seo K, Hoh RA, et al. IgH sequences in common variable immune
deficiency reveal altered B cell development and selection. *Science translational medicine* (2015) 7(302):302ra135-
302ra135. doi: 10.1126/scitranslmed.aab1216. PubMed PMID: 26311730; PubMed Central PMCID: PMCPMC4584259.
11.   Palanichamy A, Apeltsin L, Kuo TC, Sirota M, Wang S, Pitts SJ, et al. Immunoglobulin class-switched B cells
form an active immune axis between CNS and periphery in multiple sclerosis. *Science translational medicine* (2014)
6(248):248ra106-248ra106. doi: 10.1126/scitranslmed.3008930. PubMed PMID: 25100740; PubMed Central PMCID:
PMCPMC4176763.
12.   Meng W, Zhang B, Schwartz GW, Rosenfeld AM, Ren D, Thome JJC, et al. An atlas of B-cell clonal distribution
in the human body. *Nature Biotechnology* (2017) 35(9):879-84. doi: 10.1038/nbt.3942. PubMed PMID: 28829438;
PubMed Central PMCID: PMCPMC5679700.
13.   Lee YN, Frugoni F, Dobbs K, Tirosh I, Du L, Ververs FA, et al. Characterization of T and B cell repertoire
diversity in patients with RAG deficiency. *Sci Immunol* (2016) 1(6). doi: 10.1126/sciimmunol.aah6109. PubMed PMID:
28783691; PubMed Central PMCID: PMCPMC5586490.
14.   Friedensohn S, Khan TA, Reddy ST. Advanced Methodologies in High-Throughput Sequencing of Immune
Repertoires. *Trends in biotechnology* (2017) 35(3):203-14. doi: 10.1016/j.tibtech.2016.09.010. PubMed PMID: 28341036.
15.   Georgiou G, Ippolito GC, Beausang J, Busse CE, Wardemann H, Quake SR. The promise and challenge of
high-throughput sequencing of the antibody repertoire. *Nature Biotechnology* (2014) 32(2):158-68. doi:
10.1038/nbt.2782. PubMed PMID: 24441474; PubMed Central PMCID: PMCPMC4113560.
16.   Jiang L, Schlesinger F, Davis CA, Zhang Y, Li R, Salit M, et al. Synthetic spike-in standards for RNA-seq
experiments. *Genome Research* (2011) 21(9):1543-51. doi: 10.1101/gr.121095.111. PubMed PMID: 21816910; PubMed
Central PMCID: PMCPMC3166838.
17.   Shugay M, Britanova OV, Merzlyak EM, Turchaninova MA, Mamedov IZ, Tuganbaev TR, et al. Towards error-
free profiling of immune repertoires. *Nature Methods* (2014) 11(6):653-5. doi: 10.1038/nmeth.2960. PubMed PMID:
24793455.
18.   Khan TA, Friedensohn S, Gorter de Vries AR, Straszewski J, Ruscheweyh H-J, Reddy ST. Accurate and
predictive antibody repertoire profiling by molecular amplification fingerprinting. *Science advances* (2016) 2(3):e1501371.
doi: 10.1126/sciadv.1501371. PubMed PMID: 26998518; PubMed Central PMCID: PMCPMC4795664.
19.   Kinde I, Wu J, Papadopoulos N, Kinzler KW, Vogelstein B. Detection and quantification of rare mutations with
massively parallel sequencing. *Proceedings of the National Academy of Sciences* (2011) 108(23):9530-5. doi:
10.1073/pnas.1105422108. PubMed PMID: 21586637; PubMed Central PMCID: PMCPMC3111315.
20.   Shiroguchi K, Jia TZ, Sims PA, Xie XS. Digital RNA sequencing minimizes sequence-dependent bias and
amplification noise with optimized single-molecule barcodes. *Proceedings of the National Academy of Sciences* (2012)
109(4):1347-52. doi: 10.1073/pnas.1118018109. PubMed PMID: 22232676; PubMed Central PMCID:
PMCPMC3268301.
21.   Kivioja T, Vähärautio A, Karlsson K, Bonke M, Enge M, Linnarsson S, et al. Counting absolute numbers of
molecules using unique molecular identifiers. *Nature Methods* (2011) 9(1):72-4. doi: 10.1038/nmeth.1778. PubMed
PMID: 22101854.
22.   Vollmers C, Sit RV, Weinstein JA, Dekker CL, Quake SR. Genetic measurement of memory B-cell recall using
antibody repertoire sequencing. *Proceedings of the National Academy of Sciences* (2013) 110(33):13463-8. doi:
10.1073/pnas.1312146110. PubMed PMID: 23898164; PubMed Central PMCID: PMCPMC3746854.
23.   Turchaninova MA, Davydov A, Britanova OV, Shugay M, Bikos V, Egorov ES, et al. High-quality full-length
immunoglobulin profiling with unique molecular barcoding. *Nature Protocols* (2016) 11(9):1599-616. doi:
10.1038/nprot.2016.093. PubMed PMID: 27490633.
24.   Cole C, Volden R, Dharmadhikari S, Scelfo-Dalbey C, Vollmers C. Highly Accurate Sequencing of Full-Length
Immune Repertoire Amplicons Using Tn5-Enabled and Molecular Identifier-Guided Amplicon Assembly. *The Journal of*
Immunology (2016) 196(6):2902-7. doi: 10.4049/jimmunol.1502563. PubMed PMID: 26856699.
25.   Lefranc M-P, Giudicelli V, Duroux P, Jabado-Michaloud J, Folch G, Aouinti S, et al. IMGT®, the international
ImMunoGeneTics information system® 25 years on. *Nucleic acids research* (2015) 43(Database issue):D413-22. doi:
10.1093/nar/gku1056. PubMed PMID: 25378316; PubMed Central PMCID: PMCPMC4383898.
26.   Wang C, Liu Y, Xu LT, Jackson KJL, Roskin KM, Pham TD, et al. Effects of aging, cytomegalovirus infection,
and EBV infection on human B cell repertoires. *The Journal of Immunology* (2014) 192(2):603-11. doi:
10.4049/jimmunol.1301384. PubMed PMID: 24337376; PubMed Central PMCID: PMCPMC3947124.
27.   Menzel U, Greiff V, Khan TA, Haessler U, Hellmann I, Friedensohn S, et al. Comprehensive evaluation and
optimization of amplicon library preparation methods for high-throughput antibody sequencing. *PLoS ONE* (2014)
9(5):e96727. doi: 10.1371/journal.pone.0096727. PubMed PMID: 24809667; PubMed Central PMCID:
PMCPMC4014543.
28.   Greiff V, Menzel U, Haessler U, Cook SC, Friedensohn S, Khan TA, et al. Quantitative assessment of the
robustness of next-generation sequencing of antibody variable gene repertoires from immunized mice. *BMC Immunology*
(2014) 15(1):40. doi: 10.1186/s12865-014-0040-5. PubMed PMID: 25318652; PubMed Central PMCID:
PMCPMC4233042.
29.   Laserson U, Vigneault F, Gadala-Maria D, Yaari G, Uduman M, Vander Heiden JA, et al. High-resolution
antibody dynamics of vaccine-induced immune responses. *Proceedings of the National Academy of Sciences* (2014)
111(13):4928-33. doi: 10.1073/pnas.1323862111. PubMed PMID: 24639495; PubMed Central PMCID:
PMCPMC3977259.
30.   Greiff V, Miho E, Menzel U, Reddy ST. Bioinformatic and Statistical Analysis of Adaptive Immune Repertoires.
Trends in immunology (2015) 36(11):738-49. doi: 10.1016/j.it.2015.09.006. PubMed PMID: 26508293.

31.   Colwell RK, Chao A, Gotelli NJ, Lin SY, Mao CX, Chazdon RL, et al. Models and estimators linking individual-
based and sample-based rarefaction, extrapolation and comparison of assemblages. *Journal of Plant Ecology* (2012)
5(1):3-21. doi: 10.1093/jpe/rtr044.
32.   Reddy ST, Ge X, Miklos AE, Hughes RA, Kang SH, Hoi KH, et al. Monoclonal antibodies isolated without
screening by analyzing the variable-gene repertoire of plasma cells. *Nature Biotechnology* (2010) 28(9):965-9. doi:
10.1038/nbt.1673. PubMed PMID: 20802495.
33.   Carlson CS, Emerson RO, Sherwood AM, Desmarais C, Chung M-W, Parsons JM, et al. Using synthetic
templates to design an unbiased multiplex PCR assay. *Nature communications* (2013) 4:2680. doi:
10.1038/ncomms3680. PubMed PMID: 24157944.
34.   Lee J, Boutz DR, Chromikova V, Joyce MG, Vollmers C, Leung K, et al. Molecular-level analysis of the serum
antibody repertoire in young adults before and after seasonal influenza vaccination. *Nature medicine* (2016)
22(12):1456-64. doi: 10.1038/nm.4224. PubMed PMID: 27820605; PubMed Central PMCID: PMCPMC5301914.
35.   Xu GJ, Kula T, Xu Q, Li MZ, Vernon SD, Ndung'u T, et al. Viral immunology. Comprehensive serological
profiling of human populations using a synthetic human virome. *Science (New York, NY)* (2015) 348(6239):aaa0698. doi:
10.1126/science.aaa0698. PubMed PMID: 26045439; PubMed Central PMCID: PMCPMC4844011.
36.   DeWitt WS, Lindau P, Snyder TM, Sherwood AM, Vignali M, Carlson CS, et al. A Public Database of Memory
and Naive B-Cell Receptor Sequences. *PLoS ONE* (2016) 11(8):e0160853. doi: 10.1371/journal.pone.0160853. PubMed
PMID: 27513338; PubMed Central PMCID: PMCPMC4981401.
37.   Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: from structure to effector functions.
Frontiers in immunology (2014) 5(16):520. doi: 10.3389/fimmu.2014.00520. PubMed PMID: 25368619; PubMed Central
PMCID: PMCPMC4202688.
38.   Heeringa JJ, Karim AF, van Laar JAM, Verdijk RM, Paridaens D, van Hagen PM, et al. Expansion of blood
IgG4(+) B, TH2, and regulatory T cells in patients with IgG4-related disease. *J Allergy Clin Immunol* (2017). doi:
10.1016/j.jaci.2017.07.024. PubMed PMID: 28830675.
39.   Torres M, Casadevall A. The immunoglobulin constant region contributes to affinity and specificity. *Trends
Immunol* (2008) 29(2):91-7. doi: 10.1016/j.it.2007.11.004. PubMed PMID: 18191616.
40.   Masella AP, Bartram AK, Truszkowski JM, Brown DG, Neufeld JD. PANDAseq: paired-end assembler for
illumina sequences. *BMC bioinformatics* (2012) 13(1):31. doi: 10.1186/1471-2105-13-31. PubMed PMID: 22333067;
PubMed Central PMCID: PMCPMC3471323.
809

Synthetic standards combined with error and bias correction improves the accuracy and quantitative resolution of antibody repertoire sequencing in human naïve and memory B cells

Simon Friedensohn*[1], John M. Lindner*[2], Vanessa Cornacchione[2], Mariavittoria Iazeolla[2], Enkelejda Miho[1], Andreas Zingg[1], Simon Meng[1], Elisabetta Traggiai[2], and Sai T. Reddy[1]

[1]*Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland*
[2]*Novartis Institutes for BioMedical Research, Basel, Switzerland*
*equal contribution

Main Figures
Supplementary Material

Figure 1. A comprehensive set of human synthetic spike-in standards for Ig-seq. **(A)** Schematic showing the prototypical spike-in with the following regions (5' to 3'): a conserved non-coding region, ATG start codon, IGHV region with FR1 specific for multiplex-PCR, IGHJ regions, non-coding synthetic sequence identifier (specific for ddPCR probes), and downstream heavy chain constant domain sequences (IGHG, IGHM, IGHA) containing primer binding sites used for cDNA synthesis. Each spike-in contains a complete VDJ open reading frame, including nucleotides upstream of the ATG start codon, and downstream constant domain sequences containing primer binding sites used for sample cDNA synthesis. **(B)** Pairwise comparisons based on a.a. Levenshtein edit distance of all 85 standard CDR3 sequences. The germline IGHV and IGHJ segment family usage and IgG subclasses are denoted. 39 spike-ins contain rationally designed nt SHM (black circles) across the IGHV regions.

Figure 2. Library preparation of immunoglobulin (Ig) heavy chain genes for high-throughput sequencing (Ig-seq) using molecular amplification fingerprinting (MAF). **(A)** In step 1, reverse transcription (RT) is performed to generate first-strand cDNA with a gene-specific (IgM or IgG) primer which includes a unique reverse molecular identifier (RID) and partial Illumina adapter (IA) region. This results in single-molecule labeling of each cDNA with an RID. In step 2, several cycles of multiplex-PCR are performed using a forward primer set with gene-specific regions targeting heavy chain variable (V_H) framework region 1 (FR1), with overhang regions comprised of a forward unique molecular identifier (FID) and partial IA. In step 3, singleplex-PCR is used to extend the partial IAs. The result (Step 4) is the generation of antibody amplicons with FID, RIDs, and full IA ready for Ig-seq and subsequent MAF-based error and bias correction. **(B)** List of oligonucleotides sequences annealing to the V_H FR1 used in multiplex-PCR (Step 2) of the MAF library preparation protocol. The nearest germline IGHV segment(s) likely to be amplified by the respective primer are listed in the rightmost column. **(C)** The estimated melting temperature distribution of the V_H FR1 forward primer set.

Figure 3. Synthetic standards used to validate performance of error and bias correction of Ig-seq data by MAF. **(A)** The number of erroneous CDR3 sequences (at least one a.a. difference from the correct CDR3) per 100,000 reads is plotted against the relative concentration of each standard (from a master stock). Color-coded diamonds correspond to germline IGHV segment family of the respective standard and show the number of erroneous variants in uncorrected (raw) data; gray diamonds indicate the number of variants remaining after MAF error correction. **(B)** The number of erroneous VDJ variants derived from each standard was calculated by finding all variants that carried the correct CDR3 a.a. sequence, but differed by at least one nt across the entire VDJ region. Colored diamonds represent uncorrected data; gray diamonds indicate variants remaining after MAF error correction. **(C)** Sequencing bias introduced by multiplex-PCR using the FR1 primer set was assessed by plotting the measured frequencies of each standard against its relative concentration (from a master stock). Dashed line represents a bias-free ideal scenario ($R^2 = 1$). The left and right plots show observed frequencies before and after MAF bias correction, respectively. **(D)** Phylogenetic trees visualizing the CDR3 a.a. variants present for a selected standard with the CDR3 a.a. sequence *CARGINGERALEW* and IGHV1-8 and IGHJ1 segment usage. Prior to error correction, 39 erroneous CDR3 a.a. variants (branches) and 218 VDJ nt variants (black circles) were observed. Following MAF error correction, only the original correct CDR3 a.a. and two VDJ nt variants remain. The Ig-seq data sets used in A-C consisted of ~300,000 preprocessed full-length antibody reads from each of the synthetic spike-in only samples. IgG1_D1 dataset was used for panel **(D)** (see **Table S3**).

Figure 4. Ig-seq analysis of human naïve (CD27⁻IgM⁺) and memory (CD27⁺IgG⁺) B cells. **(A)** The flow cytometric workflow for isolating CD27⁻IgM⁺ and CD27⁺IgG⁺ B cells from peripheral blood. Boxed-in values indicate the frequency of each sorted subset as a percentage of the total B cell (CD19⁺) population from each of three donors (1-3 from left to right). **(B)** Experimental and Ig-seq based quantitation of antibody diversity; points represent cDNA molecule counts (using ddPCR) or unique reads (before and after MAF error correction) from cellular replicates (with mean and standard deviation shown) isolated from each donor and B cell subset. Unique read counts were based on the VDJ nt seqeunce. Dashed line represents the number of B cells isolated per cellular replicate (2×10^5 cells). **(C)** Phylogenetic trees illustrating CDR3 a.a. and nt variants present for the selected clonotype with the consensus CDR3 sequence *CARAAGSQYYYMDVW* and IGHV1-8 to IGHJ1 recombination. Prior to error correction, 70 erroneous CDR3 a.a. variants and 249 VDJ nt variants (black circles) were observed. Following MAF error correction, only 6 CDR3 a.a. and 15 VDJ nt variants remain. The Ig-seq data sets used in **(B)** are described in **Table S3**; IgG1_D1 was used for the tree in panel **(C)**.

Figure 5. Ig-seq analysis of molecular features highlight global differences between naïve (CD27⁻IgM⁺) and memory (CD27⁺IgG⁺) B cells. **(A)** Clonotype size (calculated as the total number of variants within a clonotype) for naïve IgM (blue lines) and memory IgG (red lines) B cells. Each pair of lines represents a single donor. **(B)** Graph showing the distribution of average SHM frequencies for VDJ nt variants per clonotype. The CD27⁻IgM⁺ B cell repertoires (blue lines) have a median SHM value of zero, whereas only a small fraction of clonotypes (approximately 8%) contain an average of one or more mutations. CD27⁺IgG⁺ repertoires (red lines) have a median of 20 to 24 SHM per nt variant within each clonotype. **(C)** CDR3 a.a. length distribution across clonotypes from naïve (blue lines) and memory (red lines) B cell subsets.

Figure 6. Clonotype diversity analysis across cellular replicates of naïve (CD27⁻IgM⁺) and memory (CD27⁺IgG⁺) B cells. **(A)** Venn diagrams show the presence of clonotypes (80% CDR3 a.a. similarity to least one clone in the cluster, same CDR3 a.a. length, same IGHV and IGHJ gene segment usage) across cellular replicates (2 x 10⁵ cells each) from each donor B cell subset. **(B)** Bar graph showing the fraction of clonotypes containing at least one variant with at least one nt SHM. Blue and red bars indicate clonotypes identified either in only one aliquot (unique) or in several aliquots (shared), respectively. The p-values represent significance using Fisher's exact test. **(C)** Species accumulation curves for CD27⁺IgG⁺ B cells: the number of newly discovered clonotypes from each additional cellular replicate (black circles) is plotted. Extrapolating the observed overlap provides an estimate for the total number of distinct clonotypes (Chao2 estimator: D1 = 164,268 ± 2,365; D2 = 38,034 ± 1,302; D3 = 76,904 ± 1,409) and the approximate amount of cellular replicates needed to discover all clonotypes present in the peripheral blood CD27⁺ IgG⁺ population.

Figure 7. Genetic features of naïve and IgG memory BCR repertoires. **(A)** Block map shows IGHV gene segment usage sorted by family (color-coded) across donors and B cell subset; blocks are normalized to the total number of clonotypes within each group. **(B)** IgG subclass usage in CD27+IgG+ donor repertoires. IgG4 sequences (dark blue bars) were virtually absent among three donors; a small fraction of sequences could not be unambiguously mapped based on the sequencing read (gray bars). **(C)** Venn diagrams showing the overlap of clonotypes (80% CDR-H3 amino acid similarity to least one clone in the cluster, same CDR3 a.a. length, same IGHV and IGHJ gene segment usage) between the naïve (CD27-IgM+) and memory (CD27+IgG+) BCR heavy chain repertoire in each of three donors. **(D)** Each plot shows the clonal composition of each shared clonotype (from panel **(C)**) in terms of its IgG and IgM intraclonal variants. Red triangles indicate clonotypes found in multiple IgM cellular aliquots; blue triangles show clonotypes which could only be found in one IgM cellular aliquot. The total number of clonotypes found are depicted in the corresponding contingency table. Fisher's exact test was used to quantitatively analyze enrichment of expanded IgM clonotypes in the shared IgM/IgG subset. **(E)** Ternary plots comprised of three axes representing the IgG1, IgG2, and IgG3 isotype subclasses. The relative subclass composition of intraclonal variants per IgG clonotype (each represented by a circle colored according to the number of variants belonging to that clonotype) is depicted by the position of the circle within the triagonal space. Red circles represent the average subclass composition for all IgG variants of each donor (cf. **Figure 6C**).

CDR3 amino acid sequence	CDR3 length	Variable (V) gene segment	Joining (J) gene segment	VH family	JH	IgG subclass	total mismatches	FR1 mismatches	CDR1 mismatches	FR2 mismatches	CDR2 mismatches	FR3 mismatches	spike-in pool contribution (%)
ARYANDREASY	11	IGHV3-23*01	IGHJ4*01	VH3	JH4	IgG3	0	0	0	0	0	0	7.00
ARCAFELATTE	11	IGHV3-23*01	IGHJ6*01	VH3	JH6	IgG2	9	0	4	0	5	0	0.15
ARSIMNFRIED	11	IGHV3-23*01	IGHJ1*01	VH3	JH1	IgG3	9	0	4	1	3	1	1.11
ARDRSAIREDDY	12	IGHV4-34*01	IGHJ4*01	VH4	JH4	IgG4	0	0	0	0	0	0	0.21
ARCRAIGVENTER	13	IGHV4-34*12	IGHJ6*01	VH4	JH6	IgG3	6	0	2	0	3	1	5.92
ARCHEETAHTEN	12	IGHV4-34*12	IGHJ3*01	VH4	JH3	IgG2	8	0	4	0	4	0	0.57
ARTHERESAPESCH	14	IGHV1-69*06	IGHJ4*01	VH1	JH4	IgG3	0	0	0	0	0	0	0.18
ARSSQIRREL	10	IGHV1-69*06	IGHJ6*01	VH1	JH6	IgG4	5	0	1	1	3	0	0.80
ARTARIKKHAN	11	IGHV1-69*06	IGHJ1*01	VH1	JH1	IgG3	10	0	4	0	5	1	3.04
ARCAMELLIATREE	14	IGHV1-18*01	IGHJ4*01	VH1	JH4	IgG2	0	0	0	0	0	0	0.25
ARCRISTINAPA	12	IGHV1-18*01	IGHJ6*01	VH1	JH6	IgG3	10	0	4	3	2	1	0.68
ARTETRACYCLINE	14	IGHV1-18*01	IGHJ3*01	VH1	JH3	IgG4	12	0	4	4	4	0	2.57
ARNICEECHIDNA	13	IGHV4-61*08	IGHJ4*01	VH4	JH4	IgG3	0	0	0	0	0	0	0.21
ARGIANTPANDA	12	IGHV4-61*08	IGHJ6*01	VH4	JH6	IgG2	7	0	3	0	4	0	0.94
ARGIRAFFEAFFE	13	IGHV4-61*08	IGHJ1*01	VH4	JH1	IgG3	15	0	4	3	8	0	3.04
ARIMPALAPALA	12	IGHV4-59*01	IGHJ5*01	VH4	JH5	IgG4	0	0	0	0	0	0	0.25
ARRATSEATMICE	13	IGHV4-59*01	IGHJ4*01	VH4	JH4	IgG3	10	0	4	0	4	2	0.48
ARRATTLESNAKE	13	IGHV4-59*01	IGHJ3*01	VH4	JH3	IgG2	10	1	5	0	3	1	2.17
ARREINDEER	10	IGHV3-30*03	IGHJ4*01	VH3	JH4	IgG3	0	0	0	0	0	0	0.18
ARSALAMANDER	12	IGHV3-30*03	IGHJ6*01	VH3	JH6	IgG3	9	0	3	2	4	0	1.32
ARTIGERSEVEN	12	IGHV3-30*03	IGHJ1*01	VH3	JH1	IgG3	12	0	4	0	5	3	2.17
ARTREESHREW	11	IGHV4-39*01	IGHJ4*01	VH4	JH4	IgG2	0	0	0	0	0	0	0.29
ARMANHATTAN	11	IGHV4-39*01	IGHJ6*01	VH4	JH6	IgG4	5	0	1	0	4	0	0.94
ARIRISHCAFE	11	IGHV4-39*01	IGHJ3*01	VH4	JH3	IgG4	11	2	3	2	4	0	1.84
ARNEWWHISKEY	12	IGHV3-48*02	IGHJ4*01	VH3	JH4	IgG3	0	0	0	0	0	0	0.35
ARSWEETWINE	11	IGHV3-48*02	IGHJ6*01	VH3	JH6	IgG2	7	0	3	0	4	0	1.56
ARGINMADRAS	11	IGHV3-48*02	IGHJ2*01	VH3	JH2	IgG3	11	0	4	1	6	0	1.84
ARAPPLETINIS	12	IGHV1-3*02	IGHJ4*01	VH1	JH4	IgG4	0	0	0	0	0	0	0.35
ARCAIPIRINHA	12	IGHV1-3*02	IGHJ4*01	VH1	JH4	IgG3	6	0	2	0	4	0	1.32
ARNECTARINE	11	IGHV3-21*01	IGHJ1*01	VH3	JH1	IgG2	0	0	0	0	0	0	0.41
ARPHYSALIS	10	IGHV3-21*01	IGHJ2*01	VH3	JH2	IgG3	10	0	5	0	3	2	0.80
ARAAMARANTH	11	IGHV1-2*02	IGHJ6*01	VH1	JH6	IgG4	0	0	0	0	0	0	0.57
ARGRAPESALAD	12	IGHV1-2*02	IGHJ4*01	VH1	JH4	IgG3	11	0	7	0	3	1	0.21
ARFIDDLEHEAD	12	IGHV4-31*02	IGHJ4*01	VH4	JH4	IgG2	0	0	0	0	0	0	0.48
ARFRIEDGARLIC	13	IGHV4-31*02	IGHJ1*01	VH4	JH1	IgG3	8	0	2	0	6	0	0.15
ARFATCHICKEN	12	IGHV3-33*01	IGHJ2*01	VH3	JH2	IgG4	0	0	0	0	0	0	0.48
ARDRIEDSEAWEED	14	IGHV3-33*01	IGHJ6*01	VH3	JH6	IgG3	8	0	3	0	1	4	1.56
ARSWISSCHARD	12	IGHV5-51*01	IGHJ4*01	VH5	JH4	IgG2	0	0	0	0	0	0	0.29
ARWATERCRESS	12	IGHV5-51*01	IGHJ4*01	VH5	JH4	IgG3	2	0	2	0	0	0	1.11
AREGGPLANT	10	IGHV1-46*01	IGHJ1*01	VH1	JH1	IgG4	0	0	0	0	0	0	0.41
ARARGENTINA	11	IGHV1-46*01	IGHJ6*01	VH1	JH6	IgG3	5	0	3	0	1	1	0.18
ARAEINSTEIN	11	IGHV3-7*01	IGHJ4*01	VH3	JH4	IgG2	0	0	0	0	0	0	0.29
ARKASACHSTAN	12	IGHV3-7*01	IGHJ4*01	VH3	JH4	IgG3	8	0	4	0	4	0	0.94
ARAPICCARD	10	IGHV4-38-2*02	IGHJ1*01	VH4	JH1	IgG2	0	0	0	0	0	0	0.25
ARNETHERLANDS	13	IGHV4-38-2*02	IGHJ6*01	VH4	JH6	IgG3	6	0	1	1	4	0	0.80
ARSELFIESTICK	13	IGHV3-11*01	IGHJ4*01	VH3	JH4	IgG2	0	0	0	0	0	0	0.41
ARPEKINGENTE	12	IGHV3-11*01	IGHJ3*01	VH3	JH3	IgG3	6	0	3	1	2	0	1.32
ARCHAIMASALA	12	IGHV1-8*01	IGHJ1*01	VH1	JH1	IgG2	2	0	0	2	0	0	0.21
ARGINGERALE	11	IGHV1-8*01	IGHJ6*01	VH1	JH6	IgG3	6	1	0	4	1	0	0.68
ARGREATREDMEAT	14	IGHV3-66*03	IGHJ5*01	VH3	JH5	IgG2	0	0	0	0	0	0	0.68
ARRACLETTE	10	IGHV3-66*03	IGHJ3*01	VH3	JH3	IgG3	15	0	3	2	6	4	0.25
ARAMYSMART	10	IGHV3-30*04	IGHJ1*01	VH3	JH1	IgG2	0	0	0	0	0	0	0.29
ARENDLESSWAY	12	IGHV3-30*04	IGHJ6*01	VH3	JH6	IgG3	11	1	4	4	2	0	0.80
ARLANADELREY	12	IGHV2-5*01	IGHJ4*01	VH2	JH4	IgG2	0	0	0	0	0	0	4.24
ARCDELEVIGNE	13	IGHV2-5*01	IGHJ3*01	VH2	JH3	IgG3	3	0	3	0	0	0	1.11
ARTHEISLAND	11	IGHV3-64*02	IGHJ2*01	VH3	JH2	IgG3	0	0	0	0	0	0	3.59
ARRFEDERER	10	IGHV3-64*02	IGHJ6*01	VH3	JH6	IgG3	8	0	4	1	3	0	0.57
ARVANILLACAKE	13	IGHV4-4*07	IGHJ4*01	VH4	JH4	IgG3	0	0	0	0	0	0	0.35
ARWALLSTREET	12	IGHV4-4*07	IGHJ3*01	VH4	JH3	IgG3	7	0	2	0	5	0	1.11
ARGRAYALDER	11	IGHV3-15*01	IGHJ2*01	VH3	JH2	IgG3	0	0	0	0	0	0	1.84
ARCHERRYPIE	11	IGHV3-15*01	IGHJ6*01	VH3	JH6	IgG3	4	0	0	2	2	0	0.25
ARSPICYSHRIMP	13	IGHV2-26*01	IGHJ5*01	VH2	JH5	IgG3	0	0	0	0	0	0	0.68
ARFRESHTHYME	12	IGHV2-26*01	IGHJ3*01	VH2	JH3	IgG3	6	0	4	0	2	0	0.41
ARTRAVELLING	12	IGHV3-74*01	IGHJ2*01	VH3	JH2	IgG3	0	0	0	0	0	0	0.94
ARFMAGELLAN	11	IGHV3-74*01	IGHJ6*01	VH3	JH6	IgG3	7	0	1	2	4	0	0.35
ARSWISSNESS	11	IGHV2-70*13	IGHJ1*01	VH2	JH1	IgG3	0	0	0	0	0	0	0.15
ARMILKSHAKE	11	IGHV4-30-4*01	IGHJ2*01	VH4	JH2	IgG3	0	0	0	0	0	0	0.48
ARGREENPEPPER	13	IGHV7-4-1*02	IGHJ4*01	VH7	JH4	IgG3	0	0	0	0	0	0	0.57
ARANGELFALLS	12	IGHV3-53*01	IGHJ6*01	VH3	JH6	IgG3	0	0	0	0	0	0	0.29
ARDEATHVALLEY	13	IGHV6-1*01	IGHJ1*01	VH6	JH1	IgG3	0	0	0	0	0	0	0.41
ARVARANASI	10	IGHV4-30-2*03	IGHJ3*02	VH4	JH3	IgG3	0	0	0	0	0	0	0.35
ARSTARWARS	10	IGHV4-28*01	IGHJ4*01	VH4	JH4	IgG3	0	0	0	0	0	0	4.24
ARFISHNCHIPS	12	IGHV3-9*01	IGHJ6*01	VH3	JH6	IgG3	0	0	0	0	0	0	5.92
ARITSATEST	10	IGHV3-20*01	IGHJ2*01	VH3	JH2	IgG3	0	0	0	0	0	0	2.57
ARHERAKLES	10	IGHV1-24*01	IGHJ3*01	VH1	JH3	IgG3	0	0	0	0	0	0	4.24
ARWILLIAMKELT	13	IGHV3-49*03	IGHJ5*01	VH3	JH5	IgG3	0	0	0	0	0	0	3.04
ARMINERVATHECAT	15	IGHV1-69-2*01	IGHJ6*01	VH1	JH6	IgG3	2	0	0	0	2	0	1.56
ARTARDIGRADA	9	IGHV5-10-1*02	IGHJ2*01	VH5	JH2	IgG3	0	0	0	0	0	0	1.32
ARRFRANKLIN	11	IGHV1-58*02	IGHJ3*01	VH1	JH3	IgG3	0	0	0	0	0	0	0.80
ARSELACHII	10	IGHV3-72*01	IGHJ4*01	VH3	JH4	IgG3	0	0	0	0	0	0	0.57
ARTEHRANINIRAN	14	IGHV3-73*02	IGHJ6*01	VH3	JH6	IgG3	0	0	0	0	0	0	0.18
ARDAMAVAND	10	IGHV3-13*01	IGHJ2*01	VH3	JH2	IgG3	0	0	0	0	0	0	0.21
ARNEMERTEA	10	IGHV1-45*02	IGHJ4*01	VH1	JH4	IgG3	0	0	0	0	0	0	0.48
ARKANAMYCIN	11	IGHV3-43*01	IGHJ4*01	VH3	JH4	IgG3	0	0	0	0	0	0	0.68
ARTRAGGIAI	10	IGHV7-81*01	IGHJ6*01	VH7	JH6	IgG3	0	0	0	0	0	0	0.48

Supplementary table 1. Molecular characteristics of 85 synthetic human IgH gene standards ('spike-ins') used in this study. See Figure 1A for spike-in construction schematic.

Sample	Total RNA extracted (ng/uL)	Total RNA (total amount)	Average number of cDNA transcripts per uL	Total number of cDNA transcripts	Sequencing depth (number of reads prior to pre-procssing)	Number of reads after pre-processing
IgM1_D1	6.6	132	5'109	62'581	676'557	403'271
IgM2_D1	59.1	1182	4'691	57'469	781'452	508'998
IgM3_D1	10.1	202	3'503	42'916	2'236'338	1'039'415
IgM4_D1	30.6	612	405	4'955	792'330	437'560
IgM5_D1	19	380	3'293	40'333	1'519'422	529'933
IgM1_D2	3.7	74	3'390	41'528	1'925'305	1'111'250
IgM2_D2	53.3	1066	4'330	53'043	673'482	379'811
IgM3_D2	34.4	688	2'048	25'088	992'456	488'230
IgM4_D2	2.2	44	3'075	37'669	981'274	452'817
IgM5_D2	47.4	948	3'855	47'220	2'279'624	440'083
IgM1_D3	7.7	154	4'631	56'734	545'947	333'492
IgM2_D3	29.5	590	3'568	43'702	1'448'340	636'432
IgM3_D3	2.6	52	2'860	35'035	649'626	407'914
IgM4_D3	7.2	144	3'567	43'696	986'419	529'860
IgG1_D1	61.2	1224	1'277	15'643	4'282'524	1'005'440
IgG2_D1	8.7	174	1'289	15'784	2'513'370	2'099'114
IgG3_D1	12.1	242	1'549	18'969	720'804	635'650
IgG4_D1	9.4	188	1'577	19'312	1'057'258	919'252
IgG5_D1	30.7	614	2'884	35'325	1'497'219	1'306'234
IgG1_D2	52.4	1048	421	5'157	1'049'571	913'558
IgG2_D2	3.7	74	181	2'211	959'085	700'808
IgG3_D2	8.5	170	268	3'277	910'336	793'379
IgG4_D2	6.5	130	214	2'622	747'740	662'657
IgG5_D2	5.6	112	386	4'722	998'038	826'427
IgG1_D3	7	140	1'552	19'012	604'835	490'991
IgG2_D3	5.4	108	683	8'367	804'122	658'452
IgG3_D3	5.9	118	972	11'901	756'943	649'224
IgG4_D3	6.9	138	1'639	20'072	2'724'771	1'179'968

Sample	Aligned reads (consensus build)	Raw CDR3 variants (AA)	Raw CDR3 variants w.o. singletons (AA)	Raw total variants (whole VDJ, nt)	MAF corrected CDR3 variants (AA)	MAF corrected clonotypes (Same V/J Gene, same	MAF corrected variants (whole VDJ, nt)	Donor
IgM1_D1	201'889	40'584	19'815	109'404	14'891	14'629	18'120	D1
IgM2_D1	260'867	37'987	14'952	129'770	11'257	11'088	13'629	D1
IgM3_D1	393'067	47'447	17'357	155'913	11'759	11'639	14'106	D1
IgM4_D1	203'127	46'001	21'116	128'936	15'139	14'919	18'001	D1
IgM5_D1	175'215	32'357	14'870	96'251	11'375	11'185	14'088	D1
IgM1_D2	534'227	65'874	24'260	234'442	18'087	17'646	23'645	D2
IgM2_D2	214'469	37'837	16'146	117'594	12'255	12'055	14'567	D2
IgM3_D2	244'601	35'706	14'292	120'009	10'643	10'439	13'289	D2
IgM4_D2	204'029	37'669	16'697	111'390	12'491	12'269	15'254	D2
IgM5_D2	135'195	26'236	12'557	66'889	9'397	9'292	10'797	D2
IgM1_D3	179'806	32'421	12'987	97'529	9'960	9'808	12'518	D3
IgM2_D3	285'749	42'979	18'424	138'162	14'098	13'866	17'610	D3
IgM3_D3	197'051	33'712	15'530	101'684	11'748	11'567	14'455	D3
IgM4_D3	248'200	31'542	12'514	103'246	9'693	9'582	11'799	D3
IgG1_D1	207'903	32'401	9'126	82'538	7'161	6'756	8'963	D1
IgG2_D1	1'484'045	159'576	36'883	520'865	8'549	7'744	13'393	D1
IgG3_D1	489'704	83'795	19'308	246'780	10'986	9'957	15'014	D1
IgG4_D1	712'466	105'073	24'949	322'613	11'320	10'286	17'581	D1
IgG5_D1	1'025'422	162'369	38'540	489'854	18'690	16'575	28'718	D1
IgG1_D2	644'950	56'789	13'373	199'164	3'233	3'011	4'954	D2
IgG2_D2	429'902	24'874	6'368	80'817	1'650	1'456	2'767	D2
IgG3_D2	541'960	38'086	8'957	134'139	1'965	1'849	2'895	D2
IgG4_D2	472'428	25'158	5'943	83'147	1'398	1'288	2'043	D2
IgG5_D2	566'689	40'047	9'394	132'147	2'501	2'323	3'603	D2
IgG1_D3	367'556	74'198	16'725	207'180	10'665	9'808	14'843	D3
IgG2_D3	485'151	55'285	12'829	182'680	4'559	4'266	6'327	D3
IgG3_D3	502'248	60'245	14'155	206'903	5'430	5'082	7'687	D3
IgG4_D3	693'622	96'133	22'466	325'211	9'239	8'556	14'028	D3

Supplementary table 2. Overview over our experimental results

Primer	Sequence	Notes
IgG_1r	TTGGCACCCGAGAATTCCACTGHHHHHACAHHHHHACAHHHHHNATTGTTCTGGGAAGTAGTCCTTGACCAG	Red part indicates primer binding site, black part contains unique identifier and overhang
IgM_1r	TTGGCACCCGAGAATTCCACTGHHHHHACAHHHHHACAHHHHNATTACGAGGGGGAAAAGGGTTGG	see above
VH1a	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGGCAGCTGGTGCAGTCTGGGG	see above
VH1b	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTCAGCTGGTGCAGTCTGGAG	see above
VH1c	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGCCAGCTTGTGCAGTCTGGGG	see above
VH1d	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGGCAGCTGGTGCAGTCTGGGC	see above
VH2a	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGCCCAGGTCACCTTGAAGGAGTCTG	see above
VH2b	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGCCCAGATCACCTTGAAGGAGTCTG	see above
VH3a	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGTGAGGTGCAGCTGGTGGAGTC	see above
VH3b	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGTGAAGTGCAGCTGGTGGAGTC	see above
VH3c	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGTCAGGTGCAGCTGGTGGAGTC	see above
VH3d	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGTGAGGTGCAGCTGGTGGAGAC	see above
VH4a	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGCAGCTGCAGGAGTCGGG	see above
VH4b	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTACAGCTGCAGGAGTCGGG	see above
VH4c	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGCAGCTGCAGGAGTCCGG	see above
VH4d	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGCAGCTACAGCAGTGGGG	see above
VH5	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGGCAGCTGGTGCAGTCTGGAG	see above
VH6	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTACAGCTGCAGCAGTCAGG	see above
VH7	CGTTCAGAGTTCTACAGTCCGACGATCHHHHACHHHHACHHHNGCAGTGCAGCTGGTGCAATCTGG	see above
PCR1_1r	ACTGGAGTTCCTTGGCACCCGAGAATTCCACT*G	'_*' indicates phosphorothioate bond
PCR2_f	AATGATACGGCGACCACCGAGATCTACACGTTCAGAGTTCTACAGTCCGACGAT*C	'_*' indicates phosphorothioate bond
PCR2_r	CAAGCAGAAGACGGCATACGAGATXXXXXXGTGACTGGAGTTCCTTGGCACCC	Red X indicates Illumina index
IgG_probe	/56-FAM/T+CTT+CCCC+C+TG+G/3IABkFQ/	'+_' indicate locked ribonucleic acids, '/56'-FAM = 5' 6-FAM (Fluorescein), '/3IABkFQ/' = 3' Iowa Black FQ
IgM_probe	/56-FAM/C+CCC+AA+CC+C+TTT/3IABkFQ/	'+_' indicate locked ribonucleic acids, '/56'-FAM = 5' 6-FAM (Fluorescein), '/3IABkFQ/' = 3' Iowa Black FQ
Spike_probe	/5HEX/CG+T C+T+G ACT +AGA +ACT +C/3IABkFQ/	'+_' indicate locked ribonucleic acids, '/56'-FAM = 5' HEX (Hexachlorofluorescein),'/3IABkFQ/' = 3' Iowa Black FQ
ddPCR_f	GGTCACYGTCTCYTCAG	
ddPCR_r	TGGCACCCGAGAATTC	

Supplementary table 3. Overview over all primers and probes used in this study

Figure S1. The sequencing bias of the multiplexed primer set was assessed by plotting the measured frequencies of each standard versus its actual, pipetted concentration in the pool. In an ideal case, the measured frequencies would fall onto the dashed line. The deviation from this line was used to calculate the R2 value, which decreases with greater deviation. (A) The upper panel shows the divergence of the measured frequencies obtained in the singleplex experiments versus pipetted concentration. Panels (B) and (C) show the measured frequencies and their deviations for one single experiment (IgG, Donor 1, Aliquot 1) before and after MAF correction.

Figure S2. (A) Repeatedly observed sequences within the first 5 cellular aliquots allow determination of the number of distinct clonotypes in the IgG CD27+ subset (Chao2 estimator, D1 = 164 268 ± 2 365, D2 = 38 034 ± 1 302, D3 = 76 904 ± 1 409), as well as the approximate amount of cDNA transcripts / RIDs required to discover all clonotypes. **(B)** Expanded view of panel A. **(C)** The average number of newly discovered clonotypes for each additional replicate (for both the IgM and the IgG subset).

Fig. S3. **Linear discriminant analysis distinguishes CD27+ IgG repertoires and CD27- IgM repertoires based on their V-Gene family usage.** (**A**) All aliquots from donors 1 and 2 were used to fit an LDA classifier based on the centered log ratio transformed V-Gene family frequencies. Afterwards, the aliquots from donor 3 are projected to the fitted component axis. Positive and negative values predict IgM and IgG repertoires, respectively. Arrows below the plot indicate the contribution of each V-Gene family to the prediction. Colored dots show true class membership and their positions are smoothed using kernel desnity estimators. Panels (**B**-**D**) shows the same procedure for different splits of the data.

Figure S4. Clonal composition of each clonotype that is shared between the IgG and IgM subsets in terms ofits IgG and IgM variants. The red, upward pointing triangle indicates clonotypes that are expaned in the IgM repertoire, whereas the blue triangle highlights clonotypes which could only be found in one IgM aliquot. Panels (**A**) and (**B**) show the results for donors 1 and 3, respectively.