Y chromosomal noncoding RNAs regulate autosomal gene expression via piRNAs in mouse testis

Hemakumar M. Reddy[1,2,16], Rupa Bhattacharya[1,3,16], Zeenath Jehan[1,4], Kankadeb Mishra[1,5], Pranatharthi Annapurna[1,6], Shrish Tiwari[1], Nissankararao Mary Praveena[1], Jomini Liza Alex[1], Vishnu M Dhople[1,7], Lalji Singh[8], Mahadevan Sivaramakrishnan[1,9], Anurag Chaturvedi[1,10], Nandini Rangaraj[1], Thomas Michael Shiju[1,11], Badanapuram Sridevi[1], Sachin Kumar[1], Ram Reddy Dereddi[1,12], Sunayana M Rayabandla[1,13], Rachel A. Jesudasan[1,14*, 15*]

[1]Centre for Cellular and Molecular Biology (CCMB), Uppal Road, Hyderabad, Telengana – 500007, India.

Present address:

[2]Brown University BioMed Division, Department of Molecular Biology, Cell Biology and Biochemistry, 185 Meeting Street room 260, Sidney Frank Life Sciences Building, Providence, RI 02912, USA.

[3]Pennington NJ 08534, USA.

[4]Department of Genetics and Molecular Medicines, Vasavi Medical and Research Centre, 6-1-91 Khairatabad, Hyderabad 500 004 India.

[5]Department of Cell Biology, Memorial Sloan Kettering Cancer Centre, Rockefeller Research Laboratory, 430 East 67th Street, RRL 445, New York, NY 10065, USA.

[6]Departments of Orthopaedic Surgery & Bioengineering, University of Pennsylvania, 376A Stemmler Hall, 36th Street & Hamilton Walk, Philadelphia, PA 19104.USA.

[7]Ernst-Moritz-Arndt-University of Greifswald Interfaculty Institute for Genetics and Functional Genomics, Department of Functional Genomics, Friedrich-Ludwig-Jahn-Straße 15 a, 17487 Greifswald, Germany.

[8]Deceased.

[9]Jubilant Biosystems Ltd., #96, Industrial Suburb, 2nd Stage, Yeshwantpur, Bangalore-560022, Karnataka, India.

[10]Department of Ecology and Evolution, University of Lausanne, Biophore Building, 1015, Lausanne, Switzerland.

[11]Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio 44120. USA

[12] Institute for Anatomy and Cell Biology, building-307, Heidelberg, Germany.

[13]Telangana Social Welfare Residential Degree college for Women, Suryapet-508213. Telangana. India.

[14]Department of Genetics, Osmania University, Hyderabad – 500007. Telangana. India.

[15]Inter University Centre for Genomics & Gene Technology, Karyavattom Campus, University of Kerala, Trivandrum, Kerala, India.

[16]These authors contributed equally to the work.

*Correspondence: Email ID: rachellike@gmail.com

Running title: **Y-ncRNAs regulate autosomal gene expression**

Key Words: Mouse Y chromosome; noncoding RNA; alternative splicing; piRNA; *Pirmy*; male sterility.

ABSTRACT

Majority of the genes expressed during spermatogenesis are autosomal. Mice with different

deletions of Yq show sub-fertility, sterility and sperm abnormalities. The connection

between Yq deletion and autosomal gene regulation is not well understood. We describe a

novel mouse Yq-derived long noncoding RNA, *Pirmy*, which shows unprecedented number

of splice variants in testis. Further, *Pirmy* transcript variants act as templates for several

piRNAs. We identified ten differentially expressed autosome-encoded sperm proteins in

mutant mice. *Pirmy* transcript variants have homology to 5'/3'UTRs of these deregulated

autosomal genes. Thus, subfertility in Y-deleted mice appears to be a polygenic

phenomenon that is partially regulated epistatically by the Y-chromosome. Our study

provides novel insights into possible role of MSY-derived ncRNAs in male fertility and

reproduction. Finally, sperm phenotypes from the Y-deleted mice seem to be similar to that

reported in inter-specific male-sterile hybrids. Taken together, this study provides novel

insights into possible role of Y-derived ncRNAs in male sterility and speciation.

INTRODUCTION

Y chromosome has come a long way from a single-gene male determining chromosome to

one that houses a few protein-coding genes besides sequences crucial for spermatogenesis

and fertility (Bellott et al., 2014; Jehan et al., 2007; Kuroda-Kawaguchi et al., 2001; Moretti

et al., 2017; Piergentili, 2010; Soh et al., 2014; Tiepolo & Zuffardi, 1976). Earlier studies have

shown that genes involved in sex determination and spermatogenesis are present on the

short arm. Several lines of evidence indicate that the male-specific region on the long arm of

the Y chromosome (MSYq) in mouse is replete with highly repetitive mouse-specific

sequences that are expressed in spermatids (Burgoyne & Mitchell, 2007; Conway et al.,

1994; Prado, Lee, Zahed, Vekemans, & Nishioka, 1992; Toure et al., 2005).

Previously published data have described two different strains of mice with partial deletions

of the long arm of the Y chromosome (Yq) (Conway et al., 1994; Józefa Styrna, Bili, &

Krzanowska, 2002). Mice from both the genetic backgrounds exhibit male sterile

phenotypes such as subfertility, sex ratio skewed towards females, reduced number of

motile sperms, aberrant sperm motility and sperm head morphological abnormalities

(Conway et al., 1994). Mice with partial deletions of Yq show sperm abnormalities with less

severe phenotype whereas mice with total deletion of the Yq have extensive sperm

morphological aberrations and are sterile (Toure et al., 2004). This suggested the presence

of multicopy spermiogenesis gene(s) on mouse Yq (Burgoyne, Mahadevaiah, Sutcliffe, &

Palmer, 1992; Conway et al., 1994). Subsequently multicopy transcripts such as Y353/B,

spermiogenesis-specific transcript on the Y *(Ssty)* and Sycp3-like, Y-linked *(Sly)* from the

mouse Yq were projected as putative candidate genes for male sterility and spermiogenesis

in mice (Cocquet et al., 2009; Conway et al., 1994; Riel et al., 2013; Toure et al., 2004). As

mice with partial deletions of Yq (XY^{RIII}qdel, 2/3rd interstitial deletion of Yq) show reduced

expression of *Ssty* and impaired fertility, this gene (present on Yq) was implicated in

spermatogenesis (Toure et al., 2004). The next major gene to be discovered on mouse Yq

was the multicopy *Sly*. As SLY interacts with a histone acetyl transferase and is an acrosomal

protein, the authors suggested that *Sly* could control transcription and acrosome functions

(Reynard, Cocquet, & Burgoyne, 2009). Further, Cocquet and colleagues observed major

problems in sperm differentiation when they disrupted functions of the *Sly* gene by

transgenic delivery of siRNA to the gene. Therefore, *Sly* was conjectured as a putative

candidate gene for spermiogenesis (Reynard et al., 2009). However, subsequently Ward and

colleagues showed that *Sly* expression alone is not sufficient for spermiogenesis (Riel et al.,

2019).

Vast majority of the genes required for spermatogenesis and spermiogenesis are non-Y-

linked (Choi et al., 2007; Schultz, Hamra, & Garbers, 2003; Xiao, Tang, Yu, Gui, & Cai, 2008).

Deletions of the Y-chromosome leading to different degrees of male infertility prompted us

to hypothesise interactions between Y-derived transcripts and autosomal genes in male

fertility. Earlier, studies in the lab reported a testis-specific chimeric transcript, generated by

trans-splicing between the *CDC2L2* mRNA from chromosome number 1 and a long

noncoding RNA from the Y chromosome in human (Jehan et al., 2007). We hypothesized

more such interactions between protein coding genes on autosomes and noncoding RNAs

from the Y chromosome. In this context, we studied a mutant mouse, which had a partial

deletion of the Y-chromosome heterochromatic block XY^{RIII}qdel (Conway et al., 1994) to look

for regulatory elements, if any, in the deleted region.

Previous studies in the lab identified 300-400 copies of a mouse Y chromosome specific

genomic clone, M34 (DQ907163.1) using slot blot, sequencing and bioinformatics analyses

(Bajpai, Sridhar, Reddy, & Jesudasan, 2007; Singh, Panicker, Nagaraj, & Majumdar, 1994).

There is a reduction of M34 copies in the XY^{RIII}qdel mice that exhibit multiple sperm

abnormalities. As deletions of the Y chromosome show sperm abnormalities, we reasoned

that these repeat sequences could have important functional role(s) in the multistep

developmental process of sperm production. The XY^{RIII}qdel mice also showed reduced

transcription of M34 compared to its wild type XY^{RIII} mice in testis. In order to understand

putative functions of this sequence, first of all we identified a transcript corresponding to

M34, *Pirmy,* from mouse testis. Subsequent experiments identified multiple splice variants

of this transcript. Parallel experiments identified deregulated proteins in the sperm

proteome of the XY^{RIII}qdel mice. Interestingly, genes corresponding to all these proteins

localized to different autosomes. Further, we showed that the UTRs of these genes bear

homology to piRNAs derived from *Pirmy*. Thus, our results demonstrate for the first time (i)

a novel noncoding RNA (*Pirmy*) on mouse Y long arm (ii) large number of splice variants of

Pirmy, and the generation of piRNAs from it in mouse testis and (iii) their putative role in

regulation of autosomal genes involved in male fertility and reproduction.

RESULTS

M34 is transcribed in mouse testis

To address the precise function of the M34 transcript we confirmed the localization of the

sex and species-specific repeat M34 on mouse Y long arm again by fluorescence in situ

hybridization (FISH) (Fig. 1*A*). BLAST analysis of M34 sequence against mouse whole genome

also showed maximum similarity to Y chromosome (>97% identity) with few hits on the X

(Fig. 1*B,* NCBI build 38.1). M34 was then analysed for expression in adult mouse testis. FISH

using M34 as a probe revealed abundant transcription in testis (Fig. 1*C*). Pretreatment with

RNase abolished these signals confirming the presence of RNA (Fig. 1*D*). Expression profiling

by FISH in testes showed the presence of M34 transcripts in 18-day embryos, newborns and

1-month old mice (30 days postpartum) (Fig. S1), suggesting transcription of this repeat in

mouse testis from early developmental stages.

Isolation of a novel polyadenylated noncoding transcript from mouse Y chromosome

Sequence analysis of the 9.51 kb M34 clone using Tandem Repeats Finder (TRF) and

RepeatMasker identified different simple sequence repeats and partial mid-repetitive

sequences like LINEs, SINEs and LTR elements which constitute ~35% of the total M34

sequence (Fig. 1*E*). A number of gene prediction programs like GENSCAN, GrailEXP, MZEF,

and GeneMark did not predict any genes within M34 with consistency. BLAST analysis of

M34 sequence against the EST database of NCBI (NCBI build 36.1) identified 5 ESTs at the 3'

end of M34 sequence (Fig. 1*E*). Expression of these ESTs was observed in embryos from at

least 13.5d onwards (data not shown). Two of these ESTs showed male-specific expression

by Reverse transcription PCR (RT-PCR) analysis.

In order to identify a cDNA corresponding to M34 in testis, one of the male specific ESTs

(CA533654) was used to screen a mouse testis cDNA library. A 1395 nt long polyadenylated

Y-specific cDNA was isolated and this was named *Pirmy* - piRNA from mouse Y chromosome

(DQ907162.1). FISH on to mouse metaphase spreads showed that *Pirmy* is present only on

the Y chromosome in multiple copies (Fig. 1*F*), similar to that of the genomic clone M34.

BLAST of *Pirmy* against the nucleotide database of NCBI picked up only mouse sequences

with statistically significant alignments (e-value <4e^{-04}). This suggests that *Pirmy* is specific to

mouse. BLAST analysis using *Pirmy* against the mouse genome plus transcriptome database

showed homology to *Sly* transcript in exons 1-5. The exons 2-5 were identical in *Pirmy* and

Sly (Fig. 1*G*). The region of homology is less than a quarter of *Pirmy*, and is confined to the

5'end. The entire sequence of *Pirmy* localizes to the Y chromosome at 4341127-4381724

(GRCm39). BLAST search against the nucleotide database identified a reference sequence

NR164186, which was annotated as a long noncoding RNA using evidence data for transcript

exon combination from *Pirmy*. The exon-intron junctions of *Pirmy* are shown in Figure 1H.

Splice variants of *Pirmy* in mouse testis

Pirmy was analyzed further by RT-PCR. Two rounds of amplification using primers to the two

ends of *Pirmy* yielded multiple products in testis and brain (Fig. 2*B*). PCR products were

cloned. Sanger sequencing of more than 1000 clones from testis yielded 107 transcripts

(NCBI accession numbers <u>FJ541075-FJ541181)</u>, besides the one obtained by screening testis-

cDNA library (Figs. 2A, 3). BLAST analysis of these transcripts against the NCBI genome

database (Build 39) showed that 28 of these transcripts (FJ541075- FJ541102; Fig. 3) localise

to multiple regions on the mouse Y chromosome. The remaining 79 transcripts (FJ541103-

FJ541181) and *Pirmy* were present at a single locus on the Y chromosome at 4341127-

4381724 (GRCm39) (Fig. 2). Thus, the intron-exon organization, found in *Pirmy* and the 79

splice variants is present only at a single locus on the Y chromosome. These 79 transcripts

could therefore be alternatively spliced isoforms of *Pirmy*. Each exon from *Pirmy* is present

in multiple copies on the Y chromosome. The 28 transcript variants reveal that different

combinations of these exons and introns are there at multiple loci on the Y. The splice

variants of *Pirmy* exhibited the full spectrum of splicing patterns like exon skipping,

alternative 5' and 3' splice sites, mutually exclusive alternative exons, intron retention and

combination of different splicing events (Fig. S2). The splice variants of *Pirmy* contained

consensus splice signal sequences at all the intron-exon junctions (Data sheet).

Splice variants of *Pirmy* in mouse brain

Many transcripts that are expressed in testis are also known to be expressed in brain;

therefore, we studied the expression of *Pirmy* in mouse brain also. RT-PCR amplification

(Fig. 2*B*), cloning and sequencing of *Pirmy* products from mouse brain yielded 12 transcripts.

All these transcripts localize to the single locus, to which *Pirmy* and its splice variants

localized (4341127-4381724 (GRCm39)). In fact, comparison of the exons of the splice

variants from the brain and testis showed that the same splice isoforms were present in

both the tissues (Fig. 4).

Expression of M34 in XY^{RIII}qdel mice

Metaphase spreads from the XY^{RIII}qdel mice showed a reduction in copy number of M34 on

the Y chromosome showing that it localizes to the deleted region of the Y chromosome in

the above mice (Fig. S3*B*). Therefore, expression of M34 was then checked in testis and

sperms of XY^{RIII} and XY^{RIII}qdel mice by FISH. Dramatic reduction in fluorescence intensity was

observed in testis and sperm of XY^{RIII}qdel mice. However, sperms from epididymis of both

XY^{RIII} and XY^{RIII}qdel mice showed faint fluorescence intensity (Fig. S3*B*). The subclones of

M34 (Fig. S3A) when used as probes on testis sections showed reduction in fluorescence

intensity in XY^{RIII}qdel mice (Fig. 3*C*). We also checked the copy number of *Pirmy* in genomic

DNA isolated from the wild type and XY^{RIII}qdel mice by Real-Time PCR using primers to exon

7; a significant reduction in copy number was observed in the XY^{RIII}qdel genome (Fig. S4C).

Many proteins coded by autosomal genes are deregulated in XY^{RIII}qdel sperm proteome

We then analysed the motility profile of sperms from XY^{RIII} and XY^{RIII}qdel mice and identified

a stark difference in motility patterns (Movies S1, S2 respectively). Spermatozoa from XY^{RIII}

mice show linear progressive motion whereas sperms from XY^{RIII}qdel mice show rapid

flagellar movement with non-linear and non-progressive motion. Most of the spermatozoa

from XY^{RIII}qdel mice stall at the same position with no linear displacement. Studies from

Burgoyne's lab reported morphological abnormalities in sperms from XY^{RIII}qdel mice

(Conway et al., 1994). In order to understand the connection between the Y-deletion and

sperm abnormalities, we performed comparative sperm proteome analysis between normal

mice and the XY^{RIII}qdel mice by 2D-PAGE and mass spectrometry using protocols

standardized in the laboratory (Bhattacharya, Devi, Dhople, & Jesudasan, 2013).

This analysis identified 8 protein spots that were differentially expressed in the pI ranges of

4-7 and 5-8 (Fig. 5*A*). Surprisingly four of these i.e. calreticulin (D1), Cu/Zn superoxide

dismutase (SOD (D4)), fatty acid binding protein 9 (FABP9 (D5)), Serine Peptidase Inhibitor

(Kazal type II (SPINK2)/Acrosin-Trypsin inhibitor variant 2 (D2) were upregulated in XY^{RIII}qdel

sperms compared to XY^{RIII} sperms (Fig. 5*A*). A novel shorter isoform of SPINK2 - SPINK2

variant3 (D3, Q8BMY) which was shorter by 27 amino acids at the N-terminal end was

downregulated in XY^{RIII}qdel sperms (Fig. 5*A*, Fig. S4B). Three proteins were not detectable in

XY^{RIII}qdel sperms in the pI range of 5-8 (Fig. 5A). Two of these were reported as hypothetical

proteins in the NCBI database (1700001L19 Riken cDNA and 1700026L06 Riken cDNA. As we

have identified these proteins, they have now been deposited in the Uniprot database with

accession numbers Q9DAR0 and, Q7TPM5 (MAST (Bhattacharya et al., 2013) respectively.

The third one was Stromal cell derived factor 2 like 1 (SDF2L1 (B)). Expression of four of the

eight differentially expressed proteins was also confirmed by western blotting in testis and

sperms (Fig. 5C).

Calreticulin, SOD and FABP9 showed upregulation in XY^{RIII}qdel sperms by both 2D PAGE and

western blot analyses; MAST was not detectable by both the techniques in sperms. Protein

expression of calreticulin, SOD, FABP9 and MAST did not vary significantly between testes of

XY^{RIII} and XY^{RIII}qdel mice. Thus, in sperms from the XY^{RIII}qdel, four proteins were upregulated

while one was downregulated. Three proteins were not visible in the XY^{RIII}qdel sperm

proteome. Surprisingly, all the eight genes corresponding to the differentially expressed

sperm proteins localized to different autosomes (Fig. 5B).

Next, we analysed the expression of the transcripts corresponding to the protein spots in

testis by Real-Time PCR/Northern blot analysis (Fig. S4A, B). Although the proteins Q9DAR0,

SDF2L1 and MAST were not detectable in sperms of X^{YRIII}qdel, the corresponding RNAs were

present in testis. The transcripts of SDF2L1, MAST, calreticulin and SPINK2 variant 2 proteins

were upregulated in XY^{RIII}qdel mice testis. In contrast transcripts of Q9DAR0 and Spink2

variant 3 did not show significant quantitative difference between the two (Fig. S4B).

UTRs of deregulated autosomal genes show homology to 108 transcripts

The fact that a few autosomal genes were deregulated when there was a deletion on the Y

chromosome prompted us to investigate the mechanism behind this puzzling observation.

We hypothesized that *Pirmy* transcript variants that show reduction in copy number in the

genome and reduced transcription in XY^{RIII}qdel, could be regulating autosomal gene

expression in testis. BLAST analysis of these transcripts against the 3' and 5' UTRs of the

deregulated genes revealed short stretches of homology ranging in size from 10-16

nucleotides, in either +/+ or +/- orientations. BLAST analysis of the 108 transcripts against

the UTRs of deregulated genes identified 21 different hits. Of these 11 were from *Pirmy*

splice variants and 10 from the other transcript variants respectively (Fig 6). There are as

many as 7 *Pirmy* hits in the 3' UTR of Q9DAR0 (Fig. 6). Homology between Y-derived

transcripts and UTRs of deregulated autosomal genes indicates interactions between genes

on the Y chromosome and autosomes in mouse testis. A BLAST against the entire transcripts

of the deregulated genes showed homology to the coding regions also; however, more

stringent BLAST parameters, showed homology to the UTRs alone.

Furthermore, we performed BLAST analysis of all the 108 transcripts (ncRNAs) against the

entire UTR database. This identified small stretches of homology in the UTRs of a number of

genes across different species. The homologous sequences (10-22 nt) localized to both

exons and exon-exon junctions of the 108 transcripts. Representation of some UTR

homologies are shown in Fig. 7A. We identified 372 unique homologous stretches in the 108

transcripts. Of these 302 (81.19%) localized to the exons and 70 (18.82%) to exon-exon

junctions (Fig. 7B). The autosomal genes bearing homology to these ncRNAs in their UTRs,

are expressed in multiple tissues including testis/epididymis (Table S1).

Identification of ~30nt RNAs from *Pirmy* transcript variants

To check if these short stretches of homologies corresponded to small RNAs, few

representative oligonucleotide sequences from the ncRNAs with homology to different

UTRs were used as probes (Fig. S5) on small RNA northern blots. Two genes were chosen

from the deregulated proteins Q9DAR0 (Spot A) and superoxide dismutase (SOD); Four

genes (butyrylcholinesterase (*Bche*), phospholipase A2, group XIIB *(PLA2G12B) and*

sialophorin *(Spn)*)were chosen from the BLAST output against UTR database. All the above

probes elicited approximately 26-30 nt long testis-specific signals, of the size of piRNAs (Fig.

7*C*).

To confirm that these homologous sequences are indeed piRNAs, different experiments

were designed. As the antiparallel strands of DNA are reported to express different levels of

piRNA, differential expression from the antiparallel strands were studied using Sense (S) and

antisense (AS) probes designed to homologous stretches in the 3' UTRs of *Sod* and *Bche*.

Identical experimental conditions showed differential expression of these 30 nt species of

RNAs from the two strands of DNA (Fig. 7*D*), further indicating that these short RNAs could

be piRNAs.

As piRNAs are PIWI/MIWI binding small RNAs, Electrophoretic Mobility Shift Assay (EMSA)

using the *Pirmy*-derived oligonucleotides and recombinant MIWI protein was done to check

if the sequences with homologies to UTRs of different genes are indeed piRNAs.

Representative gel shifts using oligonucleotides from UTRs of Q9DAR0 and *Sod* are depicted

in Fig. 7E and F respectively. piR1 (Girard, Sachidanandam, Hannon, & Carmell, 2006), a

known piRNA served as the positive control. Specificity of binding was indicated by the use

of corresponding cold competitors as described in the legend to Fig. 7E and F. The piRNA

derived oligonucleotides competed out binding of piR1 to MIWI protein and vice versa. This

confirmed that these oligonucleotides are indeed MIWI-binding RNAs and therefore piRNAs.

The mobility shift was also competed out by MIWI antibody while Argonaute 3 antibody did

not alter the mobility of the gel-shifted band obtained with MIWI indicating specificity of

binding. These experiments provide further evidence that these short RNA sequences are

piRNAs.

Pirmy **identifies piRNAs from NCBI Archives database**

Next line of evidence to the fact these short stretches of homologies are piRNAs came from

the NCBI Archives database for piRNAs. BLAST analysis using the 108 transcripts identified a

total of 1445 small RNAs in the Sequence Read Archives SRP001701 and SRP000623 with

>95% identity. These transcripts identified 352 MIWI2 - MILI-associated reads in SRP000623.

Of these 310 had homology to midrepetitive sequences (SINEs, LINEs and LTRs) and 5 had

homology to X chromosome. Thus, the remaining 37 piRNAs could be specifically derived

from the Y chromosome.

Antagopirs downregulate reporter gene expression

Complementary oligonucleotides synthesized to piRNA sequences present in UTRs of *Sod*

and *PLA2G12B* were designated as antagopirs (Fig. S5). Cloning of these UTRs 3' to the

Luciferase gene reduced its expression. Treatment with increasing concentrations of

antagopirs, i. e. 5 nM, 10 nM and 20 nM caused further concentration dependent reduction

in Luciferase expression (Fig. 7*H*). The antagopirs to *Sod* and *PLA2G12B* did not have an

effect when the UTR from a non-target gene (*Cdc2l1)* was cloned 3' to the Luciferase gene.

Fig. 7G is a schematic representation of 3'UTR reporter construct. Thus, the use of

antagopirs to piRNAs could modulate gene expression in vitro.

This study therefore, paved the way for a series of novel and exciting observations. We have

identified, a novel, polyadenylated lncRNA (*Pirmy*) expressed from mouse Yq in testis, that

shows the large number of splice variants in testis. These Y-derived transcripts harbour

piRNAs that have homology to the UTRs of a few autosomal genes expressed in mouse

testis. The proteins expressed from these autosomal genes are deregulated in sperms of Y-

deleted mice and appear to be regulated by piRNAs generated from MSYq derived lncRNAs.

Consolidation of the results from the proteomics analysis and the molecular studies

suggested that piRNAs generated from male-specific mouse Yq regulate expression of

multiple autosomal genes in testis. Partial deletion of Yq resulted in deregulation of these

proteins leading to sperm anomalies and subfertility. Thus, subfertility in mice appears to be

a polygenic phenomenon that is regulated epistatically by the Y chromosome.

DISCUSSION

Y-chromosomes harbour genes for male determination and male fertility. Yet the role of Y

chromosomal repeats in male fertility and spermatogenesis remains enigmatic. In this study,

we elucidate the role in male fertility of a species-specific repeat, M34, from mouse Y long

arm, which is transcribed in mouse testis. Transcription from repeats on mouse Y

chromosome has been reported earlier. Testis-specific transcription of a family of

poly(A)RNAs from the mouse Y chromosome was first reported by Bishop and Hatat using a

multicopy Y-derived probe, pY353/B (Bishop & Hatat, 1987; Conway et al., 1994).

Subsequently more transcripts were identified in mouse testis, using repeat sequences

localizing to mouse Y chromosome (Prado et al., 1992; Toure et al., 2005). The report of

Pirmy transcript variants described in this study adds to the repertoire.

The 108 *Pirmy* transcript variants in the present study were discovered serendipitously by

cloning and sequencing of the multiple RT-PCR products obtained using primers to the initial

and terminal exons. As the primers were restricted to just two of the exons, it is possible

that we might discover more transcript variants using primers to different combinations of

exons for RT-PCR amplification. Alternative splicing has been reported in ncRNAs (Pang et

al., 2005), yet, such extensive splicing as observed in our study has not been reported for

any of them. Very few polymorphically spliced genes have been described earlier from sex

chromosomes, particularly the Y (Ellis, Ferguson, Clemente, & Affara, 2007; Wegmann,

Dupanloup, & Excoffier, 2008). Thus, the 80 splice variants of *Pirmy* in this study appears to

be by far the maximum number of isoforms characterized from a single ncRNA by

alternative splicing. Elucidation of identical splice isoforms in mouse brain and testis in this

study (Figs. 2, 4) shows that these alternative splicing events are precise and not random

because the same splicing events take place in both the tissues. The consensus splice signal

sequences present at the intron-exon junctions in all the splice variants further reaffirm

programmed splicing events.

The identification of deregulated proteins, for which the corresponding genes localise to

autosomes, in a Y-deletion mutant was a surprise. Identification of small stretches of

homology in the UTRs of these transcripts from the *Pirmy* transcript variants established the

connection between the two. Observation of 27-32 nt long signals on small RNA northern

blots using the short homologous sequences as probes, which also bind to MIWI protein and

show differential expression from the antiparallel strands of DNA indicated that these

probes identify piRNAs. Some of these also identified piRNAs in the SRA databases. Small

RNAs of the size of ~27-32 nucleotides that bind PIWI protein are classified as piRNAs. These

are also known to be differentially transcribed from the two strands of DNA. The presence

of piRNAs in the UTRs of genes corresponding to deregulated proteins, suggests the

regulation of these autosomal genes by Y-derived piRNAs. The concentration dependent

reduction of Luciferase expression by antagopirs corroborates the regulation of these genes

by Y-derived piRNAs. Proteins from three other autosomal genes are also deregulated in

different strains of Y-deleted mice besides the ones identified in the proteomics screen;

caldendrin is upregulated in XY^{RIII}qdel sperm (Bhattacharya et al., 2013) and acrosin is

downregulated in B10.BR-Y^{del} sperm. (J. Styrna, Klag, & Moriwaki, 1991). Aromatase is

overexpressed in both the Y-deleted strains of mice i.e. XY^{RIII}qdel (our unpublished

observation), and B10BR-Y^{del} (Kotula-Balak, Grzmil, Styrna, & Bilinska, 2004). Therefore, it is

not surprising to find *Pirmy* homologous sequences in the UTRs of caldendrin, acrosin and

aromatase genes, suggesting Y- mediated regulation for these genes as well. Ellis and

colleagues also observed up or down regulation of genes from the X-chromosome and

autosomes in testes of mice with deletions of Y heterochromatin using a microarray

approach (Ellis et al., 2005). Homology between UTRs of some of the above genes and the

108 ncRNAs (Table S2) further strengthens the hypothesis of putative regulation of genes

located elsewhere in the genome by Y chromosomal repeats.

piRNAs are known to regulate gene expression at the levels of both transcription and

translation. Our study also suggests regulation at both the transcriptional and post-

transcriptional levels by the piRNAs derived from the 108 transcript variants. Genes

corresponding to three proteins that were missing from the mutant sperm proteome, were

indeed transcribed in testis. The fact that there is comparable expression of calreticulin and

SPINK2 variant2 proteins in XY^{RIII} and XY^{RIII}qdel testes by western blot analysis, despite

significantly upregulated expression of these transcripts in testis, could indicate regulation

at the level of both transcription and translation for these genes. piRNAs regulate

translation in early embryos and gonads besides containing transposable elements (Aravin,

Hannon, & Brennecke, 2007; Deng & Lin, 2002).

We also propose that the deregulated proteins identified in the current study are at least

partially responsible for the sperm phenotype observed in XY^{RIII}qdel mice. For example,

FABP9 that is upregulated in XY^{RIII}qdel spermatozoa, localizes to perforatorium, the

subacrosomal region in spermatozoa with falciform head shapes (Korley, Pouresmaeili, &

Oko, 1997; Oko & Morales, 1994; Pouresmaeili, Morales, & Oko, 1997). It is the most

abundant protein of the perforatorium, and putatively has a role in shaping the unique

sperm head structure of rodents (Oko & Morales, 1994; Pouresmaeili et al., 1997). The

deregulated expression of Spink2 isoforms also could contribute to sperm head

morphological abnormalities and reduced sperm motility; Kherraf and colleagues (Kherraf et

al., 2017) observed grossly misshapen sperm heads and reduced motility in Spink2 knock

out mice.

Calreticulin overexpressed in XY^{RIII}qdel spermatozoa is a calcium store associated with

sperm functions such as hyperactivated motility, capacitation and acrosome reaction

(Yanagimachi, 1982). The subsequent cascade of events could result in subfertility.

Caldendrin yet another protein that is upregulated in XY^{RIII}qdel sperm (Bhattacharya et al.,

2013), localizes to acrosome in rats and is considered to be a stimulus dependent regulator

of calcium (Redecker, Kreutz, Bockmann, Gundelfinger, & Boeckers, 2003). SPINK2 variant3

localizes to the acrosome in mouse spermatozoa (our unpublished results). The

physiological function of SPINK2 is believed to be blocking of deleterious degradation of

proteins released by acrosin from spermatozoa during acrosome reaction (Moritz, Lilja, &

Fink, 1991). Being a putative sperm acrosin inhibitor a role in fertility can be envisaged.

Sdf2l1 is an ER stress-inducible gene, induced by the unfolded protein response pathway

(Fukuda et al., 2001). The XY^{RIII}qdel spermatozoa may be more susceptible to stress induced

damages as they lack stress response protein, SDF2L1.

MAST a novel protein that is not visible in the XY^{RIII}qdel sperm proteome, but localizes to

both the acrosome and sperm tail, indicating a role in sperm motility and penetration of the

egg (Bhattacharya et al., 2013). Bioinformatic analysis of the novel protein Q9DAR0 predicts

it as a cilia related gene (McClintock, Glasser, Bose, & Bergman, 2008). Based on this

information putative role of this protein in sperm motility can be envisaged. SOD, a protein

upregulated in XY^{RIII}qdel, has been found to be positively associated with sperm count and

overall motility (Lu, Huang, & Lu, 2010). The irreversible conversion of androgens to

estrogens is catalyzed by aromatase transcribed by *Cyp19* gene (Simpson et al., 1994).

Biology behind the skewed sex ratio towards females in the progeny sired by XY^{RIII}qdel

males could be explained by the upregulated expression of aromatase in testes of these

mice. Acrosin plays a crucial role in acrosome exocytosis and egg zona pellucida penetration

(Yamagata et al., 1998). Spermatozoa lacking acrosin (acrosin$^{-/-}$) exhibit delayed fertilization

as both the processes of acrosome exocytosis and egg penetration are deferred (Adham,

Nayernia, & Engel, 1997; Mao & Yang, 2013).

Functions of Y chromosome have been elucidated using different deletions of the

chromosome in the past. Naturally occurring deletions in the euchromatic long arm of Y

chromosome in azoospermic men showed the involvement of this region in human male

infertility (Vogt, 1998). *Drosophila melanogaster* males with deletions of different regions of

the Y chromosome show absence of several sperm axoneme proteins (Goldstein, Hardy, &

Lindsley, 1982). Mice with partial or total deletions of Y heterochromatin show deregulation

of testicular gene expression and subfertility/sterility (Cocquet et al., 2009; Ellis et al., 2005).

Previous studies in the lab elucidated another example of an intronless Y-derived ncRNA

mediated regulation of an autosomal gene, *CDC2L2* via trans-splicing in human testis (Jehan

et al., 2007). The role of mouse Y heterochromatin in the current study therefore reveals a

novel pathway for the regulation of autosomal genes by Y chromosome, mediated by

piRNAs, in male reproduction. Therefore, consolidation of the observations from mouse and

human studies therefore shows that Y chromosome regulates autosomal genes expressed in

testis using distinct mechanisms in different species.

Comparative sperm proteomics analysis in our study portrays involvement of multiple

autosomal genes in subfertility. The regulation of autosomal gene expression appears to be

relaxed in sperms of Yq-deleted mice. This reflects a connection between the Y

chromosome and autosomes. In fact, as suggested by Piergentili, Y chromosome could be a

major modulator of gene expression (Piergentili, 2010). Our results seem to provide

explanation for some of the earlier classical observations of mice with different Y

chromosomal deletions exhibiting subfertility/sterility along with sperm morphological

abnormalities, fewer motile sperms, sex ratio skewed toward females etc. Similar

phenotypes are also observed in cross-species male sterile hybrids of *Drosophila* and mouse

(Albrechtova et al., 2012; Campbell, Good, Dean, Tucker, & Nachman, 2012; Heikkinen &

Lumme, 1998; Johnson, Hollocher, Noonburg, & Wu, 1993; Piergentili, 2010; Tao, Zeng, Li,

Hartl, & Laurie, 2003; Vigneault & Zouros, 1986; White, Stubbings, Dumont, & Payseur,

2012). Y-chromosome has also been implicated in the male sterility phenotype of these

interspecies hybrids (Campbell et al., 2012; Carvalho, Vaz, & Klaczko, 1997; Lamnissou,

Loukas, & Zouros, 1996; Vigneault & Zouros, 1986; Zouros, Lofdahl, & Martin, 1988). Thus,

the phenotypes observed in cross species male sterile hybrids and the Y-deletion mutants

are comparable. Introduction of Y-chromosomes into different genetic backgrounds of

Drosophila resulted in deregulated expression of hundreds of genes localizing to the X-

chromosome and autosomes (David et al., 2005; Lemos, Araripe, & Hartl, 2008). It has also

been proposed that incompatibility between the Y-chromosomes and different autosomes

could result in the hybrid dysgenesis of sperm related phenotypes observed in *Drosophila*

(Zouros et al., 1988). Zouros and colleagues also suggested the presence of epistatic

networks in interspecies hybrids, based on the fact that homospecific combination of alleles

at a given set of loci could sustain normal development, but heterospecific combinations

could not (Davis, Noonburg, & Wu, 1994; Lamnissou et al., 1996; Wu & Palopoli, 1994). This

early hypothesis seems to be amply supported by our study. Further, our results elucidate

the Y - derived piRNAs as the genetic basis of epistatic interactions between Y chromosome

and autosomes in mouse. Our results also suggest for the first time, the mechanism of

piRNA mediated regulation of autosomal genes involved in spermiogenesis and male

fertility. This is, to our knowledge is the first report on possible regulation of autosomal

genes involved in male fertility and spermiogenesis, mediated by Y-encoded small

RNAs/piRNAs in any species.

In brief, the XY^{RIII}qdel mutant strain of mouse, where there is a partial deletion of long arm

of the Y chromosome, exhibit sperm morphological and motility related aberrations and

subfertility (Conway et al., 1994). A comparative sperm proteomic profiling of the XY^{RIII} and

XY^{RIII}qdel mice captured few differentially expressed proteins that could partially account for

the aberrant sperm phenotype. Surprisingly, genes corresponding to the deregulated

proteins localized to autosomes and not to the deleted region of the Y chromosome. Earlier

we demonstrated an event of trans-splicing between a Y-ncRNA and a protein coding

autosomal mRNA in human testis for putative translational regulation. A search for the Y-

autosome connections in mouse led to the identification of novel ncRNAs from mouse Y

long arm that subsequently was shown to regulate the genes expressed in testis via piRNAs.

Thus, adopting a top-down approach we have established a novel mode of regulation of

autosomal genes expressed in mouse testis by the Y chromosome and the biology behind

the aberrant sperm phenotype in Y-deleted mice.

Finally, evolutionary impact of novel genetic interactions or regulatory mechanisms such as

those reported in this study could be significant. The generation of piRNAs from species-

specific repeats on mouse Y-chromosome that apparently regulate autosomal gene

expression in testis raises more questions in the field of speciation and evolution. Do

mutations in the Y chromosomal repeats collapse the poise of the species? Are species-

specific repeats on the Y chromosome the fulcrum on which rests the fine balance between

species identity and evolution?

METHODS

Animals and Reagents

The XY^{RIII} strain (wild type) and the XY^{RIII}qdel strain (Y-deletion mutant) of mice used in the

study was a gift from Prof. Paul S Burgoyne, MRC, UK. All the animals used in the

experiments were bred and reared in the in-house animal facility of our institute (CCMB), in

accordance with the guidelines from Indian Science Academy under CPCSEA (Committee for

the Purpose of Control and Supervision of Experimental Animals). This study was approved

by the Institutional Animal Ethics Committee (IAEC 65/28/2006). The recombinant construct

in pAAV-IRES-hrGFP from which MIWI protein was isolated, was a gift from Arvind Kumar,

CCMB, Hyderabad.

Identification of cDNA using M34 (DQ907163):

An amplified mouse testis cDNA library (Mouse testis MATCHMAKER cDNA Library –

Clonetech) was screened with the male specific ES cell EST (CA533654) with sequence

homology to the genomic clone DQ907163 (Fig. 1*E*) as per manufacturer's protocol. Mouse

testis cDNA Library was screened according to standard protocol at a stringency of 2 x SSC,

0.1% SDS at 65^{0}C for 10 min. A total of 2 x 10^5 colonies were screened to obtain 18 clones

after tertiary screening. Male-specificity of the positive clones from the library was

determined using Southern blots containing mouse male and female DNA. All 18 clones

gave the same sequence. The sequence was submitted to the NCBI database and is named

as *Pirmy* (DQ907162).

Identification of Splice Variants: Total RNA (1 µg) isolated from brain and testes tissues

each of XY^{RIII} strain of mouse were reverse transcribed with the (SuperscriptII, Reverse

Transcriptase enzyme, Invitrogen), using oligo (dT) primers and random hexamers. The RT-

mixes were concentration equalized using GAPDH primers. DQ907162 was amplified after

two rounds of PCR using primers to the first and last exons. For the first round, forward

(GTGTGACAGGGTGGGGAATC) and reverse primers (TTCCTGAAGATAGCACTTGTG), and the

following conditions were used - initial denaturation 95^{0}C, 1min, cycle denaturation 95^{0}C, 1

min, annealing 62^{0}C, 30 sec and extension 72^{0}C for 2min (35cycles), final extension 72^{0}C for

7 min. The second round of amplification was done using nested primers

GAGGACCGTATTCATGGAAGAG (forward) and GCAAATGGCTCACATCAGTGG (reverse) using

initial denaturation at 95^{0}C for 1 min, cycle denaturation at 95^{0}C, 1 min, annealing at 66^{0}C

for 30 sec and extension at 72^{0}C for 2 min (38 cycles) and final extension at 72^{0}C for 7 min.

Annealing temperatures up to 70^{0}C yielded multiple products. The multiple products

obtained after two rounds of PCR from both testis and brain (Fig. 2*B*) were cloned into pCR

TOPO vector using Invitrogen TOPO TA cloning kit. Approximately 1000 clones were

sequenced on 3730 DNA Analyser (ABI Prism) using sequencing kit BigDye Terminator V3.1

Cycle Sequencing kit. These yielded 108 unique transcripts. BLAST analysis of these

transcripts against the genomic sequences (GRCm39) at a stringency of >97% localised 80 of

these to a single locus on the Y chromosome (4341127-4381724) and the rest to multiple

sites on the Y.

Therefore, the 80 transcripts that localised to the single locus are splice variants of *Pirmy.*

Sperm Proteome Analysis: Sperm lysates (1 mg of cell weight per 5 µl of lysis buffer – [Urea

8 M, CHAPS 4% (w/v), Tris 40 mM, Biolyte (3-10) 0.2% (w/v) and TBP (1µl/100 µl]). The

suspension was incubated on ice for one hour to allow buffer to permeabilise and lyse the

sample. Further, the sample was briefly sonicated on ice. The lysate was centrifuged for 15

min at 13,000 rpm at 4^0C. The supernatant was collected and was further taken for ultra-

centrifugation at 55,000 rpm for one hour at 4^0C. The clear lysate was collected in fresh

tube and PMSF added to a final concentration of 1mM. The protein concentration in the cell

lysate was estimated by Bicinchoninic acid assay (Pierce, Rockford IL), following

manufacturer's instructions in a micro titre plate. BSA was used as the standard for

estimation. The proteins were separated in the first dimension on 4-7 and 5-8 IPG strips

(Bio-Rad). The strips were then loaded onto 8-20% gradient PAGE for separation on the

second dimension. Protein spots were visualized by Coomassie Blue staining. Spot to spot

matches were done to identify differences. Analysis of five sets of gels after normalization

with control spots using PDQUEST software version 6.0 (Bio-Rad) identified the differential

proteins. Measuring the optical density of these differentially expressed proteins in arbitrary

units validated the quantitative differences (Fig. S7). These values were subjected to

nonparametric Kruskal-Wallis H test and the levels of confidence determined by the Chi-

squared test (75-95% degrees of confidence). Trypsin digested spots were processed to

obtain the protein tags by MS analysis on Hybrid Quadrupole TOF mass spectrometer (API

QSTAR PULSAR i, PE SCIEX).

Small RNA Isolation

Total RNA was extracted from male kidney, testis and brain of mice using TRIZOL reagent (Invitrogen). Total RNA was denatured at 65°C for 10 min, incubated with 10% PEG-8000 (Sigma-Aldrich) and 5 M NaCl for 30 min on ice and centrifuged at 7,000 rpm for 7-10 min. The supernatant containing small RNA was precipitated overnight with 3 volumes of absolute alcohol and centrifuged at 13,000 rpm for 30 min. The small RNA pellet was washed with 80% ethanol and resuspended in RNase free Gibco water. The quality of small RNA was checked on 12% Urea PAGE (Biorad) and quantitated using Nanodrop V-1000 (Thermo Fisher Scientific).

Small RNA Northern Blotting

20-50 μg of small RNA from each tissue was resolved on a 12% Urea PAGE gel in 0.5 x TBE running buffer and transferred onto Hybond N$^+$ membrane. Decade marker (Ambion) was labelled and loaded according to the manufacturer's instructions. 10-25 μM of each LNA-oligonucleotide probe (Exiqon), was end labelled for use as probes (hybridization buffer - 5 x SSC, 5 x Denhardt's and 1% SDS). Blots were hybridized at 37^{0}C and washed from 37^{0}C to 65^{0}C in 2 x SSC, 0.2% SDS depending on the intensity of the signal. U6 was used as the loading control. Fig. S6 shows the location of the LNA (locked nucleic acid) probes used for small RNA northern blots on the corresponding splice variants.

Real-time PCR analysis

Copy number estimation of *Pirmy*

Genomic DNA was isolated from testes of XY^{RIII} and XY^{RIII}qdel mice using phenol-chloroform method and quantified using a Nanodrop (NANODROP 2000, ThermoScientific). Quantitative Real-time PCR (LightCycler 480, Roche) was performed using SYBR green master mix (Qiagen) with 2 ng of genomic DNA and a primer concentration of 200 nM per

reaction. The primers used were as follows: *Pirmy* (exon 7): forward 5'-GTG CGG TTG TGA

AGG TGT TC– 3', reverse 5'-CCT CCA CCT TCC ATT CAC CC-3'; *Gapdh:* forward 5'-ACG GGA

AGC TCA CTG GCA TGG -3', reverse 5'-CAA CAG CGA CAC CCA CTC CTC–3'. PCR conditions

included an initial denaturation for 5 min at 95°C followed by 45 cycles of denaturation at

95°C for 10 secs, annealing at 58°C for 20 sec and elongation at 72°C for 30 sec. The

amplification of specific product was confirmed by melting curve profile (cooling the sample

to 65°C for 1 min and heating slowly with an increase in temperature of 5°C at each step till

95°C, with continuous measurement of fluorescence). The relative fold change in *Pirmy* was

analyzed based on Livak method ($2^{-\Delta\Delta Ct}$).

Real-time PCR analysis of differentially expressed genes

The total RNA was extracted from testes using Trizol (Invitrogen). One μg of RNA was

reverse transcribed to cDNA using Verso cDNA synthesis kit (ThermoScientific). qPCR was

performed using SYBR green master mix (Qiagen) and analysed in Roche Light Cycler LC480.

The primer sequences corresponding to mouse cDNA used were as follows: forward 5'-CGA

GGG CCA GAC AGG GAT TG–3' and reverse 5'-CCC ATA GAC AGA GGA CAT CAG- 3' for Riken

cDNA 1700001L19; forward 5'-ACT TCC CGT CGC CGC TAT C-3' and reverse 5'-TGA CCG ACA

GGA ACA CAG AGG-3' for *Sdf2l1*; forward 5'-CAG CAT CGA GCA GAA GTA TAA GC-3' and

reverse 5'-TGG GTG GAG TTA TTG CAG TAG-3' for *Mast*; forward 5'-GGA AAC CAC GTC AAA

TTG -3' and reverse 5'-GGT GAT GAG GAA ATT GTC-3' for Calreticulin; forward 5'-GGC TAC

TTG ACC ACT GC-3' and reverse 5'-TTT GAG AAT CGG AAG AGT C-3' for *Spink2* Variant 2;

forward 5'-TTC CGA ACA CCA GAC TG-3' and reverse 5'-ATG GCT ACC GTC CTC C-3' for

Spink2 Variant 3; forward 5'-TGA AGT CGC AGG AGA CAA CCT-3' and reverse 5'-ATG GCC

TTC CGT GTT CCT A-3' for *Gapdh*. PCR conditions included an initial denaturation for 5 min

at 95°C followed by 45 cycles of denaturation at 95°C for 10 secs, annealing at 58°C for 20

sec and elongation at 72°C for 30 sec. The amplification of specific product was confirmed

by melting curve profile. The relative fold change in expression was estimated based on

Livak method ($2^{-\Delta\Delta Ct}$).

Electrophoretic mobility shift assay RNA oligonucleotides corresponding to

GAAGCAGAUGAGUAUAUG from *Sod* and UCAUUGGACAUAAACUGAAUUUUCCA from the

gene for hypothetical protein spot A (Q9DAR0) were end labelled with γ-^{32}P ATP and column

purified using G-25 Sephadex (Sigma-Aldrich) and quantitated on a scintillation counter.

EMSA reactions were set up in a total volume of 25 µl using binding buffer (20 mM HEPES, 3

mM MgCl2, 40 mM KCl, 5% Glycerol, 2 mM DTT and RNase inhibitor (4U)), with MIWI

protein (5 µg per reaction). MIWI was over-expressed from a recombinant construct in

pAAV-IRES-hrGFP vector and purified using the FLAG tag. Competitors i.e., unlabeled

oligonucelotides (30 x concentration of hot oligo), MIWI antibody (90 ng) and Argonaute 3

antibody (100 ng) were added to the reaction, incubated for 1 h on ice, before addition of

radio-labeled oligonucleotide (7000-10,000 cpm) and the entire mix was incubated on ice

for another 30 min. EMSA was done on 5% native PAGE and image captured using FUJI

phosphor Imager (FUJIFILM FLA-3000). A known piRNA, piR1, was used as the positive

control, Argonaute 3 antibody served as the antibody control.

Luciferase assay

Either Luciferase gene alone or luciferase along with the UTR was cloned into pcDNA3.1

expression vector for assaying the effect of antagopirs (Figure 7H) on Luciferase expression.

Co-transfection experiments were done using the GC-1spg cell line (ATCC CRL-2053) and

lipofectamine 2000 (Invitrogen) using protocols specified by the manufacturer. Cells were

seeded in 48-well plates, 24 hrs prior to transfection to obtain approximately 80%

confluency. Each well was transfected with 50 ng of pcDNA3.1 plasmid containing either

Luciferase gene alone or along with the cloned UTRs, 50 ng of the β-gal plasmid and varying

concentrations of oligonucleotides complementary to the piRNA along with 0.5µl of

lipofectamine 2000 in antibiotic and serum free DMEM (GIBCO). The complementary

oligonucleotide to the piRNA has been designated as antagopirs. The antagopirs (*Sod* -

5'GAAGCAGAUGAGUAUAUG3'; *PLA2G12B* - 5'CCAAACUGUUGGAAGAAGGAAU3') were

procured as RNA oligonucleotides from Eurofins Genomics India Pvt. Ltd, Bangalore, India.

Different concentrations of antagopirs (0 nM, 10 nM, 20 nM and 40 nM/ well) were tested

in the assay for their effect on the UTRs. Five hrs post transfection, the medium was

replaced with complete growth medium. The cell extracts were prepared 24 hrs post

transfection using Reporter Lysis Buffer (Promega) and assayed for Luciferase activity in

EnSpire 2300 multimode plate reader (Perkin Elmer). The Luciferase activity was normalized

using β galactosidase. Three independent sets of experiments were done in triplicates.

DATA ACCESS

All sequences from this study have been deposited in NCBI database. With the accession

numbers: DQ907162.1, FJ541075-FJ541181, Q7TPM5, Q9DAR0, Q8BMY7.

ACKNOWLEDGEMENTS

We would like to dedicate this manuscript to Prof. Lalji Singh and Prof. Burgoyne both of

whom have left us for their heavenly abode and whom we miss greatly at this point of time.

We gratefully acknowledge the gift of the XY[RIII] and XY[RIII]qdel mice by Prof. Paul S Burgoyne,

MRC, Edinburgh, UK. This study could not have been done without the gift of these mice.

The GC-1spg cell line was gifted by Prof. MRS Rao, JNCASR, Bangalore, India. We thank Dr.

Dinesh Kumar and Professor B. K. Thelma for reading the manuscript and giving useful

inputs and Mr. Sivarajan Karunanithi for help with partial *in silico* analysis.

The funding by Department of Science and Technology, India (SP/SO/B70/2001) and

Department of Biotechnology, India (BT/PR 10707/AGR/36/596/2008), intramural funding

from Council of Scientific and Industrial Research (CSIR), India to RAJ and fellowships by

CSIR, India to HMR, RB are acknowledged.

Author contributions

HMR, KM did experiments and partial bioinformatics analysis. RB, ZJ, PA, NMP, VMD, MS,

BS, JLA, SMT, RRD, SMR performed experiments. ST, AC, SK contributed to *in silico* analysis.

NR helped with confocal imaging. LS gave the probe and inputs. RAJ conceived, guided the

work and wrote the manuscript.

DISCLOSURE DECLARATION

The authors declare that they have no competing interests

REFERENCES:

Adham, I. M., Nayernia, K., & Engel, W. (1997). Spermatozoa lacking acrosin protein show delayed fertilization. *Mol Reprod Dev, 46*(3), 370-376. doi:10.1002/(SICI)1098-2795(199703)46:3<370::AID-MRD16>3.0.CO;2-2

Albrechtova, J., Albrecht, T., Baird, S. J., Macholan, M., Rudolfsen, G., Munclinger, P., . . . Pialek, J. (2012). Sperm-related phenotypes implicated in both maintenance and breakdown of a natural species barrier in the house mouse. *Proc Biol Sci, 279*(1748), 4803-4810. doi:10.1098/rspb.2012.1802

Aravin, A. A., Hannon, G. J., & Brennecke, J. (2007). The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science, 318*(5851), 761-764. doi:10.1126/science.1146484

Bajpai, A., Sridhar, S., Reddy, H. M., & Jesudasan, R. A. (2007). BRM-Parser: a tool for comprehensive analysis of BLAST and RepeatMasker results. *In Silico Biol, 7*(4-5), 399-403. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/18391232

Bellott, D. W., Hughes, J. F., Skaletsky, H., Brown, L. G., Pyntikova, T., Cho, T. J., . . . Page, D. C. (2014). Mammalian Y chromosomes retain widely expressed dosage-sensitive regulators. *Nature, 508*(7497), 494-499. doi:10.1038/nature13206

Bhattacharya, R., Devi, M. S., Dhople, V. M., & Jesudasan, R. A. (2013). A mouse protein that localizes to acrosome and sperm tail is regulated by Y-chromosome. *BMC Cell Biol, 14*(1), 50. doi:10.1186/1471-2121-14-50

Bishop, C. E., & Hatat, D. (1987). Molecular cloning and sequence analysis of a mouse Y chromosome RNA transcript expressed in the testis. *Nucleic Acids Res, 15*(7), 2959-2969. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/3562242

Burgoyne, P. S., Mahadevaiah, S. K., Sutcliffe, M. J., & Palmer, S. J. (1992). Fertility in mice requires XY pairing and a Y-chromosomal "spermiogenesis" gene mapping to the long arm. *Cell, 71*(3), 391-398.

Burgoyne, P. S., & Mitchell, M. J. (2007). The roles of mouse Y chromosome genes in spermatogenesis. *Y chromosome and male germ cell biology. Hackensack (New Jersey): World Scientific Publishers*, 27-45.

Campbell, P., Good, J. M., Dean, M. D., Tucker, P. K., & Nachman, M. W. (2012). The contribution of the Y chromosome to hybrid male sterility in house mice. *Genetics, 191*(4), 1271-1281. doi:10.1534/genetics.112.141804

Carvalho, A. B., Vaz, S. C., & Klaczko, L. B. (1997). Polymorphism for Y-linked suppressors of sex-ratio in two natural populations of Drosophila mediopunctata. *Genetics, 146*(3), 891-902. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9215895

Choi, E., Lee, J., Oh, J., Park, I., Han, C., Yi, C., . . . Cho, C. (2007). Integrative characterization of germ cell-specific genes from mouse spermatocyte UniGene library. *BMC Genomics, 8*, 256. doi:10.1186/1471-2164-8-256

Cocquet, J., Ellis, P. J., Yamauchi, Y., Mahadevaiah, S. K., Affara, N. A., Ward, M. A., & Burgoyne, P. S. (2009). The multicopy gene Sly represses the sex chromosomes in the male mouse germline after meiosis. *PLoS Biol, 7*(11), e1000244. doi:10.1371/journal.pbio.1000244

Conway, S. J., Mahadevaiah, S. K., Darling, S. M., Capel, B., Rattigan, A. M., & Burgoyne, P. S. (1994). Y353/B: a candidate multiple-copy spermiogenesis gene on the mouse

Y chromosome. *Mamm Genome, 5*(4), 203-210. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7516747

David, J. R., Araripe, L. O., Chakir, M., Legout, H., Lemos, B., Petavy, G., . . . Moreteau, B. (2005). Male sterility at extreme temperatures: a significant but neglected phenomenon for understanding Drosophila climatic adaptations. *Journal of evolutionary biology, 18*(4), 838-846. doi:10.1111/j.1420-9101.2005.00914.x

Davis, A. W., Noonburg, E. G., & Wu, C. I. (1994). Evidence for complex genic interactions between conspecific chromosomes underlying hybrid female sterility in the Drosophila simulans clade. *Genetics, 137*(1), 191-199. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8056310

Deng, W., & Lin, H. (2002). miwi, a murine homolog of piwi, encodes a cytoplasmic protein essential for spermatogenesis. *Developmental cell, 2*(6), 819-830. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/12062093

Ellis, P. J., Clemente, E. J., Ball, P., Toure, A., Ferguson, L., Turner, J. M., . . . Burgoyne, P. S. (2005). Deletions on mouse Yq lead to upregulation of multiple X- and Y-linked transcripts in spermatids. *Hum Mol Genet, 14*(18), 2705-2715. doi:ddi304 [pii]

10.1093/hmg/ddi304

Ellis, P. J., Ferguson, L., Clemente, E. J., & Affara, N. A. (2007). Bidirectional transcription of a novel chimeric gene mapping to mouse chromosome Yq. *BMC Evol Biol, 7*, 171. doi:1471-2148-7-171 [pii]

10.1186/1471-2148-7-171

Fukuda, S., Sumii, M., Masuda, Y., Takahashi, M., Koike, N., Teishima, J., . . . Kamiya, K. (2001). Murine and human SDF2L1 is an endoplasmic reticulum stress-inducible gene and encodes a new member of the Pmt/rt protein family. *Biochem Biophys Res Commun, 280*(1), 407-414. doi:10.1006/bbrc.2000.4111

Girard, A., Sachidanandam, R., Hannon, G. J., & Carmell, M. A. (2006). A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature, 442*(7099), 199-202. doi:10.1038/nature04917

Goldstein, L. S., Hardy, R. W., & Lindsley, D. L. (1982). Structural genes on the Y chromosome of Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America, 79*(23), 7405-7409. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/6818544

Heikkinen, E., & Lumme, J. (1998). The Y chromosomes of Drosophila lummei and D. novamexicana differ in fertility factors. *Heredity, 81 (Pt 5)*, 505-513. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9881450

Jehan, Z., Vallinayagam, S., Tiwari, S., Pradhan, S., Singh, L., Suresh, A., . . . Jesudasan, R. A. (2007). Novel noncoding RNA from human Y distal heterochromatic block (Yq12) generates testis-specific chimeric CDC2L2. *Genome Res, 17*(4), 433-440. doi:10.1101/gr.5155706

Johnson, N. A., Hollocher, H., Noonburg, E., & Wu, C. I. (1993). The effects of interspecific Y chromosome replacements on hybrid sterility within the Drosophila simulans clade. *Genetics, 135*(2), 443-453. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8244006

Kherraf, Z. E., Christou-Kent, M., Karaouzene, T., Amiri-Yekta, A., Martinez, G., Vargas, A. S., . . . Arnoult, C. (2017). SPINK2 deficiency causes infertility by inducing sperm defects in heterozygotes and azoospermia in homozygotes. *EMBO Mol Med, 9*(8), 1132-1149. doi:10.15252/emmm.201607461

Korley, R., Pouresmaeili, F., & Oko, R. (1997). Analysis of the protein composition of the mouse sperm perinuclear theca and characterization of its major protein constituent. *Biol Reprod, 57*(6), 1426-1432. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/9408250

Kotula-Balak, M., Grzmil, P., Styrna, J., & Bilinska, B. (2004). Immunodetection of aromatase in mice with a partial deletion in the long arm of the Y chromosome. *Acta histochemica, 106*(1), 55-64. doi:10.1016/j.acthis.2003.09.001

Kuroda-Kawaguchi, T., Skaletsky, H., Brown, L. G., Minx, P. J., Cordum, H. S., Waterston, R. H., . . . Page, D. C. (2001). The AZFc region of the Y chromosome features massive palindromes and uniform recurrent deletions in infertile men. *Nat Genet, 29*(3), 279-286. doi:10.1038/ng757

Lamnissou, K., Loukas, M., & Zouros, E. (1996). Incompatibilities between Y chromosome and autosomes are responsible for male hybrid sterility in crosses between Drosophila virilis and Drosophila texana. *Heredity, 76 (Pt 6)*, 603-609. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8801221

Lemos, B., Araripe, L. O., & Hartl, D. L. (2008). Polymorphic Y chromosomes harbor cryptic variation with manifold functional consequences. *Science, 319*(5859), 91-93. doi:10.1126/science.1148861

Lu, J. C., Huang, Y. F., & Lu, N. Q. (2010). [WHO Laboratory Manual for the Examination and Processing of Human Semen: its applicability to andrology laboratories in China]. *Zhonghua Nan Ke Xue, 16*(10), 867-871. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/21243747

Mao, H. T., & Yang, W. X. (2013). Modes of acrosin functioning during fertilization. *Gene, 526*(2), 75-79. doi:10.1016/j.gene.2013.05.058

McClintock, T. S., Glasser, C. E., Bose, S. C., & Bergman, D. A. (2008). Tissue expression patterns identify mouse cilia genes. *Physiol Genomics, 32*(2), 198-206. doi:10.1152/physiolgenomics.00128.2007

Moretti, C., Serrentino, M. E., Ialy-Radio, C., Delessard, M., Soboleva, T. A., Tores, F., . . . Cocquet, J. (2017). SLY regulates genes involved in chromatin remodeling and interacts with TBL1XR1 during sperm differentiation. *Cell Death Differ, 24*(6), 1029-1044. doi:10.1038/cdd.2017.32

Moritz, A., Lilja, H., & Fink, E. (1991). Molecular cloning and sequence analysis of the cDNA encoding the human acrosin-trypsin inhibitor (HUSI-II). *FEBS Lett, 278*(1), 127-130. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/1704312

Oko, R., & Morales, C. R. (1994). A novel testicular protein, with sequence similarities to a family of lipid binding proteins, is a major component of the rat sperm perinuclear theca. *Dev Biol, 166*(1), 235-245. doi:10.1006/dbio.1994.1310

Pang, K. C., Stephen, S., Engstrom, P. G., Tajul-Arifin, K., Chen, W., Wahlestedt, C., . . . Mattick, J. S. (2005). RNAdb--a comprehensive mammalian noncoding RNA database. *Nucleic acids research, 33*(Database issue), D125-130. doi:10.1093/nar/gki089

Piergentili, R. (2010). Multiple roles of the Y chromosome in the biology of Drosophila melanogaster. *The Scientific World Journal, 10*, 1749-1767.

Pouresmaeili, F., Morales, C. R., & Oko, R. (1997). Molecular cloning and structural analysis of the gene encoding PERF 15 protein present in the perinuclear theca of the rat spermatozoa. *Biol Reprod, 57*(3), 655-659. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/9283004

Prado, V. F., Lee, C. H., Zahed, L., Vekemans, M., & Nishioka, Y. (1992). Molecular characterization of a mouse Y chromosomal repetitive sequence that detects

transcripts in the testis. *Cytogenet Cell Genet, 61*(2), 87-90. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1395731

Redecker, P., Kreutz, M. R., Bockmann, J., Gundelfinger, E. D., & Boeckers, T. M. (2003). Brain synaptic junctional proteins at the acrosome of rat testicular germ cells. *J Histochem Cytochem, 51*(6), 809-819. doi:10.1177/002215540305100612

Reynard, L. N., Cocquet, J., & Burgoyne, P. S. (2009). The multi-copy mouse gene Sycp3-like Y-linked (Sly) encodes an abundant spermatid protein that interacts with a histone acetyltransferase and an acrosomal protein. *Biol Reprod, 81*(2), 250-257. doi:10.1095/biolreprod.108.075382

Riel, J. M., Yamauchi, Y., Ruthig, V. A., Malinta, Q. U., Blanco, M., Moretti, C., . . . Ward, M. A. (2019). Rescue of Sly Expression Is Not Sufficient to Rescue Spermiogenic Phenotype of Mice with Deletions of Y Chromosome Long Arm. *Genes (Basel), 10*(2). doi:10.3390/genes10020133

Riel, J. M., Yamauchi, Y., Sugawara, A., Li, H. Y., Ruthig, V., Stoytcheva, Z., . . . Ward, M. A. (2013). Deficiency of the multi-copy mouse Y gene Sly causes sperm DNA damage and abnormal chromatin packaging. *J Cell Sci, 126*(Pt 3), 803-813. doi:10.1242/jcs.114488

Schultz, N., Hamra, F. K., & Garbers, D. L. (2003). A multitude of genes expressed solely in meiotic or postmeiotic spermatogenic cells offers a myriad of contraceptive targets. *Proc Natl Acad Sci U S A, 100*(21), 12201-12206. doi:10.1073/pnas.1635054100

Simpson, E. R., Mahendroo, M. S., Means, G. D., Kilgore, M. W., Hinshelwood, M. M., Graham-Lorence, S., . . . et al. (1994). Aromatase cytochrome P450, the enzyme responsible for estrogen biosynthesis. *Endocr Rev, 15*(3), 342-355. doi:10.1210/edrv-15-3-342

Singh, L., Panicker, S. G., Nagaraj, R., & Majumdar, K. C. (1994). Banded krait minor-satellite (Bkm)-associated Y chromosome-specific repetitive DNA in mouse. *Nucleic acids research, 22*(12), 2289-2295. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8036156

Soh, Y. Q., Alfoldi, J., Pyntikova, T., Brown, L. G., Graves, T., Minx, P. J., . . . Page, D. C. (2014). Sequencing the mouse Y chromosome reveals convergent gene acquisition and amplification on both sex chromosomes. *Cell, 159*(4), 800-813. doi:10.1016/j.cell.2014.09.052

Styrna, J., Bili, B., & Krzanowska, H. (2002). The effect of a partial Y chromosome deletion in B10. BR-Ydel mice on testis morphology, sperm quality and efficiency of fertilization. *Reproduction, Fertility and Development, 14*(2), 101-108.

Styrna, J., Klag, J., & Moriwaki, K. (1991). Influence of partial deletion of the Y chromosome on mouse sperm phenotype. *Journal of reproduction and fertility, 92*(1), 187-195. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/2056490

Tao, Y., Zeng, Z. B., Li, J., Hartl, D. L., & Laurie, C. C. (2003). Genetic dissection of hybrid incompatibilities between Drosophila simulans and D. mauritiana. II. Mapping hybrid male sterility loci on the third chromosome. *Genetics, 164*(4), 1399-1418. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/12930748

Tiepolo, L., & Zuffardi, O. (1976). Localization of factors controlling spermatogenesis in the nonfluorescent portion of the human Y chromosome long arm. *Hum Genet, 34*(2), 119-124. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1002136

Toure, A., Clemente, E. J., Ellis, P., Mahadevaiah, S. K., Ojarikre, O. A., Ball, P. A., . . . Affara, N. A. (2005). Identification of novel Y chromosome encoded transcripts by testis

transcriptome analysis of mice with deletions of the Y chromosome long arm. *Genome Biol, 6*(12), R102. doi:gb-2005-6-12-r102 [pii]

10.1186/gb-2005-6-12-r102

Toure, A., Szot, M., Mahadevaiah, S. K., Rattigan, A., Ojarikre, O. A., & Burgoyne, P. S. (2004). A new deletion of the mouse Y chromosome long arm associated with the loss of Ssty expression, abnormal sperm development and sterility. *Genetics, 166*(2), 901-912. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/15020475

Vigneault, G., & Zouros, E. (1986). The genetics of asymmetrical male sterility in Drosophila mojavensis and Drosophila arizonensis hybrids: interactions between the Y-chromosome and autosomes. *Evolution*, 1160-1170.

Vogt, P. H. (1998). Human chromosome deletions in Yq11, AZF candidate genes and male infertility: history and update. *Molecular human reproduction, 4*(8), 739-744. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/9733430

Wegmann, D., Dupanloup, I., & Excoffier, L. (2008). Width of gene expression profile drives alternative splicing. *PLoS One, 3*(10), e3587. doi:10.1371/journal.pone.0003587

White, M. A., Stubbings, M., Dumont, B. L., & Payseur, B. A. (2012). Genetics and evolution of hybrid male sterility in house mice. *Genetics, 191*(3), 917-934. doi:10.1534/genetics.112.140251

Wu, C. I., & Palopoli, M. F. (1994). Genetics of postmating reproductive isolation in animals. *Annual review of genetics, 28*, 283-308. doi:10.1146/annurev.ge.28.120194.001435

Xiao, P., Tang, A., Yu, Z., Gui, Y., & Cai, Z. (2008). Gene expression profile of 2058 spermatogenesis-related genes in mice. *Biol Pharm Bull, 31*(2), 201-206. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/18239273

Yamagata, K., Murayama, K., Okabe, M., Toshimori, K., Nakanishi, T., Kashiwabara, S., & Baba, T. (1998). Acrosin accelerates the dispersal of sperm acrosomal proteins during acrosome reaction. *J Biol Chem, 273*(17), 10470-10474. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/9553106

Yanagimachi, R. (1982). Requirement of extracellular calcium ions for various stages of fertilization and fertilization related phenomena in the hamster. *Gamete Research, 5*(4), 323-344. doi:10.1002/mrd.1120050404

Zouros, E., Lofdahl, K., & Martin, P. (1988). Male hybrid sterility in Drosophila: interactions between autosomes and sex chromosomes in crosses of D. mojavensis and D. arizonensis. *Evolution*, 1321-1331.

FIGURE LEGENDS

Figure 1. Analysis of M34 (DQ907163) and identification of a novel noncoding RNA. (*A*) Localization by FISH of the genomic clone, M34 to Mouse Y long arm in multiple copies spanning its entire length. (*B*) Mouse genome map view of M34 BLAST hits, showing Y-chromosomal localization further indicating male-specificity of these repeats. (*C*) Fluorescent in situ hybridization (FISH) using M34 shows intense signals in adult mouse

testis. (*D*) Hybridization of M34 onto RNase treated testis sections does not elicit signals, indicating that the signals in panel C are due to the presence of M34 derived RNA (*E*) Sequence analysis of the 9.5kb M34 shows presence of incomplete copies of different repeats like long terminal repeats (LTRs), long interspersed nuclear elements (LINEs), short interspersed nuclear elements (SINEs), endogenous retroviral sequences (ERVK) and simple sequence repeats in both direct and reverse orientations in the clone. ESTs matching to M34 are marked as dotted arrows at the 3'end. (*F*) Shows the Y-specific localization of *Pirmy* cDNA on a mouse metaphase spread by FISH. (*G*) Partial homology between *Sly* and *Pirmy* (DQ907162) indicating identification of a novel cDNA. Homology region is highlighted in green rectangles. (*H*) Shows the consensus splice signal sequences at all intron- exon junctions. Since DQ907162 localizes to the complementary strand, the splice site consensus is seen as CT/AC instead AG/GT (see also Data Sheet).

Figure 2. Identification of multiple splice-variants of *Pirmy*. Autoassembler program of ABI-Prism identified cDNAs that differed from one another. (*A*) Color coded line diagram showing extensive alternative splicing of *Pirmy*. The splice variants depicted here localise to Y: 4341127-4381724 (GRC m39). Each exon is represented by the same colour in different isoforms. Sizes of the exons are to scale. Top two lines show the representation of all exons present at this locus according to their order in the genomic sequence as e1, e2 etc. Line 2 indicates the nucleotide positions of each exon in a scenario where all the exons present. The exons have been arranged in linear order. (*B*) RT-PCR amplification of *Pirmy* showed multiple amplicons in both testis and brain, with many more amplicons in testis compared to brain. The RT-PCR products from brain and testis were cloned and sequenced to identify the splice variants. NTC is the non-template control. (*C*) BLAST analysis against mouse genome localizes the splice variants of *Pirmy* to NT_166343.2.

Figure 3. Localization of 28 transcript variants. (*A*) Top two lines show the representation of all exons present together according to their order in the transcript variants as e1, e2 etc. Putative nucleotide positions in a scenario wherein all the exons are present are indicated. Different splice variants have been arranged in linear order. Exons in dashed lines are specific to these 28 transcript variants, whereas other exons are common to *Pirmy* splice variants (Figure 2) and the transcript variants.

Figure 4. Comparison of splice-variants from testis and brain. The sequential exons from *Pirmy* splice-variants in testis have been represented in an Excel data sheet. Highlighted in red are the splice variants that are transcribed from male brain, which were also found in testis. The representation of e1, e2 etc. and the nucleotide positions is same as in Figure 2A.

Figure 5. Sperm proteins are deregulated in XY^{RIII}qdel mice. (*A*) Sperm lysates from the wild type XY^{RIII} strain and the mutant XY^{RIII}qdel mice were separated by 2D-PAGE in the pI ranges of 4-7 and 5-8 on 8-20% gradient gels. Five differentially expressed proteins (D1 to D5) were identified by Mass Spectrometry analysis in the 4-7 pI range, of which 4 were upregulated (upward arrow) and one downregulated (downward arrow). Three proteins (A, B & C) were not detectable in the 5-8 pI range in XY^{RIII}qdel compared to the XY^{RIII} sperm lysate. (*B*) The list of all the differentially expressed proteins identified in the proteomics screen is given in the table. The genes corresponding to all the differentially expressed proteins in XY^{RIII}qdel localized to autosomes. (*C*) confirms the expression levels of four proteins (D1, D4, D5 and C) identified on 2D-gels by western blotting; the differential expression is observed in sperms but not in testis (RT - XY^{RIII} testis, YT - XY^{RIII}qdel testis, RS - XY^{RIII} sperm, YS - XY^{RIII}qdel sperm). The lower sub-panel for all blots is the loading control using Tubulin.

Figure 6. Localization of *Pirmy* homologous sequences to UTRs of deregulated genes. Panel (*A*) shows the UTR regions of the deregulated genes with the sequences homologous to *Pirmy* highlighted in red. Both +/+ and +/- homologies are observed. The splice isoforms of *Pirmy* are indicated in brown and the gene names in green color. (*B*) Couple of transcripts (aromatase and caldendrin) were identified independent of the proteomics screen (*C*) acrosin (literature survey) also harbor homologies to *Pirmy.*

Figure 7. Identification of piRNAs in the transcript variants. (*A*) Short stretches of homology identified in the UTRs by BLAST analysis against UTR database are represented on the set of 28 transcript variants. The boxes highlighted in black match within exons and boxes highlighted in red match at the exon-exon junctions. Line 1 indicates the nucleotide positions as in Fig. 3*A*. (*B*) Pie chart shows ~81% of the matches in exons and ~19% at exon-exon junctions. (*C*) Use of sequences homologous to the 3' UTRs of a hypothetical protein (SpotA4), *Sod, Bche, PLA2G12B* and *Spn* as probes on small RNA northern blots shows testis-specific signals of ~30 nt size (indicated by arrows), which correspond to the size of piRNAs. Control blot using a scrambled oligonucleotide as probe shows no signal at ~30 nt size. (*D*) Hybridization using probes from the sense (S) and antisense (AS) strands of *Sod* and *Bche* shows differential transcription from the two strands under the same conditions. Lower panels (C, D) show loading control using U6 probe. (*E, F*) EMSA using chemically synthesized RNA oligonucleotide sequences from *FJ541077* (E) and *FJ541118* (F) that have homology to the UTRs of the genes of hypothetical protein spot A (A4) and *Sod* respectively. These oligonucleotides and piR1 (Girard et al. 2006), a known piRNA, showed the shift in mobility with recombinant MIWI protein. The gel-shifts obtained with RNA-oligonucleotides from FJ541077 (spotA4) and FJ541118 (*Sod*) were competed out by cold piR1 and vice versa, i.e., the gelshift observed with piR1 was competed out by cold RNA-oligonucleotides from

FJ541077 and FJ541118. Pre-incubation with MIWI antibody abolished the gelshift whereas pre-incubation with argonaute 3 antibody (ARG) did not - indicating specificity of binding. (*G*) Schematic representation of 3'UTR reporter constructs. (*H*) Results of reporter assay: Antagopirs to piRNA homologous sequences present in 3'UTRs of two genes, *Sod* and *PLA2G12B*, bring about concentration dependent reduction in luciferase expression. Use of these oligos against a non-target 3'UTR (*Cdc2l1*) did not affect luciferase expression.

E

1 — 2000
LTR LTR AT-rich T-rich (T)n (TG)n T-rich B1 SINE (GATA)n AT-rich

2001 — 4000
T-rich AT-rich L1 LINE MaLR LTR ERVK LTR B2 SINE

4001 — 6000
L1 LINE L1 LINE B4 SINE (CATA)n (CA)n GA-rich ES cell

6001 — 8000
Soares thymus Stratagene-

8001 — 9510
-skin B6-Dendritic cell NIA 7.4K cDNA

M34 (DQ907163)

F *Pirmy*

G

37nt 219nt 519nt *Pirmy* (DQ907162.1) 1395nt

(Exons 2-5)

1nt 5nt 41nt 45nt 345nt

Sly (NM_201530.2)
Synaptonemal complex 3 like Y

918nt

H

Query DQ907162.1 (1374 bp); Subject AC213374.4 (161276 bp; +/-)

Exon1 (2-219) end
CTCTATGGCCCTGAC**CT**GTTCAATCCTC

Exon2 (220-351)
ACTCAGTAGT**CT**TCTTAGTT...CTAT**AC**CTTCCGCAAGA

Exon 3 (352-418)
GCTCTTTA**CT**GTATTGA...AAAATGTT**AC**CCCATATATTCAT

Exon 4 (419-453)
CTTCATCT**CT**GTGAAGA...AACACAT**AC**CTCCAGATT

Exon 5 (454-520)
CTTTA**CT**GTATTGAA...AATATT**AC**CCCATGTCTT

Exon 6 (521-597)
ACTCTATTGAA**CT**GGCAAA...GCATCTG**AC**CTTGAAATGCT

Exon 7 (598-632)
AATTTCATCTCT**CT**GTGAAGACA...AAAATACAT**AC**CTCCAGATGTAG

Exon 8 (633-686)
AGAAAAGAC**CT**AGAGAATTAG...ACTCAC**AC**CTCCAAGAG

Exon 9 (687-814)
AAACCC**CT**GGAGGAGGAGAAT...TCTCTT**AC**CAGCATTTG

Exon 10 (815-934)
GGATC**CT**TTGGC...AGCATT**AC**CTTGGGC

Exon 11 (935-1106)
GATAAT**CT**ATAAAGG...TAAGT**AC**CAGAATAAT

Exon 12 (1107-1374)
ATGATGCAGC**CT**GGAATATAGT

218 332 467 534 602 635 746 827 896 934 1032 1180 1281 1381 1480 2516

218 332 467 534 602 635 746 827 896 934 1032 1180 1281 1381 1480 2516
963 1061 1351 2222 2283

FJ541075
FJ541076
FJ541077
FJ541078
FJ541079
FJ541080
FJ541126
FJ541093
FJ541081
FJ541082
FJ541083
FJ541084
FJ541085
FJ541086
FJ541087
FJ541088
FJ541089
FJ541090
FJ541091
FJ541092
FJ541094
FJ541095
FJ541096
FJ541097
FJ541098
FJ541099
FJ541100
FJ541101
FJ541102

#	Splice variants	e1	e1a	e2	e3	e3a	e4	e5	e6	e7	e8	e8a	e8b	e9	e9a	e10	e11	e12	e13	e13a	e14	e15	e15a	e15b	e15c
		218	416	548	594	615	649	713	790	816	828	882	998	1026	1126	1244	1518	1625	1705	1728	1804	1893	1977	2560	2851
1	FJ541103																								
2	FJ541105																								
3	FJ541106																								
4	FJ541107																								
5	FJ541108																								
6	FJ541109																								
7	FJ541111																								
8	FJ541112																								
9	FJ541113																								
10	FJ541114																								
11	FJ541115																								
12	FJ541116																								
13	FJ541117																								
14	FJ541118																								
15	FJ541119																								
16	FJ541120																								
17	FJ541121																								
18	FJ541122																								
19	FJ541123																								
20	FJ541124																								
21	FJ541104																								
22	FJ541125																								
23	FJ541126																								
24	FJ541142																								
25	FJ541127																								
26	FJ541128																								
27	FJ541129																								
28	FJ541130																								
29	FJ541131																								
30	FJ541132																								
31	FJ541133																								
32	FJ541134																								
33	FJ541135																								
34	FJ541136																								
35	FJ541137																								
36	FJ541138																								
37	FJ541140																								
38	FJ541141																								
39	FJ541143																								
40	FJ541144																								
41	FJ541145																								
42	FJ541146																								
43	FJ541147																								
44	FJ541148																								
45	FJ541149																								
46	FJ541150																								
47	FJ541151																								
48	FJ541152																								
49	FJ541153																								
50	FJ541154																								
51	FJ541155																								
52	FJ541156																								
53	FJ541157																								
54	FJ541158																								
55	FJ541159																								
56	FJ541160																								
57	FJ541139																								
58	FJ541161																								
59	FJ541162																								
60	FJ541163																								
61	FJ541164																								
62	FJ541165																								
63	FJ541166																								
64	FJ541167																								
65	FJ541110																								
66	FJ541168																								
67	FJ541169																								
68	FJ541170																								
69	FJ541171																								
70	FJ541172																								
71	FJ541173																								
72	FJ541174																								
73	FJ541175																								
74	FJ541176																								
75	FJ541177																								
76	FJ541178																								
77	FJ541179																								
78	FJ541180																								
79	FJ541181																								

XY^RIII **XY^RIII qdel** **KDa**

A

pI 4 - 7

D1

66
43
29
20

D4
D2
D3   D5

pI 5 - 8

A
B
C

18.4

6.5

B

Serial number	Protein spot	MS-tags and N-terminal sequences	Protein identity & Gene ID	Protein expression in XY^RIII qdel	Chromosome localization
1	D1	**DPAIY** (N-terminal sequence)	CALRETICULIN **12317**	Upregulated	8
2	D2	(198.0975) **DCGHFDFPACPR** (156.1027) (269.1519) **EDGSHINII** (629.1989)	SPINK2 **69982**	Upregulated	5
3	D3	(198.08) **DCGHFDFPACPR** (159.10) **SQFRTPDCGH** (N-terminal sequence)	Q8BMY7 **69982**	Down regulated	5
4	D4	(115.0652) **GVANVSIED** (156.1003) (99.1182) **ISLSGEGSIIG** (156.0945)	SOD **20655**	Upregulated	16
5	D5	(1469.6) **ISSENFENYV** (156.0826) (131.0266) **DIQAGSAC** (156.0985) (113.0616) **GEEFEETTADN** (156.1044) (230.1680) **VECTMNNVVST** (156.1044)	FABP9 **21884**	Upregulated	3
6	A	(398.4527) **HINEEEQE** (155.9080)	Q9DAR0 **69315**	Absent	13
7	B	(218.1635) **EGLFLQ** (1192.3747)	**SDF2L1**	Absent	16
8	C	(214.1065) **VSQSNDLSSL** (156.0965)	Q7TPM5 **69987**	Absent	2

C

CALRETICULIN (D1)   SOD (D4)   FABP9 (D5)   MAST (C)

RT YT RS YS   RT YT RS YS   RT YT RS YS   RT YT RS YS

Tubulin

A

Calreticulin_3'UTR

5' CTGGTGTTTGTCTTTAATTCTCCTTCAGCCCTCATCTGGTTTCTCATTTTTGAATCAACATCTTTTCC 3'
187   3'|||||||||||5'   254
(FJ541150) AAGAGTAAAAA

Spink2_3'UTR

5' GCATGATACTTGTTCACAGAACTTCTCTGGGTAGAGGCTGGAAGACAGTGATGTAAAGCGGTTTATA 3'
93   3'|||||||||||5'   158
ATCAAGTGTCTT (FJ541142)

Spink2 variant 3_5'UTR

            +/+ (FJ541120)
5' ATGGAAGGTGAATGTGGGTCAAGCATCCTCTGCCGTTATAGTTGGGGCTTCCATTCTAGCTCCAACT
1   10   21
AGATGTCACCCAGCCATCTGTTCTACCTTTCTTAAGCAGATTTTTCTGGACCTATATACGCAGTT........3'
                                         132
                            3'|||||||||||5'(FJ541101)
                            CTAAAAAGACCT
                            106   117

Sod_3'UTR

5' CATTCCCTGTGTGGTCTGAGTCTCAGACTCATCTGCTACCCTCAAACCATTAAACTGTAATCTGAAA 3'
3   3'|||||||||||5'   69
                       TAAGTAGACGA (FJ541118)

Fabp9_5'UTR

                    +/+ (DQ907162)
5' GCCCACACTTCATGGTTTTCGGTTGTGAATGCCTGGTCTGCCCAGAAAGCTGTGCACG 3'
1   58

Spot A Hypothetical protein_3'UTR

                246   1   259   420   2   434
5' AGAA....................................GAGACCCTGAATAA   AAAAATGGCTCTCAA
1   +/+ (FJ541097)   +/+ (FJ541075)

    469   3   482   506   4   518   689   5   704
..ATAAACTGAATTTT.........CTGTACATCCATA.....................AGCACTGAAACTGTCC
    +/+ (FJ541077)   3'|||||||||||5' (FJ541077)   +/+ (FJ541078)
                         GACATGTAGGTAT

    814   6   826   1083   7   1094
....CTGGAAGCAGCTG.................AAATACTTATAT.................TTCGCT 3'
   +/+ (FJ541083)   3'|||||||||||5'   1223
                            TTTATGAATATA (DQ907162)

Sdf2l1_3'UTR

5' GTATTAAAGGTGTACACCACCACAACTGGCCCCAACCTGTTCAATAAACTCATTTTTATTACTAA 3'
306   370
                             (FJ541120) 3'GAGTAAAAATA5'

B

Aromatase_3'UTR

5' GTGTGGCCTATCTACTCTGGGGCATTACTTATCCGTAATTACATGGAAACCATCCATTTTATCGAGTAG 3'
5   73
                          (FJ541089) 3'|||||||||||5'
                                       GGTAGGTAAAA

Caldendrin_3'UTR

5' GGGGAGAAGAGCTGAGCTCGCCTCACCCCCGCAGTAGCTGCCGAAAGCCCAGGATGTACTGGCGGA 3'
43   3'|||||||||||5' (FJ541087)   108
                       CGTCATCGACG
5' CCCAGGGCAGAGCCTCTCAGCGCCTTCCTGATGCCCGTCCCAGCTCGCCTAGCCCTTTGATCTCAGA 3'
340   3'|||||||||||5'   406
                 CGGAAGGACTA (FJ541087)

C

Acrosin_3'UTR

         +/+ (FJ541142)   +/+ (FJ541082)
5' GAAAAGAGAGATGAGTGAAATAAAGTATATATTTGTATATAAATATATATGTACACTCGAGGACATGAAC 3'
1   70
         +/+ (FJ541168)
5' CTTCAACTCAACCCAAATCTTCCTCCATCAAATTTACATCCTCCCTGCTCAGCTCCAATAAAATGGAAT 3'
80   148
                                        3'|||||||||||5'
                                         GACGAGTCGA (FJ541135)