Title: Large-scale gene losses underlie the genome evolution of parasitic plant

Cuscuta australis

Authors: Guiling Sun[1,4‡], Yuxing Xu[1,2,3‡], Hui Liu[1‡], Ting Sun[4], Jingxiong Zhang[1],

Christian Hettenhausen[1], Guojing Shen[1], Jinfeng Qi[1], Yan Qin[1], Jing Li[1], Lei Wang[1], Wei

Chang[1], Zhenhua Guo[2], Ian T. Baldwin[5], and Jianqiang Wu[1,*]

Affiliations:

[1]Department of Economic Plants and Biotechnology, Yunnan Key Laboratory for Wild

Plant Resources and [2]the Germplasm Bank of Wild Species, Kunming Institute of

Botany, Chinese Academy of Sciences, Kunming 650201, China.

[3]University of Chinese Academy of Sciences, Beijing 100049, China.

[4]Institute of Plant Stress Biology, State Key Laboratory of Cotton Biology, Department

of Biology, Henan University, Kaifeng 475001, China.

[5]Max Planck Institute for Chemical Ecology, Jena 07745, Germany.

[‡]These authors contributed equally.

* Correspondence and requests for materials should be addressed to J.W.

(wujianqiang@mail.kib.ac.cn).

Dodders (*Cuscuta* spp., Convolvulaceae) are globally distributed root- and leafless parasitic plants that parasitize a wide range of hosts. The physiology, ecology, and evolution of these obligate parasites are still poorly understood. A high-quality reference genome (size 266.74 Mb and contig N50 of 3.63 Mb) of *Cuscuta australis* was assembled. Our analyses reveal that *Cuscuta* experienced accelerated evolution, and *Cuscuta* and the convolvulaceous morning glory (*Ipomoea*) shared a common whole-genome triplication event before their divergence. Importantly, *C. australis* genome harbors only 19805 protein-coding genes, and 11.7% of the conserved orthologs in autotrophic plants are lost in *C. australis*. Many of these gene loss events likely result from the plant's parasitic lifestyle and large changes in its body plan. Moreover, comparison of the gene expression patterns in *Cuscuta* prehaustoria/haustoria and various tissues of closely related autotrophic plants suggests that *Cuscuta* haustorium genes largely evolved from roots. The *C. australis* genome provides important resources for studying the evolution of parasitism, regressive evolution, and evo-devo in plant parasites.

About 1% of the flowering plants are haustorial parasites[1], and some are responsible for severe yield losses in many crops. Although the evolutionary history of plant parasitism remains elusive, all parasitic plants, except the mycoheterotrophs, use a specialized haustorial organ to extract water and nutrients through vascular connections with the hosts[2]. The *Cuscuta* spp. (dodders) are typical obligate shoot parasites widely distributed worldwide comprising ~ 194 species, and is the only genus of parasites in Convolvulaceae (Solanales). *Cuscuta* plants have large changes of body plans, being leaf- and rootless throughout their lifecycles (Fig. 1a). They contain trace amounts of

chlorophyll, but cannot sustain themselves from their own photosynthesis. Although

whether *Cuscuta* plants are hemi- or holoparasites remains debatable, they could be

considered to be transitioning from hemiparasitism to holoparasitism. Through haustoria,

dodders not only obtain water and nutrients, but secondary metabolites, mRNAs, and

proteins from their host plants[3-5]. These features make *Cuscuta* an important model to

elucidate plant-parasite interactions and the evolution of plant parasites.

Results

Genome assembly

The genome size of *Cuscuta australis* was estimated to be 272.57 Mb from *k*-mer

analysis. We next generated 26.6 Gb (97.6-fold coverage) of *C. australis* genome

sequences from a single-molecule real-time (SMRT) sequencing platform. The *C.*

australis genome sequence includes 249 contigs (N50 = 3.63 Mb), and these contigs were

further assembled to form 103 scaffolds (N50 = 5.95 Mb; Supplementary Table 1). In

total, 264.83 Mb (219 contigs) of nuclear sequences (97.16% of the estimated genome

size) and 1.9 Mb (30 contigs) of organellar sequences were acquired (Supplementary

Table 2). The accuracy and heterozygosity were estimated to be 99.99% and 0.013%,

respectively (Supplementary Table 3). A total of 155 Mb of repetitive elements were

identified in the *C. australis* genome (Supplementary Table 4). Comparison with

Ipomoea nil (Japanese morning glory; also Convolvulaceae), which is relatively closely

related, indicated similar proportions of different types of repetitive elements between the

two genomes (Supplementary Table 4), although the *I. nil* genome contains many more

repeats than does the *C. australis* genome (Supplementary Table 4), consistent with its

larger genome size (734 Mb). LTRs are the most dominant type of repeats in both *C.*

australis and *I. nil* (Supplementary Table 4), and we specifically inspected the

LTR/Gypsy and *LTR/Copia* superfamilies. These two genomes exhibit large differences in

LTR/Gypsy (84,083 in *I. nil* and 24,453 in *C. australis*) and *LTR/Copia* (96,355 in *I. nil*

and 43,853 in *C. australis*) copy numbers; moreover, the sequences of *LTR/Gypsy* and

LTR/Copia members have also diverged as revealed by phylogenetic analysis

(Supplementary Fig. 1).

Based on *de novo* gene structure prediction, homology comparison, and transcript

data of both *C. australis* and the closely-related relative *C. pentagona*[6], 19671 genes

could be annotated.

Phylogeny analysis

The phylogenetic position of *Cuscuta* was determined using 1796 one-to-one

orthogroups (Supplementary Data 1b) identified from *C. australis*, *Arabidopsis thaliana*,

and six lamiids plants – the Solanales *Ipomoea nil*, *Solanum tuberosum* (potato), *Solanum*

lycopersicum (tomato), and *Capsicum annuum* (pepper), the Gentianales *Coffea*

canephora (coffee), and the Lamiales *Mimulus guttatus* (monkey flower) (for simplicity,

these seven autotrophic species are collectively named 7Ref-Species). Consistent with

their phylogenetic relationship, *Cuscuta* forms a sister group with *Ipomoea*, and we

estimated that these two lineages split ~ 33 million years ago (Supplementary Fig. 2).

Notably, *Cuscuta* shows a much longer branch than does *Ipomoea* (Fig. 1b), providing

genome-wide evidence consistent with the hypothesis that parasitic plants evolve

rapidly[7,8]. This result is statistically significant by both two cluster analysis and relative

rate test (Supplementary Table 5 and Supplementary Table 6). Previously, a whole-

genome duplication (WGD) event was detected in *Ipomoea*[9]. The syntenic blocks and

trees of the syntenic gene groups of *Cuscuta* and *Ipomoea* vs *Coffea* genome indicate that

Cuscuta and *Ipomoea* experienced a whole-genome triplication event before their

divergence from a common ancestor (Fig. 1c, d, Supplementary Fig. 3 to 5,

Supplementary Table 7).

Contractions and expansions of gene families

Parasitism and large changes of body plan in *Cuscuta* suggest that many gene

families might have experienced substantial alterations in sizes, including those that

function in leaf and root physiology. To study gene family expansion and contraction, a

rigorous bioinformatic pipeline was adopted to identify gene families (details see

Supplementary Note 2). In addition, the genome of *Utricularia gibba* (Lentibulariaceae;

an aquatic carnivorous bladderwort plant) was included in the analysis, given that *U.*

gibba also exhibits large changes in body plan (e. g. no true roots). We identified a total

of 13981 gene families in *C. australis*, *U. gibba*, and the 7Ref-Species (Supplementary

Data 1a); among these, 1256 and 478 families in *C. australis* and 605 and 848 families in

U. gibba were found to have had significant contractions and expansions, respectively,

revealed by a maximum-likelihood analysis (Fig. 2a; Supplementary Data 1a). Moreover,

box plots of the *F*-indices (details see Supplementary Note 2), which describe the

differences among the gene numbers of the conserved gene families in the 7Ref-Species

(namely, in *Arabidopsis* and at least five of the six remaining autotrophic species),

indicate that in *C. australi*s and *U. gibba*, gene numbers in 72 and 62% of the conserved

gene families are below the averages, respectively (Fig. 2b).

Overall gene losses

The drastic contractions of gene families in *C. australi*s and *U. gibba* suggest

strong gene losses in their genomes. To identify specific orthologous genes that are lost

in *C. australis* and *U. gibba*, we developed a stringent analysis pipeline to divide each

gene family into small orthogroups using a method combining phylogenetic and syntenic

analysis (details see Supplementary Note 3) and the functional annotations were assigned

using *Arabidopsis* as the reference (Supplementary Fig. 6). This analysis resulted in

21487 orthogroups (Supplementary Data 1b). Among the 11995 conserved orthogroups

in the 7Ref-Species, there are 1402 and 1555 orthogroups whose *C. australis* and *U.*

gibba members are absent, respectively (Supplementary Data 1b and 1c). Strikingly, 563

orthogroups have no *C. australis* and *U. gibba* orthologs, whose functions include

phytohormone pathways, nutrient uptake, defense response, and root hair development

(Supplementary Data 2a); 839 orthogroups specifically lost their members in *C. australis*,

and these genes are mainly involved in response to light, photosynthesis, chloroplast

RNA processing, and adventitious root development (Supplementary Data 2b); 992

orthogroups specifically have no members of *U. gibba*, and genes in these orthogroups

are involved in signaling, response to stimuli, and protein modifications, among others

(Supplementary Data 2c). In addition to the lost genes, in *C. australis* genome we

identified 1168 pseudogenes, which contain frame-shifts or premature stop codons.

Among these, we found five flowering time-related genes and 13 photosynthesis-related

genes (Supplementary Data 3a).

We next inspected the tissue-specific expression patterns of the *S. lycopersicum*

and *I. nil* (the closest autotrophs to *Cuscuta* among the 7Ref-Species) orthologs in the

1402 orthogroups, whose *Cuscuta* members are lost (Supplementary Data 1d). It was

found that these orthologs' principally expressed tissues (the tissues, in which the

expression levels of a given gene are at least 1-fold greater than the averages of its

expression levels in other tissues, are defined as the principally expressed tissues for that

gene) are the leaves and root of both *S. lycopersicum* and *I. nil* (Fig. 3a). These data are

consistent with the leaf- and rootless body plan of *Cuscuta*.

Loss of genes important for leaf and root development and nutrient uptake

Next, the *C. australis* genome was specifically searched for genes that mediate

leaf and root development, and it was found that a number of important genes involved in

leaf and/or root development are absent (Supplementary Data 1c): The orthologs of 1)

LCR (leaf shape and vein formation); 2) *TRN1/LOP1* (leaf patterning and lateral root

development); 3) *PLT1*, *2*, and *5* (specification of root stem cell niche); 4) *SMB* (lateral

root cap maturation); 5) all the *CASP* genes which are required for Casparian strip

formation; 6) *WAK1* to *5* (root cell expansion); 7) *WOX3*, *5*, and *7* (embryonic patterning,

stem cell maintenance, and organ formation). Moreover, many genes involved in

potassium (K), phosphate (P), and nitrate (N) uptake from soil are lost in *C. australis*. For

K^+ uptake: the orthologs of *HAK5* (a high affinity K^+ transporter), *KAT3/AtKC1* (a

general regulatory negatively modulating many inward Shaker K^+ channels), and *CHX20*

(a $Na^+(K^+)/H^+$ antiporter) are absent in *C. australis*. The following orthologs important

for P uptake are also lost: *PHO2* (P uptake and translocation), *PHT2;1* (a chloroplast P

transporter), and *SPX3* (P signaling). Several genes involved in N uptake are missing as

well, such as the orthologs of *NAXT1* (a nitrate efflux transporter), and *NLP7* (a positive

regulator for nitrate-induced gene expression); moreover, while the low-affinity nitrate

transporters *NRT1.1* and *1.2* are retained, the high-affinity nitrate transporters *NRT2.1*

and *2.2* and *NRT3.1* are absent.

Loss of photosynthesis genes

C. australis has very limited photosynthetic capability (Supplementary Fig. 7).

Among 248 photosynthesis-related genes in *Arabidopsis* annotated by GO, 38 *C.*

australis orthologs are missing (Supplementary Data 1c). The plastome of *C. australis*

(Supplementary Fig. 8) appears to have gene contractions (81 genes remain, while the

plastome of *I. nil* harbors 104 genes), and is similar to those of the previously sequenced

Cuscuta species[10,11] (Supplementary Data 4), especially *C. obtusiflora* and *C. gronovii*,

which belong to the subgenus *Grammica* with *C. australis*. Notably, all these *Cuscuta*

plastomes lack of *ndh* genes encoding proteins for forming the NADH dehydrogenase

complex functioning in electron cycling around photosystem I under stress. These data

concur with the finding that *ndh* genes are among the genes that are lost in the initial

stage in the evolution of parasitism in Orobanchaceae parasites[12], apparently due to

relaxed selective constraint (Supplementary Data 4).

Loss of genes controlling flowering time

Leaves play a critical role in perceiving environmental signals and thereby

modulate the physiology of their own as well as other plant parts, such as activating

flowering[13]. We speculate that the leafless dodders may have unique means of regulating

flowering time and many flowering time-control genes may have been lost. A database of

flowering time gene networks in *Arabidopsis* was recently constructed[14]. We found that

among the 295 coding genes listed in this database, 26 are lost (Supplementary Data 1c),

including the well-known *FLC*, *FRI*, *SVP*, *AGL17*, and *CO* (Fig. 3b). Moreover, the

circadian clock genes *ELF3* and *4*, *ARR3* and *4*, and *CDF1* and *3* are also missing. *FKF1*

and *CIB1*, which are essential in the photoperiod pathway, are also lost (Fig. 3b).

Flowering time is controlled by multiple pathways[15], and it appears that the vernalization,

temperature, autonomous, circadian clock, and photoperiod pathway all seem to be

nonfunctional (Fig. 3b). It is conceivable that dodders may eavesdrop on a host flowering

signal(s), hypothetically FT/florigen, to modulate their own flowering time, as much

earlier or later flowering times of *Cuscuta* than those of the hosts could conceivably

reduce *Cuscuta* fitness.

Loss of defense genes

Leaves and roots are the most common sites of pathogen and insect infestation.

We speculate that after the ancestor of *Cuscuta* became leaf- and rootless, relaxed

selection from pathogens and insects resulted in losses of defense genes. Specifically,

four gene families were inspected in detail (Fig. 3c). *R* (*resistance*) genes are the critical

components of plant immunity. Even though there is a possibility that the ancestor of

Convolvulaceae experienced a reduction of *R* genes after the split between

Convolvulaceae and other Solanales lineages, as *I. nil* has the smallest number of *R* genes

(148; Supplementary Data 3b) among the 7Ref-Species, the reduction of *R* genes in

Cuscuta is still drastic: *C. australis* genome harbors only 15 *R* genes (Fig. 3c and

Supplementary Data 3b). Terpenes function as defenses against insects and pathogens.

We predicted 9 terpene synthase genes (*TPS*s) in *C. australis*, and there are 31to 60 *TPS*s

in the 7Ref-Species (Supplementary Data 3c). Many P450 enzymes are involved in the

biosynthesis of plant secondary metabolites , which are required for adaptation to biotic

and abiotic stresses[16]. While more than 240 P450s are encoded by the genomes of the

7Ref-Species, *C. australis* genome harbors only 89 (Supplementary Data 3d). Receptor-

like kinases (RLKs) are important in plant development and resistance to abiotic and

biotic stresses. *C. australis* genome comprises only 339 *RLK* genes, much less than in the

7Ref-Species (at least 655 in potato) (Supplementary Data 3e). Consistently, several

well-studied genes in plant resistance to pathogens are also absent in *C. australis*,

including *EDS1*, *EDS5*, *FMO1*, *SAG101*, and *PAD4*, which are essential for plant

resistance to diseases, and *ALD1*, which is critical for systemic acquired resistance.

Origin of haustorium

The haustorium is a parasitic plant-specific organ, playing a critical role in

establishing parasitism. *C. australis* and *C. pentagona* RNA-seq data[6] were assembled

using the *C. australis* genome as the reference (mapping ratios 96 and 75%, respectively;

Supplementary Table 8). In *Cuscuta* prehaustoria and haustoria, we identified 2466

principally expressed genes (PEGs; at least one-fold greater than the average of the

expression levels in all the other tissues) belonging to 1299 orthogroups (Supplementary

Data 1d, Supplementary Data 3f). GO analysis on these PEGs indicated enrichment of

biological processes of metabolic process, transport, lignin and xyloglucan metabolism,

and transcriptional regulation genes (Supplementary Data 2d). This is consistent with the

haustorial function of transporting host substances and the findings that dynamic cell wall

remodeling in parasite haustoria is important in the establishment of parasitism[17-19].

These 1299 corresponding orthologs' principally expressed tissues in *S. lycopersicum* and

I. nil were mostly found to be the roots (Supplementary Fig. 9, Supplementary Data 1d).

These data imply that the evolution of *Cuscuta* haustorium may be related to expression

changes of genes involved in root development. Similarly, comparative transcriptome

analyses on three root parasites, *Triphysaria versicolor*, *Striga hermonthica*, and

Phelipanche aegyptiaca, revealed that parasitism genes are derived primarily from root

and floral tissues[17].

Next, we performed HYPHY analysis to obtain the genes that underwent positive

selection and relaxed purifying selection after the divergence between the ancestors of

Cuscuta and *Ipomoea* lineage, as these genes might be associated with the speciation of

Cuscuta and evolution of parasitism. GO terms including "response to hormones", "DNA

methylation", "regulation of transcription", and "cell wall-related metabolism" were

enriched from the 1124 positively selected genes (Supplementary Data 3g,

Supplementary Data 2e), and among these, 115 are principally expressed in

prehaustoria/haustoria (Supplementary Data 3f), including a pectin esterase, receptor-like

kinases, transcription factors, a serine carboxypeptidase, and transporters. We also found

that 3890 genes (Supplementary Data 3h) exhibited signatures of relaxed purifying

selection. The enriched GO terms include "terpenoid biosynthetic process", "nitrate

assimilation", "photosystem II assembly", and "regulation of signal transduction"

(Supplementary Data 2f), and 504 genes with relaxed purifying selection (Supplementary

Data 3f) are principally expressed in prehaustoria/haustoria, such as genes encoding

subtilisin-like proteases and ABC transporter B family member 15-like protein. Gene

family expansion, mainly caused by gene duplication, often leads to neofunctionalization

among the gene family members and is thought to be an important driving force in the

acquisition of novel phenotypes. Thus, we performed GO analysis on the 3099 expanded

gene family members of *C. australis* (Supplementary Data 3i, Supplementary Data 2g). It

was found that "response to auxin" and "DNA methylation" were enriched from the

members of expanded gene families; among them, 109 are principally expressed in

prehaustoria/haustoria (Supplementary Data 3f). These data are consistent with the

finding that many haustorial genes in Orobanchaceae parasites also experienced relaxed

purifying selection and/or positive evolution and may play an important role in the

evolution of the parasitic lifestyle[17].

Five positively selected genes from the expanded gene families were found to be

principally expressed in haustoria (Supplementary Fig. 10. and Supplementary Data 3f).

Among these, one encodes a putative α/ß-hydrolase highly similar to *Nicotiana sylvestris*

DAD2/DWARF14 (84% identity), which is the strigolactone receptor in autotrophic

plants, implying that neofunctionalization of α/ß-hydrolase genes might also be involved

in the parasitization process in *Cuscuta*, as they are in the root parasites *Striga* and

Orobanche [20].

Discussions

How plants evolved parasitism is still unclear. Our transcriptomic data and

molecular evolution analysis suggest convergent evolution in *Cuscuta* and

Orobanchaceae root parasites[17]. The haustorium of *Cuscuta* probably evolved from root

tissues and that of Orobanchaceae parasites is derived from both root and floral tissues.

Moreover, a relatively large fraction of the genes that experienced positive selection

and/or relaxed selection are principally expressed in prehausotria/haustoria in both

Cuscuta and Orobanchaceae, and these genes may be related to parasitization and/or

evolution of parasitism.

Our comparative genomic analyses indicate that *C. australis* genome experienced

remarkably high levels of contraction, and this is consistent with *Cuscuta*'s parasitic

lifestyle and large changes in body plan – leaf- and rootless as well as intensive

dependence on host-derived metabolites and signals. Given the importance of new genes

that bring about novel phenotypes, it is possible that the autotrophic ancestor evolved

haustorium from root tissues through the neofunctionalization of duplicated genes and

transcriptional reprogramming in the *Cuscuta* lineage. *Cuscuta* is transitioning from

hemiparasites to holoparasites. Among the recognized major parasite lineages, three

contain only hemiparasites, while eight are solely holoparasites[1], implying that

holoparasitization may have additional selective advantages over hemiparasitization.

Hypothetically, the dramatic changes in body plans, including the degeneration of leaf

and root could allow these parasites to reallocate carbon and nitrogen resources, that are

required for their development and growth, to reproduction and could also enable the

holoparasites to better adjust their physiology according to that of the hosts by

eavesdropping on host signaling molecules.

Wicke et al.[12] analyzed the plastomes of nonparasitic, hemiparasitic, and

nonphotosynthetic parasitic plants in Orobanchaceae, and found that the transition from

autotrophic plant to obligate parasites relaxes functional constraints on plastid genes in a

stepwise manner. Similar analyses could not be done in *Cuscuta*, as it is the only parasitic

lineage in the Convolvulaceae. The large body plan changes that were associated with the

parasitic lifestyle relaxed selective constraints on several core pathways, such as

photosynthesis[12], flowering time control, root and leaf development, nutrient acquirement,

and defense against pathogens and insects. Under relaxed selection pressure, some genes

of these pathways were pseudogenized and melted into the background of surrounding

DNA because of accumulation of recurrent mutations, or were even deleted from the

genome, finally leading to gene losses. In *C. australis* genome, 1168 genes were

identified as pseudogenes, although some may still be functional or even have gained

new functions[21], it is likely that the majority of these pseudogenes are degenerated or in

the process of being deleted, following the fate of the lost genes. Indeed, *C. australis*

genes undergone relaxed purifying selection include those related to terpenoid

biosynthetic process (defense), nitrate assimilation (nutrient acquirement), and

photosystem II assembly (photosynthesis). Resequencing of other *Cuscuta* species may

shed light on the recent gene loss events and the underlying mechanisms (e.g., transposon

insertion, fragment deletion, and rapid accumulation of mutations) in *Cuscuta*.

Large scale gene loss is also evident from the genomic data of human parasites

Pediculus humanus humanus (body louse)[22], *Trichinella spiralis* (a zoonotic nematode)[23],

and *Giardia lamblia* (an intestinal protist)[24], and from an *Arabidopsis* pathogen, the

obligate biotrophic oomycete *Hyaloperonospora arabidopsidis*[25]. Here, we also detected

that *U. gibba* genome also experienced a large number of gene loss events during its

evolution, likely resulting from its body plan changes (mainly loss of root) and lifestyle

alteration (carnivory). It would also be of importance to compare the genomes of *Cuscuta*

and those of Orobanchaceae root parasites. We expect that Orobanchaceae parasites,

especially holoparasites, such as *Orobanche* species, may also have a large number of

lost nuclear genes, some of which may bear the same functions as those in *Cuscuta*, such

as the genes that function in leaf development and photosynthesis. The *C. australis*

genomic data strongly support the notion that regressive evolution, which is associated

with extensive gene loss, is likely to be pervasive and adaptive during the evolution of

holo- /obligate parasites[26,27]. Comparative genomics between the genome data of *C.*

australis and autotrophic plants provides an important resource for studying genome

reduction, regressive evolution, parasitism, and evo-devo in plant parasites.

Methods

DNA sample preparation and sequencing. The seeds of *Cuscuta australis* were

originally purchased from a Chinese traditional medicinal herbs store in Kunming, China,

in 2011, and had been cultivated for five generations in a glasshouse at the Kunming

Institute of Botany, Chinese Academy of Sciences. Voucher specimens of *C. australis*

can be accessed at the Herbarium of the Kunming Institute of Botany, Chinese Academy

of Sciences (accessions Nos. WJQ-001-1 and WJQ-001-3). Seedlings were prepared from

seeds of the fifth generation and were infested on soybean plants (for the germination

procedure, see Li et al.[28]). DNA isolated from young vines collected from one individual

Cuscuta australis was isolated for genomic library construction. Three genomic DNA

libraries with 350-bp, 2-kb, or 5-kb insertions were constructed for Illumina sequencing.

For PacBio sequencing, five DNA libraries with 20-kb insertions were sequenced on a

PacBio RS II instrument using the P6v2 polymerase binding and C4 chemistry kits (P6-

C4). A total of 26.6 Gb from 1,953,966 reads were obtained by processing 24 single-

molecule real-time (SMRT) cells. The average and N50 of SMRT subread length were

9.6 kb and 13.6 kb, respectively.

Transcriptome sample preparation and sequencing. *Cuscuta australis* tissues of

seeds, just-germinated seeds (one day after imbibing), seedlings (five days after

imbibing), prehaustoriua, haustoria, stems far from haustoria, stems near haustoria,

flower buds, flowers, and seed capsules were collected and RNA samples were extracted

from these tissues using the TRI Reagent (Sigma). Libraries were constructed for each

tissue according to the TrueSeq® RNA Sample Preparation protocol, and sequenced on

an Illumina HiSeq-2500. Sequences are deposited in NCBI under BioProject

PRJNA394036.

Cuscuta pentagona transcriptomic short reads dataset from Ranjan et al.[6] were retrieved

from the NCBI Short Read Archive under accession numbers SRR965929, SRR965963,

SRR966236,   SRR966405,   SRR966412,   SRR966513,   SRR966542,   SRR966549,

SRR966619 to SRR966622, SRR967154, SRR967164, SRR967181 to SRR967190,

SRR967275 to SRR967289, and SRR967291.

Genome survey. We used 23.29 Gb of HiSeq reads to estimate the genome features

using the GCE[29] (v1.0.0) software based on k-mer depth-frequency distribution. A total

of 11,700,637,064 17-mers were counted. Given the unique k-mer depth of 42, we

calculated that the genome size = KmerCount/Depth = 272.57 Mb. The repeat content

was estimated to be 58.94% based on k-mer depth distribution.

De novo **assembly.** SMRT reads were corrected, trimmed, and assembled using CANU[30]

(v1.3), a genome assembler built on the basis of Celera assembler. Briefly, SMRT reads

were firstly self-corrected based on an overlap-layout algorithm. Erroneous regions of

error-corrected SMRT reads were then trimmed to increase accuracy. We constructed an

initial assembly using CANU with those trimmed error-corrected SMRT reads following

the parameters "genomeSize=273m errorRate=0.025" and then used Quiver[31] (v4.0) to

generate consensus sequences by aligning SMRT reads to correct the errors in the

assembly. Lastly, we used Pilon[32] (v1.18) to perform the second round of error correction

with HiSeq reads from the 350-bp-insert library. The resulted error-corrected assembly

was named version 1.0 and used in the subsequent analyses. A hierarchical method was

also applied to concatenate adjacent contigs: SSPACE[33] (v 3.0) was first used to build

scaffolds using HiSeq data from the mate-pair libraries and the contigs built from the

PacoBio data. N50 of the resulted scaffolds reached 5.9 Mb. SSPACE-LongRead[34] (v1-1)

was further applied to build superscaffolds with PacBio long reads, nevertheless, no

linking information between scaffolds were found and no improvements were acquired

(Supplementary Table 1).

Assembly completeness, accuracy, and heterozygosity assessment. The completeness

of the PacBio assembly was estimated by checking a set of genes widely conserved in

eukaryotic genomes with CEGMA[35] (v2.5). The accuracy and heterozygosity rate of the

genome assembly were estimated using the following procedure: Adaptors were removed

from the short paired-end reads obtained from the 350-bp insert size library, and then the

reads were aligned to the PacBio assembly, which had been corrected with Pilon[32]

(v1.18), using bowtie2[36] (v 2.2.4) to generate a bam file. A total of 97.9% reads could be

mapped. Samtools[37] (v 1.3.1) were used to sort bam file, and Freebayes[38] (v1.1.0) was

applied to call variants. Compared with PacBio assembly as the reference genome,

homozygous SNPs or indels were considered to be assembly errors, and heterozygous

SNPs or indels were regarded as heterozygous sites. The accuracy was estimated to be

99.99% and heterozygosity was estimated to be 0.013% (Supplementary Table 3). Error

rate of the genome assembly before Pilon correction was also estimated using the same

set of methods. A total of 980 SNPs and 101825 Indels were corrected after polishing.

Exclusion of organelle-derived contigs. We excluded the organelle-derived sequences

based on the read mapping depths and the characteristic genes and structures of

organellar genomes. Firstly, the Illumina pair-end reads were mapped to the PacBio

genome assembly for calculating the sequencing depths in each contigs, and the 60

contigs with more than 100 × depths were selected. The previously reported plastidial

and/or mitochondrial genomes of *Cuscuta gronovii*, *Cuscuta reflexa*, *Ipomonea nil*, and

Arabidopsis thaliana (accession Nos. NC_009765.1, NC_009766.1, NC_031159.1,

NC_031158.1, KX551970.1, Y08501.2 respectively) were retrieved and used as queries

to search these 60 contigs using blastn[39] (v2.4.0) and tblastx[39] (v2.4.0) to mark the coding

regions, and the RepBase database was employed to mark noncoding and repetitive

regions using Repeatmasker[40] (v4.0.6). The contigs with the characteristic genes and

genome structures of plastids and mitochondria were manually inspected and removed

from the PacBio assembly.

In total, 30 contigs (Supplementary Table 2) comprising 1,906,600 bp were

identified as organellar sequences, each with sequencing depths ranging from 582.3 to

3267.5 ×, and one contig C141C with a length of 85,214 bp shows the characteristics of a

complete circular plastid genome (Supplementary Fig. 8).

Illumina reads cleanup and assembly. *de novo* assembling the Illumina short reads was

performed based on the previously reported pipeline Phusion-meta (v2.3) with several

minor modifications[41]. Briefly, after removing low-quality reads containing ten or more

unique *k*-mers, the paired-end reads were clustered into thousands of groups by

Phusion2[42] (v2.3) with *k*-mer at 51 bp. The reads within each cluster were assembled into

contigs with N50 of 17 kb using the Fermi[43] (v1.1) and SOAPdenovo[44] (v2.04) software.

All the obtained contigs from above were merged by the aligner GAP5[45] (v2.0.0b11),

which were then used to build the scaffolds with the mate-paired reads from the 2-k and

5-k libraries hierarchically and iteratively with SOAPdenovo[44] (*k*-mer at 81 bp), resulting

in an assembly with the N50 of 214 kb and length of 266.9 Mb.

Identification of repetitive elements. To identify repetitive elements in *Cuscuta*

australis, we first used *de novo* prediction to build a custom database of repeats in

Cuscuta australis with RepeatModeler[40] (v1.0.4). Then we used Repeatmasker[40] (v4.0.6)

to annotate the repetitive regions in *Cuscuta australis* genome based on our custom

database and the models of repeats from RepBase[46] (release 22.10). TRF[47] (v4.0.7b) was

used to improve finding of simple tandem repeats. After the redundant results were

removed, the repetitive elements were annotated using Repeatmasker (Supplementary

Table 4). In this manner, 58% of the genome regions were identified as the repetitive

regions. To facilitate comparison of the repeats between *Cuscuta australis* and *Ipomoea*

nil, the same pipeline was also used for the genome of *Ipomoea nil*. A total of 2138 and

2143 families of repetitive elements were identified in *Cuscuta australis* and *Ipomoea nil*,

respectively. To compare the evolutionary dynamics of LTRs, the phylogenetic trees for

two largest families, *LTR/Copia* and *LTR/Gypsy*, were reconstructed (Supplementary Fig.

1). Sequences of *LTR/Copia* and *LTR/Gypsy* were first aligned using famsa[48] (v1.2.1),

and the phylogenetic trees were then reconstructed using Fasttree[49] (v2.1.9).

Gene model prediction. Three different methods were employed for gene model

prediction:

1. Homology-based prediction

Protein sequences of 7Ref-Species (*Arabidopsis thaliana*, *Ipomoea nil*, *Solanum*

tuberosum, *Solanum lycopersicum*, *Capsicum annuum*, *Coffea canephora*, and *Mimulus*

guttatus) were first aligned to the masked genome of *Cuscuta australis* using tblastn[39]

(v2.4.0) with e-value < 1e-5 to find corresponding homologous regions in *Cuscuta*

australis. Genewise[50] (v2.0) was then utilized to generate accurate gene structures in

these regions using the protein sequences of 7Ref-Species as the references.

2. Transcriptome-based prediction

RNA-seq data from different tissues of *Cuscuta australis* were aligned to the masked

genome using TopHat[51] (v2.0.8). Cufflinks[52] (v2.1.1) was then used to construct genes.

To find genes supported by *Cuscuta pentagona* transcript evidence, we assembled

transcripts of *Cuscuta pentagona* and aligned against the genome of *Cuscuta australis*

using blat[53] (v. 35).

3. *De novo* gene prediction

Gene models of *Cuscuta australis* were first predicted using *de novo* prediction

methods Augustus[54] (v2.5.5), GlimmerHMM[55] (v3.0.1), GeneScan[56] (v1.0), SNAP[57]

(version 2006-07-28), and Geneid[58] (v1.4) . Training set for *de novo* prediction included

the complete gene model information from 7Ref-Species and the transcriptome data

obtained from *Cuscuta australis* and *Cuscuta pentagona* (assembled using Trinity[59]

(r20140717) and structurally annotated with PASA[60] (v2.0.2)).

De novo prediction, homology-based prediction, and transcriptomic evidence were

further integrated using EVM[61] (v1.1.1), resulting in a non-redundant complete gene-set.

UTR regions were identified by integrating transcript evidence with gene annotation

results using PASA[60] (v2.0.2).

Pseudogene identification. Given that some of pseudogenes may be mistakenly

identified as retrotransposons, we firstly remasked the genome with a lowered standard:

Retrotranspons and DNA transposons were identified with RepeatMasker[40] (v4.0.6)

based on the RepBase[46] database (release 22.10), other than *de novo* prediction as

previously described. Thereafter, low-complexity repeats were identified using Tandem

Repeats Finder[47] (v4.0.7b). The interspersed repeats were hard-masked (masked as N).

Considering some proteins also contain tandem repeats, low complexity repeats were

soft-masked (masked as lowercase) so that a blast hit can extend from a seed hit to the

low-complexity repetitive regions.

The procedure of pseudogene identification was adopted from Zou et al[62]. To

identify the locations of the homologs of all the gene models of 7Ref-Species in *Cuscuta*

australis genome, all protein sequences from the 7Ref-Species genomes were used as

queries to search against the remasked genome of *Cuscuta australis* using tblastn[39]

(v2.4.0). A total of 18,823 homologous regions were found. After removing those

overlapping with those previously identified gene loci, structures of genes missed in the

previous annotation were predicted using GeneWise[50] (v2.0), which built fine alignments

between query protein sequences and the homologous regions in *Cuscuta australis*.

Predicted gene structures with mutations leading to frame-shifts and premature stop

codons were selected as candidate pseudogenes. As some of the query sequences might

be mistakenly annotated in structures, false positives of pseudogenes could be identified.

To minimize this possibility, we manually examined if the mutations could be supported

by most of the query sequences in multiple sequence alignments. Genes passed the

manual inspections were considered to be pseudogenes. Newly annotated genes without

evidence of being pseudogenes were added to the previously annotated gene-set to

improve the completeness for gene loss analysis later. To this end, we obtained 19,805

protein-coding genes and 1,283 pseudogenes.

Functional annotation. The annotated protein sequences were searched against KEGG

(release 53), NR (version 20150810), and Swiss-Prot (version 20150821) using blastp[39]

(v2.4.0) (e-value < 1e-5), and the best hits were retained as putative function annotations.

InterproScan (v4.8) were used to identify domain information in *Cuscuta australis* genes.

In total, 90.2% of all the genes were functionally annotated.

URL. Genome website: http://www.dodderbase.org.

Data availability. The genome assembly, gene models, and sequence reads are available

at the NCBI under the BioProject PRJNA394036, and the former two can also be

accessed at http://www.dodderbase.org.

References

1   Westwood, J. H., Yoder, J. I., Timko, M. P. & dePamphilis, C. W. The evolution
of parasitism in plants. *Trends Plant Sci* **15**, 227-235,
doi:10.1016/j.tplants.2010.01.004 (2010).
2   Yoshida, S., Cui, S., Ichihashi, Y. & Shirasu, K. The haustorium, a specialized
invasive organ in parasitic plants. *Annu Rev Plant Biol* **67**, 643-667,
doi:10.1146/annurev-arplant-043015-111702 (2016).
3   Kim, G., LeBlanc, M. L., Wafula, E. K., Depamphilis, C. W. & Westwood, J. H.
Genomic-scale exchange of mRNA between a parasitic plant and its hosts.
Science **345**, 808-811, doi:10.1126/science.1253122 (2014).
4   Birschwilks, M., Haupt, S., Hofius, D. & Neumann, S. Transfer of phloem-mobile
substances from the host plants to the holoparasite *Cuscuta* sp. *J Exp Bot* **57**, 911-
921, doi:10.1093/Jxb/Erj076 (2006).
5   Smith, J. D., Woldemariam, M. G., Mescher, M. C., Jander, G. & De Moraes, C.
M. Glucosinolates from host plants influence growth of the parasitic plant
Cuscuta gronovii and its susceptibility to aphid feeding. *Plant Physiol* **172**, 181-
197, doi:10.1104/pp.16.00613 (2016).
6   Ranjan, A. *et al.* De novo assembly and characterization of the transcriptome of
the parasitic weed dodder identifies genes associated with plant parasitism. *Plant*
Physiol **166**, 1186-1199, doi:10.1104/pp.113.234864 (2014).
7   Estep, M. C., Gowda, B. S., Huang, K., Timko, M. P. & Bennetzen, J. L.
Genomic characterization for parasitic weeds of the genus *Striga* by sample
sequence analysis. *Plant Genome-Us* **5**, 30-41,
doi:10.3835/plantgenome2011.11.0031 (2012).
8   Bromham, L., Cowman, P. F. & Lanfear, R. Parasitic plants have increased rates
of molecular evolution across all three genomes. *BMC Evol Biol* **13**, 126,
doi:10.1186/1471-2148-13-126 (2013).
9   Hoshino, A. *et al.* Genome sequence and analysis of the Japanese morning glory
Ipomoea nil. *Nat Commun* **7**, 13295, doi:10.1038/ncomms13295 (2016).
10   Funk, H. T., Berg, S., Krupinska, K., Maier, U. G. & Krause, K. Complete DNA
sequences of the plastid genomes of two parasitic flowering plant species,
Cuscuta reflexa and Cuscuta gronovii. *BMC Plant Biol* **7**, 45, doi:10.1186/1471-
2229-7-45 (2007).
11   McNeal, J. R., Arumugunathan, K., Kuehl, J. V., Boore, J. L. & Depamphilis, C.
W. Systematics and plastid genome evolution of the cryptically photosynthetic
parasitic plant genus Cuscuta (Convolvulaceae). *BMC Biol* **5**, 55,
doi:10.1186/1741-7007-5-55 (2007).
12   Wicke, S. *et al.* Mechanistic model of evolutionary rate variation en route to a
nonphotosynthetic lifestyle in plants. *Proc Natl Acad Sci USA* **113**, 9045-9050,
doi:10.1073/pnas.1607576113 (2016).
13   Putterill, J. & Varkonyi-Gasic, E. FT and florigen long-distance flowering control
in plants. *Curr Opin Plant Biol* **33**, 77-82, doi:10.1016/j.pbi.2016.06.008 (2016).
14   Bouche, F., Lobet, G., Tocquin, P. & Perilleux, C. FLOR-ID: an interactive
database of flowering-time gene networks in Arabidopsis thaliana. *Nucleic Acids*
Res **44**, D1167-1171, doi:10.1093/nar/gkv1054 (2016).

Romera-Branchat, M., Andres, F. & Coupland, G. Flowering responses to seasonal cues: what's new? *Curr Opin Plant Biol* **21**, 120-127, doi:10.1016/j.pbi.2014.07.006 (2014).

Morant, M., Bak, S., Moller, B. L. & Werck-Reichhart, D. Plant cytochromes P450: tools for pharmacology, plant protection and phytoremediation. *Curr Opin Biotechnol* **14**, 151-162 (2003).

Yang, Z. *et al.* Comparative transcriptome analyses reveal core parasitism genes and suggest gene duplication and repurposing as sources of structural novelty. *Mol Biol Evol* **32**, 767-790, doi:10.1093/molbev/msu343 (2014).

Johnsen, H. R. *et al.* Cell wall composition profiling of parasitic giant dodder (*Cuscuta reflexa*) and its hosts: a priori differences and induced changes. *The New phytologist* **207**, 805-816, doi:10.1111/nph.13378 (2015).

Olsen, S. *et al.* Getting ready for host invasion: elevated expression and action of xyloglucan endotransglucosylases/hydrolases in developing haustoria of the holoparasitic angiosperm Cuscuta. *J Exp Bot* **67**, 695-708, doi:10.1093/jxb/erv482 (2016).

Conn, C. E. *et al.* PLANT EVOLUTION. Convergent evolution of strigolactone perception enabled host detection in parasitic plants. *Science* **349**, 540-543, doi:10.1126/science.aab1140 (2015).

Balakirev, E. S. & Ayala, F. J. Pseudogenes: Are They "Junk" or Functional DNA? *Annual Review of Genetics* **37**, 123-151, doi:10.1146/annurev.genet.37.040103.103949 (2003).

Kirkness, E. F. *et al.* Genome sequences of the human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle. *P Natl Acad Sci USA* **107**, 12168-12173, doi:10.1073/pnas.1003379107 (2010).

Mitreva, M. *et al.* The draft genome of the parasitic nematode *Trichinella spiralis*. *Nat Genet* **43**, 228-U274, doi:10.1038/ng.769 (2011).

Morrison, H. G. *et al.* Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science* **317**, 1921-1926, doi:10.1126/science.1143837 (2007).

Baxter, L. *et al.* Signatures of adaptation to obligate biotrophy in the *Hyaloperonospora arabidopsidis* Genome. *Science* **330**, 1549-1551, doi:10.1126/science.1195203 (2010).

Albalat, R. & Canestro, C. Evolution by gene loss. *Nat Rev Genet* **17**, 379-391, doi:10.1038/nrg.2016.39 (2016).

Wolf, Y. I. & Koonin, E. V. Genome reduction as the dominant mode of evolution. *Bioessays* **35**, 829-837, doi:10.1002/bies.201300037 (2013).

Li, J. *et al.* The Parasitic Plant Cuscuta australis Is Highly Insensitive to Abscisic Acid-Induced Suppression of Hypocotyl Elongation and Seed Germination. *Plos One* **10**, e0135197, doi:10.1371/journal.pone.0135197 (2015).

Liu, B. *et al.* Estimation of genomic characteristics by analyzing k-mer frequency in de novo genome projects. Preprint at http://arXiv:1308.2012 (2012).

Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* **27**, 722-736, doi:10.1101/gr.215087.116 (2017).

31 Chin, C. S. *et al.* Nonhybrid, finished microbial genome assemblies from long-
read SMRT sequencing data. *Nat Methods* **10**, 563-+, doi:10.1038/Nmeth.2474
(2013).
32 Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial
Variant Detection and Genome Assembly Improvement. *Plos One* **9**,
doi:10.1371/journal.pone.0112963 (2014).
33 Boetzer, M., Henkel, C. V., Jansen, H. J., Butler, D. & Pirovano, W. Scaffolding
pre-assembled contigs using SSPACE. *Bioinformatics* **27**, 578-579,
doi:10.1093/bioinformatics/btq683 (2011).
34 Boetzer, M. & Pirovano, W. SSPACE-LongRead: scaffolding bacterial draft
genomes using long read sequence information. *BMC Bioinformatics* **15**, 211,
doi:10.1186/1471-2105-15-211 (2014).
35 Parra, G., Bradnam, K. & Korf, I. CEGMA: a pipeline to accurately annotate core
genes in eukaryotic genomes. *Bioinformatics* **23**, 1061-1067,
doi:10.1093/bioinformatics/btm071 (2007).
36 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat
Methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).
37 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
38 Garrison E, M. G. Haplotype-based variant detection from short-read sequencing.
arXiv **preprint arXiv:1207.3907 [q-bio.GN]** (2012).
39 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic Local
Alignment Search Tool. *J Mol Biol* **215**, 403-410, doi:DOI
10.1006/jmbi.1990.9999 (1990).
40 Smit, A., Hubley, R & Green, P. *RepeatMasker Open-4.0*,
<http://www.repeatmasker.org> (2013-2015).
41 Peng, Z. *et al.* The draft genome of the fast-growing non-timber forest species
moso bamboo (Phyllostachys heterocycla). *Nat Genet* **45**, 456-461, 461e451-452,
doi:10.1038/ng.2569 (2013).
42 Mullikin, J. C. & Ning, Z. The phusion assembler. *Genome Res* **13**, 81-90,
doi:10.1101/gr.731003 (2003).
43 Li, H. FermiKit: assembly-based variant calling for Illumina resequencing data.
Bioinformatics **31**, 3694-3696, doi:10.1093/bioinformatics/btv440 (2015).
44 Luo, R. *et al.* SOAPdenovo2: an empirically improved memory-efficient short-
read de novo assembler. *Gigascience* **1**, 18, doi:10.1186/2047-217X-1-18 (2012).
45 Bonfield, J. K. & Whitwham, A. Gap5--editing the billion fragment sequence
assembly. *Bioinformatics* **26**, 1699-1703, doi:10.1093/bioinformatics/btq268
(2010).
46 Bao, W. D., Kojima, K. K. & Kohany, O. Repbase Update, a database of
repetitive elements in eukaryotic genomes. *Mobile DNA-Uk* **6**,
doi:10.1186/s13100-015-0041-9 (2015).
47 Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic
Acids Research* **27**, 573-580, doi:DOI 10.1093/nar/27.2.573 (1999).
48 Deorowicz, S., Debudaj-Grabysz, A. & Gudys, A. FAMSA: Fast and accurate
multiple sequence alignment of huge protein families. *Sci Rep* **6**, 33964,
doi:10.1038/srep33964 (2016).

49   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2-Approximately Maximum-
Likelihood Trees for Large Alignments. *Plos One* **5**,
doi:10.1371/journal.pone.0009490 (2010).
50   Birney, E., Clamp, M. & Durbin, R. GeneWise and Genomewise. *Genome Res*
14, 988-995, doi:10.1101/gr.1865504 (2004).
51   Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions
with RNA-Seq. *Bioinformatics* **25**, 1105-1111, doi:10.1093/bioinformatics/btp120
(2009).
52   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-578,
doi:10.1038/nprot.2012.016 (2012).
53   Kent, W. J. BLAT - The BLAST-like alignment tool. *Genome Research* **12**, 656-
664, doi:10.1101/gr.229202 (2002).
54   Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: a web
server for gene finding in eukaryotes. *Nucleic Acids Research* **32**, W309-W312,
doi:10.1093/nar/gkh379 (2004).
55   Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two
open source ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878-2879,
doi:10.1093/bioinformatics/bth315 (2004).
56   Burge, C. & Karlin, S. Prediction of complete gene structures in human genomic
DNA. *J Mol Biol* **268**, 78-94, doi:DOI 10.1006/jmbi.1997.0951 (1997).
57   Leskovec, J. & Sosic, R. SNAP: A General-Purpose Network Analysis and
Graph-Mining Library. *Acm T Intel Syst Tec* **8**, doi:10.1145/2898361 (2016).
58   Parra, G., Blanco, E. & Guigo, R. GeneID in Drosophila. *Genome Research* **10**,
511-515, doi:DOI 10.1101/gr.10.4.511 (2000).
59   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data
without a reference genome. *Nature Biotechnology* **29**, 644, doi:10.1038/nbt.1883
(2011).
60   Haas, B. J. *et al.* Improving the Arabidopsis genome annotation using maximal
transcript alignment assemblies. *Nucleic Acids Research* **31**, 5654-5666,
doi:10.1093/nar/gkg770 (2003).
61   Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using
EVidenceModeler and the program to assemble spliced alignments. *Genome Biol*
9, doi:10.1186/gb-2008-9-1-r7 (2008).
62   Zou, C. *et al.* Evolutionary and expression signatures of pseudogenes in
Arabidopsis and rice. *Plant Physiol* **151**, 3-15, doi:10.1104/pp.109.140632
(2009).
660

Acknowledgments This work was supported by the Strategic Priority Research Program

of the Chinese Academy of Sciences (CAS) (No. XDB11050200; J.W.). We also thank

Drs. Lei Wang (Institute of Botany, CAS), Hongtao Liu (Institute of Plant Physiology

and Ecology, CAS), Jinyong Hu and Chengjun Zhang (Kunming Institute of Botany,

CAS), and Zhaojun Ding (Shangdong University) for valuable comments.

Author Contributions J.W., G.S., and Y.X. designed and conceived research; T.S., C.H.,

G.S., prepared plant tissues; W.C. performed photosynthesis experiments; T.S., and Y.Q.

prepared DNA samples; J.L. and J.Q. prepared RNA samples; Y.X., and H.L. performed

genome assembly and genome annotation; Y.X., H.L, G.S., and T.S. analyzed data; J.W.,

G.S., Y.X., H.L., J.Z., L.W., I.T.B., and Z.G. wrote the paper. All authors read and

approved the final manuscript.

Materials & Correspondence

C. australis seeds can be distributed by Dr. Jianqiang Wu (Kunming Institute of

Botany, Chinese Academy of Sciences, email: wujianqiang@mail.kib.ac.cn) up request.

Figure 1 | Morphological traits and genome structure of *C. australis*. a, Photographs of *C. australis* seed (1), seedling (2), vines twining around the wild tomato *Solanum pennellii* (3 & 4; partial haustoria can be seen in 3), flowers (5), and seed capsules (6). **b**, Phylogenetic tree generated from genome-wide one-to-one orthogroups (bootstrap values for all clades are 100%). **c**, Circos plot of a set of syntenic genome segments of *C. australis*, Japanese morning glory, and coffee. Numbers besides terminals of each karyotype denote the start and end of chromosome segment or contigs with unit of Mb. **d**, Numbers of gene clades (shown on top of the trees)

703   supporting different hypotheses on the order of speciation and whole-genome triplication event

704   in the *Cuscuta* and *Ipomoea* lineage.

Figure 2 | Expansion and contraction in *C. australis* gene families. **a**, Significantly expanded

and contracted gene families. Brackets above each branch indicate numbers of expanded (in

black, before comma) and contracted (in red, after comma) gene families. **b**, Box plot overview

of the differences among the gene numbers of the conserved gene families, based on the *F*-index

values (*F*-indices range from 0 to 1; when *F* =, <, or > 0.5, the gene number in the given gene

family is equal to, smaller, or greater than the average size of this gene family in all species).

c

	R	TPS	P450	RLK
Cuscuta australis	15	9	89	339
Ipomoea nil	148	60	334	992
Solanum tuberosum	390	31	410	655
Solanum lycopersicum	264	34	255	685
Capsicum annuum	311	53	455	701
Coffea canephora	784	43	451	785
Mimulus guttatus	336	53	346	720
Arabidopsis thaliana	170	32	249	670

Figure 3 | Gene losses in *C. australis*. a, The principally expressed tissues (PETs) of the

respective orthologs of *C. australis* lost genes in *S. lycopersicum* and *I. nil*. The boxes represent

different tissues. The respective PETs of the orthogroups, which have no *C. australis* members,

were identified in *S. lycopersicum* and *I. nil*. The numbers of orthogroups, which have PETs in

leaves, roots, flowers, and other tissues, are shown in the boxes, and the respective percentages

(proportional to the areas of the boxes) indicate the ratios between these indicated numbers and

the numbers of all orthogroups whose PETs were identified to be the corresponding tissues of *I.*

nil and *S. lycopersicum.* **b**, Simplified gene network controlling flowering time. Genes in green

boxes are retained in *C. australis*, and the lost ones are in red boxes. Arrows and T-ends

represent promoting and inhibiting genetic interactions, respectively, and round dots at both ends

symbolize genetic interactions with unknown directions. **c**, Numbers of genes in the gene

families of *R* genes, *TPS*s, *P450*s, and *RLK*s.