Quantitative variation in female sensory structures supports species recognition and

intraspecific mate choice functions in damselflies

Alexandra A. Barnard[1,2] and John P. Masly[1]

[1]Ecology & Evolutionary Biology Program, Department of Biology, University of Oklahoma, Norman, OK, USA
[2]Email: alex.barnard@ou.edu

Keywords: sensilla, reproductive character displacement, reproductive isolation, Odonata, sexual signal

Data archived: The data included in this manuscript will be submitted to Dryad Digital Repository.

Author contributions: AAB and JPM planned the study; AAB collected the data and performed the analyses; AAB and JPM wrote the paper

Abstract

Males and females exchange signals prior to mating that convey information such as sex, species identity, or individual condition. Tactile signals relayed during physical contact between males and females before and during mating appear to be important for mate choice and reproductive isolation in some animals. However, compared to our understanding of visual, auditory, and chemical signals, we know little about the importance of tactile signals in mating decisions. Among North American damselflies in the genus *Enallagma* (Odonata: Coenagrionidae) species-specific tactile stimulation contributes to reproductive isolation between species and may also be important for intraspecific mate choice. We quantified several mechanosensory sensilla phenotypes on the female thorax among multiple sympatric and allopatric populations of two

Enallagma species that occasionally interbreed in nature. Although each species differed in

features of sensilla distribution within the thoracic plates, we found no strong evidence of

reproductive character displacement among the sensilla traits we measured in regions of

sympatry. However, substantial variation of sensilla traits was observed within populations of

both species. Our results suggest that species-specific placement of female mechanoreceptors

appears sufficient for species recognition, but mechanosensor variation among females within

species may be important for mate choice.

For sexual species, maintenance of species boundaries relies on reproductive isolation

(RI) between recently diverged species (Mayr 1942). Premating reproductive isolating barriers,

including behavioral or sexual isolation, often evolve earlier in the speciation process than

postmating barriers in a variety of animal taxa (*e.g.*, McMillan et al. 1997; Price and Bouvier

2002; Mendelson and Wallis 2003; Dopman et al. 2010; Sánchez-Guillén et al. 2012; Williams

and Mendelson 2014; Castillo et al. 2015; Barnard et al. 2017). Behavioral isolation requires

that mate recognition signals and/or preferences diverge between populations, which ultimately

results in the ability for individuals to discriminate conspecifics from heterospecifics. Species

recognition signals may rely on a variety of sensory modalities such as color (Wiernasz and

Kingsolver 1992; Sætre et al. 1997; Jiggins et al. 2001; Boughman et al. 2005; Kronforst et al.

2006; Williams and Mendelson 2014), courtship behavior (Stratton and Uetz 1986),

sound/vibration (Ewing and Bennet-Clark 1968; Wells and Henry 1998; Shaw 2000; Gerhardt

and Huber 2002; Arthur et al. 2013), and volatile chemicals (Coyne et al. 1994; Noor and Coyne

1996; Trabalon et al. 1997; Rafferty and Boughman 2006). Often, multiple signals act in concert

to affect species recognition (*e.g.*, Costanzo and Monteiro 2007; Girard et al. 2015).

Although much is known about the importance of visual, chemical, and auditory signals

and responses in sexual communication and species recognition, we know relatively little about

other sensory modalities that may have strong effects on individual mating decisions. Tactile

signals have been hypothesized as a likely contributor to mating decisions (Mendelson and Shaw

2012), but it is unclear whether touch could represent a primary species recognition signal, given

that visual and auditory cues usually act earlier during the mating sequence. Research on the

prevalence of tactile signals in mating decisions is limited (Coleman 2008) because of the

experimental challenge it poses: whereas other sensory modalities present male signals to a focal

female from a distance, studying female preference for tactile cues requires contact between

males and females, which is not always easily achieved or quantified under controlled

conditions.

Despite this challenge, the role of tactile signals along the continuum between

intraspecific mate choice and interspecific RI deserves attention because it broadens our

understanding of the causes and consequences of a common pattern in nature— the rapid

divergence of male genital morphology between species. It has been suggested that rapid genital

differentiation can cause RI (Dufour 1844), although mechanical incompatibilities between

heterospecific male and female genitalia do not appear to be a common cause of RI (Shapiro and

Porter 1989; Masly 2012; Simmons 2014). However, observations both within (Eberhard 1994;

Edvardsson and Göran 2000; Briceño and Eberhard 2009a; Briceño and Eberhard 2009b; Frazee

and Masly 2015) and between species (Patterson and Thaeler Jr 1982; Robertson and Paterson

1982; Eberhard 1992; Coyne 1993) suggest that male genitalia may convey tactile information to

females that affects their subsequent behavior and/or physiology. Although female genital

structures often appear invariant among closely related species (Shapiro and Porter 1989), subtle

morphological differences (*e.g.*, Kamimura and Mitsumoto 2011; Yassin and Orgogozo 2013)

could enable females to detect variation among males' genital morphology. This variation could

occur in signal processing at the level of neurons and neural networks and/or in the distribution

and morphology of sensory structures that receive male tactile signals.

Female sensory structures that reside in body regions that contact species-specific male

structures during mating have been documented in several arthropods, including flies (Eberhard

2001; Ingram et al. 2008) and damselflies (Cordoba-Aguilar 2005; Robertson and Paterson

1982). Other studies have demonstrated that tactile cues from male grasping organs influence

female mating responses, either via experimental manipulation of male structures and/or

desensitization of females (Eberhard 2002; Briceño et al. 2007; Briceño and Eberhard 2009a;

Eberhard 2010; Myers et al. 2016) or comparison of female behavior when grasped by males

with varying genital morphologies (Sánchez-Guillén et al. 2012; Sánchez-Guillén et al. 2014;

Barnard et al. 2017). Premating tactile isolation may also be important in vision-limited

vertebrates. For example, contact cues via the lateral line system may influence female mate

choice in a cavefish (Plath et al. 2004; but see Rüschenbaum and Schlupp 2013).

Tactile signals appear to be a significant cause of RI in Zygoptera, the damselfly suborder

of Odonata (Krieger and Krieger-Loibl 1958; Loibl 1958; Robertson and Paterson 1982; Corbet

1999). Concentrations of cuticular mechanoreceptors (sensilla) on the female thorax have been

described in several coenagrionid damselfly genera. These sensilla reside in areas where male

grasping appendages contact the female thorax before and during mating, which has led to

speculation that they allow females to evaluate male morphologies and identify conspecifics

(Jurzitza 1974, 1975; Tennessen 1975; Robertson and Paterson 1982; Battin 1993a,b). Each

mechanoreceptor is associated with a single sensory neuron (McIver 1975; Kiel 1997). The

thoracic sensilla thus represent a spatial matrix that can transmit signals to the female central

nervous system based on the pattern in which the sensilla are stimulated. Greater numbers of

these receptors enhance a female's sensory resolution by increasing the combinatorial

complexity of tactile signals that a female can perceive. For example, if a female possesses 25

sensilla, and each sensillum has two response states ("on" if contacted and "off" if not

contacted), then the number of unique tactile patterns that the female could distinguish is $2^{25} =$

3.4×10^7. A female that possesses just one additional sensillum would be able to distinguish

among roughly twice as many patterns ($2^{26} = 6.7 \times 10^7$). Should individual sensilla respond to

quantitative variation in touch (rather than a binary response), this would dramatically increase

the number of response states and therefore further enhance tactile acuity (Gaffin and Brayfield

2017). Female damselfly thoracic sensilla thus present an external, quantifiable phenotype to

investigate the mechanistic basis of tactile stimuli and female mating decisions.

The North American damselfly genus *Enallagma* includes several recently diverged

species that often co-occur in the same habitats (Johnson and Crowley 1980; McPeek 1998), and

do not engage in premating courtship (Fincke et al. 2007; Barnard et al. 2017) or use chemical

cues for mate selection (Rebora et al. 2018). A female's first opportunity to assess a potential

mate occurs when the male uses his terminal appendages to grasp the mesostigmal plates on the

female's thorax to form tandem, the premating position. The males' superior appendages (cerci)

have species-specific morphologies, and differences in genital morphology are the primary cause

of RI in this genus (Paulson 1974; Barnard et al. 2017). Two species, *E. anna* and *E.*

carunculatum, occasionally hybridize in nature to produce males and females with morphologies

that are intermediate to each of the pure species (Donnelly 2008; Johnson 2009; Barnard et al.

2017). Females of both pure species discriminate strongly against both heterospecific and

interspecific hybrid males that take them in tandem, which shows that not only can *E. anna* and

E. carunculatum females detect large differences in male cercus morphologies, but also more

subtle differences such as those between conspecific and hybrid males (Barnard et al. 2017).

Because it appears that mesostigmal sensilla are used to mediate species recognition, they

might be expected to show signs of reproductive character displacement (RCD) in regions where

species co-occur (Brown and Wilson 1956; Howard 1993; Pfennig and Pfennig 2009). RCD can

evolve via direct selection on adult prezygotic phenotypes or via reinforcement, in which direct

selection against interspecific hybrids gives rise to selection for enhanced premating isolation

between species (Dobzhansky 1937). *Enallagma anna* and *E. carunculatum* can interbreed, but

these species produce hybrids with significantly reduced fitness (Barnard et al. 2017). This

suggests that species-specific sensilla phenotypes might show patterns consistent with RCD in

regions of sympatry, where females are known to experience frequent mating attempts from

heterospecific males (Paulson 1974; Fincke et al. 2007; Barnard et al. 2017). Here, we test the

hypothesis that variation in female sensilla phenotypes supports a function in species

recognition. We test this hypothesis by quantifying sensilla number, density, and location

phenotypes on the mesostigmal plates of a large set of *E. anna* and *E. carunculatum* females

from multiple populations across the western United States and comparing phenotypes of each

pure species from sympatric and allopatric populations to identify patterns consistent with RCD.

Methods

Population sampling

We measured the sensilla traits of 29 *E. anna* females across 13 populations, and 74 *E.*

carunculatum females across 19 populations (Figure 1, Table 1). We classified each population

as either allopatric, locally allopatric, or sympatric. Sympatric populations are those where *E.*

anna and *E. carunculatum* co-occur temporally as well as spatially. Because *E. anna*'s

geographic range falls completely within *E. carunculatum*'s range, only *E. carunculatum* has

completely allopatric populations. We designated populations that exist at sites within the area

of range overlap, but where only one species is known to occur, as locally allopatric. Some

specimens were collected as early as 1945, but the majority of samples (82 of 103) we studied

were collected between 2012 and 2016.

Figure 1. Sampling sites and species ranges. *Enallagma anna*'s geographic range (red) occurs

within *E. carunculatum*'s geographic range (orange). Names of sites associated with each

number are described in Table 1. Symbol color indicates the species sampled and symbol shape

indicates the population type. (Species ranges adapted from Johnson 2009; Paulson 2009, 2011).

Trait imaging and quantification

We photographed each damselfly using a Nikon D5100 camera (16.2 MP; Nikon

Corporation, Tokyo, Japan). We dissected the ventral thoracic cuticle from each female using

forceps and imaged the mesostigmal plates using scanning electron microscopy. Specimens

were mounted on aluminum stubs with carbon tape, sputter-coated with gold-palladium, and

imaged at 200X magnification and 3kV using a Zeiss NEON scanning electron microscope.

To avoid any potential bias during measurements, we blind-coded image files before

measuring traits. We measured abdomen length (abdominal segments 1-10, excluding terminal

appendages) on the full-body photos as a proxy for body size using the segmented line tool in

ImageJ (Abramoff et al. 2004). We quantified sensilla traits on the right mesostigmal plate of

each female damselfly unless the right plate was dirty or damaged, in which case we quantified

the left plate. Sensilla counts on a subset of 57 females showed that left plate and right plate

sensilla counts are highly correlated ($r = 0.85$). In cases where we quantified the left plate, we

flipped the image horizontally, so it was in the same orientation as a right plate. We

standardized the position of the mesostigmal plate in each image by cropping and rotating so that

the lower medial corner of the plate was in line with the lower left corner of each image. We

counted sensilla and obtained their x and y coordinates in ImageJ using the multi-point selection

tool. We traced an outline around the plate image, excluding the lateral carina (Figure S1), using

a Wacom Cintiq 12WX tablet and stylus (Wacom, Saitama, Japan) and the freehand selection

tool in ImageJ. This procedure provided x and y coordinates that describe the plate outline. We

performed all measurements twice for each specimen. Measurements across the two technical

replicates were highly correlated ($r_{abdomen} = 0.96$, n $= 155$; $r_{count} = 0.97$, n $= 183$; $r_{plate\ area} = 0.98$, n

$= 157$), so we used the mean trait values of the two replicates in subsequent analyses.

Sensilla trait analyses

We conducted all morphometric and statistical analyses using R v. 3.4.1 (R Core Team

2015). We used the plate outline coordinates to calculate each plate's two-dimensional area.

To calculate the area of the sensilla-covered region of each plate, we generated a polygon

connecting the coordinates of the outermost sensilla and calculated the area within this outline.

We determined the proportion of each plate that is covered by sensilla by dividing the sensilla

area by total plate area. We calculated sensilla density in two ways. First, we divided sensilla

number by the area of the sensilla-covered region. This measures the number of sensilla that

occur in a particular area but does not capture the relative arrangement of sensilla within that

area. Second, we computed the nearest neighbor distances among all sensilla within each plate

based on their *x* and *y* coordinates and then calculated the mean and median nearest neighbor

distances between the sensilla for each female. Nearest neighbor mean and median distances

were highly correlated ($r_{E.\ carunculatum}$ = 0.83; $r_{E.\ anna}$ = 0.88), so we report only the analyses using

the mean values.

To determine whether larger females possess more sensilla, we regressed sensilla number

against abdomen length. We found no significant relationship between these traits in either

species (*E. anna*: R^2_{adj} = -0.02, $F_{1,43}$ = 0.13, P = 0.73; *E. carunculatum*: R^2_{adj} = 0.005, $F_{1,65}$ =

1.35, P = 0.25). We thus present the results that compare sensilla counts without correcting for

differences in body size.

Sensilla spatial analyses

To quantify sensilla distributions within each plate, we generated kernel density estimates

(KDEs) for populations with at least four sampled individuals (two *E. anna* and six *E.*

carunculatum populations) using the R package ks (Duong 2016). First, we randomly selected

one of the two replicate sets of sensilla and plate outline coordinates for each female. To prepare

the coordinate data for KDE analyses, we concatenated the sensilla and plate coordinates for

each female and adjusted all plate outlines to have an area of one. This standardized each set of

sensilla coordinates for size while maintaining their relative positions within each plate. Next,

we translated each set of coordinates to place the origin of the coordinate system at the plate

outline's centroid. We concatenated sensilla coordinates for all females sampled within each

population to compute a representative KDE for each population.

Within each species, we conducted pairwise tests to compare each population's KDE

against every other population using the function kde.test() with the default settings. This test

returns a *P*-value that reflects the probability of generating the two sets of from the same

distribution of points. Because we performed multiple pairwise tests among *E. carunculatum*

populations, we adjusted the resulting *P*-values using the false discovery rate (Benjamini &

Hochberg 1995).

We generated an average plate outline for each population on which to visualize the

KDEs. The total number of coordinates that describe each plate outline varied among females,

ranging from 647-1078 for *E. anna* and 688-1028 for *E. carunculatum*. We standardized the

number of coordinates representing each plate by retaining the points for each of the upper and

lower medial corners and randomly sampled 198 points in between. We then treated each of

these 200 points as landmarks (the corners represented fixed landmarks and the remaining points

were designated as sliding semilandmarks) and used the R package geomorph (Adams and

Otarola-Castillo 2013) to perform general Procrustes analysis (Rohlf 1999) and obtain an

average two-dimensional plate shape for each population.

Statistical analyses

Some populations were well-sampled and other populations were represented by a single

female (Table 1). To avoid psedoreplication, for each population with N > 1, our analyses used

population means of trait values, so that each population was represented by a single

measurement. We arcsin transformed proportion data prior to analysis. We pooled data for

locally allopatric and fully allopatric *E. carunculatum* after *t*-tests showed that these groups did

not significantly differ with respect to sensilla number ($t_{2.7} = 0.80$, $P = 0.49$), sensilla density ($t_{9.2}$

= -1.62, P = 0.13), or the proportion of the plate that contained sensilla (t_{10} = 0.06, P = 0.95). To

compare traits between sympatric and allopatric populations of each species, we used t-tests or

Wilcoxon rank sum tests. We combined data for the two locally allopatric *E. carunculatum*

populations with the data from completely allopatric populations, after determining that these

data were similar enough to pool (sensilla number: $t_{2.1}$ = -0.91, P = 0.46; proportion plate with

sensilla: t_{11} = -1.24, P = 0.24; sensilla density: $t_{5.8}$ = 0.51, P = 0.63). To understand the

relationships between sensilla number, sensilla density, and the area of the plate occupied by

sensilla, we performed linear regressions between pairs of these traits.

Results

***E. anna* and *E. carunculatum* females possess distinct sensilla traits**

246   *Enallagma anna* females possessed significantly more sensilla per plate ($\bar{x}$ = 46.2 $\pm$ 1.4)

than *E. carunculatum* females ($\bar{x}$ = 28.7 $\pm$ 0.6; $t_{19.4}$ = 7.37, P = 4.9 $\times$ 10^{-7}; Figure 2A).

Enallagma anna females also had sensilla distributed over a larger proportion of each plate (W =

240, P = 2.6 $\times$ 10^{-7}; Figure 2B), and larger mean distances between sensilla (t_{30} = 5.2, P = 1.3 $\times$

10^{-5}; Figure 2C), which made *E. anna*'s overall sensilla distributions less dense than *E.*

carunculatum's (W = 239.5, P = 9.2.$\times$ 10^{-6}; Figure 2D). The sensilla occurred in different

locations on the mesostigmal plates of each species: they were more medial in *E. anna* and more

lateral in *E. carunculatum* (Figures 3, S2).

254
Figure 2. *E. anna* and *E. carunculatum* sensilla traits by population type. **(A)** The number of sensilla on one mesostigmal plate. **(B)** Proportion of the plate that contains sensilla. **(C)** Mean nearest neighbor distances between sensilla. **(D)** Sensilla density in the region of the plate that contains sensilla. Within each panel, each open circle represents the mean of one population. Boxplots show the interquartile range. The line within the box shows the median and whiskers extend to the most extreme observation within 1.5 times the interquartile range.

Both species showed a strong positive relationship between sensilla number and the

absolute area of the plate occupied by sensilla (*E. anna*: $R^2_{adj} = 0.33$, $F_{1,27} = 14.71$, $P < 0.0007$;

E. carunculatum: $R^2_{adj} = 0.33$, $F_{1,72} = 37.68$, $P = 4.1 \times 10^{-8}$). Consistent with this result, linear

regressions also revealed that females with more sensilla also had a larger proportion of the plate

occupied by sensilla (*E. anna*: $R^2_{adj} = 0.26$, $F_{1,27} = 10.65$, $P = 0.003$; *E. carunculatum*: $R^2_{adj} =$

0.20, $F_{1,65} = 18.93$, $P = 4.4 \times 10^{-5}$). Females with more sensilla had smaller mean distances

between neighboring sensilla (*E. anna*: $R^2_{adj} = 0.11$, $F_{1,27}$ 4.34, $P = 0.046$; *E. carunculatum*: R^2_{adj}

$= 0.09$, $F_{1,72} = 3.80$, $P = 0.01$). Overall, these results showed that a greater number of sensilla

was more strongly associated with a sensilla distribution that covers a larger area of the

mesostigmal plate rather a greater concentration sensilla within in a smaller area.

***E. carunculatum* sensilla traits do not show a strong pattern of reproductive character**

displacement

We made several non-mutually exclusive predictions expected under RCD for sensilla

traits in sympatric populations relative to allopatric populations. In particular, we predicted to

observe at least one of the following phenotypic differences in sympatric females relative to

allopatric females: (1) more numerous sensilla (2) denser sensilla, (3) sensilla concentrated in

different regions of the mesostigmal plates. We did not find significant differences in any of

these traits between sympatric and locally allopatric *E. anna* females (Table S1). However,

because our *E. anna* samples included only four females from three locally allopatric

populations, we could not perform a robust comparison of *E. anna* sensilla traits between

populations that do, or do not encounter *E. carunculatum*. We thus focus our analysis on

comparisons between sympatric and allopatric *E. carunculatum* populations, for which we had

larger sample sizes.

286   Sympatric *E. carunculatum* populations did not differ significantly from allopatric

287 populations in sensilla number ($t_{16.3} = 0.98$, $P = 0.35$), proportion of the mesostigmal plate

288 covered by sensilla ($t_{16.8} = 1.33$, $P = 0.20$), or sensilla density (overall density: $t_{9.7} = -0.26$, $P =$

289 0.80; mean distance between sensilla: $t_{18} = -1.31$, $P = 0.21$). In addition to divergence of mean

290 trait values, RCD can also result in reduced trait variance in sympatry without affecting the mean

291 (Pfennig and Pfennig 2009). Sympatric *E. carunculatum* populations displayed less variance in

292 both sensilla number (Figure 2A) and the proportion of the plate covered by sensilla (Figure 2B).

293 However, these trends were not statistically significant (sensilla number with locally allopatric

294 outlier removed: Bartlett's $K^2_1 = 0.75$, $P = 0.39$; proportion of plate covered by sensilla: Bartlett's

295 $K^2_1 = 2.5$, $P = 0.11$). KDE comparisons also did not reveal significant differences in sensilla

296 distributions between sympatric and allopatric *E. carunculatum* populations (Table S2).

297 However, the analysis revealed significant differences in sensilla distributions between several

298 pairs of populations that are not sympatric with *E. anna* (Figure 3E). This result is consistent

299 with those described above that indicated higher variance in sensilla traits among allopatric

300 populations compared to sympatric populations.

301
Figure 3. Sensilla locations. **(A)** White box indicates the location of right mesostigmal plate on
the thorax. **(B)** Ultrastructural details of individual sensilla. Scale bar represents 10 μm. **(C, D)**
Scanning electron micrographs show the locations of sensilla (yellow) on the mesostigmal plates
of *E. anna* (C) and *E. carunculatum* (D). Scale bars represent 100 μm. **(E, F)** Population kernel
density estimates for *E. carunculatum* (E) and *E. anna* (F) sensilla. The shading indicates
different regions of sensilla density: red represents the 75-99th percentile, orange represents the
50-74th percentile of sensilla density, and yellow represents the 25th-49th percentile. Each outline
represents the average mesostigmal plate shape for the population. Asterisks indicate *E.*
carunculatum populations whose KDEs were determined to be significantly different (* indicates
$P < 0.05$, *** $P < 0.001$).

Interestingly, although mean trait values did not differ significantly between sympatric

and allopatric populations, sensilla traits displayed considerable variation within the populations

we sampled. For example, within a single population, a particular female might have twice as

many sensilla than another female (Figure S2). This pattern was also observed in the *E. anna*

populations we studied.

Discussion

Enallagma anna and *E. carunculatum* females possessed different numbers of sensilla in

species-specific distributions on their mesostigmal plates. This result supports the idea that

receptors that receive male stimuli will occur in patterns that correspond to the male organs

during contact (Eberhard 2010). An association between male morphology and female sensilla

has been described for African *Enallagma* species (Robertson and Paterson 1982), and our

results show a similar pattern. *Enallagma anna* male cerci are considerably larger than *E.*

carunculatum cerci, and the observation that *E. anna* females had a larger number of sensilla

compared to *E. carunculatum* females is consistent with the difference in species-specific male

genital morphology.

When species make secondary contact after initial divergence in allopatry, the possible

outcomes are increased species divergence (*e.g.*, Sætre et al. 1997; Noor 2000; Naisbit et al.

2001; Yukilevich 2012; Dyer et al. 2014), decreased divergence (*e.g.*, Ritchie et al. 1989;

Shurtliff et al. 2013; Yang et al. 2016), one species goes locally extinct due to reproductive

exclusion (Hochkirch et al. 2007, Groning and Hochkirch 2008), or no change in either direction

(Abbott et al. 2013). Because *E. anna* and *E. carunculatum* produce reproductively

disadvantaged hybrids (Barnard et al. 2017), selection is expected to favor increased premating

isolation. Within each species, we predicted that female sensilla traits would show character

displacement in sympatric populations, which could indicate a shift in female preferences to

avoid mating with heterospecifics. Contrary to this prediction, *E. carunculatum* sympatric and

allopatric populations were not significantly different in mean sensilla trait values (Figure 2) or

sensilla density distributions (Figure 3E). Although we observed a trend toward more sensilla in

sympatric *E. anna* populations relative to allopatric populations (Figure 2A), it is difficult to

conduct a robust comparison for this species because (1) *E. anna*'s entire geographic range

overlaps with *E. carunculatum*'s range and (2) *E. anna* are often relatively rare (Acorn 2004; A.

Barnard, personal obs.). It was therefore difficult to collect sufficient *E. anna* samples from

populations that do not co-occur with *E. carunculatum*. We do, however, expect a stronger

pattern of RCD in sympatric *E. anna* females because *E. carunculatum* males can take them in

tandem relatively easily, whereas *E. anna* males are typically unsuccessful at taking *E.*

carunculatum females in tandem (Barnard et al. 2017). This means that *E. anna* females may

have more opportunities for mating mistakes than *E. carunculatum* females, which can result in

stronger asymmetric RCD (Lemmon 2009; Pfennig and Pfennig 2009).

There are at least four potential explanations for the absence of RCD in the form of

significant differences in the sensilla traits we measured between sympatric and allopatric

populations of *E. carunculatum*. First, species-specific sensilla distributions may be sufficiently

different to allow females to recognize when they are taken in tandem by heterospecific or

conspecific males. If this is true, small degrees of variation within the overall species pattern

among females might not affect females' species-recognition abilities. RCD is most easily

facilitated when the trait under selection already differs between species (Pfennig and Pfennig

2009). However, these traits may have already diverged enough sufficiently to preclude strong

selection for further divergence.

Second, it is possible that the external sensilla phenotypes we measured are not

representative of proximate female sensory traits, and the important variation lies deeper within

the female nervous system. For example, individual sensilla might differ in response rate or

ability to distinguish different levels of pressure applied by the cerci, and grasping pressure

might differ between males of each species. The direction of mechanosensor deflection is also

important for stimulus detection (Keil 1997), and different species' cercus morphologies may

contact sensilla from different angles. Female mate preferences may also be influenced by male

exposure and sexual experience (Svensson et al. 2014).

Third, the thoracic sensilla may not be a target of strong selection. For example, earlier

acting forms of RI may prevent most heterospecific interactions in the sympatric populations we

sampled. In one region where *E. anna* and *E. carunculatum* co-occur, habitat and temporal

isolation were close to zero (Barnard et al. 2017), but the strength of these isolating barriers may

vary geographically.

Finally, although we did not detect a statistically significant difference between group

means, the small differences we observed may still have biological relevance. If gaining just one

additional mechanosensor can (at least) double a female's tactile discriminatory power (Gaffin

and Brayfield 2017), then females in a population with a seemingly minor upward shift in

sensilla number could gain a remarkable increase in their ability to detect and avoid mating with

heterospecifics. Similarly, it is difficult to determine the features of sensilla density distributions

that may influence female preference solely by conducting statistical tests between KDEs. The

human eye can visually detect differences in the KDE plots shown in Figure 3, and it is thus

possible that these spatial differences reflect salient variation in the way females might receive

tactile stimuli.

These possible explanations highlight the interesting avenues that female damselfly

sensilla provide for investigating how females evaluate male tactile signals to make mating

decisions. The ability to quantify the number and locations of female mechanoreceptors in a

region contacted by male reproductive structures complements our understanding of patterns of

variation in male morphologies (McPeek et al. 2008; McPeek et al. 2009; McPeek et al. 2011;

Barnard et al. 2017). Females of both species display substantial intrapopulation variation in

sensilla traits (Figure S2) and this variation may play a role in sexual selection and female

preferences within species. Behavioral studies will be crucial to link mechanoreceptor

phenotypes to female mating decisions and clarify how sensilla traits influence both species

recognition and sexual selection. For example, do females with more sensilla make fewer

mating mistakes than females with fewer sensilla (Lemmon 2009)? Another outstanding

question of this system is how the cerci stimulate individual sensilla during tandem. This might

be determined by flash-freezing male-female tandem pairs and using micro-CT scanning to

understand how the male and female structures interact, similar to a recent approach used in seed

beetles (Dougherty and Simmons 2017). Once we understand how cerci contact the sensilla,

functional tests of sensilla electrophysiology could reveal how individual sensilla respond to

stimulation and indicate whether certain sensilla make greater contributions to reproductive

decision-making than others.

Female preference can drive sexual selection, promote trait divergence, and cause RI

between species (Ritchie 1996). A longstanding presumption in the literature on genital

evolution and speciation has been that female reproductive morphologies are less variant or

species-specific than male genitalia (Shapiro and Porter 1989). However, recent studies of

variation in female reproductive structures suggest that variation does exist among individuals

and species (Ah-King et al. 2014), and our data support the need to look beyond the visible

external morphologies. When male genital morphologies are obviously divergent, but female

morphologies are not, females may possess important variation at neurophysiological levels that

affects how they evaluate male tactile signals, similar to the way females evaluate signals in

other sensory modalities.

References cited

Abramoff, M. D., P. J. Magalhaes, and S. J. Ram. 2004. Image processing with ImageJ.
413   Biophotonics International 11:36-42.
414   Acorn, J. 2004. Damselflies of Alberta: flying neon toothpicks in the grass. The University of
415   Alberta Press, Edmonton, Alberta, Canada.
416   Adams, D. C. and E. Otarola-Castillo. 2013. geomorph: an R package for the collection and
417   analysis of geometric morphometric shape data. Methods in Ecology and Evolution
418   4:393-399.
419   Ah-King, M., A. B. Barron, and M. E. Herberstein. 2014. Genital evolution: why are females
420   still understudied? PLoS Biology 12:e1001851.
421   Arthur, B. J., T. Sunayama-Morita, P. Coen, M. Murthy, and D. L. Stern. 2013. Multi-channel
422   acoustic recording and automated analysis of *Drosophila* courtship songs. BMC Biology
423   11:1-11.
424   Barnard, A. A., O. M. Fincke, M. A. McPeek, and J. P. Masly. 2017. Mechanical and tactile
425   incompatibilities cause reproductive isolation between two young damselfly species.
426   Evolution 71:2410–2427.
427   Battin, T. J. 1993a. The odonate mating system, communication, and sexual selection: A review.
428   Bulletin of Zoology 60:353-360.
429   Battin, T. J. 1993b. Revision of the puella group of the genus Cuenagrion Kirby, 1890 (Odonata,
430   Zygoptera), with emphasis on morphologies contributing to reproductive isolation.
431   Hydrobiologia 262:13-29.
432   Boughman, J. W., H. D. Rundle, and D. Schluter. 2005. Parallel evolution of sexual isolation in
433   sticklebacks. Evolution 59:361-373.
434   Briceño, R., W. Eberhard, and A. Robinson. 2007. Copulation behaviour of *Glossina pallidipes*
435   (Diptera: Muscidae) outside and inside the female, with a discussion of genitalic
436   evolution. Bulletin of entomological research 97:471-488.
437   Briceño, R. D. and W. G. Eberhard. 2009a. Experimental demonstration of possible cryptic
438   female choice on male tsetse fly genitalia. Journal of Insect Physiology 55:989-996.

Briceño, R. D. and W. G. Eberhard. 2009b. Experimental modifications imply a stimulatory
function for male tsetse fly genitalia, supporting cryptic female choice theory. Journal of
evolutionary biology 22:1516-1525.
Brown, W. L. and E. O. Wilson. 1956. Character displacement. Systematic zoology 5:49-64.
Castillo, D. M., M. K. Burger, C. M. Lively, and L. F. Delph. 2015. Experimental evolution:
Assortative mating and sexual selection, independent of local adaptation, lead to
reproductive isolation in the nematode *Caenorhabditis remanei*. Evolution 69:3141-3155.
Coleman, S. W. 2008. Taxonomic and sensory biases in the mate-choice literature: there are far
too few studies of chemical and multimodal communication. Acta Ethologica 12:45-48.
Corbet, P. S. 1999. Dragonflies: Behaviour and ecology of Odonata. Cornell University Press,
Ithaca, NY.
Cordoba-Aguilar, A. 2005. Possible coevolution of male and female genital form and function in
a calopterygid damselfly. J Evol Biol 18:132-137.
Costanzo, K. and A. Monteiro. 2007. The use of chemical and visual cues in female choice in the
butterfly *Bicyclus anynana*. Proceedings of the Royal Society of London B: Biological
Sciences 274:845-851.
Coyne, J. A. 1993. The genetics of an isolating mechanism between two sibling species of
Drosophila. Evolution 47:778-788.
Coyne, J. A., A. Crittenden, and K. Mahi. 1994. Genetics of a pheromonal difference
contributing to reproductive isolation in Drosophila. Science 265:1461–1464.
Dobzhansky, T. 1937. Genetics and the Origin of Species. Columbia University press.
Donnelly, N. T. 2008. A Hybrid Complex in *Enallagma*. Argia 20:10-11.
Dopman, E. B., P. S. Robbins, and A. Seaman. 2010. Components of reproductive isolation
between North American pheromone strains of the European corn borer. Evolution
64:881-902.
Dougherty, L. R. and L. W. Simmons. 2017. X-ray micro-CT scanning reveals temporal
separation of male harm and female kicking during traumatic mating in seed beetles.
Proceedings of the Royal Society of London B: Biological Sciences 284: 20170550.
Dufour, L. 1844. Anatomie générale des diptéres. Annales des Sciences Naturelles 1:244-264.
Duong, T. 2016. ks: Kernel Smoothing. R package version 1.10.4.
Dyer, K. A., B. E. White, J. L. Sztepanacz, E. R. Bewick, and H. D. Rundle. 2014. Reproductive
character displacement of epicuticular compounds and their contribution to mate choice
in Drosophila subquinaria and Drosophila recens. Evolution 68:1163-1175.
Eberhard, W. G. 1992. Species isolation, genital mechanics, and the evolution of species-specific
genitalia in three species of *Macrodactylus* Beetles (Coleoptera, Scarabeidae,
Melolonthinae). Evolution 46:1774-1783.
Eberhard, W. G. 1994. Evidence for widespread courtship during copulation in 131 species of
insects and spiders, and implications for cryptic female choice. Evolution 48:711-733.
Eberhard, W. G. 2001. The functional morphology of species-specific clasping structures on the
front legs of male sepsid flies. Zoological Journal of the Linnean Society 133:335–368.
Eberhard, W. G. 2002. Physical restraint or stimulation? The function(s) of the modified front
legs of male *Archisepsis diversiformis* (Diptera, Sepsidae). Journal of Insect Behavior
15:831-850.
Eberhard, W. G. 2010. Evolution of genitalia: theories, evidence, and new directions. Genetica
138:5-18.

Edvardsson, M. and A. Göran. 2000. Copulatory courtship and cryptic female choice in red flour
beetles Tribolium castaneum. Proceedings of the Royal Society of London B: Biological
Sciences 267:559-563.
Ewing, A. W. and H. Bennet-Clark. 1968. The courtship songs of *Drosophila*. Behaviour
31:288-301.
Fincke, O. M., A. Fargevieille, and T. D. Schultz. 2007. Lack of innate preference for morph and
species identity in mate-searching *Enallagma* damselflies. Behavioral Ecology and
Sociobiology 61:1121-1131.
Frazee, S. R. and J. P. Masly. 2015. Multiple sexual selection pressures drive the rapid evolution
of complex morphology in a male secondary genital structure. Ecology and Evolution
5:4437-4450.
Gaffin, D. and B. Brayfield. 2017. Exploring the chemo-textural familiarity hypothesis for
scorpion navigation. Journal of Arachnology 45:265-270.
Gerhardt, H. C. and F. Huber. 2002. Acoustic Communication in Insects and Anurans: Common
Problems and Diverse Solutions. University of Chicago Press, Chicago, IL.
Girard, M. B., D. O. Elias, and M. M. Kasumovic. 2015. Female preference for multi-modal
courtship: multiple signals are important for male mating success in peacock spiders.
Proceedings of the Royal Society of London B: Biological Sciences 282.
Howard, D. J. 1993. Reinforcement: origin, dynamics, and fate of an evolutionary hypothesis.
Pp. 46–69 *in* R. G. Harrison, ed. Hybrid zones and the evolutionary process. Oxford
University Press, New York.
Ingram, K. K., T. Laamanen, N. Puniamoorthy, and R. Meier. 2008. Lack of morphological
coevolution between male forelegs and female wings in *Themira* (Sepsidae: Diptera:
Insecta). Biological Journal of the Linnean Society 93:227-238.
Jiggins, C. D., R. E. Naisbit, R. L. Coe, and J. Mallet. 2001. Reproductive isolation caused by
colour pattern mimicry. Nature 411:302–305.
Johnson, D. M. and P. H. Crowley. 1980. Habitat and seasonal segregation among coexisting
odonate larvae. Odonatologica 9:297–308.
Johnson, J. 2009. Presumed *Enallagma anna* Williamson × *carunculatum* Morse Hybrids from
Oregon and California. Bulletin of American Odonatology 11:8-10.
Jurzitza, G. 1974. Rasterelektronenmikroskopische Untersuchungen des Zangen- griffes und der
Laminae mesostigmales einiger Coenagrionidae (Odonata, Zygoptera). Forma Functio
7:377-392.
Jurzitza, G. 1975. Rasterelektronenmikroskopische Untersuchungen an den Appendices und den
Laminae mesotigmales einiger *Enallagma*-Arten (Odonata. Zygoptera). Forma Functio
8:33-48.
Kamimura, Y. and H. Mitsumoto. 2011. Comparative copulation anatomy of the *Drosophila*
melanogaster species complex (Diptera: Drosophilidae). Entomological science 14:399-
410.
Keil, T. A. 1997. Functional morphology of insect mechanoreceptors. Microscopy Research and
Technique 39:506–531.
Krieger, F. and E. Krieger-Loibl. 1958. Beiträge zum Verhalten von *Ischnura elegans* und
Ischnura pumilio (Odonata) 1. Zeitschrift für Tierpsychologie 15:82-93.
Kronforst, M. R., L. G. Young, D. D. Kapan, C. McNeely, R. J. O'Neill, and L. E. Gilbert. 2006.
Linkage of butterfly mate preference and wing color preference cue at the genomic
location of wingless. Proc Natl Acad Sci U S A 103:6575-6580.

Lemmon, E. M. 2009. Diversification of conspecific signals in sympatry: geographic overlap drives multidimensional reproductive character displacement in frogs. Evolution 53:1155-1170.

Loibl, E. 1958. Zur Ethologie und Biologie der deutschen Lestiden (Odonata) 1. Zeitschrift für Tierpsychologie 15:54-81.

Masly, J. P. 2012. 170 Years of "Lock-and-Key": Genital Morphology and Reproductive Isolation. International Journal of Evolutionary Biolog 2012:247352.

Mayr, E. 1942. Systematics and the origin of species, from the viewpoint of a zoologist. Harvard University Press, Cambridge, MA.

McIver, S. B. 1975. Structure of cuticular mechanoreceptors of arthropods. Annual Review of Entomology 20:381-397.

McMillan, W. O., C. Jiggins, and J. Mallet. 1997. What initiates speciation in passion-vine butterflies? Proceedings of the National Academy of Sciences USA 94:8628–8633.

McPeek, M. A. 1998. The consequences of changing the top predator in a food web: a comparative experimental approach. Ecological Monographs 68:1-23.

McPeek, M. A., L. Shen, and H. Farid. 2009. The correlated evolution of three-dimensional reproductive structures between male and female damselflies. Evolution 63:73-83.

McPeek, M. A., L. Shen, J. Z. Torrey, and H. Farid. 2008. The tempo and mode of three-dimensional morphological evolution in male reproductive structures. Am Nat 171:E158-178.

McPeek, M. A., L. B. Symes, D. M. Zong, and C. L. McPeek. 2011. Species recognition and patterns of population variation in the reproductive structures of a damselfly genus. Evolution 65:419-428.

Mendelson, T. C. and K. L. Shaw. 2012. The (mis) concept of species recognition. Trends in ecology & evolution 27:421-427.

Mendelson, T. C. and G. Wallis. 2003. Sexual isolation evolves faster than hybrid inviability in a diverse and sexually dimorphic genus of fish (Percidae: *Etheostoma*). Evolution 57:317-327.

Myers, S. S., T. R. Buckley, and G. I. Holwell. 2016. Male genital claspers influence female mate acceptance in the stick in sect *Clitarchus hookeri*. Behavioral Ecology and Sociobiology 70:1547-1556.

Naisbit, R. E., C. D. Jiggins, and J. Mallet. 2001. Disruptive sexual selection against hybrids contributes to speciation between *Heliconius cydno* and *Heliconius melpomene*. Proceedings of the Royal Society of London B: Biological Sciences 268:1849-1854.

Noor, M. and J. A. Coyne. 1996. Genetics of a difference in cuticular hydrocarbons between *Drosophila pseudoobscura* and *D. persimilis*. Genetics Research 68:117-123

Noor, M. F. 2000. On the evolution of female mating preferences as pleiotropic byproducts of adaptive evolution. Adaptive Behavior 8:3-12.

Patterson, B. D. and C. S. Thaeler Jr. 1982. The mammalian baculum: hypotheses on the nature of bacular variability. Journal of Mammalogy 63:1-15.

Paulson, D. R. 2009. Dragonflies and Damselflies of the West. Princeton University Press, Princeton, New Jersey.

Paulson, D. R. 1974. Reproductive isolation in damselflies. Systematic Zoology 23:40-49.

Paulson, D. R. 2011. Dragonflies and Damselflies of the East. Princeton University Press, Princeton, New Jersey.

Pfennig, K. S. and D. W. Pfennig. 2009. Character displacement: ecological and reproductive responses to a common evolutionary problem. The Quarterly Review of Biology 84:253-276.

Plath, M., J. Parzefall, K. E. Körner, and I. Schlupp. 2004. Sexual selection in darkness? Female mating preferences in surface-and cave-dwelling Atlantic mollies, *Poecilia mexicana* (Poeciliidae, Teleostei). Behavioral Ecology and Sociobiology 55:596-601.

Price, T. D. and M. M. Bouvier. 2002. The evolution of F1 postzygotic incompatibilities in birds. Evolution 56:2083-2089.

Rafferty, N. E. and J. W. Boughman. 2006. Olfactory mate recognition in a sympatric species pair of three-spined sticklebacks. Behavioral Ecology 17:965-970.

Rebora, M., F. Frati, S. Piersanti, G. Salerno, R. Selvaggini, and O. M. Fincke. 2018. Field tests of multiple sensory cues in sex recognition and harassment of a colour polymorphic damselfly. Animal Behaviour 136:127-136.

Ritchie, M. G. 1996. The shape of female mating preferences. Proceedings of the National Academy of Sciences USA 93:14628-14631.

Ritchie, M. G., R. K. Butlin, and G. M. Hewitt. 1989. Assortative mating across a hybrid zone in *Chorthippus parallelus* (Orthoptera: Acrididae). Journal of Evolutionary Biology 2:339–352.

Robertson, H. M. and H. E. H. Paterson. 1982. Mate Recognition and Mechanical Isolation in *Enallagma* Damselflies (Odonata: Coenagrionidae). Evolution 36:243-250.

Rohlf, F. J. 1999. Shape statistics: Procrustes superimpositions and tangent spaces. Journal of Classification 16:197-223.

Rüschenbaum, S. and I. Schlupp. 2013. Non-visual mate choice ability in a cavefish (Poecilia mexicana) is not mechanosensory. Ethology 119:368-376.

Sætre, G.-P., T. Moum, S. Bureš, M. Král, M. Adamjan, and J. Moreno. 1997. A sexually selected character displacement in flycatchers reinforces premating isolation. Nature 387:589-592.

Sánchez-Guillén, R. A., A. Cordoba-Aguilar, A. Cordero Rivera, and M. Wellenreuther. 2014. Rapid evolution of prezygotic barriers in non-territorial damselflies. Biological Journal of the Linnean Society 113:485-496.

Sánchez-Guillén, R. A., M. Wellenreuther, and A. Cordero Rivera. 2012. Strong asymmetry in the relative strengths of prezygotic and postzygotic barriers between two damselfly sister species. Evolution 66:690-707.

Shapiro, A. M. and A. H. Porter. 1989. The lock-and-key hypothesis: Evolutionary and biosystematic interpretation of insect genitalia. Annual Review of Entomology 34:321-345.

Shaw, K. L. 2000. Interspecific genetics of mate recognition: inheritance of female acoustic preference in Hawaiian crickets. Evolution 54:1303–1312.

Shurtliff, Q. R., P. J. Murphy, J. D. Yeiter, and M. D. Matocq. 2013. Experimental evidence for asymmetric mate preference and aggression: behavioral interactions in a woodrat (*Neotoma*) hybrid zone. BMC Evolutionary Biology 13:220-233.

Simmons, L. W. 2014. Sexual selection and genital evolution. Austral Entomology 53:1-17.

Stratton, G. E. and G. W. Uetz. 1986. The Inheritance of Courtship Behavior and Its Role as a Reproductive Isolating Mechanism in Two Species of *Schizocosa* Wolf Spiders (Araneae; Lycosidae). Evolution 40:129-141.

Svensson, E. I., A. Runemark, M. N. Verzijden, and M. Wellenreuther. 2014. Sex differences in
developmental plasticity and canalization shape population divergence in mate
preferences. Proceedings of the Royal Society of London B: Biological Sciences 281:1-8.
Team, R. C. 2015. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
Tennessen, K. J. 1975. Reproductive behavior and isolation of two sympatric Coenagrionid
damselflies in Florida. University of Florida, Gainesville, FL.
Trabalon, M., A. G. Bagnères, and C. Roland. 1997. Contact sex signals in two sympatric spider
species, *Tegenaria domestica* and *Tegenaria pagana*. Journal of Chemical Ecology
23:747–758.
Wells, M. M. and C. S. Henry. 1998. Songs, reproductive isolation, and speciation in cryptic
species of insects: A case study using green lacewings. Pp. 217-233. Endless forms:
species and speciation. Oxford University Press, New York, NY.
Wiernasz, D. C. and J. G. Kingsolver. 1992. Wing melanin pattern mediates species recognition
in *Pieris occidentalis*. Animal Behaviour 43:89–94.
Williams, T. H. and T. C. Mendelson. 2014. Quantifying reproductive barriers in a sympatric
pair of darter species. Evolutionary Biology 41:212-220.
Yang, Y., C. L. Richards-Zawacki, A. Devar, and M. B. Dugas. 2016. Poison frog color morphs
express assortative mate preferences in allopatry but not sympatry. Evolution 70:2778–
2788.
Yassin, A. and V. Orgogozo. 2013. Coevolution between male and female genitalia in the
Drosophila melanogaster species subgroup. PLoS One 8:e57158.
Yukilevich, R. 2012. Asymmetrical patterns of speciation uniquely support reinforcement in
Drosophila. Evolution 66:1430-1446.

Table 1. Sampling sites for *E. anna* and *E. carunculatum* populations. N refers to the number of females that were imaged and measured for this study. Sources: A. Barnard(AB), Ola Fincke (OF), Bill Mauffray (BM), and Dennis Paulson (DP).

Type	Site (site number)	Species	Year collected	N	Source
Sympatric	Big Spring, UT (1)	*E. anna*	2016	10	AB
		E. carunculatum		4	
	Big Sandy Creek, MT (2)	*E. carunculatum*	2015	1	AB
	Creston, MT (3)	*E. anna*	1972	1	BM
	Dry Sheep Creek, NE (4)	*E. anna*	2012	1	BM
	Fish Springs Run, CA (5)	*E. anna*	1998	2	BM
	Grace Coolidge Creek, SD (6)	*E. anna*	1969	1	BM
	Horseshoe Springs, UT (7)	*E. anna*	2016	1	AB
		E. carunculatum		1	
	Long Valley Creek, CA (8)	*E. anna*	1973	5	DP
	Murray Creek, NV (9)	*E. anna*	2001	1	
	Malad River, UT (10)	*E. carunculatum*	1983	2	BM
	Niwot Ditch, CO (11)	*E. anna*	2015	2	AB
		E. carunculatum		1	
	Pondera Coulee, MT (12)	*E. anna*	2015	1	AB
		E. carunculatum		1	
Locally allopatric	Beaver Creek, WY (13)	*E. anna*	2015	1	AB
	Indian Road Camp, MT (14)	*E. carunculatum*	2015	4	AB
	Jackson, WY (15)	*E. anna*	1971	2	BM
	Muddy Creek, MT (16)	*E. anna*	2015	1	AB
	Strawberry River, UT (17)	*E. carunculatum*	2016	1	AB
	West Greenbelt, CO (18)	*E. carunculatum*	2014	9	AB
Allopatric	Bull Lake, MT (19)	*E. carunculatum*	2015	1	AB
	Crab Creek, WA (20)	*E. carunculatum*	2016	20	DP
	Clear Lake, IN (21)	*E. carunculatum*	1945	1	BM
	Columbia River, WA (22)	*E. carunculatum*	1952	2	BM
	Douglas Lake, MI (23)	*E. carunculatum*	2016	17	OF
	Flathead River, MT (24)	*E. carunculatum*	2015	4	AB
	Home Lake, CO (25)	*E. carunculatum*	2015	1	AB
	Little Lake, CA (26)	*E. carunculatum*	1967	1	DP
	Drumond Island, MI (27)	*E. carunculatum*	2002	1	BM
	Snake River, ID (28)	*E. carunculatum*	1983	2	BM