Title:

Integrative cross-species analyses identify deficits in habituation learning as a widely affected mechanism in Autism

Short title:

Habituation deficits in Autism models

Authors:

Michaela Fenckova[1], Lenke Asztalos[2,3], Pavel Cizek[4], Euginia L. Singgih[1,5], Laura E.R. Blok [1], Jeffrey C. Glennon[5], Joanna IntHout[6], Christiane Zweier[7], Evan E. Eichler[8,9], Raphael A. Bernier[10], Zoltan Asztalos[2,3,11], Annette Schenck[1,*]

Affiliations:

[1]Department of Human Genetics, Donders Institute for Brain, Cognition and Behaviour, Radboud university medical center, 6525 GA, Nijmegen, the Netherlands

[2]Aktogen Ltd., Department of Genetics, University of Cambridge, CB2 3EH, Cambridge, United Kingdom

[3]Aktogen Hungary Ltd., Bay Zoltán Nonprofit Ltd. for Applied Research, Institute for Biotechnology (BAY-BIO), H-6726, Szeged, Hungary

[4]Centre for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboud university medical center, Nijmegen, 6525 GA, the Netherlands

[5]Department of Cognitive Neuroscience, Donders Institute for Brain, Cognition and Behaviour, Radboud university medical center, 6525 EN, Nijmegen, the Netherlands

[6]Department for Health Evidence, Radboud university medical center, 6525 EZ, Nijmegen, The Netherlands

[7]Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg, 91054, Erlangen, Germany

[8]Department of Genome Sciences, University of Washington School of Medicine, Seattle, WA 98195,

USA

[9]Howard Hughes Medical Institute, University of Washington, Seattle, WA 98195, USA

[10]Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, WA 98105,

USA

[11]Institute of Biochemistry, Biological Research Centre, Hungarian Academy of Sciences, H-6726,

Szeged, Hungary

***Corresponding author:**

Annette Schenck

Radboudumc

Dept. of Human Genetics

Geert Grooteplein 10

6525 GA, Nijmegen

The Netherlands

Annette.Schenck@radboudumc.nl

Tel: +31243610868

<u>Keywords:</u>

Habituation learning, Intellectual Disability, Autism Spectrum Disorder, *Drosophila*, Ras-MAPK,

GABAergic neurons

Number of words in abstract: 245

Number of words in the article body: 3995

Number of figures: 7

Number of tables: 0

Number of supplemental files: 2

Abstract:

Background: Although habituation is one of the most ancient and fundamental forms of learning, its regulators and relevance for human disease are poorly understood.

Methods: We manipulated the orthologs of 286 genes implicated in intellectual disability (ID) with or without comorbid autism spectrum disorder (ASD) specifically in *Drosophila* neurons, and tested these models in light-off jump habituation. We dissected neuronal substrates underlying the identified habituation deficits and integrated genotype-phenotype annotations, gene ontologies and interaction networks to determine the clinical features and molecular processes that are associated with habituation deficits.

Results: We identified more than 100 genes required for habituation learning. For the vast majority of these, 93 genes, a role in habituation learning was previously unknown. These genes characterize ID disorders with overgrowth/macrocephaly and comorbid ASD. Moreover, ASD individuals from the Simons Simplex Collection carrying disruptive *de novo* mutations in these genes exhibit increased rates of specific aberrant behaviors including stereotypic speech, hyperactivity and irritability. At the molecular level, ID genes required for normal habituation are enriched in synaptic function and converge on Ras-MAPK signaling. Both increased Ras-MAPK signaling in GABAergic and decreased Ras-MAPK signaling in cholinergic neurons specifically inhibit the adaptive habituation response.

Conclusions: Our work demonstrates the relevance of habituation learning to autism, identifies an unprecedented number of novel habituation players, supports an emerging role for inhibitory neurons in habituation and reveals an opposing, circuit-level-based mechanism for Ras-MAPK signaling. This establishes habituation as a possible, widely applicable target for pharmacologic intervention in ID/ASD.

Introduction:

Habituation is one of the most ancient and fundamental forms of learning, conserved across the animal kingdom (1). It causes an organism's initial response to repeated not meaningful stimuli to gradually decline. Learning to ignore irrelevant stimuli as a result of habituation is thought to represent a filter mechanism that prevents information overload, allow for selective attention, and thus to focus cognitive resources on relevant matters. Habituation learning has been proposed to represent an important element and prerequisite for higher cognitive functions (2–4). In line with this, habituation in infants correlates better than other measures with later cognitive abilities (5). Habituation has also emerged as a good proxy for synaptic plasticity (6). However, key players and molecular mechanisms underlying habituation are poorly understood (7).

In humans, deficits in habituation have been reported in a number of neuropsychiatric and behavioral disorders. In particular, defective cortical filtering of sensory stimuli and information overload, as expected to arise from habituation deficits, are thought to represent a critical mechanism underlying autism spectrum disorder (ASD) (8, 9). Indeed, a decreased ability to habituate has been described in a fraction of ASD individuals (10–12), but has not been connected yet to specific genetic defects, with a single exception. Recently, two independent studies demonstrated habituation deficits in patients with Fragile X syndrome, the most common monogenic cause of intellectual disability (ID) and ASD (13, 14), confirming previously reported habituation deficits in Fmr1 KO mice (15, 16). Habituation deficits have also been reported in a limited number of other ID/ASD disease models (17–20).

We apply light-off jump habituation in *Drosophila* to increase our knowledge on the genetic control of habituation and, at the same time, to assess the relevance of decreased habituation in ID and in comorbid ASD disorders. *Drosophila* is a well-established model for ID (21–23) and offers genome-wide resources to study gene function in large scale (24, 25). Several forms of habituation have been established in *Drosophila* (26–30). Deficits in light-off jump habituation have already been reported in several ID models (22, 31–35) and in classical learning and memory mutants (27, 30).

Moreover, this form of habituation can be assessed in a high-throughput manner. In the light-off

jump paradigm, the initial jump response to repeated light-off stimuli gradually wanes, as has been

demonstrated not due to sensory adaptation (a decrease in detecting the stimulus) or motor fatigue

(a decrease in the ability to execute the response) but as a result of learned adaptation of the startle

circuit (30). This behavior meets all habituation criteria (3), including spontaneous recovery and

dishabituation with a novel stimulus (30, 36). The rich genetic toolbox of *Drosophila* allows for

dissection of neuronal substrates and mechanisms of light-off jump habituation.

Here, we use inducible RNA interference (RNAi) in *Drosophila* to systematically assess the

role of 286 genes that are well-established to cause ID in humans when mutated (hereinafter

referred to as ID genes). 69 of them (24%) have also been implicated in ASD (37, 38) (hereinafter

referred to as ID plus ASD-associated genes). We report more than a hundred novel genes required

for habituation learning. This substantially exceeds the number of previously known regulators of

habituation across paradigms and species, and identifies habituation learning as a widely affected

mechanism in ID/ASD disorders. We further systematically reveal the relation of these genes to their

associated clinical phenotypes and molecular functions, and determine neuronal substrates of Ras-

MAPK signaling, which we identify as a central key pathway controlling habituation.

Methods and Materials

Investigated ID genes

A systematic source of ID genes and their *Drosophila* orthologs is available online (SysID database,

sysid.cmbi.umcn.nl (39)). We investigated the *Drosophila* orthologs of 286 human ID genes from the

SysID category primary ID genes (**Table S1**) (containing mutations with robust published evidence for

causality, see **Supplemental Methods (SM)**). SysID inclusion criteria and in/exclusion criteria of

experimentally investigated genes are indicated in the **SM**. In brief, the vast majority of genes are

from the first data freeze of the SysID database (all ID genes with proven causality, status of mid

2010). Genes have been included based on conservation in *Drosophila*, available tools (RNAi) from

large-scale resources and viability as a prerequisite for behavioral testing. No selection was

performed.

Information about the identification of *Drosophila* orthologs, proposed disease mechanism, light-off

habituation assay and analysis, phenotype reproducibility, *Drosophila* stocks, fatigue assay, quality

criteria for RNAi lines, annotation of ID plus ASD associated genes, comparison of behavior and

cognition in ASD SSC, molecular interaction network, clustering, physical interaction enrichment

(PIE), enrichment analysis, data visualization and statistics are described in the **SM**.

Results:

Systematic identification of habituation deficits in *Drosophila* models of ID

To identify novel genes implicated in habituation, we systematically investigated the role of 278 *Drosophila* orthologs representing 286 human ID genes in the light-off jump habituation paradigm. We induced neuron-specific knockdowns of each ID gene ortholog by RNAi (24) using 513 RNAi lines fulfilling previously established quality criteria (39, 40), with two independent constructs per gene whenever available. These were crossed to the panneuronal elav-Gal4 driver line (see **SM**). Knockdown is a suitable approach for modeling of the here-investigated human disease conditions since (partial) loss of function is considered to be the underlying mechanism in the vast majority of these disorders (40) (**Table S1**). Restricting gene knockdown to neurons eliminates potential effects on viability or organismal fitness originating from an essential role of genes in other tissues and establishes neuron-autonomous mechanisms.

Knockdown and control flies of identical genetic background were subjected to a series of 100 light-off stimuli, hereinafter referred to as trials, in the light-off jump habituation paradigm. The screening procedure and paradigm allowed us to distinguish the following parameters: viability, initial jump response (percentage of flies that jumped in at least one of the first five trials), and premature and reduced habituation, with the latter representing the learning-defective phenotype category of main interest. Genotypes with a good initial jump response (≥50% initial jumpers) but premature habituation were subjected to a secondary assay to evaluate their organismal fitness and exclude fatigue as a confounder of premature habituation (see **SM** and **Figure S1**). Based on these parameters, genes were assigned to at least one of four phenotype categories (**Figure A1**): (1) "not affected": (both) tested RNAi lines targeting such genes were viable, showed good initial jump response, and had no significant effect on habituation (based on the FDR-corrected p-value (p_{adj}), see **SM**); (2) "reduced organismal fitness": at least one RNAi line led to lethality, poor jump response (<50% initial jumpers), or premature habituation because of increased fatigue; (3) "habituation

deficient": at least one RNAi line showed good initial jump response but failed to suppress their

response with the increasing number of light-off trials (based on p_{adj}); and (4) "premature

habituation": at least one RNAi line showed good initial jump response followed by faster decline

(based on p_{adj}), without fatigue being detectable in the secondary assay. Still, this later phenotype

category can result from other defects than improved habituation, and will be further investigated

elsewhere. In this study we focus on habituation deficits (3), corresponding to the phenotype that

has been shown in ID and ASD (10–14).

We validated the experimental approach to identify genes which, if manipulated, cause

habituation deficits (hereinafter referred to as habituation deficient genes) by recapitulating

published habituation deficits of *Drosophila* ID null mutant models *G9a* (22) and *Synapsin* (41), and of

the classical learning and memory mutant *dunce* (27, 42, 43) (**Figure 1B,C,D**). This demonstrates that

light-off jump habituation in combination with RNAi can efficiently identify genetic regulators of

habituation learning.

In our screen, we found that the *Drosophila* orthologs of 98 human ID genes (35% of all

investigated orthologs) are required, in neurons, for habituation learning. This phenotype represents

a highly specific defect in behavioral adaptation to the stimulus; flies keep on jumping in response to

the repetitive light-off stimulus, illustrating that they do not suffer from broad neuronal transmission

deficits (which would disable jumping), fatigue, sensory or other deficiencies. 27% of ID gene

orthologs had no effect on habituation, whereas 41% of genes fell into the category of "reduced

organismal fitness", and 8% of genes showed "premature habituation" without detectable fatigue.

The complete list of habituation screen results and distribution of human ID genes in phenotype

categories can be found in **Table S2, S3**. The screen thus identified nearly a hundred disease genes

controlling habituation learning.

Drosophila **habituation deficits characterize ID genes associated with macrocephaly**

To understand whether habituation deficits in *Drosophila* represent a proxy of specific phenotypes in human individuals, we performed enrichment analysis among ID-associated clinical core features (39). We found that orthologs of ID genes associated with habituation deficits in *Drosophila* are specifically enriched among ID genes associated with macrocephaly/overgrowth (**Figure 2A**, E=2.21, p=0.0095, **Table S4**) and vegetative anomalies (**Figure 2A**, E=2.14, p=0.0495, **Table S4**). In contrast, ID genes associated with the severe "reduced organismal fitness" phenotype category show enrichment in different, severe ID-associated features such as limb anomalies, brain malformations, endocrine and eye anomalies (**Figure S2**, **Table S4**). Moreover, ID genes not giving rise to habituation deficits ("not affected" category) did not show any enrichment among ID-associated clinical features (**Figure 2A**, **Table S4**).

Habituation deficits characterize ID genes associated with ASD and deficits in specific ASD-relevant behavioral domains

There is a long-known relationship between macrocephaly and autism (44). For this reason and because of the potential relevance of habituation deficits to ASD (10–12), we decided to further investigate the relationship of *Drosophila* habituation and human ASD. We used the Simons Simplex Collection (SSC) (37), a genetically and phenotypically well-characterized cohort of sporadic ASD individuals. We matched genes with likely gene-disrupting (LGD) *de novo* mutations (45) in this ASD cohort to those included in our experimental *Drosophila* habituation approach. 86 ASD individuals carried mutations in 59 of the investigated genes (**Table S5**). We first asked whether these ID plus ASD-associated genes preferentially fall into a specific *Drosophila* phenotype category. Strikingly, they are significantly enriched among the genes that in *Drosophila* caused habituation deficits (**Figure 3A,** E=1.57, p=0.014, **Table S4**, ASD SSC). Independently, significant enrichment was obtained for high-confidence ID plus ASD-associated genes identified from the SFARI database (38) (41 investigated genes, **Figure 3B,** E=1.58, p=0.023, **Table S4**, ASD SFARI), confirming a relationship between *Drosophila* habituation deficits and human ASD.

To further characterize the relationship between *Drosophila* habituation and human phenotypes, we divided the SSC individuals into two distinct clusters based on their habituation phenotype in the corresponding fly models: habituation deficits (N=41 individuals, 29 genes) and no habituation deficits (N=23 individuals, 16 genes) (**Table S5**). We compared both groups across five broad quantitative measures of behavior and cognition: cognitive ability (full-scale IQ); Social Responsiveness Scale (SRS); depression and anxiety (Child Behavior Checklist Internalizing Disorders, CBCL-Int); impulsivity, attention and conduct (Child Behavior Checklist Externalizing Disorders, CBCL-Ext); and atypical behavior (Aberrant Behavior Checklist, ABC). There was no significant difference for IQ (p=0.80), SRS (p=0.49), or CBCL-Int (p=0.07), but a significant difference for CBCL-Ext (p=0.04) and ABC (p=0.004; **Figure 3C**, **Table S6)**. The latter persists even in case of multiple testing correction (p<0.01). This effect is mainly driven by the ABC subdomains of inappropriate, stereotypic speech (p=0.003), with a nominal contribution from the subdomains of hyperactivity (p=0.016) and irritability (p=0.03) but not by lethargy (p=0.11) or stereotypy (p=0.16) (**Table S6**). In summary, these data provide a direct indication that habituation deficits in *Drosophila* are relevant to ASD-implicated genes. They also suggest that SSC individuals carrying LGD mutations in genes associated with habituation deficits in *Drosophila* show a higher rate and/or severity of atypical behaviors associated with stereotypic speech, hyperactivity and irritability.

Habituation deficits characterize ID genes with synaptic function

Specificity of "habituation deficient" genes in distinct human disease phenotypes suggests that these may converge on specific mechanisms also at the molecular level. ID genes are known to be enriched in a number of biological processes, but which are important for habituation? Performing an enrichment analysis of ID-enriched Gene Ontology-based (GO) categories (see **SM**) against the stringent background of the investigated ID genes, we found that "habituation deficient" genes are significantly enriched in a sole GO-based category: processes related to the synapse (22/44 ID genes, E=1.59, p=0.024, **Figure 4**, **Table S4**). No enriched GO terms were found in the "not affected"

category. Together, our results support synaptic processes to be crucial for habituation, as previously

shown for other forms of this behavior (46, 47).

Molecular networks and modules underlying habituation

With the rich repertoire of nearly a hundred genes required for habituation that moreover show

specificity for ASD and synapse function, we set out to determine the molecular pathways these

genes are operating in. ID gene products are significantly interconnected via protein-protein

interactions (48, 49). Consistent with previously published findings (39), ID genes investigated in our

screen are 1.69 times enriched in interactions compared to 1000 randomly chosen protein sets of the

same size and number of known interactions (physical interaction enrichment (PIE) score (50) =1.69;

p<0.001). To identify biologically relevant modules, we resolved this network into communities with

even tighter interconnectivity using unsupervised community clustering (51). This analysis resulted in

26 communities containing 109 proteins (**Figure 5A, Table S7**). Their proximity and specificity for ID-

enriched GO-based processes are depicted in **Figure S3**. Mapping "habituation deficient" genes onto

the communities (**Figure 5A, red circles**) highlighted modules with high incidence of habituation

deficits (**Figure 5A**).

A key role for ID and ASD-associated Ras signaling in habituation

Five communities form a large, interconnected module with high incidences of habituation deficits.

However, the tightly interconnected hub at its center is characterized by the absence of habituation

deficits (**Figure 5A, square)**. This hub represents the key proteins of Ras-MAPK signaling (**Figure 5B**).

This pathway, best known for its role in cancer, underlies a group of disorders collectively referred as

Rasopathies. Importantly, while 92% of the modeled ID disorders are thought to result from loss of

function of the underlying genes, Rasopathies are caused by gain-of-function mutations in the core

pathway (**Figure 5C, Table S1**). The RNAi approach used in this screen, despite addressing gene

function, did thus not recapitulate the molecular pathology of these specific cognitive disorders.

However, Rasopathies can also result from loss of function in negative regulators of the pathway. We

therefore asked whether exactly the same genetic mechanisms that cause Rasopathies in humans

also hold true for habituation deficits in *Drosophila*. In our screen, we tested habituation of three

negative regulators of Ras: NF1 (*Drosophila* Nf1), which can directly inactivate Ras through the GAP

enzymatic activity (52), SPRED1 (*Drosophila* Spred) (53, 54), and CASK (*Drosophila* CASK) (55).

Panneuronal knockdown of either negative regulator caused strong habituation deficits (**Figure 5D, in**

red). We therefore tested a constitutively active *Ras* mutant, *Ras1^{R68Q}* (56). Heterozygous *Ras1^{R68Q}*

flies showed strong habituation deficits compared to the control flies with the same genetic

background (p=3.56x10^{-9}; **Figure 5D, in green**). Moreover, the same was true when we

overexpressed, specifically in neurons, the same mutant *Ras* allele from an inducible transgenic

construct (p=1.96x10^{-6}; **Figure 5D, in green**). We conclude that increased activity of Ras, causing

Rasopathies and associated cognitive deficits in humans, causes habituation deficits in *Drosophila*.

Habituation-inhibiting function of increased Ras-MAPK signaling maps to inhibitory/GABAergic

neurons

We next aimed to identify in which type of neurons the habituation-inhibiting function of Ras-MAPK

signaling resides. Because the well-characterized neurons of the giant fiber circuit controlling the

light-off jump response in *Drosophila* are cholinergic (57), just as the majority of excitatory neurons

in *Drosophila*, we first tested whether increased Ras-MAPK signaling activity would induce

habituation deficits when directed to cholinergic neurons. For this, we adopted the knockdown of

negative Ras regulators (*Nf1*, *Spred* and *CASK*) and expressed constitutively active *Ras1* (*Ras1^{R68Q}*). In

addition, we tested expression of a gain-of-function allele of *Raf* (*Raf^{GOF}*), a downstream mediator of

Ras signaling. We crossed these inducible UAS alleles to the cholinergic Cha-Gal4 driver and assessed

habituation. None of these conditions recapitulated the panneuronally evoked habituation deficits

(**Figure 6A**).

Because of the recently established role of GABAergic neurons in *Drosophila* olfactory and

proboscis extension reflex habituation (28, 58, 59) and the emerging importance of GABA inhibition

in autism (60), we next targeted GABA neurons using the Gad1-Gal4 driver and the same toolbox to

increase Ras-mediated signaling activity. Notably, this consistently induced habituation deficits in all

tested conditions (**Figure 6B**), recapitulating the observed panneuronally evoked deficits. We

conclude that the habituation-inhibiting function of increased Ras-MAPK signaling maps to

GABAergic neurons.

Ras-MAPK signaling in cholinergic neurons is essential for habituation learning

In the panneuronal screen, RNAi-mediated knockdown of the Ras pathway core components *Ras, Raf*

and *Mek* resulted, likely due to their promiscuous role and fundamental importance, in impaired

jump response or increased fatigue ("reduced organismal fitness" category). This could potentially

mask an essential role for Ras signaling in habituation, in addition to the habituation-inhibiting

function of increased Ras-MAPK signaling. In fact, in our panneuronal RNAi screen not only

knockdown of negative regulators but also of two established positive Ras-MAPK regulators, *Sos* and

Csw, resulted in habituation deficits. We therefore addressed whether decreased Ras-MAPK signaling

could also be critical for habituation learning. We downregulated Ras-MAPK activity by crossing the

UAS-based RNAi lines targeting *Sos* and *Csw*, but also RNAi lines targeting *Ras*, *Raf* and *Mek,* to the

GABAergic driver Gad1-Gal4. We did not observe any detrimental effect on habituation (**Figure 6D**).

In contrast, downregulating Ras-MAPK signaling using the same toolset in cholinergic neurons

consistently prevented normal habituation learning (**Figure 6C**). We conclude that Ras-MAPK

signaling is essential in cholinergic but not in GABAergic neurons. Thus, Ras-MAPK signaling plays a

dual, opposing role in inhibitory versus excitatory neurons in habituation learning.

Discussion:

Drosophila **screen identifies deficits in habituation learning as a widely affected mechanism in**

autism

To systematically address the genetic basis of habituation deficits associated with

neurodevelopmental disorders, we investigated 286 ID/ASD genes with a clear *Drosophila* ortholog in

light-off jump habituation. Panneuronal knockdown of 98 ID/ASD genes specifically suppressed the

adaptive habituation response to repeated stimulation without affecting organismal health or

fitness. Follow-up work on the Ras-MAPK pathway raised this number to 104. 93 of these are novel

regulators of habituation, substantially exceeding the sum of previously recognized habituation

genes across species and paradigms. Our data strongly suggest that deficits in habituation learning

are a widely affected mechanism in ID/ASD. Stringent criteria for RNAi specificity and correction for

multiple testing (see **SM**) in our experiments ensured a minimal level of potential false positive

discoveries. Of eleven previously identified ID/ASD genes with habituation deficits, our screen

confirmed eight (**Table S8**). Habituation deficits might therefore be a hallmark of even more ID/ASD

genes than determined here. In particular, the phenotype category of "reduced organismal fitness" is

likely to contain genes with promiscuous functions masking a specific role in habituation learning.

Enrichment analysis of ID-associated clinical features revealed that "habituation deficient" ID

genes are preferentially characterized by macrocephaly/overgrowth. Macrocephaly has been since

long associated with ASD (44). Strikingly, we found that mutations in genes associated with

Drosophila habituation deficits are significantly overrepresented among ASD individuals (SFARI

database, 50%; SSC cohort: 48% of genes). SSC individuals carrying mutations in these genes show a

high rate and/or severity of aberrant behaviors associated with stereotypic speech, hyperactivity and

irritability, suggesting that habituation deficits may contribute to specific behavioral anomalies in

ASD. Habituation deficits thus represent a common phenotypic signature of ASD in *Drosophila* and

highlight specific behavioral subdomains affected in ASD.

349 **Synapse-related processes and Ras-MAPK signaling play a key role in habituation**

350 Synapse biology has been proposed to play a central role in ASD (61). Our data show that among the

351 investigated disease genes, "habituation deficient" genes are specifically enriched in genes with

352 synaptic function. This is in line with habituation representing a measurable form of synaptic

353 plasticity (8, 47).

354 Analyzing the distribution of "habituation deficient" genes in ID-specific molecular

355 interaction networks, we discovered that they accumulate in a multiple-community module and

356 connect to the Ras-MAPK pathway core proteins Ras, Raf and Mek (**Figure 5A,B**). We observed

357 habituation deficits upon panneuronal knockdown of Ras negative regulators and panneuronal

358 expression of the constitutively active *Ras* allele *Ras1^{R68Q}* (**Figure 5C**), demonstrating that increased

359 Ras-mediated signaling causes habituation deficits. Moreover, proteins encoded by "habituation

360 deficient" genes form a significantly interconnected module (**Figure 7**). The coherence of this module

361 further supports the validity of the chosen RNAi approach to identify genes and molecular processes

362 regulating habituation learning. The module contains a number of synaptic proteins (**Figure 7**) with

363 not yet investigated roles in Ras signaling. It would be interesting to determine whether some of

364 these enlarge the spectrum of diseases caused by deregulated Ras signaling.

365

366 **Ras-MAPK signaling exerts a dual but opposing role in inhibitory versus excitatory neurons, a novel**

367 **systems-level mechanism**

368 Identification of neuronal substrates in which specific ID genes are required to warrant habituation

369 learning is important for understanding of the neuronal mechanisms underlying habituation in flies

370 and cognitive dysfunction in individuals with ID/ASD. Restoring the function of affected neurons

371 might represent a suitable treatment strategy. The light-off jump startle circuit of *Drosophila* is

372 relatively simple and well described. The chemical component is *Drosophila*'s major excitatory

373 neurotransmitter, acetylcholine (57). However, it is not known how habituation of this circuit is

374 regulated. The commonly accepted view regards synaptic depression in excitatory neurons, induced

by repetitive stimulation, as the underlying mechanism (46, 62). This has recently been challenged by

Ramaswami and colleagues who showed that plasticity of inhibitory, GABAergic neurons drives two

non-startle types of habituation (58, 59). We found that increased activity of our identified key

pathway, Ras-MAPK, in GABAergic but not in cholinergic neurons causes deficits in light-off jump

habituation. Our results thus support inhibitory circuits as crucial components of habituation learning

across different paradigms and sensory modalities. Further experiments are needed to establish the

direct involvement of GABAergic signaling. At the same time, we identified that also decreased Ras-

MAPK signaling activity can lead to habituation deficits. Yet, the neuronal substrates of these deficits

are different and map to excitatory, cholinergic neurons. Although our experiments do not

distinguish between developmental effects and acute circuit plasticity, the opposing role for Ras-

MAPK signaling on habituation may provide new insights into mechanisms of neural plasticity in

health and disease. It may also have crucial implications for treatment of Rasopathies. Future clinical

trials, as opposed to those that broadly decreased Ras activity and failed (63), may need more

attention towards restoring circuit function and balance.

Translational value and application of cross-species habituation measures for diagnosis and

treatment of ID and ASD

Based on our findings that habituation is widely affected in *Drosophila* models of ID, and that

habituation deficits are particularly common among genes also implicated in ASD, we propose that

disrupted habituation may contribute to ID/ASD pathology.

The emerging importance of inhibitory inputs for habituation ((28, 58) and this study) and

sensory information filtering in the cortical centers of the brain (64, 65) suggests the existence of an

overarching circuit-based mechanism responsible for prevention of inappropriate behavioral

responses (8). Though our findings that habituation deficits in *Drosophila* correlate with increased

rate and/or severity of atypical behaviors compared to ID/ASD genes without habituation deficits

should be replicated, we speculate that disrupted habituation arising from GABAergic defects

represents a mechanism that contributes to these core ID/ASD features. As such, it has potential to

serve as a cross-species, mechanism-specific functional readout—a pressing need for efficient

personalized (pharmacological) treatment in the field of neurodevelopmental disorders.

Implementing suitable low-burden protocols for habituation measures in clinical research and

diagnostics of ID/ASD, such as those developed for investigation of habituation deficits in Fragile X

syndrome (13), will help to further delineate the affected cognitive domains that arise from deficient

habituation. In future clinical trials, these could serve as objective and quantitative readouts for

patient stratification in mechanism-based treatment strategies and for monitoring of drug efficacy.

Dissection of the underlying defective mechanisms in *Drosophila* can at the same time identify novel

targets for treatment, with high-throughput light-off jump habituation serving as a translational

pipeline for drug testing.

Acknowledgements:

We acknowledge the Vienna *Drosophila* Resource Center and Bloomington *Drosophila* Stock Center (NIH P40OD018537) for providing *Drosophila* strains. We thank the anonymous expert referees for their help to improve our manuscript. This research was supported in part by the European Union's FP7 large-scale integrated network Gencodys (HEALTH-241995) to Z.A. and A.S., by FP7 TACTICS, OPTIMISTIC, Aggressotype and MATRICS (HEALTH grant agreement numbers n°278948, n°305697, n°602805 and n°603016) to J.C.G., by a TOP grant (912-12-109) from The Netherlands Organization for Scientific Research (NWO), by a Horizon 2020 Marie Sklodowska-Curie European Training Network grant (MiND, 643051), by a grant from the Jérôme Léjeune foundation to A.S., and by U.S. National Institute for Mental Health (NIMH) funding (R01MH101221 to E.E.E. & R01MH100047 to R.B.). E.E.E is an investigator of the Howard Hughes Medical Institute. We are grateful to all of the families at the participating Simons Simplex Collection (SSC) sites, as well as the principal investigators (A. Beaudet, R. Bernier, J. Constantino, E. Cook, E. Fombonne, D. Geschwind, R. Goin-Kochel, E. Hanson, D. Grice, A. Klin, D. Ledbetter, C. Lord, C. Martin, D. Martin, R. Maxim, J. Miles, O. Ousley, K. Pelphrey, B. Peterson, J. Piggot, C. Saulnier, M. State, W. Stone, J. Sutcliffe, C. Walsh, Z. Warren, E. Wijsman). We appreciate obtaining access to phenotypic data on SFARI Base. Approved researchers can obtain the SSC population dataset described in this study (http://sfari.org/resources/simons-simplex-collection) by applying at https://base.sfari.org.

Financial disclosures:

In the past 3 years, J.C.G. has acted as a consultant to Boehringer Ingelheim GmbH but is not an

employee, stock- or share-holder of this company. He has no other financial or material support to

declare, including expert testimony, patents and royalties. E.E.E. is on the scientific advisory board

(SAB) of DNAnexus, Inc.. Z.S. is a director of Aktogen ltd.. M.F., L.E., P.C., E.L.S., L.R.B., J.IH., C.Z.,

R.A.B. and A.S. declare that they have no conflict of interests.

References:

1. Peeke H (1973): *Habituation: Behavioral studies*. (Vol. 1), Academic Press.

2. Colombo J, Mitchell DW (2009): Infant visual habituation. *Neurobiol Learn Mem*. 92: 225–234.

3. Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton D, Colombo J, *et al.* (2009): Habituation Revisited: An Updated and Revised Description of the Behavioral Characteristics of Habituation. *Neurobiol Learn Mem*. 92: 135–138.

4. Barron HC, Vogels TP, Behrens TE, Ramaswami M (2017): Inhibitory engrams in perception and memory. *Proc Natl Acad Sci*. 201701812.

5. Kavšek M (2004): Predicting later IQ from infant visual habituation and dishabituation: A meta-analysis. *J Appl Dev Psychol*. 25.

6. Glanzman DL (2010): Common Mechanisms of Synaptic Plasticity in Vertebrates and Invertebrates. *Curr Biol*. 20. doi: 10.1016/j.cub.2009.10.023.

7. Schmid S, Wilson DA, Rankin CH (2015): Habituation mechanisms and their importance for cognitive function. *Front Integr Neurosci*. 77: 419–450.

8. Ramaswami M (2014): Network plasticity in adaptive filtering and behavioral habituation. *Neuron*. 82.

9. Sinha P, Kjelgaard MM, Gandhi TK, Tsourides K, Cardinaux AL, Pantazis D, *et al.* (2014): Autism as a disorder of prediction. *Proc Natl Acad Sci*. 111: 15220–5.

10. Pellicano E, Rhodes G, Calder AJ (2013): Reduced gaze aftereffects are related to difficulties categorising gaze direction in children with autism. *Neuropsychologia*. 51: 1504–1509.

11. Ewbank MP, Rhodes G, Von Dem Hagen EAH, Powell TE, Bright N, Stoyanova RS, *et al.* (2015): Repetition suppression in ventral visual cortex is diminished as a function of increasing autistic traits. *Cereb Cortex*. 25: 3381–3393.

12. Swartz JR, Wiggins JL, Carrasco M, Lord C, Monk CS (2013): Amygdala habituation and prefrontal functional connectivity in youth with autism spectrum disorders. *J Am Acad Child Adolesc Psychiatry*. 52: 84–93.

13. Ethridge LE, White SP, Mosconi MW, Wang J, Byerly MJ, Sweeney JA (2016): Reduced habituation of auditory evoked potentials indicate cortical hyper-excitability in Fragile X Syndrome. *Transl Psychiatry*. 6: e787.

14. Rigoulot S, Knoth IS, Lafontaine M-P, Vannasing P, Major P, Jacquemont S, *et al.* (2017): Altered visual repetition

suppression in Fragile X Syndrome: New evidence from ERPs and oscillatory activity. *Int J Dev Neurosci*. 59: 52–59.

15. Restivo L, Ferrari F, Passino E, Sgobio C, Bock J, Oostra BA, *et al.* (2005): Enriched environment promotes behavioral and

morphological recovery in a mouse model for the fragile X syndrome. *Proc Natl Acad Sci U S A*. 102: 11557–11562.

16. Lovelace JW, Wen TH, Reinhard S, Hsu MS, Sidhu H, Ethell IM, *et al.* (2016): Matrix metalloproteinase-9 deletion rescues

auditory evoked potential habituation deficit in a mouse model of Fragile X Syndrome. *Neurobiol Dis*. 89: 126–135.

17. Wolman MA, deGroh ED, McBride SM, Jongens TA, Granato M, Epstein JA (2014): Modulation of cAMP and Ras

Signaling Pathways Improves Distinct Behavioral Deficits in a Zebrafish Model of Neurofibromatosis Type 1. *Cell Rep*.

8: 1265–1270.

18. Kirshenbaum GS, Clapcote SJ, Duffy S, Burgess CR, Petersen J, Jarowek KJ, *et al.* (2011): Mania-like behavior induced by

genetic dysfunction of the neuron-specific Na+,K+-ATPase α3 sodium pump. *Proc Natl Acad Sci U S A*. 108: 18144–9.

19. Cheli VT, Adrover MF, Blanco C, Verde ER, Guyot-Revol V, Vidal R, *et al.* (2002): Gene transfer of NMDAR1 subunit

sequences to the rat CNS using herpes simplex virus vectors interfered with habituation. *Cell Mol Neurobiol*. 22: 303–

314.

20. Walsh J, Desbonnet L, Clarke N, Waddington JL, O'Tuathaigh CMP (2012): Disruption of exploratory and habituation

behavior in mice with mutation of DISC1: An ethologically based analysis. *J Neurosci Res*. 90: 1445–1453.

21. McBride SMJ, Choi CH, Wang Y, Liebelt D, Braunstein E, Ferreiro D, *et al.* (2005): Pharmacological rescue of synaptic

plasticity, courtship behavior, and mushroom body defects in a Drosophila model of Fragile X syndrome. *Neuron*. 45:

753–764.

22. Kramer JM, Kochinke K, Oortveld MAW, Marks H, Kramer D, de Jong EK, *et al.* (2011): Epigenetic regulation of learning

and memory by Drosophila EHMT/G9a. *PLoS Biol*. 9. doi: 10.1371/journal.pbio.1000569.

23. van der Voet M, Nijhof B, Oortveld MAW, Schenck A (2014): Drosophila models of early onset cognitive disorders and

their clinical applications. *Neurosci Biobehav Rev*. 46.

24. Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, *et al.* (2007): A genome-wide transgenic RNAi library for

conditional gene inactivation in Drosophila. *Nature*. 448: 151–6.

25. Bellen HJ, Tong C, Tsuda H (2010): 100 years of Drosophila research and its impact on vertebrate neuroscience: a history

lesson for the future. *Nat Rev Neurosci*. 11: 514–22.

26. Duerr JS, Quinn WG (1982): Three Drosophila mutations that block associative learning also affect habituation and

sensitization. *Proc Natl Acad Sci U S A*. 79: 3646–3650.

27. Asztalos Z, Arora N, Tully T (2007): Olfactory Jump Reflex Habituation in Drosophila and Effects of Classical Conditioning

Mutations. *J Neurogenet*. 21: 1–18.

28. Larkin A, Karak S, Priya R, Das A, Ayyub C, Ito K, *et al.* (2010): Central synaptic mechanisms underlie short-term olfactory

habituation in Drosophila larvae. *Learn Mem*. 17: 645–653.

29. Acevedo SF, Froudarakis EI, Kanellopoulos A, Skoulakis EMC (2007): Protection from premature habituation requires

functional mushroom bodies in Drosophila. *Learn Mem*. 14: 376–84.

30. Engel JE, Wu CF (1996): Altered habituation of an identified escape circuit in Drosophila memory mutants. *J Neurosci*.

16: 3486–99.

31. van Bon BWM, Oortveld MAW, Nijtmans LG, Fenckova M, Nijhof B, Besseling J, *et al.* (2013): CEP89 is required for

mitochondrial metabolism and neuronal function in man and fly. *Hum Mol Genet*. 22: 3138–3151.

32. Willemsen MH, Nijhof B, Fenckova M, Nillesen WM, H F Bongers EM, Castells-Nobau A, *et al.* (2013): GATAD2B loss-of-

function mutations cause a recognisable syndrome with intellectual disability and are associated with learning

deficits and synaptic undergrowth in Drosophila. *J Med Genet*. 50: 507–514.

33. Lugtenberg D, Reijnders MRF, Fenckova M, Bijlsma EK, Bernier R, van Bon BWM, *et al.* (2016): De novo loss-of-function

mutations in WAC cause a recognizable intellectual disability syndrome and learning deficits in Drosophila. *Eur J Hum*

Genet. 1–9.

34. Stessman HAF, Willemsen MH, Fenckova M, Penn O, Hoischen A, Xiong B, *et al.* (2016): Disruption of POGZ Is Associated

with Intellectual Disability and Autism Spectrum Disorders. *Am J Hum Genet*. 98: 541–552.

35. Esmaeeli-Nieh S, Fenckova M, Porter IM, Motazacker MM, Nijhof B, Castells-Nobau A, *et al.* (2016): BOD1 Is Required

for Cognitive Function in Humans and Drosophila. *PLoS Genet*. 12. doi: 10.1371/journal.pgen.1006022.

36. Kelly LE (1983): An altered electroretinogram transient associated with an unusual jump response in a mutant of

Drosophila. *Cell Mol Neurobiol*. 3: 143–9.

37. Fischbach GD, Lord C (2010): The simons simplex collection: A resource for identification of autism genetic risk factors.

Neuron. 68.

38. Basu SN, Kollu R, Banerjee-Basu S (2009): AutDB: A gene reference resource for autism research. *Nucleic Acids Res*. 37.
doi: 10.1093/nar/gkn835.

39. Kochinke K, Zweier C, Nijhof B, Fenckova M, Cizek P, Honti F, *et al.* (2016): Systematic Phenomics Analysis Deconvolutes
Genes Mutated in Intellectual Disability into Biologically Coherent Modules. *Am J Hum Genet*. 98: 149–164.

40. Oortveld MAW, Keerthikumar S, Oti M, Nijhof B, Fernandes AC, Kochinke K, *et al.* (2013): Human Intellectual Disability
Genes Form Conserved Functional Modules in Drosophila. *PLoS Genet*. 9. doi: 10.1371/journal.pgen.1003911.

41. Sadanandappa MK, Blanco Redondo B, Michels B, Rodrigues V, Gerber B, VijayRaghavan K, *et al.* (2013): Synapsin
function in GABA-ergic interneurons is required for short-term olfactory habituation. *J Neurosci*. 33: 16576–85.

42. Dudai Y, Jan YN, Byers D, Quinn WG, Benzer S (1976): dunce, a mutant of Drosophila deficient in learning. *Proc Natl
Acad Sci U S A*. 73: 1684–8.

43. Tempel BL, Bonini N, Dawson DR, Quinn WG (1983): Reward learning in normal and mutant Drosophila. *Proc Natl Acad
Sci*. 80: 1482–1486.

44. Lainhart JE, Piven J, Wzorek M, Landa R, Santangelo SL, Coon H, Folstein SE (1997): Macrocephaly in children and adults
with autism. *J Am Acad Child Adolesc Psychiatry*. 36: 282–290.

45. Iossifov I, O'roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, *et al.* (2014): The contribution of de novo coding
mutations to autism spectrum disorder. *Nature*. 13: 216–221.

46. Castellucci V, Pinsker H, Kupfermann I, Kandel ER (1970): Neuronal Mechanisms of Habituation and dishabituation of
the gill withdrawal reflex in Aplysia. *Science (80-)*. 167: 1745–1748.

47. Poon C-S, Young DL (2006): Nonassociative learning as gated neural integrator and differentiator in stimulus-response
pathways. *Behav Brain Funct*. 2: 29.

48. Cristino a S, Williams SM, Hawi Z, An J-Y, Bellgrove M a, Schwartz CE, *et al.* (2014): Neurodevelopmental and
neuropsychiatric disorders represent an interconnected molecular system. *Mol Psychiatry*. 19: 294–301.

49. Honti F, Meader S, Webber C (2014): Unbiased Functional Clustering of Gene Variants with a Phenotypic-Linkage
Network. *PLoS Comput Biol*. 10. doi: 10.1371/journal.pcbi.1003815.

50. Sama IE, Huynen MA (2010): Measuring the physical cohesiveness of proteins using Physical Interaction Enrichment
(PIE). *Bioinformatics*. 26: 2737–2743.

51. Kalinka AT, Tomancak P (2011): linkcomm: An R package for the generation, visualization, and analysis of link

communities in networks of arbitrary size and type. *Bioinformatics*. 27: 2011–2012.

52. Martin GA, Viskoohil D, Bollag G, McCabe PC, Crosier WJ, Haubruck H, *et al.* (1990): The GAP-related domain of the

neurofibromatosis type 1 gene product interacts with ras p21. *Cell*. 63: 843–849.

53. Dunzendorfer-Matt T, Mercado EL, Maly K, McCormick F, Scheffzek K (2016): The neurofibromin recruitment factor

Spred1 binds to the GAP related domain without affecting Ras inactivation. *Proc Natl Acad Sci*. 201607298.

54. Hirata Y, Brems H, Suzuki M, Kanamori M, Okada M, Morita R, *et al.* (2016): Interaction between a domain of the

negative regulator of the ras-ERK pathway, SPRED1 protein, and the GTPase-activating protein-related domain of

neurofibromin is implicated in legius syndrome and neurofibromatosis type 1. *J Biol Chem*. 291: 3124–3134.

55. Hsueh YP, Roberts AM, Volta M, Sheng M, Roberts RG (2001): Bipartite interaction between neurofibromatosis type I

protein (neurofibromin) and syndecan transmembrane heparan sulfate proteoglycans. *J Neurosci*. 21: 3764–3770.

56. Gafuik C, Steller H (2011): A gain-of-function germline mutation in drosophila ras1 affects apoptosis and cell fate during

development. *PLoS One*. 6. doi: 10.1371/journal.pone.0023535.

57. Allen MJ, Murphey RK (2007): The chemical component of the mixed GF-TTMn synapse in Drosophila melanogaster uses

acetylcholine as its neurotransmitter. *Eur J Neurosci*. 26: 439–445.

58. Das S, Sadanandappa MK, Dervan A, Larkin A, Lee JA, Sudhakaran IP, *et al.* (2011): Plasticity of local GABAergic

interneurons drives olfactory habituation. *Proc Natl Acad Sci U S A*. 108: E646-54.

59. Paranjpe P, Rodrigues V, VijayRaghavan K, Ramaswami M (2012): Gustatory habituation in Drosophila relies on rutabaga

(adenylate cyclase)-dependent plasticity of GABAergic inhibitory neurons. *Learn Mem*. 19: 627–635.

60. Robertson CE, Ratai EM, Kanwisher N (2016): Reduced GABAergic Action in the Autistic Brain. *Curr Biol*. 26: 80–85.

61. Bourgeron T (2015): From the genetic architecture to synaptic plasticity in autism spectrum disorder. *Nat Rev Neurosci*.

16: 551–563.

62. Weber M, Schnitzler HU, Schmid S (2002): Synaptic plasticity in the acoustic startle pathway: The neuronal basis for

short-term habituation? *Eur J Neurosci*. 16: 1325–1332.

63. van der Vaart T, Plasschaert E, Rietman AB, Renard M, Oostenbrink R, Vogels A, *et al.* (2013): Simvastatin for cognitive

deficits and behavioural problems in patients with neurofibromatosis type 1 (NF1-SIMCODA): A randomised, placebo-

controlled trial. *Lancet Neurol*. 12: 1076–1083.

64. Isaacson JS, Scanziani M (2011): How inhibition shapes cortical activity. *Neuron*. 72.

65. Geramita MA, Burton SD, Urban NN (2016): Distinct lateral inhibitory circuits drive parallel processing of sensory

information in the mammalian olfactory bulb. *Elife*. 5. doi: 10.7554/eLife.16039.001.

Supplemental information

Document S1.

Figures S1-S3, Tables S5, S8, Supplemental Methods

Document S2.

Table S1

List of investigated human ID genes

Table S2. List of investigated human ID genes, corresponding Drosophila orthologs, RNAi lines, and

habituation results

Table S3

Distribution of human ID genes in four phenotype categories identified in Drosophila light-off jump

habituation screen

Table S4. Enrichments

Table S5. List of investigated SSC and SFARI genes

Table S7. Interaction communities

Figure 1. Habituation screen of intellectual disability genes, phenotype distribution and proof of principle

(A) Procedure, phenotype categories and phenotype distribution of the light-off jump habituation screen. Knockdowns that resulted in lethality, no jumper phenotype (defined as less than 50% flies jumping in at least one of the first five light-off trials) or premature habituation plus increased fatigue were assigned to the category "reduced organismal fitness" and their habituation was not further analyzed. Other phenotype categories are "habituation deficient", "not affected", and "premature habituation" (the latter if no fatigue was detected in secondary assay, see example in **Figure S1**). Drosophila orthologs of 34% of the investigated human ID genes were associated with defects in habituation learning. See also T**able S2, S3**. (B, C, D) Defective habituation upon neuron-specific RNAi-mediated knockdown of G9a, Synapsin (syn), and dunce (dnc) (2xGMR-wIR/+; UAS-RNAi/elav-Gal4, UAS-Dicer-2, in red) compared to their respective genetic background controls (2xGMR-wIR/+; elav-Gal4, UAS-Dicer-2/+, in gray). Jump response curves show the average jump response (% of jumping flies) over 100 light-off trials at 1 s inter-trial interval). Mean TTC: the mean number of trials that flies needed to reach the no-jump criterion (see Methods and Materials) presented as Mean TTC ± SEM. *** padj<0.001, ** padj<0.01, based on FDR-corrected lm analysis. A complete list of ID/ASD genes with previously identified habituation defects is provided as **Table S8**, adding further proof of principle.

Figure 2. Habituation defects in Drosophila characterize ID genes associated with macrocephaly in humans
Enrichment of Drosophila phenotype categories across 27 ID-accompanying clinical features (39). "Habituation deficient" genes show specificity for macrocephaly and/or overgrowth (E=2.21, p=0.0095) and vegetative anomalies (E=2.14, p=0.049) ** p<0.01, * p<0.05, based on Fisher's Exact test. For enrichment of "reduced organismal fitness" category, see Figure S2. Enrichment scores, p-values and enriched genes are listed in **Table S4**.

Figure 3. Habituation defects in Drosophila characterize ID genes associated with ASD and deficits in specific behavioral domains

(A,B) Enrichment of Drosophila phenotype categories "habituation deficient" and "not affected" in ID plus ASD-associated genes identified in SFARI database (ASD SFARI, E=1.58, p=0.023, (A)) and SSC cohort (ASD SSC, E=1.57, p=0.014 (B)). Circles represent total number of tested ID plus ASD-associated genes. (C) Genes associated with "habituation deficient" versus "not affected" phenotype categories in Drosophila show significantly more aberrant behaviors on the ABC (F(1,62)=8.9, p=0.004) and more CBCL externalizing symptoms (F(1,62)=4.3, with suggestive p-value, p=0.04) in the ASD SSC cohort. Data presented as mean score ± SEM. ** p<0.01, * p<0.05, based on MANOVA. See also **Table S5** (list of ASD SFARI and ASD SSC genes) and **Table S6** (complete MANOVA results).

Figure 4. Habituation defects in Drosophila characterize ID genes with synapse-related functions

Of 25 gene ontology (GO)-based processes, "habituation deficient" genes are specifically and significantly enriched in processes related to synapse (E=1.59, p=0.024). Genes with no effect on habituation do not show significant enrichment in any GO process. * p<0.05, based on Fisher's exact test. All enrichment scores, p-values and enriched genes are listed in **Table S4**.

Figure 5. A central role for Ras-MAPK signaling in habituation learning

(A) Highly connected communities identified by unbiased community clustering, colored by their functional proximity (**Figure S3**). Red circles and gene names highlight nodes representing "habituation deficient" genes. For complete list of communities and genes see **Table S7**. (B) Nodes connecting four communities from the central module represent the core components of Ras-MAPK signaling. (C) Schematic representation of Ras-MAPK signaling and associated mechanisms in ID disorders called 'Rasopathies'.

(D) Increasing Ras signaling by inducing either loss of function of negative Ras regulators (left side of pathway scheme) or by constitutively activating Ras (right side) disrupts habituation learning. Left: Defective habituation upon neuron-specific knockdown of negative Ras regulators, Nf1 (2xGMR-wIR/+; Nf1-RNAivdrc35877/elav-Gal4, UAS-Dicer-2, N=72, in red), Spred (2xGMR-wIR/+; Spred-RNAivdrc18024/elav-Gal4, UAS-Dicer-2, N=73, in red), and CASK (2xGMR-wIR/+; CASK-RNAivdrc34184/elav-Gal4, UAS-Dicer-2, N=75, in red) compared to their corresponding genetic background controls (2xGMR-wIR/+; elav-Gal4, UAS-Dicer-2/+. N: 55, 20, 25, in gray). *** padj<0.001, based on lm analysis and FDR correction in the screen (see **Methods and Materials**). Mean TTC ± SEM.Right: Defects in habituation learning in a heterozygous, constitutively active Ras mutant (Ras1R68Q/+, N=55, in green) compared to its genetic background control (N=43 in gray), and upon neuron-specific expression of Ras1R68Q (elav>Ras1R68Q: UAS-Ras1R68Q/2xGMR-wIR; elav-Gal4, UAS-Dicer-2/+, N=52, in green) compared to its genetic background control (2xGMR-wIR/+; elav-Gal4, UAS-Dicer-2/+, N=34, in gray). *** p<0.001, based on lm analysis. Data presented as Mean TTC ± SEM

Figure 6. Dual, opposing role of Ras-MAPK signaling in GABAergic and cholinergic neurons in the regulation of habituation learning

(A) No effect on habituation of Ras1R68Q (N=51, in green), Nf1-RNAi (N=38, in red), Spred1-RNAi (N=55, in red) and CASK-RNAi (N=39, in red) upon expression in cholinergic neurons compared to their respective genetic background controls (Cha-Gal4/+; 2xGMR-wIR/+, N: 54, 45, 54, 52, in gray). Expression of RafGOF in cholinergic neurons resulted in lethality. (B) Defective habituation of Ras1R68Q (N=52, in green), RafGOF (N=57, in green), Nf1-RNAi (N=55, in red), Spred1-RNAi (N=37, in red) and CASK-RNAi (N=54, in red) on habituation upon expression in GABAergic neurons compared to their respective genetic background controls (Gad1-Gal4/+; 2xGMR-wIR/+, N: 50, 50, 39, 58, 46, in gray). (C) Defective habituation of Csw-RNAi (UAS-Csw-RNAivdrc21756/Y; Cha-Gal4/+; 2xGMR-wIR/+, N=58), Sos1-RNAi (UAS-Sos1-RNAivdrc42848/Cha-Gal4; 2xGMR-wIR/+, N=56), Ras1-RNAi (UAS-Ras1-RNAivdrc106642/Cha-Gal4; 2xGMR-wIR/+, N=55), Raf-RNAi (UAS-Raf-RNAivdrc20909/Cha-Gal4; 2xGMR-wIR/+, N=59) and Mek-RNAi (Cha-Gal4/+; UAS-Mek-RNAivdrc40026/2xGMR-wIR, N=58) in cholinergic neurons (in green) compared to their respective genetic background controls (Cha-Gal4/+; 2xGMR-wIR/+, N: 62, 54, 34, 46, 46, in gray). (D) No effect on habituation of Csw-RNAi (N=58), Sos1-RNAi (N=51), Ras1-RNAi (N=53), Raf-RNAi (N=52) and Mek-RNAi (N=54) in GABAergic neurons (in green) compared to their respective genetic background controls (Gad1-Gal4/+; 2xGMR-wIR/+, N: 60, 46, 54, 39, 39, in gray). Data presented as Mean TTC ± SEM. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, based on lm analysis.

Figure 7. Connections between "habituation deficient" genes

Connections between "habituation deficient" genes, including Ras, identified in the reference network used for community clustering (See **SM)** with significantly increased connectivity (PIE score =1.89, p<0.001). Nodes are colored based on the community, to which they belong. Nodes that represent "habituation deficient" genes but are not members of a community are labeled in black.