1 **Divergent gene expression among phytoplankton taxa in response to upwelling**
2

3 Robert H. Lampe,[1] Natalie R. Cohen,[1] Kelsey A. Ellis,[1] Kenneth W. Bruland,[2] Maria T.
4 Maldonado,[3] Tawnya D. Peterson,[4] Claire P. Till,[2,5] Mark A. Brzezinski,[6] Sibel Bargu,[7]
5 Kimberlee Thamatrakoln,[8] Fedor I Kuzminov,[8] Benjamin S. Twining[9] and Adrian Marchetti[1*]
6
[1]Department of Marine Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC,
USA
[2]Department of Ocean Sciences, University of California, Santa Cruz, CA, USA
[3]Department of Earth, Ocean, and Atmospheric Sciences, University of British Columbia,
Vancouver, BC, Canada
[4]Institute of Environmental Health, Oregon Health & Science University, Portland, OR, USA
[5]Department of Chemistry, Humboldt State University, Arcata, CA, USA
[6]The Marine Science Institute and the Department of Ecology Evolution and Marine Biology,
University of California, Santa Barbara, CA, USA
[7]Department of Oceanography and Coastal Sciences, School of the Coast and Environment,
Louisiana State University, Baton Rouge, LA, USA
[8]Department of Marine and Coastal Sciences, Rutgers, the State University of New Jersey, New
Brunswick, NJ, USA
[9]Bigelow Laboratory for Ocean Sciences, East Boothbay, ME, USA
21
*For correspondence:
Dr. Adrian Marchetti
Department of Marine Sciences
University of North Carolina at Chapel Hill
3202 Venable Murray Hall CB 3300
Chapel Hill, NC 27599-3300, USA
Email: amarchetti@unc.edu
Tel. (+1) 919 843 3473
Fax (+1) 919 962 1254
31

Abstract

Frequent blooms of phytoplankton occur in coastal upwelling zones creating hotspots of biological productivity in the ocean. As cold, nutrient-rich water is brought up to sunlit layers from depth, phytoplankton are also transported upwards to seed surface blooms that are often dominated by diatoms. The physiological response of phytoplankton to this process, commonly referred to as shift-up, is characterized by rapid growth rates and increases in nitrate assimilation. To examine the molecular underpinnings behind this phenomenon, metatranscriptomics was applied to a simulated upwelling experiment using natural phytoplankton communities from the California Upwelling Zone. An increase in diatom growth following five days of incubation was attributed to the genera *Chaetoceros* and *Pseudo-nitzschia*. Here we show that certain bloom-forming diatoms exhibit a distinct transcriptional response that coordinates shift-up where diatoms exhibited the greatest transcriptional change following upwelling; however, comparison of co-expressed genes exposed overrepresentation of distinct sets within each of the dominant phytoplankton groups. The analysis revealed that diatoms frontload genes involved in nitrogen assimilation likely in order to outcompete other groups for available nitrogen during upwelling events. We speculate that the evolutionary success of diatoms may be due, in part, to this proactive response to frequently encountered changes in their environment.

Introduction

Wind-driven coastal upwelling associated with eastern boundary currents delivers rich supplies of nutrients to illuminated surface waters. This phenomenon provides ideal conditions for blooms of phytoplankton that render coastal upwelling regimes centers of new production even though their relative ocean area is small (Capone and Hutchins, 2013). Typically dominated by large chain-forming diatoms, phytoplankton blooms in upwelling zones rapidly sequester carbon dioxide and are the base of short, efficient food chains that comprise a significant percentage of the global fish catch (Ryther, 1969; Estrada and Blasco, 1985; Lassiter et al., 2006; Lachkar and Gruber, 2013).

The phytoplankton community in upwelling zones is postulated to undergo a 'conveyer belt cycle' in which viable cells are upwelled into sunlit waters to seed a surface bloom. The community is then advected away from the upwelled source, and some cells eventually sink out of the photic zone. Surviving cells at depth and positioned in future upwelled waters are able to act as seed stock once winds are favorable for upwelling (Wilkerson and Dugdale, 1987; Wilkerson and Dugdale, 2008). This continuity between a subsurface population and surface bloom during an upwelling event has been observed through a combination of glider and remote sensing techniques (Seegers et al., 2015).

The physiological response of phytoplankton to being vertically transported into a higher light and nutrient-rich environment, commonly referred to as shift-up, includes rapid growth rates and strong increases in nitrate uptake and assimilation (MacIsaac et al., 1985; Wilkerson and Dugdale, 1987). This nitrate-related activity has been repeatedly observed in simulated upwelling mesocosm experiments (Dugdale and Wilkerson, 1989; Fawcett and Ward, 2011) and in a laboratory experiment on the diatom *Skeletonema costatum* (Smith et al., 1992). Shift-up, as expressed through rapid nitrate assimilation, is hypothesized to be linked to the success of diatoms in upwelling regions; it is believed that diatoms respond quickest to available nitrate once conditions are optimal (Fawcett and Ward, 2011). Characterization of this physiological response at a molecular level, however, is lacking. Only upregulation of the nitrogen assimilation gene, nitrate reductase, has been observed in *Skeletonema costatum* under lab-simulated upwelling conditions, indicating that there is a molecular basis for the shift-up response (Smith et al., 1992).

Metatranscriptomics is increasingly being applied to eukaryotic phytoplankton communities to provide a deeper understanding of molecular responses among resident phytoplankton groups (Caron et al., 2017). With the growing availability of reference

transcriptomes and genomes of eukaryotic phytoplankton, unprecedented levels and confidence in

gene annotation are being obtained from environmental sequences (Keeling et al., 2014; Alexander

et al., 2015b). Here we apply comparative metatranscriptomics to a simulated upwelling event in

a shipboard incubation experiment to characterize the phytoplankton community's response and

investigate the molecular basis for shift-up. Our results indicate that phytoplankton functional

groups exhibit highly distinct transcriptional responses to being upwelled in which diatoms

constitutively express genes involved in nitrogen assimilation. This strategy possibly allows

diatoms to outcompete other groups for available nitrogen once physical conditions are optimal

for growth.

Results and Discussion

Experimental overview and physiological observations

Results from the simulated upwelling experiment indicate that a bloom of large phytoplankton (>5

µm) was induced with observations of shift-up in terms of growth, primary production, and nitrate

uptake within these large cells. Macronutrient concentrations in the upwelled waters remained high

throughout the incubations; however, significant growth in the large (>5 µm) phytoplankton

community was observed (Fig. 1A and Supporting Information Fig. S1A). The initial dissolved

iron concentration was approximately 1.28 nmol L^{-1} which is marginally higher than the normal

values (<1 nmol L^{-1}) observed in the region. For complete drawdown of nitrate, an iron to nitrate

ratio of 8 nmol L^{-1}:20 µmol L^{-1} is typically required (Bruland et al., 2001). The initial ratio of 1.28

nmol L^{-1}:21.86 µmol L^{-1} in the incubations therefore indicates that iron had the potential to be a

limiting nutrient which resulted in 15 µmol L^{-1} of unused nitrate after 120 hours.

In addition to chlorophyll *a*, the upwelling simulation produced notable increases in

biogenic silica suggesting that the phytoplankton growth may be attributed to diatoms (Fig. 1A).

The presence of domoic acid also suggests a considerable presence of the diatom genus *Pseudo-*

nitzschia (Lelong et al., 2012). Concentrations were below the detection limit until 120 hours when

the average concentration was 1.34 µg L^{-1} (Supporting Information Fig. S1B). Although this

concentration is lower than peak concentrations observed during blooms in California coastal

waters (Schnetzer et al., 2013), it nevertheless indicates that *Pseudo-nitzschia* spp. were abundant

within the incubations by 120 hours.

This success of large phytoplankton is consistent with previous studies showing
phytoplankton from large size fractions as the significant contributors to growth and new
production during upwelling. Large phytoplankton consistently have greater increases in biomass
and outcompete small cells for nutrients during a bloom (Wilkerson et al., 2000; Fawcett and Ward,
2011). These large phytoplankton are commonly characterized as chain-forming colonial diatoms
such as *Chaetoceros* spp. but include *Pseudo-nitzschia* spp. (Estrada and Blasco, 1985; Lassiter et
al., 2006).

The large phytoplankton community also exhibited clear physiological responses to being
upwelled. Maximum photochemical yields (F_v:F_m) of the whole community increased from 0.25
to 0.51 within the first 72 hours (Supporting Information Fig. S1C). Dissolved inorganic carbon
and nitrate uptake in the large cells increased throughout the experiment and was significantly
higher than the small cells (Fig. 1B and Fig. 1C). The particulate carbon-to-nitrogen ratio (C:N)
was initially 31.5:1 in the >5 µm size fraction but decreased to approach the expected elemental
composition of 6.6:1 (Redfield et al., 1963), while C:N values remained fairly constant and above
the Redfield ratio for the small size fraction (Fig. 1D). This return to Redfield stoichiometry for
the larger community was coupled with increasing biomass-specific NO_3^- uptake rates (V_{NO3}) that
were approximately double that of biomass-specific carbon uptake rates (V_{DIC}) (Fig. 1E and Fig.
1F).

These data are also a clear indication of a positive response from light limitation, or
potentially a resting stage, to high growth for the larger phytoplankton. The initial low F_v:F_m
signifies that the community was stressed but quickly able to return to higher photosynthetic
efficiencies. A high initial C:N ratio that approaches the Redfield-predicted value has also been
observed in similar mesocosm experiments (Kudela and Dugdale, 2000; Fawcett and Ward, 2011).
These studies suggest that the initial high C:N ratio indicates severe N limitation which likely
occurred as the phytoplankton cells in aged upwelling water began to sink to depth. Once released
from light limitation, the community is able to stabilize with large phytoplankton controlling the
total C:N as time progresses. It is also possible that there was C-rich detrital material elevating the
initial measurement (Fawcett and Ward, 2011), but acceleration of nitrate uptake, especially in
relation to carbon uptake, to drive the community toward the Redfield-predicated ratio is clear.
The larger cells are able to take advantage of nitrate as conditions become optimal and dominate
the community since they uptake nitrate at higher rates than the smaller cells.

Taxonomic Composition

Metatranscriptome assembly resulted in 3.1 million contigs with levels of annotation similar to previous studies utilizing KEGG and reference transcriptomes from the Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP; Supporting Information Table S1)(Keeling et al., 2014; Alexander et al., 2015b; Cohen et al., 2017). Obtaining taxonomically-annotated mRNA read counts allows for inference of relative taxonomic composition which is supported by microscopic cell counts and observations from an imaging flow cytometer (FlowCAM; Fig. 2A and Supporting Information Table S2). The initial community biomass and cell abundance were relatively low but quite diverse (Fig. 1A and Supporting Information Table S2). Relative abundances of transcripts suggests that dinoflagellates were the dominant phytoplankton group although microscopic cell counts suggest chlorophytes may have been more abundant. Satellite-derived sea surface temperature and ship-board wind data indicate that upwelling-favorable conditions were not present for 13 days before the incubations (Supporting Information Fig. S2 and S3). Although multiple factors contribute to the residence time of cells at depth, these data suggest an upper limit of 13 days prior to sampling. By 72 and 120 hours following incubation, there was an overwhelming increase in the abundance of mRNA reads attributable to diatoms (Fig. 2A) consistent with the bulk measurements, FlowCAM, and previous studies (Estrada and Blasco, 1985): diatoms were unequivocally the dominant group within the simulated upwelling event.

The taxonomic composition of diatoms followed a similar trend as the whole community with an initially more diverse diatom community that transitioned into one dominated by just two genera: *Chaetoceros* and *Pseudo-nitzschia* (Fig. 2B). *Chaetoceros* appeared to make rapid early gains but the community became mostly *Pseudo-nitzschia* by 120 hours. These two genera were also dominant within a previous mesocosm experiment examining shift-up at a nearby coastal California site (Kudela and Dugdale, 2000). *Chaetoceros* spp. were found as resting spores and may quickly germinate following upwelling to make early gains in cell abundance (Pitcher, 1990). Although a resting stage for *Pseudo-nitzschia* spp. is not known (Lelong et al., 2012), they are significant members of the phytoplankton community throughout the upwelling cycle and dominated after 120 hours consistent with a peak in the *Pseudo-nitzschia* produced toxin, domoic acid (Supporting Information Fig. S1B). The presence of *Pseudo-nitzschia* is unsurprising considering the reports that members of this genus often dominate subsurface chlorophyll maxima

(Ryan et al., 2005), thin layers (Rines et al., 2002; McManus et al., 2008), and upwelled

communities (Seegers et al., 2015) that often result in harmful algal blooms in this region.

Comparative Gene Expression of Phytoplankton Groups

Examining shifts in the total transcript pool provides a broad depiction of the responsiveness of

different groups. By comparing expression levels at 0 and 72 hours, these shifts reveal the initial

whole transcriptome responses to simulated upwelling by the main detected groups of

phytoplankton (Fig. 3A). Diatoms had a high proportion of overrepresented genes after upwelling

compared to other groups, over 950 (20%) of which were significantly overrepresented (*P*-value

< 0.05). Dinoflagellates showed an opposite pattern with gene expression skewed towards

overrepresentation in the pre-upwelled condition while haptophytes had an even distribution of

overrepresented genes in both phases. Interestingly, chlorophytes also had a higher number of

significantly overrepresented genes post-upwelling, and they were able to maintain their relative

proportion of the overall transcript pool unlike the dinoflagellates and haptophytes.

Smaller changes across all four groups were observed when examining shifts from 72 to

120 hours (Supporting Information Fig. S4). Relatively minor changes in the whole transcript pool

and a less pronounced taxonomic shift from 72 to 120 hours indicates that most of the activity in

relation to diatom dominance likely occurred in the first 72 hours. This timing and slowing of

response also corresponds to field observations that predict a 5-7 day window for cells to achieve

balanced growth and transition from shift-up to a low nutrient shift-down (Dugdale and Wilkerson,

1989; Wilkerson et al., 2006). It has been speculated that these shifts, or variable transcript

allocation, are a reflection of r- and K-type growth strategies (Alexander et al., 2015b). Our

observations appear to follow this paradigm with diatoms exhibiting r-type growth and the highest

transcript reallocation in terms of gene count.

Analysis of the expression of genes with shared KEGG Orthology (KO) annotation allows

for direct comparisons between taxonomic groups as orthologs normally retain the same function

throughout evolutionary history. Similar or different expression of a gene among groups may

signify correspondingly similar or different investments in cellular processes at given time points.

We detected 1,476 orthologous genes expressed by all four taxonomic groups at 0 or 72 hours

(Fig. 3B and Fig. 3C). Only 18 genes were binned as highly overrepresented at 72 hours by all

four phytoplankton groups, of which many were related to chlorophyll synthesis. Over 550 genes

had low absolute fold change values, many of them positive, across all four groups. These included more photosynthesis-related genes such as photosystem II constituents, photosynthesis electron transport proteins, light-harvesting chlorophyll protein complex proteins, and most of the genes associated with the Calvin cycle. The shared expression of these genes across groups is unsurprising considering the community is transitioning from a deep and dark environment to a sunlit environment, and would benefit from investing in photosynthetic machinery. Other genes that were highly expressed but showed little change in expression across all four groups were associated with other predictable cellular functions such as ribosomal proteins, translation initiation factors, and all constituents of the citric acid cycle.

Of particular interest is the clear overrepresentation at 72 hours of approximately 200 genes per taxonomic group that show little or negative fold change in the other three groups (Fig. 3C). It is important to note that although differences in shifts in the total transcript pool were observed (Fig, 3A), all groups are still responding and highly increasing their expression of a distinct set of genes compared to the other groups. This pattern continues to hold when examining the genes that were shared between diatoms and just one or two of the other groups (Supporting Information Fig. S5). The genes highly expressed by each group appear to be of diverse function as they do not cluster into certain categories or modules but can be broadly interpreted as investments in different metabolic processes (Supporting Information Fig. S6). These unique responses may reflect fundamental differences in life strategies and ecological traits among functional groups.

To further explain the dominance of diatoms in these systems, expression of diatom annotated genes was investigated. 1,132 KOs were found solely in diatoms, likely due to the abundance of diatoms in our samples resulting in an improved metatranscriptome assembly for that group (Fig. 3B). However, only 173 of these KOs were significantly overrepresented at either 0 or 72 h. It is difficult to determine the importance of the remaining genes that were expressed in low abundances.

Diatom taxa, however, were not found to respond equivalently to being upwelled; clear differences were noted between *Chaetoceros*, *Pseudo-nitzschia*, and other diatoms (Fig. 4a). Expression of 2,807 orthologs was detected in the genera *Chaetoceros*, *Pseudo-nitzschia*, and all other diatom genera combined mostly consisting of *Thalassiosira*. Similar to what was observed for major taxonomic groups, there was large overrepresentation of distinct sets of genes,

particularly in *Chaetoceros* spp., also potentially reflecting transcriptional investments in different
processes at different times.

Eighty-five genes showed significantly opposite fold-changes in *Chaetoceros* and *Pseudo-*
nitzschia when compared to other diatoms. This highlights that gene expression may not be as
accurately assessed by combining genes at high level taxonomic groupings as done in previous
metatranscriptomics studies (Alexander et al., 2015b; Bertrand et al., 2015). The high expression
of a gene at one time point or treatment by one group may be cancelled out by another group with
opposing expression leading to the incorrect conclusion for the group as a whole. Additionally,
one genus could be driving expression of many genes rather than being distributed across the entire
group.

Molecular Characterization of the Nitrogen Assimilation Response
Gene expression was assessed among specific diatom genera and other phytoplankton groups to
investigate nitrogen assimilation and utilization. Querying nitrogen-related genes for these groups
and partitioning by k-means clustering revealed differences in gene expression for the diatoms
compared to other phytoplankton (Fig. 4B, Supporting Information Fig. S7). The genes that
clustered as highly expressed both pre- and post-simulated upwelling, referred to here as
frontloaded, were almost all from diatoms and related to nitrogen assimilation: nitrate transporter,
nitrate reductase, nitrite reductase, and ammonium transporter (Fig. 4C). The only constitutively
and highly expressed nitrogen-related gene by other groups was glutamine synthetase within
dinoflagellates and haptophytes. Within diatoms, the nitrate assimilation genes all had a positive
fold change suggesting slightly greater abundance of these genes post-upwelling when compared
to pre-upwelling matching our observations of increased nitrate uptake at 72 and 120 hours (Fig.
1C and Fig. 1F). The change in expression in nitrate reductase was very low which contrasts a
simulated upwelling experiment with a *Skeletonema* species (Smith et al., 1992). *Skeletonema*,
however, was not found to be an abundant genus within this study, and this variation further
highlights potential genera-specific differences in the upwelling response.

The urea cycle is believed to facilitate recovery from prolonged nitrogen limitation for
diatoms (Allen et al., 2011), but may also be important for the shift-up response. The urea cycle
genes carbamoyl-phosphate synthetase and argininosuccinate synthase were also frontloaded by
diatoms (Fig. 4B, Fig. 4C, and Supporting Information Fig. S7). Several others were significantly

overrepresented post-upwelling including ornithine carbamoyltransferase, arginosuccinate lyase,
and urease. The exception in diatoms was arginase, the final enzyme in the urea cycle which was
significantly overrepresented pre-upwelling by diatoms not including *Chaetoceros* (Supporting
Information Fig. S7). Low expression of arginase post-upwelling is similar to the diatom response
to iron enrichment (Marchetti et al., 2012) and may suggest that in both of these scenarios, there
are alternative fates for urea cycle intermediates such as nitrogen storage or silica precipitation
(Kröger et al., 2001; Llácer et al., 2008).

High relative expression of almost all of these nitrogen-related genes in diatoms compared
to most of the other phytoplankton groups at both 0 and 72 hours suggests that investing in nitrogen
assimilation and utilization is a priority even when conditions are not optimal for growth.
Constitutively high expression of the primary genes for nitrate assimilation such as nitrate
transporters and nitrate reductase pre-upwelling may contribute to the rapid response of diatoms
as part of their shift-up process. By maintaining elevated pools of nitrogen-related gene transcripts
or expressed proteins, upwelled cells are set up to rapidly assimilate available nitrogen whereas
other phytoplankton groups appear to wait to upregulate these genes once upwelled into the
euphotic zone. These results further support the hypothesis that one reason diatoms dominate
upwelling regions is because they have the ability to take up and assimilate nitrate more quickly
than other phytoplankton groups (Fawcett and Ward, 2011).

This transcriptionally proactive approach to abiotic changes, termed 'frontloading', has
been characterized with environmental stress response genes in coral and yeast (Berry and Gasch,
2008; Barshis et al., 2013). It is also similar to what has been observed in diatoms within a previous
metatranscriptomic study in relation to iron stress. Iron-enrichment experiments in the northeastern
Pacific Ocean demonstrated oceanic diatoms continued expressing genes encoding for iron-free
photosynthetic proteins rather than substituting genes encoding for iron-containing functionally
equivalent proteins which was in contrast to other phytoplankton groups (Marchetti et al., 2012;
Cohen et al., 2017). This strategy is speculated to provide oceanic diatoms with the ability to
rapidly acclimate to the inevitable return to iron-limited conditions just as our observations show
a strategy that provides certain diatoms with the ability to rapidly take up nitrogen following
upwelling. Constitutive frontloading is suggested to provide organisms with resilience to such
stressors (Barshis et al., 2013). Along similar lines, bloom-forming diatoms such as *Chaetoceros*
and *Pseudo-nitzschia* may have evolved to frontload transcripts of particular genes depending on

frequently encountered environmental fluctuations such as conditions associated with the
upwelling conveyor belt cycle rather than simply reactively responding to these changes.
297

Chaetoceros and Pseudo-nitzschia Expressed Genes
Analyzing genes assigned to two of the most dominant diatom genera, *Chaetoceros* and *Pseudo-*
nitzschia, provides further insight into the molecular mechanisms these genera use at depth and as
part of their shift-up response. From KOs with module annotations, it is evident that the
significantly overrepresented genes at both time points fall into a diverse set of functional
categories even at a high-level grouping (Supporting Information Fig. S8). To obtain finer
resolution, gene expression among all KOs for these genera was examined.

Many genes overrepresented in the pre-upwelling condition, such as those involved in
proteolysis, stabilization of DNA or RNA, defense against reactive oxygen species (superoxide
dismutases)(Fridovich, 1998), and TEP production, suggest that diatoms are responding to stress
(Fig. 4D and Supporting Information). *Pseudo-nitzchia* in particular expressed a set of distinctive
genes as part of its shift-up response. Ferritin was highly expressed post-upwelling, possibly
providing a method of storing the essential micronutrient iron. As iron availability in the California
upwelling regime can be sporadic and potentially growth limiting, ferritin may provide an
advantage to *Pseudo-nitzschia* by concentrating iron for longer-term storage (Bruland et al., 2001;
Marchetti et al., 2009) although it may also be used for iron homeostasis (Pfaffen et al., 2015).

At 72 hours, *Pseudo-nitzschia* also highly expressed a subunit of soluble guanylate cyclase
(GUCY1B, sGC, Fig. 4D). sGC is the only proven receptor of nitric oxide (Denninger and
Marletta, 1999) and synthesizes cyclic guanosine monophostphate (cGMP), a second messenger
related to many physiological responses (Delledonne, 2005). cGMP activates protein kinase G
(PRKG1) which was also significantly expressed (Fig. 4D). Although nitric oxide has been
hypothesized to be an infochemical for intercellular signaling and monitoring of stress in diatoms
(Vardi, 2008; Amin et al., 2012), *Pseudo-nitzschia* are generally not believed to have a nitric oxide
synthase gene as a putative sequence was detected in only one species, *P. multistriata* (Di Dato et
al., 2015). *Pseudo-nitzschia* may be using sGC to monitor exhibition of stress from other genera
which could allow them to rapidly adapt to changing conditions or respond to sexual cues (Basu
et al., 2017). Nitric oxide is also produced by activation of nitrate reductase (Sakihama et al.,
2002). As increased nitrate reductase activity occurs as part of the shift-up response, sGC may be

used to monitor the continuation of that response and promote certain cellular functions such as gliding of pennates or binary fission (Thompson et al., 2008). Inhibition of sGC prevents the germination of *Leptocyclindrus danicus* resting spores suggesting that it may be involved in transitioning from a resting stage in certain diatoms (Shikata et al., 2011).

Examination of this gene in our reference database reveals that it is highly conserved among *Pseudo-nitzschia* spp. but not ubiquitously present among diatoms (Supporting Information Fig. S9). The high expression of sGC as part of the upwelling response, evolutionary conservation of this gene, and potential to act as an important signaling device among *Pseudo-nitzschia* spp. may play a role in their success. Additionally, uncharacterized genes that have been previously observed to also increase in expression in comparable microarray studies (Ashworth et al., 2013; Nymark et al., 2013) may elucidate additional mechanisms for the shift-up response in *Chaetoceros* and *Pseudo-nitzschia* (Supporting Information Data Set S1).

Conclusions

Our simulated upwelling experiment in the California Upwelling Zone is consistent with previous physiological observations of the shift-up response in upwelled phytoplankton: growth of large chain-forming diatoms and increased nitrate assimilation rates. The application of metatranscriptomics to the entire phytoplankton community highlights the divergent transcriptional response of major phytoplankton groups and diatom genera, potentially reflecting variations in their life history strategies. By frontloading, diatoms exhibit the potential for abundant nitrogen-related genes in their enzyme pool allowing them to respond to available nitrate more rapidly than other phytoplankton. This trait is not unlike the response of oceanic diatoms to iron enrichment and may indicate that diatoms have evolved to frontload transcripts in response to frequently encountered changes in their environment. Although the response to upwelling has largely been focused on nitrogen-related pathways, it is likely that other uncharacterized genes and pathways are also important to diatom success.

Experimental procedures

Sample Collection

On 17 July 2014, upwelling conditions were not present at a site within the California Upwelling Zone (35° 56.071' N, 121° 44.022' W; Supporting Information Fig. S2 and Fig. S3). At 05:00 PDT

(12:00 GMT) at the same location, viable phytoplankton cells were detected via imaging flow

cytometry (FlowCAM, Fluid Imaging Technologies Inc., Scarborough, ME, USA) at 96 m which

corresponded to the 10°C isotherm (Supporting Information Fig. S10). Seawater from this depth

was processed immediately for the initial time point. To simulate upwelling, additional seawater

from the same depth was filled into a large acid-rinsed HDPE barrel for homogenization, dispensed

to triplicate 10L Cubitainer® (Hedwin Corporation, Newark, DE, USA), and incubated in an on-

deck plexiglass incubator with flow-through seawater at 33% incident irradiance. Seawater

collection and incubations followed trace metal clean techniques as they were conducted as part

of a larger study to examine diatom responses to iron addition or removal (R. H. Lampe,

unpublished) although for the purposes of this study, only the unamended control incubations are

considered. Temperature and on deck irradiance values throughout the incubation are provided in

Supporting Information Fig. S11.

Based on macronutrient drawdown, triplicate cubitainer were harvested following 72 hours

and 120 hours of incubation. Subsamples from each cubitainer were preserved or measured for

chlorophyll *a*, species composition (by microscopy and FlowCAM), biogenic silica, $F_v:F_m$, domoic

acid, nutrients, particulate carbon and nitrogen, carbon and nitrogen uptake, and RNA. Chlorophyll

a, particulate carbon and nitrogen, and nitrate uptake rates were size fractionated using a series

filter cascade. Caron uptake rates were size fractionated using a mesh spacer. Additional methods

are described in the Supporting Information.

Chlorophyll

Four hundred mL of seawater was gravity-filtered through a 5 µm polycarbonate filter (47 mm)

followed by a GF/F filter (25 mm) under gentle vacuum pressure (<100 mm Hg). Filters were

rinsed with 0.45 µm filtered seawater and immediately frozen at -80°C until analysis. Chlorophyll

a extraction was performed using 90% acetone at -20°C for 24 h and measured via *in vitro*

fluorometry on a 10-AU fluorometer (Turner Designs, San Jose, CA, USA) using the acidification

method (Parsons et al., 1984).

Biogenic silica

Biogenic silica was determined via filtration of 335 mL onto 1.2 µm polycarbonate filters (45 mm). Concentrations were measuring using a NaOH digestion in teflon tubes (Krause et al., 2009) and a colorimetric ammonium molybdate method (Brzezinski and Nelson, 1995).

Particulate Carbon, Particulate Nitrogen, and Nitrate Uptake

Particulate nitrogen (PN), carbon (PC) and nitrate (NO_3^-) uptake was obtained by spiking 618 mL of seawater with ^{15}N-$NaNO_3$ at no more than 10% of ambient nitrate concentration and incubating for 8 h in the flow-through plexiglass incubator. Following incubation, seawater filtration commenced immediately and was performed by gravity through a 5 µm polycarbonate filter (47 mm), and with an in-line vacuum (<100 mm Hg) onto a precombusted (450°C for 5 h) GF/F filter (25 mm). Cells on the 5 µm polycarbonate filter were then rinsed onto an additional precombusted GF/F filter (25 mm) using an artificial saline solution. Filters were then stored at -20°C. Prior to analysis, filters were dried at 50°C for 24 h, wrapped in tin capsules, and pelletized in preparation for analysis of the atom % ^{15}N, PN, and PC using an elemental analyzer paired with an isotope ratio mass spectrometer (EA-IRMS). Absolute uptake rates (ρ, NO_3 taken up per unit time) were calculated using a constant transport model (Eq. (3) from Dugdale and Wilkerson (1986)). Biomass-specific NO_3 uptake rates (V, NO_3 taken up per unit PN per unit time) were also calculated according to the constant specific uptake mode (Eq. (6) Dugdale and Wilkerson (1986)). $^{15}NO_3$ uptake rates were not corrected for possible losses of ^{15}N in the form of dissolved organic nitrogen (Bronk et al., 1994); therefore, the reported values are considered conservative estimates or net uptake.

Dissolved Inorganic Carbon Uptake

Sixty mL samples from each cubitainer were distributed into light and dark bottles cleaned with 1.2 mol L^{-1} HCl. For each bottle, 1.2 µCi of $NaH^{14}CO_3$ was added and mixed. A 1 mL subsample was taken and added to vials containing 6 mol L^{-1} NaOH to trap and validate the initial inorganic $H^{14}CO_3$ quantities. The light and dark bottles were incubated on-deck for 6.5-8 h. Samples were filtered onto stacked polycarbonate filters (5 µm and 1 µm) separated with a mesh spacer. Blank control bottles also containing 1.2 µCi of $NaH^{14}CO_3$ were filtered onto a GF/F filter after 5 min and had counts similar to dark bottles. Filters were vacuumed dried, placed in scintillation vials

with 0.5 mL of 6 mol L^{-1} HCl, permitted to degas for 24 h, and counted using a Beckman Coulter

LS 6500 scintillation counter. Reported values are light bottles minus dark bottles. Biomass-

specific dissolved inorganic carbon (DIC) uptake rates (V_{DIC}) were calculated by normalizing DIC

uptake to PC.

RNA Extraction and Sequencing

Seawater was filtered onto 0.8 µm Pall Supor filters (142 mm) using a peristaltic pump then

immediately flash frozen in liquid nitrogen. RNA was extracted using the ToTALLY RNA Total

RNA Isolation Kit and treated with DNase 1 (Life Technologies, Grand Island, NY, USA). The

extraction procedure was followed according to the manufacturer's instructions with additional

first step of glass bead addition to assist with organic matter disruption. RNA quantity and purity

was assessed prior to sequencing on an Agilent Bioanalyzer 2100. Total RNA from the triplicate

samples for the initial time point (T0) and the first time point (T72) were pooled into one sample

due to low RNA yields. Triplicate samples were maintained for the second time point (T120).

Library prep was conducted with the Illumina TruSeq Stranded mRNA Library Preparation Kit

and HiSeq v4 reagents. Sequencing of barcoded samples was performed on an Illumina HiSeq

2000 (125bp, paired-end).

Metatranscriptome assembly, annotation, and read quantification

Reads were trimmed for quality and adapter removal using Trimmomatic v0.32 (paired-end mode,

adaptive quality trim with 40 bp target length and strictness of 0.6, minimum length of 36

bp)(Bolger et al., 2014). Trimmed paired reads that overlap were merged into single reads with

BBMerge v8.0. Merged pairs and non-overlapping paired-end reads were then used to assemble

contigs using ABySS v1.5.2 with varied k-mer sizes (Birol et al., 2009). Assemblies were merged

using Trans-ABySS v1.5.3 to remove redundant contigs (Robertson et al., 2010), and those shorter

than 125 bp were discarded. Read counts were obtained by mapping raw reads to contigs with

Bowtie2 v2.2.6 (Langmead and Salzberg, 2012) and filtered by mapping quality with SAMtools

v1.2 (Li et al., 2009). Mapping percentages are provided in the Supporting Information Table S3.

Annotation was assigned by best homology (lowest E-value) to protein databases using

BLASTX v2.2.31 (E-value $\leq 10^{-5}$). For taxonomic identification, MarineRefII, a custom reference

database was used. MarineRefII contains predicted protein sequences of marine microbial

eukaryotes and prokaroytes including all sequenced transcriptomes from the Marine Microbial

Eukaryote Transcriptome Sequencing Project (Keeling et al., 2014). MarineRefII was

supplemented with transcriptomes of isolated phytoplankton from these incubations adding

increased confidence in the taxonomic annotation of some contigs (Supporting Information Table

S4). To assign gene function to contigs, the same methodology with the Kyoto Encyclopedia of

Genes and Genomes (KEGG; Release 75) was used (Kanehisa et al., 2017). The best hit with a

KEGG Ortholog (KO) number from the top 10 hits was chosen. Similarly, analysis of module

annotations (MO) was conducted by selecting the top BLASTX hit with a KEGG MO number

from the top 10 hits. A summary of annotation results is provided in Supporting Information Table

S2.

Differential expression analysis

Differential expression was assessed by summing read counts of contigs within a taxonomic group

(phylum-based or genus for only the diatoms, *Chaetoceros* and *Pseudo-nitzschia*) by KEGG Gene

Definition or KEGG Orthology (KO) annotation. edgeR v3.12.0 was used to calculate normalized

fold change and counts-per-million (CPM) from pairwise comparisons within each taxonomic

group using the exactTest function (Robinson and Smyth, 2008; Robinson et al., 2010).

Significance (P-value < 0.05) was calculated by using edgeR's estimate of tagwise dispersions

utilizing the available replication of all treatments from the second time point (T120; Supporting

Information Fig. S12)(Chen et al., 2014). ExactTest output in combination with the taxonomic

distributions per gene were plotted using a custom plotting function available at

https://github.com/marchettilab/mantaPlot.

Shared expression of gene was considered when a gene was detected in at least one of the

libraries under comparison for each taxonomic group. For binning of genes displayed in heatmaps

(Fig. 3C and Fig. 4A), a positive or negative fold change, variance greater than the number of

taxonomic groups, and fold change greater than or less than all other groups were used. Genes

with a $\log_2$ fold change greater than 2 or less than -2 but had a variance less than the number of

taxonomic groups were considered similarly overrepresented by all groups. Otherwise, the

expression level was considered similar on the basis of fold change. These data were visualized

with pheatmap v1.0.8.

Acknowledgements

We thank the captain and crew of the R/V Melville and the participants of the IRNBRU cruise: T.

Coale (UCSD) performed dissolved nutrient measurements, H. McNair (UCSB) and J. Jones

(UCSB) assisted with biogenic silica measurements, M. Maier (OHSU) acquired FlowCAM data,

and C. Duckham (UBC) assisted with primary productivity measurements. We also thank W.

Burns (UNC), K. Delong (UNC), and C. Payne (UBC) for assistance with sample analyses as well

as S. Davies (UNC), W. Gong (UNC), and S. Haines (UNC) for bioinformatic assistance. This

work was funded by the National Science Foundation OCE-1334935 (to A.M.), OCE-1334632 (to

B.S.T.), OCE-1333929 (to K.T.), OCE-1334387 (to M.A.B.), OCE-1259776 (to K.W.B.).

Availability of the isolate transcriptomes used in this study was made possible by funding from

National Science Foundation OCE-1341479 (to A.M.). R.H.L was partially supported by a

fellowship from the UNC Graduate School.

Conflict of Interest

The authors declare no conflict of interest.

References

Alexander, H., Jenkins, B.D., Rynearson, T.A., and Dyhrman, S.T. (2015a) Metatranscriptome
analyses indicate resource partitioning between diatoms in the field. *P Natl Acad Sci USA*
112: E2182-E2190.
Alexander, H., Rouco, M., Haley, S.T., Wilson, S.T., Karl, D.M., and Dyhrman, S.T. (2015b)
Functional group-specific traits drive phytoplankton dynamics in the oligotrophic ocean. *P*
Natl Acad Sci USA **112**: E5972-E5979.
Allen, A.E., Dupont, C.L., Obornik, M., Horak, A., Nunes-Nesi, A., McCrow, J.P. et al. (2011)
Evolution and metabolic significance of the urea cycle in photosynthetic diatoms. *Nature*
473: 203-207.
Amin, S.A., Parker, M.S., and Armbrust, E.V. (2012) Interactions between Diatoms and
Bacteria. *Microbiol Mol Biol R* **76**: 667-684.
Ashworth, J., Coesel, S., Lee, A., Armbrust, E.V., Orellana, M.V., and Baliga, N.S. (2013)
Genome-wide diel growth state transitions in the diatom Thalassiosira pseudonana. *P Natl*
Acad Sci USA **110**: 7518-7523.
Barshis, D.J., Ladner, J.T., Oliver, T.A., Seneca, F.O., Traylor-Knowles, N., and Palumbi, S.R.
(2013) Genomic basis for coral resilience to climate change. *P Natl Acad Sci USA* **110**: 1387-
1392.
Basu, S., Patil, S., Mapleson, D., Russo, M.T., Vitale, L., Fevola, C. et al. (2017) Finding a
partner in the ocean: molecular and evolutionary bases of the response to sexual cues in a
planktonic diatom. *New Phytologist* **215**: 140-156.
Berry, D.B., and Gasch, A.P. (2008) Stress-activated Genomic Expression Changes Serve a
Preparative Role for Impending Stress in Yeast. *Mol Biol Cell* **19**: 4580-4587.
Bertrand, E.M., McCrow, J.P., Moustafa, A., Zheng, H., McQuaid, J.B., Delmont, T.O. et al.
(2015) Phytoplankton–bacterial interactions mediate micronutrient colimitation at the coastal
Antarctic sea ice edge. *P Natl Acad Sci USA* **112**: 9938-9943.
Birol, I., Jackman, S.D., Nielsen, C.B., Qian, J.Q., Varhol, R., Stazyk, G. et al. (2009) De novo
transcriptome assembly with ABySS. *Bioinformatics* **25**: 2872-2877.
Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: A flexible trimmer for Illumina
Sequence Data. *Bioinformatics* **30**: 2114-2120.
Bronk, D.A., Glibert, P.M., and Ward, B.B. (1994) Nitrogen Uptake, Dissolved Organic
Nitrogen Release, and New Production. *Science* **265**: 1843-1846.
Bruland, K.W., Rue, E.L., and Smith, G.J. (2001) Iron and macronutrients in California coastal
upwelling regimes: Implications for diatom blooms. *Limnol Oceanogr* **46**: 1661-1674.
Brzezinski, M.A., and Nelson, D.M. (1995) The annual silica cycle in the Sargasso Sea near
Bermuda. *Deep-Sea Res Pt I* **42**: 1215-1237.
Capone, D.G., and Hutchins, D.A. (2013) Microbial biogeochemistry of coastal upwelling
regimes in a changing ocean. *Nature Geosci* **6**: 711-717.
Caron, D.A., Alexander, H., Allen, A.E., Archibald, J.M., Armbrust, E.V., Bachy, C. et al.
(2017) Probing the evolution, ecology and physiology of marine protists using
transcriptomics. *Nat Rev Micro* **15**: 6-20.
Chen, Y., Lun, A., and Smyth, G. (2014) Differential expression analysis of complex RNA-seq
experiments using edgeR. In *Statistical Analysis of Next Generation Sequence Data*. Datta,
S., and Nettleton, D.S. (eds). New York: Springer.

Cohen, N.R., Ellis, K.A., Lampe, R.H., McNair, H.M., Twining, B.S., Brzezinski, M.A. et al.
(2017) Variations in diatom transcriptional responses to changes in iron availability across
ocean provinces. *Front Mar Sci* **4**: 360.
Delledonne, M. (2005) NO news is good news for plants. *Curr Opin Plant Biol* **8**: 390-396.
Denninger, J.W., and Marletta, M.A. (1999) Guanylate cyclase and the ·NO/cGMP signaling
pathway. *BBA-Bioenergetics* **1411**: 334-350.
Di Dato, V., Musacchia, F., Petrosino, G., Patil, S., Montresor, M., Sanges, R., and Ferrante, M.I.
(2015) Transcriptome sequencing of three Pseudo-nitzschia species reveals comparable gene
sets and the presence of Nitric Oxide Synthase genes in diatoms. *Sci Rep* **5**: 12329.
Dugdale, R.C., and Wilkerson, F.P. (1986) The use of 15N to measure nitrogen uptake in
eutrophic oceans; experimental considerations. *Limnol Oceanogr* **31**: 673-689.
Dugdale, R.C., and Wilkerson, F.P. (1989) New production in the upwelling center at Point
Conception, California: temporal and spatial patterns. *Deep-Sea Res* **36**: 985-1007.
Estrada, M., and Blasco, D. (1985) Phytoplankton assemblages in coastal upwelling areas. In
International Symposium on the Most Important Upwelling Areas off Western Africa. Bas,
C., Margalef, R., and Rubies, P. (eds). Barcelona, pp. 379-402.
Fawcett, S., and Ward, B. (2011) Phytoplankton succession and nitrogen utilization during the
development of an upwelling bloom. *Mar Ecol-Prog Ser* **428**: 13-31.
Fridovich, I. (1998) Oxygen toxicity: a radical explanation. *J Exp Biol* **201**: 1203-1209.
Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., and Morishima, K. (2017) KEGG: new
perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**: D353-D361.
Keeling, P.J., Burki, F., Wilcox, H.M., Allam, B., Allen, E.E., Amaral-Zettler, L.A. et al. (2014)
The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP):
Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through
Transcriptome Sequencing. *PLoS Biol* **12**: e1001889.
Krause, J.W., Nelson, D.M., and Lomas, M.W. (2009) Biogeochemical responses to late-winter
storms in the Sargasso Sea, II: Increased rates of biogenic silica production and export. *Deep-*
Sea Res Pt I **56**: 861-874.
Kröger, N., Deutzmann, R., and Sumper, M. (2001) Silica-precipitating Peptides from Diatoms:
The Chemical Structure of Silaffin-1A from Cylindrotheca Fusiformis. *J Biol Chem* **276**:
26066-26070.
Kudela, R.M., and Dugdale, R.C. (2000) Nutrient regulation of phytoplankton productivity in
Monterey Bay, California. *Deep-Sea Res Pt II* **47**: 1023-1053.
Lachkar, Z., and Gruber, N. (2013) Response of biological production and air–sea CO2 fluxes to
upwelling intensification in the California and Canary Current Systems. *J Marine Syst* **109–**
110: 149-160.
Langmead, B., and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Meth* **9**:
357-359.
Lassiter, A.M., Wilkerson, F.P., Dugdale, R.C., and Hogue, V.E. (2006) Phytoplankton
assemblages in the CoOP-WEST coastal upwelling area. *Deep-Sea Res Pt II* **53**: 3063-3077.
Lelong, A., Hégaret, H., Soudant, P., and Bates, S.S. (2012) *Pseudo-nitzschia*
(Bacillariophyceae) species, domoic acid and amnesic shellfish poisoning: revisiting previous
paradigms. *Phycologia* **51**: 168-216.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N. et al. (2009) The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078-2079.

Llácer, J.L., Fita, I., and Rubio, V. (2008) Arginine and nitrogen storage. *Curr Opin Struct Biol* **18**: 673-681.

MacIsaac, J.J., Dugdale, R.C., Barber, R.T., Blasco, D., and Packard, T.T. (1985) Primary production cycle in an upwelling center. *Deep-Sea Res* **32**: 503-529.

Marchetti, A., Parker, M.S., Moccia, L.P., Lin, E.O., Arrieta, A.L., Ribalet, F. et al. (2009) Ferritin is used for iron storage in bloom-forming marine pennate diatoms. *Nature* **457**: 467-470.

Marchetti, A., Schruth, D.M., Durkin, C.A., Parker, M.S., Kodner, R.B., Berthiaume, C.T. et al. (2012) Comparative metatranscriptomics identifies molecular bases for the physiological responses of phytoplankton to varying iron availability. *P Natl Acad Sci USA* **109**: E317-E325.

McManus, M.A., Kudela, R.M., Silver, M.W., Steward, G.F., Donaghay, P.L., and Sullivan, J.M. (2008) Cryptic Blooms: Are Thin Layers the Missing Connection? *Estuar Coast* **31**: 396-401.

Nymark, M., Valle, K.C., Hancke, K., Winge, P., Andresen, K., Johnsen, G. et al. (2013) Molecular and Photosynthetic Responses to Prolonged Darkness and Subsequent Acclimation to Re-Illumination in the Diatom Phaeodactylum tricornutum. *PLoS ONE* **8**: e58722.

Parsons, T.R., Maita, Y., and Lalli, C.M. (1984) In *A Manual of Chemical & Biological Methods for Seawater Analysis*. Amsterdam: Pergamon.

Pfaffen, S., Bradley, J.M., Abdulqadir, R., Firme, M.R., Moore, G.R., Le Brun, N.E., and Murphy, M.E.P. (2015) A Diatom Ferritin Optimized for Iron Oxidation but not Iron Storage. *J Biol Chem*.

Pitcher, G.C. (1990) Phytoplankton seed populations of the Cape Peninsula upwelling plume, with particular reference to resting spores of Chaetoceros (bacillariophyceae) and their role in seeding upwelling waters. *Estuar Coast Shelf S* **31**: 283-301.

Redfield, A.C., Ketchum, B.H., and Richards, F.A. (1963) The influence of organisms on the composition of sea water. In *The Sea*. Hill, M. (ed). New York: Interscience, pp. 26-77.

Rines, J.E.B., Donaghay, P.L., Dekshenieks, M.M., Sullivan, J.M., and Twardowski, M.S. (2002) Thin layers and camouflage: hidden Pseudo-nitzschia spp. (Bacillariophyceae) populations in a fjord in the San Juan Islands, Washington, USA. *Mar Ecol-Prog Ser* **225**: 123-137.

Robertson, G., Schein, J., Chiu, R., Corbett, R., Field, M., Jackman, S.D. et al. (2010) De novo assembly and analysis of RNA-seq data. *Nat Meth* **7**: 909-912.

Robinson, M.D., and Smyth, G.K. (2008) Small-sample estimation of negative binomial dispersion, with applications to SAGE data. *Biostatistics* **9**: 321-332.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139-140.

Ryan, J.P., Chavez, F.P., and Bellingham, J.G. (2005) Physical-biological coupling in Monterey Bay, California: topographic influences on phytoplankton ecology. *Mar Ecol-Prog Ser* **287**: 23-32.

Ryther, J.H. (1969) Photosynthesis and Fish Production in the Sea. *Science* **166**: 72-76.

Sakihama, Y., Nakamura, S., and Yamasaki, H. (2002) Nitric Oxide Production Mediated by Nitrate Reductase in the Green Alga Chlamydomonas reinhardtii: an Alternative NO Production Pathway in Photosynthetic Organisms. *Plant Cell Physiol* **43**: 290-297.

Schnetzer, A., Jones, B.H., Schaffner, R.A., Cetinic, I., Fitzpatrick, E., Miller, P.E. et al. (2013)
Coastal upwelling linked to toxic Pseudo-nitzschia australis blooms in Los Angeles coastal
waters, 2005–2007. *J Plankton Res* **35**: 1080-1092.
Seegers, B.N., Birch, J.M., Marin, R., Scholin, C.A., Caron, D.A., Seubert, E.L. et al. (2015)
Subsurface seeding of surface harmful algal blooms observed through the integration of
autonomous gliders, moored environmental sample processors, and satellite remote sensing
in southern California. *Limnol Oceanogr* **60**: 754-764.
Shikata, T., Iseki, M., Matsunaga, S., Higashi, S.-i., Kamei, Y., and Watanabe, M. (2011) Blue
and Red Light-Induced Germination of Resting Spores in the Red-Tide Diatom
Leptocylindrus danicus†. *Photochem Photobiol* **87**: 590-597.
Smith, G.J., Zimmerman, R.C., and Alberte, R.S. (1992) Molecular and physiological responses
of diatoms to variable levels of irradiance and nitrogen availability: growth of Skeletonema
costatum in simulated upwelling conditions. *Limnol Oceanogr* **37**: 989-1007.
Thompson, S.E.M., Taylor, A.R., Brownlee, C., Callow, M.E., and Callow, J.A. (2008) The Role
of Nitric Oxdie in Diatom Adhesion in Relation to Substratum Properties. *J Phycol* **44**: 967-
976.
Vardi, A. (2008) Cell signaling in marine diatoms. *Commun Integr Biol* **1**: 134-136.
Wilkerson, F.P., and Dugdale, R.C. (1987) The Use of Large Shipboard Barrels and Drifters to
Study the Effects of Coastal Upwelling on Phytoplankton Dynamics. *Limnol Oceanogr* **32**:
368-382.
Wilkerson, F.P., and Dugdale, R.C. (2008) *Nitrogen in the Marine Environment (2nd Edition).*
Burlington, MA, USA: Academic Press.
Wilkerson, F.P., Dugdale, R.C., Kudela, R.M., and Chavez, F.P. (2000) Biomass and
productivity in Monterey Bay, California: contribution of the large phytoplankton. *Deep-Sea*
Res Pt II **47**: 1003-1022.
Wilkerson, F.P., Lassiter, A.M., Dugdale, R.C., Marchi, A., and Hogue, V.E. (2006) The
phytoplankton bloom response to wind events and upwelled nutrients during the CoOP
WEST study. *Deep-Sea Research Part II* **53**: 3023-3048.
668

 Figure Legends

Fig. 1. Measurements from the initial upwelled water and incubations at 72 and 120 hours for the
community >5 µm (red) and <5 µm (blue). (A) Chlorophyll *a* (closed circles), particulate nitrogen
(open circles), and biogenic silica (green). (B) Primary productivity expressed as inorganic carbon
uptake (µmol C L^{-1} d^{-1}). (C) Absolute nitrate (NO$_3^-$) uptake rates (ρ, NO$_3^-$ taken up per unit time).
(D) Ratios of particulate carbon to particulate nitrogen. (E) Biomass-specific dissolved inorganic
carbon uptake rates (V$_{DIC}$), i.e. inorganic carbon uptake normalized to biomass as particulate
carbon (µmol C L^{-1} d^{-1} / µmol C L^{-1} or d^{-1}) (F) Biomass-specific nitrate uptake rates (V$_{NO3}$), i.e.
nitrate uptake rates normalized to biomass as particulate nitrogen (µmol N L^{-1} d^{-1} / µmol N L^{-1} or
d^{-1}). Error bars indicate standard deviation of the mean (n = 3).

Fig. 2. Average taxonomic distribution by mapped reads from each time point. (A) Percentage of
reads from the whole community. (B) Percentage of reads for diatom genera within all reads
assigned as diatoms.

Fig. 3. KEGG Ortholog (KO) gene expression comparison among the four main detected
phytoplankton groups: diatoms (blue), dinoflagellates (red), chlorophytes (green), and haptophytes
(orange). (A) Histograms of KO counts binned by log$_2$ fold change intervals of 1 for 0 and 72
hours. Dashed vertical lines indicate a log$_2$ fold change of -1 or 1. The number and percentage of
only the significantly (*P*-value < 0.05) overrepresented genes at 0 hours (pre-upwelling; left) and
72 hours (post-upwelling; right) are annotated on each plot. (B) Venn diagram of expressed KOs
at 72 and 0 hours for each group. (C) Heatmap for the 1,476 commonly expressed KOs at 0 and
72 hours. Each row indicates an expressed KO with darker red (positive fold change) indicating
overrepresentation at 72 hours and darker blue (negative fold change) indicating
overrepresentation at 0 hours.

Fig. 4. Diatom gene expression. (A) Heatmap for the 2,807 commonly expressed KOs at 0 and 72
hours for *Chaetoceros*, *Pseudo-nitzschia*, and all other diatoms. Each row indicates an expressed
KO with darker red (positive fold change) indicating overrepresentation at 72 hours and darker
blue (negative fold change) indicating overrepresentation at 0 hours. (B) MA plot of nitrogen-
related genes for the four main detected phytoplankton groups: diatoms (blue), dinoflagellates
(red), chlorophytes (green), and haptophytes (orange). Genes are grouped into clusters using k-
means clustering with confidence ellipses at the 90% level. Select genes are labelled within clusters
that represent the most highly frontloaded genes (black ellipse) and additional frontloaded genes
(medium gray ellipse). These are labelled as follows: NRT, nitrate transporter; GLNA, glutamine
synthetase; NR, nitrate reductase; AMT, ammonium transporter; NIRA, nitrite reductase
(ferredoxin); ASSY, arginosuccinate synthase; CPS, carbamoyl-phosphate synthetase; OTC,
ornithine carbamoyltransferase; ASL, argininosuccinate lysase; URE, urease, ARG, arginase. (C)
Cell schematic depicting frontloading of genes associated with nitrogen assimilation and
utilization for diatoms. The model is based on Alexander et al. (2015a) and utilizes the same KO
numbers. Color indicates the average abundance of the genes (log$_2$ CPM) minus the absolute value
of the log$_2$ fold change to highlight the most abundant, lowest change (i.e., frontloaded) genes.
Labels are as indicated in Fig. 4B and as follows: GLT1, glutamate synthase (NADP/NADH);
GLT2, glutamate synthase (ferredoxin); GLTD, glutamate synthase; small chain. (D) Differential
transcript abundance between 0 and 72 hours for *Chaetoceros* (red) and *Pseudo-nitzschia* (blue)

for expressed KEGG Orthologs (KO). Each pie represents a KO and increases in size with absolute
values of its coordinates to optimize visibility. Gene circles that are shaded and have grey borders
are not significantly represented in either library (P-value ≥ 0.05). Select gene names discussed in
the text are labeled as follows: CHSY, chondroitin sulfate synthase; FTN, ferritin; GUCY1B,
guanylate cyclase soluble subunit beta; HERC1, ubiquitin-protein ligase; MRT4; mRNA turnover
protein, POT1, protection of telomeres protein 1; PRKG1, cGMP-dependent protein kinase 1;
SOD, superoxide dismutases.
721
722

Supporting Information

724 **Data Set S1.** Genes of unknown function significantly expressed in our study and with positive
725 fold-changes in a similar study on *Thalassiosira pseudonana* (tps) and *P. tricornutum* (pti;
726 Supporting Information).
727