Sex-specific changes in the aphid DNA methylation landscape

Thomas C. Mathers[1], Sam T. Mugford[1], Lawrence Percival-Alwyn[2, a], Yazhou Chen[1], Gemy Kaithakottil[2], David Swarbreck[2], Saskia A. Hogenhout[1, *] and Cock van Oosterhout[3, *]

[1]Department of Crop Genetics, John Innes Centre, Norwich Research Park, Norwich, United Kingdom

[2]Earlham Institute, Norwich Research Park, Norwich, United Kingdom

[2]School of Environmental Sciences, University of East Anglia, Norwich, United Kingdom

[a]Current address: The John Bingham Laboratory, NIAB, Huntingdon Road, Cambridge, United Kingdom

*Corresponding authors

E-mail: c.van-oosterhout@uea.ac.uk

E-mail: saskia.hogenhout@jic.ac.uk

Running title

Aphid DNA methylation

Keywords

Dosage compensation, epigenetic regulation, *Myzus persicae*, sex-biased gene expression, sex chromosome.

Abstract

Aphids present an ideal system to study epigenetics as they can produce diverse, but genetically identical, morphs in response to environmental stimuli. Here, using whole genome bisulphite sequencing and transcriptome sequencing of the green peach aphid (*Myzus persicae*), we present the first detailed analysis of cytosine methylation in an aphid and investigate differences in the methylation and transcriptional landscapes of male and asexual female morphs. We find that methylation primarily occurs in a CG dinucleotide (CpG) context and that exons are highly enriched for methylated CpGs, particularly at the 3' end of genes. Methylation is positively associated with gene expression, and methylated genes are more stably expressed than un-methylated genes. Male and asexual female morphs have distinct methylation profiles. Strikingly, these profiles are divergent between the sex chromosome and the autosomes; autosomal genes are hypo-methylated in males compared to asexual females, whereas genes belonging to the sex chromosome, which is haploid in males, are hyper-methylated. Overall, we find correlated changes in methylation and gene expression between males and asexual females, and this correlation is particularly strong for genes located on the sex chromosome. Our results suggest that differential methylation of sex-biased genes plays a role in *M. persicae* sexual differentiation.

Introduction

Cytosine methylation is an epigenetic mark found in many eukaryotic organisms (Bewick et al. 2016, 2017; Feng et al. 2010; Zemach and Zilberman 2010). In mammals, cytosine methylation mainly occurs in a CG dinucleotide context (CpG) (Suzuki and Bird 2008). However, in human embryonic stem cells (Guo et al. 2014), and human and mouse oocytes (Guo et al. 2014; Okae et al. 2014), cytosines are methylated in other sequence contexts (non-CpG). Plants also have high levels of non-CpG methylation that is maintained by a set of specialised CHROMOMETHYLASE enzymes not found in other eukaryotes (Bewick et al. 2017). CpG methylation is extensively detected throughout mammalian and plant genomes and often associated with suppression of gene, or transposable element, expression. In contrast, insect genomes have sparse cytosine methylation that is almost exclusively restricted to CpG sites in gene bodies (Zemach et al. 2010). Furthermore, rather than supressing gene expression, insect CpG methylation is associated with high and stable gene expression (Wang

et al. 2013; Patalano et al. 2015; Xiang et al. 2010; Libbrecht et al. 2016).

Social Hymenoptera have been used as a model system to study the function of insect DNA

methylation and its role caste determination (Yan et al. 2014). However, replicated

experimental designs have recently shown high between-sample variation (low repeatability)

and no evidence of statistically significant differences in CpG methylation between social

insect castes (Libbrecht et al. 2016). Furthermore, DNA methylation has a patchy distribution

across the insect phylogeny, having been lost in many species, and appears to be dispensable

for the evolution of sociality (Bewick et al. 2016). Development of additional model systems

is therefore desirable to gain a deeper understanding of the function of cytosine methylation

in insects.

Aphids have a functional DNA methylation system (Bewick et al. 2016; Hunt et al. 2010; Walsh

et al. 2010) and are an outgroup to holometabolous insects (Misof et al. 2014), which have

been the main focus of research into insect DNA methylation to date. Furthermore, aphids

display extraordinary phenotypic plasticity during their life cycle (Dixon 1977), in the absence

of confounding genetic variation, making them ideal for studying epigenetics (Srinivasan and

Brisson 2012). During the summer months, aphids are primarily found as alate, asexually

reproducing, females. These asexual females are able to produce morphologically distinct

morphs in response to environmental stimuli. This can include the induction of winged

individuals in response to crowding (Müller et al. 2001), or the production of sexually

reproducing forms in response to changes in temperature and day length (Blackman 1971b).

In the case of the production of sexually reproducing individuals, sex is determined by an XO

chromosomal system, where males are genetically identical to their mothers apart from the

random loss of one copy of the X chromosome (Wilson et al. 1997). Differences between

aphid morphs are known to be associated with large changes in gene expression (Jaquiéry et

al. 2013; Purandare et al. 2014), but whether or not changes in cytosine methylation are also

involved is unknown.

Here, we performed the first in-depth, genome-wide, analysis of aphid DNA methylation. We

find that asexual females and males have distinct expression and methylation profiles and

that changes in methylation differ between the X chromosome and autosomes. In males, the

autosomes are hypo-methylated relative to asexual females whilst the X chromosome is

hyper-methylated. Changes in gene expression and methylation between asexual females

and males are correlated, and this correlation is strongest for X-linked genes. Taken together

our findings suggest a role for DNA methylation in the regulation of aphid gene expression

and the establishment of sexual dimorphism.

Results and Discussion

Extensive sex-biased expression between asexual females and males

To identify genes with sex-biased expression in *M. persicae* clone O, we sequenced the

transcriptomes of asexual females and males (six biological replicates each) using RNA-seq

(**Supplementary Table 1**). After mapping these reads to the *M. persicae* clone O genome

(Mathers et al. 2017), we conducted differential expression analysis with edgeR (Robinson et

al. 2009). Genes were classified based on whether their expression was significantly biased

(edgeR; Benjamini-Hochberg (BH) corrected $p < 0.05$ and absolute fold change (FC) > 1.5)

towards asexual females (FAB genes) or males (MB genes). We also considered the magnitude

of sex bias, classifying genes as either moderately sex-biased ($1.5 \leq FC < 10$, for FAB or MB) or

extremely sex-biased ($FC \geq 10$, for FAB+ or MB+). In total, 3,433 genes exhibited sex-biased

expression (**Figure 1a**; **Supplementary Table 2**), representing 19 % of all annotated *M.*

persicae genes and 33 % of all genes with detectable expression (> 2 counts-per-million in at

least 3 samples, n = 10,334). MB genes outnumbered FAB genes by 18 % (1,778 vs 1,505,

binomial test; $p = 1.02 \times 10^{-6}$) and only a handful of FAB+ genes (15) were observed compared

to 135 MB+ genes (binomial test; $p = 1.28 \times 10^{-25}$; **Figure 1b**). This is consistent with patterns

of sex-biased gene expression in the pea aphid (Purandare et al. 2014) and other

invertebrates such as *Caenorhabditis* (Thomas et al. 2012; Albritton et al. 2014) and

Drosophila (Zhang et al. 2007), which also show a tendency towards an excess of male-biased

genes.

Figure 1: Differential gene expression between *M. persicae* asexual females and males. (a) Male (M; x-axis) and asexual female (FA; y-axis) gene expression expressed as $\log_{10}$ fragments per kilobase of transcript per million mapped reads (FPKM) averaged over 6 biological replicates for genes retained for differential expression (DE) analysis with edgeR (n = 10,334). DE genes are coloured according to the direction and magnitude of sex-bias (see main text). UB = unbiased expression (edgeR; Benjamini-Hochberg (BH) corrected $p > 0.05$ and absolute fold change (FC) > 1.5). **(b)** Male-biased (MB) genes significantly outnumber asexual female-biased (FAB) genes. **(c)** Asexual females and males differ significantly in expression at two out of five DNA methyltransferase genes (DNMT1a and DNMT3a; edgeR; BH corrected $p < 0.05$ and FC > 1.5). DNMT1b and DNMT3b are also significantly down-regulated in males (edgeR; BH corrected $p = 6.35 \times 10^{-6}$ and 0.039, respectively). However, the absolute FC of these genes falls below our cut-off of absolute FC > 1.5 (FC = 1.42 and 1.35, respectively).

Methylation genes are differentially expressed

Next, we used our transcriptome data to investigate expression patterns of known methylation genes in *M. persicae* asexual females and males. Genome-wide patterns of DNA methylation in animals are maintained by a toolkit of DNA methyltransferase genes (Schübeler 2015). *De novo* DNA methylation is established by DNMT3 and DNA methylation patterns are maintained by DNMT1 (Law and Jacobsen 2010). An additional homolog of DNMT1 and DNMT3, DNMT2, is responsible for tRNA methylation (Goll et al. 2006) and not involved in DNA methylation. Conservation of the DNA methylation toolkit varies across insects (Bewick et al. 2016) with DNMT1 being associated with the presence of detectable levels of DNA methylation. Aphid genomes contain a full complement of DNA methylation genes with two copies of DNMT1, a single copy of DNMT2, and two copies of DNMT3 (Mathers et al. 2017; Nicholson et al. 2015; Walsh et al. 2010). We find that DNMT1a is down-regulated in males, relative to asexual females (edgeR; BH corrected $p = 5.84 \times 10^{-40}$, abs. FC = 2.25), and DNMT3a is up-regulated in males (edgeR; BH corrected $p = 3.18 \times 10^{-14}$, abs. FC = 2.44) (**Figure 1c**). DNMT1b and DNMT3b are also down-regulated in males (edgeR; BH corrected $p = 6.35 \times 10^{-6}$ and 0.039, respectively), however the FC of these genes falls below our 1.5-fold threshold. In contrast, the tRNA methyltransferase DNMT2 is uniformly expressed (edgeR; BH corrected $p = 0.067$). These results suggest that changes in DNA methylation may be involved in the establishment of sexual dimorphism in *M. persicae*.

Genome-wide methylation patterns in *M. persicae*

DNA methylation has been poorly studied in insects outside of Holometabola and has only been superficially described in Hemiptera as part of a broad scale comparative analysis (Bewick et al. 2016). We therefore first sought to characterise genome-wide patterns of methylation in *M. persicae* before going on to investigate sex-specific changes in DNA methylation levels between asexual female and male morphs. To characterise genome-wide DNA methylation levels at base-level resolution, we sequenced bisulphite-treated DNA extracted from whole bodies of asexual females and males (three biological replicates each) derived from the same clonally reproducing population (clone O), and mapped these reads to the *M. persicae* clone O genome (Mathers et al. 2017) using Bismark (Krueger and Andrews 2011). After removal of ambiguously mapped reads and PCR duplicates, each replicate was

sequenced to between 24x and 37x average read depth (**Supplementary Table 3**), resulting

in 7,836,993 CpG sites covered by at least 5 reads in all samples.

M. persicae individuals harbour an obligate endosymbiont, *Buchnera aphidicola*, which lacks

a functional DNA methylation system (van Ham et al. 2003). We made use of *Buchnera*

derived reads in each sample to establish rates of false positive methylation calls caused by

incomplete cytosine conversions by mapping each sample to the *M. persicae Buchnera*

genome (Mathers et al. 2017) and quantifying methylation levels (**Supplementary Table 4**).

The average methylation level in *Buchnera* for Cs in any sequence context was 0.45% ± 0.68

(mean ± SD), indicating that bisulphite treatment of the aphid DNA was 99.55% efficient and

was consistent across samples. Based on this, we assessed methylation levels in *M. persicae*

for C's in a CpG, CHH and CHG context. Only Cs in a CpG context had methylation levels higher

than the false positive rate in *B. aphidicola*, indicating that CpG methylation is the

predominant form of DNA methylation in *M. persicae* (**Figure 2a**). Overall, global CpG

methylation levels (2.93% ± 0.32% of Cs; mean ± SD) were similar to those reported in other

hemipteran insects (2 – 4 %) and higher than in Hymenoptera (0.1 – 2.2 %) (Bewick et al.

2016). Exons were highly enriched for methylated CpGs relative to the rest of the genome (χ^2

= 1.07 x 10^8, d.f. = 1, p < 2.2 x 10^{-16}), with only 7.7% of methylated CpGs occurring in intergenic

regions (**Figure 2b and c**). Identification of significantly methylated CpG sites using a binomial

test that incorporates the false positive methylation rate (derived from *Buchnera*) showed

that methylation is non-randomly distributed across *M. persicae* gene bodies, with

methylated CpG sites biased towards the 3' end of genes despite the total number of CpG

sites being much higher at the 5' ends of genes (**Figure 2d**). This is likely driven by high rates

of methylation in 3' UTRs (**Figure 2c**). Interestingly, methylation bias towards the 3' end of

genes is potentially a unique feature of aphid or hemipteran CpG methylation as insects from

other orders show an opposite bias, with higher methylation at the 5' end of genes (Bonasio

et al. 2012; Hunt et al. 2013; Zemach et al. 2010).

Figure 2: The *M. persicae* methylome. (a) *Boxplots* showing the proportion of methylated cytosines (mC) by sequence context (CpG, CHG and CHH) for *M. persicae* and its obligate endosymbiont *Buchnera aphidicola*, which lacks a functional methylation system. **(b)** Example genome browser view showing the distribution of CpG methylation in asexual females and males across the first 100Kb of scaffold_93. **(c)** The distribution of methylated CpGs across genomic features and the proportion of methylated CpGs in each feature. Methylated and un-methylated CpG counts were summed across all replicates. **(d)** The distribution of all covered CpG sites (min. 5 reads per sample) and significantly methylated CpG sites (*binomial test*, BH-corrected p < 0.05) across *M. persicae* gene bodies. -1 = 1Kb upstream, 0 = transcription start site, 1 = transcription stop site, 2 = 1Kb upstream. **(e)** The distribution of RNA-seq expression levels in asexual females (log$_{10}$ FPKM) for un-methylated (0 - 1% CpG methylation) and methylated genes (FPKM = Fragments Per Kilobase of transcript per Million). Expression values were averaged across six biological replicates and methylation levels averaged across three biological replicates. Only genes with average expression levels of at least 1 FPKM in males and asexual females were included. Dots and whiskers inside the *violin plots* indicate median and interquartile range respectively. **(f)** As for **(e)** but showing the distribution of variation in expression between the six asexual female RNA-seq replicates (measured as the log$_{10}$ transformed coefficient of variation (log$_{10}$ CV) of FPKM) for un-methylated (0 - 1% CpG methylation) and methylated genes. **(g)** The relationship between the mean and the CV of gene expression for un-methylated and methylated genes with a trend line for each methylation level shown as a LOESS-

smoothed line with shaded areas indicating the 95% CI. The difference between the grey (un-
methylated; 0 - 1% CpG methylation) and pink/red lines (methylated; > 1% CpG methylation) shows
that methylation reduces the between-replicate variation in gene expression, particularly in highly
expressed genes. The negative correlation and downwards slope of trend lines shows that higher
expressed genes are better canalized, showing less between-individual variation in gene expression.

Next, we investigated the relationship between genome-wide patterns of DNA methylation

and gene expression using data for asexual females (**Supplementary Table 5**). We find that

the presence DNA methylation is positively associated with gene expression, with methylated

genes having significantly higher expression than un-methylated genes (**Figure 2e**). We also

find that methylated genes are more stably expressed than un-methylated genes (**Figure 2f**),

even after accounting for the higher expression of methylated genes (**Figure 2g**). The same

patterns were also observed using male methylation and gene expression data

(**Supplementary Figure 1**). Taken together, these data suggest that DNA methylation in

aphids may be involved in establishing and stabilising high gene expression, as has been

suggested in corals (Liew et al. 2017) and holometabolous insects (Wang et al. 2013; Patalano

et al. 2015; Xiang et al. 2010; Libbrecht et al. 2016).

Asexual females and males have distinct methylation profiles

To gain an overview of methylation differences between asexual female and male *M. persicae*

morphs, we conducted principle component analysis based on methylation levels of 350,782

CpG sites significantly methylated (*binomial test*, BH-corrected $p < 0.05$) in at least one

sample. Male and asexual female morphs clearly form distinct clusters, indicating

reproducible differences in global CpG methylation (**Figure 3a**). To further characterise

methylation differences between asexual females and males we conducted site-wise

differential methylation (DM) analysis, identifying 20,964 DM CpG sites (> 15% methylation

difference, BH corrected $p < 0.05$; **Supplementary Table 6**), 79% of which show a reduction

in methylation (hypo-methylation) in males relative to asexual females and 21% the opposite

(**Figure 3b**). This is significantly higher than expected by chance (see **Supplementary Figure

2**), and indicates that differences in methylation between asexual female and male morphs

are unlikely to be due to random between-sample variation. Rather, alterations in CpG

methylation appear to underpin differentiation between sexual morphs in aphids. These

findings are striking given the absence of significant levels of sex-biased or caste-biased

methylation in other insect systems (Libbrecht et al. 2016; Patalano et al. 2015; Herb et al.

2012; Wang et al. 2015).

Figure 3: Differential methylation between *M. persicae* asexual female and male morphs. (**a**) Principle component analysis (PCA) based on methylation levels at 350,782 CpG sites significantly methylated in at least one sample. PC1 seperates the samples based on sex (45% of the variation), whilst PC2 and PC3 seperate male and asexual female replicates, respectively (explaining 18 to 17 % of the variation). (**b**) *Volcano plot* showing results of MethylKit (Akalin et al. 2012) site-wise tests of differential methylation between asexual females (FA) and males (M). Methylation differences are shown for M relative to FA. Only CpG sites showing significant differential methylation (DM) (BH corrected *p* < 0.05) are shown. A minimum methylation difference threshold of 15% per site was applied to define a site DM between FA and M. (**c**) The number of differentially methylated sites per gene (±1Kb flanking region). (**d**) The distribution of DM CpG sites along *M. persicae* gene bodies. -1 = 1Kb upstream, 0 = transcription start site, 1 = transcription stop site, 2 = 1Kb upstream.

Overlap analysis revealed that the majority (92%) of DM CpG sites between asexual females

and males were located in gene bodies (± 1 Kb), with genes having between 1 and 107 DM

CpG sites (**Figure 3c**). These DM CpG sites were non-randomly distributed along gene bodies,

being biased towards the 5' end of genes (**Figure 3d**). As such, whilst overall methylation

levels are biased towards the 3' end of genes, sites with variable methylation are more likely

to be at the 5' end. To directly correlate gene body methylation levels with gene expression,

we also performed DM analysis at the gene level (**Supplementary Table 7**). This identified

1,344 DM genes with > 10% methylation difference (BH corrected p < 0.05), of which 205

showed significant male-biased methylation and 1,129 asexual female-biased methylation

(**Figure 4a and b**). Considering genes with variable methylation, males have undergone a

global loss of gene body methylation relative to asexual females (Wilcoxon signed-rank test,

p < 2.2 x 10^{-22}; **Figure 4c**).

Figure 4: Genome-wide changes in gene body methylation between asexual female and male

morphs. (**a**) Male (M; x-axis) and asexual female (FA; y-axis) gene-wise methylation levels averaged

over 3 biological replicates for genes methylated > 1% in at least one of the two morphs (n = 6,699).

Differentially methylated (DM) genes (MethylKit; > 10% methylation difference, BH corrected p <

0.05) are coloured according to the direction of sex-bias: MBm = male biased methylation, FABm =

female-biased methylation, UB = unbiased methylation. (**b**) FABm genes outnumber MBm genes. (**c**)

Violin plot showing the distribution of mean methylation level in FA and M for DM genes. Dots and

whiskers indicate median and interquartile range respectively; **** = Wilcoxon signed-rank test p <

0.0001. (**d**) Enriched GO terms relating to molecular function plotted in semantic space for FABm

genes and MBm genes (for terms relating to biological process see **Supplementary Figure 3**). A full

list of enriched GO terms for each DM class and functional category is given in **Supplementary Table**

8).

Gene ontology (GO) term enrichment analysis showed that asexual female-biased

methylation and male-biased methylation genes were both enriched for GO terms relating to

core biological processes, including metabolism and regulation of gene expression (**Figure 4d**;

Supplementary Figure 3; **Supplementary Table 8**). Protein SUMOylation is enriched among

genes with male-biased methylation. This is interesting because protein SUMOylation is

essential for dosage compensation of the *C. elegans* sex chromosome (Pferdehirt and Meyer

2013) and plays a key role in insect development and metamorphosis (Ureña et al. 2015).

Changes in methylation may therefore underpin core processes involved in aphid

polyphenism and sex determination. Consistent with this, we also find enrichment of

hormone signalling amongst genes with male-biased methylation, with 3 insulin genes hyper-

methylated in males (2 not expressed, 1 has male-specific expression). Insulin receptors

determine alternative wing morphs in planthoppers (Xu et al. 2015) and have been shown to

interact with the core sex determination gene TRANSFORMER-*2* (Zhuo et al. 2017).

The X chromosome has distinct patterns of expression and methylation

We identified X-linked scaffolds in the *M. persicae* genome assembly based on the ratio of

male to asexual female bisulphite sequencing coverage. This approach takes advantage of the

hemizygous condition of the X chromosome in males, which should result in X-linked scaffolds

having half the read depth of autosomal scaffolds (Jaquiéry et al. 2017). As expected, we

observe a bimodal distribution in the ratio of male to asexual female scaffold coverage, with

the lower coverage peak falling at approximately half the relative coverage of the higher

coverage peak (**Figure 5a**; **Supplementary Table 9**). Scaffolds in this lower coverage peak are

putatively derived from the X chromosome. To validate the coverage results, we mapped

known X-linked (n=4) and autosomal (n=8) microsatellite loci to the clone O genome and

retrieved the male to asexual female coverage ratios of their corresponding scaffolds. The

coverage of these known sex-linked scaffolds also exactly matches expectations (**Figure 5a**;

Supplementary Table 10). Using a cut-off of adjusted coverage, we identified 748 X-linked

scaffolds and 1,852 autosomal scaffolds, totalling 68.7 and 239.7 Mb of sequence respectively

(**Supplementary Figure 4**). Scaffolds assigned to the X chromosome therefore account for

22.3% of the assembled (scaffolds ≥ 20Kb) *M. persicae* clone O genome. This is in line with

expectations given the most common *M. persicae* karyotype of 2n = 12 and that the X

chromosome is larger than the autosomes (Blackman 1971a). Using the chromosomal

assignment of scaffolds, we were able to assign 3,110 gene models to the X chromosome and

10,768 to autosomes, leaving 4,555 (24.7%) gene models on unassigned scaffolds shorter

than 20 Kb. The number of identified X-linked genes is not different to expectations based on

the assembled size of the respective chromosomal regions (binomial test, p = 0.65). However,

we find that the X chromosome is depleted in coding sequence (CDS) compared to the

autosomes (6.3% vs 6.5%; χ^2 = 5,821.5, d.f. = 1, p < 2.2 x 10^{-16}). This is due to the reduced CDS

length of X-linked genes (Wilcoxon signed-rank test, p = 4.2 x 10^{-4}; **Supplementary Figure 5**).

Figure 5: Distinct patterns of methylation and expression between the *M. persicae* X chromosome

and autosomes. (**a**) X-linked and autosomal scaffolds (≥ 20Kb) in the *M. persicae* genome were

identified based on the relative coverage of BS-seq reads in males (M) compared to asexual females

(FA). Given the XO sex determination system of aphids, X-linked scaffolds are predicted to have half

autosomal coverage in males. A bimodal distribution in the ratio of M to FA coverage is clearly visible

(upper panel). We considered scaffolds falling in the lower coverage peak (ratio of adjusted coverage

< 1) as X-linked and scaffolds in the second, higher coverage peak (ratio of adjusted coverage > 1), as
autosomal. The assignment of scaffolds to the X chromosome or autosomes was validated by
comparing the M : FA ratio of coverage for scaffolds containing microsatellite markers on the X-
chromosome (blue dots) and autosome (red dots) (lower panel). (**b**) The distribution of gene body
methylation levels for X-linked and autosomal genes. (**c**) Observed / expected (odds ratio) counts of
DM and DE genes on the X chromosome by expression or methylation bias category. The X
chromosome is significantly enriched for MB+ genes (≥ 10-fold upregulation in M) and genes with
male-biased methylation (MBm). (**d**) The distribution of mean methylation levels in asexual females
(FA) and males (M) for X-linked and autosomal DM genes (MethylKit; > 10% methylation difference,
BH corrected p < 0.05). Methylation levels are significantly higher in FA than M for autosomal genes,
whereas M have a higher methylation than FA in X-linked genes (**d**) dots and whiskers inside the *violin*
plots indicate median and interquartile range respectively; *** = Wilcoxon signed-rank test p < 0.001,
**** = p < 0.0001.

Strikingly, the X chromosome has a distinct methylation landscape compared to the
autosomes (Anderson-Darling k-sample test, p = 1.35 x 10^{-65}; **Figure 5b**), with fewer highly
methylated genes. We also find opposing dynamics of sex-biased methylation between the X
chromosome and the autosomes. The X chromosome is significantly enriched for genes with
male-biased methylation and depleted for genes with female-biased methylation (χ^2 =
176.65, d.f. = 2, p < 2.2 x 10^{-16}; **Figure 5c**). Overall, X chromosome genes are hyper-methylated
in males (Wilcoxon signed-rank test, p = 8.6 x 10^{-4}; **Figure 5d**) compared to the genome-wide
pattern of hypo-methylation (Wilcoxon signed-rank test, p < 2.2 x 10^{-16}). Mirroring differences
in methylation between the X chromosome and the autosomes, we also find that the X
chromosome is enriched for genes with extreme male-biased expression (χ^2 = 42.38, d.f. = 1,
p = 7.5 x 10^{-11}; **Figure 5c**), a phenomenon also observed in the pea aphid (Jaquiéry et al. 2013).
Male-biased expression of X-linked genes is therefore conserved across two distantly related
aphid species, and, at least in the case of *M. persicae*, this also extends to patterns of DNA
methylation.

Finally, we investigated whether changes in methylation between *M. persicae* asexual
females and males are associated with changes in gene expression. The relationship between
gene expression and gene body methylation is an open question in invertebrates and few
studies have directly tested for changes in expression and methylation. We find that DM
genes are significantly enriched for DE (χ^2 = 7.84, d.f.= 1, p = 0.005), suggesting that
methylation changes may be involved in the regulation of at least a subset of sex-biased

genes. In support of this, we find a weak but significant positive correlation between changes
in gene expression and methylation between asexual females and males when considering
genes methylated (> 1%) and expressed (> 1 FPKM) in at least one of the sexes (n = 6,699;
Spearman's ρ = 0.089, *p* = 2.7 x 10^{-13}; **Figure 6a**). Interestingly, this correlation is driven by X-
linked genes which show a significantly stronger correlation between changes in expression
and methylation than autosomal genes (GLM: $F_{1,6185}$ = 93.07, *p* < 0.0001; **Figure 6b**). Combined
with recent results demonstrating a role for chromatin accessibility in the sex-specific
regulation of genes on the X chromosome and dosage compensation in the pea aphid (Richard
et al. 2017), our findings suggest a key role for epigenetics in establishing patterns of X-linked
gene expression in aphids.

Figure 6: Correlated changes in expression and methylation between asexual females and males.
(**a**) *Scatter plot* showing the relationship between fold-change (FC) in gene expression and
methylation between asexual females (FA) and males (M) for genes expressed (> 1 FPKM) and
methylated (> 1%) in at least one of the sexes (n = 6,699). Methylation levels of genes were estimated
across the whole gene body and averaged across replicates. Positive values indicate increased
expression or methylation in males, relative to asexual females; negative values indicate increased
expression or methylation in asexual females, relative to males. (**b**) The correlation between gene
expression changes and methylation changes between FA and M is significantly stronger for X-linked
genes (X; n = 925) than autosomal genes (A; n = 5,272). Spearman's ρ was used to assess significance
and strength of the relationship between change in expression and methylation for each set of genes.
The trend lines indicate linear fit with shaded areas indicating 95% confidence intervals.

Conclusions

We presented the first detailed analysis of genome-wide methylation patterns in an aphid,
evaluating its importance for gene expression and sexual differentiation. We found that 3,433

genes (19 % of the annotated genome) were differentially expressed between the males and
asexual females, and that there was a significant excess of male-biased genes. We also found
evidence suggesting that methylation plays an important role in sexual differentiation of
aphids, showing that DNMT1a and b are significantly downregulated in males, whereas
DNMT3a is upregulated in males. CpG methylation is the predominant form of DNA
methylation in *M. persicae* and, in contrast to other insects, exons were highly enriched for
methylated CpGs at the 3' end rather than the 5' end of genes. Methylation is positively
associated with gene expression, and in addition, methylated genes are more stably
expressed than un-methylated genes. Methylation was significantly reduced in males
compared to asexual females, yet remarkably, the X chromosome genes of males were hyper-
methylated. Given that differentially methylated genes were also significantly differentially
expressed between the sexes, we propose that changes in DNA methylation play a role in *M.*
persicae sexual differentiation. Our findings pave the way for future functional studies of DNA
methylation in aphids, and its potential role in the remarkable evolutionary potential of these
insects, and their extraordinary phenotypic plasticity.

Methods

Aphid rearing and sample preparation

An asexual colony of *M. persicae* clone O derived from a single apterous asexual female
(Mathers et al. 2017) was maintained on *Brassica rapa* plants in long-day conditions (14h
light, 22° C day time, and 20° C night time, 48% relative humidity). Male morphs were induced
by transferring the colony to short-day conditions (8h light, 18° C day time, and 16° C night
time, 48% relative humidity) and samples were collected 2 months after transfer. Replicate
samples were harvested from the same populations, with each replicate consisting of 20
adults, with apterous asexual females collected from the long-day population, and males from
the short-day population. Samples were immediately frozen in liquid nitrogen prior to RNA or
DNA extraction. DNA (three biological replicates per morph) was extracted using the CTAB
protocol (Marzachi et al. 1998), with the addition of a proteinase K digestion step during the
initial extraction. RNA (six biological replicates per morph) was extracted using the Trizol
reagent according to the manufacturers' protocol (Sigma), and further purified using the
RNeasy kit with on-column DNAse treatment (Qiagen).

Transcriptome sequencing

RNA samples were sent for sequencing at the Earlham Institute (Norwich, UK) where twelve non-orientated libraries were constructed using the TruSeq RNA protocol v2 (Illumina #15026495 Rev.F). 1ug of total RNA was enriched for mRNA using oligo(dT) beads. The RNA was then fragmented and first strand cDNA synthesised. Following end repair and adapter ligation, each library was subjected to a bead based size selection using Beckman Coulter XP beads (Beckman Coulter Inc., Brea, CA, USA) before performing PCR to enrich for fragments containing TruSeq adapter sequences. Libraries were then pooled and sequenced on the Illumina HiSeq 2000 platform (Illumina Inc., San Diego, CA, USA) (v3 chemistry; 2 x 100 bp), generating between 15 and 57 million paired-end reads per sample. RNA-seq reads have been deposited in the NCBI short read archive (SRA) under accession number PRJNA437622.

Gene expression analysis

Raw RNA-seq reads for each sample were trimmed for low quality bases and adapter contamination with Trim Galore! v0.4.0 using default settings for paired end reads (www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Gene-level expression quantification was then performed for each sample based on the *M. persicae* clone O reference genome and gene annotation (Mathers et al. 2017), using RSEM v1.2.31 (Li and Dewey 2011) with STAR v2.5.2a (Dobin et al. 2013). Average expression and the coefficient of variation was calculated per gene for asexual females and males separately using FPKM (fragments per kilobase of transcript per million) values estimated by RSEM. We also identified differentially expressed (DE) genes between asexual females and males using edgeR (Robinson et al. 2009) based on gene-level expected counts estimated by RSEM. Only genes with greater than 2 counts-per-million in at least three samples were retained for DE analysis and we considered genes DE if they had a fold-change (FC) ≥ 1.5 and $p < 0.05$ after adjusting for multiple testing using the Benjamini-Hochberg (BH) procedure (Benjamini and Hochberg 1995).

Bisulphite sequencing

Bisulphite sequencing library construction was performed using 500 ng genomic DNA per sample with a BIOO Scientific NEXTflex™ Bisulfite-Seq Kit (Bioo Scientific Corporation, Austin,

TX, USA) according to the manufacturer's instructions with the following modifications:
genomic DNA was sheared to 200-400 bp with a Covaris S2 sonicator (Covaris Inc., Woburn,
MA) using the following settings: duty cycle 10%, intensity 5, 200 cycles per burst for 120
seconds. The power mode was frequency sweeping, temperature 5-6°C and water level 12.
Libraries either received NEXTflex™ barcode #24 (GGTAGC) or #31 (CACGAT). All purified
libraries were QC checked with the Bioanalyzer DNA HS assay and further quantified by Qubit
dsDNA HS Assay Kit (Life Technologies, Carlsbad, CA, USA) before pooling as pairs. Pooled
libraries were further quantified by qPCR using a KAPA Library Quantification Kit - Illumina/ABI
Prism (Kapa Biosystems Inc., Wilmington, MA, USA) on a StepOnePlus™ Real-Time PCR System
(Life Technologies, Carlsbad, CA, USA). Sequencing was performed at the Earlham Institute
(Norwich, UK) on an Illumina HiSeq 2500 (Illumina Inc., San Diego, CA, USA) using paired-end
sequencing (v4 chemistry; 2 x 126 bp) with a 15% PhiX spike in, clustering to 650 K/mm^2. In
total, we generated between 70 and 127 million paired-end reads per sample.

DNA methylation analysis

Bisulphite treated reads for each sample were trimmed for low quality bases and adapter
contamination using Trim Galore! v0.4.0 with default settings
(www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Read pairs where one or both
reads were shorter than 75bp after trimming were discarded. We then mapped the trimmed
reads to the *M. persicae* clone O reference genome (Mathers et al. 2017) using Bismark
v0.16.1 (Krueger and Andrews 2011). Trimmed reads were also mapped to the genome of the
M. persicae strain of the obligate aphid endosymbiont *Buchnera aphidicola* (Mathers et al.
2017) to estimate the error rate of the C to T conversion. Reads derived from PCR duplicates
and that mapped to multiple locations in the genome were removed from downstream
analysis. The distribution of methylation across selected scaffolds was visualised using Sushi
(Phanstiel et al. 2014).

Overall levels of methylation in a CpG, CHG and CHH sequence context were estimated
directly from mapped reads with Bismark (Krueger and Andrews 2011). We also characterised
CpG methylation levels of features in the *M. persicae* clone O genome based on the reference
annotation (Mathers et al. 2017). Average CpG methylation levels of introns, exons, 5' UTRs,
3' UTRs and intergenic regions were calculated with bedtools v2.25.0 (Quinlan and Hall 2010),

pooling data from all replicates and counting overlapping methylated and unmethylated

CpGs. We also calculated per-gene methylation levels for asexual females and males

independently in the same way. To assess the genome-wide distribution of methylated CpGs,

we filtered CpG sites to those covered by at least five reads in all samples and used a binomial

test to identify significantly methylated sites in each sample using the C to T conversion error

rate (derived from mapping to *Buchnera*) as the probability of success and corrected for

multiple testing using the BH procedure (Benjamini and Hochberg 1995), setting the FDR at

5% (BH adjusted $p < 0.05$).

Methylation differences between asexual females and males were assessed using a principle

component analysis (PCA) and by identifying differentially methylated (DM) sites and genes.

PCA was carried out with prcomp, implemented in R v3.2.2 (R Core Team 2017), using the

methylation levels of CpG sites significantly methylated in a least one sample (binomial test,

BH adjusted $p < 0.05$). We identified DM sites and genes using logistic regression implemented

in MethylKit (Akalin et al. 2012) which accepts input directly from Bismark. *p* values were

adjusted to Q-values using the SLIM method (Wang et al. 2011) to account for multiple

testing. For the site-level analysis, we discarded CpG sites covered by less than 5 reads and

those that fell into the top 0.1% of coverage. We considered sites significantly DM if they had

at least a 15% methylation difference at a 5% FDR (Q < 0.05). At the gene level, we discarded

genes covered by less than 20 reads which fell into the top 1% of coverage, and called genes

as DM if they had at least 10% methylation difference and at a 5% FDR (Q < 0.05). A less

stringent percent methylation difference was used at the gene-level as the signal of DM may

be diluted over the length of the gene body. To assess the rate of false positive methylation

calls caused by random variation between samples we generated a null distribution of DM

calls at Q < 0.05 for a range of percentage methylation difference cut-offs using all possible,

non-redundant, pairs of samples where an asexual female sample is grouped with a male

sample (n = 19). Overall, these random pairings resulted in a low number of DM calls,

indicating our results are robust (**Supplementary Figure 2a and b**).

X chromosome identification

We used our whole-genome bisulphite sequencing data for males and asexual females to

identify X–linked scaffolds in the *M. persicae* clone O genome assembly based on the ratio of

male to asexual female coverage using a procedure similar to Jaquiéry et. al (2017). BAM files
generated by MethylKit were merged for each morph using Picard v2.1.1
(http://broadinstitute.github.io/picard/) to maximise the depth of coverage. We then
calculated per site sequence depth with SAMtools v1.3 (Li et al. 2009). The average depth of
the pooled asexual female and male samples was 79x and 90x, respectively. We then
calculated the ratio of male median depth of coverage to asexual female median depth of
coverage for all scaffolds longer than 20 Kb, normalising male coverage to that of asexual
female coverage (multiplying male median coverage by 79 / 90). This resulted in a clear
bimodal distribution with modes at ~0.75 and ~1.5 (**Figure 5a**). We applied a cut-off of male
to asexual female normalised median coverage ratio < 1 to assign scaffolds to the X
chromosome and > 1 to assign scaffolds to the autosomes. To validate the coverage results,
we mapped known X-linked (n=4) and autosomal (n=8) microsatellite loci from Sloane et. al.
(2001) and Wilson et. al. (2004) to the clone O genome with blastn and retrieved coverage
ratios for their respective scaffolds.

Testing for correlation between changes in methylation and expression

To investigate the relationship between changes in gene expression and methylation we
compared expression and methylation levels of genes in males and asexual females. Using
average expression (FPKM) and methylation levels, we calculated the $\log_2$ FC in expression
(FC_{Expr}) and methylation (FC_{Meth}), and tested for correlation using Spearman's ρ (rho). We also
investigated the effect of chromosomal location (X chromosome vs. autosomes) on the
relationship between gene expression and methylation using a general linear model (GLM).
The GLM was formulated with FC_{Expr} as the response variable, and FC_{Meth} as a covariate,
crossed with chromosome (as fixed factor). This interaction term tests whether the slopes of
the regression lines of the X chromosome and autosomes run parallel.

Annotation of methyltransferase genes

Amino acid sequences of human DNA methyltransferase genes were blasted against
annotated protein sequences of *Myzus persicae* Clone O (Mathers et al., 2017). The top *M.*
persicae clone O hit for each gene was then used to blast against the *M. persicae* protein set
in an iterative fashion until no additional genes were identified. The E value were set as 1E-
10.

GO term enrichment analysis

Go term enrichment analysis of specific gene sets was performed with BINGO (Maere et al.

2005) using the complete *M. persicae* clone O proteome as the reference set. Redundant

terms were then removed with REVIGO (Supek et al. 2011).

Data availability

524 Raw RNA-seq and BS-seq data generated for this study have been deposited in the NCBI short

525 read archive under accession number PRJNA437622.

Acknowledgements

This work was supported by a BBSRC Industrial Partnership Award (BB/L002108/1) to S.H.,

D.S. and C.v.O. and co-funded by Syngenta, the Plant Health Institute Strategy Programme

(BB/P012574/1) awarded to the John Innes Centre, Gatsby Charitable Foundation funding to

S.H., a BBSRC award (BB/N02317X/1) to C.v.O., as well as support by the Earth & Life

Systems Alliance (ELSA). Next-generation sequencing and library construction was delivered

via the Biotechnology and Biological Sciences Research Council (BBSRC) National Capability

in Genomics and Single Cell (BB/CCG1720/1) at Earlham Institute by members of the

Genomics Pipelines Group. We thank Anna Jordan at the JIC insectary for assistance with

aphid rearing and identification of aphid morphs.

Author contributions

537 T.C.M., S.T.M., Y.C., D.S., S.H. and C.v.O. conceived the study. T.C.M. performed

538 bioinformatics analysis with additional analysis performed by C.v.O, G.K. and Y.C.. S.T.M.

539 performed aphid morph phenotyping and extracted DNA and RNA. L.P.A. constructed

540 bisulphite sequencing libraries. T.C.M, C.v.O and S.H. wrote the manuscript. All authors read,

541 edited and approved the final manuscript.

Competing interests

543 The authors declare no competing financial interests.

References

Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, Mason CE. 2012. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol* **13**: R87.

Albritton SE, Kranz AL, Rao P, Kramer M, Dieterich C, Ercan S. 2014. Sex-biased gene expression and evolution of the X chromosome in nematodes. *Genetics* **197**: 865–883.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc* **57**: 289–300.

Bewick AJ, Niederhuth CE, Ji L, Rohr NA, Griffin PT, Leebens-Mack J, Schmitz RJ. 2017. The evolution of CHROMOMETHYLASES and gene body DNA methylation in plants. *Genome Biol* **18**: 65.

Bewick AJ, Vogel KJ, Moore AJ, Schmitz RJ. 2016. Evolution of DNA methylation across insects. *Mol Biol Evol*.

Blackman RL. 1971a. Chromosomal abnormalities in an anholocyclic biotype of *Myzus persicae* (Sulzer). *Experientia* **27**: 704–706.

Blackman RL. 1971b. Variation in the photoperiodic response within natural populations of *Myzus persicae* (Sulz.). *Bull Entomol Res* **60**: 533–46.

Bonasio R, Li Q, Lian J, Mutti NS, Jin L, Zhao H, Zhang P, Wen P, Xiang H, Ding Y, et al. 2012. Genome-wide and caste-specific DNA methylomes of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Curr Biol* **22**: 1755–1764.

Dixon AFG. 1977. Aphid ecology: life cycles, polymorphism, and population regulation. *Annu Rev Ecol Syst* **8**: 329–353.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Feng S, Cokus SJ, Zhang X, Chen P-Y, Bostick M, Goll MG, Hetzel J, Jain J, Strauss SH, Halpern ME, et al. 2010. Conservation and divergence of methylation patterning in plants and animals. *Proc Natl Acad Sci U S A* **107**: 8689–94.

Goll MG, Kirpekar F, Maggert KA, Yoder JA, Hsieh C, Zhang X, Golic KG, Jacobsen SE, Bestor TH. 2006. Methylation of tRNA[Asp] by the DNA methyltransferase homolog Dnmt2. *Science* **311**: 395–398.

Guo H, Zhu P, Yan L, Li R, Hu B, Lian Y, Yan J, Ren X, Lin S, Li J, et al. 2014. The DNA

methylation landscape of human early embryos. *Nature* **511**: 606–610.

Herb BR, Wolschin F, Hansen KD, Aryee MJ, Langmead B, Irizarry R, Amdam G V, Feinberg

AP. 2012. Reversible switching between epigenetic states in honeybee behavioral

subcastes. *Nat Neurosci* **15**: 1371–1373.

Hunt BG, Brisson JA, Yi S V., Goodisman MAD. 2010. Functional conservation of DNA

methylation in the pea aphid and the honeybee. *Genome Biol Evol* **2**: 719–728.

Hunt BG, Glastad KM, Yi S V., Goodisman MAD. 2013. The function of intragenic DNA

methylation: Insights from insect epigenomes. *Integr Comp Biol* **53**: 319–328.

Jaquiéry J, Peccoud J, Ouisse T, Legeai F, Gouin A. 2017. Disentangling the causes for faster-X

evolution in aphids. *bioRxiv* 1–33.

Jaquiéry J, Rispe C, Roze D, Legeai F, Le Trionnaire G, Stoeckel S, Mieuzet L, Da Silva C,

Poulain J, Prunier-Leterme N, et al. 2013. Masculinization of the X chromosome in the

pea aphid. *PLoS Genet* **9**.

Krueger F, Andrews SR. 2011. Bismark: A flexible aligner and methylation caller for Bisulfite-

Seq applications. *Bioinformatics* **27**: 1571–1572.

Law J a, Jacobsen SE. 2010. Establishing, maintaining and modifying DNA methylation

patterns in plants and animals. *Nat Rev Genet* **11**: 204–220.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics* **12**: 323.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.

2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–

2079.

Libbrecht R, Oxley PR, Keller L, Kronauer DJC. 2016. Robust DNA methylation in the clonal

raider ant brain. *Curr Biol* **26**: 1–5.

Liew YJ, Zoccola D, Li Y, Tambutté E, Venn AA, Craig T. 2017. Epigenome-associated

phenotypic acclimatization to ocean acidification in a reef-building coral. *bioRxiv*.

Maere S, Heymans K, Kuiper M. 2005. BiNGO: A Cytoscape plugin to assess

overrepresentation of gene ontology categories in biological networks. *Bioinformatics*

21: 3448–3449.

Marzachi C, Veratti F, Bosco D. 1998. Direct PCR detection of phytoplasmas in

experimentally infected insects. *Ann Appl Biol* **133**: 45–54.

Mathers TC, Chen Y, Kaithakottil G, Legeai F, Mugford ST, Baa-Puyoulet P, Bretaudeau A,

Clavijo B, Colella S, Collin O, et al. 2017. Rapid transcriptional plasticity of duplicated
gene clusters enables a clonally reproducing aphid to colonise diverse plant species.
Genome Biol **18**: 27.

Misof B, Liu S, Meusemann K, Peters R. 2014. Phylogenomics resolves the timing and
pattern of insect evolution. *Science (80-)* **346**: 763–768.

Müller CB, Williams IS, Hardie J. 2001. The role of nutrition, crowding and interspecific
interactions in the development of winged aphids. *Ecol Entomol* **26**: 330–340.

Nicholson SJ, Nickerson ML, Dean M, Song Y, Hoyt PR, Rhee H, Kim C, Puterka GJ. 2015. The
genome of *Diuraphis noxia*, a global aphid pest of small grains. *BMC Genomics* **16**: 429.

Okae H, Chiba H, Hiura H, Hamada H, Sato A. 2014. Genome-wide analysis of DNA
methylation dynamics during early human development. *PLoS Genet* **10**: 1–12.

Patalano S, Vlasova A, Wyatt C, Ewels P, Camara F, Ferreira PG, Asher CL, Jurkowski TP,
Segonds-pichon A, Bachman M, et al. 2015. Molecular signatures of plastic phenotypes
in two eusocial insect species with simple societies. *PNAS*.

Pferdehirt RR, Meyer BJ. 2013. SUMOylation is essential for sex-specific assembly and
function of the *Caenorhabditis elegans* dosage compensation complex on X
chromosomes. *Proc Natl Acad Sci* **110**: E3810–E3819.

Phanstiel DH, Boyle AP, Araya CL, Snyder MP. 2014. Sushi.R: Flexible, quantitative and
integrative genomic visualizations for publication-quality multi-panel figures.
Bioinformatics **30**: 2808–2810.

Purandare SR, Bickel RD, Jaquiery J, Rispe C, Brisson J a. 2014. Accelerated evolution of
morph-biased genes in pea aphids. *Mol Biol Evol* **31**: 2073–2083.

Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842.

R Core Team. 2017. R: A Language and Environment for Statistical Computing.

Richard G, Legeai F, Prunier-Leterme N, Bretaudeau A, Tagu D, Jaquiéry J, Le Trionnaire G.
2017. Dosage compensation and sex-specific epigenetic landscape of the X
chromosome in the pea aphid. *Epigenetics Chromatin* **10**: 30.

Robinson MD, McCarthy DJ, Smyth GK. 2009. edgeR: A Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

Schübeler D. 2015. Function and information content of DNA methylation. *Nature* **517**: 321–
326.

Sloane M a, Sunnucks P, Wilson a C, Hales DF. 2001. Microsatellite isolation, linkage group

identification and determination of recombination frequency in the peach-potato

aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae). *Genet Res* **77**: 251–60.

Srinivasan DG, Brisson J a. 2012. Aphids: a model for polyphenism and epigenetics. *Genet*

Res Int **2012**: 431531.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. Revigo summarizes and visualizes long lists of

gene ontology terms. *PLoS One* **6**.

Suzuki MM, Bird A. 2008. DNA methylation landscapes: Provocative insights from

epigenomics. *Nat Rev Genet* **9**: 465–476.

Thomas CG, Li R, Smith HE, Woodruff GC, Oliver B, Haag ES. 2012. Simplification and

desexualization of gene expression in self-fertile nematodes. *Curr Biol* **22**: 2167–2172.

Ureña E, Pirone L, Chafino S, Pérez C, Sutherland JD, Lang V, Rodriguez MS, Lopitz-Otsoa F,

Blanco FJ, Barrio R, et al. 2015. Evolution of SUMO function and chain Formation in

insects. *Mol Biol Evol* **33**: msv242.

van Ham RCHJ, Kamerbeek J, Palacios C, Rausell C, Abascal F, Bastolla U, Fernández JM,

Jiménez L, Postigo M, Silva FJ, et al. 2003. Reductive genome evolution in *Buchnera*

aphidicola. *Proc Natl Acad Sci U S A* **100**: 581–586.

Walsh TK, Brisson J a., Robertson HM, Gordon K, Jaubert-Possamai S, Tagu D, Edwards OR.

2010. A functional DNA methylation system in the pea aphid, *Acyrthosiphon pisum*.

Insect Mol Biol **19**: 215–228.

Wang HQ, Tuominen LK, Tsai CJ. 2011. SLIM: A sliding linear model for estimating the

proportion of true null hypotheses in datasets with dependence structures.

Bioinformatics **27**: 225–231.

Wang X, Werren JH, Clark AG. 2015. Genetic and epigenetic architecture of sex-biased

expression in the jewel wasps *Nasonia vitripennis* and *giraulti*. *Proc Natl Acad Sci*

201510338.

Wang X, Wheeler D, Avery A, Rago A, Choi JH, Colbourne JK, Clark AG, Werren JH. 2013.

Function and Evolution of DNA Methylation in *Nasonia vitripennis*. *PLoS Genet* **9**.

Wilson A, Sunnucks P, Hales D. 1997. Random loss of X chromosome at male determination

in an aphid, *Sitobion* near *fragariae*, detected using an X-linked polymorphic

microsatellite marker. *Genet Res* **69**: 233–236.

Wilson ACC, Massonnet B, Simon J-C, Prunier-Leterme N, Dolatti L, Llewellyn KS, C FC, C RC,

Blackman RL, Estoup A, et al. 2004. Cross-species amplification of microsatellite loci in
aphids: assessment and application. *Mol Ecol …* **4**: 104–109.

Xiang H, Zhu J, Chen Q, Dai F, Li X, Li M, Zhang H, Zhang G, Li D, Dong Y, et al. 2010. Single
base–resolution methylome of the silkworm reveals a sparse epigenomic map. *Nat*
Biotechnol **28**: 516–520.

Xu H-J, Xue J, Lu B, Zhang X-C, Zhuo J-C, He S-F, Ma X-F, Jiang Y-Q, Fan H-W, Xu J-Y, et al.
2015. Two insulin receptors determine alternative wing morphs in planthoppers.
Nature **519**: 464–467.

Yan H, Simola DF, Bonasio R, Liebig J, Berger SL, Reinberg D. 2014. Eusocial insects as
emerging models for behavioural epigenetics. *Nat Rev Genet* **15**: 677–688.

Zemach A, McDaniel IE, Silva P, Zilberman D. 2010. Genome-wide evolutionary analysis of
eukaryotic DNA methylation. *Science (80-)* **328**: 916–919.

Zemach A, Zilberman D. 2010. Evolution of Eukaryotic DNA methylation and the pursuit of
safer sex. *Curr Biol* **20**: R780–R785.

Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. 2007. Constraint and turnover in sex-biased
gene expression in the genus *Drosophila*. *Nature* **450**: 233–237.

Zhuo JC, Lei C, Shi JK, Xu N, Xue WH, Zhang MQ, Ren ZW, Zhang HH, Zhang CX. 2017. Tra-2
mediates cross-talk between sex determination and wing polyphenism in female
Nilaparvata lugens. *Genetics* **207**: 1067–1078.