Distinct Functional Connectivity Mode during Viewing Natural Scenes Revealed by Principal Component Analysis

Murat Demirtaş[1,2], Adrian Ponce-Alvarez[2], Matthieu Gilson[2], Patric Hagmann[4],Dante Mantini[5,6], Viviana Betti[7], Gian Luca Romani[8], Karl Friston[9], Maurizio Corbetta[10], Gustavo Deco[2,3]

[1]N3 Division, Department of Psychiatry, Yale University, 40 Temple Street, New Haven, 06511, Connecticut, United States

[2]Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Roc Boronat 138, Barcelona, 08018, Spain

[3]Institució Catalana de la Recerca i Estudis Avançats (ICREA), Universitat Pompeu Fabra, Passeig Lluís Companys 23, Barcelona, 08010, Spain

[4]Department of Radiology, Lausanne University Hospital and University of Lausanne (CHUV-UNIL), Rue du Bugnon 46, 1011 Lausanne, Switzerland.

[5]Research Center for Motor Control and Neuroplasticity, KU Leuven, Tervuursevest 101, 3001 Leuven, Belgium.

[6]Department of Neurorehabilitation, IRCCS San Camillo Hospital Foundation, via Alberoni 70, 30126 Venice Lido, Italy.

[7]Department of Psychology, Sapienza University of Rome, via dei Marsi 78, 00185 Rome; Fondazione Santa Lucia and Istituto Di Ricovero e Cura a Carattere Scientifico, 00142, Rome, Italy.

[8]Institute for Advanced Biomedical Technologies, "G. d'Annunzio" University of Chieti-Pescara, 66100, Chieti, Italy.

[9]Wellcome Trust Centre for Neuroimaging, Institute of Neurology, University College London, 12 Queen Square, London WC1N 3BG, United Kingdom.

[10]Departments of Neurology, Radiology, Anatomy of Neurobiology, School of Medicine, Washington University, St. Louis, St Louis, USA

Corresponding author:

Murat Demirtaş, Yale University, 40 Temple Street, New Haven, 06511, Connecticut, United States. murat.demirtas@yale.edu

Abstract

A fundamental question in systems neuroscience is how spontaneous activity at rest is reorganized during task performance. Recent studies suggest a strong relationship between resting and task FC. Furthermore, the relationship between resting and task FC has been shown to reflect individual differences. Particularly, various studies have demonstrated that the FC has higher reliability and provides enhanced detection of individual differences while viewing natural scenes. Although the large-scale organization of FC during rest and movie-viewing conditions have been well studied in relation to individual variations, the re-organization of FC during viewing natural scenes have not been studied in depth. In this study, we used principal component analysis on FC during rest and movie-viewing condition to characterize the dimensionality of FC patterns across conditions and subjects. We found that the variations in FC patterns related to viewing natural scenes can be explained by a single component, which enables identification of the task over subjects with 100% accuracy. We showed that the FC mode associated to viewing natural scenes better reflects individual variations. Furthermore, we investigated the signatures of movie-viewing-specific functional modes in dynamic FC based on phase-locking values between brain regions. We found that the movie-specific functional mode is persistent across time; suggesting the emergence of a stable processing mode. To explain the reorganization of whole-brain FC through the changes in local dynamics, we appeal to a large-scale computational model. This modelling suggested that the reorganization of whole-brain FC is associated to the interaction between frontal-parietal and frontal-temporal activation patterns.

Introduction

Neural dynamics underwrite information processing at multiple spatiotemporal scales. The neural correlates of information processing at a local scale have been widely studied. However, the integration of information in whole-brain level is also crucial for understanding brain function (Baars, 1993; Tononi, 2004). Long-range synchronization of oscillatory activity has been proposed as dynamical mechanism for mediating the interaction between brain areas in a task-dependent manner (Engel et al., 2001; Fries, 2005). Recent studies showed that neuronal synchronization mediates neuronal communication in large-scale cortical networks during task performance (Betti et al., 2013) and resting state (de Pasquale et al., 2010).

Resting state functional connectivity (rs-FC) is a powerful technique to characterize large-scale organization of brain activity based on the temporal correlations between blood oxygen level-dependent (BOLD) signals (Biswal et al., 1995). Rs-FC patterns have been shown to provide 'fingerprints' for the functional brain organization of individuals (Finn et al., 2015). Furthermore, recent studies showed strong relationship between the FC during resting state and task performance (Betti et al., 2013; Cole et al., 2014, 2016; Rosenberg et al., 2015). The relationship between resting state and task FC has been shown to reflect individual differences (Tavor et al., 2016). In particular, experimental paradigms such as viewing natural scenes (i.e. movie watching) are of interest due to their ecological validity (Betti et al., 2013). Several studies found that FC has higher reliability and provides enhanced detection of individual differences while viewing natural scenes (Kim et al., 2017; Vanderwal et al., 2015, 2017). However, the features underlying the enhanced reliability of FC are not clear given that the large-scale organization of FC is very similar during rest and viewing natural scenes. Theoretical studies have proposed that entropy of the cortical activity space is reduced during task (Ponce-Alvarez et al., 2015). Therefore, the functional reorganization during viewing natural scenes is expected to reflect not only individual subject-specific variations but also task-specific variations. Nevertheless, the task-dependent reorganization of FC during viewing natural scenes and its relationship to individual subject-specific variations are poorly understood.

Understanding the organization of whole-brain FC during distinct conditions is challenging because of enormous dimensionality, which increases quadratically with the number of brain regions. Principal component analysis (PCA) is widely known and intuitive mathematical procedure to characterize the dimensionality of data. PCA transforms a set of observations into orthogonal components and allows characterizing the relationship between these orthogonal components and the projections of individual observations. PCA and associated techniques have been used to characterize resting-state fluctuations (Carbonell et al., 2011), whole-brain connectivity dynamics (Allen et al., 2012) and disease-related rs-FC states (Craddock et al., 2009). In this paper, we used PCA to investigate the dimensionality of FC during rest and movie-viewing condition. Based on the projections of individual subjects and sessions on principal components, we identified FC-states specific to natural viewing condition. Then, we

63 studied the consistency of this principal component across sessions, subjects, and under
64 different preprocessing approaches.
65
Another important question related to the task-dependent reorganization of FC is whether
alterations in grand-average FC (over the whole session) reflect a persistent (temporally stable)
functional state or they reflect the emergence of various functional states fluctuating over time
(Gonzalez-Castillo et al., 2015). To answer this question, we extended our analysis beyond
grand-average FC states and investigated the temporal fluctuations in FC states based on the
dynamics of phase-coupling among brain regions.
72
Although empirical findings provide insights on the task-dependent reorganization of whole-
brain FC, these results may not offer a mechanistic understanding. We therefore adopted a
mechanistic approach for task-dependent reorganization of whole-brain FC using large-scale
by physically plausible modelling framework. We constraint the long-range interactions
between brain regions by diffusion weight imaging-derived (DWI) structural connectivity, and
studied the alterations in local dynamics of each brain regions during natural viewing conditions.
79

Results

Principal component analysis reveals distinctive FC mode during movie watching condition

The grand average FC during resting state and movie exhibited similar patterns (r = 0.8, p-value < 0.0001) (Figure 1A). However, the similarity across FCs of individual subjects was substantially higher under the same condition (resting state r = 0.46 ± 0.06; movie r = 0.49 ± 0.06) than across conditions (r = 0.40 ± 0.07). To characterize the variations in FC during rest-state and movie viewing conditions, we performed principal component analysis (PCA) across FCs of individual subjects (Figure 1B-C).

We concatenated FCs of 21 individuals during 2 separate sessions of resting state and movie watching conditions and then employed PCA. We found that the first principal component (PC-1) explaining 25.8% of the variance (Figure 2A) reflected the dominant FC pattern that were conserved over conditions. The projections of PC-1 were significantly correlated with global signal standard deviations of individual subjects/session (r = 0.99, p-value < 0.0001) (Figure 2J). This result suggested that the principal mode of variation in FC reflects the global signal variance and associated overall synchronization levels. However, the second principal component (PC-2) (Figure 2C) explaining 7.2% of the variance clearly distinguished the movie condition from resting state. Based on the projections of two principal components (i.e. 2-dimensional projections of subject/session data on principal components), we found that movie-specific PC-2 separates the two conditions with 100% accuracy (Figure 2D).

Previous studies have shown restricted subject movements and increased arousal while watching natural scenes (Vanderwal et al., 2015). Therefore, the movie-specific PC may also reflect the contributions of artefactual signal changes. To rule out possible artefactual contributions, we repeated the analyses after regressing out the global signal (Figure 2E-H). We found that after global signal regression (GSR) the first principal component (PC-1) explaining 9.69% of the variance reflected movie-specific variations in FC. The projections of PC-1 separated two conditions with 100% accuracy (Figure 2H). Furthermore, the topologies of movie-specific component modes were highly consistent with and without GSR (r = 0.81, p-value < 0.0001). These results showed that the movie-specific variations in FC patterns can be explained by a single dimension.

Movie-specific FC variations are consistent across sessions and they reflect individual variations better than non-specific principal components

We repeated the PCA (with and without GSR) using 2 sessions separately. For both sessions we found the movie-specific and non-specific components without GSR (Figure 3A-B) and with GSR (Figure 3C-D). Without GSR, PC-1 (global signal component) of session 1 and session 2 was significantly correlated (r = 0.75, p-value < 0.0001). However, the similarity between movie-specific PC-2 of session 1 and session 2 was substantially higher than that of PC-1 (r = 0.83,

120 p < 0.0001). Similarly, with GSR, we found higher similarity between session 1 and session 2
121 for movie-specific PC-1 (r = 0.82, p-value < 0.0001) than for non-specific PC-2 (r = 0.54, p-
122 value < 0.0001). These results show that the movie-specific principal component is highly
123 consistent across sessions.
124
To investigate the individual variations associated to each principal component, we compared
the PC projections across sessions. We found that the projections of movie-specific
components were highly consistent across sessions (without GSR r = 0.92, p-value < 0.0001;
with GSR r = 0.93, p-value < 0.0001), and the correlations were substantially higher than those
of non-specific components (without GSR r = 0.76, p-value < 0.0001; with GSR r = 0.52, p-
value < 0.0001). These results show that the FC topography related to movie-watching
conditions reflects individual variations better than that related to resting condition.
132
Movie-specific FC patterns are temporally stable
We investigated whether the movie-specific FC patterns are temporally stable in time or
whether they emerge as a consequence of the fluctuations in FC (i.e. collection of single or
multiple transient states). We constructed dynamic FC (dFC) based on the time-dependent
fluctuations in phase-locking values (PLVs) between brain regions. First, we band-pass filtered
the BOLD time-series in 0.04-0.07Hz narrow-band. After employing Hilbert transform, we
calculated the PLVs at each time point using instantaneous phases of each brain region (Figure
4A). Then, we calculated the correlation between the instantaneous PLVs and average PLVs
of resting state and movie sessions. To avoid any session-specific bias, we calculated the
correlations between average PLVs of resting state and movie from session 1 and
instantaneous PLVs of session 2 (and vice versa). We also repeated the PCA on average PLV
matrices for session 1 and session 2 (Figure 4C, F). We found that the principal components
based on PLVs exhibit movie condition specificity for both sessions. Furthermore, movie-
specific PC of PLVs were highly correlated with that of Pearson correlation based FC (r = 0.88,
p-value < 0.0001).
148
For resting state sessions, the correlations between PLVs and average PLVs of movie sessions
were significantly lower than that for resting state sessions across time points (Wilcoxon signed-
rank test p-value < 0.0001, for both session 1 and session 2). For movie sessions, the
correlations between PLVs and average PLVs of movie sessions were significantly higher than
that for resting state sessions across time points (Wilcoxon signed-rank test p-value < 0.0001,
for both session 1 and session 2) (Figure 4D-E). The results were the same when the analyses
were repeated using movie-specific and non-specific principal components instead of average
PLVs (Supplementary Figure 1). In addition, we constructed the average dFC across subjects
and estimated the correlation between the average PLVs at each time-point and average PLVs
of resting-state and movie sessions. We found that for both resting state and movie sessions,
the PLVs at each time-point exhibited higher correlations with grand-average PLVs of matching

condition (Supplementary Figure 1). We further employed PCA over the PLV fluctuations
concatenated across sessions for each subject. The trajectories of the principle components
revealed emergence of movie-specific FC as a distinct mode for most of the subjects
(Supplementary Figure 2). Nevertheless, we did not find task-specific components for PLVs
concatenated across all subjects and all sessions. These results suggest that the movie-
specific FC pattern emerges as a temporally stable mode during movie watching sessions,
although the time-resolved FC states is difficult to estimate.
167
Large-scale computational modelling of the regional dynamics underlying movie-
watching FC
We used a large-scale computational model to characterize the alterations in regional dynamics
associated to the movie-watching condition. We used Hopf normal model to characterize the
BOLD activity of each region (Deco et al., 2017). The regions were coupled to each other via
DWI-derived structural connectivity scaled by a global coupling parameter (Figure 5A). The
dynamics of each region were governed by local bifurcation parameter (a). The local bifurcation
parameters (a) reflects whether an individual region is dominantly in a noise-driven regime (a
< 0), oscillatory regime (a > 0), or alternates between the two regimes (a ~ 0) (Figure 5A). We
estimated the optimal global coupling and local bifurcation parameters of each subject/session
by maximizing the similarity (i.e. Pearson correlation) between empirical and model FCs using
gradient-descent optimization. There was no significant difference in the model fit for resting-
state (r = 0.518 ± 0.057) and movie sessions (r = 0.497 ± 0.045) (T = 1.256, p-value = 0.219,
permutation t-test).
182
To characterize the overall topology underlying each condition, first we estimated the optimal
global coupling parameter (g) and optimal bifurcation parameters (a) for resting state and movie
watching condition based on the similarity between average empirical and model FC. At rest,
the average optimal bifurcation parameters were low in parietal and temporal regions, whereas
they were higher in occipital and frontal regions (Figure 5B). For movie condition, the bifurcation
parameters were increased in parietal and temporal regions and decreased in anterior
cingulate, lateral prefrontal cortices and in supramarginal gyrus (Figure 5C). There was no
significant difference between the mean optimal bifurcation parameters of rest and movie
conditions (Figure 5D).
192
To quantify the group differences, we compared regional optimal bifurcation parameters of
resting state and movie sessions (Figure 6A). We found no significant differences in global
coupling parameters between rest and movie conditions (Figure 6B). In movie condition, the
local bifurcation parameters were significantly decreased towards negative values in bilateral
caudal anterior and posterior cingulate, right supramarginal gyrus, and left post- and para-
central cortices (Figure 6D). In contrast, the bifurcation parameters were significantly increased
in bilateral orbital frontal cortex, right lateral orbital frontal cortex, right middle rostral frontal

cortex, right superior parietal cortex, right fusiform gyrus, and left frontal pole and left medial
temporal cortex (Figure 6D).
202
Finally, we repeated the PCA on optimal bifurcation parameters across subjects and conditions
(Figure 6E-G). The first principal component (PC-1) explaining 41.77% of the variation and the
second principal component (PC-2) explaining 10.25% of the variation were both significantly
correlated with the contrast between average bifurcation parameters of movie and resting-state
sessions (PC-1 r = 0.38, p-value = 0.002; PC-2 r = 0.73, p-value < 0.0001). The first principal
component (PC-1) exhibited a strong positive peak in precuneus and isthmus of cingulate,
slightly higher values in medial frontal and temporal regions, which is very similar to default
mode network (DMN) topography (Figure 6E). The second principal component (PC-2) showed
higher values in temporal and frontal regions (Figure 6F). Furthermore, the second principal
component (PC-2) exhibited movie specificity (Figure 6G). This result shows that the alterations
in regional dynamics in the movie-condition reflect at enhanced influence of frontal-temporal
brain regions on whole-brain dynamics.

Discussion

We characterized the reorganization of FC during natural viewing condition compared to resting state. Using principal component analysis, we found that the alterations in FC during natural viewing condition can be explained along a single dimension or mode of variation (i.e. a condition-specific pattern of connectivity that captures the variations across subjects). The projections of the FCs of each subject on the movie-specific principal component provided a clear separation between conditions with classification accuracy of 100%.

FC signatures of viewing natural scenes

The movie-specific FC topology exhibited enhanced connectivity between occipital-temporal regions and frontal-parietal regions (Supplementary Figure 3). This result can be interpreted as enhanced communication between sensory and association regions during natural viewing condition. Previous studies reported that the subjects show decreased head movements and higher arousal while natural viewing condition (Vanderwal et al., 2015). Therefore, the exceptionally high classification power can also reflect systematic artifacts. To rule of this possibility we repeated the analysis after global signal regression (GSR), separate sessions, bandpass filtered signals, and phase-locking values. We found that the results are invariant to different preprocessing approaches. Furthermore, the movie-specific FC mode was highly consistent before and after GSR, and across sessions. The correlations between movie-specific components across sessions were higher than those for non-specific components. These findings confirmed previous studies showing the enhanced reliability and individual subject detection during natural viewing conditions (Kim et al., 2017; Vanderwal et al., 2017). In addition, we showed that the enhanced reliability and individual subject detection during natural viewing conditions are driven by task-specific reorganization of FC, which is distinct from the resting-state related individual variations.

Before GSR, the first principal component reflected the overall synchronization in FC (as well as global signal variance). This was an expected result, since the global signal was consistently reported as explaining the large amount of variation in BOLD signals (Carbonell et al., 2011). However, our results showed that the global signal component is not related to the movie condition. After GSR the global synchronization component disappeared, whereas the movie-specific component remained intact. It is important to note that the global signal component reflects the common FC topology across subjects and it was strongly correlated across sessions. The high correlations between the projections of global signal components across sessions suggest that the global signal component also explains individual variations. In other words, the variations and similarities across subjects during resting-state are mostly driven by the differences in global signal variance. However, in this study, it cannot be known to what extend the global signal-related component and individual variations reflect neural, hemodynamic and/or artefactual differences. The relationship between global signal

255 fluctuations and whole-brain synchronization is essential to understand individual subject
256 variations at resting-state.
257
258 **Stable task-specific FC patterns in dynamic FC**
259
We also investigated how movie-specific FC topography relates to the dynamic FC. We used
Hilbert transform on narrowband filtered time-series and characterized dynamic FC based on
phase-locking values across time. We found that the movie-specific principal components also
appear in FC and average PLVs of the narrowband filtered time-series. Then, we calculated
the similarity between grand-average PLVs of resting state and movie during one session and
the PLV fluctuations during the other session. We found that for movie sessions the PLV of
each time point was persistently more similar to the average PLVs of the movie sessions than
to those of the resting state sessions. In contrast, for resting sessions the PLV of each time
point was more similar to the resting state FC than to the movie FC. Overall, these results
showed that during natural viewing condition the FC is persistently reorganized into its
associated mode. One limitation of this approach is the substantial observation noise of
instantaneous PLVs in conjunction with limited variance explained by the PCs, which leads to
very low correlations between instantaneous PLVs. The stability of task-related FC patterns is
important for the assumptions behind large-scale computational modeling. Our results suggest
that whole-brain FC (at the time-scale of BOLD signals) is persistently reconfigured into a
distinct mode rather than an epiphenomenon reflecting activation of several transient states.
276
277 **Computational modeling of task-specific alterations**
278
Although variations related to the natural viewing condition can be explained in a single
dimension, it is very difficult interpret a whole-brain connectivity pattern. We proposed a
computational model to link the alterations in local dynamics to reorganization of whole-brain
FC. We used Hopf normal model to characterize BOLD signals. The motivation behind using
this model was that noise-driven and oscillatory dynamics can be modeled using a single
parameter (local bifurcation parameter). When the local bifurcation parameter of a particular
region is negative, the region exhibits noise-driven dynamics. For positive bifurcation parameter
values, the region exhibits sustained oscillations. Therefore, higher parameters values of a
region in the model indicate that the region has higher influence on its connected regions. The
model revealed significant decreases in bifurcation parameters particularly in cingulate cortex
(anterior and posterior cingulate) and in supramarginal gyrus. In contrast, the bifurcation
parameters significantly increased in lateral prefrontal cortex, medial temporal cortex and
superior parietal regions. These results showed that during movie condition, the influence of
frontal, temporal and parietal association regions on the whole-brain FC is enhanced.
Nevertheless, it is important to note that the model describes the BOLD signals in the
associated low-frequency narrow-band. Therefore, higher values of the bifurcation parameters

can also be interpreted as the contribution of low-frequency fluctuations in the regions is
enhanced since hemodynamic responses act as a low-pass filter. Furthermore, BOLD signals
are known to have negative relationship with the neural activity in middle frequency ranges
(Schölvinck et al., 2013). Therefore, the increased/decreased local bifurcation parameters
should be interpreted only in relation to low-frequency fluctuations.
300
Since the empirical data provided a clear separation of rest and movie conditions, we also
employed PCA on the bifurcation parameters of the model. The first PC mode showed a typical
pattern associated to the default mode network (DMN), in which the high values were observed
in isthmus cingulate, precuneus, medial frontal and temporal cortices. The second principal
component mode, characterized by higher bifurcation parameter values in temporal and frontal
regions, showed movie-specificity. These results suggested that the variations in bifurcation
parameters in frontal-parietal and frontal-temporal networks have exclusive contributions to the
organization of whole-brain FC during movie-viewing condition. Furthermore, we found that the
projections of the first and second principal components on movie sessions were negatively
correlated (spearman rank r = -0.496, p-value= 0.02). Therefore, the variations across both
task-specific and DMN-like activation patterns, and the antagonistic relationship between these
patterns are associated to task-related organization of FC. Nevertheless, based on these
results, it is not possible to draw conclusions on the causal mechanisms that drive the
relationship between DMN and task-related networks. The results may indicate that several
regions of DMN (particularly precuneus) having a role in mediating the switch between distinct
functional states, which is consistent with previous studies showing that precuneus dynamically
binds to distinct functional networks (Utevsky et al., 2014). An alternative explanation may
involve the variations of arousal and vigilance levels. This explanation is consistent with a
selective neuromodulatory enabling of intrinsic synaptic connections by ascending modulatory
neurotransmitter systems (e.g., noradrenaline). This is particularly relevant in light of the
systematic changes in the local bifurcation parameter that showed regionally-specific and
condition-sensitive effects in our modelling analyses.
323
324 **Limitations**
325
One limitation of our modeling results is that the model relies on DWI-derived SC, which has
limited performance on detecting interhemispheric connections, individual variations, and also
directions of the connections. In addition, the changes in directed effective connectivity may
also play role in defining the reorganization of FC during task (Gilson et al., 2017). We re-
analyzed the alterations in EC (Gilson et al., 2017) based on our current findings: the PCA
analysis of EC exhibited a better and clearer separation between resting-state and movie
sessions than FC (Supplementary Figure 4). Furthermore, the movie-specific principal
component of EC revealed enhanced connectivity from frontal regions towards parietal and
occipital brain regions, and from occipital and temporal regions toward parietal and frontal

regions (Supplementary Figure 3). Therefore, during natural viewing condition lower sensory
regions in occipital and temporal regions project to frontal and parietal higher-order association
regions, and that this pattern is completed as a recurrent loop by the frontal regions projecting
towards parietal regions and parietal regions projecting towards temporal and occipital regions.
These results are consistent with the alterations that we found in the local dynamics (i.e. under
fixed-connectivity assumption) such that frontal and parietal regions play crucial role in re-
organization of FC during task. In contrast, the model based on local dynamics cannot resolve
the alterations in sensory regions. Effective connectivity – as assessed using dynamic causal
modelling studies of the resting state – again point to a modulation of regional excitability by
different components of the default mode. For example, previous studies revealed that the
influence of the SN (salience network) and DAN (dorsal attention network) on the DMN (default
mode network) regions is inhibitory; whereas the DMN exerted an excitatory influence on the
SN and DAN regions (Zhou et al., 2018). Therefore, we speculate that the alterations in higher-
order association regions are better captured by the local dynamics because these regions
contribute to whole-brain organization through strong bidirectional connections across the
cortex, whereas the influence of sensory regions is dominantly unidirectional.

Conclusion and future directions

Current experimental paradigms are optimal to study task-dependent changes in BOLD signals,
but these are not optimal to study task-dependent re-organization of whole-brain FC.
Naturalistic condition, such as movie watching, that is comparable to the resting-state, may
have important implications on understanding the dynamic organization of whole-brain activity.
Here, we proposed a novel approach to link task-dependent functional organization and
dynamic functional connectivity. Nevertheless, it may not be possible to exploit the full potential
of dynamic organization of the brain through natural viewing paradigms. The major limitation of
this approach is that the natural viewing condition is a trivial task. Future studies may explore
other natural experimental paradigms that involve more challenging conditions such as problem
solving, navigation, social interactions, task engagement during present distractors. We
speculate that sophisticated natural tasks that require enhanced cognitive control may reveal
richer dynamical manifestation of functional reorganization. For example, certain challenging
tasks may show a better picture of dynamic reorganization of the whole-brain such as
consolidation of particular functional states in time (i.e. slow adaptation) and/or emergence of
observable transient functional states (i.e. multistability).

Materials and Methods

Study design

The fMRI imaging data used in this paper have been described in details elsewhere (Betti et al., 2013; Mantini et al., 2012). Twenty-four right-handed young, healthy volunteers (15 females, 20–31 years old) participated in the study. They were informed about the experimental procedures, which were approved by the Ethics Committee of the Chieti University, and signed a written informed consent. The study included a resting state and a natural vision condition. In the resting state, participants fixated a red target with a diameter of 0.3 visual degrees on a black screen. In the natural-vision condition, subjects watched (and listened) to 30 minutes of the movie "The Good, the Bad and the Ugly" in a window of 24x10.2 visual degrees. Visual stimuli were projected on a translucent screen using an LCD projector, and viewed by the participants through a mirror tilted by 45 degrees. Auditory stimuli were delivered using MR-compatible headphones.

Data acquisition

Functional imaging was performed with a 3T MR scanner (Achieva; Philips Medical Systems, Best, The Netherlands) at the Institute for Advanced Biomedical Technologies in Chieti, Italy. The functional images were obtained using T2*-weighted echo-planar images (EPI) with BOLD contrast using SENSE imaging. EPIs comprised of 32 axial slices acquired in ascending order and covering the entire brain (32 slices, 230 x 230 in-plane matrix, TR/TE=2000/35, flip angle = 90°, voxel size=2.875×2.875×3.5 mm3). For each subject, 2 and 3 scanning runs of 10 minutes duration were acquired for resting state and natural vision, respectively. Each run included 5 dummy volumes – allowing the MRI signal to reach steady state, and an additional 300 functional volumes that were used for analysis. Eye position was monitored during scanning using a pupil-corneal reflection system at 120 Hz (Iscan, Burlington, MA, USA). A three-dimensional high-resolution T1-weighted image, for anatomical reference, was acquired using an MP-RAGE sequence (TR/TE=8.1/3.7, voxel size=0.938x0.938x1 mm3) at the end of the scanning session.

Data preprocessing

Data preprocessing was performed using SPM5 (Wellcome Department of Cognitive Neurology, London, UK) running under MATLAB (The Mathworks, Natick, MA). The preprocessing steps involved the following: (1) correction for slice-timing differences (2) correction of head-motion across functional images, (3) coregistration of the anatomical image and the mean functional image, and (4) spatial normalization of all images to a standard stereotaxic space (Montreal Neurological Institute, MNI) with a voxel size of 3×3×3 mm3. Furthermore, the BOLD time series in MNI space were subjected to spatial independent component analysis (ICA) for the identification and removal of artifacts related to blood pulsation, head movement and instrumental spikes (Smith et al., 2010). This BOLD artifact

removal procedure was performed by means of the GIFT toolbox (Medical Image Analysis Lab,

University of New Mexico). No global signal regression or spatial smoothing was applied.

For each recording session (subject and run), we extracted the mean BOLD time series from

the 66 regions of interest (ROIs) of the brain atlas (Hagmann et al., 2008)(see Supplementary

Table 1). 2 subjects were excluded due to signal dropout and 1 subject was excluded due to

substantial spikes in the time-series.

Anatomical Connectivity

Anatomical connectivity was estimated from Diffusion Spectrum Imaging (DSI) data collected

in five healthy right-handed male participants (Hagmann et al., 2008; Honey et al., 2009). The

grey matter was first parcellated into 66 ROIs, using the same low-resolution atlas used for the

FC analysis. For each subject, we performed white matter tractography between pairs of

cortical areas to estimate a neuroanatomical connectivity matrix. The coupling weights between

two brain areas were quantified using the fiber tract density, and were proportional to a

normalized number of detected tracts. The structural matrix (SC) was then obtained by

averaging the matrices over subjects.

Principal component analysis

For all subjects and sessions (i.e. 21 subjects, 2 resting state and 2 movie sessions) the

functional connectivity matrices were constructed based on Pearson correlation coefficient

between all pairs of ROIs.

The upper triangular parts of FC (i.e. $66(66-1)/2$ connections) matrices were concatenated

across subjects/sessions (21x4 session/subjects) leading to the feature matrix with dimensions

2145 x 84 (number of connections x number of sessions/subjects). Then, principal component

analysis was applied to the resulting feature matrix. The analyses were repeated for 1000

surrogates time-series with preserved power-spectrum based on each session/subject. The

dimensionality of the data was characterized by explained variance of the principal components

that are larger than those of the surrogates.

To quantify the consistency of principal components, we repeated the analysis using 2 separate

sessions. For both sessions, the feature matrices comprised the concatenated upper triangular

FC matrices of 1 resting state session and 1 movie session (i.e. 2145 x 42 matrices). The

consistency across sessions was quantified as Pearson correlation coefficients of the

components and the projection of components between sessions.

Since during natural viewing condition the individuals are shown to have restricted movements

and increase arousal (Vanderwal et al., 2015), the differences in FC can be substantially

affected by underlying artifacts. For this reason, we repeated all the analyses after regressing

out global signal from the time-series of each ROI.

Dynamic functional connectivity

Preprocessed time series were band-pass filtered in 0.04-0.07Hz range in order to reduce the

effects of low-frequency drift and high-frequency noise (Glerean et al., 2012). Then, Hilbert
transform was used for the assessment of dynamic functional connectivity (Demirtaş et al.,
2016; Glerean et al., 2012). The Hilbert transform, $S(t) = A\cos(\varphi(t))$ of the preprocessed
BOLD time series broke the signal down to an analytical signal *S(t)* with an instantaneous phase
φ(t) and amplitude *A*. For each time point *t*, the difference $\Delta\varphi_{ij}(t)$ between the phases of the
respective ROIs was calculated, where *i* and *j* are the indices of each ROI. The phase
differences were adjusted between 0 and π such that:
458

$$\Delta\varphi_{ij}(t) = \begin{matrix} |\varphi_i(t) - \varphi_j(t)|, & if \; |\varphi_i(t) - \varphi_j(t)| \le \pi \\ 2\pi - |\varphi_i(t) - \varphi_j(t)|, & otherwise \end{matrix} \qquad 1$$

Then, the phase-locking values (PLVs), $\mathrm{PLV}_{ij}(t)$ were constructed using the phase differences
normalized between 0 and 1, thereby representing perfect anti-synchronization and perfect
synchronization respectively, such that: $\mathrm{PLV}_{ij}(t) = 1 - \Delta\varphi_{ij}(t)/\pi.$
We computed the grand-average PLVs for each session (i.e. resting state and movie watching).
Then, for each subject, we calculated the similarity (Pearson correlation coefficient) between
grand-average PLVs and time-resolved PLVs. In brief, for each subject two quantities were
estimated and compared as a function of time: the correlation between instantaneous PLVs
and grand-average PLVs for resting state and that for movie condition. To avoid the bias caused
by shared variability within each session, we calculated the grand-average PLVs of one
session, and then the similarities were calculated for the other session. Therefore, the analysis
was performed twice: the grand-average PLVs of session 1 projected to the PLVs of session 2
and the grand-average PLVs of session 2 projected to the PLVs of session 1. To provide a
better illustration for the magnitudes of correlations, we performed a same analysis averaging
the time-resolved PLVs across subjects (Supplementary Figure 1).
The principal component analysis was repeated for band-pass filtered time-series and average
PLVs. Since the PLVs were much more sensitive to global synchronization levels, we
subtracted the mean from each average PLV matrix. We also repeated the time-resolved
analysis using movie-specific and non-specific principal component instead of grand-average
PLVs. After identifying the principal component whose projections distinguish the resting
condition from movie condition, we re-projected each component on the instantaneous PLVs
using Pearson correlation coefficient (Supplementary Figure 1). We also performed PCA over
the concatenated time courses (i.e. 2 resting state and 2 movie sessions) of individual subject
PLVs. The trajectories of the first 2 principal components were plotted to illustrate the switch
between resting and movie watching conditions (Supplementary Figure 2).
485

Computational modeling
We modeled the whole-brain rs-fMRI BOLD signals using 66 nodes. Each node was coupled
with each other via DWI-derived structural connectivity (SC) matrix. We described the local
dynamics of each individual node using normal form of a supercritical Hopf bifurcation (Deco

et al., 2017). The advantage of this model is that it allows transitions between asynchronous
noise activity and oscillations. Where ω is the intrinsic frequency of each node, a is the local
bifurcation parameter, η is additive Gaussian noise with standard deviation β, the temporal
evolution of the activity, z, in node j is given in complex domain as:

494
$$\frac{dz_j}{dt} = \left[a_j + i\omega_j - |z_j^2|\right] + \beta\eta_j(t) \qquad 2$$

496 and,

497
$$z_j = \rho_j e^{i\theta_j} = x_j + iy_j \qquad 3$$

499 This system shows a supercritical bifurcation at $a_j = 0$. Being specific, if a_j is smaller than 0, the
500 local dynamics has a stable fixed point at $z_j = 0$, and for a_j values larger than 0, there exists a
501 stable limit cycle oscillation with a frequency $f = \omega/2\pi$. Finally, the whole-brain dynamics is
502 described by the following coupled equations:

503
$$\frac{dx_j}{dt} = [a_j - x_j^2 - y_j^2]x_j - \omega_j y_j + g \sum_i C_{ij}(x_i - x_j) + \beta\eta_{xj}(t) \qquad 4$$

505
$$\frac{dy_j}{dt} = [a_j - x_j^2 - y_j^2]y_j + \omega_j x_j + g \sum_i C_{ij}(y_i - y_j) + \beta\eta_{yj}(t) \qquad 5$$

Where C_{ij} is the Structural Connectivity (SC) between nodes i and j, g is the global coupling
factor, and the standard deviation of Gaussian noise, $\beta = 0.02$. The natural frequency (f) of
each region was taken as the peak frequency in the given narrowband of the corresponding
region in the empirical time-series.
Following a similar approach previously employed on biophysically-based computational model
(Deco et al., 2014), we analytically estimated the model FC using linearization of the system
around a stable fix point. Where $\delta\mathbf{u} = \{\delta x_1 \dots \delta x_{66}, \delta y_1 \dots \delta y_{66}\}$ represents the Taylor expansion
of the system, **A** is the Jacobian matrix, and $\varepsilon(t)$ is the noise term, the fluctuations around the
fix point can be described as:

516
$$\frac{d\delta\mathbf{u}}{dt} = \mathbf{A}\delta\mathbf{u} + \varepsilon(t) \qquad 6$$

517 Where the deterministic parts of right-hand side of equations 4 and 5 are described by $-F_j$ and
518 $-G_j$, respectively, the Jacobian matrix of the system evaluated at the fixed point $\widehat{x_j}, \widehat{y_j}, j \in$
519 $\{1 \dots 66\}$ can be constructed as:

$$\mathbf{A} = \begin{bmatrix} \dfrac{\partial F_1}{\partial x_1} & \cdots & \dfrac{\partial F_1}{\partial x_N} & \dfrac{\partial F_1}{\partial y_1} & \cdots & \dfrac{\partial F_1}{\partial y_N} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \dfrac{\partial F_N}{\partial x_1} & \cdots & \dfrac{\partial F_N}{\partial x_N} & \dfrac{\partial F_N}{\partial y_1} & \cdots & \dfrac{\partial F_N}{\partial y_N} \\ \dfrac{\partial G_j}{\partial x_1} & \cdots & \dfrac{\partial G_1}{\partial x_N} & \dfrac{\partial G_1}{\partial y_1} & \cdots & \dfrac{\partial G_N}{\partial y_1} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \dfrac{\partial G_N}{\partial x_1} & \cdots & \dfrac{\partial G_N}{\partial x_N} & \dfrac{\partial G_1}{\partial y_N} & \cdots & \dfrac{\partial G_N}{\partial y_N} \end{bmatrix}$$

521 Where $i, j \in \{1 \dots 66\}$, each element of matrix **A** can be calculated as:

522
$$\frac{\partial F_j}{\partial x_j} = \frac{\partial G_j}{\partial y_j} = a - g \sum_k C_{jk} \qquad\qquad 7$$

524
$$\frac{\partial F_j}{\partial x_l} = \frac{\partial G_j}{\partial y_l} = g C_{jl} \qquad\qquad 8$$

526
$$\frac{\partial F_j}{\partial y_j} = -\omega_0 \qquad\qquad 9$$

528
$$\frac{\partial G_j}{\partial y_j} = \omega_0 \qquad\qquad 10$$

530
$$\frac{\partial F_j}{\partial x_l} = \frac{\partial G_j}{\partial y_l} = 0 \qquad\qquad 11$$

532 Where **Q** is the noise covariance matrix, the covariance matrix of the system **P** can be estimated

533 by solving Lyapunov equation:

534
$$\mathbf{AP} + \mathbf{PA^T} = -\mathbf{Q} \qquad\qquad 12$$

537 Finally, the model correlation matric (FC) can be extracted from the covariance matrix as:

538
$$mFC_{ij} = \frac{P_{ij}}{\sqrt{P_{ii}P_{jj}}}, i, j \in \{1 \dots 66\} \qquad\qquad 13$$

We estimated the model optimal parameters a and g by maximizing the similarity between

model FC (equation 13) and empirical FC using gradient descent optimization. To avoid the

solutions reflecting a local minimum, for each subject/session we estimated the best solution

after repeating the optimization with 100 random initial conditions. The similarity between model

FC and empirical FC was quantified as Pearson correlation coefficient. For each subject, the

empirical functional connectivity was calculated as the average FC across the corresponding

conditions (i.e. resting state or movie sessions) of the corresponding subject.

The group comparisons for resting state versus movie sessions (optimal bifurcation parameters

and global coupling parameter) were done using permutation t-test (5000 permutations). For

optimal bifurcation parameters, the p-values were corrected using FDR approach with
Benjamini&Hochberg algorithm if necessary (Hochberg and Benjamini, 1990).
551

552 **Acknowledges**
553
554 We thank Alan Anticevic, John Murray, Markus Helmer and Joshua Burt for the insightful
555 discussions and their comments during the preparation of this paper.
556

References
Allen, E.A., Damaraju, E., Plis, S.M., Erhardt, E.B., Eichele, T., Calhoun, V.D., 2012. Tracking
Whole-Brain Connectivity Dynamics in the Resting State. Cereb. Cortex bhs352.
https://doi.org/10.1093/cercor/bhs352
Baars, B.J., 1993. A cognitive theory of consciousness., A cognitive theory of consciousness.
Cambridge University Press, New York, NY, US.
Betti, V., Della Penna, S., de Pasquale, F., Mantini, D., Marzetti, L., Romani, G.L., Corbetta,
M., 2013. Natural scenes viewing alters the dynamics of functional connectivity in the human
brain. Neuron 79, 782–797. https://doi.org/10.1016/j.neuron.2013.06.022
Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor
cortex of resting human brain using echo-planar MRI. Magn Reson Med 34, 537–541.
Carbonell, F., Bellec, P., Shmuel, A., 2011. Global and System-Specific Resting-State fMRI
Fluctuations Are Uncorrelated: Principal Component Analysis Reveals Anti-Correlated
Networks. Brain Connect 1, 496–510. https://doi.org/10.1089/brain.2011.0065
Cole, M.W., Bassett, D.S., Power, J.D., Braver, T.S., Petersen, S.E., 2014. Intrinsic and Task-
Evoked Network Architectures of the Human Brain. Neuron 83, 238–251.
https://doi.org/10.1016/j.neuron.2014.05.014
Cole, M.W., Ito, T., Bassett, D.S., Schultz, D.H., 2016. Activity flow over resting-state networks
shapes cognitive task activations. Nat Neurosci 19, 1718–1726.
https://doi.org/10.1038/nn.4406
Craddock, R.C., Holtzheimer, P.E., Hu, X.P., Mayberg, H.S., 2009. Disease state prediction
from resting state functional connectivity. Magn. Reson. Med. 62, 1619–1628.
https://doi.org/10.1002/mrm.22159
de Pasquale, F., Della Penna, S., Snyder, A.Z., Lewis, C., Mantini, D., Marzetti, L., Belardinelli,
P., Ciancetta, L., Pizzella, V., Romani, G.L., Corbetta, M., 2010. Temporal dynamics of
spontaneous MEG activity in brain networks. Proceedings of the National Academy of Sciences
107, 6040–6045. https://doi.org/10.1073/pnas.0913863107
Deco, G., Kringelbach, M.L., Jirsa, V.K., Ritter, P., 2017. The dynamics of resting fluctuations
in the brain: metastability and its dynamical cortical core. Scientific Reports 7.
https://doi.org/10.1038/s41598-017-03073-5
Deco, G., Ponce-Alvarez, A., Hagmann, P., Romani, G.L., Mantini, D., Corbetta, M., 2014. How
Local Excitation-Inhibition Ratio Impacts the Whole Brain Dynamics. The Journal of

Neuroscience 34, 7886–7898. https://doi.org/10.1523/JNEUROSCI.5068-13.2014

Demirtaş, M., Tornador, C., Falcón, C., López-Solà, M., Hernández-Ribas, R., Pujol, J.,
Menchón, J.M., Ritter, P., Cardoner, N., Soriano-Mas, C., Deco, G., 2016. Dynamic functional
connectivity reveals altered variability in functional connectivity among patients with major
depressive disorder. Hum. Brain Mapp. 37, 2918–2930. https://doi.org/10.1002/hbm.23215

Engel, A.K., Fries, P., Singer, W., 2001. Dynamic predictions: Oscillations and synchrony in
top–down   processing.   Nature   Reviews   Neuroscience   2,   704–716.
https://doi.org/10.1038/35094565

Finn, E.S., Shen, X., Scheinost, D., Rosenberg, M.D., Huang, J., Chun, M.M., Papademetris,
X., Constable, T.R., 2015. Functional connectome fingerprinting: identifying individuals using
patterns of brain connectivity 18, 1664–1671. https://doi.org/10.1038/nn.4135

Fries, P., 2005. A mechanism for cognitive dynamics: neuronal communication through
neuronal   coherence.   Trends   in   Cognitive   Sciences   9,   474–480.
https://doi.org/10.1016/j.tics.2005.08.011

Gilson, M., Deco, G., Friston, K.J., Hagmann, P., Mantini, D., Betti, V., Romani, G.L., Corbetta,
M., 2017. Effective connectivity inferred from fMRI transition dynamics during movie viewing
points   to   a   balanced   reconfiguration   of   cortical   interactions.   NeuroImage.
https://doi.org/10.1016/j.neuroimage.2017.09.061

Glerean, E., Salmi, J., Lahnakoski, J.M., Jaaskelainen, I.P., Sams, M., 2012. Functional
Magnetic Resonance Imaging Phase Synchronization as a Measure of Dynamic Functional
Connectivity. Brain Connect 2, 91–101. https://doi.org/10.1089/brain.2011.0068

Gonzalez-Castillo, J., Hoy, C.W., Handwerker, D.A., Robinson, M.E., Buchanan, L.C., Saad,
Z.S., Bandettini, P.A., 2015. Tracking ongoing cognition in individuals using brief, whole-brain
functional connectivity patterns. Proceedings of the National Academy of Sciences 112, 8762–
8767. https://doi.org/10.1073/pnas.1501242112

Hagmann, P., Cammoun, L., Gigandet, X., Meuli, R., Honey, C.J., Wedeen, V.J., Sporns, O.,
2008. Mapping the Structural Core of Human Cerebral Cortex. PLoS Biol 6, e159.
https://doi.org/10.1371/journal.pbio.0060159

Hochberg, Y., Benjamini, Y., 1990. More powerful procedures for multiple significance testing.
Statist. Med. 9, 811–818. https://doi.org/10.1002/sim.4780090710

Honey, C.J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J.P., Meuli, R., Hagmann, P.,
2009. Predicting human resting-state functional connectivity from structural connectivity. PNAS
106, 2035–2040. https://doi.org/10.1073/pnas.0811168106

Kim, D., Kay, K., Shulman, G.L., Corbetta, M., 2017. A New Modular Brain Organization of the
BOLD   Signal   during   Natural   Vision.   Cerebral   Cortex   1–17.
https://doi.org/10.1093/cercor/bhx175

Mantini, D., Corbetta, M., Romani, G.L., Orban, G.A., Vanduffel, W., 2012. Data-driven analysis
of analogous brain networks in monkeys and humans during natural vision. NeuroImage 63,
1107–1118. https://doi.org/10.1016/j.neuroimage.2012.08.042

Ponce-Alvarez, A., He, B.J., Hagmann, P., Deco, G., 2015. Task-Driven Activity Reduces the

Cortical Activity Space of the Brain: Experiment and Whole-Brain Modeling. PLOS
Computational Biology 11, e1004445. https://doi.org/10.1371/journal.pcbi.1004445

Rosenberg, M.D., Finn, E.S., Scheinost, D., Papademetris, X., Shen, X., Constable, R.T.,
Chun, M.M., 2015. A neuromarker of sustained attention from whole-brain functional
connectivity. Nature Neuroscience 19, 165–171. https://doi.org/10.1038/nn.4179

Schölvinck, M.L., Leopold, D.A., Brookes, M.J., Khader, P.H., 2013. The contribution of
electrophysiology to functional connectivity mapping. NeuroImage, Mapping the Connectome
80, 297–306. https://doi.org/10.1016/j.neuroimage.2013.04.010

Smith, J.F., Pillai, A., Chen, K., Horwitz, B., 2010. Identification and validation of effective
connectivity networks in functional magnetic resonance imaging using switching linear dynamic
systems. NeuroImage 52, 1027–1040. https://doi.org/10.1016/j.neuroimage.2009.11.081

Tavor, I., Jones, O.P., Mars, R.B., Smith, S.M., Behrens, T.E., Jbabdi, S., 2016. Task-free MRI
predicts individual differences in brain activity during task performance. Science 352, 216–220.
https://doi.org/10.1126/science.aad8127

Tononi, G., 2004. An information integration theory of consciousness. BMC Neuroscience 5,
42. https://doi.org/10.1186/1471-2202-5-42

Utevsky, A.V., Smith, D.V., Huettel, S.A., 2014. Precuneus Is a Functional Core of the Default-
Mode Network. J Neurosci 34, 932–940. https://doi.org/10.1523/JNEUROSCI.4227-13.2014

Vanderwal, T., Eilbott, J., Finn, E.S., Craddock, R.C., Turnbull, A., Castellanos, F.X., 2017.
Individual differences in functional connectivity during naturalistic viewing conditions.
NeuroImage 157, 521–530. https://doi.org/10.1016/j.neuroimage.2017.06.027

Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L.C., Castellanos, F.X., 2015. Inscapes : A movie
paradigm to improve compliance in functional magnetic resonance imaging. NeuroImage 122,
222–232. https://doi.org/10.1016/j.neuroimage.2015.07.069

Zhou, Y., Friston, K.J., Zeidman, P., Chen, J., Li, S., Razi, A., 2018. The Hierarchical
Organization of the Default, Dorsal Attention and Salience Networks in Adolescents and Young
Adults. Cerebral Cortex 28, 726–737. https://doi.org/10.1093/cercor/bhx307

656
657

Figures and captions

Figure 1. Conceptual overview. **A** Mean FC across subjects during resting state (top) and movie (bottom) sessions. For convenience the matrices were ordered by pre-defined networks involving visual, auditory, motor, integration and frontal regions. **B-C** Overview of principal component analysis (PCA). The FCs of the subjects at each session were concatenated into a single feature matrix. Then PCs of the feature matrix was calculated **(B)**. Note that the (vectorized) FC features pertain to correlations over the number of regions squared. However, the principal components of these FC features can be reorganized into a connectivity matrix that has the same size as the number of connections – as shown on the right (of panel B).The projections of each subject/session of the first two principal components were plotted against each other **(C)**. The second principal component projections revealed a clear distinction between rest and movie sessions (top) leading to representation of FC variations in two dimensions (bottom).

Figure 2. Principal component analysis of FC based on 2 resting state and 2 movie sessions

concatenated across 21 subjects. **A-D.** Results without global signal regression (GSR). **A**

Explained variance by each PC (black) and random surrogates (gray) without GSR. Compared

to 1000 random surrogates the dimensionality of FCs without GSR is 13. The first PC **(B)**

explains 25.8% of the variation, whereas second PC **(C)** explains 7.2% of the variation. The

projections of first two PCs reveals that the second component is specific to movie sessions

(D). The first PC of the FCs without GSR reflects global signal standard deviation **(J)**. **E-H.**

Results with global signal regression (GSR). **E** Explained variance by each PC (black) and

random surrogates (gray) with GSR. Compared to random surrogates the dimensionality of

FCs with GSR is 22. The first PC, which is specific to movie sessions explains 9.69% of the

variation **(F)**. The movie-specific components with and without GSR is highly consistent. *** p-

value < 0.0001.

Figure 3. The consistency of principal components across sessions. The second PC is specific

to movie session in both session 1 **(A)** and session 2 **(B)** without GSR, whereas the first PC is

movie-specific in session 1 **(C)** and session 2 **(D)** with GSR. The consistency of the movie-

specific component (**F** without GSR; **G** with GSR) is substantially higher than that of the non-

specific components (**E** without GSR; **H** with GSR). Similarly, the consistency of the projections

for movie-specific component (**J** without GSR; **K** with GSR) is substantially higher than that of

the non-specific components (**I** without GSR; **L** with GSR). *** p-value < 0.0001.

Figure 4. Stability of movie-specific components in dynamic FC. **A** The schematic describing
the derivation of phase-locking value (PLV) dynamics. **B** Kernel density estimates of the
correlations across PLVs. **C** Components projections of session 1. **D** Correlations between
average PLVs at rest (black), during movie condition (red) and PLVs of resting state session 2.
E Correlations between average PLVs at rest (black), during movie condition (red) and PLVs
of movie session 2. **F-H** The same procedure as C-E, where the average PLVs were calculated
for session 2 and projected on session 1 PLVs. The similarity between average resting-state
PLVs and instantaneous PLVs of resting state sessions are significantly higher than those of
movie sessions (Wilcoxon signed-rank test, p-value < 0.0001) **(D,G)**. Conversely, The similarity
between average movie PLVs and instantaneous PLVs of movie sessions are significantly
higher than those of resting state sessions (Wilcoxon signed-rank test, p-value < 0.0001) **(E,H)**.
Shaded regions indicate standard error of mean. *** indicates p-values < 0.0001.

Figure 5. Large-scale computational modeling. **A** The schematic of the modeling framework.

The BOLD activity of each region was described using Hopf normal model, where the local

bifurcation parameters (a) mediate the local dynamics. Negative values of bifurcation

parameter, a, indicates noise-driven activity, whereas positive values indicate oscillatory activity

with increasing amplitude. Brain regions are coupled each other through DWI-derived SC

matrix. The optimal model parameters were estimated using gradient descent optimization,

which maximizes the similarity between empirical and model FC. **B** Mean optimal bifurcation

parameter topography at resting state. **C** Mean optimal bifurcation parameter topography

during movie condition. **D** The distributions of the bifurcation parameters during movie condition

and resting state.

Figure 6. Modeling results for individual subject fitting. **A** The schematic of individual subject fitting. **B** The group differences for global coupling parameters did not show significant difference. **C-D** The group differences between optimal bifurcation parameters at rest (black) and during movie condition (red) (permutation t-test, 5000 permutations). **C** The topography of the group differences (T-statistics; hot colors indicate larger values during movie condition). **D** Boxplots of the regions showing significantly difference after FDR correction. **E-G** Principal component analysis applied to optimal bifurcation parameters in the model. **E** The topography of the first principal component. **F** The topography of the second principal component. PC-1 has higher values in precueus, posterior cingulate, medial temporal and frontal regions, exhibiting typical pattern associated to default mode network. PC-2 exhibit increased values in frontal and temporal regions, and decreased values particularly in supramarginal gyri and in medial brain regions. **G** The projections of the principal components on rest and movie sessions. *** indicate p-value < 0.0001.

 Supplementary Figures

Supplementary Figure 1. A-D The similarity between mean PLVs and instantaneous PLVs

of each condition. The mean-PLVs were computed for resting-state and movie sessions, and

then the correlation coefficient between condition-specific mean-PLVs and instantaneous

PLVs (averaged across subjects) were calculated. Since higher correlations are expected

between average and instantaneous PLVs for the same sessions, the analyses were done

cross-sessions: We calculated the correlation between mean-PLVs of rest/movie sessions 1

and instantaneous PLVs of rest/movie sessions 2, and vice versa. E-F The similarity between

movie-specific and non-specific principal components and instantaneous PLVs.

749
750 **Supplementary Figure 2.** The trajectories extracted from the PCA on dynamic functional
751 connectivity of each subject. Gray colors indicate the time course of the components at rest,
752 red colors indicate time course of the components while watching movie.
753

Supplementary Figure 3. The visualization of the highest 1% of the connections of PC-1 **(A)**

and PC-2 **(B)** of FC, and PC-1 of EC **(C-D)**. **A** The non-specific component of FC exhibits

larger connectivity strengths across posterior cingulate and precuneus, medial temporal and

frontal regions as well as occipital cortex, which suggests default-mode network connectivity.

B The movie-specific component of FC shows high connectivity between occipital and

parietal regions, and between temporal and frontal regions. For effective connectivity (EC),

movie-specific component reveals enhanced connectivity from frontal regions towards parietal

and occipital brain regions **(C)**, and enhanced connectivity from occipital and temporal

regions toward parietal and frontal regions **(D)**.

Supplementary Figure 4. The PCA on functional connectivity (left) and effective connectivity

(right). The separation between resting-state and movie conditions are clearer in EC.

Furthermore, there is a strong correlation between the projections of movie-specific and non-

specific components only for movie sessions for effective connectivity.

770