Title: Efficacy and acceptability of non-invasive brain stimulation for the treatment of adult

unipolar and bipolar depression: A systematic review and meta-analysis of randomised sham-

controlled trials

Authors: Julian Mutz[1] MSc, Daniel R. Edgcumbe[2] MSc, Andre R. Brunoni[3,4] MD PhD,

Cynthia H.Y. Fu[2,5] MD PhD

Affiliations:

1. Department of Epidemiology and Biostatistics, School of Public Health, Faculty of

Medicine, Imperial College London, United Kingdom

2. School of Psychology, College of Applied Health and Communities, University of East

London, United Kingdom

3. Service of Interdisciplinary Neuromodulation, Laboratory of Neurosciences (LIM-27) and

National Institute of Biomarkers in Psychiatry (INBioN), Department and Institute of

Psychiatry, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de

Sao Paulo, Sao Paulo, SP, Brazil.

4. Department of Psychiatry and Psychotherapy, Ludwig-Maximilians-University, Munich,

Germany.

5. Centre for Affective Disorders, Department of Psychological Medicine, Institute of

Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom

Author for Correspondence:

Julian Mutz, Department of Epidemiology and Biostatistics, School of Public Health, Faculty

of Medicine, Imperial College London, United Kingdom; Norfolk Place, London W2 1PG,

United Kingdom. Email: julian.mutz@gmail.com; phone: +44(0)2075943686.

Abstract

We examined the efficacy and acceptability of non-invasive brain stimulation in adult

unipolar and bipolar depression. Randomised sham-controlled trials of transcranial direct

current stimulation (tDCS), transcranial magnetic stimulation (TMS) and theta-burst

stimulation (TBS), without co-initiation of another treatment, were included. We analysed

response, remission and all-cause discontinuation rates, and depression severity scores. Fifty-

four studies were included (N = 2,959, mean age = 44.94 years, 61.98% female). Response

rates demonstrated efficacy of high-frequency rTMS over the left DLPFC (OR = 3.94, 95%

CI [2.52; 6.15]), right-sided low-frequency rTMS (OR = 7.44, 95% CI [2.06; 26.83]) bilateral

rTMS (OR = 3.68, 95% CI [1.66; 8.13]), deep TMS (OR = 1.69, 95% CI [1.003; 2.85]),

intermittent TBS (OR = 4.70, 95% CI [1.14; 19.38]) and tDCS (OR = 4.32, 95% CI [2.02;

9.29]); but not for continuous TBS, bilateral TBS or synchronised TMS. There were no

differences in all-cause discontinuation rates. The strongest evidence was for high-frequency

rTMS over the left DLPFC. Intermittent TBS provides an advance in terms of reduced

treatment duration. tDCS is a potential treatment for non-resistant depression.

Keywords: transcranial magnetic stimulation, theta burst stimulation, transcranial direct

current stimulation, major depression, meta-analysis

Introduction

Major depression is prevalent[1] and associated with considerable disease burden[2]. Its course is

often recurrent and may become chronic with relapse rates within one year of remission

ranging from 35% to 80%[3,4]. The most common treatments are pharmacological and

psychological therapies. Yet, even with a full course of treatment, at least one third of patients

fail to achieve remission[5]. Non-invasive neurostimulation therapies, such as transcranial

magnetic stimulation (TMS) and transcranial electrical stimulation (tES), offer a potential

alternative or add-on treatment strategy.

TMS was originally introduced as a tool for investigating and mapping cortical functions and

connectivity[6]. TMS utilises intense, rapidly-changing electromagnetic fields, which are

generated by a coil of wire near the scalp, and allows for a mostly undistorted induction of an

electrical current to alter neural activity in relatively focal, superficial areas of the brain.

Standard TMS typically involves single or paired pulses, and repetitive transcranial magnetic

stimulation (rTMS) involves the delivery of repeated pulses which enable the prolonged

modulation of neural activity. Depending on the stimulation frequency, rTMS can increase or

decrease cortical excitability. The prevailing hypothesis is that high-frequency (usually 10 Hz

or higher) stimulation is excitatory and causes neural depolarisation, whereas low-frequency

(≤ 1 Hz) stimulation inhibits neural firingRosa and Lisanby [7].

The rationale for using rTMS to treat depressive illness comes from clinical symptomatology

and neuroanatomy as well as neuroimaging studies indicating functional impairments in

prefrontal cortical and limbic regions[8]. In 2008, the US Food and Drug Administration (FDA)

approved the first rTMS device for the treatment major depressive disorder (MDD) in which

there was poor response to at least one pharmacological agent in the current episode[9], and its

clinical utilisation has increased since[10].

As stimulation at high frequencies can be uncomfortable during the initial stimulation period,

low-frequency rTMS may minimise the occurrence of undesired side effects, namely

headaches and scalp discomfort, and may be associated with fewer adverse events, for

instance by lowering the risk for seizures[11].

Bilateral applications of rTMS have also been developed: simultaneous stimulation over the

left and right DLPFC (rDLPFC) or stimulation over one side followed by stimulation of the

other side. These applications were hypothesised to be potentially additive or synergistic to

reinstate any imbalance in prefrontal neural activity[12]. Moreover, there may be a selective

unilateral response and the likelihood for a clinical response may increase by providing both

types of stimulation[13].

Technical and methodological efforts to improve the antidepressant efficacy of TMS have led

to several alternative treatment protocols. Deep TMS (dTMS) was FDA-approved in 2013,

which is able to stimulate larger brain volumes and deeper structures[14] that could be more

directly relevant in the pathophysiology of depression (e.g., reward-mediating pathways and

areas connected to the subgenual cingulate cortex)[8,15,16].

Another recent modification is theta burst stimulation (TBS)[17], which is a patterned form of

TMS pulse delivery that utilises high and low frequencies in the same stimulus train. TBS

delivers bursts of three at a high frequency (50 Hz) with an inter-burst interval of 5Hz in the

theta range at 5 Hz. Two different protocols are utilised: continuous theta burst stimulation

(cTBS), which delivers 300 or 600 pulses without interruption, and intermittent theta burst

stimulation (iTBS), which delivers 30 pulses every 10 seconds for a duration of 190 seconds,

totalling 600 pulsesChung, et al. [18]. It is suggested that cTBS reduces cortical excitability

while iTBS increases it, mimicking the processes of long-term potentiation and long-term

depression, respectively[17]. Notably, there is some debate as to whether prolonged stimulation

periods reverse the hypothesised effects of TBSGamboa, et al. [19], while there is also support

for a dose-response relationship for iTBS[20].

The main advantages of TBS are its reduced administration time, which is typically less than

five minutes as opposed to 20–45 minutes for conventional rTMS, and the lower intensity

needed to produce lasting neurophysiological effects, as TBS is typically administered at 80%

of the resting motor threshold (rMT) and might be more comfortable than stimulation at

higher intensities typically used with standard rTMS.

Synchronised TMS refers to magnetic low-field synchronised stimulation (sTMS), a new

treatment paradigm that involves rotating spherical rare-earth (neodymium) magnets

positioned sagittally along the midline of the scalp, which deliver stimulation synchronised to

an individual's alpha frequency[21]. The magnets are positioned to provide a global magnetic

field distributed broadly across the midline cortical surface (one magnet over the frontal polar

region, one magnet over the top of the head, and one magnet over the parietal region). The

rationale for sTMS synchronised to an individual's alpha frequency is the observation that

one mechanism of action of rTMS is the entrainment of oscillatory activity to the

programmed frequency of stimulation, thereby resetting thalamo-cortical oscillators and

restoring normal endogenous oscillatory activity[22]. This modification of TMS may be

associated with fewer treatment-emergent adverse and side effects because it does not cause

neural depolarisation. It also uses less energy than conventional rTMS as it utilises sinusoidal

instead of pulsed magnetic fields, which require less than 1% of the energy needed for

conventional rTMS and may thus be less expensive.

Access and costs are among the major impediments to a more widespread use of rTMS,

although costs may be lower for TBS and sTMS. A less expensive technique is transcranial

electrical stimulation (tES). Its most commonly used protocol, transcranial direct current

stimulation (tDCS), was reappraised as a tool in research through the work of Priori, et al. [23]

and Nitsche and Paulus [24]. tDCS involves the application of a low-amplitude electrical direct

current through surface scalp electrodes to superficial areas of the brain. While it does not

directly trigger action potentials, it modulates cortical excitability by shifting the neural

membrane resting potential and these effects can outlast the electrical stimulation period[25].

The direction of such excitability changes may depend on the polarity of the stimulation:

anodal stimulation is hypothesised to cause depolarisation and an increase in neural

excitability, whereas cathodal stimulation causes hyperpolarisation and a decrease in cortical

excitability[26,27].

The advantages of tDCS compared to TMS include its ease of administration, being much less

expensive, its more benign side effect profile, and its portability which could potentially be

used in the home environment.

We sought to perform a systematic review and meta-analysis of the antidepressant efficacy

and acceptability of non-invasive neuromodulation in treating a current depressive episode in

unipolar and bipolar depression from randomised sham-controlled trials. The only study to

date that evaluated the efficacy of a range of rTMS techniques is Brunoni, et al. [28] network

meta-analysis[28]. However, the analysis had included trials that had co-initiated other

treatments (e.g. sleep deprivation and TMS); trials which had not included a sham treatment;

had not separated the TBS modifications; and had not included any age-related exclusion

criteria. Also, tDCS trials were not included in that meta-analysis. We sought to address these

limitations by including only trials with randomised allocation to active or sham treatments,

excluding studies which had co-initiated another treatment, and limiting our sample to the

adult age range as geriatric depression may impact on efficacy.

Materials and Methods

Search strategy and selection criteria

We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

(PRISMA) guidelines[29]. A systematic search of the Embase, Medline, and PsychINFO

databases was performed from the first date available to 1[st] November 2017 (Figure 1,

Supplementary Materials).

Inclusion criteria were: 1) adults aged 18 – 70 years; 2) DSM or ICD diagnosis of MDD or

bipolar disorder currently in a major depressive episode; 3) randomised sham-controlled

trials, which utilised a parallel-group or cross-over design; 4) clinician-administered

depression rating scale, Hamilton Depression Rating Scale (HDRS; Hamilton [30]) or

Montgomery-Åsberg Depression Rating Scale (MADRS; Montgomery and Åsberg [31]).

Exclusion criteria were: 1) primary diagnoses other than MDD or bipolar depression; 2)

studies limited to a specific subtype of depression (e.g., postpartum depression or vascular

depression) or in which a major depressive episode was a secondary diagnosis (e.g.,

fibromyalgia and major depression); 3) co-initiation of any other form of treatment, such as

pharmacotherapy or cognitive control training.

Data analysis

The following sample characteristics were extracted: sex, age, hospitalisation status, whether

patients with psychotic symptoms were excluded from the study, diagnosis, treatment

strategy, and treatment resistance.

The following treatment-related parameters were extracted. For TMS: type of coil and sham

procedure, magnet location, stimulation frequency (Hz) for each site, stimulation intensity

(percentage of the rMT), total number of pulses delivered, and number of treatment sessions.

For TBS: data on the treatment protocol (iTBS, cTBS or bilateral TBS) were also recorded.

For tDCS: location of the anode and cathode, electrode size (cm^2), current intensity (mA) and

density (mA/cm^2), session duration, number of sessions, and duration of active stimulation in

the sham condition.

The primary outcome measure was clinical response, defined as a $\geq$ 50% reduction in

symptom scores at the primary study endpoint. Remission rates were the secondary outcome

measure based on the definition provided by each study. If response or remission rates were

reported for both HDRS and MADRS, data for the HDRS was selected to facilitate

comparability between trials. If data for multiple versions of the HDRS were reported, the

original 17-item version was selected. We extracted baseline and post-treatment depression

scores. If available, the intention-to-treat (ITT) or modified intention-to-treat (mITT) data

were preferred over data based only on completers. For cross-over trials, only data from the

initial randomisation were used to avoid carry-over effects. Data presented in figures were

extracted with WebPlotDigitizer (http://arohatgi.info/WebPlotDigitizer/app/). All-cause

discontinuation rates were recorded separately for active and sham groups.

Data that could not be directly retrieved from the original publications were requested from

the authors or searched for in previous systematic reviews and meta-analyses. For trials with

more than two groups that could not be included as separate treatment comparisons, we

combined groups to create single pair-wise comparisons.

Analyses were conducted using the 'meta' package[32] for RStudio (Version 0.98.932) and

STATA (Version 13.1; StataCorp, 2013) was used for data processing.

Contour-enhanced funnel plots[33] were visually inspected to assess whether potential funnel

asymmetry is likely to be due to statistical significance-based publication bias.

A random-effects model was chosen as it was assumed that the underlying true effect size

would vary between studies. A random-effects model provides wider confidence intervals

than a fixed-effects model if there is significant heterogeneity among studies and thus tends to

be more conservative in estimating summary effect sizes.

For dichotomous outcome data, odds ratios (Mantel-Haenszel method) were used as an index

of effect size. We also computed Hedge's g to estimate the effect sizes for continuous post-

treatment depression scores.

Heterogeneity between studies was assessed with the Q_T statistic, which estimates whether the

variance of effect sizes is greater than what would be expected due to sampling error. A p

value smaller than .01 provides an indication for significant heterogeneity[34]. The I^2 statistic

was computed for each analysis to provide a descriptive measure of inconsistency across the

results of individual trials included in our analyses. It provides an indication of what

percentage of the observed variance in effect sizes reflects real differences in effect sizes as

opposed to sampling error. Higgins, et al. [35] suggested that 25%, 50%, and 75% represent

little, moderate, and high heterogeneity, respectively.

The Cochrane tool for assessing risk of bias in randomised trials[36] was used to evaluate

included studies. Each trial received a score of low, high, or unclear risk of bias for each of

the potential sources of bias. Two raters independently conducted the assessment of risk of

bias.

Results

Fifty-four RCTs, consisting of 127 treatment arms, met our inclusion criteria (Figure 1,

Supplementary Materials). Overall, 64 treatment comparisons were included, total $N = 2,959$

patients (mean age = 44.94 years, 61.98% female) of whom $n = 1,548$ were randomised to

active and $n = 1,411$ to sham treatments (Tables 1-4).

Visual inspection of the contour-enhanced funnel plots did not suggest small study effects

(Figure 2; Supplementary Figure 1-2). However, due to the small number of studies, we only

computed funnel plots for all neurostimulation techniques combined, which might mask small

study effects pertaining to a specific treatment modality.

Sixty comparisons of experimental and sham treatment arms met the inclusion criteria for the

meta-analysis of response rates (Table 5; Figure 3), and 46 treatment comparisons for the

meta-analysis of remission rates (Table 6; Figure 4).

High-frequency rTMS over the left DLPFC (lDLPFC) was associated with improved rates of

response as well as remission in comparison with sham treatment. The odds ratio of response

was OR = 3.94 compared to sham ($k = 31$, 95% CI [2.52; 6.15]). There was little evidence

that the heterogeneity between trials exceeded that expected by chance ($I^2 = 27.1\%$; $Q_{30} =$

41.15, $p = .08$). Sensitivity analyses suggested similar effect sizes in trials that had recruited

patients with unipolar depression only and those that had recruited both patients with unipolar

and bipolar depression (Supplementary Figure 3a). Only one pilot study[37] had recruited

patients with bipolar depression only, but provided no support for antidepressant efficacy (OR

= 1.14, 95% CI [0.21; 6.37]). Response rates were greater in trials that (i) excluded patients

with psychotic features, (ii) recruited outpatients only, and (iii) recruited either treatment

resistant patients only or both treatment resistant patients and those that were not treatment

resistant (Supplementary Figure 3b,c,d).

The odds of achieving remission were over twice that of sham ($k = 25$, OR = 2.74, 95% CI

[1.75; 4.28]). There was no evidence for significant heterogeneity ($I^2 = 0.0\%$; $Q_{24} = 22.67$, $p =$

.54). Sensitivity analyses for remission rates were in line with those for response rates,

although we did not find left-sided high-frequency rTMS to be effective in samples that had

recruited both treatment resistant and non-treatment resistant patients (Supplementary Figure

6a,b,c,d).

Low-frequency rTMS over the rDLPFC was also associated with significantly greater

response and remission rates than sham stimulation. There was a sevenfold improvement in

response rates compared to sham ($k = 3$, OR= 7.44 (95% CI [2.06; 26.83]), with no indication

for significant heterogeneity between trials ($I^2 = 0.0\%$; $Q_2 = 1.59$, $p = .45$). No sensitivity

analyses were conducted due to the small number of treatment comparisons.

The odds of remission were greater than those of sham ($k = 2$, OR = 14.10 (95% CI [2.79;

71.42]). Heterogeneity between trials was not greater than expected due to sampling error (I^2

$= 0.0\%$; $Q_1 = 0.50$, $p = .48$). No sensitivity analyses were conducted due to the small number

of treatment comparisons.

Low-frequency rTMS over the lDLPFC was not associated with any significant

improvements in rates of response or remission. There were no significant differences in

response rates compared to sham ($k = 3$, OR = 1.41, 95% CI [0.15; 12.88]). The heterogeneity

between trials did not exceed that expected by chance ($I^2 = 0.0\%$; $Q_2 = 0.14$, $p = .93$), and no

sensitivity analyses were conducted due to the small number of treatment comparisons. There

were no significant differences in remission rates compared to sham ($k = 3$, OR = 0.86, 95%

CI [0.08; 9.11]). The variance in effect sizes between trials was no greater than expected due

to sampling error ($I^2 = 0.0\%$; $Q_2 = 0.03$, $p = .98$). No sensitivity analyses were conducted due

to the small number of treatment comparisons.

Bilateral rTMS was associated with significant improvement in response but not remission

rates compared to sham. There was a significant improvement in response rates compared to

sham ($k = 6$, OR = 3.68 (95% CI [1.66; 8.13]), and the variance in effect sizes between trials

did not exceed that expected due to sampling error ($I^2 = 0.0\%$; $Q_5 = 3.45$, $p = .63$). Sensitivity

analyses suggested subgroup differences according to whether trials had excluded psychotic

patients or had recruited patients with diagnosis of MDD only, bipolar depression only, or

both MDD and bipolar depression (Supplementary Figure 4a,b). We found no evidence for a

significant improvement in rates of remission associated with bilateral TMS compared to

sham ($k = 5$, OR = 3.05, 95% CI [0.87; 10.67]). There was no evidence for significant

heterogeneity between trials ($I^2 = 10.7\%$; $Q_4 = 4.48$, $p = .34$), and sensitivity analyses

suggested no differences according to any patient characteristics tested (Supplementary

Figure 7a,7b).

There were significant improvements in both response and remission rates for dTMS

compared to sham. The response rates were marginally higher while statistically significant

for dTMS relative to sham ($k = 2$, OR = 1.69, 95% CI [1.003; 2.85]). The variance in effect

sizes between trials did not exceed that expected due to sampling error ($I^2 = 0.0\%$; $Q_1 = 0.97$,

$p = .33$). No sensitivity analyses were conducted due to the small number of treatment

comparisons. The remission rates were greater for dTMS compared to sham ($k = 2$, OR =

2.24, 95% CI [1.24; 4.06]). There was no evidence for significant heterogeneity between trials

$(I^2 = 0.0\%; Q_1 = 0.02, p = 0.88)$, and no sensitivity analyses were conducted due to the small

number of treatment comparisons.

Neither response nor remission rates for sTMS were significantly higher than for sham. There

was no evidence for increased response rates compared to sham ($k = 2$, OR = 2.71, 95% CI

[0.44; 16.86]). There was significant heterogeneity between these two studies ($I^2 = 75.9\%$;

$Q_1 = 4.15$, $p = .04$). No sensitivity analyses were conducted due to the small number of

treatment comparisons. There were also no significant improvements in remission rates for

sTMS compared to sham ($k = 2$, OR = 2.51 (95% CI [0.23; 26.76]). There was evidence for

significant heterogeneity between the two studies though ($I^2 = 75.7\%$; $Q_1 = 4.12$, $p = .04$). No

sensitivity analyses were conducted due to the small number of treatment comparisons.

iTBS over the lDLPFC was associated with a fivefold improvement in response rates

compared to sham ($k = 2$, OR = 4.70 (95% CI [1.14; 19.38]). The heterogeneity between trials

did not exceed that expected by chance ($I^2 = 0.0\%$; $Q_1 = 0.02$, $p = .89$). No sensitivity

analyses were conducted due to the small number of treatment comparisons. For only one

trial[38] was data on remission rates for iTBS available, with no evidence for antidepressant

efficacy compared to sham.

Neither cTBS over the rDLPFC nor bilateral TBS were statistically different from sham in

terms of response rates ($k = 1$, OR = 1.63, 95% CI [0.23; 11.46] and $k = 2$, OR = 4.28, 95% CI

[0.54; 34.27]). For bilateral TBS there was evidence that the variance in effect sizes between

studies was greater than what would be expected due to sampling error ($I^2 = 65.7\%$; $Q_1 =$

2.91, $p = .09$). No sensitivity analyses were conducted due to the small number of treatment

comparisons. The only trial of bilateral TBS for which remission rates were available[39] found

no evidence for its antidepressant efficacy compared to sham. No remission rates were

available for cTBS.

tDCS was associated with significant improvement in both response and remission rates in

comparison to sham stimulation. There was a significant improvement in response rates

relative to sham (k = 8, OR = 4.32, 95% CI [2.02; 9.29]). There was little evidence for

significant heterogeneity between studies (I^2 = 34.2%; Q_7 = 10.64, p = .16) and sensitivity

analyses suggested tDCS to be effective only in patients with non-treatment resistant

depression and in trials that had recruited patients with both treatment resistant and non-

treatment resistant depression (Supplementary Figure 5).

The analysis of remission rates showed a statistically significant advantage of tDCS compared

to sham (k = 7, OR = 3.07, 95% CI [1.58; 5.99]). There was no indication for significant

heterogeneity between trials (I^2 = 4.4%; Q_6 = 6.27, p = .39), and sensitivity analyses found

that only trials that had recruited patients with both treatment resistant and non-treatment

resistant depression provided evidence for antidepressant efficacy.

Forty-six treatment comparisons reported post-intervention continuous depression scores.

There was evidence for the antidepressant efficacy of high-frequency rTMS over the lDLPFC

compared to sham (k = 28, Hedge's g = -0.75, 95% CI [-1.02; -0.47]), dTMS compared to

sham (k = 2, Hedge's g = -0.29, 95% CI [-0.55; -0.03]), and tDCS compared to sham (k = 6,

Hedge's g = -0.76, 95% CI [-1.42; -0.10]). There was evidence for significant heterogeneity

between trials for several treatment modalities (Table 7; Figure 5).

Sixty-four treatment comparisons were available for all-cause discontinuation rates. There

were no significant differences in drop-out rates for any treatment modalities (Table 8; Figure

6).

Discussion

The present systematic review and meta-analysis examined the efficacy and acceptability of non-invasive brain stimulation techniques for a current depressive episode in unipolar and bipolar depression. We sought to investigate the efficacy of the brain stimulation techniques without the potential confound of co-initiation of another treatment and in trials which had included randomised allocation to a sham stimulation treatment arm in order to account for potential placebo effects.

The largest evidence base to date is for high-frequency rTMS over the lDLPFC which is associated with 3.94 times greater odds of response than sham stimulation as well as odds of remission that are 2.74 times greater than sham. These findings are consistent with previous systematic reviews and meta-analysesBerlim, et al. [40] and have led to the consensus review and treatment guideline by the *Clinical TMS Society* for daily high-frequency rTMS over the lDLPFC for the treatment of medication-resistant or medication-intolerant depressive episodes[41].

Additional support for treatment efficacy was revealed for low-frequency rTMS over the rDLPFC, which was associated with improved rates of response as well as remission. Bilateral rTMS was associated with higher rates of response but not remission. It is unclear whether any advantages of bilateral rTMS compared to left-sided high-frequency or right-sided low-frequency rTMS would be due to the treatment protocol. As bilateral stimulation delivers a greater number of pulses than unilateral stimulation, unless the number of treatment sessions or the treatment duration are adjusted for accordingly, it is difficult to reliably assess whether the difference in stimulation protocol (bilateral vs. unilateral stimulation) or the difference in the number of stimuli delivered leads to differences in clinical effects[42].

To date, no studies have directly compared dTMS and standard rTMS protocols. In an

exploratory meta-analysis of nine open-label trials, including a total of 150 patients, Kedzior,

et al. [43] provided evidence for the antidepressant efficacy of dTMS. The present meta-analysis

found that dTMS was associated with 1.69 times greater odds of response and 2.24 greater

odds of remission than sham which were statistically significant. While the open-label trials

included in Kedzior, et al. [43] may have overestimated the true efficacy of dTMS, we provide

initial support for the clinical efficacy of dTMS that was greater than for sham treatment but

less than for high-frequency rTMS over the lDLPFC, low-frequency rTMS over the rDLPFC

or bilateral rTMS.

The meta-analytic estimates did not indicate significant treatment effects associated with low-

frequency rTMS over the lDLPFC or with sTMS. However, these have been trialled in only

three[44-46] and two studies[21,47], respectively. Specific treatment effects of TMS that depend on

side and frequency of stimulation have been proposed but it may be possible that low-

frequency rTMS over the lDLPFC has a marginal effect in at least a small number of

patients[46]. Leuchter, et al. [47] found sTMS to only be effective when administered at the

individual's alpha frequency and with a minimum of 80% treatment adherence, suggesting a

dose-response relationship.

With theta burst stimulation, the duration of each treatment session is reduced to a few

minutes. Our meta-analysis did demonstrate almost five times greater odds of response

compared to sham for iTBS over the lDLPFC. However, this estimate is based on two trials

only. One trial had examined remission rates as well[38], reporting remission rates of 0% for

sham and 9.1% for active stimulation. The meta-analytic estimates for cTBS and the bilateral

modification of TBS did not show any advantage over sham in terms of response rates. The

only trial that reported remission rates for bilateral TBS did not provide evidence for its

antidepressant efficacy either and no data were available to evaluate remission rates following

cTBS.

Transcranial direct current stimulation is a form of neurostimulation that offers greater

portability and lower costs relative to TMS. The meta-analysis revealed significant

improvements in response and remission rates following tDCS treatment in comparison to

sham, which was 4.32 times greater for response rates and 3.07 times greater for remission

rates. We have been able to identify the effects of tDCS without potential confounds of co-

initiation of another treatment, revealing significantly greater odds of response as well as

remission[48]. The clinical efficacy of tDCS is evident also in the non-treatment resistant form

of depression, in contrast to most rTMS trials, suggesting that tDCS is a potential initial

therapeutic option for depression.

The finding that there were no differences in terms of drop-out rates at study end between the

active treatment and sham conditions for any treatment modality suggests that non-invasive

brain stimulation is generally well tolerated by patients. We chose all-cause discontinuation

rates based on the intention-to-treat sample, representing the most conservative estimate of

treatment acceptability.

We chose response and remission rates as our main outcome measures, which arguably

constitute clinically-useful estimates of the antidepressant efficacy of non-invasive brain

stimulation techniques. However, response and remission rates were not reported for each

trial, and some missing data could not be obtained. Studies have also suggested that the

antidepressant efficacy of active stimulation may separate from sham only after multiple

weeks of treatment, for both rTMSO'Reardon, et al. [9,]Chistyakov, et al. [49] and cTBS[49]. We

only looked at the acute antidepressant effects at primary study endpoint, and we cannot

estimate the long-term effects.

A significant number of TMS studies used active magnetic stimulation with the coil being

angulated at 45 or 90 degrees to the scalp surface as sham condition. Because differences in

coil orientation may produce considerably different sensations on the scalp and coil

angulation might still produce a limited degree of intracortical activity[50], ensuring a valid

control condition constitutes a methodological challenge. One study placed an inactive coil on

the patient's head while discharging an active coil at least one meter away in order to mimic

the auditory effects of rTMS[51].

A more recent approach is to use a specifically designed sham coil that does not generate a

magnetic field but is visually and auditorily indistinguishable from an active coil. A meta-

analysis by Berlim, et al. [52] found no significant differences between the number of patients

who correctly guessed their treatment allocation when comparing active high-frequency left-

sided or bilateral rTMS and sham. There were also no significant differences between studies

that utilised angulated coils and sham coils. Blinding integrity is less of a methodological

hurdle for sTMS trials because neither active stimulation nor sham procedure produce any

physical sensation, they look identical, and are comparable in terms of acoustic artefacts.

Only few of the more recent modifications of TMS reported on the adequacy of their blinding

procedure. Given that cross-over designs are particularly prone to unblinding after cross-over,

we included only data corresponding to the initial randomisation in our analyses.

For tDCS, the sham condition typically involves delivering active stimulation for up to 30

seconds, which mimics the initial somatic sensations without inducing a therapeutic effect.

However, the adequacy of blinding of tDCS sham has also been called into question[53].

The clinical trials had enrolled patients based on a diagnostic assessment of clinical symptoms

rather than underlying brain pathology. The potential for biological heterogeneity might mask

the clinical efficacy of non-invasive brain stimulation in some trials but could not be assessed

in the present analysis. We implemented reasonably strict inclusion criteria to limit the

influence of a range of potential confounders, for example we excluded RCTs that co-initiated

treatment with medication. However, potential effects of specific medications on the clinical

efficacy of brain stimulation could not be adequately controlled for as patients often had a

large number of heterogeneous treatments prior to enrolling, which might have distorted the

clinical effects of brain stimulation.

Finally, compared to the network meta-analysis on TMS[28], we were not able to compare the

active treatments. In the NMA priming rTMS seemed most effective. However, the two RCTs

that used this treatment modality compared it with another active stimulation and could not be

included in the present meta-analysis.

Conclusion

The present systematic review and meta-analysis supports the efficacy and acceptability of

non-invasive brain stimulation techniques in adult unipolar and bipolar depression. The

strongest evidence was for high-frequency rTMS over the lDLPFC, followed by low-

frequency rTMS over the rDLPFC and bilateral rTMS. Intermittent TBS provides a potential

advance in terms of reduced treatment duration and the meta-analysis did find support for

improved rates of response. tDCS is a potential treatment for non-resistant depression which

has demonstrated efficacy in terms of response as well as remission. All the trials included in

the present meta-analysis had included randomised allocation to a sham treatment arm and we

had excluded trials in which there was co-initiation of another treatment. Some of the more

recent treatment modalities though require additional trials and more direct comparisons

between different treatment modalities are warranted.

Authorship contributions

J.M. and C.H.Y.F. conceived the project; J.M. performed the systematic literature search with

supervision by C.H.Y.F; J.M. extracted and analysed the data; D.R.E. confirmed the quality

of the extracted data; J.M. wrote the initial draft of the paper; C.H.Y.F. and A.R.B. critically

revised the paper, including interpretation of the data. All authors read and approved the final

version of this paper.

Funding and disclosure

The authors declare no conflict of interest.

Acknowledgments

We thank Dr Angelo Alonzo & Professor Colleen Loo, Professor Ian Anderson, Dr Lysianne

Beynel, Dr Noomane Bouaziz & Dr Dominique Januel, Dr Poul E Buchholtz, Dr Romain

Duprat & Professor Chris Baeken, Professor Mark S George, Professor Ehud Klein, Professor

Berthold Langguth & Dr Martin Schecklmann, Dr Giuseppe Lanza, Professor Andy Leuchter,

Dr Alessandra Minelli, Professor William M McDonald, Professor Declan McLoughlin, Dr

Marie-Laure Paillère Martinot, Dr Bill Phillips, Professor Rober M Post, Dr Mohammad Ali

Salehinejad, Dr Christos Theleritis, and Professor Abraham Zangen for providing additional

data. We are also grateful to those authors who could not provide additional data but

responded to our inquiry. The views expressed in this article represent those of the authors

and not necessarily those of the individuals who have provided additional data for the

analyses.

Supplementary material

Supplementary information is available online.

References

1　Kessler, R. C. *et al.* The epidemiology of major depressive disorder: results from the
National Comorbidity Survey Replication (NCS-R). *Jama* **289**, 3095-3105 (2003).

2　Murray, C. J. *et al.* Global, regional, and national disability-adjusted life years
(DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188
countries, 1990–2013: quantifying the epidemiological transition. *The Lancet* **386**,
2145-2191 (2015).

3　Fekadu, A. *et al.* What happens to patients with treatment-resistant depression? A
systematic review of medium to long term outcome studies. *Journal of affective*
disorders **116**, 4-11 (2009).

4　Eaton, W. W. *et al.* Population-based study of first onset and chronicity in major
depressive disorder. *Archives of General Psychiatry* **65**, 513-520 (2008).

5　Rush, A. J. *et al.* Acute and longer-term outcomes in depressed outpatients requiring
one or several treatment steps: a STAR* D report. *American Journal of Psychiatry*
163, 1905-1917 (2006).

6　Barker, A. T., Jalinous, R. & Freeston, I. L. Non-invasive magnetic stimulation of
human motor cortex. *The Lancet* **325**, 1106-1107 (1985).

7　Rosa, M. A. & Lisanby, S. H. Somatic treatments for mood disorders.
Neuropsychopharmacology **37**, 102-116 (2012).

8　Atkinson, L., Sankar, A., Adams, T. M. & Fu, C. H. Recent advances in neuroimaging
of mood disorders: structural and functional neural correlates of depression, changes
with therapy, and potential for clinical biomarkers. *Current Treatment Options in*
Psychiatry **1**, 278-293 (2014).

9　O'Reardon, J. P. *et al.* Efficacy and safety of transcranial magnetic stimulation in the
acute treatment of major depression: a multisite randomized controlled trial.
Biological psychiatry **62**, 1208-1216 (2007).

10　Janicak, P. G., Sackett, V., Kudrna, K. & Cutler, B. Advances in transcranial magnetic
stimulation for managing major depressive disorders: The utility of TMS for treating
depression continues to widen, as the technology is refined. *Current Psychiatry* **15**,
49-56 (2016).

11　Rossi, S., Hallett, M., Rossini, P. M., Pascual-Leone, A. & Group, S. o. T. C. Safety,
ethical considerations, and application guidelines for the use of transcranial magnetic
stimulation in clinical practice and research. *Clinical neurophysiology* **120**, 2008-2039
(2009).

12 Conca, A. *et al.* Combining high and low frequencies in rTMS antidepressive
treatment: preliminary results. *Human Psychopharmacology: Clinical and*
Experimental **17**, 353-356 (2002).

13 Fitzgerald, P. B. *et al.* A randomized, controlled trial of sequential bilateral repetitive
transcranial magnetic stimulation for treatment-resistant depression. *American Journal*
of Psychiatry **163**, 88-94 (2006).

14 Roth, Y., Amir, A., Levkovitz, Y. & Zangen, A. Three-dimensional distribution of the
electric field induced in the brain by transcranial magnetic stimulation using figure-8
and deep H-coils. *Journal of Clinical Neurophysiology* **24**, 31-38 (2007).

15 Greicius, M. D. *et al.* Resting-state functional connectivity in major depression:
abnormally increased contributions from subgenual cingulate cortex and thalamus.
Biological psychiatry **62**, 429-437 (2007).

16 Costafreda, S. G. *et al.* Modulation of amygdala response and connectivity in
depression by serotonin transporter polymorphism and diagnosis. *Journal of affective*
disorders **150**, 96-103 (2013).

17 Huang, Y.-Z., Edwards, M. J., Rounis, E., Bhatia, K. P. & Rothwell, J. C. Theta burst
stimulation of the human motor cortex. *Neuron* **45**, 201-206 (2005).

18 Chung, S., Hoy, K. & Fitzgerald, P. Theta-burst stimulation: a new form of TMS
treatment for depression? *Depression and anxiety* **32**, 182-192 (2015).

19 Gamboa, O. L., Antal, A., Moliadze, V. & Paulus, W. Simply longer is not better:
reversal of theta burst after-effect with prolonged stimulation. *Experimental brain*
research **204**, 181-187 (2010).

20 Nettekoven, C. *et al.* Dose-dependent effects of theta burst rTMS on cortical
excitability and resting-state connectivity of the human motor system. *The Journal of*
Neuroscience **34**, 6849-6859 (2014).

21 Jin, Y. & Phillips, B. A pilot study of the use of EEG-based synchronized Transcranial
Magnetic Stimulation (sTMS) for treatment of Major Depression. *BMC psychiatry* **14**,
1 (2014).

22 Leuchter, A. F., Cook, I. A., Jin, Y. & Phillips, B. The relationship between brain
oscillatory activity and therapeutic effectiveness of transcranial magnetic stimulation
in the treatment of major depressive disorder. *Frontiers in Human Neuroscience* **7**
(2013).

23 Priori, A., Berardelli, A., Rona, S., Accornero, N. & Manfredi, M. Polarization of the
human motor cortex through the scalp. *Neuroreport* **9**, 2257-2260 (1998).

24   Nitsche, M. A. & Paulus, W. Excitability changes induced in the human motor cortex
by weak transcranial direct current stimulation. *The Journal of physiology* **527**, 633-
639 (2000).

25   Nitsche, M. A. *et al.* Transcranial direct current stimulation: state of the art 2008.
Brain stimulation **1**, 206-223 (2008).

26   Nitsche, M. A. & Paulus, W. Sustained excitability elevations induced by transcranial
DC motor cortex stimulation in humans. *Neurology* **57**, 1899-1901 (2001).

27   Merzagora, A. C. *et al.* Prefrontal hemodynamic changes produced by anodal direct
current stimulation. *Neuroimage* **49**, 2304-2310 (2010).

28   Brunoni, A. R. *et al.* Repetitive transcranial magnetic stimulation for the acute
treatment of major depressive episodes: A systematic review with network meta-
analysis. *JAMA psychiatry* **74**, 143-152 (2017).

29   Moher, D., Liberati, A., Tetzlaff, J. & Altman, D. G. Preferred reporting items for
systematic reviews and meta-analyses: the PRISMA statement. *Annals of internal*
medicine **151**, 264-269 (2009).

30   Hamilton, M. A rating scale for depression. *Journal of Neurology, Neurosurgery &*
Psychiatry **23**, 56-62 (1960).

31   Montgomery, S. A. & Åsberg, M. A new depression scale designed to be sensitive to
change. *The British journal of Psychiatry* **134**, 382-389 (1979).

32   Schwarzer, G. Meta: An R package for meta-analysis. *R news* **7**, 40-45 (2007).

33   Peters, J. L., Sutton, A. J., Jones, D. R., Abrams, K. R. & Rushton, L. Contour-
enhanced meta-analysis funnel plots help distinguish publication bias from other
causes of asymmetry. *Journal of clinical epidemiology* **61**, 991-996 (2008).

34   Cochran, W. G. The combination of estimates from different experiments. *Biometrics*
10, 101-129 (1954).

35   Higgins, J. P., Thompson, S. G., Deeks, J. J. & Altman, D. G. Measuring
inconsistency in meta-analyses. *British Medical Journal* **327**, 557-560 (2003).

36   Higgins, J. P. *et al.* The Cochrane Collaboration's tool for assessing risk of bias in
randomised trials. *BMJ* **343**, 889-893 (2011).

37   Nahas, Z., Kozel, F. A., Li, X., Anderson, B. & George, M. S. Left prefrontal
transcranial magnetic stimulation (TMS) treatment of depression in bipolar affective
disorder: a pilot study of acute safety and efficacy. *Bipolar disorders* **5**, 40-47 (2003).

38   Duprat, R. *et al.* Accelerated intermittent theta burst stimulation treatment in
medication-resistant major depression: A fast road to remission? *Journal of affective*
disorders **200**, 6-14 (2016).

39   Prasser, J. *et al.* Bilateral prefrontal rTMS and theta burst TMS as an add-on treatment
for depression: a randomized placebo controlled trial. *The World Journal of Biological*
Psychiatry **16**, 57-65 (2015).

40   Berlim, M. T., Van den Eynde, F., Tovar-Perdomo, S. & Daskalakis, Z. Response,
remission and drop-out rates following high-frequency repetitive transcranial magnetic
stimulation (rTMS) for treating major depression: a systematic review and meta-
analysis of randomized, double-blind and sham-controlled trials. *Psychological*
medicine **44**, 225-239 (2014).

41   Perera, T. *et al.* The Clinical TMS Society Consensus Review and Treatment
Recommendations for TMS Therapy for Major Depressive Disorder. *Brain*
stimulation **9**, 336-346 (2016).

42   Chen, J.-j. *et al.* Bilateral vs. unilateral repetitive transcranial magnetic stimulation in
treating major depression: a meta-analysis of randomized controlled trials. *Psychiatry*
Research **219**, 51-57 (2014).

43   Kedzior, K. K., Gellersen, H. M., Brachetti, A. K. & Berlim, M. T. Deep transcranial
magnetic stimulation (DTMS) in the treatment of major depression: an exploratory
systematic review and meta-analysis. *Journal of Affective Disorders* **187**, 73-83
(2015).

44   Padberg, F. *et al.* Repetitive transcranial magnetic stimulation (rTMS) in
pharmacotherapy-refractory major depression: comparative study of fast, slow and
sham rTMS. *Psychiatry research* **88**, 163-171 (1999).

45   Kimbrell, T. A. *et al.* Frequency dependence of antidepressant response to left
prefrontal repetitive transcranial magnetic stimulation (rTMS) as a function of
baseline cerebral glucose metabolism. *Biological psychiatry* **46**, 1603-1613 (1999).

46   Speer, A. M., Wassermann, E. M., Benson, B. E., Herscovitch, P. & Post, R. M.
Antidepressant efficacy of high and low frequency rTMS at 110% of motor threshold
versus sham stimulation over left prefrontal cortex. *Brain stimulation* **7**, 36-41 (2014).

47   Leuchter, A. F. *et al.* Efficacy and safety of low-field synchronized transcranial
magnetic stimulation (sTMS) for treatment of major depression. *Brain stimulation* **8**,
787-794 (2015).

48   Meron, D., Hedger, N., Garner, M. & Baldwin, D. S. Transcranial direct current
646   stimulation (tDCS) in the treatment of depression: systematic review and meta-
647   analysis of efficacy and tolerability. *Neuroscience & Biobehavioral Reviews* **57**, 46-62
648   (2015).

649   49   Chistyakov, A. V. *et al.* Preliminary assessment of the therapeutic efficacy of
650   continuous theta-burst magnetic stimulation (cTBS) in major depression: a double-
651   blind sham-controlled study. *Journal of affective disorders* **170**, 225-229 (2015).

652   50   Lisanby, S. H., Gutman, D., Luber, B., Schroeder, C. & Sackeim, H. A. Sham TMS:
653   intracerebral measurement of the induced electrical field and the induction of motor-
654   evoked potentials. *Biological psychiatry* **49**, 460-463 (2001).

655   51   Loo, C. K., Mitchell, P. B., McFarquhar, T. F., Malhi, G. S. & Sachdev, P. S. A sham-
656   controlled trial of the efficacy and safety of twice-daily rTMS in major depression.
657   *Psychological medicine* **37**, 341-349 (2007).

658   52   Berlim, M. T., Broadbent, H. J. & Van den Eynde, F. Blinding integrity in randomized
659   sham-controlled trials of repetitive transcranial magnetic stimulation for major
660   depression: a systematic review and meta-analysis. *International Journal of*
661   *Neuropsychopharmacology* **16**, 1173-1181 (2013).

662   53   Blumberger, D. M., Tran, L. C., Fitzgerald, P. B., Hoy, K. E. & Daskalakis, Z. J. A
663   randomized double-blind sham-controlled study of transcranial direct current
664   stimulation for treatment-resistant major depression. *Frontiers in Psychiatry* **3**, 119-
665   126 (2012).

Figure captions

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)

flow diagram of literature search.

Figure 2. Contour-enhanced funnel plot of all RCTs included in the meta-analysis of response

rates.

Figure 3. Forest plot of response rates.

Figure 4. Forest plot of remission rates.

Figure 5. Forest plot of post-treatment continuous depression scores.

Figure 6. Forest plot of all-cause discontinuation rates.

Table 1

Treatment characteristics: TMS studies

Authors	Location	Frequency (Hz)		% rMT	Total pulses	Sessions	Treatment strategy	Active group	Sham group
HF-L		Left	Right						
Anderson et al., 2007	LDLPFC	10	-	110[a]	12,000	12	Mixed	Figure-of-eight	Sham-coil
Avery et al., 2006	LDLPFC	10	-	110[b]	24,000	15	Mixed	Figure-of-eight	90°
Avery et al., 1999	LDLPFC	10	-	80	NR	10	Mixed	NR	45°
Baeken et al., 2013[*]	LDLPFC	20	-	110	31,200	20	Monotherapy	Figure-of-eight	90°
Bakim et al., 2012[1]	LDLPFC	20	-	80; 100	24,000	30	Augmentation	Figure-of-eight	45°
Berman et al., 2000	LDLPFC	20	-	80	NR	10	Monotherapy	Figure-of-eight	30-45°
Bortolomasi et al., 2007	LDLPFC	20	-	90	4,000	5	Mixed	Circular	90°
Boutros et al., 2002	LDLPFC	20	-	80	8,000	10	Mixed	Figure-of-eight	90°
Chen et al., 2013	LDLPFC	20	-	90	NR	10	Augmentation	Figure-of-eight	90°
Concerto et al., 2015	LDLPFC	10	-	120	60,000	20	Augmentation	Figure-of-eight	45°
Eschweiler et al., 2000[*]	LDLPFC	10	-	90	NR	5	Augmentation	Figure-of-eight	90°

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Fitzgerald et al., 2012 (1)	LDLPFC	10	-	120	NR	15	Mixed	Figure-of-eight	45°
Fitzgerald et al., 2003 (1)	LDLPFC	10	-	100	10,000	10	Augmentation	Figure-of-eight	45°
Garcia-Toro et al., 2001	LDLPFC	20	-	90	NR	10	Augmentation	Figure-of-eight	90°
George et al., 2010	LDLPFC	10	-	120	45,000	15	Monotherapy	Figure-of-eight	Sham-coil
George et al., 2000[2]	LDLPFC	5; 20[c]	-	100[d]	16,000	10	Monotherapy	Figure-of-eight	45°
George et al., 1997[*]	LDLPFC	20	-	80	8000	10	Mixed	Figure-of-eight	45°
Hansen et al., 2004	LDLPFC	10	-	90	30,000	15	Augmentation	Figure-of-eight	90°
Hernández-Ribas et al., 2013	LDLPFC	15	-	100	22,500	15	Augmentation	Figure-of-eight	90°
Holtzheimer et al., 2004	LDLPFC	10	-	110	16,000	10	Monotherapy	Figure-of-eight	45°[oe]
Jakob et al., 2008 (1)	LDLPFC	20	-	100	20,000	10	Mixed	Figure-of-eight	Sham-coil
Jakob et al., 2008 (2)	LDLPFC	50	-	100	20,000	10	Mixed	Figure-of-eight	Sham-coil
Kimbrell et al., 1999[*]	LDLPFC	20	-	80	8,000	10	Monotherapy	Figure-of-eight	45°
Kreuzer et al., 2015	LDLPFC	10	-	110	30,000	15	Augmentation	Figure-of-eight	Sham-coil
Lingeswaran et al., 2011	LDLPFC	10	-	100	NR	12	NR	Figure-of-eight	90°
Loo et al., 1999[*]	LDLPFC	10	-	110	NR	10	Mixed	Figure-of-eight	90°

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Nahas et al., 2003	LDLPFC	5	-	110	16,000	10	Monotherapy	Figure-of-eight	45°
O'Reardon et al., 2007	LDLPFC	10	-	120[g]	60,000	20	Monotherapy	Figure-of-eight	Sham-coil
Paillère-Martinot et al., 2010	LDLPFC	10	-	90	16,000	10	Augmentation	Figure-of-eight	Sham-coil
Speer et al., 2014	LDLPFC	20	-	110	24,000	15	Monotherapy	Figure-of-eight	45°
Su et al., 2005[3]	LDLPFC	5; 20	-	100	16,000	10	Augmentation	Figure-of-eight	90°
Theleritis et al., 2017 (1)	LDLPFC	20	-	100	24,000	15	Mixed	Figure-of-eight	90°
Theleritis et al., 2017 (2)	LDLPFC	20	-	100	48,000	30[f]	Mixed	Figure-of-eight	90°
Zheng et al., 2010	LDLPFC	15	-	110[g]	60,000	20	Augmentation	Figure-of-eight	90°
LF-R									
Fitzgerald et al., 2003 (2)	RDLPFC	-	1	100	3,000	10	Augmentation	Figure-of-eight	45°
Januel et al., 2006	RDLPFC	-	1	90	1,920	16	Monotherapy	Figure-of-eight	Sham-coil
Pallanti et al., 2010 (1)	RDLPFC	-	1	110	6,300	15	Augmentation	Figure-of-eight	Sham-coil
LF-L									
Kimbrell et al., 1999[*]	LDLPFC	1	-	80	8,000	10	Monotherapy	Figure-of-eight	45°
Padberg et al., 1999	LDLPFC	0.3	-	90	1,250	5	Mixed	Figure-of-eight	90°

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Speer et al., 2014	LDLPFC	1	-	110	24,000	15	Monotherapy	Figure-of-eight	45°
BL									
Fitzgerald et al., 2006	DLPFC	10	1	110(R); 100(L)	7,200	10	Mixed	Figure-of-eight	45°
Fitzgerald et al., 2016	DLPFC	10	1	110	40,000	20	Mixed	Figure-of-eight	45°
Fitzgerald et al., 2012 (2)	DLPFC	10	1	120	NR	15	Mixed	Figure-of-eight	45°
McDonald et al., 2006[4]	DLPFC	10	1	110	16,000	10	Monotherapy	Figure-of-eight	90°
Pallanti et al., 2010 (2)	DLPFC	10	1	110(R); 100(L)	21,300	15	Augmentation	Figure-of-eight	Sham-coil
Prasser et al., 2015 (1)	DLPFC	10	1	110	30,000	15	Augmentation	Figure-of-eight	Sham-coil
iTBS									
Duprat et al., 2016[*]	LDLPFC	50	-	110	32,400	20[i]	Monotherapy	Figure-of-eight	Sham-coil
Li et al., 2014 (1)	LDLPFC	50	-	80[j]	18,000	10	Mixed	Figure-of-eight	90°
cTBS									
Li et al., 2014 (2)	RDLPFC	50	-	80[j]	18,000	10	Mixed	Figure-of-eight	90°
BLTBS									
Li et al., 2014 (3)	DLPFC	50	50	80[j]	36,000	10	Mixed	Figure-of-eight	90°

Prasser et al., 2015 (2)	DLPFC	50	50	80	36,000	15	Augmentation	Figure-of-eight	Sham-coil

dTMS

Levkovitz et al., 2015	LDLPFC	18	-	120[h]	39,600	20	Monotherapy	H1	Sham-coil
Tavares et al., 2017	LDLPFC	18	-	120	39,600	20	Augmentation	H1	Sham-coil

sTMS

Jin et al., 2014[5]	Midline	IAF; 8-13		-	-	20	Augmentation	sTMS	NMRS
Leuchter et al., 2015	Midline	IAF		-	-	30	Monotherapy	sTMS	NMRS

Note. Numbers in parentheses behind authors indicate that multiple active treatment arms of the same study are reported. Hz = hertz; rMT = resting motor threshold; LDLPFC = left dorsolateral prefrontal cortex; RDLPFC = right dorsolateral prefrontal cortex; TMS = transcranial magnetic stimulation; HF-L = high-frequency, left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency, right-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; iTBS = intermittent theta burst stimulation; cTBS = continuous theta burst stimulation; BLTBS = bilateral theta burst stimulation; dTMS = deep transcranial magnetic stimulation; sTMS = synchronised transcranial magnetic stimulation; IAF = individual alpha frequency; NMRS = non-magnetic rotating shaft; NR = not reported. [*]Cross-over design. [1-5]Two active treatment groups were combined. [a]Two patients received active stimulation at 100% rMT. [b]Stimulation delivered at estimated prefrontal threshold. [c]During the 5th session, stimulation was delivered for 2min at 10Hz. [d]During the 5th session, stimulation was delivered for 2min at 60% rMT. [e]Two patients received sham treatment with the coil angulated at 90°. [f]Received treatment twice daily. [g]During the first week, 110% rMT could be used for tolerability. [h]During the first three treatment session, rMT could be titrated from 100% to 120%. [i]Received treatment five times daily. [j]Stimulation delivered at active motor threshold.

Table 2

Sample characteristics: TMS studies

Authors	Number of participants (female)		Age		Diagnosis	HDRS / MADRS		Excluded psychosis	Status	Treatment resistance
	Active	Sham	Active	Sham		Active	Sham			
HF-L										
Anderson et al., 2007[1]	13 (7)	16 (9)	48.0 (8.0)	46.0 (12.0)	MDD	26.7 (3.6)[M]	27.7 (7.1)[M]	No	Outpatient	Mixed
Avery et al., 2006[2]	35 (21)	33 (16)	44.3 (10.3)	44.2 (9.7)	MDD	23.5 (3.9)[a]	23.5 (2.9)[a]	Yes	NR	TRD
Avery et al., 1999	4 (4)	2 (1)	44.3 (10.1)	45.0 (7.1)	Mixed	21.3 (6.7)[b]	19.5 (8.1)[b]	Yes	Outpatient	TRD
Baeken et al., 2013	9 (7)	11 (5)	51.8 (12.1)	47.3 (13.7)	MDD	24.8 (7.1)[a]	26.5 (8.7)[a]	Yes	Mixed	TRD
Bakim et al., 2012[3]	23 (20)	12 (11)	40.8 (10.0)	44.4 (10.2)	MDD	23.6 (3.6)[a]	25.6 (3.8)[a]	Yes	Outpatient	TRD
Berman et al., 2000[2]	10 (2)	10 (4)	45.2 (9.5)	39.4 (10.8)	Mixed	37.1 (9.7)[c]	37.3 (8.5)[c]	No	Mixed	TRD
Bortolomasi et al., 2007	12 (7)	7 (4)	NR	NR	Mixed	25.17 (7.84)[d]	21.57 (2.15)[d]	No	Inpatient	TRD
Boutros et al., 2002[6]	12 (4)	9 (1)	49.5 (8.0)	52.0 (7.0)	MDD	34.4 (10.1)[c]	31.7 (4.9)[c]	No	Outpatient	TRD

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Study										
Chen et al., 2013	10 (7)	10 (4)	44.1 (4.4)	47.3 (3.5)	MDD	23.5 (1.9)[a]	24.9 (1.9)[a]	No	Inpatient	TRD
Concerto et al., 2015	15 (6)	15 (7)	51.0 (6.5)	53.0 (6.7)	MDD	22.0 (21.0; 24.0)[b]	21.0 (20.0; 22.0)[b]	Yes	Outpatient	TRD
Eschweiler et al., 2000	5 (NR)	5 (NR)	NR	NR	MDD	27.4 (4.6)[b]	20.2 (3.8)[b]	No	NR	non-TRD
Fitzgerald et al., 2012 (1)[2]	24 (15)	20 (8)	43.4 (12.7)	44.9 (15.7)	MDD	23.7 (3.8)[a]	22.8 (2.1)[a]	No	NR	TRD
Fitzgerald et al., 2003 (1)	20 (8)	20 (11)	42.2 (9.8)	49.2 (14.2)	Mixed	36.1 (7.5)[M]	35.7 (8.1)[M]	No	Outpatient	TRD
Garcia-Toro et al., 2001	17 (7)	18 (8)	51.5 (15.9)	50.0 (11.0)	MDD	27.1 (6.7)[b]	25.6 (4.9)[b]	No	NR	TRD
George et al., 2010[2]	92 (58)	98 (50)	47.7 (10.6)	46.5 (12.3)	MDD	26.3 (5.0)[d]	26.5 (4.8)[d]	Yes	Outpatient	TRD
George et al., 2000[4]	20 (13)	10 (6)	42.4 (10.5)	48.5 (8.0)	Mixed	28.2 (5.9)[b]	23.8 (4.1)[b]	Yes	Outpatient	Mixed
George et al., 1997	7 (6)	5 (5)	42.4 (15.5)	41.0 (8.3)	Mixed	30.0 (4.0)[b]	26.0 (3.0)[b]	Yes	Outpatient	non-TRD
Hansen et al., 2004[6]	6 (2)	7 (2)	42.5 (38; 58)[13]	46 (44; 62)[13]	Mixed	26.5 (21.5; 27.6)[a]	23.8 (19.4; 28.0)[a]	No	Inpatient	NR
Hernández-Ribas et al., 2013	10 (8)	11 (8)	42.6 (5.6)	50.1 (8.1)	Mixed	19.7 (3.8)[b]	16.6 (2.4)[b]	Yes	Outpatient	TRD
Holtzheimer et al., 2004	7 (4)	8 (3)	40.4 (8.5)	45.4 (4.9)	MDD	22.7 (5.3)[a]	20.8 (6.3)[a]	Yes	Outpatient	TRD
Jakob 2008 (1)	12 (6)	12 (5)	NR	NR	MDD	27.2 (NR)[a]	23.9 (NR)[a]	NR	NR	NR

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Study										
Jakob 2008 (2)	12 (7)	12 (5)	NR	NR	MDD	24.1 (NR)[a]	23.9 (NR)[a]	NR	NR	NR
Kimbrell et al., 1999	5 (2)	3 (1)	40.2 (15.1)	43.7 (19.1)	Mixed	25.0 (6.6)[b]	24.3 (6.8)[b]	No	Mixed	TRD
Kreuzer et al., 2015	15 (8)	12 (8)	46.1 (9.5)	43.8 (10.5)	Mixed	22.3 (4.7)[b]	22.3 (4.7)[b]	No	Inpatient	NR
Lingeswaran et al., 2011	9 (6)	14 (8)	34 (10.5)	37.2 (11.8)	MDD	22.8 (3.7)[a]	22.0 (3.1)[a]	Yes	Mixed	NR
Loo et al., 1999	9 (NR)	9 (NR)	45.7 (14.7)	50.9 (14.7)	Mixed	21.5 (NR)[a]	25.1 (NR)[a]	No	Mixed	TRD
Nahas et al., 2003	11 (7)	12 (7)	42.4 (7.3)	43.4 (9.3)[11]	BD[12]	32.5 (4.3)[e]	32.8 (7.6)[e]	NA	Outpatient	NR
O'Reardon et al., 2007[6]	155 (86)	146 (74)	47.9 (11.0)	48.7 (10.6)	MDD	22.6 (3.3)[a]	22.9 (3.5)[a]	Yes	Outpatient	TRD
Paillère-Martinot et al., 2010	18 (11)	14 (10)	48.2 (7.8)	46.6 (10.3)	Mixed	26.0 (6.4)[b]	25.9 (6.7)[b]	Yes	Inpatient	TRD
Speer et al., 2014[2]	8 (5)	8 (11)	41.3 (14.5)	44.9 (9.1)	Mixed	35.8 (10.6)[e]	24.0 (4.6)[e]	No	Mixed	TRD
Su et al., 2005[5]	20 (15)	10 (17)	43.4 (11.3)	42.6 (11.0)	Mixed	24.9 (6.4)[b]	22.7 (4.7)[b]	Yes	NR	TRD
Theleritis et al., 2017 (1)[6]	26 (15)	20 (10)	39.1 (10.1)	38.0 (9.9)	MDD	30.6 (3.2)[a]	29.4 (3.2)[a]	Yes	Outpatient	TRD
Theleritis et al., 2017 (2)[6]	26 (11)	24 (10)	38.9 (13.9)	39.4 (8.9)	MDD	29.7 (4.6)[a]	30.3 (3.6)[a]	Yes	Outpatient	TRD
Zheng et al., 2010	19 (7)	15 (5)	26.9 (6.2)	26.7 (4.3)	MDD	24.6 (3.0)[a]	24.6 (2.8)[a]	Yes	NR	TRD

BRAIN STIMULATION DEPRESSION META-ANALYSIS

LF-R										
Fitzgerald et al., 2003 (2)	20 (7)	20 (11)	45.6 (11.5)	49.2 (14.2)	Mixed	37.7 (8.4)[M]	35.7 (8.1)[M]	No	Outpatient	TRD
Januel et al., 2006[2]	11 (9)	16 (12)	38.6 (11.2)	37.2 (11.7)	MDD	21.7 (3.5)[a]	22.5 (2.7)[a]	Yes	Inpatient	non-TRD
Pallanti et al., 2010 (1)	20 (12)	20 (12)	51.2 (12.5)	47.9 (9.1)	MDD	28.0 (5.9)[a]	29.1 (3.5)[a]	Yes	Outpatient	TRD
LF-L										
Kimbrell et al., 1999 (2)[2]	5 (4)	3 (1)	44 (15.92)	43.67 (19.14)	Mixed	34.4 (7.99)[b]	24.33 (6.81)[b]	No	Mixed	TRD
Padberg et al., 1999	6 (5)	6 (4)	46.7 (14.7)	43.3 (11.6)	MDD	26.7 (9.4)[b]	22.2 (8.8)[b]	NR	NR	TRD
Speer et al., 2014	8 (5)	8 (3)	39.6 (9)	44.9 (9.1)	Mixed	28.6 (7.6)[e]	24 (4.6)[e]	No	Mixed	TRD
BL										
Fitzgerald et al., 2006[2]	25 (15)	25 (16)	46.8 (10.7)	43.7 (10.2)	Mixed	22.5 (7.4)[a]	19.8 (4.4)[a]	No	Outpatient	TRD
Fitzgerald et al., 2016[7]	23 (13)	23 (13)	46.3 (12.6)	49.7 (11.0)	BD	23.2 (4.0)[a]	23.0 (5.1)[a]	NA	Outpatient	TRD
Fitzgerald et al., 2012 (2)[2]	22 (14)	20 (8)	40.5 (15.5)	44.9 (15.7)	MDD	24.3 (3.6)[a]	22.8 (2.1)[a]	No	NR	TRD
McDonald et al., 2006[8]	50 (27)	12 (5)	NR	NR	Mixed	26.4 (1.38)[b]	27.33 (2.86)[b]	Yes	Outpatient	TRD

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Pallanti et al., 2010 (2)	20 (11)	20 (12)	47.6 (12.3)	47.9 (9.1)	MDD	28.8 (6.0)[a]	29.1 (3.5)[a]	Yes	Outpatient	TRD
Prasser et al., 2015 (1)	17 (8)	17 (9)	50.4 (9.9)	42.6 (12.4)	Mixed	25.0 (4.4)[b]	25.3 (5.4)[b]	No	Mixed	Mixed
iTBS										
Duprat et al., 2016	22 (16)	25 (17)	40.09 (11.45)	43.16 (12.15)	MDD	21.14 (4.99)[a]	21.52 (6.21)[a]	Yes	Mixed	TRD
Li et al., 2014 (1)	15 (8)	15 (11)	42.4 (NR)	46.9 (NR)	MDD	23.1 (3.9)[a]	23.8 (3.2)[a]	Yes	NR	TRD
cTBS										
Li et al., 2014 (2)	15 (10)	15 (11)	49.2 (NR)	46.9 (NR)	MDD	24.3 (5.5)[a]	23.8 (3.2)[a]	Yes	NR	TRD
BLTBS										
Li et al., 2014 (3)	15 (11)	15 (11)	42.5 (NR)	46.9 (NR)	MDD	25.4 (5.1)[a]	23.8 (3.2)[a]	Yes	NR	TRD
Prasser et al., 2015 (2)	20 (10)	17 (9)	48.2 (10.9)	42.6 (12.4)	Mixed	27.4 (6.5)[b]	25.3 (5.4)[b]	No	Mixed	Mixed
dTMS										
Levkovitz et al..2015[6]	101 (48)	111 (53)	45.1 (11.7)	47.6 (11.6)	MDD	23.5 (4.3)[b]	23.4 (3.7)[b]	Yes	Outpatient	TRD
Tavares et al., 2017[6]	25 (17)	25 (18)	43.5 (12)	41.2 (8.9)	BD	25.32 (3.76)[a]	25.8 (5.25)[a]	NA	Outpatient	TRD

sTMS										
Jin et al., 2014[6,9,10]	29 (16)	16 (9)	42.5 (15.0)	46.3 (12.7)	MDD	21.3 (4.0)[a]	19.4 (4.1)[a]	No	Outpatient	non-TRD
Leuchter et al., 2015	59 (NR)	61 (NR)	46.7 (11.2)	45.7 (12.6)	MDD	21.8 (3.8)[a]	21.2 (2.9)[a]	Yes	Mixed	Mixed

Note. Mean ages are reported in years with standard deviation in parentheses for each of the active and sham treatment arms. The mean Hamilton Depression Rating Scale (HDRS) score at baseline is reported for each study with standard deviation in parentheses (except for Concerto et al., 2015 and Hansen et al., 2004 for which median, first quartile, and third quartile are reported). The Montgomery-Åsberg Depression Rating Scale (MADRS) score, denoted with superscript [M], is reported when the HDRS was not recorded. Means and standard deviations are rounded to the first figure after the decimal. Status refers to whether patients were outpatients, inpatients in a hospital admission, or whether there were both outpatients and inpatients (mixed). TMS = transcranial magnetic stimulation; HF-L = high-frequency left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency right-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; iTBS = intermittent theta burst stimulation; cTBS = continuous theta burst stimulation; BLTBS = bilateral theta burst stimulation; dTMS = deep transcranial magnetic stimulation; sTMS = synchronised transcranial magnetic stimulation; NR = not reported; NA = not applicable; MDD = major depressive disorder; BD = bipolar depression; TRD = treatment resistant depression. [1]MADRS based on the intention-to-treat sample who received ≥ 1 session of active stimulation. [2]Numbers are based on the intention-to-treat sample. [3,4,5,8,9]Two active treatment groups were combined. [6]Numbers based on the intention-to-treat sample who received ≥ 1 session of active stimulation. [7]HDRS based on the intention-to-treat sample. [10]Age based on the intention-to-treat sample who received ≥ 1 session of active stimulation. [11]Age based on 11 patients. [12]Two patients had mixed features. [13]Indicates Median and IQR. [a]HDRS-17. [b]HDRS-21.[c]HDRS-25. [d]HDRS-24. [e]HDRS-28.

Table 3

Treatment characteristics: tDCS studies

Authors	Location		Electrode size	Current strength	Current density	Session duration	Number of sessions	Treatment strategy	Sham stimulation
	Anode	Cathode/Reference							
Fregni et al., 2006a	F3	FP2	35cm^2	1mA	0.028	20min	5	Monotherapy	05sec
Fregni et al., 2006b	F3	FP2	35cm^2	1mA	0.028	20min	5	Monotherapy	05sec
Boggio et al., 2008[1]	F3	FP2; Midline	35cm^2	2mA	0.057	20min	10	Monotherapy	30sec
Loo et al., 2010	pF3	F8	35cm^2	1mA	0.028	20min	5	Mixed	30sec
Blumberger et al., 2012	F3	F4	35cm^2	2mA	0.057	20min	15	Mixed	30sec
Brunoni et al., 2013[2]	F3	F4	25cm^2	2mA	0.080	30min	12	Monotherapy	60sec
Salehinejad et al., 2015	F3	F4	35cm^2	2mA	0.057	20min	22	Monotherapy	30sec
Salehinejad et al., 2017	F3	F4	35cm^2	2mA	0.057	30min	10	Monotherapy	30sec
Brunoni et al., 2017[2]	F3	F4	25cm^2	2mA	0.080	30min	10	Monotherapy	30sec

Note. Electrode locations are reported according to the EEG 10/20 system. Current densities are reported in mA/cm^2. Sham stimulation indicates the duration of time that current was applied for giving an initial sensation of tDCS on the scalp. tDCS = transcranial direct current stimulation. [1]Two sham treatment groups were combined. [2]Patients in sham group also received an oral placebo tablet.

Table 4

Sample characteristics: tDCS studies

Authors	Number of participants (female)		Age		Diagnosis	HDRS		Excluded psychosis	Status	Treatment resistance
	Active	Sham	Active	Sham		Active	Sham			
Fregni et al., 2006a	5 (NR)	5 (NR)	NR	NR	MDD	NR	NR	NR	NR	NR
Fregni et al., 2006b	9 (5)	9 (6)	47.6 (10.4)	45.3 (9.3)	MDD	23,6 (5,0)	25,9 (4,3)	Yes[a]	Outpatient	NR
Boggio et al., 2008[1]	21 (14)	19 (13)	51.6 (7.7)	46.4 (7.1)	MDD	21,1 (4,4)[b]	21,8 (4,8)[b]	Yes	NR	Mixed
Loo et al., 2010[2]	20 (11)	20 (11)	49.0 (10.0)	45.6 (12.5)	MDD	18,3 (5,8)[c]	17,3 (4,7)[c]	Yes[a]	Outpatient	Mixed
Blumberger et al., 2012[3,6]	13 (10)	11 (10)	45.3 (11.6)	49.7 (9.4)	MDD	24,9 (3,1)[c]	24,1 (2,9)[c]	Yes	Outpatient	TRD
Brunoni et al., 2013[4]	30 (21)	30 (20)	41.0 (12.0)	46.4 (14.0)	MDD	21,0 (3,8)[c]	22,0 (4,2)[c]	Yes	Outpatient	Mixed
Salehinejad et al., 2015	15 (8)	15 (9)	28.7 (5.87)	27.9 (5.84)	MDD	24.7 (3.05)[d]	22.8 (2.06)[d]	Yes	Outpatient	TRD
Salehinejad et al., 2017	12 (7)	12 (8)	26.8 (7.1)	25.5 (4.6)	MDD	24,6 (2,6)[d]	22,6 (1,9)[d]	Yes	Outpatient	non-TRD
Brunoni et al., 2017[5,6,7]	91 (64)	60 (41)	44 (11.19)	40.88 (12.87)	MDD	21.93 (3.89)[c]	22.7 (4.27)[c]	Yes	Outpatient	Mixed

BRAIN STIMULATION DEPRESSION META-ANALYSIS

Note. Mean ages are reported in years with standard deviation in parentheses for each of the active and sham treatment arms. The mean Hamilton Depression Rating Scale (HDRS) score at baseline is reported for each study with standard deviation in parentheses. Means and standard deviations are rounded to the first figure after the decimal. Status refers to whether patients were outpatients, inpatients in a hospital admission, or whether there were both outpatients and inpatients (mixed). tDCS = transcranial direct current stimulation; MDD = major depressive disorder; TRD = treatment resistant depression; NR = not reported. [1]Two sham treatment groups were combined. [2,3,4,7]Numbers are based on the intention-to-treat sample. [5]Numbers based on participants of age ≤ 70 years. [6]Patients in sham group also received an oral placebo tablet. [a]Excluded "other psychiatric disorders." [b]HDRS-21. [c]HDRS-17. [d]HDRS-24.

Table 5

Random-Effects Meta-Analysis of Response Rates

Treatment Modality	k	Odds Ratio	95% Confidence Interval		Q	I^2
HF-L	31	**3.94**	**2.52**	**6.15**	41.15	27.1%
LF-R	3	**7.44**	**2.06**	**26.83**	1.59	0.0%
LF-L	3	1.41	0.15	12.88	0.14	0.0%
BL	6	**3.68**	**1.66**	**8.13**	3.45	0.0%
cTBS[*]	1	1.63	0.23	11.46	-	-
iTBS	2	**4.70**	**1.14**	**19.38**	0.02	0.0%
blTBS	2	4.28	0.54	34.27	2.91	65.7%
dTMS	2	**1.69**	**1.003**	**2.85**	0.97	0.0%
sTMS	2	2.71	0.44	16.86	4.15	75.9%
tDCS	8	**4.32**	**2.02**	**9.29**	10.64	34.2%

Note. HF-L = high-frequency, left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency, right-sided repetitive transcranial magnetic stimulation; LF-L = low-frequency, left-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; dTMS = deep transcranial magnetic stimulation; cTBS = continuous theta burst stimulation; iTBS = intermittent theta burst stimulation; blTBS = bilateral theta burst stimulation; sTMS = synchronised transcranial magnetic stimulation; tDCS = transcranial magnetic stimulation. [*]inverse variance method used.

Table 6

Random-Effects Meta-Analysis of Remission Rates

Treatment Modality	k	Odds Ratio	95% Confidence Interval		Q	I^2
HF-L	25	**2.74**	**1.75**	**4.28**	22.67	0.0%
LF-R	2	**14.10**	**2.79**	**71.42**	0.50	0.0%
LF-L	3	0.86	0.08	9.11	0.03	0.0%
BL	5	**3.05**	**0.87**	**10.67**	4.48	10.7%
cTBS	-	-	-	-	-	-
iTBS[*]	1	6.22	0.28	136.90	-	-
blTBS[*]	1	1.32	0.19	9.02	-	-
dTMS	2	**2.24**	**1.24**	**4.06**	0.02	0.0%
sTMS	2	2.51	0.23	26.76	4.12	75.7%
tDCS	7	**3.07**	**1.58**	**5.99**	6.27	4.4%

Note. HF-L = high-frequency, left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency, right-sided repetitive transcranial magnetic stimulation; LF-L = low-frequency, left-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; dTMS = deep transcranial magnetic stimulation; cTBS = continuous theta burst stimulation; iTBS = intermittent theta burst stimulation; blTBS = bilateral theta burst stimulation; sTMS = synchronised transcranial magnetic stimulation; tDCS = transcranial magnetic stimulation. [*]inverse variance method used.

Table 7

Random-Effects Meta-Analysis of Continuous Treatment Effects

Treatment Modality	k	g	95% Confidence Interval		Q	I^2
HF-L	28	**-0.75**	**-1.02**	**-0.47**	101.64	73.4%
LF-R	2	-0.77	-1.64	0.09	2.72	63.3%
LF-L	2	-0.33	-1.18	0.51	0.76	0.0%
BL	4	-0.07	-0.38	0.25	0.25	0.0%
cTBS	-	-	-	-	-	-
iTBS	1	-0.44	-1.02	0.14	0.00	-
blTBS	1	-0.03	-0.65	0.56	-	-
dTMS	2	-0.29	-0.55	-0.03	0.75	0.0%
sTMS	2	-0.55	-1.13	0.02	3.24	69.1%
tDCS	6	-0.76	-1.42	-0.10	32.65	84.7%

Note. HF-L = high-frequency, left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency, right-sided repetitive transcranial magnetic stimulation; LF-L = low-frequency, left-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; dTMS = deep transcranial magnetic stimulation; cTBS = continuous theta burst stimulation; iTBS = intermittent theta burst stimulation; blTBS = bilateral theta burst stimulation; sTMS = synchronised transcranial magnetic stimulation; tDCS = transcranial magnetic stimulation. [*]inverse variance method used.

Table 8

Random-Effects Meta-Analysis of All-cause Discontinuation Rates

Treatment Modality	k	Odds Ratio	95% Confidence Interval		Q	I^2
HF-L	34	0.85	0.59	1.23	14.54	0.0%
LF-R	3	0.48	0.12	1.99	0.35	0.0%
LF-L	3	0.84	0.11	6.73	0.71	0.0%
BL	6	0.90	0.33	2.43	3.03	0.0%
cTBS[*]	1	1.00	0.02	53.66	-	-
iTBS	2	1.06	0.06	17.66	0.00	0.0%
BLTBS	2	0.47	0.04	5.88	0.23	0.0%
dTMS	2	1.03	0.32	3.36	2.10	52.3%
sTMS	2	0.72	0.36	1.44	0.32	0.0%
tDCS	9	1.34	0.68	2.66	6.66	0.0%

Note. HF-L = high-frequency, left-sided repetitive transcranial magnetic stimulation; LF-R = low-frequency, right-sided repetitive transcranial magnetic stimulation; LF-L = low-frequency, left-sided repetitive transcranial magnetic stimulation; BL = bilateral repetitive transcranial magnetic stimulation; dTMS = deep transcranial magnetic stimulation; cTBS = continuous theta burst stimulation; iTBS = intermittent theta burst stimulation; blTBS = bilateral theta burst stimulation; sTMS = synchronised transcranial magnetic stimulation; tDCS = transcranial magnetic stimulation. [*]inverse variance method used.

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of literature search.

Figure 2. Contour-enhanced funnel plot of all RCTs included in the meta-analysis of response rates.

Figure 3. Forest plot of response rates.

Neurostimulation	Active		Sham		Odds Ratio	OR	95%−CI
	n	N	n	N			
BL rTMS	38	154	10	114		3.68	[1.66; 8.13]
BL TBS	15	35	5	32		4.28	[0.54; 34.27]
cTBS	3	15	2	15		1.62	[0.23; 11.46]
dTMS	49	126	37	136		1.69	[1.00; 2.85]
HF−L rTMS	215	718	65	653		3.94	[2.52; 6.15]
iTBS	10	37	3	40		4.70	[1.14; 19.38]
LF−L rTMS	1	19	0	17		1.41	[0.15; 12.88]
LF−R rTMS	15	51	3	56		7.44	[2.06; 26.83]
sTMS	36	89	20	77		2.71	[0.44; 16.86]
tDCS	86	216	24	181		4.33	[2.02; 9.29]

0.01   0.1   0.5 1 2   10   200

Favours sham treatment   Favours active treatment

Figure 4. Forest plot of remission rates.

Neurostimulation	Active		Sham		Odds Ratio	OR	95%-CI
	n	N	n	N			
BL rTMS	19	154	3	114		3.05	[0.87; 10.67]
BL TBS	3	35	2	32		1.32	[0.19; 9.02]
dTMS	38	126	22	136		2.24	[1.24; 4.06]
HF−L rTMS	103	718	25	653		2.74	[1.75; 4.28]
iTBS	2	37	0	40		6.22	[0.28; 136.90]
LF−L rTMS	0	19	0	17		0.86	[0.08; 9.11]
LF−R rTMS	13	51	2	56		14.10	[2.79; 71.42]
sTMS	21	89	11	77		2.51	[0.23; 26.76]
tDCS	52	216	14	181		3.08	[1.58; 5.99]

0.01 0.1 0.5 1 2 10 200

Favours sham treatment   Favours active treatment

Figure 5. Forest plot of post-treatment continuous depression scores.

Neurostimulation	Active N	Sham N	Standardised mean difference	SMD	95%-CI
BL rTMS	154	111		-0.07	[-0.38; 0.25]
BL TBS	35	35		-0.03	[-0.65; 0.59]
dTMS	114	117		-0.29	[-0.55; -0.03]
HF-L rTMS	707	645		-0.75	[-1.02; -0.47]
iTBS	37	40		-0.44	[-1.02; 0.14]
LF-L rTMS	19	16		-0.33	[-1.18; 0.51]
LF-R rTMS	51	56		-0.77	[-1.64; 0.09]
sTMS	89	89		-0.55	[-1.13; 0.02]
tDCS	216	181		-0.76	[-1.42; -0.10]

-4   -2   0   2   4

Favours active treatment   Favours sham treatment

Figure 6. Forest plot of all-cause discontinuation rates.

Neurostimulation	Active		Sham		Odds Ratio	OR	95%−CI
	n	N	n	N			
BL rTMS	8	158	9	118		0.90	[0.33; 2.43]
BL TBS	0	35	1	33		0.47	[0.04; 5.88]
cTBS	0	15	0	15		1.00	[0.02; 53.66]
dTMS	18	137	22	148		1.03	[0.32; 3.36]
HF−L rTMS	56	744	65	685		0.85	[0.59; 1.23]
iTBS	0	37	0	40		1.06	[0.06; 17.66]
LF−L rTMS	1	19	1	17		0.84	[0.11; 6.73]
LF−R rTMS	3	51	8	56		0.48	[0.12; 2.00]
sTMS	18	136	22	118		0.72	[0.36; 1.44]
tDCS	28	216	16	181		1.34	[0.68; 2.66]

0.01 0.1 0.5 1 2 10 200

Favours sham treatment   Favours active treatment