1 **Title:**
2 *Fusarium graminearum*-Induced Shoot Elongation and Root Reduction in Maize
3 Seedlings Correlate with Later Seedling Blight Severity
4

Authors:
Shaoqun Zhou[1,2,*], Justin S. Bae[1], Gary C. Bergstrom[3], Georg Jander[1]
[1]Boyce Thompson Institute, 533 Tower Road, Ithaca, NY 14853
[2]Plant Biology Section, School of Integrated Plant Science, Cornell University, Ithaca,
NY14853
[3]Plant Pathology and Plant-Microbe Biology Section, School of Integrated Plant Science,
Cornell University, Ithaca, NY14853
12
13 ORCID IDs: 0000-0002-3643-7875 (S.Z.); 0000-0002-7166-9030 (J.S.B.); 0000-0001-
14 9613-270X (G.C.B.); 0000-0002-9675-934X (G.J.)
15
16 *To whom correspondence should be addressed, sz357@cornell.edu

ABSTRACT

        Fusarium graminearum seedling blight is a common disease of maize (*Zea mays*). Development of genetic resistance to seedling blight in maize germplasm requires efficient and accurate quantitative assessment of disease severity. Through artificial inoculation experiments under controlled growth conditions, we determined that host genotype, pathogen genotype, and infection dose influence the extent to which *F. graminearum* induces shoot elongation and inhibits root growth in maize seedlings. A comparison of fifteen maize inbred lines showed independent variation of these two fungus-induced morphological changes. In a broader survey with nine commercial maize hybrids and three field-collected fungal isolates, there was significant correlation between these seedling morphological responses, as well as with later seedling blight severity. Analysis of variance suggested that this variation and the observed correlative relationships were primarily driven by differing pathogenicity of the three fungal isolates. Together, our results indicate that *F. graminearum*-induced shoot elongation and root reduction in maize seedlings have distinct underlying physiological mechanisms, and that early observations of seedling morphological changes can serve as a proxy for investigating natural variation in host resistance and pathogen aggressiveness at later growth stages.

Keywords: *Fusarium graminearum* seedling blight, maize, shoot elongation, root reduction.

INTRODUCTION

Fusarium graminearum Schwabe (teleomorph: *Gibberella zeae*) is a widespread fungal phytopathogen that can cause seedling blight, root rot, stem rot, or ear rot in maize (*Zea mays*) depending on the timing and tissue of infection (Munkvold and White 2016). The severity of maize diseases caused by *F. graminearum* in the United States varies significantly from year to year, causing 15 to 80 million bushels of annual yield loss due to ear rot, and another 30 to 85 million bushels due to stalk rot (Mueller 2016a, b, c, 2017). Unlike Gibberella ear rot and stem rot diseases caused by *F. graminearum*, which can be tracked by the characteristic pink-reddish pigmentation of fungal mycelium, seedling blight and root rot diseases are often caused by a consortium of soil-dwelling fungal and oomycete pathogens, including *F. graminearum*. While the dominant species/strains of these consortia are likely to vary considerably across geographic locations, collectively these seedling blight and root rot diseases can account for over 200 million bushels of annual yield loss, making them among the most damaging maize diseases in the northern states (Mueller 2016b). Management of *F. graminearum* seedling blight can be difficult because standard seed fungicide treatment is not always sufficient and may promote the development of resistance in the fungal population. Genetic sources of seedling blight resistance have yet to be established in commercial maize germplasm. In addition to causing direct yield loss through tissue rotting, *F. graminearum* can also cause crop contamination with mycotoxins such as deoxynivalenol and zearalenone, leaving the grain and stover unsafe for human and livestock consumption, and imposing further financial burdens on the growers (Arunachalam and Doohan 2013; Maresca 2013). Furthermore, *F. graminearum* can also cause destructive scab and head blight diseases in barley and wheat (McMullen et al. 2012; Trail 2009). Considering the

overwinter persistence and airborne infectious ascospores of this pathogen, unchecked *F.*

graminearum diseases in maize could have a far-reaching negative impact on other crop

species in the vicinity.

Breeders and researchers interested in improving resistance against *F.*

graminearum in maize have been doing artificial inoculation experiments in both field

and controlled environments, primarily focused on infected maize ears and stalks.

Genetic mapping of Gibberella ear rot resistance based on symptom severity has

identified a large number of quantitative trait loci (QTL), each with small effect size.

Moreover, these QTL tend to be sensitive to experimental methods and environmental

conditions, making them difficult to validate across different studies (Ali et al. 2005;

Brauner et al. 2017; Kebede et al. 2016). In contrast, major genetic loci contributing to

Gibberella stalk rot resistance have been identified with bi-parental mapping populations

(Chen et al. 2017; Ma et al. 2017; Yang et al. 2010; Zhang et al. 2012). In a near-isogenic

line population segregating for Gibberella stalk rot resistance, Ye et al. (2013) observed

significant differences in seedling root browning and shrinkage after *F. graminearum*

inoculation. This suggests that resistance against Gibberella stalk rot and *F. graminearum*

seedling blight, both diseases in vegetative tissues, are likely controlled by similar genetic

and physiological mechanisms, and are distinct from those affecting Gibberella ear rot

resistance. This is consistent with the known influence of tissue type and developmental

stage on the interactions between maize and *F. graminearum* (Zhang et al. 2016).

Genetic improvement of *F. graminearum* resistance depends on accurate

quantitative assessment of resistant and susceptible phenotypes. Here we show that *F.*

graminearum inoculation of maize seedlings not only inhibits root growth, but can also

promote shoot elongation in a dosage- and genotype-dependent manner. Shoot elongation

in response to *F. graminearum* root infection is significantly correlated with seedling

survival, suggesting that these traits could be used to assess seedling resistance, as well as

serving as a quantitative measure of the aggressiveness of different *F. graminearum*

isolates.

MATERIALS AND METHODS

Plant and fungal materials. Fifteen parental lines from the maize nested association

mapping population (McMullen et al. 2009) were chosen for *F. graminearum* infection

assays. For a broader survey, nine organic commercial maize hybrids were kindly

provided by the Blue River Organic Seed Company (Table 1). The maize hybrids were

selected to cover different maturation time and provider-assessed seedling vigor levels.

The accession numbers of the tested hybrids are masked per the provider's request. Three

field-collected *F. graminearum* isolates (Table 2) that have shown different levels of

aggressiveness in a previous laboratory test (Kuhnem et al. 2015) were used for

inoculations (Table 2). Fungal chemotype, namely the type of major mycotoxin

accumulated, was not considered during isolate selection, since it was not significantly

associated with aggressiveness in the previous test.

Plant growth and fungal inoculation methods. Maize seeds were germinated in

moisturized rolls of germination paper after surface sterilization with 10% household

bleach for 30 minutes. Germinated seedlings were transplanted to 7.5 cm × 7.5 cm plastic

pots with Turface ® MVP ® calcined clay (Profile Products LLC, Buffalo Grove, IL)

when their primary roots reached approximately 9 cm. For inoculation experiments,

seedling roots were immersed in *F. graminearum* spore suspension or mock solution

(0.03% Phytagar) for one hour prior to transplanting. Fungal spore suspensions were

freshly prepared by flooding and scraping 7-day-old fungal cultures maintained on Potato

Dextrose Agar plates under black lights with 0.03% Phytagar suspension. The spore

concentration was estimated with a hemocytometer (Hausser Scientific, Horsham, PA)

and light microscopy. Fungal hyphae fragments were observed in spore suspension by

light microscopy.

For the concentration gradient experiment, twenty-four seedlings of each maize

hybrid were grown, and six were inoculated with each tested fungal spore concentration

and mock inoculum. For each pair of maize hybrid and fungal isolate included in the

diversity screening, at least 15 seedlings were inoculated with fungal spores, and 5 were

mock-inoculated. Transplanted seedlings were kept at 26 °C (Day)/23 °C (Night), 60%

humidity, and a 16 hour long-day light cycle. Seedlings inoculated with different *F.*

graminearum isolates or mock inoculum were kept in separated growth chambers set to

identical growth conditions.

Phenotypic data collection and analysis. The heights of fungus- and mock-inoculated

seedlings were measured nine days post-inoculation. On the same day, seedlings were

removed from pots to measure their total root length using an image analysis algorithm

implemented in the RootReader 2D software (Clark et al. 2013). After measurement,

seedlings were re-potted in the same Turface particles. At fourteen days post-inoculation,

the number of surviving seedlings within each fungus-inoculated maize hybrid population

was counted as a measurement of *F. graminearum* seedling blight severity. Seedlings

were considered dead when they broke at the soil line, with browning and rot at the

breakpoint.

For seedling height and total root length, an average ratio of mock- and fungus-

inoculated seedlings for each maize hybrid-fungus isolate combination was calculated to

shown variation in these fungus-induced responses. Measurements from fungus- and

mock-inoculated seedlings were compared using Student's t-tests. Average ratios were

rounded to one when measurements from the two groups were not significantly different

from one another. For two-way ANOVA, maturation times of maize hybrids were

grouped into four bins, as indicated in parentheses in Table 1. The average ratios of

seedling height and total root length between fungus- and mock-treated seedlings were

used as quantitative measurements to calculate linear correlation with seedling survival

rate measured at 14 days after infection.

RESULTS

Fusarium graminearum **inoculation at low concentration induces shoot elongation**

and root reduction in maize seedlings.

Different concentrations of spore suspensions have been used to study natural variation in

maize seedling responses to *F. graminearum* inoculation (Ye et al., 2013; Kuhnem et al.,

2015). To determine the optimal concentration of *F. graminearum* spores that leads to the

largest change in plant morphology, two concentration gradient experiments were

performed with Gz014, a field-collected *F. graminearum* strain (Table 2), and two

commercial maize hybrids with differing seedling vigor levels (Hybrid IV and V in Table

1, as assessed by the provider, Blue River Organic Seed Company).

At 1.2 x 10^6 spores/mL, a concentration that is comparable to a previous study,

the seedling stunting phenotype that had been reported was not observed in either of the

tested maize hybrids (Kuhnem et al. 2015). Instead, the hybrid with high seedling vigor

showed no significant change in seedling height, whereas the moderately vigorous hybrid

seedlings grew significantly taller when inoculated with *F. graminearum* spores. At

lower concentrations, fungus-induced shoot elongation became obvious in both highly

and moderately vigorous hybrid seedlings (Figure 1A). Underground, *F. graminearum*

infection significantly reduced root growth in the moderately vigorous hybrid with no

evidence of dosage response. This fungus-induced change was not observed in the highly

vigorous hybrid, which showed no significant change in total root length at all tested

doses (Figure 1B).

For both fungus-induced morphological changes, inoculation with the lowest

fungal spore concentration (3 x 10^5 count/mL) resulted in a response that was stronger

than or comparable to that observed with the two higher inoculum concentrations.

Therefore, the 3 x 10^5 spores/mL *F. graminearum* dose was used for all subsequent

experiments.

**Fusarium graminearum*-induced shoot elongation and root reduction vary*

independently across maize inbred lines.

To confirm the *F. graminearum*-induced shoot elongation and root reduction observed in

the concentration gradient experiment, the artificial inoculation experiment was repeated

with the reference maize genetic inbred line B73 with the same fungal isolate following

the same protocol. Consistently, the same morphological changes were observed (Figure

2). To investigate how *F. graminearum*-induced seedling shoot elongation and root

reduction may be related, seedlings of fifteen maize inbred lines were inoculated with

Gz014 or mock inoculum and their shoot and root growth were quantified 9 days after

inoculation. Although *F. graminearum* inoculation reduced total root length in most

inbred lines by approximately 50%, no significant difference was observed between

mock- and *F. graminearum*-inoculated NC350, NC358, and Mo17 seedlings, (Figure

3A).

In the same experiment, *F. graminearum*-induced shoot elongation also varied,

ranging from 0 to over 25% increase (Figure 3B). Noticeably, the two fungus-induced

morphological changes do not correlate with one another across the fifteen tested inbred

lines tested (Pearson's $R^2 = 0.0079$, $p > 0.05$). Two inbred lines showing no significant *F.*

graminearum-induced root reduction, NC350 and NC358, grew 16% and 24% taller,

respectively, after *F. graminearum* inoculation. Conversely, Tx303 only showed obvious

root reduction (40%) but no significant shoot elongation (Figure 3). Inbred line Mo17

showed neither the tested fungus-induced morphological changes.

Fusarium graminearum*-induced morphological changes and *F. graminearum

seedling blight severity are primarily determined by the fungal genotype.

To further survey the natural variation in *F. graminearum*-induced seedling shoot

elongation and root reduction, the artificial inoculation experiment was expanded to the

nine commercial maize hybrids, including the two used in the concentration gradient

experiment above (Table 1), and three *F. graminearum* isolates (Table 2). Both *F.*

graminearum-induced morphological changes were highly variable across this panel

(Figure 4). Two-way analysis of variance (ANOVA) performed on both traits showed

that fungal isolate, rather than maize maturation time or provider-assessed seedling vigor,

is the main source of assay variance (Table 3).

After quantifying *F. graminearum*-induced morphological changes, the fungus-

infected seedlings were kept under the same growth conditions, and *F. graminearum*

seedling blight severity was quantified as the percentage of seedlings surviving in each

maize hybrid population fourteen days after inoculation. As in the case of the

morphological data, ANOVA of the seedling survival rate data identified the *F.*

graminearum isolate as the most important contributor. Whereas both Gz014 and Gz829

infection left less than 50% of inoculated seedlings surviving after fourteen days, more

than 85% of Gz835-inoculated seedlings survived in the same time period (Figure 5). The

maturation time and seedling vigor of maize hybrids were not significant sources of

variance (Table 3).

Fusarium graminearum*-induced shoot elongation is positively correlated with *F.

***graminearum* seedling blight severity.**

To investigate the relationship between *F. graminearum*-induced morphological changes

and *F. graminearum* seedling blight severity, linear regression analysis was performed on

the extent of these changes and seedling survival. This showed that *F. graminearum*-

induced seedling elongation has a significant negative correlation with seedling survival

rate at 14 days after inoculation (Figure 6A; Pearson's $R^2 = 0.244$; $p < 0.005$). Thus,

stronger fungus-induced shoot elongation may be predictive of more severe *F.*

graminearum seedling blight. Unexpectedly, the extents of *F. graminearum*-induced

shoot elongation showed strong positive correlation with root length (Figure 6B;

Pearson's $R^2 = 0.676$; $p < 0.005$), and a significant negative correlation between root

reduction and seedling survival rate (Figure 6C; Pearson's $R^2 = 0.263$; p < 0.005). This

leads to the counter-intuitive inference that resistance to fungus-induced root growth

inhibition co-occurs with stronger shoot elongation and more severe *F. graminearum*

seedling blight.

DISCUSSION

Under field conditions, maize seedlings infected by *F. graminearum* as well as

other fungal and oomycete pathogens are often characterized by stunted growth and

dehydration in the aboveground tissues before death. Similar seedling stunting symptoms

have also been observed in a previous artificial inoculation test with *F. graminearum*

spores under laboratory conditions (Kuhnem et al., 2015). However, in our concentration

gradient experiment with the same fungal isolates, low dosage *F. graminearum*

inoculation on maize seedling root induced shoot elongation (Figure 1A; Figure 2A).

Based on the natural variation in this *F. graminearum*-induced response across different

maize inbred lines and hybrids reported in this study (Figure 3A; Figure 4A), it is likely

that the inconsistent observations between studies could arise from inoculum

concentration, maize genotype, or unintended differences between experimental

conditions.

Interestingly, the observed *F. graminearum*-induced shoot elongation phenotype

is reminiscent of the foolish seedling disease in rice, which is caused by a related fungal

pathogen *Gibberella fujikuroi*. Rice seedlings affected by foolish seedling disease

demonstrate abnormal seedling elongation resulting from the production of gibberellic

acid (GA), a growth-promoting phytohormone, *G. fujikuroi* (Yabuta and Sumiki 1938).

Fungus-derived GA has been hypothesized to be a virulence factor by negatively

influencing the signaling transduction pathway of the immunity-regulating

phytohormone, jasmonic acid (Navarro et al. 2008). However, unlike *G. fujikuroi, F.*

graminearum is not known to produce GA, and its genome does not contain biosynthetic

genes required for GA biosynthesis (Cuomo et al. 2007). However, we cannot rule out

the possibility that *F. graminearum* infection induces endogenous maize GA production.

Consistent with previous reports (Ye et al. 2013; Kuhnem et al., 2015), *F.*

graminearum infection induced maize seedling root growth reduction in a host genotype-

dependent manner (Figure 1B; Figure 2B; Figure 3B; Figure 4B). Application of a

computer image-processing tool, RootReader 2D (Clark et al. 2013), in our study enabled

more accurate, quantitative, and efficient measurement of fungus-induced morphological

change, and its natural variation with host and pathogen genotype. Fungus-induced root

reduction and shoot elongation measurements from the same experiment on fifteen maize

inbred lines showed no correlation in the extent of these two traits, with some inbred lines

showing only one or the other fungus-induced change (Figure 3A&B). This observation

suggests that these two responses may be governed by distinct physiological mechanisms.

As one of the most widespread fungal phytopathogens, *F. graminearum* is known

to be highly diverse, and different strains are highly variable in their aggressiveness

(Backhouse 2014; Boutigny et al. 2011; Lee et al. 2010; Sampietro et al. 2011; Wang et

al. 2011). Our survey of nine commercial maize hybrids and three fungal isolates (Figure

4 & 5) is consistent with prior observations that the prevalence of *F. graminearum*-

related diseases is highly variable across multiple locations, depending on the

aggressiveness of the predominant isolate and environmental permissiveness (Backhouse

2014; Boutigny et al. 2011; Lee et al. 2010; Sampietro et al. 2011; Wang et al. 2011).

Unexpectedly, maize hybrid-fungal isolate pairs with stronger fungus-induced

root growth inhibition tended to show a lower level of induced shoot elongation (Figure

6C) and better survival (Figure 6B). This was surprising, because severe reduction in root

growth was expected to significantly impair the seedlings' ability to acquire water and

nutrients from the soil, which could lead to seedling death. Two-way ANOVA and a

closer look at the data distribution suggest that these correlative relationships are

primarily driven by differences between the fungus isolates. Specifically, the isolate that

induced significantly stronger shoot elongation, Gz835, induced less root reduction and

seedling death (Figure 5). This suggests that these fungus-induced morphological

changes can be used for quantitative assessment of the aggressiveness of *F. graminearum*

isolates, and that different *F. graminearum* isolates may have distinct pathogenic

mechanisms when interacting with maize seedlings.

ACKNOWLEDGEMENT

The authors would like to thank the Blue River Organic Seed Company for

providing the maize hybrids used in this study, Jon Shaff and Eric Craft for training and

assistance in using the RootReader 2D platform, and Jaime Cummings for training and

assistance in fungal culture maintenance. This work was funded by the Northeast

Sustainable Agriculture and Research Education Graduate Student Grant to S.Z.

(GNE14-092), USDA-NIFA Hatch Grant NYC1537436 to G.C.B., and US National

Science Foundation IOS-1339237 to G.J. J.S.B. was supported by the Plant Genome

Research Program at the Boyce Thompson Institute, a Research Experience for

Undergraduate program funded by the US National Science Foundation award DBI-

1358843.

Author Contribution

S.Z. conceived, designed, and performed the experiments, analyzed the data, and

wrote the manuscript. J.S.B. assisted in the natural variation experiments and related data

analyses. G.C.B. provided the fungal isolates used in the experiments, and advice in

experimental design and manuscript revision. G.J. provided infrastructural support to this

study, and advice in experimental design and manuscript revision.

Figure Legends

Figure 1. *Fusarium graminearum* **inoculation induces shoot elongation and root reduction in maize seedlings in a dosage- and host genotype-dependent manner.** Shoot height (A) and total root length (B) measurements on seedlings inoculated at each fungal spore concentration were compared to mock-treated seedlings with Dunnett's tests relative to mock-infected controls (N = 6 for each group; * p < 0.05). All measurements were taken 9 days post-inoculation. Fungal spore concentration is in count/milliliter. s.d. = standard deviation.

Figure 2. *Fusarium graminearum* **inoculation induces shoot elongation and root reduction in maize inbred line B73.** Representative photos of shoots (A) and roots (C) mock- and fungus-inoculated B73 seedlings were taken 9 days post-inoculation. Shoot height (B) and total root length (D) were measured on the same day and compared with Student's *t*-tests (N = 6 for each group; * p < 0.05). s.d. = standard deviation.

Figure 3. *Fusarium graminearum*-**induced morphological changes in maize seedlings vary across maize inbred lines.** Root length (A) and shoot height (B) of each maize inbred line were compared to measurement from a corresponding mock treatment group with Student's *t*-test (* p < 0.05). The ratio of fungus- and mock-inoculated shoot height and root length were plotted, with the dotted line denoting the expected ratio if there is no significant difference between the two groups. All measurements were taken at 9 days post-inoculation. s.d. = standard deviation.

Figure 4. *Fusarium graminearum*-**induced morphological changes in maize seedlings vary with both plant and fungal genotype.** Nine maize hybrid lines were inoculated with three *F. graminearum* isolates (Gz014, Gz829, and Gz835). Root length (A) and shoot height (B) of each maize hybrid-fungus isolate pair were compared to measurement from a corresponding mock treatment group with Student's *t*-test (* p < 0.05). The ratios of fungus- and mock-inoculated root length and shoot height were plotted, with the dotted line denoting the expected ratio if there is no significant difference between the two groups. All measurements were taken at 9 days post-inoculation. s.d. = standard deviation.

Figure 5. *Fusarium graminearum*-**induced morphological changes and seedling blight severity vary significantly between fungal isolates.** The extent of *F.* graminearum-induced shoot elongation and root reduction at 9 days post-inoculation (A & B) and survival rates of maize seedlings inoculated with different *F. graminearum* isolates at 14 days post-inoculation (C) were compared using two-way analysis of variance with Tukey's HSD. Significantly different groups are denoted with different letters. s.d. = standard deviation.

Figure 6. *Fusarium graminearum*-**induced morphological changes and seedling blight severity are significantly correlated with one another.** Each mark represents a maize hybrid-fungal isolate pair, with maize hybrids inoculated with different fungal isolates represented by different mark shapes

340   **References**

341   Ali ML, Taylor JH, Jie L, Sun G, William M, Kasha KJ, Reid LM, Pauls KP (2005)
342   Molecular mapping of QTLs for resistance to Gibberella ear rot, in corn,
343   caused by *Fusarium graminearum*. Genome 48 (3):521-533.
344   doi:10.1139/g05-014
345   Arunachalam C, Doohan FM (2013) Trichothecene toxicity in eukaryotes: Cellular
346   and molecular mechanisms in plants and animals. Toxicol Lett 217 (2):149-
347   158. doi:10.1016/j.toxlet.2012.12.003
348   Backhouse D (2014) Global distribution of *Fusarium graminearum*, *F. asiaticum* and
349   *F. boothii* from wheat in relation to climate. Eur J Plant Pathol 139 (1):161-
350   173. doi:10.1007/s10658-013-0374-5
351   Boutigny AL, Ward TJ, Van Coller GJ, Flett B, Lamprecht SC, O'Donnell K, Viljoen A
352   (2011) Analysis of the *Fusarium graminearum* species complex from wheat,
353   barley and maize in South Africa provides evidence of species-specific
354   differences in host preference. Fungal Genet Biol 48 (9):914-920.
355   doi:10.1016/j.fgb.2011.05.005
356   Brauner PC, Melchinger AE, Schrag TA, Utz HF, Schipprack W, Kessel B, Ouzunova M,
357   Miedaner T (2017) Low validation rate of quantitative trait loci for Gibberella
358   ear rot resistance in European maize. Theor Appl Genet 130 (1):175-186.
359   doi:10.1007/s00122-016-2802-3
360   Chen Q, Song J, Du WP, Xu LY, Jiang Y, Zhang J, Xiang XL, Yu GR (2017) Identification,
361   mapping, and molecular marker development for Rgsr8.1: a new quantitative
362   trait locus conferring resistance to Gibberella stalk rot in maize (Zea mays
363   L.). Front Plant Sci 8:1355. doi:10.3389/fpls.2017.01355
364   Clark RT, Famoso AN, Zhao KY, Shaff JE, Craft EJ, Bustamante CD, Mccouch SR,
365   Aneshansley DJ, Kochian LV (2013) High-throughput two-dimensional root
366   system phenotyping platform facilitates genetic analysis of root growth and
367   development. Plant Cell Environ 36 (2):454-466. doi:10.1111/j.1365-
368   3040.2012.02587.x
369   Cuomo CA, Guldener U, Xu JR, Trail F, Turgeon BG, Di Pietro A, Walton JD, Ma LJ,
370   Baker SE, Rep M, Adam G, Antoniw J, Baldwin T, Calvo S, Chang YL, Decaprio
371   D, Gale LR, Gnerre S, Goswami RS, Hammond-Kosack K, Harris LJ, Hilburn K,
372   Kennell JC, Kroken S, Magnuson JK, Mannhaupt G, Mauceli E, Mewes HW,
373   Mitterbauer R, Muehlbauer G, Munsterkotter M, Nelson D, O'Donnell K,
374   Ouellet T, Qi W, Quesneville H, Roncero MI, Seong KY, Tetko IV, Urban M,
375   Waalwijk C, Ward TJ, Yao J, Birren BW, Kistler HC (2007) The *Fusarium*
376   *graminearum* genome reveals a link between localized polymorphism and
377   pathogen specialization. Science 317 (5843):1400-1402.
378   doi:10.1126/science.1143708
379   Kebede AZ, Woldemariam T, Reid LM, Harris LJ (2016) Quantitative trait loci
380   mapping for Gibberella ear rot resistance and associated agronomic traits
381   using genotyping-by-sequencing in maize. Theor Appl Genet 129 (1):17-29.
382   doi:10.1007/s00122-015-2600-3
383   Kuhnem PR, Del Ponte EM, Dong YH, Bergstrom GC (2015) *Fusarium graminearum*
384   ssolates from wheat and maize in New York show similar range of

aggressiveness and toxigenicity in cross-species pathogenicity tests.
Phytopathology 105 (4):441-448. doi:10.1094/Phyto-07-14-0208-R
Lee SH, Lee J, Nam YJ, Lee S, Ryu JG, Lee T (2010) Population structure of *Fusarium*
graminearum from maize and rice in 2009 in Korea. Plant Pathology J 26
(4):321-327. doi:10.5423/Ppj.2010.26.4.321
Ma C, Ma X, Yao L, Liu Y, Du F, Yang X, Xu M (2017) qRfg3, a novel quantitative
resistance locus against Gibberella stalk rot in maize. Theor Appl Genet 130
(8):1723-1734. doi:10.1007/s00122-017-2921-5
Maresca M (2013) From the gut to the brain: journey and pathophysiological effects
of the food-associated trichothecene mycotoxin deoxynivalenol. Toxins 5
(4):784-820. doi:10.3390/toxins5040784
McMullen M, Bergstrom G, De Wolf E, Dill-Macky R, Hershman D, Shaner G, Van
Sanford D (2012) A unified effort to fight an enemy of wheat and barley:
Fusarium head blight. Plant Dis 96 (12):1712-1728. doi:10.1094/Pdis-03-12-
0291-Fe
McMullen MD, Kresovich S, Villeda HS, Bradbury P, Li H, Sun Q, Flint-Garcia S,
Thornsberry J, Acharya C, Bottoms C, Brown P, Browne C, Eller M, Guill K,
Harjes C, Kroon D, Lepak N, Mitchell SE, Peterson B, Pressoir G, Romero S,
Oropeza Rosas M, Salvo S, Yates H, Hanson M, Jones E, Smith S, Glaubitz JC,
Goodman M, Ware D, Holland JB, Buckler ES (2009) Genetic properties of the
maize nested association mapping population. Science 325 (5941):737-740.
doi:10.1126/science.1174320
Mueller D (2016a) Corn disease loss estimates from the United States and Ontario,
Canada - 2013. Purdue University Extension, Retrieved from
http://cropprotectionnetwork.org/crop-loss-estimates/corn-disease-loss-
estimates-2013/.
Mueller D (2016b) Corn disease loss estimates from the United States and Ontario,
Canada - 2014. Purdue University Extension, Retrieved from
http://cropprotectionnetwork.org/crop-loss-estimates/corn-disease-loss-
estimates-2014/.
Mueller D (2016c) Corn disease loss estimates from the United States and Ontario,
Canada - 2015. Purdue University Extension, Retrieved from
http://cropprotectionnetwork.org/crop-loss-estimates/corn-disease-loss-
estimates-2015/.
Mueller D (2017) Corn disease loss estimates from the United States and Ontario,
Canada - 2016. Purdue University Extension, Retrieved from
http://cropprotectionnetwork.org/crop-loss-estimates/corn-disease-loss-
estimates-2016/.
Munkvold GP, White DG (2016) Compendium of Corn Diseases. Fourth edn. The
American Phytopathological Society,
Navarro L, Bari R, Achard P, Lison P, Nemri A, Harberd NP, Jones JDG (2008) DELLAs
control plant immune responses by modulating the balance and salicylic acid
signaling. Curr Biol 18 (9):650-655. doi:10.1016/j.cub.2008.03.060
Sampietro DA, Diaz CG, Gonzalez V, Vattuone MA, Ploper LD, Catalan CAN, Ward TJ
(2011) Species diversity and toxigenic potential of *Fusarium graminearum*

430   complex isolates from maize fields in northwest Argentina. Int J Food
431   Microbiol 145 (1):359-364. doi:10.1016/j.ijfoodmicro.2010.12.021
432 Trail F (2009) For Blighted Waves of Grain: *Fusarium graminearum* in the
433   Postgenomics Era. Plant Physiol 149 (1):103-110.
434   doi:10.1104/pp.108.129684
435 Wang JH, Ndoye M, Zhang JB, Li HP, Liao YC (2011) Population structure and Genetic
436   diversity of the *Fusarium graminearum* species complex. Toxins 3 (8):1020-
437   1037. doi:10.3390/toxins3081020
438 Yabuta T, Sumiki Y (1938) On the crystal of gibberellin, a substance to promote
439   plant growth. Journal of the Agricultural Chemical Society of Japan 14 (1526)
440 Yang Q, Yin GM, Guo YL, Zhang DF, Chen SJ, Xu ML (2010) A major QTL for resistance
441   to Gibberella stalk rot in maize. Theor Appl Genet 121 (4):673-687.
442   doi:10.1007/s00122-010-1339-0
443 Ye JR, Guo YL, Zhang DF, Zhang N, Wang C, Xu ML (2013) Cytological and molecular
444   characterization of quantitative trait locus qRfg1, which confers resistance to
445   Gibberella stalk rot in maize. Mol Plant Microbe In 26 (12):1417-1428.
446   doi:10.1094/Mpmi-06-13-0161-R
447 Zhang DF, Liu YJ, Guo YL, Yang Q, Ye JR, Chen SJ, Xu ML (2012) Fine-mapping of
448   qRfg2, a QTL for resistance to Gibberella stalk rot in maize. Theor Appl Genet
449   124 (3):585-596. doi:10.1007/s00122-011-1731-4
450 Zhang Y, He J, Jia LJ, Yuan TL, Zhang D, Guo Y, Wang Y, Tang WH (2016) Cellular
451   Tracking and gene profiling of *Fusarium graminearum* during maize stalk rot
452   disease development elucidates its strategies in confronting phosphorus
453   limitation in the host apoplast. PLoS Pathog 12 (3):e1005485.
454   doi:10.1371/journal.ppat.1005485
455

456 **Table 1.** Description of hybrid lines from the Blue River Organic Seeds Company
457

458
459
460

Hybrid ID	Days to Flowering (Maturation Time)	Seedling Vigor Score (Out of 5)
Hybrid I	86 (<90)	4
Hybrid II	88 (<90)	4
Hybrid III	94 (90-100)	3
Hybrid IV	97 (90-100)	4
Hybrid V	99 (90-100)	3
Hybrid VI	106 (100-110)	4
Hybrid VII	108 (100-110)	4
Hybrid VIII	114 (>110)	3
Hybrid IX	116 (>110)	3

474
475
476
477
478 **Table 2.** Sources of tested *F. graminearum* isolates
479

Isolate ID	Host of Origin	Collection Location (Year)
Gz014	Wheat	Varna, NY (1998)
Gz829	Maize	Union Springs, NY (2011)
Gz835	Maize	Union Springs, NY (2011)

480
481
482

483 **Table 3.** Two-way ANOVA result tables of *Fusarium graminearum*-induced
484 morphological changes and seedling survival rates, by fungal isolates and different
485 properties of the maize hybrids.
486

Shoot Height Ratio (9 DPI)

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.9316	0.4658	15.271	2.41E-04
Maize Maturation Time	0.1248	0.0416	1.364	0.292
Interactive	0.0415	0.0069	0.227	0.962
Residuals	0.4575	0.0305		

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.9316	0.4658	17.03	4.02E-05
Seedling Vigor	0.032	0.032	1.172	0.291
Interactive	0.0174	0.0087	0.319	0.731
Residuals	0.5744	0.0274		

Total Root Length Ratio (9 DPI)

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.4435	0.22174	26.486	1.20E-05
Maize Maturation Time	0.0353	0.01175	1.404	0.280
Interactive	0.0523	0.00872	1.042	0.437
Residuals	0.1256	0.00837		

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.4435	0.22174	23.923	3.85E-06
Seedling Vigor	0	0.00001	0.001	0.979
Interactive	0.0185	0.00926	0.999	0.385
Residuals	0.1947	0.00927		

Seedling Survival Rate (14 DPI)

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.7954	0.3977	4.247	3.46E-02
Maize Maturation Time	0.066	0.022	0.235	0.871
Interactive	0.4366	0.0728	0.777	0.6
Residuals	1.4048	0.0937		

	Sum Sq.	Mean Sq.	F value	P value
Fungal Isolate	0.7954	0.3977	5.518	1.19E-02
Seedling Vigor	0.0802	0.0802	1.112	0.3035
Interactive	0.3136	0.1568	2.176	0.1384
Residuals	1.5136	0.0721		

Figure 1. ***Fusarium graminearum* inoculation induces shoot elongation and root reduction in maize seedlings in a dosage- and host genotype-dependent manner.** Shoot height (A) and total root length (B) measurements on seedlings inoculated at each Gz014 fungal spore concentration were compared to mock-treated seedlings with Dunnett's tests relative to mock-infected controls (N = 6 for each group; * p < 0.05). All measurements were taken 9 days post-inoculation. Fungal spore concentration is in count/milliliter. s.d. = standard deviation.

Figure 2. *Fusarium graminearum* **inoculation induces shoot elongation and root reduction in maize inbred line B73.** Representative photos of shoots (A) and roots (B) mock- and Gz014-inoculated B73 seedlings were taken 9 days post-inoculation. Shoot height (C) and total root length (D) were measured on the same day and compared with Student's *t*-tests (N = 6 for each group; * p < 0.05). s.d. = standard deviation.

Figure 3. ***Fusarium graminearum*-induced morphological changes in maize seedlings vary across maize inbred lines.** Root length (A) and shoot height (B) of each maize inbred line were compared to measurement from a corresponding mock treatment group with Student's *t*-test (* $p < 0.05$). The ratio of fungus- and mock-inoculated shoot height and root length were plotted, with the dotted line denoting the expected ratio if there is no significant difference between the two groups. All measurements were taken at 9 days post-inoculation. s.d. = standard deviation.

Figure 4. *Fusarium graminearum*-**induced morphological changes in maize seedlings vary with both plant and fungal genotype.** Nine maize hybrid lines were inoculated with three *F. grmainarum* isolates (Gz014, Gz829, and Gz835). Root length (A) and shoot height (B) of each maize hybrid-fungus isolate pair were compared to measurement from a corresponding mock treatment group with Student's *t*-test (* p < 0.05). The ratios of fungus- and mock-inoculated root length and shoot height were plotted, with the dotted line denoting the expected ratio if there is no significant difference between the two groups. All measurements were taken at 9 days post-inoculation. s.d. = standard deviation.

Figure 5. *Fusarium graminearum*-**induced morphological changes and seedling blight severity vary significantly between fungal isolates.** The extent of *F.* graminearum-induced shoot elongation and root reduction at 9 days post-inoculation (A & B) and survival rates of maize seedlings inoculated with different *F. graminearum* isolates at 14 days post-inoculation (C) were compared using two-way analysis of variance with Tukey's HSD. Significantly different groups are denoted with different letters. s.d. = standard deviation.

Figure 6. *Fusarium graminearum*-**induced morphological changes and seedling blight severity are significantly correlated with each other.** Each mark represents a maize hybrid-fungal isolate pair, with maize hybrids inoculated with different fungal isolates represented by different mark shapes.