A taxonomically-restricted Uric Acid transporter provides insight into antioxidant

equilibrium

Diogo Oliveira*[,1], André Machado*[,1], Tiago Cardoso[1], Mónica Lopes-Marques[1], L.

Filipe C. Castro[1,2]• and Raquel Ruivo[1] •

[1]CIIMAR – Interdisciplinary Centre of Marine and Environmental Research, U.

Porto – University of Porto, Porto, Portugal

[2]Department of Biology, Faculty of Sciences, U. Porto - University of Porto,

Portugal

*Equal contribution

•Corresponding authors at: CIIMAR, Terminal de Cruzeiros do Porto de Leixões,

Av. General Norton de Matos s/n, 4450-208 Matosinhos, Portugal. Tel.: +351 223

401 831

Running head: Novel uric acid transporter in fish and amphibians

Abstract

Nucleobase-Ascorbate Transporter (NAT) family includes ascorbic acid, nucleobases and uric acid transporters, with a broad evolutionary distribution. In vertebrates, four members have been previously recognized, the ascorbate transporters Slc23a1 and Slc3a2, the nucleobase transporter Slc23a4 and an orphan transporter SLC23A3. Here we identify a fifth member of the vertebrate *slc23* complement (*slc23a5*), expressed in neopterygians (gars and teleosts) and amphibians, and clarify the evolutionary relationships between the novel gene and known *slc23* genes. Further comparative analysis puts forward uric acid as the preferred substrate for Slc23a5. Gene expression quantification suggests kidney and testis as major expression sites in *Xenopus tropicalis* (western clawed frog) and *Danio rerio* (zebrafish). Additional expression in brain was detected in *D. rerio*, while in the Neoteleostei *Oryzias latipes* (medaka) *slc23a5* expression is restricted to brain. The biological relevance of the retention of an extra transporter in fish and amphibians is examined: with respect to the (1) antioxidant role of uric and ascorbic acid in seminal fluid and brain, (2) the ability to endogenously synthesize ascorbic acid and (3) the morphological adaptations of the male urogenital system.

Keywords: Nucleobase-Ascorbate Transporters, Slc23a5, Uric acid, Antioxidant, Gene duplication, Molecular evolution, Comparative studies

Introduction

Introduction

Ascorbic acid (or its salt form ascorbate) is an essential enzyme cofactor,

participating in collagen and norepinephrine synthesis, as well as a potent antioxidant

and free radical scavenger (Burzle et al., 2013, Linster & Van Schaftingen, 2007). Despite

its importance, anthropoid primates, teleost fish, Guinea pigs, some bats and

Passeriformes bird species, lack a functional L-gulono-ϒ-lactone oxidase (*GLO*) gene and,

thus, the ability to catalyse the last step of ascorbic acid synthesis (Drouin et al., 2011,

Dutta Gupta et al., 1973). The scattered loss of the *GLO* gene was suggested to evolve

neutrally, with ascorbic acid requirements being met by dietary intake (Drouin et al.,

2011). Unlike ascorbate, uric acid is largely regarded as end-product of purine

metabolism and metabolic waste (Wright, 1995). Yet, human-biased research has

suggested alternative physiological functions for uric acid or urate (uric acid salts)

(Bobulescu & Moe, 2012). Due to loss of the *uricase* (UOX) enzyme in hominids, which

converts uric acid from hepatic nucleotide metabolism into the excretion product

allantoin, humans have the highest uric acid levels among animals (Wu et al., 1992,

Bobulescu & Moe, 2012, Anzai et al., 2012, Alvarez-Lario & Macarrón-Vicente, 2001, Oda

et al., 2002). In hominids, the fate of *UOX* followed *GLO* silencing, suggestive of an

increased selective pressure for an alternative anti-oxidant system (Bobulescu & Moe,

2012, Alvarez-Lario & Macarrón-Vicente, 2001). In humans, uric acid is thus proposed to

act as antioxidant and free radical scavenger, similarly to ascorbic acid, counterbalancing

disease-related oxidative stress (Bobulescu & Moe, 2012, Alvarez-Lario & Macarrón-

Vicente, 2001, Oda et al., 2002)

Ascorbate occurs in two forms. Uptake and cellular trafficking of the oxidized form

(dehydroascorbic acid) involves the facilitative hexose transporters comprising the

glucose transporters (GLUT1, GLUT3 and GLUT4), belonging to the Solute Carrier family

2 (SLC2) (Burzle et al., 2013, Corti et al., 2010). Ascorbic acid levels (reduced form), on

the other hand, are maintained by an active mechanism that requires sodium-

dependent transporters from the SLC23 family: Slc23a1 responsible for renal and

intestine ascorbic acid absorption across brush border epithelia and Slc23a2, which

exhibits a widespread expression and mediates ascorbic acid supply, from blood and

fluids, to target tissues and cells, including neurons (Burzle et al., 2013, Corti et al., 2010,

Savini et al., 2008).   Slc23a1 and Slc23a2 belong to the Nucleobase-Ascorbate

Transporter (NAT) family, which has a very broad evolutionary distribution, from

bacteria to metazoans, and displays a highly conserved signature motif defining

substrate specificity: [Q/E/P]-N-x-G-x-x-x-x-T-[R/K/G] (Burzle et al., 2013, Koukaki et al.,

2005, Papakostas & Frillingos, 2012). In vertebrates, two additional transporters were

identified, the orphan *transporter* Slc23a3 and the nucleobase transporter Slc23a4,

described in rats but found pseudogenized in humans (Burzle et al., 2013, Yamamoto et

al., 2010). Rat Slc23a4 is restricted to the small intestine and was shown to

predominantly mediate sodium-dependent uracil transport (Yamamoto et al., 2010).

Human Slc23a3, on the other hand, is expressed in proximal renal tubules but is

unresponsive to both ascorbic acid and nucleobases (Burzle et al., 2013). Besides the

described animal transporters, the NAT family includes several transport proteins from

bacteria, fungi and plants, which can be divided into three functional groups (Figure 1).

Slc23a1, Slc23a2 and Slc23a3 belong to the ascorbate group and harbour a conserved

proline (P) in the first amino acid position of the signature motif (Burzle et al., 2013). A

glutamate (E) residue defines the uracil group, including bacterial anion symporters and

Slc23a4, while a glutamine (Q) is present in the bacterial uric acid and/or xanthine anion

symporters (Koukaki et al., 2005, Yamamoto et al., 2010, Burzle et al., 2013). Here we

identify a novel member of the NAT family, *slc23a5*, expressed in neopterygian fishes

(gars and teleosts) and amphibians. Combining relative gene expression patterns and

comparative analyses of protein motifs, we put forward the functional relevance of this

taxonomically restricted gene in the context of oxidative stress in the aquatic

environment.

Materials and Methods

Sequence mining

Slc23 protein sequences were obtained from the GenBank database by BLASTp

using human Slc23a1, Slc23a2 and Slc23a3 sequences as query. The sequences from

Lethenteron japonica (Japanese lamprey) were obtained from the Japanese Lamprey

Genome Project. The sequences from *Leucoraja erinacea* and *Scyliorhinus canicula* were

obtained as nucleotide sequences from SkateBase and translated. Accession numbers

for all the retrieved sequences are listed in Supplementary Table 1 (Table S1).

Synteny and paralogy

Synteny and paralogy maps were constructed based on information about the

genomic location of the *Slc23* members' flanking genes from the GenBank database.

Paralogs were identified according to the information available on Ensembl, and

ancestral chromosomal partitioning was established as per reference (Nakatani et al.,

2007, Putnam et al., 2008).

Phylogenetic Analysis

The retrieved sequences were aligned using the MAFFT software web service

(http://mafft.cbrc.jp/alignment/software/) in default automatic settings, with L-INS-I

refinement method (Katoh & Standley, 2013). The output alignment was used to

construct a phylogenetic tree using PhyML 3.0 with Smart Model Selection web service

(http://www.atgc-montpellier.fr/phyml-sms/) in default settings, which selected the LG

+G+I+F model (Guindon et al., 2010).

Membrane topology prediction

Slc23a1 and Slc23a5 membrane topology prediction was performed using the

CCTOP prediction web server (http://cctop.enzim.ttk.mta.hu/) (Dobson et al., 2015).

Filtering and quality control of SRA datasets

For Rna-seq analysis, we included 39 SRA datasets from brain, intestine, kidney,

liver, and testis tissues of Human (*Homo sapiens*), mouse (*Mus musculus*), chicken

(*Gallus gallus*), lizard (*Anolis carolinensis*), western clawed frog (*Xenopus tropicalis*),

zebrafish (*Danio rerio*), spotted gar (*Lepisosteus oculatus*) and Japanese medaka (*Oryzias*

latipes), which were obtained from the National Center for Biotechnology (NCBI)

Sequence Read Archive (*SRA)* (https://www.ncbi.nlm.nih.gov/sra/) (Table S2). The read

quality assessment of each SRA dataset was performed using FastQC software

(https://www.bioinformatics. Babraham.ac.uk/projects/fastqc/). Trimmomatic 0.36

(Bolger et al., 2014) was used to trim and drop reads with quality scores below 5, at

leading and trailing ends, with an average quality score below 20 in a 4 bp sliding window

and with less than 36 bases length.

Relative gene expression levels

To assess the relative gene expression of *slc23* genes from the selected species,

reference sequences and the corresponding annotations were collected from NCBI and

Ensemble (Yates et al., 2016) databases. For human, mouse, lizard, zebrafish and

spotted gar the GTF and DNA files (genome) were retrieved from Ensemble (Release 89)

(Yates et al., 2016). For chicken and western clawed frog transcript sequences (coding

and non-coding) and mapping (gene/transcript) from NCBI Refseq database were used

after filtering (O'Leary et al., 2016) (for each species the rna.fasta file from

ftp://ftp.ncbi.nih.gov/genomes/refseq/ was used – See Table S3 for detail). The SRA

reads of each specimen were aligned to the reference using Bowtie2 (default

parameters)(Langmead & Salzberg, 2012), and gene expression was quantified as

transcript per million (TPM) using RSEM v.1.2.31 (RNA-seq by Expectation Maximization)

software package (Li & Dewey, 2011). Isoform expression levels for each gene were

summed to derive the TPM values. For genes with low relative expression values TPM

values were $\log_2$-transformed after adding a value of one; TPM values < 0.5 were

considered unreliable and substituted with zero.

PCR-based Slc23a5 expression

Gene expression of *Slc23a5* was further analyzed by PCR for *Xenopus tropicalis* and

Danio rerio using the primer pairs 5'-AGACGTTGAAGCCGTTACTGA-3' and 5'-

TGCTATTATGCCTCCAAAGGCT-3' in the former and 5'-GGTGGAATGTTCCTCGTCAT-3' and

5'-GATGAACATGTGGGTGGTGA-3' in the latter. These primers map to a small gene

region flanking an intron. PCR reactions were carried out on a tissue panel containing

heart, skin, stomach, intestine, liver, kidney, ovary and testis tissues for *X. tropicalis* and

heart, skin, gut, liver, kidney, ovary and testis tissues for *D. rerio*.

Results and Discussion

Through database mining and phylogenetic analysis, we detected the presence of

a novel gene, *slc23a5*, found in neopterygians (Holostei and Teleostei) and amphibians

(Fig 1). No *slc23a5* sequence was retrieved from chondrichthyans, chondrosteans (basal

actinopterygians) nor from the sarcopterygian *Latimeria chalumnae* (coelacanth).

Secondary loss of *slc23a5* in chondrichthyans was further supported by the conservation

of the genomic *locus* in *Callorhynchus milii* (Australian ghostshark) (Fig S1). Regarding

coelacanth, two scaffolds, denoting conservation of the flanking genes, were found;

while, no information was retrieved regarding the chondrostean *loci* (Fig S1). The

remaining genes of the *slc23* family were retrieved and genomic *loci* were found

generally conserved across species, with exception of the *slc23a4 loci* in fish (Fig S2-S5).

The branching pattern of our phylogenetic analysis suggests that this novel *slc23a5* gene

emerged, along with the *slc23a4* paralogue, from an ancestral duplication and that

slc23a5 was secondarily lost in sharks, basal actinopterygians, coelacanth and amniotes

(Fig 1 and Fig S1). In agreement, we obtained three phylogenetic clusters possibly

corresponding to three ancestral lineages (Fig 1): one composed of typical *slc23a1* and

slc23a2 genes and a second group including the previously reported s*lc23a4* and the

novel *slc23a5* genes, both outgrouped by invertebrate sequences. A third cluster, which

outgroups the remaining *slc23* genes, included the highly divergent *slc23a3* genes,

mirrored by the long branch lengths (Fig 1). To confirm ancestral duplication events we

conducted paralogy analysis of the human *loci* using two chromosomal partitioning

approaches (Putnam et al., 2008, Nakatani et al., 2007). While some reorganization is

observed, our combined analysis seems to support a 2R origin for the *slc23a1/slc23a2*

and *slc23a4/slc23a5* gene *loci*, while *slc23a3* might stand as an independent lineage (Fig

S6). Also, human paralogues of the genes flanking an invertebrate *slc23*-like sequence,

from *Branchiostoma belcheri*, were mapped back to the same chromosomal partition as

the human *slc23a3* sequence, corroborating the existence of a *slc23a3*-like invertebrate

gene (Fig S6). Further comparative protein sequence analysis highlighted the presence

of a NAT family signature motif with a conserved Q in substrate determining position,

placing Slc23a5 in the uric acid and/or xanthine group (Fig 2). Besides the typical

signature motif, a carboxyl terminus tripeptide motif was identified in some SLC23

family members: a S/T-X-Ø tripeptide shown to interact PDZ domains of scaffolding

proteins belonging to the Sodium/Hydrogen Exchanger Regulatory Factor, NHERF,

family (Anzai et al., 2012). NHERF family members, such as PDZ-Containing Kidney

Protein 1 (PDZK1), are mostly expressed in the apical side of absorptive epithelia of

kidney, intestine and liver, and establish protein-protein interaction networks with a

wide range of proteins, including receptors, transporters and channels (Dev, 2004,

Clapéron et al., 2011). Besides its scaffolding role, NHERF family members were also

shown to modulate the activity of protein assemblages (Clapéron et al., 2011, Raghuram

et al., 2001, Wang et al., 2000). Curiously, S/T-X-Ø are found highly conserved across

species in Slc23a1 and Slc23a4, in agreement with their renal and intestinal expression

and role in ascorbic acid and nucleobase absorption. Slc23a5, on the other hand, exhibits

a highly conserved tripeptide motif in Otocephala (Cypriniformes, Characiformes and

Siluriformes) but not Osteoglossomorpha (Fig 3). The Euteleostei Esociformes and

Salmoniformes exhibit an inverted motif while in the Neoteleostei no conservation was

found. Finally, topology predictions suggest a conserved cellular import topology, similar

to other family members (Fig S7).

Our findings indicate a restricted distribution of *slc23a5* to *taxa* that share an

aquatic nature of their habitat. Why would these retain a transporter for uric acid, or its

precursor, xanthine? Here we hypothesize that, beyond its metabolic waste nature, uric

acid may act as an antioxidant in fish and amphibians: notably in *GLO*-deficient teleost.

To gain further insight into the functional relevance of this additional gene, we collected

RNA-seq data and determined relative expression levels of *Slc23* genes. Gene expression

analysis suggested that, in *Danio rerio* (zebrafish) and *Xenopus tropicalis* (western

clawed frog), *slc23a5* is mostly expressed in kidney and testis (Fig 3 and Tables S4 and

S5). Expression in brain was also detected in *D. rerio* (zebrafish) and was found to be the

exclusive expression site in the Neoteleostei *Oryzias latipes* (medaka). The holostean

Lepisosteus oculatus (spotted gar), on the other hand, yielded no significant expression

in the tested tissues. Gene expression levels were further confirmed by PCR in *D. rerio*

and *X. tropicalis*. The gonadal expression provides a relevant clue for the retention of

the extra isoform and supports the antioxidant role of uric acid. In external fertilizers,

such as most fish and amphibians, reactive oxygen species were suggested to decrease

motility and thus fertilisation success (Lahnsteiner & Mansour, 2010). In agreement,

high concentrations of uric acid were detected, and suggested to act as the main non-

enzymatic antioxidant mechanism, in seminal fluid of several fish species: including

Cypriniformes, Esociformes, Salmoniformes and the Neoteleostei Gadiformes and

Perciformes (Ciereszko et al., 1999, Dzyuba et al., 2014b, Lahnsteiner & Mansour, 2010,

Lahnsteiner et al., 2010). Yet, regarding amphibians, no information on seminal fluid

composition was retrieved (Keogh et al., 2017). Nonetheless, *in vitro* assays suggested

that other antioxidant species, such as ascorbic acid or vitamin E, were unsuitable at

offsetting oxidative stress and improving sperm motility in *Litoria booroolongensis*

(Booroolong frog) and suggested uric acid as a pertinent candidate (Keogh et al., 2017).

In this context, the absence of *slc23a5* sequences in Chondrichthyes is possibly due to

their internal fertilization strategies derived from ovoviviparous or viviparous

reproductive modes (Goodwin et al., 2002). Accordingly, viviparity is also observed in

the sarcopterygian *L. chalumnae* (Goodwin et al., 2002).

Besides reproductive categories, other morphological and metabolic traits can

justify the seemingly scattered retention of *slc23a5* among fish; for instance, specific

anatomical features such as the presence of a urogenital connection: described in

Elasmobranchii and in most non-teleost Osteichthyes, yet absent in Cyclostomata and

Teleostei (Blün, 1986). By connecting testicular and renal ducts, antioxidant

requirements could be met by renal secretion into urine. In agreement, in the

Chondrostei *Acipenser ruthenus*, full sperm maturation and motility was shown to

require dilution by urine (Dzyuba et al., 2014a). An urogenital connection is also present

in amphibians (Blün, 1986). Additionally, other antioxidant mechanisms could exist.

Besides uric acid, ascorbic acid is also present in seminal fluid of *Oncorhynchus mykiss*

(rainbow trout) (Ciereszko & Dabrowski, 1995). Seminal ascorbate was reported to be

affected by seasonal variations and semen quality was shown to improve upon dietary

ascorbic acid supplementation (Ciereszko & Dabrowski, 2000, Ciereszko & Dabrowski,

1995), contrasting with the amphibian *L. booroolongensis* (Keogh et al., 2017). In the

case of non-teleost fish, such as Chondrostei (*A. ruthenus*) and Holostei (*L. oculatus*), a

functional *GLO* gene is retained, providing endogenously synthesised ascorbate

(Moreau & Dabrowski, 2000). Curiously, no gonadal expression was detected in the

Neoteleostei *O. latipes*. Neoteleosts exhibit varied reproduction modes ranging from

external fertilizers to internal fertilizers with viviparity (Forsman et al., 2015, Bone &

Moore, 2008, Goodwin et al., 2002, Scott & Fuller, 1976). Also, neoteleosts were shown

to exhibit a distinctive gonadal morphology, the lobular testis, whereas the remaining

bony fish exhibit tubular testis (Parenti & Grier, 2004). Although still obscure, the

absence of testicular *slc23a5* expression in the externally fertilizer *O. latipes* could be

due to specific morphological adaptations of neoteleosts.

Besides the gonadal expression, we detected significant expression of *slc23a5* in

teleost brain (Fig 3). Again, an antioxidant role can be postulated for neuronal uric acid

(Uttara et al., 2009). Active swimming behaviour requires a high metabolic rate,

mirrored by an increased oxygen consumption, subsequently leading to reactive oxygen

species production (Lopez-Cruz et al., 2011). In agreement, fish antioxidant levels were

shown to be correlated with their swimming activity (Lopez-Cruz et al., 2011). Neuronal

cell populations are particularly sensitive to oxidative stress, thus, uric acid could

participate in the neutralization of oxidative species to by-pass oxidative stress, notably

in *GLO*-deficient fish. Yet, information on neuronal uric acid levels in fish is currently

absent.

Additionally, renal Slc23a5, detected in *D. rerio* and *X. tropicalis* could participate

in the maintenance of uric acid, or xanthine, levels for antioxidant purposes, osmotic

regulation and/or excretion. Kidneys play a crucial role in the maintenance of both

ascorbate and urate levels through the excretion and reabsorptions processes (Anzai et

al., 2012, Bobulescu & Moe, 2012). Regarding urate, several transporters, localized to

specific cellular regions (apical or basolateral), seem to cooperate for efficient urate

handling in the human proximal renal tubules: the urate transportome (Anzai et al.,

2012, Bobulescu & Moe, 2012). In kidney absorptive epithelium, transporters located on

the filtrate-epithelial interface (apical side) interact with the scaffolding protein PDZK1

(Anzai et al., 2012). A PDZK1 interacting motif was found in Slc23a5 highly conserved in

Cypriniformes, Characiformes and Siluriformes, but not in other fish groups or

amphibians (Fig 4). No relationship between salt and freshwater environments was

found. Thus, the presence/absence of this peptide signal might reflect distinct uric acid

flows in the kidney: an apically expressed Slc23a5 likely participates in absorptive uric

acid flows, while a basolateral expression would be related to uric acid secretion (Fig 4).

In agreement, *X. tropicalis*, *D. rerio* and *L. oculatus* appear to lack the full complement

of renal urate transporters (Table S6). A similar scenario is observed for birds and

reptiles; however, their excretory system represents a particular adaptation for uric acid

excretion (Braun, 1998, Singer, 2003). With this in mind, *X. tropicalis* and *D. rerio*

illustrate the two opposing mechanisms: while *D. rerio* captures, *X. tropicalis* secretes

uric acid into renal filtrate. Besides simple excretion, and given the renal-testicular

morphology of amphibians, uric acid secretion into urine could also promote antioxidant

build-up for reproductive purposes.

Curiously, despite its inability to mediate transport, Slc23a3 is also expressed in

proximal renal tubules, along Slc23a1 and Slc23a5 (Burzle et al., 2013). Comparative

sequence analysis places this orphan *transporter* in the ascorbic acid group, due to

conservation of the P in the first amino acid position of the NAT family signature motif,

crucial for substrate selection. Yet, significant sequence variation was observed in

Teleostei and Sauropsida (Fig 1). A conserved, yet apparently disruptive, Q was also

evident in the last amino acid position of the signature motif: as a mirror image of the

conserved P (Fig 1). An interesting, yet untested, hypothesis suggests Slc23a3 as a sensor

for downstream regulation of transportome expression. Slc23a1 expression is, indeed,

modulated by ascorbic acid levels (Burzle et al., 2013). Transporter to transreceptor

shifts, leading to nutrient sensing by transporter homologues, were previously

acknowledged in yeast (Kriel et al., 2011, Karhumaa et al., 2010, Poulsen et al., 2005,

Wu et al., 2006). Thus, renal ascorbic and uric acid handling could share common

mechanistic features and future studies should address the cooperative handling, if any,

of uric and ascorbic acid in kidney proximal tubules.

Taken together, these observations put forward the novel Slc23a5 as a uric acid

transporter and support the antioxidant role of uric acid, notably in GLO-deficient fish.

The retention of an additional SLC23 gene, with presumed uric acid transport function,

seems to parallel antioxidant requirements for (1) aquatic reproductive strategies, by

maintaining semen integrity in external fertilizers, and (2) to counterbalance increasing

metabolic rates, notably in the brain of active swimmers. These observations further

support the (1) role of both ascorbic and uric acids in the maintenance of an optimum

body redox potential, and the (2) need to equilibrate antioxidant scavenging

requirements. Additionally, our results emphasize the morphological and physiological

diversity observed in kidneys and gonads, as well as the distinct levels of elaboration and

interaction between these ontogenically related organs across evolutionary groups. In

aquatic animals, and developing amniotes, excretory functions are mainly orchestrated

by the mesonephric kidney (Blün, 1986). During amniote ontogeny, the mesonephros

relays its excretory function to the metanephros, with the mesonephros giving rise to a

separate seminal duct (Blün, 1986). Within aquatic animals, distinct levels of complexity

were described regarding urine and semen transportation: with contrasting urogenital

connections between teleosts and non-teleosts (Blün, 1986). Thus, although uric acid

does seem beneficial for sperm motility and survival in aquatic animals, the means by

which this antioxidant, and other antioxidant mechanisms, are maintained and

conveyed to the seminal fluid appears diverse and should be further explored.

Supplementary material: Supplementary tables and figures including synteny and

paralogy analysis and relative gene expression levels, as well as reference tables.

Acknowledgments

This work was supported by the MarInfo – Integrated Platform for Marine Data

Acquisition and Analysis (reference NORTE-01-0145-FEDER-000031), a project by the

North Portugal Regional Operational Program (NORTE 2020), under the PORTUGAL 2020

Partnership Agreement, through the European Regional Development Fund (ERDF) and

by the European Regional Development Fund (ERDF) through the COMPETE -

Operational Competitiveness Programme and POPH - *Operational Human Potential*

Programme.

References

Alvarez-Lario, B. & Macarrón-Vicente, J. (2001) Evolution of Uric Acid Metabolism in Humans.
347   In: *eLS.* pp. John Wiley & Sons, Ltd.
348   Anzai, N., Jutabha, P., Amonpatumrat-Takahashi, S. & Sakurai, H. 2012. Recent advances in renal
349   urate transport: characterization of candidate transporters indicated by genome-wide
350   association studies. *Clin. Exp. Nephrol.* **16**: 89-95.
351   Blün, V. (1986) Comparative Anatomy of the Urogenital System. In: *Vertebrate Reproduction, a*
352   *Textbook.* pp. 43-74. Springer-Verlag, Berlin Heidelberg New York Tokyo.
353   Bobulescu, I. A. & Moe, O. W. 2012. Renal transport of uric acid: evolving concepts and
354   uncertainties. *Adv. Chronic Kidney Dis.* **19**: 358-71.
355   Bolger, A. M., Lohse, M. & Usadel, B. 2014. Trimmomatic: a flexible trimmer for Illumina
356   sequence data. *Bioinformatics* **30**: 2114-20.
357   Bone, Q. & Moore, R. H. 2008. *Biology of Fishes* Taylor & Francis Group, New York.
358   Braun, E. J. 1998. Comparative renal function in reptiles, birds, and mammals. *Semin. Avian*
359   *Exotic. Pet. Med.* **7**: 62-71.
360   Burzle, M., Suzuki, Y., Ackermann, D., Miyazaki, H., Maeda, N., Clemencon, B., et al. 2013. The
361   sodium-dependent ascorbic acid transporter family SLC23. *Mol. Aspects Med.* **34**: 436-
362   54.
363   Ciereszko, A. & Dabrowski, K. 1995. Sperm quality and ascorbic acid concentration in rainbow
364   trout semen are affected by dietary vitamin C: an across-season study. *Biol. Reprod.* **52**:
365   982-8.
366   Ciereszko, A. & Dabrowski, K. 2000. Effect of ascorbic acid supplement in vitro onrainbow trout
367   sperm viability. *Aquacult. Int.* **8**: 1-8.
368   Ciereszko, A., Dabrowski, K., Kucharczyk, D., Dobosz, S., Goryczko, K. & Glogowski, J. 1999. The
369   presence of uric acid, an antioxidantive substance, in fish seminal plasma. *Fish Physiol.*
370   *Biochem.* **21**: 313-315.
371   Clapéron, A., Mergey, M. & Fouassier, L. 2011. Roles of the scaffolding proteins NHERF in liver
372   biology. *Clin. Res. Hepatol. Gastroenterol.* **35**: 176-181.
373   Corti, A., Casini, A. F. & Pompella, A. 2010. Cellular pathways for transport and efflux of
374   ascorbate and dehydroascorbate. *Arch. Biochem. Biophys.* **500**: 107-15.
375   Dev, K. K. 2004. Making protein interactions druggable: targeting PDZ domains. *Nat. Rev. Drug*
376   *Discov.* **3**: 1047-1056.
377   Dobson, L., Remenyi, I. & Tusnady, G. E. 2015. CCTOP: a Consensus Constrained TOPology
378   prediction web server. *Nucleic Acids Res.* **43**: W408-12.
379   Drouin, G., Godin, J. R. & Page, B. 2011. The genetics of vitamin C loss in vertebrates. *Curr.*
380   *Genomics* **12**: 371-8.
381   Dutta Gupta, S., Choudhury, P. K. & Chatterjee, I. B. 1973. Synthesis of l-ascorbic acid from d-
382   glucurono-1,4-lactone conjugates by different species of animals. *Int. J. Biochem.* **4**: 309-
383   314.
384   Dzyuba, B., Cosson, J., Boryshpolets, S., Bondarenko, O., Dzyuba, V., Prokopchuk, G., et al. 2014a.
385   In vitro sperm maturation in sterlet, Acipenser ruthenus. *Reprod. Biol.* **14**: 160-3.
386   Dzyuba, V., Dzyuba, B., Cosson, J., Boryshpolets, S., Yamaner, G., Kholodniy, V., et al. 2014b. The
387   antioxidant system of sterlet seminal fluid in testes and Wolffian ducts. *Fish Physiol.*
388   *Biochem.* **40**: 1731-9.
389   Forsman, A., Tibblin, P., Berggren, H., Nordahl, O., Koch-Schmidt, P. & Larsson, P. 2015. Pike Esox
390   lucius as an emerging model organism for studies in ecology and evolutionary biology:
391   a review. *J. Fish Biol.* **87**: 472-479.
392   Goodwin, N. B., Dulvy, N. K. & Reynolds, J. D. 2002. Life-history correlates of the evolution of live
393   bearing in fishes. *Philos. Trans. R. Soc. Lond. B. Biol. Sci* **357**: 259-267.

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. 2010. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst. Biol.* **59**: 307-21.
Karhumaa, K., Wu, B. & Kielland-Brandt, M. C. 2010. Conditions with high intracellular glucose
inhibit sensing through glucose sensor Snf3 in Saccharomyces cerevisiae. *J. Cell.*
Biochem. **110**: 920-5.
Katoh, K. & Standley, D. M. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**: 772-80.
Keogh, L. M., Byrne, P. G. & Silla, A. J. 2017. The effect of antioxidants on sperm motility
activation in the Booroolong frog. *Anim. Reprod. Sci.* **183**: 126-131.
Koukaki, M., Vlanti, A., Goudela, S., Pantazopoulou, A., Gioule, H., Tournaviti, S., et al. 2005. The
nucleobase-ascorbate transporter (NAT) signature motif in UapA defines the function of
the purine translocation pathway. *J. Mol. Biol.* **350**: 499-513.
Kriel, J., Haesendonckx, S., Rubio-Texeira, M., Van Zeebroeck, G. & Thevelein, J. M. 2011. From
transporter to transceptor: signaling from transporters provokes re-evaluation of
complex trafficking and regulatory controls: endocytic internalization and intracellular
trafficking of nutrient transceptors may, at least in part, be governed by their signaling
function. *Bioessays* **33**: 870-9.
Lahnsteiner, F. & Mansour, N. 2010. A comparative study on antioxidant systems in semen of
species of the Percidae, Salmonidae, Cyprinidae, and Lotidae for improving semen
storage techniques. *Aquaculture* **307**: 130-140.
Lahnsteiner, F., Mansour, N. & Plaetzer, K. 2010. Antioxidant systems of brown trout (Salmo
trutta f. fario) semen. *Anim. Reprod. Sci.* **119**: 314-21.
Langmead, B. & Salzberg, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9: 357-9.
Li, B. & Dewey, C. N. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**: 323.
Linster, C. L. & Van Schaftingen, E. 2007. Vitamin C. *FEBS J.* **274**: 1-22.
Lopez-Cruz, R. I., Dafre, A. L. & Filho, D. W. (2011) Oxidative Stress in Sharks and Rays. In:
Oxidative Stress in Aquatic Ecosystems, (Abele, D., Vazquez-Medina, J. P. & Zenteno-
Savin, T., eds.). pp. 157-164. Wiley-Blackwell.
Moreau, R. & Dabrowski, K. 2000. Biosynthesis of ascorbic acid by extant actinopterygians. *J.*
Fish Biol. **57**: 733-745.
Nakatani, Y., Takeda, H., Kohara, Y. & Morishita, S. 2007. Reconstruction of the vertebrate
ancestral genome reveals dynamic genome reorganization in early vertebrates. *Genome*
Res. **17**: 1254-1265.
O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., et al. 2016.
Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion,
and functional annotation. *Nucleic Acids Res.* **44**: D733-45.
Oda, M., Satta, Y., Takenaka, O. & Takahata, N. 2002. Loss of urate oxidase activity in hominoids
and its evolutionary implications. *Mol. Biol. Evol.* **19**: 640-53.
Papakostas, K. & Frillingos, S. 2012. Substrate selectivity of YgfU, a uric acid transporter from
Escherichia coli. *J. Biol. Chem.* **287**: 15684-95.
Parenti, L. R. & Grier, H. J. 2004. Evolution and phylogeny of gonad morphology in bony fishes.
Integr. Comp. Biol. **44**: 333-48.
Poulsen, P., Wu, B., Gaber, R. F., Ottow, K., Andersen, H. A. & Kielland-Brandt, M. C. 2005. Amino
acid sensing by Ssy1. *Biochem. Soc. Trans.* **33**: 261-4.
Putnam, N. H., Butts, T., Ferrier, D. E. K., Furlong, R. F., Hellsten, U., Kawashima, T., et al. 2008.
The amphioxus genome and the evolution of the chordate karyotype. *Nature* **453**: 1064-
1071.

Raghuram, V., Mak, D.-O. D. & Foskett, J. K. 2001. Regulation of cystic fibrosis transmembrane
conductance regulator single-channel gating by bivalent PDZ-domain-mediated
interaction. *Proc. Natl. Acad. Sci. U. S. A.* **98**: 1300-1305.
Savini, I., Rossi, A., Pierro, C., Avigliano, L. & Catani, M. V. 2008. SVCT1 and SVCT2: key proteins
for vitamin C uptake. *Amino Acids* **34**: 347-55.
Scott, D. B. C. & Fuller, J. D. 1976. The reproductive biology of Scleropages formosus (Müller &
Schlegel) (Osteoglossomorpha, Osteoglossidae) in Malaya, and the morphology of its
pituitary gland. *J. Fish Biol.* **8**: 45-53.
Singer, M. A. 2003. Do mammals, birds, reptiles and fish have similar nitrogen conserving
systems? *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **134**: 543-58.
Uttara, B., Singh, A. V., Zamboni, P. & Mahajan, R. T. 2009. Oxidative Stress and
Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant
Therapeutic Options. *Curr. Neuropharmacol.* **7**: 65-74.
Wang, S., Yue, H., Derin, R. B., Guggino, W. B. & Li, M. 2000. Accessory Protein Facilitated CFTR-
CFTR Interaction, a Molecular Mechanism to Potentiate the Chloride Channel Activity.
Cell **103**: 169-179.
Wright, P. A. 1995. Nitrogen excretion: three end products, many physiological roles. *J. Exp. Biol.*
198: 273-81.
Wu, B., Ottow, K., Poulsen, P., Gaber, R. F., Albers, E. & Kielland-Brandt, M. C. 2006. Competitive
intra- and extracellular nutrient sensing by the transporter homologue Ssy1p. *J Cell Biol*
173: 327-31.
Wu, X. W., Muzny, D. M., Lee, C. C. & Caskey, C. T. 1992. Two independent mutational events in
the loss of urate oxidase during hominoid evolution. *J. Mol. Evol.* **34**: 78-84.
Yamamoto, S., Inoue, K., Murata, T., Kamigaso, S., Yasujima, T., Maeda, J. Y., et al. 2010.
Identification and functional characterization of the first nucleobase transporter in
mammals: implication in the species difference in the intestinal absorption mechanism
of nucleobases and their analogs between higher primates and other mammals. *J. Biol.*
Chem. **285**: 6522-31.
Yates, A., Akanni, W., Amode, M. R., Barrell, D., Billis, K., Carvalho-Silva, D., et al. 2016. Ensembl
2016. *Nucleic Acids Res.* **44**: D710-6.

Figures

Figure 1 – Maximum likelihood phylogenetic tree describing relationships among

Nucleobase-Ascorbate Transporter (NAT) family members. Node values represent

branch support using the aBayes algorithm. Accession numbers for all sequences are

provided in Supplementary Materials.

Figure 2 – Alignment of the signature motif of the Nucleobase-Ascorbate Transporter (NAT) family. NAT family members display a highly conserved signature motif defining substrate specificity: [Q/E/P]-N-x-G-x-x-x-x-T-[R/K/G]. Asterisks indicate conserved positions. The first amino acid position, with residues highlighted in red within a black box, defines substrate preference: Q – Uric acid/xanthine group; P – Ascorbate group and E – Uracil group. The mirror Q in the Slc23a3 group is also highlighted by a black box.

Figure 3 – Expression patterns of *Slc23* isoforms across vertebrates. (A) RNAseq-based

quantification. Relative gene expression levels are provided as log₂-transformed

transcript per million (TPM) values, from 0 to 8 TPM's, the NF and ND means gene not

found and no SRA data available to a specific tissue. (B) PCR-based amplification of

Slc23a5 transcripts in *D. rerio* and *X. tropicalis* tissues.

Figure 4 – The S/T-X-Ø tripeptide provides insight into the cellular function of Slc23a5 homologues from fish and amphibians. (A) Schematic representation of tripeptide-dependent uric acid flows in kidney epithelium and non-polarized cells. (B) Alignment of the PDZ-binding tripeptide in Slc23a5 homologues from fish and amphibians. The S/T-X-Ø tripeptide is found in Cypriniformes, Characiformes and Siluriformes while the Euteleostei Esociformes and Salmoniformes exhibit an inverted motif. Predicted cellular localization is indicated as well as morphological and physiological adaptations among the different fish groups.

A taxonomically-restricted Uric Acid transporter provides insight into antioxidant

equilibrium

Supplementary Figures:

Figure S1 - Synteny maps of *Slc23a5*.

Figure S2 - Synteny maps of *Slc23a1*.

Figure S3 - Synteny maps of *Slc23a2*.

Figure S4 - Synteny maps of *Slc23a3*.

Figure S5 - Synteny maps of *Slc23a4*.

Figure S6 – Paralogy analysis of the human *loci*.

Figure S7 – Topology prediction of Slc23a5.

Supplementary Tables:

Table S1 - Accession numbers.

Table S2 - Accession numbers of the RNAseq files.

Table S3 - Gene identifier for all genes included in gene expression analysis,
523   **collected from NCBI and ENSEMBLE Databases.**

524   **Table S4 - Genome and GTF files retrieved from Ensemble database (Release**
525   **89) and Transcriptome files retrieved from NCBI used for this study.**

526   **Table S5 - Relative gene expression levels of Slc23 family genes for eight**
527   **species. The values are presented in log2 (TPM +1) ratios**

528   **Table S6 - Renal uric acid absorption and secretion: the urate transportome.**

Figure S1 - Synteny maps of *Slc23a5*. Conservation of the *Slc23a5* locus in major fish lineages, *Xenopus tropicalis* and *Homo sapiens*. Numbering after species corresponds to chromosome or scaffold. Information is presently absent for the Chondrostei *Acipenser ruthenus*. Grey boxes indicate non-conserved genes.

Figure S2 - Synteny maps of *Slc23a1*. Conservation of the *Slc23a1* locus in major vertebrate lineages. Numbering after species corresponds to chromosome or scaffold. Grey boxes indicate non-conserved genes.

Figure S3 - Synteny maps of *Slc23a2*. Conservation of the *Slc23a2* locus in major vertebrate lineages. Numbering after species corresponds to chromosome or scaffold. Grey boxes indicate non-conserved genes.

Figure S4 - Synteny maps of *Slc23a3*. Conservation of the *Slc23a3* locus in major vertebrate lineages. Numbering after species corresponds to chromosome or scaffold. Grey boxes indicate non-conserved genes.

Figure S5 - Synteny maps of *Slc23a4*. Conservation of the *Slc23a4* locus in major vertebrate lineages. Numbering after species corresponds to chromosome or scaffold. The human homologue is indicated as pseudogenized (*SLC23A4P*). Grey boxes indicate non-conserved genes.

Figure S6 – Paralogy analysis of the human *loci*. (A-C) Reconstruction of ancestral genome partitions according to Putnam and colleagues (Putnam, et al. 2008) and (D) according to Nakatani and colleagues (Nakatani, et al. 2007). (A-C) Numbers following gene symbols correspond to predicted chromosomal partitions grouped within linkange groups, that is, groups of genes with a predicted common chromosomal origin in the vertebrates' ancestor; corresponding paralogues of neighbouring genes are indicated below. (C) An invertebrate Slc23-like locus, from

Branchiostoma belcheri is also shown with corresponding human homologues indicated below each gene. (D) Reconstruction of duplication and fission events in the predicted ancestral lineages including the Slc23a1/Slc23a2 pair (lineage B) and the Slc23a4 and Slc23a5 pair (lineage E).

Figure S7 – Topology prediction of Slc23a5.

Supplementary Tables:

Table S1 - **Accession numbers.**

Species	Gene	Accession Number	Common name
Aspergillus nidulans	UapA	CAA50681	-
Aspergillus nidulans	UapC	CAA56190	-
Bacillus subtilis	PbuX	KIX81273	-
Bacillus subtilis	PucK	KIX81727	-
Escherichia coli	YgfO	CDU41213.1	-
Escherichia coli O25b:H4	UraA	ANK02789	-
Escherichia coli O25b:H4	YicE	ANK04183	-
Zea mays	Lep1	AAB17501	Maize
Branchiostoma belcheri	Slc23-like	XP_019630437	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019636191	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019633776	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019616280	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019646793	Amphioxus
Branchiostoma floridae	Slc23-like	XP_002600710	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002597306	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002595086	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002611161	Florida lancelet
Anolis carolinensis	Slc23a1	XP_008113637	Green anole
Callorhinchus milii	Slc23a1	XP_007904441	Ghost shark
Cyprinus carpio	Slc23a1	XP_018975086	Common carp
Danio rerio	Slc23a1	NP_001166970	Zebrafish
Gallus gallus	Slc23a1	XP_004944825	Chicken
Homo sapiens	Slc23a1	NP_005838	Human
Ictalurus punctatus	Slc23a1	XP_017348904	Channel catfish
Latimeria chalumnae	Slc23a1	XP_005989382	Coelacanth
Lepisosteus oculatus	Slc23a1	XP_015205203	Spotted gar
Lethenteron japonicum	Slc23a1	JL2986	Japanese lamprey
Monodelphis domestica	Slc23a1	XP_016278852	Gray short-tailed opossum
Mus musculus	Slc23a1	NP_035527	House mouse
Oryzias latipes	Slc23a1	XP_004076777	Japanese rice fish
Poecilia formosa	Slc23a1	XP_007556555	Amazon molly
Rhincodon typus	Slc23a1	XP_020385431	Whale shark
Xenopus tropicalis	Slc23a1	XP_002932349	Tropical clawed frog
Anolis carolinensis	Slc23a2	XP_003229574	Green anole
Danio rerio	Slc23a2	XP_021327808	Zebrafish
Gallus gallus	Slc23a2	XP_015152797	Chicken
Homo sapiens	Slc23a2	NP_976072	Human
Latimeria chalumnae	Slc23a2	XP_014352307	Coelacanth
Lepisosteus oculatus	Slc23a2	XP_015197789	Spotted gar
Lethenteron japonicum	Slc23a2	JL15308	Japanese lamprey
Leucoraja erinacea	Slc23a2	ctg10612	Little Skate
Monodelphis domestica	Slc23a2	XP_007476346	Gray short-tailed opossum
Mus musculus	Slc23a2	NP_061294	House mouse
Sarcophilus harrisii	Slc23a2	XP_003757995	Tasmanian devil

Species	Gene	Accession	Common name
Scyliorhinus canicula	Slc23a2	ctg15352	Small-spotted catshark
Xenopus tropicalis	Slc23a2	NP_001120161	Tropical clawed frog
Anolis carolinensis	Slc23a3	XP_008108299	Green anole
Danio rerio	Slc23a3	XP_017213243	Zebrafish
Gallus gallus	Slc23a3	XP_015145565	Chicken
Homo sapiens	Slc23a3	NP_001138362	Human
Latimeria chalumnae	Slc23a3	XP_014343771	Coelacanth
Lepisosteus oculatus	Slc23a3	XP_015214926	Spotted gar
Monodelphis domestica	Slc23a3	XP_007501758	Gray short-tailed opossum
Mus musculus	Slc23a3	NP_919314	House mouse
Xenopus tropicalis	Slc23a3	XP_012826889	Tropical clawed frog
Anolis carolinensis	Slc23a4	XP_016849222	Green anole
Callorhinchus milii	Slc23a4	XP_007903282	Elephant shark
Danio rerio	Slc23a4	NP_001013353	Zebrafish
Gallus gallus	Slc23a4	XP_015145041	Chicken
Gekko japonicus	Slc23a4	XP_015276774	Schlegel's Japanese gecko
Latimeria chalumnae	Slc23a4	XP_005991386	Coelacanth
Lepisosteus oculatus	Slc23a4	XP_006633863	Spotted gar
Lethenteron japonicum	Slc23a4	JL7088	Japanese lamprey
Monodelphis domestica	Slc23a4	XP_007504433	Gray short-tailed opossum
Mus musculus	Slc23a4	XP_006506197	House mouse
Oryzias latipes	Slc23a4	XP_004069901	Japanese medaka
Rhincodon typus	Slc23a4	XP_020391140	Whale shark
Sarcophilus harrisii	Slc23a4	XP_012406506	Tasmanian devil
Xenopus tropicalis	Slc23a4	XP_012815547	Tropical clawed frog
Boleophthalmus pectinirostris	Slc23a5	XP_020774843	Great blue-spotted mudskipper
Cynoglossus semilaevis	Slc23a5	XP_016888339	Tongue sole
Cyprinus carpio	Slc23a5	XP_018944039	Common carp
Danio rerio	Slc23a5	XP_009296238	Zebrafish
Fundulus heteroclitus	Slc23a5	XP_021172742	Mummichog
Hippocampus comes	Slc23a5	XP_019750396	Tiger tail seahorse
Ictalurus punctatus	Slc23a5	XP_017328967	Channel catfish
Lates calcarifer	Slc23a5	XP_018548144	Barramundi perch
Lepisosteus oculatus	Slc23a5	XP_015208244	Spotted gar
Maylandia zebra	Slc23a5	XP_014264577	Zebra mbuna
Monopterus albus	Slc23a5	XP_020448997	Swamp eel
Nanorana parkeri	Slc23a5	XP_018412235	-
Neolamprologus brichardi	Slc23a5	XP_006782389	Fairy cichlid
Nothobranchius furzeri	Slc23a5	XP_015819289	Turquoise killifish
Oncorhynchus mykiss	Slc23a5	XP_021432371	Rainbow trout
Oreochromis niloticus	Slc23a5	XP_005454237	Nile tilápia
Oryzias latipes	Slc23a5	XP_004086320	Japanese medaka
Poecilia reticulata	Slc23a5	XP_008408845	Guppy
Pundamilia nyererei	Slc23a5	XP_005731628	-
Pygocentrus nattereri	Slc23a5	XP_017573200	Red-bellied piranha
Salmo salar	Slc23a5	XP_013981136	Atlantic salmon
Scleropages formosus	Slc23a5	XP_018602707	Asian bonytongue

Sinocyclocheilus rhinocerous	Slc23a5	XP_016366936	-
Stegastes partitus	Slc23a5	XP_008289900	Bicolor damselfish
Takifugu rubripes	Slc23a5	XP_003967776	Fugu rubripes
Xenopus tropicalis	Slc23a5	XP_002937113	Tropical clawed frog

Table S2 - Accession numbers of the RNAseq files. PE – portable executable format.

Spotted gar	Library	SRA Run	Read length
Brain	PE	SRR1524250	100
Intestine	PE	SRR1524257	100
Kidney	PE	SRR1524255	100
Liver	PE	SRR1524254	100
Testis	PE	SRR1524260	100
Zebrafish	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR1524238	100
Intestine	PE	SRR1524245	100
Kidney	PE	SRR1524243	100
Liver	PE	SRR1524242	100
Testis	PE	SRR1524249	100
Chicken	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR924542	100
Intestine	PE	SRR3194289	100
Kidney	PE	SRR924553	100
Liver	PE	SRR924555	100
Testis	PE	SRR924547	100
Human	**Library**	**SRA Run**	**Read length**
Brain	PE	ERR315432	100
Intestine	PE	ERR315419	100
Kidney	PE	ERR315468	100
Liver	PE	ERR315451	100
Testis	PE	ERR315456	100
Mouse	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR5171101	100
Intestine	PE	SRR5171080	100
Kidney	PE	SRR5171094	100
Liver	PE	SRR5171078	100
Testis	PE	SRR5171084	100
Western clawed frog	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR579560	75
Intestine	SE	SRR1186999	50
Kidney	PE	SRR579562	75
Liver	PE	SRR579561	75
Testis	SE	SRR943353	100
Anole lizard	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR579556	75
Kidney	PE	SRR579557	75
Liver	PE	SRR391651	100
Testis	PE	SRR5420127	50
Japanese medaka	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR1524271	100
Intestine	PE	SRR1524278	100
Kidney	PE	SRR1524276	100
Liver	PE	SRR1524275	100
Testis	PE	SRR1628839	90

Table S3 - Gene identifier for all genes included in gene expression analysis, collected from NCBI and ENSEMBLE Databases.

Organism	Gene Symbol	Gene ID NCBI	Gene ID Ensemble
Human	Slc23a1	9962	ENSG00000170482
	Slc23a2	9963	ENSG00000089057
	Slc23a3	151295	ENSG00000213901
Mouse	Slc23a1	20522	ENSMUSG00000024354
	Slc23a2	54338	ENSMUSG00000027340
	Slc23a3	22626	ENSMUSG00000026205
	Slc23a4	243753	ENSMUSG00000029847
Lizard	Slc23a1	100556489	ENSACAG00000016325
	Slc23a2	100566000	ENSACAG00000005986
	Slc23a3	100567833	ENSACAG00000003143
	Slc23a4	100563862	ENSACAG00000011027
Chicken	Slc23a1	416188	ENSGALG00000037642
	Slc23a2	419520	ENSGALG00000000195
	Slc23a3	429039	Not Found
	Slc23a4	417937	ENSGALG00000011717
Western clawed frog	Slc23a1	100493636	ENSXETG00000008402
	Slc23a2	100145200	ENSXETG00000016853
	Slc23a3	100486175	ENSXETG00000013878
	Slc23a4	100490216	ENSXETG00000020228
	Slc23a5	100486632	Not Found
Zebrafish	Slc23a1	559435	ENSDARG00000015033
	Slc23a2	100321249	ENSDARG00000017365
	Slc23a3	100333020	ENSDARG00000088891
	Slc23a4	503757	ENSDARG00000052497
	Slc23a5	798958	ENSDARG00000045792
Japanese medaka	Slc23a1	101155711	ENSORLG00000012701
	Slc23a2	101156743	ENSORLG00000002889
	Slc23a3	Not Found	Not Found
	Slc23a4[a]	101173186	ENSORLG00000009915
	Slc23a4b	101157211	ENSORLG00000015991
	Slc23a5	101175057	ENSORLG00000019882
Spotted gar	Slc23a1	102691341	ENSLOCG00000012533
	Slc23a2	102691255	ENSLOCG00000015630
	Slc23a3	102699040	ENSLOCG00000010877
	Slc23a4	102686145	ENSLOCG00000016950
	Slc23a5	2682847	ENSLOCG00000016369

1 **Table S4 - Genome and GTF files retrieved from Ensemble database (Release 89) and Transcriptome files retrieved from NCBI used for this**
2 **study.**

Organisms	Genome / Transcriptome Files	GTF FILE	Download date	Source
Mouse	Mus_musculus.GRCm38.dna.toplevel.fa	Mus_musculus.GRCm38.89.gtf	09/08/2017	Ensemble
Spotted gar	Lepisosteus_oculatus.LepOcu1.dna.toplevel.fa	Lepisosteus_oculatus.LepOcu1.89.gtf	09/08/2017	Ensemble
Anole lizard	Anolis_carolinensis.AnoCar2.0.dna.toplevel.fa	Anolis_carolinensis.AnoCar2.0.89.gtf	09/08/2017	Ensemble
Zebrafish	Danio_rerio.GRCz10.dna.toplevel.fa	Danio_rerio.GRCz10.89.gtf	09/08/2017	Ensemble
Human	Homo_sapiens.GRCh38.dna.toplevel.fa	Homo_sapiens.GRCh38.89.gtf	09/08/2017	Ensemble
Japanese medaka	Oryzias_latipes.MEDAKA1.dna.toplevel.fa	Oryzias_latipes.MEDAKA1.89.gtf	09/08/2017	Ensemble
Western clawed frog	GCF_000004195.3_Xenopus_tropicalis_v9.1_rna.fna	Xenopus_tropicalis_Gene_Transcript_Map.txt*	09/08/2017	Ncbi
Chicken	GCF_000002315.4_Gallus_gallus-5.0_rna.fna	Gallus_gallus_Gene_Transcript_Map.txt*	09/08/2017	Ncbi

6 **Table S5 - Relative gene expression levels of Slc23 family genes for eight species. The values are presented in log2 (TPM +1) ratios.**

Organism	Tissue	Slc23a1	Slc23a2	Slc23a3	Slc23a4a	Slc23a4b	Slc23a5
Human	BRAIN	0,00	5,82	0,00	-	-	-
	INTESTINE	5,85	2,64	4,61	-	-	-
	KIDNEY	3,71	3,70	4,28	-	-	-
	LIVER	4,03	3,58	2,36	-	-	-
	TESTIS	0,00	4,59	0,00	-	-	-
Mouse	BRAIN	0,00	4,91	0,00	0,00	-	-
	INTESTINE	3,73	5,24	1,02	5,89	-	-
	KIDNEY	6,91	2,82	4,08	0,00	-	-
	LIVER	5,06	2,85	0,00	0,00	-	-
	TESTIS	0,00	1,63	0,00	0,00	-	-
Lizard	BRAIN	0,91	2,82	0,72	2,49	-	-
	KIDNEY	6,76	1,39	6,10	1,19	-	-
	LIVER	3,02	0,00	0,00	0,65	-	-
	TESTIS	0,00	0,00	0,00	0,00	-	-
Chicken	BRAIN	0,00	6,27	0,96	1,82	-	-
	INTESTINE	6,44	3,25	2,96	6,09	-	-
	KIDNEY	7,78	1,49	4,17	1,60	-	-
	LIVER	5,20	1,98	1,76	0,82	-	-
	TESTIS	2,35	2,08	2,03	1,85	-	-
Western clawed frog	BRAIN	0,00	6,45	0,00	2,54	-	0,00
	INSTESTINE	0,00	2,23	0,00	5,65	-	0,00
	KIDNEY	7,63	4,08	6,53	5,00	-	5,40
	LIVER	0,00	3,41	0,00	2,14	-	0,00
	TESTIS	2,69	4,01	0,00	1,49	-	1,64
Zebrafish	BRAIN	0,85	3,73	0,71	0,00	-	1,01
	INTESTINE	2,69	1,21	0,00	4,00	-	0,00
	KIDNEY	5,35	2,27	3,42	4,78	-	0,93
	LIVER	0,00	1,26	0,00	0,00	-	0,00
	TESTIS	0,93	2,74	0,00	0,63	-	1,97
Japanese medaka	BRAIN	0,00	4,70	-	0,00	0,00	0,84
	INTESTINE	2,76	2,91	-	4,95	0,00	0,00
	KIDNEY	3,89	3,02	-	4,43	0,00	0,00
	LIVER	0,00	4,06	-	0,00	0,00	0,00
	TESTIS	0,00	3,20	-	0,60	0,00	0,00
Spotted gar	BRAIN	0,58	4,94	0,00	0,00	-	0,00
	INTESTINE	5,67	0,96	0,00	0,00	-	0,00
	KIDNEY	3,18	0,90	1,78	0,86	-	0,00
	LIVER	0,74	0,78	0,00	0,00	-	0,00
	TESTIS	1,92	7,21	0,00	0,00	-	0,00

13 **Table S6 - Renal uric acid absorption and secretion: the urate transportome.** n.f. – not found.

	Transporter	H. sapiens	M. musculus	G. gallus	A. carolinensis	X. tropicalis	D. rerio	L. oculatus
Basolateral Absorption	SLC2A9/GLUT9	✓	✓	✓	✓	✓	✓	✓
Apical Absorption	SLC22A11/OAT4	✓	n.f.	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC22A12/URAT1	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC22A13/OAT10	✓	✓	✓	✓	✓	✓	✓
Basolateral Secretion	SLC22A6/OAT1	✓	✓	✓	✓	✓	n.f.	n.f.
	SLC22A8/OAT3	✓	✓				n.f.	n.f.
Apical Secretion	SLC17A1/NPT1	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC17A3/NPT4	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.

17