Identification of a novel Nucleobase-Ascorbate Transporter family member in fish and amphibians

Diogo Oliveira*,1, André M. Machado*,1, Tiago Cardoso1, Mónica Lopes-Marques1, L. Filipe C. Castro1,2● and Raquel Ruivo1●

1CIIMAR – Interdisciplinary Centre of Marine and Environmental Research, U. Porto – University of Porto, Porto, Portugal

2Department of Biology, Faculty of Sciences, U. Porto - University of Porto, Portugal

*Equal contribution

●Corresponding authors at: CIIMAR, Terminal de Cruzeiros do Porto de Leixões, Av. General Norton de Matos s/n, 4450-208 Matosinhos, Portugal. Tel.: +351 223 401 831

Running title: Novel uric acid transporter in fish and amphibians

Abstract: Nucleobase-Ascorbate Transporter (NAT) family includes ascorbic acid, nucleobases and uric acid transporters: with a broad evolutionary distribution. In vertebrates, four members have been previously recognized, the ascorbate transporters Slc23a1 and Slc3a2, the nucleobase transporter Slc23a4 and an orphan transporter Slc23a3. Using phylogenetic and synteny analysis, we identify a fifth member of the vertebrate *slc23* complement (*slc23a5*), present in neopterygians (gars and teleosts) and amphibians, and clarify the evolutionary relationships between the novel gene and known *slc23* genes. Further comparative analysis puts forward uric acid as the preferred substrate for Slc23a5. Gene expression quantification, using available transcriptomic data, suggests kidney and testis as major expression sites in *Xenopus tropicalis* (western clawed frog) and *Danio rerio* (zebrafish). Additional expression in brain was detected in *D. rerio*, while in the Neoteleostei *Oryzias latipes* (medaka) *slc23a5* expression is restricted to brain. The biological relevance of the retention of an extra transporter in fish and amphibians is discussed.

Keywords: Nucleobase-Ascorbate Transporters (NAT); Slc23; Slc23a5; Uric acid; Antioxidant

1. Introduction

Ascorbic acid (or its salt form ascorbate) is an essential enzyme cofactor, participating in collagen and norepinephrine synthesis, as well as a potent antioxidant and free radical scavenger [1,2]. Despite its importance, anthropoid primates, teleost fish, Guinea pigs, some bats and Passeriformes bird species, lack a functional L-gulono-Y-lactone oxidase (GLO) gene and, thus, the ability to catalyse the last step of ascorbic acid synthesis [3,4]. The scattered loss of the GLO gene was suggested to evolve neutrally, with ascorbic acid requirements being met by dietary intake [3]. Unlike ascorbate, uric acid is largely regarded as end-product of purine metabolism and metabolic waste [5]. Yet, human-biased research has suggested alternative physiological functions for uric acid or urate (uric acid salts), such as antioxidant and free radical scavenger, similarly to ascorbic acid, counterbalancing disease-related oxidative stress [6-8].

Ascorbic and uric acid serve as substrates to active transporters that belong to the Nucleobase-Ascorbate Transporter (NAT) family; NAT family members, which include Solute carrier family 23 (Slc23) transporters, display a very broad evolutionary distribution, from bacteria to metazoans [1]. Sodium-dependent Slc23 transporters were suggested to maintain vertebrate ascorbic acid levels: with Slc23a1 responsible for renal and intestine ascorbic acid absorption across brush border epithelia and Slc23a2, which exhibits a widespread expression, responsible for ascorbic acid supply to target tissues and cells, including neurons [1,9,10]. In vertebrates, two additional transporters were identified, the orphan transporter Slc23a3 and the nucleobase transporter Slc23a4, described in rats but found pseudogenized in humans [1,11]. Rat Slc23a4 is restricted to the small intestine and was shown to predominantly mediate sodium-dependent uracil transport [11]. Slc23a3, on the other hand, is expressed in proximal renal tubules but is unresponsive to both ascorbic acid and nucleobases [1].

NAT family members display a highly conserved signature motif: [Q/E/P]-N-x-G-x-x-x-x-T-[R/K/G] [1,11-13]. Functional characterization of distinct member of the NAT family highlighted the importance of the first amino acid position of the motif in the definition of

substrate specificity [1,11-13]. Thus, these transporters can be divided into three functional

groups. Slc23a1, Slc23a2 and Slc23a3 belong to the ascorbate group and harbour a conserved

proline (P) in the first amino acid position of the signature motif [1]. A glutamate (E) residue

defines the uracil group, including bacterial anion symporters and Slc23a4, while a glutamine (Q)

is present in the bacterial uric acid and/or xanthine anion symporters [1,11-13]. Topology

predictions, using proteins from the distinct groups, suggest a cytosolic location of the motif [1,

11, 12]. Here we revise the portfolio of NAT family members in vertebrates.

2. Materials and Methods

2. 1Sequence mining

Slc23 protein sequences were obtained from the GenBank database by BLASTp using

human Slc23a1 (NP_005838), Slc23a2 (NP_976072) and Slc23a3 (NP_001138362) sequences as

query. The sequences from *Lethenteron japonica* (Japanese lamprey) were obtained from the

Japanese Lamprey Genome Project (http://jlampreygenome.imcb.a-star.edu.sg/). The sequences

from *Leucoraja erinacea* (little skate) and *Scyliorhinus canicula* (small-spotted catshark) were

obtained as nucleotide sequences from SkateBase (http://skatebase.org) and translated. Accession

numbers for all the retrieved sequences are listed in Supplementary Table 1 (Table S1).

2. 2 Synteny

Slc23 genes were mapped onto the respective species genomes, using the latest genome

assemblies available in GenBank database (https://www.ncbi.nlm.nih.gov/). Genes flanking

Slc23a1, *Slc23a2*, *Slc23a3*, *Slc23a4* and the novel *Slc23a*-like gene were identified and mapped using

the human loci as reference. If the target gene was not found, the syntenic genomic region was

retrieved using neighboring genes as reference.

2. 3Phylogenetic Analysis

The retrieved amino acid sequences were aligned using the MAFFT software web service

(http://mafft.cbrc.jp/alignment/software/) in default automatic settings, with L-INS-I refinement

method [14]. The alignment was stripped from columns with at least 20% of gaps resulting in an alignment with 75 sequences and 526 positions. The output alignment was used to construct a phylogenetic tree using PhyML 3.0 with Smart Model Selection web service (http://www.atgc-montpellier.fr/phyml-sms/) in default settings, which selected the LG +G+I+F model [15]. Bayesian-like transformation of aLRT was selected to assess branch support. An additional phylogenetic tree, with automatic bootstrapping (552 bootstraps) and using the default hill-climbing algorithm, was constructed with RAxML, available in CIPRES Science Gateway.

2.4Membrane topology prediction

Slc23a5 membrane topology predictions were performed for selected species, using distinct prediction web servers: TMHMM Server v. 2.0 (http://www.cbs.dtu.dk/services/TMHMM/) and TMPred (https://embnet.vital-it.ch/software/TMPRED_form.html). Human Slc23a1 was used as control.

2.5Filtering and quality control of Sequence Reading Archive (SRA) datasets

For RNA-seq analysis, we included 39 SRA datasets from brain, intestine, kidney, liver, and testis tissues of *Homo sapiens* (human), *Mus musculus* (mouse), *Gallus gallus* (chicken), *Anolis carolinensis* (American green anole), *Xenopus tropicalis* (western clawed frog), *Danio rerio* (zebrafish), *Lepisosteus oculatus* (spotted gar) and *Oryzias latipes* (Japanese medaka), which were obtained from the National Center for Biotechnology (NCBI) Sequence Read Archive (*SRA*) (https://www.ncbi.nlm.nih.gov/sra/) (Table S2). The read quality assessment of each SRA dataset was performed using FastQC software (https://www.bioinformatics. Babraham.ac.uk/projects/fastqc/). Trimmomatic 0.36 [17] was used to trim and drop reads with quality scores below 5, at leading and trailing ends, with an average quality score below 20 in a 4 bp sliding window and with less than 36 bases length.

2.6Relative gene expression levels

To assess the relative gene expression of *slc23* genes from the selected species, reference

sequences and the corresponding annotations were collected from NCBI and Ensembl [18]

databases. For human, mouse, lizard, zebrafish and spotted gar, the GTF and DNA files (genome)

were retrieved from Ensembl (Release 89) [18]. For chicken and western clawed frog, transcript

sequences (coding and non-coding) and mapping (gene/transcript) from NCBI Refseq database

were   used   after   filtering   [19]   (for   each   species   the   rna.fasta   file   from

ftp://ftp.ncbi.nih.gov/genomes/refseq/ was used – See Table S3 and S4 for detail). The SRA reads

of each specimen were aligned to the reference using Bowtie2 (default parameters)[20], and gene

expression was quantified as transcript per million (TPM) using RSEM v.1.2.31 (RNA-seq by

Expectation Maximization) software package [21]. Isoform expression levels for each gene were

summed to derive the TPM values. For genes with low relative expression values, TPM values

were log2-transformed after adding a value of one; TPM values < 0.5 were considered unreliable

and substituted with zero.

2. 7PCR-based validation of Slc23a5 expression

Total RNA extraction from *X. tropicalis* and *D. rerio* tissues was performed using the Illustra

RNAspin Mini RNA Isolation Kit animal tissues protocol with on-column DNAseI digestion (GE

Healthcare). RNA quality was visually assessed by electrophoresis and sample concentration was

determined using the Take3 micro-volume plate system (BioTek). First-strand cDNA was

synthesized from 250ng of total RNA using the iScriptTM cDNA Synthesis Kit (Bio-Rad),

according to the manufacturer's instructions. Gene expression of *Slc23a5* was analyzed by PCR

using the intro flanking expression primer pairs 5'-AGACGTTGAAGCCGTTACTGA-3' and 5'-

TGCTATTATGCCTCCAAAGGCT-3' for *X. tropicalis* and 5'-GGTGGAATGTTCCTCGTCAT-3'

and 5'-GATGAACATGTGGGTGGTGA-3' for *D. rerio*. PCR reactions were carried out on a tissue

panel containing heart, skin, stomach, intestine, liver, kidney, ovary and testis tissues for *X.*

tropicalis and brain, heart, skin, gut, liver, kidney, ovary and testis tissues for *D. rerio*. The PCR

was performed with Phusion® Flash high fidelity Master Mix (Thermofisher), 1µM of each

primer and 25ng/μl of template cDNA, with the following parameters: initial denaturation at

98°C for 1s, annealing at 58°C for *D. rerio* or 66ºC for *X. tropicalis* primers and elongation at 72°C

for 4s, with 30 cycles.

3. Results

3.1Identification of a novel member of the NAT family in neopterygians
and amphibians

Through database mining and phylogenetic analysis, we detected the presence of a novel

gene, *slc23a5*, found in neopterygians (Holostei and Teleostei) and amphibians (Figure 1 and

Figure S1). No *slc23a5* sequence was retrieved from chondrichthyans, chondrosteans (basal

actinopterygians) nor from the sarcopterygian *Latimeria chalumnae* (coelacanth). Subsequent loss

of *slc23a5* in chondrichthyans was further supported by the conservation of the genomic locus in

Callorhynchus milii (Australian ghostshark) (Figure S2). Regarding coelacanth, two scaffolds,

denoting conservation of the flanking genes, were found; while, no information was retrieved

regarding the chondrostean loci (Figure S2). The remaining genes of the *slc23* family were

retrieved and genomic loci were found generally conserved across vertebrate species, with

exception of the *slc23a4* loci in fish (Figures S3-S6). The branching pattern of our phylogenetic

analysis suggests that after the expansion of the *slc23a* gene family *slc23a5* was subsequently lost

in sharks, basal actinopterygians, coelacanth and amniotes (Figure 1 and Figure S2). In agreement,

we obtained three phylogenetic clusters possibly corresponding to three ancestral lineages

(Figure 1 and Figure S1): one composed of typical *slc23a1* and *slc23a2* genes and a second group

including the previously reported *slc23a4* and the novel *slc23a5* genes, both outgrouped by

invertebrate sequences. A third cluster, which outgroups the remaining *slc23* genes, included the

highly divergent *slc23a3* genes, mirrored by the long branch lengths (Figure 1).

Figure 1 – Maximum likelihood phylogenetic tree describing relationships among Nucleobase-Ascorbate Transporter (NAT) family members. Node values represent branch

 support using the aBayes algorithm. Accession numbers for all sequences are provided in

 Supplementary Materials.

 3.2Slc23a5 signature motifs

    Further comparative protein sequence analysis highlighted the presence of a NAT family

 signature motif with a conserved Q in the substrate determining position. According to previous

 functional characterization of NAT family members, this residue is found in uric acid and/or

 xanthine transporters, thus placing Slc23a5 in the uric acid and/or xanthine group (Figure 2)[1,11-

 13]. Besides the typical signature motif, a carboxyl terminus tripeptide motif was identified in

 some SLC23 family members: a S/T-X-Ø tripeptide. S/T-X-Ø motifs were shown to interact PDZ

 domains of scaffolding proteins belonging to the Sodium/Hydrogen Exchanger Regulatory

 Factor, NHERF, family [22]. S/T-X-Ø are found highly conserved across species in Slc23a1 and

 Slc23a4, in agreement with their renal and intestinal expression and role in ascorbic acid and

 nucleobase absorption (Figure S7). Slc23a5, on the other hand, exhibits a highly conserved

 tripeptide motif in Otocephala (Cypriniformes, Characiformes and Siluriformes) but not

 Osteoglossomorpha (Figure 3A). The Euteleostei Esociformes and Salmoniformes exhibit an

 inverted motif while in the Neoteleostei no conservation was found. Finally, topology predictions

 yielded similar transmembrane profiles for Slc23a5 sequences from *D. rerio*, *O. latipes*, *Salmo salar*

 (Atlantic salmon) and *X. tropicalis*, when compared to human Slc23a1 (Figure S8). The NAT motif

 was predicted cytosolic in all analyzed sequences. This suggests a conserved cellular import

 topology, as observed in other family members.

Figure 2 – Alignment of the signature motif of the Nucleobase-Ascorbate Transporter (NAT)

family. NAT family members display a highly conserved signature motif defining substrate

specificity: [Q/E/P]-N-x-G-x-x-x-x-T-[R/K/G]. Asterisks indicate conserved positions. The first

amino acid position, with residues highlighted in red within a black box, defines substrate

preference: Q – Uric acid/xanthine group; P – Ascorbate group and E – Uracil group. The mirror

Q in the Slc23a3 group is also highlighted by a black box.

Figure 3 – The S/T-X-Ø tripeptide provides insight into the cellular function of Slc23a5

homologues from fish and amphibians. (A) Alignment of the PDZ-binding tripeptide in Slc23a5

homologues from fish and amphibians. The S/T-X-Ø tripeptide is found in Cypriniformes,

Characiformes and Siluriformes while the Euteleostei Esociformes and Salmoniformes exhibit an

inverted motif. Predicted cellular localization is indicated as well as morphological and

physiological adaptations among the different fish groups. (B) Schematic representation of

tripeptide-dependent uric acid flows in polarized epithelia and non-polarized cells. In polarized

epithelia (i.e. kidney) the presence of the tripeptide motif suggests interaction with the scaffolding

protein PDZK1 which localizes apically, in the absorptive side.

3.3Slc23a5 expression patterns

To gain further insight into the functional relevance of this additional gene, we collected

RNA-seq data and determined relative expression levels of *Slc23* genes. Gene expression analysis

suggested that, in *D. rerio* and *X. tropicalis*, slc23a5 is mostly expressed in kidney and testis, further

confirmed by PCR (Figure 4 and Table S5). Expression in brain was also detected in *D. rerio* and

was found to be the exclusive expression site in the Neoteleostei *O. latipes*. The holostean *L.*

oculatus, on the other hand, yielded no significant expression in the tested tissues.

Figure 4 – Expression patterns of *Slc23* isoforms across vertebrates. (A) RNAseq-based

quantification. Relative gene expression levels are provided as $\log_2$-transformed transcript per

million (TPM) values, from 0 to 8 TPM's, the NF and ND means gene not found and no SRA data

available to a specific tissue. (B) PCR-based amplification of *Slc23a5* transcripts in *D. rerio* and *X.*

tropicalis tissues.

4. Discussion

Our findings indicate a restricted distribution of *slc23a5* to taxa that share an aquatic nature

of their habitat. In fact, *slc23a5* was found to be expressed in kidney and testis of both *X. tropicalis*

and *D. rerio*. In the latter, as well as in the Neoteleostei *O. latipes*, brain expression was also

detected. Comparative analysis using previously characterized NAT family transporters placed

this novel transporter in the uric acid and/or xanthine transporter group. In fact, similarly to uric

acid and/or xanthine transporters, Slc23a5 proteins harbor a Q in the first position of the NAT

family signature motif ([Q/E/P]-N-x-G-x-x-x-x-T-[R/K/G]). Yet, further functional studies are

required to fully clarify the molecular function of Slc23a5 proteins, notably its uric acid and/or

xanthine binding abilities.

Given the nature of NAT family transporter ligands, including ascorbic and uric acids,

xanthine or nucleobases, renal Slc23a5 could participate in the homeostatic maintenance of

excretion products, in osmotic regulation and/or in antioxidant processes. In fact, kidneys play a

crucial role in the maintenance of both ascorbic and uric acid levels through the excretion and

reabsorptions processes [6,22]. In kidney absorptive epithelium, transporters located on the

filtrate-epithelial interface (apical side) interact with the scaffolding protein PDZ-Containing

Kidney Protein 1 (PDZK1) [22]. PDZK1 belongs to the NHERF family of proteins which are

mostly expressed in the apical side of absorptive epithelia of kidney, intestine and liver, and

establish protein-protein interaction networks with receptors, transporters and channels [23,24].

Besides its scaffolding role, NHERF family members were also shown to modulate the activity of

protein assemblages [24-26]. Within Slc23 family members, S/T-X-Ø motifs are found in Slc23a1

and Slc23a4 proteins, which are present in renal and intestinal polarized epithelia and participate

in ascorbic acid and nucleobase absorption, and absent in Slc23a2. Curiously, a highly conserved

PDZK1 interacting motif (S/T-X-Ø) was also detected in Slc23a5 from Cypriniformes,

Characiformes and Siluriformes, but not in other fish groups or amphibians. No relationship

between salt and freshwater environments was found. Thus, the presence/absence of this peptide

signal might reflect distinct flows in the kidney: an apically expressed Slc23a5 likely participates

in absorptive flows, while a basolateral expression would be related to secretion (Figure 3B).

In the human proximal renal tubules, several transporters, localized to specific cellular

regions (apical or basolateral), seem to cooperate for efficient urate handling: named the urate

transportome [6,22]. Thus, if the predicted uric acid transport activity is confirmed, *X. tropicalis*

and *D. rerio* Slc23a5 could illustrate the two opposing mechanisms: while *D. rerio* captures, *X.*

tropicalis secretes uric acid into renal filtrate. In agreement, *X. tropicalis*, *D. rerio*, as well as *L.*

oculatus, appear to lack the full complement of renal urate transporters (Table S6). A similar

scenario is observed for birds and reptiles; however, their excretory system represents a particular

adaptation for uric acid excretion [27,28]. Besides simple excretion, and given that high

concentrations of uric acid were detected, and suggested to act as the main non-enzymatic

antioxidant mechanism, in seminal fluid of several fish species (including Cypriniformes,

Esociformes, Salmoniformes and the Neoteleostei Gadiformes and Perciformes [29-32]), this

putative uric acid transport activity could be maintained for reproductive purposes, in agreement

with the observed testicular expression. In fact, in external fertilizers, such as most fish and

amphibians, reactive oxygen species were suggested to decrease motility and thus fertilisation

success [29]. Yet, regarding amphibians, no information on seminal fluid composition was

retrieved [33]. Nonetheless, in vitro assays suggested that other antioxidant species, such as

ascorbic acid or vitamin E, were unsuitable at offsetting oxidative stress and improving sperm

motility in *Litoria booroolongensis* (Booroolong frog) and suggested uric acid as a pertinent

candidate [33]. In this context, the absence of *slc23a5* sequences in Chondrichthyes could be due

to their internal fertilization strategies derived from ovoviviparous or viviparous reproductive

modes [34]. Accordingly, viviparity is also observed in the sarcopterygian *L. chalumnae* [34].

Besides reproductive categories, other morphological and metabolic traits could justify the

seemingly scattered retention of *slc23a5* among fish in the context of the proposed uric acid

transport activity; for instance, specific anatomical features such as the presence of a urogenital

connection: described in Elasmobranchii and in most non-teleost Osteichthyes, yet absent in

Cyclostomata and Teleostei [35]. By connecting testicular and renal ducts, antioxidant

requirements could be met by renal secretion into urine. In agreement, in the Chondrostei

Acipenser ruthenus, full sperm maturation and motility was shown to require dilution by urine

[36]. An urogenital connection is also present in amphibians [35]. Additionally, other antioxidant

mechanisms could exist. Besides uric acid, ascorbic acid is also present in seminal fluid of

Oncorhynchus mykiss (rainbow trout) [37]. Seminal ascorbate was reported to be affected by

seasonal variations and semen quality was shown to improve upon dietary ascorbic acid

supplementation [37,38], contrasting with the amphibian *L. booroolongensis* [33]. In the case of non-

teleost fish, such as Chondrostei (*A. ruthenus*) and Holostei (*L. oculatus*), a functional GLO gene is

retained, providing endogenously synthesised ascorbate [39]. Curiously, no gonadal expression

was detected in the Neoteleostei *O. latipes*. Neoteleosts exhibit varied reproduction modes

ranging from external fertilizers to internal fertilizers with viviparity [34,40-42]. Also, neoteleosts

were shown to exhibit a distinctive gonadal morphology, the lobular testis, whereas the

remaining bony fish exhibit tubular testis [43]. Although still obscure, the absence of testicular

slc23a5 expression in the externally fertilizer *O. latipes* could be due to specific morphological

adaptations of neoteleosts.

A similar role could be postulated for neuronal uric acid [44]. Active swimming behaviour

requires a high metabolic rate, mirrored by an increased oxygen consumption, subsequently

leading to reactive oxygen species production [45]. In agreement, fish antioxidant levels were

shown to be correlated with their swimming activity [45]. Neuronal cell populations are particularly sensitive to oxidative stress, thus, uric acid could participate in the neutralization of oxidative species to by-pass oxidative stress, notably in GLO-deficient fish. Yet, information on neuronal uric acid levels in fish is currently absent.

5. Conclusion

Taken together, these observations put forward the retention of an additional *Slc23* gene, in neopterygians and amphibians, with restricted expression patterns and presumed uric acid transport function. Yet, extensive comparative functional analysis should be conducted to clarify the physiological relevance of this novel receptor in aquatic organisms.

Supplementary material: Supplementary tables and figures including synteny, phylogeny and topology analysis, as well as reference tables.

Acknowledgments : This work was supported by Coral—Sustainable Ocean Exploitation (Norte-01-0145-FEDER-000036), a project by the North Portugal Regional Operational Program (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement, through the European Regional Development Fund (ERDF) and by the European Regional Development Fund (ERDF) through the COMPETE - Operational Competitiveness Programme and POPH - Operational Human Potential Programme.

References

1. Burzle, M.; Suzuki, Y.; Ackermann, D.; Miyazaki, H.; Maeda, N.; Clemencon, B.; Burrier, R.; Hediger, M.A. The sodium-dependent ascorbic acid transporter family SLC23. *Mol. Aspects Med.* **2013**, *34*, 436-454, doi:10.1016/j.mam.2012.12.002.
2. Linster, C.L.; Van Schaftingen, E. Vitamin C. *FEBS J.* **2007**, *274*, 1-22, doi:10.1111/j.1742-4658.2006.05607.x.
3. Drouin, G.; Godin, J.R.; Page, B. The genetics of vitamin C loss in vertebrates. *Curr. Genomics* **2011**, *12*, 371-378, doi:10.2174/138920211796429736.
4. Dutta Gupta, S.; Choudhury, P.K.; Chatterjee, I.B. Synthesis of l-ascorbic acid from d-glucurono-1,4-lactone conjugates by different species of animals. *Int. J. Biochem.* **1973**, *4*, 309-314, doi:http://dx.doi.org/10.1016/0020-711X(73)90053-0.

5. Wright, P.A. Nitrogen excretion: three end products, many physiological roles. *J. Exp. Biol.* **1995**, *198*, 273-281.

6. Bobulescu, I.A.; Moe, O.W. Renal transport of uric acid: evolving concepts and uncertainties. *Adv. Chronic Kidney Dis.* **2012**, *19*, 358-371, doi:10.1053/j.ackd.2012.07.009.

7. Alvarez-Lario, B.; Macarrón-Vicente, J. Evolution of Uric Acid Metabolism in Humans. In *eLS*, John Wiley & Sons, Ltd: 2001; 10.1002/9780470015902.a0024618.

8. Oda, M.; Satta, Y.; Takenaka, O.; Takahata, N. Loss of urate oxidase activity in hominoids and its evolutionary implications. *Mol. Biol. Evol.* **2002**, *19*, 640-653.

9. Corti, A.; Casini, A.F.; Pompella, A. Cellular pathways for transport and efflux of ascorbate and dehydroascorbate. *Arch. Biochem. Biophys.* **2010**, *500*, 107-115, doi:10.1016/j.abb.2010.05.014.

10. Savini, I.; Rossi, A.; Pierro, C.; Avigliano, L.; Catani, M.V. SVCT1 and SVCT2: key proteins for vitamin C uptake. *Amino Acids* **2008**, *34*, 347-355, doi:10.1007/s00726-007-0555-7.

11. Yamamoto, S.; Inoue, K.; Murata, T.; Kamigaso, S.; Yasujima, T.; Maeda, J.Y.; Yoshida, Y.; Ohta, K.Y.; Yuasa, H. Identification and functional characterization of the first nucleobase transporter in mammals: implication in the species difference in the intestinal absorption mechanism of nucleobases and their analogs between higher primates and other mammals. *J. Biol. Chem.* **2010**, *285*, 6522-6531, doi:10.1074/jbc.M109.032961.

12. Koukaki, M.; Vlanti, A.; Goudela, S.; Pantazopoulou, A.; Gioule, H.; Tournaviti, S.; Diallinas, G. The nucleobase-ascorbate transporter (NAT) signature motif in UapA defines the function of the purine translocation pathway. *J. Mol. Biol.* **2005**, *350*, 499-513, doi:10.1016/j.jmb.2005.04.076.

13. Papakostas, K.; Frillingos, S. Substrate selectivity of YgfU, a uric acid transporter from Escherichia coli. *J. Biol. Chem.* **2012**, *287*, 15684-15695, doi:10.1074/jbc.M112.355818.

14. Katoh, K.; Standley, D.M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **2013**, *30*, 772-780, doi:10.1093/molbev/mst010.

15. Guindon, S.; Dufayard, J.F.; Lefort, V.; Anisimova, M.; Hordijk, W.; Gascuel, O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **2010**, *59*, 307-321, doi:10.1093/sysbio/syq010.

16. Dobson, L.; Remenyi, I.; Tusnady, G.E. CCTOP: a Consensus Constrained TOPology prediction web server. *Nucleic Acids Res.* **2015**, *43*, W408-412, doi:10.1093/nar/gkv451.

17. Bolger, A.M.; Lohse, M.; Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **2014**, *30*, 2114-2120, doi:10.1093/bioinformatics/btu170.

18. Yates, A.; Akanni, W.; Amode, M.R.; Barrell, D.; Billis, K.; Carvalho-Silva, D.; Cummins, C.; Clapham, P.; Fitzgerald, S.; Gil, L., et al. Ensembl 2016. *Nucleic Acids Res.* **2016**, *44*, D710-716, doi:10.1093/nar/gkv1157.

19. O'Leary, N.A.; Wright, M.W.; Brister, J.R.; Ciufo, S.; Haddad, D.; McVeigh, R.; Rajput, B.; Robbertse, B.; Smith-White, B.; Ako-Adjei, D., et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **2016**, *44*, D733-745, doi:10.1093/nar/gkv1189.

20. Langmead, B.; Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **2012**, *9*, 357-359, doi:10.1038/nmeth.1923.

21. Li, B.; Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **2011**, *12*, 323, doi:10.1186/1471-2105-12-323.

22. Anzai, N.; Jutabha, P.; Amonpatumrat-Takahashi, S.; Sakurai, H. Recent advances in renal urate transport: characterization of candidate transporters indicated by genome-wide

association studies. *Clin. Exp. Nephrol.* **2012**, *16*, 89-95, doi:10.1007/s10157-011-0532-
z.

23. Dev, K.K. Making protein interactions druggable: targeting PDZ domains. *Nat. Rev. Drug*
Discov. **2004**, *3*, 1047-1056.

24. Clapéron, A.; Mergey, M.; Fouassier, L. Roles of the scaffolding proteins NHERF in liver
biology. *Clin. Res. Hepatol. Gastroenterol.* **2011**, *35*, 176-181,
doi:http://dx.doi.org/10.1016/j.clinre.2010.11.009.

25. Raghuram, V.; Mak, D.-O.D.; Foskett, J.K. Regulation of cystic fibrosis transmembrane
conductance regulator single-channel gating by bivalent PDZ-domain-mediated
interaction. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98*, 1300-1305.

26. Wang, S.; Yue, H.; Derin, R.B.; Guggino, W.B.; Li, M. Accessory Protein Facilitated CFTR-
CFTR Interaction, a Molecular Mechanism to Potentiate the Chloride Channel Activity.
Cell **2000**, *103*, 169-179, doi:10.1016/S0092-8674(00)00096-9.

27. Braun, E.J. Comparative renal function in reptiles, birds, and mammals. *Semin. Avian*
Exotic. Pet. Med. **1998**, *7*, 62-71, doi:http://dx.doi.org/10.1016/S1055-937X(98)80044-
3.

28. Singer, M.A. Do mammals, birds, reptiles and fish have similar nitrogen conserving
systems? *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **2003**, *134*, 543-558.

29. Lahnsteiner, F.; Mansour, N. A comparative study on antioxidant systems in semen of
species of the Percidae, Salmonidae, Cyprinidae, and Lotidae for improving semen
storage techniques. *Aquaculture* **2010**, *307*, 130-140,
doi:http://dx.doi.org/10.1016/j.aquaculture.2010.07.011.

30. Ciereszko, A.; Dabrowski, K.; Kucharczyk, D.; Dobosz, S.; Goryczko, K.; Glogowski, J. The
presence of uric acid, an antioxidantive substance, in fish seminal plasma. *Fish Physiol.*
Biochem. **1999**, *21*, 313-315, doi:10.1023/A:1007886121663.

31. Dzyuba, V.; Dzyuba, B.; Cosson, J.; Boryshpolets, S.; Yamaner, G.; Kholodniy, V.; Rodina,
M. The antioxidant system of sterlet seminal fluid in testes and Wolffian ducts. *Fish*
Physiol. Biochem. **2014**, *40*, 1731-1739, doi:10.1007/s10695-014-9963-2.

32. Lahnsteiner, F.; Mansour, N.; Plaetzer, K. Antioxidant systems of brown trout (Salmo
trutta f. fario) semen. *Anim. Reprod. Sci.* **2010**, *119*, 314-321,
doi:10.1016/j.anireprosci.2010.01.010.

33. Keogh, L.M.; Byrne, P.G.; Silla, A.J. The effect of antioxidants on sperm motility activation
in the Booroolong frog. *Anim. Reprod. Sci.* **2017**, *183*, 126-131,
doi:http://dx.doi.org/10.1016/j.anireprosci.2017.05.008.

34. Goodwin, N.B.; Dulvy, N.K.; Reynolds, J.D. Life-history correlates of the evolution of live
bearing in fishes. *Philos. Trans. R. Soc. Lond. B. Biol. Sci* **2002**, *357*, 259-267,
doi:10.1098/rstb.2001.0958.

35. Blün, V. Comparative Anatomy of the Urogenital System. In *Vertebrate Reproduction, a*
Textbook, Springer-Verlag: Berlin Heidelberg New York Tokyo, 1986; 10.1007/978-3-
642-71074-2pp. 43-74.

36. Dzyuba, B.; Cosson, J.; Boryshpolets, S.; Bondarenko, O.; Dzyuba, V.; Prokopchuk, G.;
Gazo, I.; Rodina, M.; Linhart, O. In vitro sperm maturation in sterlet, Acipenser ruthenus.
Reprod. Biol. **2014**, *14*, 160-163, doi:10.1016/j.repbio.2014.01.003.

37. Ciereszko, A.; Dabrowski, K. Sperm quality and ascorbic acid concentration in rainbow
trout semen are affected by dietary vitamin C: an across-season study. *Biol. Reprod.*
1995, *52*, 982-988.

38. Ciereszko, A.; Dabrowski, K. Effect of ascorbic acid supplement in vitro onrainbow trout
sperm viability. *Aquacult. Int.* **2000**, *8*, 1-8, doi:10.1023/A:1009253330451.

39. Moreau, R.; Dabrowski, K. Biosynthesis of ascorbic acid by extant actinopterygians. *J.*
Fish Biol. **2000**, *57*, 733-745, doi:10.1111/j.1095-8649.2000.tb00271.x.

40.   Forsman, A.; Tibblin, P.; Berggren, H.; Nordahl, O.; Koch‑Schmidt, P.; Larsson, P. Pike
Esox lucius as an emerging model organism for studies in ecology and evolutionary
biology: a review. *J. Fish Biol.* **2015**, *87*, 472-479, doi:10.1111/jfb.12712.

41.   Bone, Q.; Moore, R.H. *Biology of Fishes* Taylor & Francis Group: New York, 2008.

42.   Scott, D.B.C.; Fuller, J.D. The reproductive biology of Scleropages formosus (Müller &
Schlegel) (Osteoglossomorpha, Osteoglossidae) in Malaya, and the morphology of its
pituitary gland. *J. Fish Biol.* **1976**, *8*, 45-53, doi:10.1111/j.1095-8649.1976.tb03906.x.

43.   Parenti, L.R.; Grier, H.J. Evolution and phylogeny of gonad morphology in bony fishes.
Integr. Comp. Biol. **2004**, *44*, 333-348, doi:10.1093/icb/44.5.333.

44.   Uttara, B.; Singh, A.V.; Zamboni, P.; Mahajan, R.T. Oxidative Stress and
Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant
Therapeutic Options. *Curr. Neuropharmacol.* **2009**, *7*, 65-74,
doi:10.2174/157015909787602823.

45.   Lopez-Cruz, R.I.; Dafre, A.L.; Filho, D.W. Oxidative Stress in Sharks and Rays. In *Oxidative*
Stress in Aquatic Ecosystems, Abele, D., Vazquez-Medina, J.P., Zenteno-Savin, T., Eds.
Wiley-Blackwell: 2011; pp. 157-164.

A taxonomically-restricted Uric Acid transporter provides insight into antioxidant equilibrium

Supplementary Figures:

Figure S1 – RaxML phylogenetic tree.

Figure S2 - Synteny maps of *Slc23a5*.

Figure S3 - Synteny maps of *Slc23a1*.

Figure S4 - Synteny maps of *Slc23a2*.

Figure S5 - Synteny maps of *Slc23a3*.

Figure S6 - Synteny maps of *Slc23a4.*

Figure S7 – S/T-X-Ø tripeptide conservation in Slc23a1 and Slc23a4 sequences.

Figure S8 – Topology prediction of Slc23a5.

Supplementary Tables:

Table S1 - Accession numbers.

Table S2 - Accession numbers of the RNAseq files.

Table S3 - Gene identifier for all genes included in gene expression analysis, collected from NCBI and ENSEMBLE Databases.

Table S4 - Genome and GTF files retrieved from Ensemble database (Release 89) and Transcriptome files retrieved from NCBI used for this study.

Table S5 - Relative gene expression levels of Slc23 family genes for eight species. The values are presented in log2 (TPM +1) ratios

Table S6 - Renal uric acid absorption and secretion: the urate transportome.

Figure S1 – RaxML phylogenetic tree. Node values represent bootstrap values.

Figure S2 - Synteny maps of *Slc23a5*. Conservation of the *Slc23a5* locus in major fish lineages, *Xenopus tropicalis* and *Homo sapiens*. Numbering after species corresponds to chromosome or scaffold. Information is presently absent for the Chondrostei *Acipenser ruthenus*. Grey boxes indicate non-conserved genes.

464 **Figure S3 - Synteny maps of *Slc23a1*.** Conservation of the *Slc23a1* locus in major vertebrate lineages. Numbering after species corresponds to

465 chromosome or scaffold. Grey boxes indicate non-conserved genes.

468   **Figure S4 - Synteny maps of *Slc23a2*.** Conservation of the *Slc23a2* locus in major vertebrate lineages. Numbering after species corresponds to

469   chromosome or scaffold. Grey boxes indicate non-conserved genes.

Figure S5 - Synteny maps of *Slc23a3*. Conservation of the *Slc23a3* locus in major vertebrate lineages. Numbering after species corresponds to chromosome or scaffold. Grey boxes indicate non-conserved genes.

477 **Figure S6 - Synteny maps of *Slc23a4*.** Conservation of the *Slc23a4* locus in major vertebrate lineages. Numbering after species corresponds to

478 chromosome or scaffold. The human homologue is indicated as pseudogenized (*SLC23A4P*). Grey boxes indicate non-conserved genes.

480 **Figure S7 – S/T-X-Ø tripeptide conservation in Slc23a1 and Slc23a4 sequences.**

Figure S8 – Topology prediction of Slc23a5 sequences: from *D. rerio*, *O. latipes*, *Salmo salar* (Atlantic salmon) and *X. tropicalis*, and human Slc23a1 used as control. TMHMM Server v. 2.0 (left) and hydropathy plots (right) are provided. Arrows detone the position of NAT family motif.

Supplementary Tables:

Table S1 - **Accession numbers.**

Species	Gene	Accession Number	Common name
Aspergillus nidulans	UapA	CAA50681	-
Aspergillus nidulans	UapC	CAA56190	-
Bacillus subtilis	PbuX	KIX81273	-
Bacillus subtilis	PucK	KIX81727	-
Escherichia coli	YgfO	CDU41213.1	-
Escherichia coli O25b:H4	UraA	ANK02789	-
Escherichia coli O25b:H4	YicE	ANK04183	-
Zea mays	Lep1	AAB17501	Maize
Branchiostoma belcheri	Slc23-like	XP_019630437	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019636191	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019633776	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019616280	Amphioxus
Branchiostoma belcheri	Slc23-like	XP_019646793	Amphioxus
Branchiostoma floridae	Slc23-like	XP_002600710	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002597306	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002595086	Florida lancelet
Branchiostoma floridae	Slc23-like	XP_002611161	Florida lancelet
Anolis carolinensis	Slc23a1	XP_008113637	Green anole
Callorhinchus milii	Slc23a1	XP_007904441	Ghost shark
Cyprinus carpio	Slc23a1	XP_018975086	Common carp
Danio rerio	Slc23a1	NP_001166970	Zebrafish
Gallus gallus	Slc23a1	XP_004944825	Chicken
Homo sapiens	Slc23a1	NP_005838	Human
Ictalurus punctatus	Slc23a1	XP_017348904	Channel catfish
Latimeria chalumnae	Slc23a1	XP_005989382	Coelacanth
Lepisosteus oculatus	Slc23a1	XP_015205203	Spotted gar
Lethenteron japonicum	Slc23a1	JL2986	Japanese lamprey
Monodelphis domestica	Slc23a1	XP_016278852	Gray short-tailed opossum
Mus musculus	Slc23a1	NP_035527	House mouse
Oryzias latipes	Slc23a1	XP_004076777	Japanese rice fish
Poecilia formosa	Slc23a1	XP_007556555	Amazon molly
Rhincodon typus	Slc23a1	XP_020385431	Whale shark
Xenopus tropicalis	Slc23a1	XP_002932349	Tropical clawed frog
Anolis carolinensis	Slc23a2	XP_003229574	Green anole
Danio rerio	Slc23a2	XP_021327808	Zebrafish
Gallus gallus	Slc23a2	XP_015152797	Chicken
Homo sapiens	Slc23a2	NP_976072	Human
Latimeria chalumnae	Slc23a2	XP_014352307	Coelacanth
Lepisosteus oculatus	Slc23a2	XP_015197789	Spotted gar
Lethenteron japonicum	Slc23a2	JL15308	Japanese lamprey
Leucoraja erinacea	Slc23a2	ctg10612	Little Skate
Monodelphis domestica	Slc23a2	XP_007476346	Gray short-tailed opossum
Mus musculus	Slc23a2	NP_061294	House mouse
Sarcophilus harrisii	Slc23a2	XP_003757995	Tasmanian devil

Scyliorhinus canicula	Slc23a2	ctg15352	Small-spotted catshark
Xenopus tropicalis	Slc23a2	NP_001120161	Tropical clawed frog
Anolis carolinensis	Slc23a3	XP_008108299	Green anole
Danio rerio	Slc23a3	XP_017213243	Zebrafish
Gallus gallus	Slc23a3	XP_015145565	Chicken
Homo sapiens	Slc23a3	NP_001138362	Human
Latimeria chalumnae	Slc23a3	XP_014343771	Coelacanth
Lepisosteus oculatus	Slc23a3	XP_015214926	Spotted gar
Monodelphis domestica	Slc23a3	XP_007501758	Gray short-tailed opossum
Mus musculus	Slc23a3	NP_919314	House mouse
Xenopus tropicalis	Slc23a3	XP_012826889	Tropical clawed frog
Anolis carolinensis	Slc23a4	XP_016849222	Green anole
Callorhinchus milii	Slc23a4	XP_007903282	Elephant shark
Danio rerio	Slc23a4	NP_001013353	Zebrafish
Gallus gallus	Slc23a4	XP_015145041	Chicken
Gekko japonicus	Slc23a4	XP_015276774	Schlegel's Japanese gecko
Latimeria chalumnae	Slc23a4	XP_005991386	Coelacanth
Lepisosteus oculatus	Slc23a4	XP_006633863	Spotted gar
Lethenteron japonicum	Slc23a4	JL7088	Japanese lamprey
Monodelphis domestica	Slc23a4	XP_007504433	Gray short-tailed opossum
Mus musculus	Slc23a4	XP_006506197	House mouse
Oryzias latipes	Slc23a4	XP_004069901	Japanese medaka
Rhincodon typus	Slc23a4	XP_020391140	Whale shark
Sarcophilus harrisii	Slc23a4	XP_012406506	Tasmanian devil
Xenopus tropicalis	Slc23a4	XP_012815547	Tropical clawed frog
Boleophthalmus pectinirostris	Slc23a5	XP_020774843	Great blue-spotted mudskipper
Cynoglossus semilaevis	Slc23a5	XP_016888339	Tongue sole
Cyprinus carpio	Slc23a5	XP_018944039	Common carp
Danio rerio	Slc23a5	XP_009296238	Zebrafish
Fundulus heteroclitus	Slc23a5	XP_021172742	Mummichog
Hippocampus comes	Slc23a5	XP_019750396	Tiger tail seahorse
Ictalurus punctatus	Slc23a5	XP_017328967	Channel catfish
Lates calcarifer	Slc23a5	XP_018548144	Barramundi perch
Lepisosteus oculatus	Slc23a5	XP_015208244	Spotted gar
Maylandia zebra	Slc23a5	XP_014264577	Zebra mbuna
Monopterus albus	Slc23a5	XP_020448997	Swamp eel
Nanorana parkeri	Slc23a5	XP_018412235	-
Neolamprologus brichardi	Slc23a5	XP_006782389	Fairy cichlid
Nothobranchius furzeri	Slc23a5	XP_015819289	Turquoise killifish
Oncorhynchus mykiss	Slc23a5	XP_021432371	Rainbow trout
Oreochromis niloticus	Slc23a5	XP_005454237	Nile tilápia
Oryzias latipes	Slc23a5	XP_004086320	Japanese medaka
Poecilia reticulata	Slc23a5	XP_008408845	Guppy
Pundamilia nyererei	Slc23a5	XP_005731628	-
Pygocentrus nattereri	Slc23a5	XP_017573200	Red-bellied piranha
Salmo salar	Slc23a5	XP_013981136	Atlantic salmon
Scleropages formosus	Slc23a5	XP_018602707	Asian bonytongue

Sinocyclocheilus rhinocerous	Slc23a5	XP_016366936	-
Stegastes partitus	Slc23a5	XP_008289900	Bicolor damselfish
Takifugu rubripes	Slc23a5	XP_003967776	Fugu rubripes
Xenopus tropicalis	Slc23a5	XP_002937113	Tropical clawed frog

490 **Table S2 - Accession numbers of the RNAseq files.** PE – portable executable format.

Spotted gar	Library	SRA Run	Read length
Brain	PE	SRR1524250	100
Intestine	PE	SRR1524257	100
Kidney	PE	SRR1524255	100
Liver	PE	SRR1524254	100
Testis	PE	SRR1524260	100
Zebrafish	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR1524238	100
Intestine	PE	SRR1524245	100
Kidney	PE	SRR1524243	100
Liver	PE	SRR1524242	100
Testis	PE	SRR1524249	100
Chicken	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR924542	100
Intestine	PE	SRR3194289	100
Kidney	PE	SRR924553	100
Liver	PE	SRR924555	100
Testis	PE	SRR924547	100
Human	**Library**	**SRA Run**	**Read length**
Brain	PE	ERR315432	100
Intestine	PE	ERR315419	100
Kidney	PE	ERR315468	100
Liver	PE	ERR315451	100
Testis	PE	ERR315456	100
Mouse	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR5171101	100
Intestine	PE	SRR5171080	100
Kidney	PE	SRR5171094	100
Liver	PE	SRR5171078	100
Testis	PE	SRR5171084	100
Western clawed frog	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR579560	75
Intestine	SE	SRR1186999	50
Kidney	PE	SRR579562	75
Liver	PE	SRR579561	75
Testis	SE	SRR943353	100
Anole lizard	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR579556	75
Kidney	PE	SRR579557	75
Liver	PE	SRR391651	100
Testis	PE	SRR5420127	50
Japanese medaka	**Library**	**SRA Run**	**Read length**
Brain	PE	SRR1524271	100
Intestine	PE	SRR1524278	100
Kidney	PE	SRR1524276	100
Liver	PE	SRR1524275	100
Testis	PE	SRR1628839	90

494 **Table S3 - Gene identifier for all genes included in gene expression analysis, collected**
495 **from NCBI and ENSEMBLE Databases.**

Organism	Gene Symbol	Gene ID NCBI	Gene ID Ensemble
Human	Slc23a1	9962	ENSG00000170482
	Slc23a2	9963	ENSG00000089057
	Slc23a3	151295	ENSG00000213901
Mouse	Slc23a1	20522	ENSMUSG00000024354
	Slc23a2	54338	ENSMUSG00000027340
	Slc23a3	22626	ENSMUSG00000026205
	Slc23a4	243753	ENSMUSG00000029847
Lizard	Slc23a1	100556489	ENSACAG00000016325
	Slc23a2	100566000	ENSACAG00000005986
	Slc23a3	100567833	ENSACAG00000003143
	Slc23a4	100563862	ENSACAG00000011027
Chicken	Slc23a1	416188	ENSGALG00000037642
	Slc23a2	419520	ENSGALG00000000195
	Slc23a3	429039	Not Found
	Slc23a4	417937	ENSGALG00000011717
Western clawed frog	Slc23a1	100493636	ENSXETG00000008402
	Slc23a2	100145200	ENSXETG00000016853
	Slc23a3	100486175	ENSXETG00000013878
	Slc23a4	100490216	ENSXETG00000020228
	Slc23a5	100486632	Not Found
Zebrafish	Slc23a1	559435	ENSDARG00000015033
	Slc23a2	100321249	ENSDARG00000017365
	Slc23a3	100333020	ENSDARG00000088891
	Slc23a4	503757	ENSDARG00000052497
	Slc23a5	798958	ENSDARG00000045792
Japanese medaka	Slc23a1	101155711	ENSORLG00000012701
	Slc23a2	101156743	ENSORLG00000002889
	Slc23a3	Not Found	Not Found
	Slc23a4[a]	101173186	ENSORLG00000009915
	Slc23a4b	101157211	ENSORLG00000015991
	Slc23a5	101175057	ENSORLG00000019882
Spotted gar	Slc23a1	102691341	ENSLOCG00000012533
	Slc23a2	102691255	ENSLOCG00000015630
	Slc23a3	102699040	ENSLOCG00000010877
	Slc23a4	102686145	ENSLOCG00000016950
	Slc23a5	2682847	ENSLOCG00000016369

499 **Table S4 - Genome and GTF files retrieved from Ensemble database (Release 89) and Transcriptome files retrieved from NCBI used for this**
500 **study.**

Organisms	Genome / Transcriptome Files	GTF FILE	Download date	Source
Mouse	Mus_musculus.GRCm38.dna.toplevel.fa	Mus_musculus.GRCm38.89.gtf	09/08/2017	Ensemble
Spotted gar	Lepisosteus_oculatus.LepOcu1.dna.toplevel.fa	Lepisosteus_oculatus.LepOcu1.89.gtf	09/08/2017	Ensemble
Anole lizard	Anolis_carolinensis.AnoCar2.0.dna.toplevel.fa	Anolis_carolinensis.AnoCar2.0.89.gtf	09/08/2017	Ensemble
Zebrafish	Danio_rerio.GRCz10.dna.toplevel.fa	Danio_rerio.GRCz10.89.gtf	09/08/2017	Ensemble
Human	Homo_sapiens.GRCh38.dna.toplevel.fa	Homo_sapiens.GRCh38.89.gtf	09/08/2017	Ensemble
Japanese medaka	Oryzias_latipes.MEDAKA1.dna.toplevel.fa	Oryzias_latipes.MEDAKA1.89.gtf	09/08/2017	Ensemble
Western clawed frog	GCF_000004195.3_Xenopus_tropicalis_v9.1_rna.fna	Xenopus_tropicalis_Gene_Transcript_Map.txt*	09/08/2017	Ncbi
Chicken	GCF_000002315.4_Gallus_gallus-5.0_rna.fna	Gallus_gallus_Gene_Transcript_Map.txt*	09/08/2017	Ncbi

504 **Table S5 - Relative gene expression levels of Slc23 family genes for eight species. The values are presented in log2 (TPM +1) ratios.**

Organism	Tissue	Slc23a1	Slc23a2	Slc23a3	Slc23a4a	Slc23a4b	Slc23a5
Human	BRAIN	0,00	5,82	0,00	-	-	-
	INTESTINE	5,85	2,64	4,61	-	-	-
	KIDNEY	3,71	3,70	4,28	-	-	-
	LIVER	4,03	3,58	2,36	-	-	-
	TESTIS	0,00	4,59	0,00	-	-	-
Mouse	BRAIN	0,00	4,91	0,00	0,00	-	-
	INTESTINE	3,73	5,24	1,02	5,89	-	-
	KIDNEY	6,91	2,82	4,08	0,00	-	-
	LIVER	5,06	2,85	0,00	0,00	-	-
	TESTIS	0,00	1,63	0,00	0,00	-	-
Lizard	BRAIN	0,91	2,82	0,72	2,49	-	-
	KIDNEY	6,76	1,39	6,10	1,19	-	-
	LIVER	3,02	0,00	0,00	0,65	-	-
	TESTIS	0,00	0,00	0,00	0,00	-	-
Chicken	BRAIN	0,00	6,27	0,96	1,82	-	-
	INTESTINE	6,44	3,25	2,96	6,09	-	-
	KIDNEY	7,78	1,49	4,17	1,60	-	-
	LIVER	5,20	1,98	1,76	0,82	-	-
	TESTIS	2,35	2,08	2,03	1,85	-	-
Western clawed frog	BRAIN	0,00	6,45	0,00	2,54	-	0,00
	INSTESTINE	0,00	2,23	0,00	5,65	-	0,00
	KIDNEY	7,63	4,08	6,53	5,00	-	5,40
	LIVER	0,00	3,41	0,00	2,14	-	0,00
	TESTIS	2,69	4,01	0,00	1,49	-	1,64
Zebrafish	BRAIN	0,85	3,73	0,71	0,00	-	1,01
	INTESTINE	2,69	1,21	0,00	4,00	-	0,00
	KIDNEY	5,35	2,27	3,42	4,78	-	0,93
	LIVER	0,00	1,26	0,00	0,00	-	0,00
	TESTIS	0,93	2,74	0,00	0,63	-	1,97
Japanese medaka	BRAIN	0,00	4,70	-	0,00	0,00	0,84
	INTESTINE	2,76	2,91	-	4,95	0,00	0,00
	KIDNEY	3,89	3,02	-	4,43	0,00	0,00
	LIVER	0,00	4,06	-	0,00	0,00	0,00
	TESTIS	0,00	3,20	-	0,60	0,00	0,00
Spotted gar	BRAIN	0,58	4,94	0,00	0,00	-	0,00
	INTESTINE	5,67	0,96	0,00	0,00	-	0,00
	KIDNEY	3,18	0,90	1,78	0,86	-	0,00
	LIVER	0,74	0,78	0,00	0,00	-	0,00
	TESTIS	1,92	7,21	0,00	0,00	-	0,00

Table S6 - **Renal uric acid absorption and secretion: the urate transportome.** n.f. – not found.

	Transporter	*H. sapiens*	*M. musculus*	*G. gallus*	*A. carolinensis*	*X. tropicalis*	*D. rerio*	*L. oculatus*
Basolateral Absorption	SLC2A9/GLUT9	✓	✓	✓	✓	✓	✓	✓
Apical Absorption	SLC22A11/OAT4	✓	n.f.	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC22A12/URAT1	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC22A13/OAT10	✓	✓	✓	✓	✓	✓	✓
Basolateral Secretion	SLC22A6/OAT1	✓	✓	✓	✓	✓	n.f.	n.f.
	SLC22A8/OAT3	✓	✓				n.f.	n.f.
Apical Secretion	SLC17A1/NPT1	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.
	SLC17A3/NPT4	✓	✓	n.f.	n.f.	n.f.	n.f.	n.f.