Full title: Molecular and electrophysiological features of spinocerebellar ataxia type seven in

induced pluripotent stem cells

Short title: Spinocerebellar ataxia type seven and induced pluripotent stem cells

RJ Burman[1,2,3¶*], LM Watson[4,5¶], DC Smith[4,5¶], JV Raimondo[1,2,5], R Ballo[1], J Scholefield[6],

SA Cowley[7], MJA Wood[8], SH Kidson[1,5], LJ Greenberg[4,5]

[1]Department of Human Biology University of Cape Town, Cape Town, Western Cape, South

Africa

[2]Neuroscience Institute, University of Cape Town, Cape Town, Western Cape, South Africa

[3]Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, Oxfordshire,

United Kingdom

[4]Department of Pathology, University of Cape Town, Cape Town, Western Cape, South

Africa

[5]Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape

Town, Western Cape, South Africa

[6]Gene Expression & Biophysics Group, Synthetic Biology ERA, CSIR Biosciences, Pretoria,

Gauteng, South Africa

[7]Sir William Dunn School of Pathology, University of Oxford, Oxford, Oxfordshire, United

Kingdom

[8]Department of Paediatrics, University of Oxford, Oxford, Oxfordshire United Kingdom

[*]Corresponding authors:

richard.burman@ndcn.ox.ac.uk (RJB)

[¶]These authors contributed equally to this work

Abstract

Spinocerebellar ataxia type 7 (SCA7) is an inherited neurodegenerative disease caused by a polyglutamine repeat expansion in the *ATXN7* gene. Patients with this disease suffer from a degeneration of their cerebellar Purkinje neurons and retinal photoreceptors that result in a progressive ataxia and loss of vision. As with many neurodegenerative diseases, studies of pathogenesis have been hindered by a lack of disease-relevant models. To this end, we have generated induced pluripotent stem cells (iPSCs) from a cohort of SCA7 patients in South Africa. First, we differentiated the SCA7 affected iPSCs into neurons which showed evidence of a transcriptional phenotype affecting components of STAGA (*ATXN7* and *KAT2A*) and the heat shock protein pathway (*DNAJA1* and *HSP70*). We then performed electrophysiology on the SCA7 iPSC-derived neurons and found that these cells show features of functional aberrations. Lastly, we were able to differentiate the SCA7 iPSCs into retinal photoreceptors that also showed similar transcriptional aberrations to the SCA7 neurons. Our findings demonstrate that iPSC-derived neurons and photoreceptors from SCA7 patients express molecular and electrophysiological differences that are indicative of impaired neuronal health. We hope that these findings will contribute towards the ongoing efforts to establish the cell-derived models of neurodegenerative diseases that are needed to develop patient-specific treatments.

Introduction

Spinocerebellar ataxia type 7 (SCA7) is an inherited neurodegenerative disease caused by a CAG repeat expansion in the *ATXN7* gene. Since the translation of this CAG repeat leads to an expanded polyglutamine (polyQ) tract within the resultant protein, SCA7 is classified as a polyQ–repeat disorder. Other diseases with a similar pathophysiology include five different SCAs (SCA 1, 2, 3, 6 and 17), as well as Huntington disease, dentatorubral-pallidoluysian atrophy and spinal bulbar muscular atrophy [1]. Clinically, SCA7 patients present with ataxia, dysarthria and visual loss. This is caused by a selective degeneration of cerebellar Purkinje neurons and retinal photoreceptors [2]. Symptoms progressively worsen over a period of 10 to 30 years, leading ultimately to brainstem dysfunction, blindness, physical disability and death.

The mechanism by which a polyQ expansion within the ubiquitously expressed ATXN7 protein leads to the selective degeneration of Purkinje neurons and photoreceptors remains to be fully elucidated. ATXN7 is known to be a component of the mammalian transcription co-activator complex, STAGA (SPT3-TAF9-ADA-GCN5 acetyltransferase) [3]. This protein has been shown to facilitate the interaction between STAGA and the cone-rod homeobox (CRX) transactivator of photoreceptor genes, linking the function of ATXN7 with the retinal phenotype observed in SCA7 patients [4]. In addition, several studies have highlighted the role of transcriptional aberrations in the neuronal cell dysfunction that precedes the onset of disease symptoms [3-7]. These gene expression changes may arise either directly from alterations in transcriptional regulation by mutant ATXN7, or indirectly, as a consequence of a generalised cellular response to the presence of mutant ATXN7. More recently,

dysfunction in cell metabolism has been proposed as an additional pathogenic mechanism

[8, 9].

As with many neurodegenerative conditions, research into the molecular pathogenesis of

SCA7 has been hindered by a lack of suitable models of human disease progression. This

is particularly relevant in cases where the genomic context of the mutation may have an

impact on gene function and might prove useful for therapeutic development. There does,

however, seem to be increasing momentum in this area with new models being proposed

and refined [10-12].

SCA7 occurs at an unusually high frequency in the South African population as a result of a

founder effect in patients of Black African ethnic origin [13]. South African SCA7 patients

also display a unique phenomenon – a single nucleotide polymorphism (SNP) within *ATXN7*

(rs3774729), which is linked to the mutation in all patients studied to date [14].

Approximately 43% of these individuals are heterozygous for the polymorphism, allowing for

allelic discrimination, and providing an ideal target for developing an allele-specific silencing

therapy. This haplotype has been shown to extend into other Southern African populations,

suggesting that such a therapy may be more widely applicable than was first thought [15].

We have previously demonstrated the efficacy of an allele-specific RNAi treatment in an

over-expression cell model of SCA7 [16], as well as in SCA7 patient fibroblasts [17].

Disease-relevant cell lines generated from these patients are thus of vital importance in the

understanding of disease pathogenesis and the development of therapies, as they carry the

patient's full genomic sequence, including SNPs which may be used as targets for gene

silencing.

The generation of induced pluripotent stem cells (iPSCs) involves reprogramming somatic

cells to a pluripotent state by means of viral transduction with the pluripotency genes *OCT4*,

SOX2, *KLF4* and *c-MYC* [18, 19]. Importantly, iPSCs can then be differentiated into any

tissue of the body through treatment with specific growth factors. This makes iPSCs a

useful starting point for the generation of disease-relevant cell models, particularly in

neurodegenerative diseases, where primary CNS cultures may only be obtained using

invasive methods.

In this study, we have generated and characterised iPSC lines from two South African

SCA7 patients and an unaffected, related control. We were then able to differentiate these

iPSCs into cells expressing markers associated with retinal photoreceptors, neural

progenitor cells (NPCs) and neurons. Thereafter, we have obtained preliminary evidence for

a disease phenotype in these cells, in the form of pathogenically relevant gene expression

changes and alterations in intrinsic neuronal properties. Our findings indicate that these

cells may be able to replicate, to some extent, the pathogenesis underlying SCA7.

Material and methods

Ethics approval and patient recruitment

Ethics approval for the study was granted by the University of Cape Town (UCT) Faculty of

Health Sciences Human Research Ethics Committee (HREC REF. 380/2009 and

434/2011), and was renewed annually, incorporating amendments to the project protocol

where necessary. All methods were carried out in accordance with the guidelines approved

by the Ethics Committee. Participants were recruited from the Neurogenetics clinic at

Groote Schuur Hospital in Cape Town, where they were counselled by a genetic counsellor

and a clinical geneticist. Written informed consent was obtained from all participants prior

to their enrolment in the study.

Establishment of primary fibroblast cultures

110 Primary dermal fibroblast cultures were established from punch skin biopsies [20] taken

111 from the inner forearm of two unrelated SCA7 patients (47Q and 41Q) and an unaffected

112 control individual (CON, sibling of 41Q) who had consented to participate in the study. CAG

113 genotypes and ages at diagnosis and biopsy are shown in S1 Fig.

Generation and characterisation of patient-derived iPSCs

Dermal fibroblasts were reprogrammed into iPSCs using a replication-defective and

persistent Sendai virus vector (SeVdp) containing OCT4, SOX2, KLF4 and c-MYC as

previously described [21, 22]. One week after infection, the cells were transferred to

mitomycin C-inactivated mouse embryonic fibroblast feeder layers in stem cell medium (KO

DMEM, 20% KOSR, 1% NEAA, 50 µM β-mercaptoethanol, glutamax, 10ng/ml bFGF). Half

of the medium was refreshed every day and colonies were expanded by manual passaging

(dissection of colonies using needles) on feeder layers as previously described by Schwartz

et al [23]. After 10 passages the expression of pluripotency markers OCT4 and TRA-1-60,

as well as silencing of the reprogramming SeVdp vector, were confirmed by

immunocytochemistry (antibodies listed in S1-2 Tables). The expression of selected

pluripotency genes (OCT4, SOX2, NANOG) was determined by quantitative real-time

reverse transcriptase PCR (qRT-PCR).   Genomic integrity was assessed by means of

karyotype analysis (G-banding, S2 Fig). To confirm the iPSC lines' capacity to differentiate

into the three embryonic germ layers, in vitro differentiation via embryoid body formation

was performed as previously described [24].

Following SeVdp-mediated reprogramming, iPSC colonies with the correct morphology (flat,

with distinct borders, containing tightly packed cells with a high nucleus-to-cytoplasm ratio)

appeared within three to four weeks. These colonies were manually picked and clonally

expanded in separate dishes on a feeder layer of inactivated mouse embryonic fibroblasts.

Two SCA7 patient iPSC lines and two control lines (representing two affected individuals

and a single related, unaffected control) were successfully generated and characterised.

Immunocytochemical analysis revealed iPSC colonies with distinct nuclear staining for the

pluripotency transcription factor, *OCT4*, compared to the surrounding feeder fibroblasts (S3

Fig, A – top panel). The iPSC colonies also stained positive for the embryonic stem cell

surface marker TRA-1-60 (S3 Fig, A – bottom panel). In addition, we confirmed the

differentiation capacity of these cells into mesoderm, ectoderm and endoderm (S4 Fig).

Co-staining of all iPSC lines with primary antibodies against OCT4 and the nucleocapsid

protein of the virus (anti-NP) showed little or no evidence of NP staining in the OCT4-

positive pluripotent cells (S3 Fig, B –bottom panel), compared to intense cytoplasmic

staining in newly infected fibroblasts (S3 Fig, B – top panel). This indicated that the Sendai

virus had been effectively silenced, and that the iPSCs had achieved self-regulating

pluripotency. All lines were assessed after passage 8. The expression of pluripotency

markers was further confirmed by qRT-PCR. All five iPSC lines expressed high levels of

OCT4, SOX2 and *NANOG*, standard markers of pluripotency (S3 Fig, C - E), compared to

donor fibroblasts or cells that had been subjected to retinal or neuronal differentiation. The

expression levels of *SOX2* and *NANOG* were similar across the five iPSC lines, but lines

41Q and CON$_B$ showed lower levels of *OCT4* expression compared to the remaining three

lines (although still significantly higher than differentiated fibroblasts and retinal cells).

Neural differentiation and characterisation

Differentiation of iPSCs into neural progenitors was performed by treatment of iPSCs with

3µM glycogen synthase kinase 3 (GSK3) inhibitor (CHIR99021) and 2µM TGFβ inhibitor

(SB431542) as previously described, with neural progenitors appearing in culture after

seven days. For neuronal differentiation, neural progenitors were seeded at a density of

150 000 cells/well onto a Matrigel-coated six-well plate in neural induction medium [25].

After two days, medium was changed to neuronal differentiation medium (DMEM/FBS

supplemented with N2, B27, 300ng/ml cyclic AMP (Sigma), 0.2mM ascorbic acid (Sigma),

10ng/ml BDNF (Peprotech) and 10ng/ml GDNF (Peprotech)) and the cells maintained in

culture for 14 to 21 days, with neurite outgrowth typically observed after one week of

culture. Medium was changed every 2-3 days. Characterisation was performed by

immunocytochemistry and qRT-PCR (for antibodies and primers see S1-3 Tables).

Electrophysiology

Neurons cultured between 2-3 weeks on glass cover slips (uncoated) were removed from

the incubator and rapidly transported to the recording chamber of a Zeiss Axioskop Upright

Microscope (Zeiss). Electrophysiological recordings were made in neuronal differentiation

medium at room temperature and were restricted to the first 5 hours following cell removal

from the incubator environment. Patch pipettes of 13-20 MOhm tip resistance were pulled

from filamental borosilicate glass capillaries (2.00 mm outer diameter, 1.58 mm inner

diameter, Hilgenberg), using a horizontal puller (Model P-1000, Sutter). The pipettes were

filled with an internal solution containing (in mM): K-gluconate (126); KCl (4); Na 2 ATP (4);

NaGTP (0.3); Na 2 -phosphocreatinine (10) and HEPES (10). Osmolarity was adjusted to

between 290 and 300 mOsM and the pH was adjusted to between 7.38 and 7.42 with KOH.

Cells were visualised using a 40x water-immersion objective (Zeiss). Digital images were

obtained using a CCD camera (VX55, TILL Photonics). Individual cells were selected for

recordings based on a small round or ovoid cell body (diameters, 5–10 µm) and typically

two or more extended processes. Recordings were made in current-clamp and voltage-

clamp mode using an Axopatch 200B amplifier (Molecular Devices). Data acquisition was

performed through an ITC-1600 board (Instrutech) connected to a PC running a custom-

written routine (PulseQ) under IGOR Pro (Wavemetrics). Analysis was performed using

custom-written scripts in MATLAB (Mathworks). Statistical analysis was performed using

GraphPad Prism (v8, GraphPad Software). Normally distributed data are presented with

mean and standard error of the mean (SEM) whilst non-normally distributed data are

presented with median and interquartile range (IQR). Appropriate parametric and non-

parametric tests were performed including unpaired Students t-test, Mann-Whitney U test

and the Kruskal-Wallis test. Significance was defined as $p < 0.05$. For a comparison of

means across all cell lines, an ANOVA test was used. Assessment of categorical variables

was done using a Chi-Squared test.

191   Retinal differentiation and characterisation

Differentiation into retinal photoreceptors was performed as previously described [26], using

iPSCs cultured in feeder-free conditions on Matrigel in mTESR™ medium. The cells were

dissociated enzymatically and plated onto Matrigel-covered dishes in neural differentiation

medium containing N2 and B27 supplements (Life Technologies). After settling for an hour,

adhered cells were covered in a 2% Matrigel solution. The following day, and thereafter

every second day, the medium was replaced with neural differentiation medium without

Matrigel. From day 10 the medium was supplemented with 3nM recombinant SHH (R&D

Systems), 50ng/µl acidic fibroblast growth factor (αFGF) (R&D Systems), 10ng/µl basic

fibroblast growth factor (bFGF) (Miltenyi), 1mM taurine and 500nM retinoic acid (both Sigma

Aldrich). After 30 days of culture characterisation was performed by immunocytochemistry

and qRT-PCR (for antibodies and primers, see S Tables 1 and 2). Each patient and control

line underwent two rounds of differentiation, with each time point analysed in biological

duplicate.

205   DNA and RNA isolation, cDNA synthesis

206   DNA extraction from cultured cells was performed using the QIAGEN DNeasy Blood and

207   Tissue Kit. RNA was isolated from cultured cells using the QIAGEN RNeasy Plus Mini Kit.

208   Synthesis of cDNA was performed using the Applied Biosystems High Capacity cDNA

209   Reverse Transcription Kit (Life Technologies) using 500ng-1µg template RNA.

210   Quantitative RT-PCR

qRT-PCR was performed on the BioRad CFX96 Real-Time PCR System, using the Power

SYBR Green PCR Master Mix (Applied Biosystems), according to manufacturer's

instructions. Primers (S3 Table) were obtained from PrimerDesign Ltd, Integrated DNA

Technologies (IDT), or Sigma-Aldrich. Relative quantities of target mRNA were determined

using the relative standard curve method [27]. Standard curves were prepared for each

primer pair, from serial dilutions of pooled sample cDNA. Universal cycling conditions were

used (95°C for 10min, followed by 40 cycles of 95°C for 15 seconds and 60°C for 1min).

PCRs were performed in technical triplicate, on at least two biological replicates, and results

were analysed using the BioRad CFX Manager software (v3.1).   Gene of interest

expression was normalised to *β Actin* (*ACTB*) expression in each case. The design and

reporting of qRT-PCR experiments aimed to comply with the Minimum Information for

publication of Quantitative real-time PCR Experiments guidelines [28, 29] wherever

possible. Statistical analysis was performed using the unpaired Students' t-test. Significance

was defined as $p < 0.05$.

CAG repeat length determination

The length of the disease-causing CAG repeat in *ATXN7* was determined from DNA by

means of PCR and automated fluorescent genotyping. The PCR reaction mix consisted of

0.4µM each, forward and reverse primer (S3 Table), 0.6units of GoTaq DNA polymerase

(Promega), 100ng of DNA, and Failsafe buffer J (Epicentre Biotechnologies) at a final

concentration of 1X, made up to a final reaction volume of 10µl. Cycling conditions were as

follows: 95°C for 5min; followed by 30 cycles of 95°C for 30 seconds, 53°C for 6 seconds

and 72°C for 40 seconds; and a final elongation step at 72°C for 7min. Automated

fluorescent genotyping was performed using the ABI 3130*xl* Genetic Analyzer (Applied

Biosystems).

The length of the CAG repeat [$(CAG)_n$] was approximated using the following equation,

adapted from (Dorschner et al., 2002): $n = (0.3063$ x Length of major PCR product in base

pairs) $- 76.475$bp. The major PCR product was defined as the product generating the

highest fluorescent peak, as detected using the ABI 3130*xl* Genetic Analyzer.

Results

SCA7 iPSC-derived neural progenitor cells show transcriptional disease-associated aberrations

SCA7 patient and control iPSCs generated neural progenitor cells (NPCs) at comparable efficiencies, when cultured in neural induction medium supplemented with SB431542 (TGFβ inhibitor) and CHIR99021 (GSK3 inhibitor), with 100% of the cells expressing the early neural marker Nestin, after five passages (Fig 1A). The cells also stained positive for the disease-causing protein, ATXN7 (Fig 1B), and demonstrated repression of the pluripotency gene *OCT4*, and upregulation of the early neural gene *PAX6* (Fig 1C).

Fig 1: SCA7 iPSC-derived NPCs show transcriptional aberrations. (A) Representative images of control (top panel) and SCA7 patient (bottom panel) NPCs, expressing the early neural marker, Nestin (green). (B) High magnification images showing nuclear expression of ATXN7 (red) in control (top) and patient (bottom) NPCs (Green: βIII-Tubulin). (C) qPCR results indicating suppression of *OCT4* (top panel) and upregulation of *PAX6* (bottom panel) expression in NPCs, compared with the iPSC lines from which they were derived. All levels shown relative to beta (β)-actin. (D) Expression of 9 selected genes in SCA7 patient NPCs ($n = 2$) shown relative to unaffected control NPCs ($n = 2$). All levels shown relative to β actin and unaffected control cells. (E) Control (top panel) and SCA7 NPC-derived neurons (bottom panel) stain positive for βIII-Tubulin (green) and ATXN7 (red) after 14 days of differentiation. (F) Both control and SCA7 patient NPCs produced a small proportion of GABA-positive neurons (red) after 14 days in culture (Green, βIII-Tubulin). *$p < 0.05$.

To determine whether any transcriptional differences could be detected between SCA7

patient- and unaffected control-derived cell types, a panel of candidate genes was selected

which had previously been shown to be dysregulated in the retinal and cerebellar tissue of

SCA7 mouse models and patient lymphoblasts [5, 6, 30, 31]. The panel included the

following genes: *ATXN7,* brain expressed, X-linked 1 (*BEX1*), DnaJ (Hsp40) homolog,

subfamily A, member 1 (*DNAJA1*), glutamate receptor, ionotropic, AMPA 2 (*GRIA2*), heat

shock protein 27 (*HSP27*), heat shock protein 70 (*HSP70*), heat shock protein 105

(*HSP105*), K (lysine) acetyltransferase 2A (*KAT2A*), Phospholipase C, Beta 3 (*PLCB3*) and

ubiquitin carboxyl-terminal esterase L1 (*UCHL1*). The expression of these genes was

determined by means of qRT-PCR in fibroblasts (S5 Fig) and NPCs (Fig 1D).

In contrast to the fibroblasts and undifferentiated iPSCs, two genes were downregulated in

SCA7 patient iPSC-derived NPCs when compared to the unaffected control lines. These

included the disease-causing gene *ATXN7* (p = 0.018 in NPCs); and *KAT2A*, encoding

GCN5, the histone acetyltransferase (HAT) component of the STAGA transcription

coactivator complex (p = 0.003 in NPCs). In addition, NPC-specific alterations in

expression were identified in the heat shock protein genes DNAJA1 (p = 0.04) and HSP70

(p = 0.04), both of which were found to be downregulated in SCA7-patient derived cells.

Lastly, BEX1, an interactor of the p75 neurotrophin receptor, which regulates neurotrophin

signalling and neuronal differentiation, was found to be downregulated in the SCA7 NPCs (p

= 0.03).

In order to confirm the size of the *ATXN7* CAG repeat alleles in mRNA from patient- and

control-derived fibroblasts, iPSCs and NPCs an RT-PCR-based assay was performed. The

results were visualised on an agarose gel (S6 Fig panel A) and confirmed by automated

fluorescent genotyping. The length of the CAG repeat did not appear to fluctuate during

reprogramming or differentiation, corresponding to previous reports in similar cell lines [32,

33]. This assay also confirmed that both the mutant and wild-type *ATXN7* alleles were

expressed by all cell types, and that there were no obvious differences in allele expression

in affected or unaffected cells.

After 14 days culture in neuronal differentiation medium containing N2/B27 supplement,

cAMP ascorbic acid, BDNF and GDNF [25], both SCA7 patient and control NPCs stained

positive for the neuronal marker βIII-Tubulin, and showed robust, diffuse nuclear localisation

of the disease-causing protein, ATXN7 (Fig 1E-F). A subset of cells (approximately 1.8%)

stained positive for gamma-aminobutyric acid (GABA), although the proportion of

GABAergic neurons did not appear to vary between SCA7 patients and controls (Fig 1E-F).

No obvious differences in morphology were observed when comparing neurons derived

from SCA7 patient iPSCs with those derived from controls. In order to test functional

differences between SCA7 neurons and controls, electrophysiological measurements were

performed on a subset of these cells.

299 SCA7 iPSC-derived neurons demonstrate functional differences

300 that may affect cell excitability

Cells were assayed for physiological properties between 14- and 23-days post induction of

neuronal differentiation. Cells were targeted for whole-cell recordings based on their

morphological properties. This included a small round or ovoid cell body with diameters

between 5–10 µm and typically two or more extended processes. Following the attainment

of a whole-cell patch, current pulses of between 0 and 10 pA were applied. Individual cells

displayed four general types of spiking responses: a purely passive (Fig 2A), abortive spike

(Fig 2B), single spike (Fig 2C) and recurrent spiking response (Fig 2D). These spiking

properties are similar to those observed in acute human fetal brain slices [34], hESC-

derived neurons [35-37] and iPSC-derived neurons [38]. A postmitotic neuron matures by

inserting voltage-gated channels into its plasma membrane [39]. Therefore, the spiking

response of a cell to current injection can be used to determine the maturation stage of a

differentiating neuron: passive (least mature) → abortive spike → single spike → recurrent

spikes (most mature).

Fig 2: Patch-clamp analyses of iPSC derived neuronal cultures. Cells could be divided

into four categories based on their spiking responses. (A-D) Differential interference

contrast images of cells targeted for patch-clamp recordings (top). Whole-cell recordings in

current clamp mode (middle) depict spiking patterns following injection of current (bottom).

Cells fell into (A) 'passive', (B) 'abortive spike', (C) 'single spike' and (D) 'recurrent spikes'

categories. (E) The fraction of cells which fell into each category for cells derived from two

control and two SCA7 patient iPSC lines. Note that although the cells derived from various

iPSC lines exhibited different distributions of spiking responses, no trend between control

and patient lines was observed. (F) The resting membrane potentials (Vm) of cells varied

significantly across each cell line. Error bars denote mean median and IQR. ** $p < 0.01$, ***

$p < 0.001$.

Spiking responses were collected in current-clamp mode from cells derived from four

separate iPSC lines: two control lines CON_A ($n = 70$) and CON_B ($n = 42$), and two patient

lines 47Q ($n = 44$) and 41Q ($n = 72$). Although the fraction of cells which fell into each

spiking response category was significantly dependent on the iPSC line from which the cells

were derived ($p < 0.0001$, *Chi-squared test*), no trend could be discerned between control

and patient lines (Fig 2E). For example, the control line CON_A had the most mature

phenotype with the highest fraction of cells in the single spike and recurrent spiking

categories whilst 47Q demonstrated a relatively immature phenotype, with the majority of

cells displaying spiking responses falling into the passive category. However, the other

lines did not corroborate this difference as the control line CON_B displayed a less mature

phenotype than patient line 41Q.

Next, we compared the resting membrane potential (Vm) of cells derived from each cell line.

This parameter was significantly dependent on the iPSC line from which the cells were

derived (Fig 2F, $p = 0.002$, *ANOVA*). The mean resting membrane potential +/- SEM for the

CON_A, CON_B, 47Q and 41Q cell lines were -57.0 +/- 1.7, -54.7 +/- 3.2, -67.3 +/- 2.7 and -

62.5 +/- 2.5 mV respectively.

Next, we performed voltage-clamp recordings of the neurons in order to measure the input

resistance as well as the voltage-gated sodium and potassium currents (Fig 3A-B). A lower

input resistance is associated with neurite outgrowth and increased numbers of ion

channels inserted into the plasma membrane during the process of neuronal maturation.

Once again, a cell's input resistance was significantly dependent on the cell line to which it

belonged (Fig 3C, $p = 0.005$, *ANOVA*). Input resistance +/- SEM was 4987 +/- 421, 5484 +/-

243, 5979 +/- 513 and 7094 +/- 455 mΩ for the CON_A, CON_B, 47Q and 41Q cell lines

respectively.

Fig 3: Sodium and potassium currents in neurons differentiated from control and

patient iPSC lines. (A) Voltage-clamp recording trace. Membrane currents (top) were

recorded following 10 mV voltage steps between -90 and 30 mV (bottom). Maximum

potassium current (I_K) indicated. Dashed gray rectangle indicates the position of voltage-

gated sodium currents (enhanced in 'B'). Maximum sodium current (I_{Na}) is indicated 'right'.

(B) Zoom-in of rectangle in '**A**' with the maximum I_{Na} highlighted. (C) Cell input resistance

was also significantly varied across cell lines. (D) Population data from the four iPSC lines

for maximum voltage-gated I_K and I_{Na} currents showing significant variability across cell

lines. Error bars denote median and IQR. *** $p < 0.001$, **** $p < 0.0001$.

As one would predict, the spiking properties of neurons are directly correlated with the size

of their currents. The maximum size of voltage-gated sodium currents and potassium

currents measured in each cell (Max. I_K and Max. I_{NA}) was significantly dependent on the

particular iPSC line concerned (Fig 3D and E, $p < 0.0001$ in both cases, *ANOVA*). The

mean Max. I_K +/- SEM was 180.1 +/- 16.5, 176 +/- 10.8, 222.5 +/- 19.3 and 105.7 +/- 8.0 pA

for the CON_A, CON_B, 47Q and 41Q cell lines respectively. The mean Max. I_{NA} +/- SEM was

-166.5 +/- 18.6, -277.1 +/- 23.0, -268.0 +/- 30.5 and -108.0 +/- 14.2 pA for the CON_A, CON_B,

47Q and 41Q cell lines.

SCA7 iPSC-derived retinal photoreceptors show transcriptional aberrations

For the retinal differentiation of iPSCs, the Matrigel "sandwich" system facilitated a rapid

self-organisation and differentiation of the pluripotent stem cells into structures containing

cells morphologically indicative of columnar neuroepithelia.   These structures lost their

integrity from day 4-5, and cells spread into an adherent monolayer.   The gradual

emergence of cells with a neuronal morphology was observed from day 10 to day 30,

particularly in areas of low confluence. Following differentiation of patient and control cells

into retinal cells, immunocytochemical analyses were performed on cells at the end of the

differentiation period (day 30), to determine whether the cells expressed retinal cell

markers. The cells were stained for either the disease-causing protein ATXN7 (Fig 4A) or

the retinal cell markers CRX and RCVRN (Fig 4B-C). No obvious differences in morphology

were observed between SCA7 patient and control iPSC-derived cells. The differentiated

cells displayed varying expression levels of the retinal genes *CRX, PAX6, RCVRN* and

OTX2 (Fig 4D-G).

Fig 4: iPSC-derived retinal cells exhibit aberrant transcriptional changes.

Immunocytochemical analysis of control (top panel, representative images from line CON_B)

and SCA7 (bottom panel, representative images from patient line 47Q) retinal cells showed

positive staining for ATXN7 (A), cone-rod homeobox (CRX, B) and recoverin (RCVRN, C)

on day 30 of the differentiation experiment (nuclei in blue). qRT-PCR results confirmed the

expression of *CRX* (D), *PAX6* (E), *RCVRN* (F) and *OTX2* (G) in all five lines, compared with

the iPSC lines from which they were derived. All levels shown relative to β actin. (H)

Expression of 23 selected genes in SCA7 patient iPSC-derived retinal cells (*n* = 2) shown

relative to unaffected control cells (*n* = 1). All levels shown relative to beta actin and

unaffected control cells. The data in this figure includes the cell lines from both patient cell

lines (47Q and 41Q) relative to the mean the control lines (CON_A and CON_B). A comparison

against both control lines was not deemed necessary as these were shown to have similar

expression profiles. *$p < 0.05$.

To determine transcriptional changes in the retinal cells, retinal cell specific genes were

added to the gene panel used in the NPCs. The retinal genes included: arrestin 3 (*ARR3*);

cone-rod homeobox (*CRX*); guanine nucleotide binding protein (G protein); alpha

transducing activity polypeptide 1 (*GNAT1*); Microphthalmia-associated transcription factor

(*MITF*); neural retina leucine zipper (*NRL*); orthodenticle homeobox 2 (*OTX2*); paired box 6

(*PAX6*); recoverin (*RCVRN*); rhodopsin (*RHO*) and retinal pigment epithelium-specific

protein 65kDa (*RPE65*). As observed in the NPC cells, both ATXN7 (p = 0.04) and KAT2A

(p = 0.02) were down regulated in the retinal cells. In addition, we also noticed expression

changes in genes that appeared to be specific to the SCA7 photoreceptors. These included

GRIA2, encoding the glutamate receptor, ionotropic, AMPA2 (GluR2) (downregulated, p =

0.04); and three retinal-specific genes, *OTX2* (upregulated, p = 0.002), *RCVRN*

(downregulated, p = 0.01) and *RPE65* (upregulated, p = 0.001). Lastly, and in contrast to

the SCA7 NPCs, BEX1 expression appear to be upregulated in SCA7 photoreceptors (p =

0.0006). The opposing BEX1 transcription changes (up in photoreceptors and down in

NPCs) may be indicative of a possible differential response by the different cell types to the

presence of mutant ATXN7. As with the NPCs, *ATXN7* CAG repeat alleles were confirmed

using an RT-PCR (S Fig 5B).

Discussion

Discussion

This study describes the generation and characterisation of the first iPSCs from the South

African SCA7 patient cohort, through the transduction of patient dermal fibroblasts with

Sendai virus vectors (17, 18). Furthermore, we show that we are able to differentiate these

iPSCs from SCA7 into neurons and retinal photoreceptors. The SCA7 patient iPSCs

generated in this study were capable of differentiation to NPC, and subsequent self-renewal

for up to 30 passages, with morphological similarities to neuroepithelial cells, as has been

previously reported (20). The generation of sustainable populations of neural progenitors

from SCA7 patients is significant, as these cells may act as a starting point for the

generation of numerous disease-affected neuronal subtypes, which can be expanded for

large-scale experiments at relatively low cost. Both patient- and control- derived NPCs

appeared capable of differentiating into neurons expressing βIII-Tubulin and GABA, with

comparable efficiencies. This corresponds with results from previous studies of

neurodegenerative conditions, including other polyQ diseases (25, 26, 33), which found no

difference in differentiation potential between patient and control iPSCs. In addition, no

obvious difference in the ability to generate GABA-positive processes could be detected

between SCA7 and control iPSC-derived neurons.

Electrophysiological studies were carried out to establish whether there were any

functional differences in the intrinsic properties of patient and control cells. The majority of

cells recorded were capable of generating spiking activity including single and multiple

action potentials, an indication of neuronal maturity [39]. Despite the presence of significant

differences in spiking responses between the four cell lines, we did not observe a reliable

trend between the control and patient derived neurons. We do acknowledge a major

limitation in these findings being significant variability across cell lines. This is likely due to

unappreciated differences in culturing conditions between cell lines, which may have

masked our ability to detect a reliable difference in spiking responses caused by the mutant

ATXN7. Alternatively, it may be that mutant ATXN7 does not affect the spiking properties of

neurons at early stages of development, suggesting that an extrinsic stressor of some kind

might be required to elicit a pathological phenotype. Indeed, similar functional analysis of

neurons derived from patients with SCA3, a related polyQ-repeat disorder, showed no

difference between control and patient cells until neurons were excited via bath application

of glutamate [33].

Whilst we acknowledge that it is not possible to make conclusive claims due to the

differences between all the cell lines used, we did observe an apparent difference in resting

membrane potential and input resistance between control and SCA7 patient derived

neurons. Patient cells had more negative resting membrane potentials and increased input

resistance compared to control cells. A more hyperpolarised membrane potential is usually

associated with neuronal maturity whist the high input resistance is usually associated with

neuronal immaturity. These conflicting findings will need to be further explored in future

studies to provide a more conclusive understanding as to how SCA7 specific changes

cause changes in neuronal maturation or drive alternative functional alterations. We can,

however, postulate that the more hyperpolarised membrane potential in the patient cells

may be indicative of a reduction in the excitability of neurons containing mutant ATXN7 in

response to synaptic input. Reductions in Purkinje cell excitability have previously been

observed in animal models of SCA1 and SCA2 [40, 41] and could represent a common

functional endpoint in several polyQ disorders [42]. Future work will involve exploring the

underlying mechanisms that might explain these differences in resting membrane potential

and input resistance. Importantly, it will be necessary to repeat this physiological functional

analysis on iPSC-derived neurons of the Purkinje cell lineage in order to fully recapitulate

the cerebellar-specific elements of the disease using the recently published protocols [43,

44].

Retinal differentiation of iPSCs yielded a heterogeneous cell population after 30 days,

containing a large proportion of cells expressing the photoreceptor markers CRX and

RCVRN, as well as retinal cell genes *PAX6, OTX2, CRX* and *NRL* in the differentiating

cells. The cells expressed varying levels of the photoreceptor genes, *RCVRN* and *RHO*,

confirming that the retinal differentiation protocol was capable of producing cells expressing

markers of "mature" photoreceptors, albeit at relatively low levels of efficiency.

A PCR-based assay was used to confirm the size of the *ATXN7* CAG repeat alleles in

cDNA from patient- and control-derived fibroblasts, iPSCs, neural and retinal cells. Besides

serving as a "fingerprinting" assay, confirming the identity of the lines, these results offered

a semi-quantitative analysis of the expression levels of mutant and wildtype *ATXN7* alleles,

confirming that both alleles were expressed in all cell lines. Despite the inherent instability

of the CAG repeat within the *ATXN7* gene [45], automated fluorescent genotyping of the

PCR products indicated that no contractions or large-scale expansions had occurred during

either reprogramming or differentiation, consistent with previous reports of other iPSC-

derived models of polyQ disorders, including SCA3 and HD [32, 33].

Immunocytochemical analysis of ATXN7 expression in SCA7 patient neural and retinal cells

showed diffuse expression within the nucleus (neurons) or nucleus and cytoplasm (retinal

photoreceptors).   No obvious aggregates were observed in either the iPSCs or the

differentiated cells (Fig 1E, F and Fig 4A). This strongly suggests that the differentiated

cells represent a population of cells at an early stage of development, rather than

recapitulating the age or disease stage of the patient from which the primary cells were

derived.   Previous studies employing similar models for the study of neurodegenerative

disease have raised concerns regarding the relevance of modelling adolescent- and adult-

onset diseases over the short lifespan of cultured neurons [46, 47]. These findings suggest

that pathological hallmarks of disease such as the formation of aggregates may take

decades to manifest, requiring the gradual accumulation of toxic proteins as a result of age-

dependent deficiencies in protein homeostasis.   Although some studies suggest that

aggregates may be detected at earlier stages, the major determinants of aggregate

formation remain the length of the polyQ expansion, and the levels of expression of the

polyQ-containing protein [48, 49]. Thus, a cell line derived from an individual expressing

endogenous levels of a moderately expanded ATXN7 protein may be less likely to

demonstrate an observable cellular phenotype. Alternatively, the aggregation of mutant

protein may require prolonged periods in culture, or an exogenous trigger, such as

exposure to oxidative stress or neurotoxins, or excitation-induced calcium influx [33].

The role of transcriptional dysregulation in the polyQ diseases has been extensively

documented, particularly in cell and animal models of SCA7 [5, 6, 30, 31, 50, 51]. The

identification of gene expression changes, which precede the onset of symptoms, suggests

strongly that alterations in transcription may be among the earliest manifestations of

disease [52]. Thus, it follows that gene expression changes may be used as a tool to

identify a disease-associated phenotype in cells representing early stages of development

[53].

In order to investigate gene expression changes in the SCA7 iPSCs and iPSC-derived

neurons generated here, a panel of candidate transcripts were selected, in which robust

changes had been previously demonstrated [4, 6, 31, 54-56]. The iPSC-derived NPCs and

retinal photoreceptors displayed changes in expression of these key transcripts, suggesting

that these cells may serve as useful models of neurodegenerative disease progression and

for the testing of potential therapies (Fig 1D and Fig 4H). It should be noted that the low cell

numbers and heterogeneity of mature neuronal cultures proved a challenge to obtaining

significant biological material to perform reliable expression analysis. Hence, the decision

was made to perform this analysis on the relatively homogeneous NPC cultures, with the

view that these cells may yield insights into the early cellular dysfunction underlying SCA7

pathology.

Of the three genes consistently downregulated across both differentiated cell types, two

(*ATXN7* and *KAT2A*) encode components of the STAGA transcriptional co-activator

complex. Previous studies in SCA7 patient fibroblasts and mouse models have

demonstrated a disease-associated increase in *ATXN7* expression, mediated by non-

coding RNAs [54, 57]. The contradictory decrease in *ATXN7* expression in SCA7 NPCs

and photoreceptors observed here could reflect the early developmental stage of the cells,

but further analysis of the regulatory pathways will be required in order to elucidate the

basis for this apparent decrease in the disease-causing protein in affected cell types.

KAT2A encodes the histone acetyltransferase GCN5. Although there are no established

links between *ATXN7* and the expression of *KAT2A*, numerous studies have identified a

functional interaction between the two proteins, which results in changes in STAGA activity

in *in vitro* and *in vivo* models of SCA7 [4, 58, 59]. Loss of GCN5 expression has been

shown to result in increased retinal degeneration in SCA7 mice [58].

The interaction between ATXN7 and CRX has been hypothesised to be a key factor behind

the development of retinal degeneration in SCA7 patients [51]. Therefore, the expression of

multiple known CRX targets, which were previously shown to be down-regulated in SCA7

mice, were included in the gene expression experiments. None of these target genes

(including *ARR3*, *GNAT1* or *RHO*) showed consistent changes in patient cells. However,

transcriptional changes in the expression of additional retinal genes, including *OTX2*

(involved in the determination of photoreceptor cell fate), *RCVRN* (expressed in

photoreceptors), and *RPE65* (expressed in retinal pigment epithelial cells), were noted in

the patient derived cells.   A significant degree of heterogeneity was observed in the

differentiated retinal cells, both in terms of morphology, and gene/protein expression (Fig 4),

therefore additional investigation will be required on additional iPSC cell lines to determine

whether these differences can be attributed to experimental differences or pathogenic

mechanisms.

Downregulation of the HSP genes *HSP70* and *DNAJA1* was observed in SCA7 patient

NPCs. A decrease in levels of these two *HSP*s has been previously reported in both SCA7

mice, and human patient lymphoblasts [6, 31]. Although this decrease in expression was

hypothesised in mice to represent an advanced stage of disease progression, the early

developmental stage recapitulated by our model suggests that decreases in certain HSP

genes may instead be an inherent defect, which could predispose certain populations of

cells to degeneration.

The generation of patient-specific, disease-relevant cell types is particularly important in

neurodegenerative diseases; as such cells provide a unique model in which to evaluate

disease pathogenesis without the complications associated with transgene overexpression

in cell or animal models. In addition, the use of cells containing the patient's own genetic

background offers the opportunity to investigate potential modifiers of disease onset and

progression [60, 61]. Perhaps most importantly to the South African context, iPSC-derived

neurons provide the first opportunity to evaluate the efficacy of the allele-specific RNAi-

based therapy developed by Scholefield et al [16], in disease-affected cells.

One significant caveat of this study remains the small number of patients assessed - a

consequence of the rare nature of the condition - and the challenges associated with patient

recruitment in a developing world setting, in which many of those affected are unable to

access tertiary healthcare. Whilst, in this study, comparisons were not made between

isogenic gene-edited lines, they do retain significantly similar background genetics, as they

are all generated from the same immediate family, minimising (to a degree) the extensive

differences between unrelated individuals. Future studies will focus on recruitment, in order

to extend these investigations in a larger patient cohort. To control for the inherent genetic

variability associated with comparisons between unrelated patients, future work will also

focus on the generation of isogenic control lines, by means of CRISPR/Cas9-mediated

genome editing. Dastidar et al [62] have previously demonstrate this approach. It remains,

however, technically challenging to target the disease-causing repeat in ataxin-7 and retain

the endogenous regulatory landscape. Due to the sequence homology of the wild -type

allele, such a strategy would likely render the wild-type protein non-functional via a

frameshift.

Nevertheless, the SCA7 iPSCs generated here serve as a resource for differentiation into a

variety of disease-associated cell types, providing an ideal model in which to study

neurodegenerative diseases. The results of this study provide evidence of a disease

phenotype in iPSC-derived cells from the South African SCA7 patient cohort. We hope that

our data will contribute to the ongoing efforts to use iPSC cells to study the pathogenesis of

neurodegenerative disorders, which are needed in development of population-specific

therapies.

577 Acknowledgements

578   The authors thank Dr Mahito Nakanishi for kind provision of the Sendai virus vector, Ms

579   Jane Vowles for assistance with iPSC generation, and Ms Theresa Ruppelt and colleagues

References

1.   Orr HT, Zoghbi HY. Trinucleotide repeat disorders. Annu Rev Neurosci.

2007;30:575-621. doi: 10.1146/annurev.neuro.29.051605.113042.

2.   Gouw L, Digre K, Harris C, Haines J, Ptacek L. Autosomal dominant cerebellar

ataxia with retinal degeneration Clinical, neuropathologic, and genetic analysis of a large

kindred. Neurology. 1994;44(8):1441. doi: 10.1212/WNL.44.8.1441.

3.   Garden GA, La Spada AR. Molecular pathogenesis and cellular pathology of

spinocerebellar ataxia type 7 neurodegeneration. Cerebellum. 2008;7(2):138-49. doi:

10.1007/s12311-008-0027-y.

4.   Palhan VB, Chen S, Peng G-H, Tjernberg A, Gamper AM, Fan Y, et al.

Polyglutamine-expanded ataxin-7 inhibits STAGA histone acetyltransferase activity to

produce retinal degeneration. Proc Natl Acad Sci USA. 2005;102(24):8472-7. doi:

10.1073/pnas.0503505102.

5.   Abou-Sleymane G, Chalmel F, Helmlinger D, Lardenois A, Thibault C, Weber C, et

al. Polyglutamine expansion causes neurodegeneration by altering the neuronal

differentiation program. Hum Mol Genet. 2006;15(5):691-703. doi: 10.1093/hmg/ddi483.

6.   Chou A-H, Chen C-Y, Chen S-Y, Chen W-J, Chen Y-L, Weng Y-S, et al.

Polyglutamine-expanded ataxin-7 causes cerebellar dysfunction by inducing transcriptional

dysregulation. Neurochem Int. 2010;56(2):329-39. doi: 10.1016/j.neuint.2009.11.003.

7.   Ström A-L, Forsgren L, Holmberg M. A role for both wild-type and expanded ataxin-7

in   transcriptional   regulation.   Neurobiol   Dis.   2005;20(3):646-55.   doi:

10.1016/j.nbd.2005.04.018.

8.   Ward JM, Stoyas CA, Switonski PM, Ichou F, Fan W, Collins B, et al. Metabolic and

organelle morphology defects in mice and human patients define spinocerebellar ataxia

type   7   as   a   mitochondrial   disease.   Cell   Rep.   2019;26(5):1189-202.   doi:

10.1016/j.celrep.2019.01.028.

9.   Stoyas CA, Bushart DD, Switonski PM, Ward JM, Alaghatta A, Tang M-b, et al.

Nicotinamide pathway-dependent SIRT1 activation restores calcium homeostasis to

achieve neuroprotection in spinocerebellar ataxia type 7. Neuron. 2020;105(4):630-44. doi:

10.1016/j.neuron.2019.11.019.

10.   Fiszer A, Wroblewska JP, Nowak BM, Krzyzosiak WJ. Mutant CAG repeats

effectively targeted by RNA interference in SCA7 cells. Genes. 2016;7(12):132. doi:

10.3390/genes7120132.

11.   Niewiadomska-Cimicka A, Trottier Y. Molecular targets and therapeutic strategies in

spinocerebellar ataxia type 7. Neurotherapeutics. 2019:1-23. doi: 10.1007/s13311-019-

00778-5.

12.   Naphade S, Tshilenge K-T, Ellerby LM. Modeling Polyglutamine Expansion Diseases

with Induced Pluripotent Stem Cells. Neurotherapeutics. 2019:1-20. doi: 10.1007/s13311-

019-00810-8.

13.   Smith D, Bryer A, Watson L, Greenberg L. Inherited polyglutamine spinocerebellar

ataxias in South Africa. S Afr Med J. 2012;102(8):683-6. doi: 10.7196/samj.5521.

14.   Greenberg J, Solomon G, Vorster A, Heckmann J, Bryer A. Origin of the SCA7 gene

mutation in South Africa: implications for molecular diagnostics. Clin Genet. 2006;70(5):415-

7. doi: 10.1111/j.1399-0004.2006.00680.x.

15.   Smith DC, Atadzhanov M, Mwaba M, Greenberg LJ. Evidence for a common founder

effect amongst South African and Zambian individuals with Spinocerebellar ataxia type 7. J

Neurol Sci. 2015;354(1):75-8. doi: 10.1016/j.jns.2015.04.053.

16.   Scholefield J, Greenberg LJ, Weinberg MS, Arbuthnot PB, Abdelgany A, Wood MJ.

Design of RNAi hairpins for mutation-specific silencing of ataxin-7 and correction of a SCA7

phenotype. PloS ONE. 2009;4(9):e7232. doi: 10.1371/journal.pone.0007232.

17.   Scholefield J, Watson L, Smith D, Greenberg J, Wood MJ. Allele-specific silencing of

mutant Ataxin-7 in SCA7 patient-derived fibroblasts. Eur J Hum Genet. 2014;22(12):1369-

75. doi: 10.1038/ejhg.2014.39.

18.   Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic

and   adult   fibroblast   cultures   by   defined   factors.   Cell.   2006;126(4):663-76.   doi:

10.1016/j.cell.2006.07.024.

19.   Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction

of pluripotent stem cells from adult human fibroblasts by defined factors. Cell.

2007;131(5):861-72. doi: 10.1016/j.cell.2007.11.019.

20.   Freshney R. Specialized Cells. In: Capes-Davis A, editor. Culture of Animal Cells: A

Manual of Basic Technique and Specialized Applications. 6th ed. Hoboken, New Jersey,

United States: Wiley-Blackwell; 2000. p. 383-432.

21.   Nishimura K, Sano M, Ohtaka M, Furuta B, Umemura Y, Nakajima Y, et al.

Development of defective and persistent Sendai virus vector a unique gene

delivery/expression system ideal for cell reprogramming. J Biol Chem. 2011;286(6):4760-

71. doi: 10.1074/jbc.M110.183780.

22.   Nakanishi M, Otsu M. Development of Sendai virus vectors and their potential

applications in gene therapy and regenerative medicine. Curr Gene Ther. 2012;12(5):410-6.

doi: 10.2174/156652312802762518.

23.   Schwartz P, Brick D, Nethercott H, Stover A. Traditional Human Embryonic Stem Cell

Culture. Methods Mol Biol. 2011;767:107-23. doi: 10.1007/978-1-61779-201-4_8.

24.   Martí M, Mulero L, Pardo C, Morera C, Carrió M, Laricchia-Robbio L, et al.

Characterization of pluripotent stem cells. Nat Protoc. 2013;8(2):223. doi:

10.1038/nprot.2012.154.

25.   Li W, Sun W, Zhang Y, Wei W, Ambasudhan R, Xia P, et al. Rapid induction and

long-term self-renewal of primitive neural precursors from human embryonic stem cells by

small molecule inhibitors. Proc Natl Acad Sci USA. 2011;108(20):8299-304. doi:

10.1073/pnas.1014041108.

26.   Boucherie C, Mukherjee S, Henckaerts E, Thrasher AJ, Sowden JC, Ali RR. Brief

report: Self-organizing neuroepithelium from human pluripotent stem cells facilitates

derivation of photoreceptors. Stem Cells. 2013;31(2):408-14. doi: doi: 10.1002/stem.1268.

27.   Larionov A, Krause A, Miller W. A standard curve based method for relative real time

PCR data processing. BMC Bioinformatics. 2005;6(1):62. doi: 10.1186/1471-2105-6-62.

28.   Taylor J, Grote SK, Xia J, Vandelft M, Graczyk J, Ellerby LM, et al. Ataxin-7 can

export from the nucleus via a conserved exportin-dependent signal. J Biol Chem.

2006;281(5):2730-9. doi: 10.1074/jbc.M506751200.

29.   Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE

guidelines: minimum information for publication of quantitative real-time PCR experiments.

Clin Chem. 2009;55(4):611-22. doi: 10.1373/clinchem.2008.112797.

30.   Yoo S-Y, Pennesi ME, Weeber EJ, Xu B, Atkinson R, Chen S, et al. SCA7 knockin

mice model human SCA7 and reveal gradual accumulation of mutant ataxin-7 in neurons

and abnormalities in short-term plasticity. Neuron. 2003;37(3):383-401. doi: 10.1016/s0896-

6273(02)01190-x.

31.   Tsai H-F, Lin SJ, Li C, Hsieh M. Decreased expression of Hsp27 and Hsp70 in

transformed lymphoblastoid cells from patients with spinocerebellar ataxia type 7. Biochem

Biophys Res Commun. 2005;334(4):1279-86. doi: 10.1016/j.bbrc.2005.06.207.

32.   Camnasio S, Carri AD, Lombardo A, Grad I, Mariotti C, Castucci A, et al. The first

reported generation of several induced pluripotent stem cell lines from homozygous and

heterozygous Huntington's disease patients demonstrates mutation related enhanced

lysosomal activity. Neurobiol Dis. 2012;46(1):41-51. doi: 10.1016/j.nbd.2011.12.042.

33.   Koch P, Breuer P, Peitz M, Jungverdorben J, Kesavan J, Poppe D, et al. Excitation-

induced ataxin-3 aggregation in neurons from patients with Machado–Joseph disease.

Nature. 2011;480(7378):543. doi: doi: 10.1038/nature10671.

34.   Moore AR, Filipovic R, Mo Z, Rasband MN, Zecevic N, Antic SD. Electrical

excitability of early neurons in the human cerebral cortex during the second trimester of

gestation. Cereb Cortex. 2008;19(8):1795-805. doi: 10.1093/cercor/bhn206.

35.   Perrier AL, Tabar V, Barberi T, Rubio ME, Bruses J, Topf N, et al. Derivation of

midbrain dopamine neurons from human embryonic stem cells. Proc Natl Acad Sci USA.

2004;101(34):12543-8. doi: 10.1073/pnas.0404700101.

36.   Vazin T, Becker KG, Chen J, Spivak CE, Lupica CR, Zhang Y, et al. A novel

combination of factors, termed SPIE, which promotes dopaminergic neuron differentiation

from   human   embryonic   stem   cells.   PloS   ONE.   2009;4(8):e6606.   doi:

10.1371/journal.pone.0006606.

37.   Belinsky GS, Moore AR, Short SM, Rich MT, Antic SD. Physiological properties of

neurons derived from human embryonic stem cells using a dibutyryl cyclic AMP-based

protocol. Stem Cells Dev. 2011;20(10):1733-46. doi: 10.1089/scd.2010.0501.

38. Belinsky GS, Rich MT, Sirois CL, Short SM, Pedrosa E, Lachman HM, et al. Patch-

clamp recordings and calcium imaging followed by single-cell PCR reveal the

developmental profile of 13 genes in iPSC-derived human neurons. Stem Cell Res.

2014;12(1):101-18. doi: 10.1016/j.scr.2013.09.014.

39. Moody WJ, Bosma MM. Ion channel development, spontaneous activity, and activity-

dependent development in nerve and muscle cells. Physiol Rev. 2005;85(3):883-941. doi:

10.1152/physrev.00017.2004.

40. Duvick L, Barnes J, Ebner B, Agrawal S, Andresen M, Lim J, et al. SCA1-like disease

in mice expressing wild-type ataxin-1 with a serine to aspartic acid replacement at residue

776. Neuron. 2010;67(6):929-35. doi: 10.1016/j.neuron.2010.08.022.

41. Hansen ST, Meera P, Otis TS, Pulst SM. Changes in Purkinje cell firing and gene

expression precede behavioral pathology in a mouse model of SCA2. Hum Mol Genet.

2012;22(2):271-83. doi: 10.1093/hmg/dds427.

42. Chopra R, Shakkottai VG. The role for alterations in neuronal activity in the

pathogenesis of polyglutamine repeat disorders. Neurotherapeutics. 2014;11(4):751-63. doi:

10.1007/s13311-014-0289-7.

43. Muguruma K. Self-organized cerebellar tissue from human pluripotent stem cells and

disease modeling with patient-derived iPSCs. Cerebellum. 2018;17(1):37-41. doi:

10.1007/s12311-017-0905-2.

44.   Watson LM, Wong MM, Vowles J, Cowley SA, Becker EB. A simplified method for

generating Purkinje cells from human-induced pluripotent stem cells. Cerebellum.

2018;17(4):419-27. doi: 10.1007/s12311-017-0913-2.

45.   Monckton DG, Cayuela ML, Gould FK, Brock GJ, De Silva R, Ashizawa T. Very large

(CAG) n DNA repeat expansions in the sperm of two spinocerebellar ataxia type 7 males.

Hum Mol Genet. 1999;8(13):2473-8. doi: 10.1093/hmg/8.13.2473.

46.   Soldner F, Hockemeyer D, Beard C, Gao Q, Bell GW, Cook EG, et al. Parkinson's

disease patient-derived induced pluripotent stem cells free of viral reprogramming factors.

Cell. 2009;136(5):964-77. doi: 10.1016/j.cell.2009.02.013.

47.   Hartl FU, Bracher A, Hayer-Hartl M. Molecular chaperones in protein folding and

proteostasis. Nature. 2011;475(7356):324. doi: 10.1038/nature10317.

48.   Miller J, Arrasate M, Shaby BA, Mitra S, Masliah E, Finkbeiner S. Quantitative

relationships between huntingtin levels, polyglutamine length, inclusion body formation, and

neuronal death provide novel insight into huntington's disease molecular pathogenesis. J

Neurosci. 2010;30(31):10541-50. doi: 10.1523/JNEUROSCI.0146-10.2010.

49.   Arrasate M, Mitra S, Schweitzer ES, Segal MR, Finkbeiner S. Inclusion body

formation reduces levels of mutant huntingtin and the risk of neuronal death. Nature.

2004;431(7010):805. doi: 10.1038/nature02998.

50.   Zijlstra M, Rujano M, Van Waarde M, Vis E, Brunt E, Kampinga H. Levels of DNAJB

family members (HSP40) correlate with disease onset in patients with spinocerebellar

ataxia type 3. Eur J Neurosci. 2010;32(5):760-70. doi: 10.1111/j.1460-9568.2010.07352.x.

51.   La Spada AR, Fu Y-H, Sopher BL, Libby RT, Wang X, Li LY, et al. Polyglutamine-

expanded ataxin-7 antagonizes CRX function and induces cone-rod dystrophy in a mouse

model of SCA7. Neuron. 2001;31(6):913-27. doi: 10.1016/s0896-6273(01)00422-6.

52.   Helmlinger D, Tora L, Devys D. Transcriptional alterations and chromatin remodeling

in   polyglutamine   diseases.   Trends   Genet.   2006;22(10):562-70.   doi:

10.1016/j.tig.2006.07.010.

53.   Feyeux M, Bourgois-Rocha F, Redfern A, Giles P, Lefort N, Aubert S, et al. Early

transcriptional changes linked to naturally occurring Huntington's disease mutations in

neural derivatives of human embryonic stem cells. Hum Mol Genet. 2012;21(17):3883-95.

doi: 10.1093/hmg/dds216.

54.   Sopher BL, Ladd PD, Pineda VV, Libby RT, Sunkin SM, Hurley JB, et al. CTCF

regulates ataxin-7 expression through promotion of a convergently transcribed, antisense

noncoding RNA. Neuron. 2011;70(6):1071-84. doi: 10.1016/j.neuron.2011.05.027.

55.   Wang H-L, Chou A-H, Lin A-C, Chen S-Y, Weng Y-H, Yeh T-H. Polyglutamine-

expanded ataxin-7 upregulates Bax expression by activating p53 in cerebellar and inferior

olivary neurons. Exp Neurol. 2010;224(2):486-94. doi: 10.1016/j.expneurol.2010.05.011.

56.   Luthi-Carter R, Strand AD, Hanson SA, Kooperberg C, Schilling G, La Spada AR, et

al. Polyglutamine and transcription: gene expression changes shared by DRPLA and

Huntington's disease mouse models reveal context-independent effects. Hum Mol Genet.

2002;11(17):1927-37. doi: 10.1093/hmg/11.17.1927.

57.   Tan JY, Vance KW, Varela MA, Sirey T, Watson LM, Curtis HJ, et al. Cross-talking

noncoding RNAs contribute to cell-specific neurodegeneration in SCA7. Nat Struct Mol Biol.

2014;21(11):955. doi: 10.1038/nsmb.2902.

58.   Chen YC, Gatchel JR, Lewis RW, Mao C-A, Grant PA, Zoghbi HY, et al. Gcn5 loss-

of-function accelerates cerebellar and retinal degeneration in a SCA7 mouse model. Hum

Mol Genet 2011;21(2):394-405. doi: 10.1093/hmg/ddr474.

59.   Helmlinger D, Hardy S, Eberlin A, Devys D, Tora L. Both normal and polyglutamine-

expanded ataxin-7 are components of TFTC-type GCN5 histone acetyltransferase-

containing complexes. Biochem Soc Symp. 2006;73:155-63. doi: 10.1042/bss0730155.

60.   Bilen J, Bonini NM. Genome-wide screen for modifiers of ataxin-3 neurodegeneration

in Drosophila. PLoS Genet. 2007;3(10):e177. doi: 10.1371/journal.pgen.0030177.

61.   Marsh JL, Thompson LM. Drosophila in the study of neurodegenerative disease.

Neuron. 2006;52(1):169-78. doi: 10.1016/j.neuron.2006.09.025.

62.   Dastidar S, Ardui S, Singh K, Majumdar D, Nair N, Fu Y, et al. Efficient

CRISPR/Cas9-mediated editing of trinucleotide repeat expansion in myotonic dystrophy

patient-derived iPS and myogenic cells. Nucleic acids research. 2018;46(16):8275-98.

Supporting information

S1 Fig: Genogram of SCA7 and control patients. Genograms illustrating two families

from which the SCA7 patient samples were taken from. A related control, sibling of 41Q,

was used. From the three patients (two SCA7 and one control), 4 iPSC clones where

derived (2 SCA7 lines and 2 Control lines).

S2 Fig: Representative karyogram from a iPSC line. Karyogram from iPSC clone P1

showing no gross abnormalities.

S3 Fig: iPSC characterisation. (A) Representative immunocytochemistry image showing

positive OCT4 staining (red, top panel) and TRA-1-60 staining (green, bottom panel) in

iPSCs clone 47Q. (B), Immunocytochemistry in newly infected fibroblasts (top panel) and

iPSC colonies (bottom panel) co-stained with primary antibodies against OCT4 (red) and

the viral nucleocapsid protein (green) showed effective silencing of the reprogramming

Sendai virus. DAPI staining shown in blue. Expression of pluripotency markers in iPSC

lines, determined by qRT-PCR. All five iPSC lines (47Q, 41Q, CON_A, CON_B) expressed

OCT4 (C), SOX2 (D) and NANOG (E), compared to low expression levels in the original

donor fibroblasts (F), or cells subjected to the retinal differentiation protocol for 10 days

(D10, pooled data from lines 41Q and CON_B). All levels shown relative to beta (β)-actin.

S4 Fig: iPSCs differentiation. iPSCs were differentiated to three germ layers *in vitro and*

validated by staining with appropriate markers: mesoderm using smooth muscle actin

(SMA), ectoderm using β-III-tubulin, and endoderm using alpa-fetoprotein (AFP). Nuclei are

counterstained with DAPI (blue). Images are representative of iPSCs from the 47Q patient

clone.

S5 Fig: Quantitative PCR results from SCA7 derived fibroblasts. There were no

significant transcriptional changes in SCA7 patient-derived fibroblasts compared to

unaffected control fibroblasts.

S6 Fig: Gel from semi-quantitative PCR of iPSC-derived NPCs and retinal cells. Gel

electrophoresis after qRT-PCR on NPCs (A) and retinal cells (B**)**. Samples from all three

individuals showed a single band at approximately 355bp, corresponding with a wild-type

allele, whilst patient cell lines 47Q and 41Q showed an additional larger band corresponding

to a mutant allele. CAG repeat length in NPCs was evaluated at varying passages

(indicated by p13, p15 or p17).

S Table 1: Primary antibodies

S Table 2: Secondary antibodies

S Table 3: Primer sequences

Data access

All data presented in this manuscript can be accessed here:

https://www.dropbox.com/sh/2t5aydvclgd9tp3/AABIEZPqWIULiCyoaym-i4iaa?dl=0

A Passive **B** Abortive spike **C** Single spike **D** Recurrent spikes

10μm

E

F

	Family 1		Family 2
Sample name	41Q	CON	47Q
CAG alleles	10/41	10/10	10/47
Age at biopsy	44	33	50
Age of SCA onset	44	N/A	37
No. of iPSC clones	1	2	1
Clone names	41Q	CON$_A$ CON$_B$	47Q