Full title: Molecular and electrophysiological features of spinocerebellar ataxia type seven in

induced pluripotent stem cells

Short title: Spinocerebellar ataxia type seven and induced pluripotent stem cells

RJ Burman[1,2,3¶*], LM Watson[4,5¶], , DC Smith[4,5¶], JV Raimondo[1,2,5], R Ballo[2], J Scholefield[6],

SA Cowley[7], MJA Wood[8], SH Kidson[1,5], LJ Greenberg[4,5]

[¶]These authors contributed equally to this work

[1]Division of Cell Biology, Department of Human Biology, Faculty of Health Sciences,

University of Cape Town, Cape Town, South Africa

[2]Neuroscience Institute, Faculty of Health Sciences, University of Cape Town, Cape Town,

South Africa

[3]Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United

Kingdom

[4]Division of Human Genetics, Department of Pathology, Faculty of Health Sciences,

University of Cape Town, Cape Town, South Africa

[5]Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences,

University of Cape Town, Cape Town, South Africa

[6]Gene Expression & Biophysics Group, Synthetic Biology ERA, CSIR Biosciences, Pretoria,

South Africa

[7]Sir William Dunn School of Pathology, University of Oxford, Oxford, United Kingdom

[8]Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United

Kingdom

[*]Corresponding author:

richard.burman@ndcn.ox.ac.uk (RJB)

Abstract

Spinocerebellar ataxia type 7 (SCA7) is an inherited neurodegenerative disease caused by a polyglutamine repeat expansion in the *ATXN7* gene. Patients with this disease suffer from a degeneration of their cerebellar Purkinje neurons and retinal photoreceptors that result in a progressive ataxia and loss of vision. As with many neurodegenerative diseases, studies of pathogenesis have been hindered by a lack of disease-relevant models. To this end, we have generated induced pluripotent stem cells (iPSCs) from a cohort of SCA7 patients in South Africa. First, we differentiated the SCA7 affected iPSCs into neurons which showed evidence of a transcriptional phenotype affecting components of STAGA (*ATXN7* and *KAT2A*) and the heat shock protein pathway (*DNAJA1* and *HSP70*). We then performed electrophysiology on the SCA7 iPSC-derived neurons and found that these cells show features of functional immaturity. Lastly, we were able to differentiate the SCA7 iPSC cells into retinal photoreceptors that also showed similar transcriptional aberrations to the SCA7 neurons. Our findings demonstrate that iPSC-derived neurons and photoreceptors from SCA7 patients express molecular and electrophysiological differences that are indicative of impaired neuronal health. We hope that these findings will contribute towards the ongoing efforts to establish the cell-derived models of neurodegenerative diseases that are needed to develop patient-specific treatments.

Introduction

Spinocerebellar ataxia type 7 (SCA7) is an inherited neurodegenerative disease caused by a CAG repeat expansion in the *ATXN7* gene. Since the translation of this CAG repeat leads to an expanded polyglutamine (polyQ) tract within the resultant protein, SCA7 is classified as a polyQ–repeat disorder. Other diseases with a similar pathophysiology include five different SCAs (SCA 1, 2, 3, 6 and 17), as well as Huntington disease, dentatorubral-pallidoluysian atrophy and spinal bulbar muscular atrophy (1). Clinically, SCA7 patients present with ataxia, dysarthria and visual loss. This is caused by a selective degeneration of cerebellar Purkinje neurons and retinal photoreceptors (2). Symptoms progressively worsen over a period of 10 to 30 years, leading ultimately to brainstem dysfunction, blindness, physical disability and death.

The mechanism by which a polyQ expansion within the ubiquitously expressed ATXN7 protein leads to the selective degeneration of Purkinje neurons and photoreceptors remains to be fully elucidated. ATXN7 is known to be a component of the mammalian transcription co-activator complex, STAGA (SPT3-TAF9-ADA-GCN5 acetyltransferase) (3). This protein has been shown to facilitate the interaction between STAGA and the cone-rod homeobox (CRX) transactivator of photoreceptor genes, linking the function of ATXN7 with the retinal phenotype observed in SCA7 patients (4). In addition, several studies have highlighted the role of transcriptional aberrations in the neuronal cell dysfunction that precedes the onset of disease symptoms (3-7). These gene expression changes may arise either directly from alterations in transcriptional regulation by mutant ATXN7, or indirectly, as a consequence of a generalised cellular response to the presence of mutant ATXN7. More recently, mitochondrial dysfunction has been proposed as an additional pathogenic mechanism (8).

As with many neurodegenerative conditions, research into the molecular pathogenesis of

SCA7 has been hindered by a lack of suitable models of human disease progression. This

is particularly relevant in cases where the genomic context of the mutation may have an

impact on gene function and might prove useful for therapeutic development. SCA7 occurs

at an unusually high frequency in the South African population as a result of a founder effect

in patients of Black African ethnic origin (9). South African SCA7 patients also display a

unique phenomenon – a single nucleotide polymorphism (SNP) within *ATXN7* (rs3774729),

which is linked to the mutation in all patients studied to date (10). Approximately 43% of

these individuals are heterozygous for the polymorphism, allowing for allelic discrimination,

and providing an ideal target for developing an allele-specific silencing therapy. This

haplotype has been shown to extend into other Southern African populations, suggesting

that such a therapy may be more widely applicable than was first thought (11). We have

previously demonstrated the efficacy of an allele-specific RNAi treatment in an over-

expression cell model of SCA7 (12), as well as in SCA7 patient fibroblasts (13). Disease-

relevant cell lines generated from these patients are thus of vital importance in the

understanding of disease pathogenesis and the development of therapies, as they carry the

patient's full genomic sequence, including SNPs which may be used as targets for gene

silencing.

The generation of iPSCs involves reprogramming somatic cells to a pluripotent state by

means of viral transduction with the pluripotency genes *OCT4*, *SOX2*, *KLF4* and *c-MYC*

(14, 15). Importantly, iPSCs can then be differentiated into any tissue of the body through

treatment with specific growth factors. This makes iPSCs a useful starting point for the

generation of disease-relevant cell models, particularly in neurodegenerative diseases,

where primary CNS cultures may only be obtained using invasive methods.

In this study, we have generated and characterised induced pluripotent stem cell (iPSC)

lines from two South African SCA7 patients and an unaffected, related control. We were

then able to differentiate these iPSCs into cells expressing markers associated with retinal

photoreceptors, neural progenitors cells (NPCs) and neurons. Thereafter, we have obtained

preliminary evidence for a disease phenotype in these cells, in the form of pathogenically

relevant gene expression changes and alterations in intrinsic neuronal properties. Our

findings indicate that these cells may be able to replicate, to some extent, the pathogenesis

underlying SCA7.

Material and methods

Ethics approval and patient recruitment

Ethics approval for the study was granted by the University of Cape Town (UCT) Faculty of

Health Sciences Human Research Ethics Committee (HREC REF. 380/2009 and

434/2011), and was renewed annually, incorporating amendments to the project protocol

where necessary. All methods were carried out in accordance with the guidelines approved

by the Ethics Committee. Participants were recruited from the Neurogenetics clinic at

Groote Schuur Hospital in Cape Town. Informed consent was obtained from all participants

prior to their enrolment in the study.

Establishment of primary fibroblast cultures

108   Primary fibroblast cultures were established from punch skin biopsies taken from the inner

109   forearm of two unrelated SCA7 patients (P1 and P2) and an unaffected control individual

110   (C1, sibling of P2) who had consented to participate in the study, as previously described

111   (16). CAG genotypes and ages at diagnosis and biopsy are shown in S1 Fig.

Generation and characterisation of patient-derived iPSCs

Dermal fibroblasts were reprogrammed into iPSCs using Sendai virus vectors (SeVdp)

containing *OCT4, SOX2, KLF4* and *c-MYC* as previously described (17, 18). One week

after infection, the cells were transferred to mitomycin C-inactivated feeder cells in stem cell

medium (KODMEM, KOSR, NEAA, glutamax, bFGF). Half of the medium was refreshed

every day and colonies were expanded by manual passaging on feeder layers. After 10

passages the expression of pluripotency markers OCT4 and TRA-1-60, as well as silencing

of the reprogramming Sendai virus, were confirmed by immunocytochemistry (antibodies

listed in S1-2 Tables). The expression of selected pluripotency genes (*OCT4, SOX2,*

NANOG) was determined by quantitative PCR. Genomic integrity was assessed by means

of karyotype analysis (G-banding, S2 Fig). To confirm the iPSC lines' capacity to

differentiate into the three embryonic germ layers, *in vitro* differentiation via embryoid body

formation was performed as previously described (19).

Following Sendai virus-mediated reprogramming, iPSC colonies with the correct

morphology (flat, with distinct borders, containing tightly packed cells with a high nucleus-to-

cytoplasm ratio) appeared within three to four weeks. These colonies were manually picked

and clonally expanded in separate dishes on a feeder layer of inactivated mouse embryonic

fibroblasts. Two SCA7 patient iPSC lines and two control lines (representing two affected

individuals and a single related, unaffected control) were successfully generated and

characterised. Immunocytochemical analysis of SCA7 patient and control iPSC lines

revealed iPSC colonies with distinct nuclear staining for the pluripotency transcription factor,

OCT4, compared to the surrounding mouse embryonic feeder fibroblasts (S3 Fig, A – top

panel). The iPSC colonies also stained positive for the embryonic stem cell surface marker

TRA-1-60 (S3 Fig, A – bottom panel).

Co-staining of all iPSC lines with primary antibodies against OCT4 and the nucleocapsid

protein of the virus (anti-NP) showed little or no evidence of NP staining in the OCT4-

positive pluripotent cells (S3 Fig, B –bottom panel), compared to intense cytoplasmic

staining in newly infected fibroblasts (S3 Fig, B – top panel). This indicated that the Sendai

virus had been effectively silenced, and that the iPSCs had achieved self-regulating

pluripotency. All lines were assessed after passage 8. The expression of pluripotency

markers was further confirmed by qPCR. All five iPSC lines expressed high levels of *OCT4,*

SOX2 and *NANOG*, standard markers of pluripotency (S3 Fig, C - E), compared to donor

fibroblasts or cells that had been subjected to retinal or neuronal differentiation. The

expression levels of *SOX2* and *NANOG* were similar across the five iPSC lines, but lines

P2b and C1b showed lower levels of *OCT4* expression compared to the remaining three

lines (although still significantly higher than differentiated fibroblasts and retinal cells).

Neural differentiation and characterisation

Differentiation of iPSCs into neural precursors was performed by treatment of iPSCs with 3µM glycogen synthase kinase 3 (GSK3) inhibitor (CHIR99021) and 2µM TGFβ inhibitor (SB431542) as previously described (20). For neuronal differentiation, neural precursors were seeded at a density of 150 000 cells/well onto a Matrigel-coated six-well plate in neural induction medium (20). After two days, medium was changed to neuronal differentiation medium (DMEM/FBS supplemented with N2, B27, 300ng/ml cyclic AMP (Sigma), 0.2mM ascorbic acid (Sigma), 10ng/ml BDNF (Peprotech) and 10ng/ml GDNF (Peprotech)) and the cells maintained in culture for 14 to 21 days. Medium was changed every 2-3 days. Characterisation was performed by immunocytochemistry and qPCR (for antibodies and primers see S1-3 Tables).

Electrophysiology

Neurons cultured between 2-3 weeks on glass cover slips (uncoated) were removed from the incubator and rapidly transported to the recording chamber of a Zeiss Axioskop Upright Microscope (Zeiss). Electrophysiological recordings were made in neuronal differentiation medium at room temperature and were restricted to the first 5 hours following cell removal from the incubator environment. Patch pipettes of 13-20 MOhm tip resistance were pulled from filamental borosilicate glass capillaries (2.00 mm outer diameter, 1.58 mm inner diameter, Hilgenberg), using a horizontal puller (Model P-1000, Sutter). The pipettes were filled with an internal solution containing (in mM): K-gluconate (126); KCl (4); Na 2 ATP (4); NaGTP (0.3); Na 2 -phosphocreatinine (10) and HEPES (10). Osmolarity was adjusted to between 290 and 300 mOsM and the pH was adjusted to between 7.38 and 7.42 with KOH.

Cells were visualised using a 40x water-immersion objective (Zeiss). Digital images were

obtained using a CCD camera (VX55, TILL Photonics). Individual cells were selected for

recordings based on a small round or ovoid cell body (diameters, 5–10 µm) and typically

two or more extended processes. Recordings were made in current-clamp and voltage-

clamp mode using an Axopatch 200B amplifier (Molecular Devices). Data acquisition was

performed through an ITC-1600 board (Instrutech) connected to a PC running a custom-

written routine (PulseQ) under IGOR Pro (Wavemetrics). Analysis was performed using

custom-written scripts in MATLAB (Mathworks). Statistical analysis was performed using

GraphPad Prism (v8, GraphPad Software). Normally distributed data is present with mean

and standard error of the mean (SEM) whilst non-normally distributed data was presented

with median and interquartile range (IQR). Appropriate parametric and non-parametric tests

were performed including unpaired Students t-test, Mann-Whitney U test and Kruskal-

Wallis. Significance was defined as $p < 0.05$.

184 Retinal differentiation and characterisation

Differentiation into retinal photoreceptors was performed as previously described

(Boucherie et al., 2013), using iPSCs cultured in feeder-free conditions on Matrigel in

mTESR^{TM} medium. The cells were dissociated enzymatically and plated onto Matrigel-

covered dishes in neural differentiation medium containing N2 and B27 supplements (Life

Technologies). After settling for an hour, adhered cells were covered in a 2% Matrigel

solution. The following day the medium was replaced with neural differentiation medium

without Matrigel, and cells were fed with fresh medium every second day. From day 10 the

medium was supplemented with 3nM recombinant SHH (R&D Systems), 50ng/µl acidic

fibroblast growth factor (aFGF) (R&D Systems), 10ng/µl basic fibroblast growth factor

(bFGF) (Miltenyi), 1mM taurine and 500nM retinoic acid (both Sigma Aldrich).

Characterisation was performed by immunocytochemistry and qPCR (for antibodies and

primers, see S Tables 1 and 2).

DNA and RNA isolation, cDNA synthesis

198 DNA extraction from cultured cells was performed using the QIAGEN DNeasy Blood and

199 Tissue Kit. RNA was isolated from cultured cells using the QIAGEN RNeasy Plus Mini Kit.

200 Synthesis of cDNA was performed using the Applied Biosystems High Capacity cDNA

201 Reverse Transcription Kit (Life Technologies) using 500ng-1µg template RNA.

Real-time quantitative PCR

Real-time quantitative PCR (qPCR) was performed on the BioRad CFX96 Real-Time PCR

System, using the Power SYBR Green PCR Master Mix (Applied Biosystems), according to

manufacturer's instructions. Primers (S3 Table) were obtained from PrimerDesign Ltd,

Integrated DNA Technologies (IDT), or Sigma-Aldrich. Relative quantities of target mRNA

were determined using the relative standard curve method (Larionov et al., 2005). Standard

curves were prepared for each primer pair, from serial dilutions of pooled sample cDNA.

Universal cycling conditions were used (95°C for 10min, followed by 40 cycles of 95°C for

15 seconds and 60°C for 1min). PCRs were performed in technical triplicate, on at least

two biological replicates, and results were analysed using the BioRad CFX Manager

software (v3.1). Gene of interest expression was normalised to *beta actin* (*ACTB*)

expression in each case. The design and reporting of qPCR experiments aimed to comply

with the Minimum Information for publication of Quantitative real-time PCR Experiments

guidelines (21, 22) wherever possible. Statistical analysis was performed using the unpaired

Students' t-test. Significance was defined as $p < 0.05$.

217 CAG repeat length determination

The length of the disease-causing CAG repeat in *ATXN7* was determined from DNA by

means of polymerase chain reaction (PCR) and automated fluorescent genotyping. The

PCR reaction mix consisted of 0.4µM each, forward and reverse primer (S3 Table), 0.6units

of GoTaq DNA polymerase (Promega), 100ng of DNA, and Failsafe buffer J (Epicentre

Biotechnologies) at a final concentration of 1X, made up to a final reaction volume of 10µl.

Cycling conditions were as follows: 95°C for 5min; followed by 30 cycles of 95°C for 30

seconds, 53°C for 6 seconds and 72°C for 40 seconds; and a final elongation step at 72°C

for 7min. Automated fluorescent genotyping was performed using the ABI 3130*xl* Genetic

Analyzer (Applied Biosystems).

The length of the CAG repeat [$(CAG)_n$] was approximated using the following equation,

adapted from (Dorschner et al., 2002): $n = (0.3063$ x Length of major PCR product in base

pairs) $- 76.475$bp. The major PCR product was defined as the product generating the

highest fluorescent peak, as detected using the ABI 3130*xl* Genetic Analyzer.

231 Results

232 SCA7 iPSC-derived neural precursor cells show transcriptional

233 disease-associated aberrations

SCA7 patient and control iPSCs generated neural precursor cells (NPCs) at comparable

efficiencies, when cultured in neural induction medium supplemented with SB431542 (TGFβ

inhibitor) and CHIR99021 (GSK3 inhibitor), with 100% of the cells expressing the early neural

marker Nestin, after five passages (Fig 1A). The cells also stained positive for the disease-causing

protein, ATXN7 (Fig 1B), and demonstrated repression of the pluripotency gene *OCT4*, and

upregulation of the early neural gene *PAX6* (Fig 1C).

Fig 1: SCA7 iPSC-derived NPCs show transcriptional aberrations. (A) Representative

images of control (top panel) and SCA7 patient (bottom panel) NPCs, expressing the early

neural marker, Nestin (green). (B) High magnification images showing nuclear expression of

ATXN7 (red) in control (top) and patient (bottom) NPCs (Green: βIII-Tubulin). (C) qPCR

results indicating suppression of *OCT4* (top panel) and upregulation of *PAX6* (bottom panel)

expression in NPCs, compared with the iPSC lines from which they were derived. All levels

shown relative to beta (β)-actin. (D) Expression of 9 selected genes in SCA7 patient NPCs

($n = 2$) shown relative to unaffected control NPCs ($n = 2$). All levels shown relative to beta

actin and unaffected control cells. (E) Control (top panel) and SCA7 NPC-derived neurons

(bottom panel) stain positive for βIII-Tubulin (green) and ATXN7 (red) after 14 days of

differentiation. (F) Both control and SCA7 patient NPCs produced a small proportion of

GABA-positive neurons (red) after 14 days in culture (Green, βIII-Tubulin). *$p < 0.05$.

To determine whether any transcriptional differences could be detected between SCA7

patient- and unaffected control-derived cell types, a panel of candidate genes was selected,

which had previously been shown to be dysregulated in the retinal and cerebellar tissue of

SCA7 mouse models and patient lymphoblasts (5, 6, 23, 24). The panel included the

following genes: *ATXN7*, brain expressed, X-linked 1 (*BEX1*), DnaJ (Hsp40) homolog,

subfamily A, member 1 (*DNAJA1*), glutamate receptor, ionotropic, AMPA 2 (*GRIA2*), heat

shock protein 27 (*HSP27*), heat shock protein 70 (*HSP70*), heat shock protein 105

(*HSP105*), K (lysine) acetyltransferase 2A (*KAT2A*), Phospholipase C, Beta 3 (*PLCB3*) and

ubiquitin carboxyl-terminal esterase L1 (*UCHL1*).   The expression of these genes was

determined by means of qPCR in fibroblasts (S4 Fig) and NPCs (Fig 1D).

In contrast to the fibroblasts and undifferentiated iPSCs, two genes were downregulated in

SCA7 patient iPSC-derived NPCs when compared to the unaffected control lines. These

included the disease-causing gene *ATXN7* (p = 0.018 in NPCs); and *KAT2A*, encoding

GCN5, the histone acetyltransferase (HAT) component of the STAGA transcription

coactivator complex (p = 0.003 in NPCs).   In addition, NPC-specific alterations in

expression were identified in the heat shock protein genes DNAJA1 (p = 0.04) and HSP70

(p = 0.04), both of which were found to be downregulated in SCA7-patient derived cells.

Lastly, BEX1, an interactor of the p75 neurotrophin receptor, which regulates neurotrophin

signalling and neuronal differentiation, was found to be downregulated in the SCA7 NPCs (p

= 0.03).

In order to confirm the size of the *ATXN7* CAG repeat alleles in mRNA from patient- and

control-derived fibroblasts, iPSCs and NPCs an RT-PCR-based assay was performed. The

results were visualised on an agarose gel (S Fig 5A), and confirmed by automated

fluorescent genotyping. The length of the CAG repeat did not appear to fluctuate during

reprogramming or differentiation, corresponding to previous reports in similar cell lines (25,

26). This assay also confirmed that both the mutant and wild-type *ATXN7* alleles were

expressed by all cell types, and that there were no obvious differences in allele expression

in affected or unaffected cells.

After culture for 14 days in neuronal differentiation medium containing N2/B27 supplement,

cAMP ascorbic acid, BDNF and GDNF (20), both SCA7 patient and control NPCs stained

positive for the neuronal marker βIII-Tubulin, and showed robust, diffuse nuclear localisation

of the disease-causing protein, ATXN7 (Fig 2E-F). A subset of cells (approximately 1.8%)

stained positive for gamma-aminobutyric acid (GABA), although the proportion of

GABAergic neurons did not appear to vary between SCA7 patients and controls (Fig 2G-H).

No obvious differences in morphology were observed when comparing neurons derived

from SCA7 patient iPSCs with those derived from controls. In order to test functional

differences between SCA7 neurons and controls, however, we then sought to obtain

electrophysiological measurements from these cells.

SCA7 iPSC-derived neurons demonstrate functional differences that may affect cell excitability

Cells were assayed for physiological properties between 14 and 23 days post induction of

neuronal differentiation. Cells were targeted for whole-cell recordings based on their

morphological properties. This included a small round or ovoid cell body with diameters

between 5–10 µm and typically two or more extended processes. Following the attainment

of a whole-cell patch, current pulses of between 0 and 10 pA were applied. Individual cells

displayed four general types of spiking responses: a purely passive (Fig 3A), abortive spike

(Fig 3B), single spike (Fig 3C) and recurrent spiking response (Fig 3D). These spiking

properties are similar to those observed in acute human fetal brain slices (27), hESC-

derived neurons (28-30) and iPSC-derived neurons (31). A postmitotic neuron matures by

inserting voltage-gated channels into its plasma membrane (32). Therefore, the spiking

response of a cell to current injection can be used to determine the maturation stage of a

differentiating neuron: passive (least mature) → abortive spike → single spike → recurrent

spikes (most mature).

Fig 2: Patch-clamp analyses of iPSC derived neuronal cultures. Cells could be divided

into four categories based on their spiking responses. (A-D) Differential interference

contrast images of cells targeted for patch-clamp recordings (top). Whole-cell recordings in

current clamp mode (middle) depict spiking patterns following injection of current (bottom).

Cells fell into (A) 'passive', (B) 'abortive spike', (C) 'single spike' and (D) 'recurrent spikes'

categories. (E) The fraction of cells which fell into each category for cells derived from two

control and two SCA7 patient iPSC lines. Note that although the cells derived from various

iPSC lines exhibited different distributions of spiking responses, no trend between control

and patient lines was observed. (F) The resting membrane potentials (Vm) of cells varied

significantly across each cell line. (G) The Vm derived from SCA7 patient lines was

significantly more hyperpolarised compared to the control cell lines. Error bars denote mean

+/- SEM in (H) and (I), and median and IQR in (F)**,** (G)**,** (J) and (K). * $p < 0.05$, ** $p < 0.01$,

*** $p < 0.001$, **** $p < 0.0001$.

Spiking responses were collected in current-clamp mode from cells derived from four

separate iPSC lines: two control lines C1a ($n = 70$) and C1b ($n = 42$), and two patient lines

P1 ($n = 44$) and P2b ($n = 72$). Although the fraction of cells which fell into each spiking

response category was significantly dependent on the iPSC line from which the cells were

derived ($p < 0.0001$, *Chi-squared test*), no trend could be discerned between control and

patient lines (Fig 2E). For example, the control line C1a had the most mature phenotype

with the highest fraction of cells in the single spike and recurrent spiking categories whilst

P1 demonstrated a relatively immature phenotype, with the majority of cells displaying

spiking responses falling into the passive category. However, the other lines did not

corroborate this difference as the control line C1b displayed a less mature phenotype than

patient line P2b.

Next, we compared the resting membrane potential (Vm) of cells derived from each cell line.

This parameter was significantly dependent on the iPSC line from which the cells were

derived (Fig 2F, p = 0.002, *ANOVA*). The mean resting membrane potential +/- SEM for the

C1a, C1b, P1a and P2b cell lines were -57.0 +/- 1.7, -54.7 +/- 3.2, -67.3 +/- 2.7 and -62.5

+/- 2.5 mV respectively.   The control lines had significantly more depolarised resting

membrane potentials as compared to SCA7 patient derived cell lines (Fig 2G, p = 0.0003,

unpaired t-test).

Next, we performed voltage-clamp recordings of the neurons in order to measure the input

resistance as well as the voltage-gated sodium and potassium currents (Fig 3A-B). A lower

input resistance is associated with neurite outgrowth and increased numbers of ion

channels inserted into the plasma membrane during the process of neuronal maturation.

Once again, a cell's input resistance was significantly dependent on the cell line to which it

belonged (Fig 3C, p = 0.005, *Kruskal-Wallis*). Input resistance +/- SEM was 4987 +/- 421,

5484 +/- 243, 5979 +/- 513 and 7094 +/- 455 mΩ for the C1a, C1b, P1a and P2b cell lines

respectively. Cells derived from SCA7 patient lines had a significantly higher mean input

resistance than cells from the control lines (Fig 3D, p = 0.01, *Mann-Whitney U test*).

Fig 3: Sodium and potassium currents in neurons differentiated from control and

patient iPS cell lines. (A) Voltage-clamp recording trace. Membrane currents (top) were

recorded following 10 mV voltage steps between -90 and 30 mV (bottom).   Maximum

potassium current (I_K) indicated. Dashed gray rectangle indicates the position of voltage-

gated sodium currents (enhanced in 'B'). Maximum sodium current (I_{Na}) is indicated 'right'.

(B) Zoom-in of rectangle in '**A**' with the maximum I_{Na} highlighted. (C) Cell input resistance

was also significantly varied across cell lines, however the input resistance of the SCA7

patient-derived cells was significantly increased in SCA7 patient lines as compared to

control lines (D). (E) Population data from the four iPSC lines for maximum voltage-gated I_K

current. (F) The I_K was significantly smaller in the cells derived from SCA7 patients. (G) The

I_{Na} was significantly varied across cell lines. (H) The I_{Na} was significantly smaller in the

SCA7 patient-derived cells. Error bars denote median and IQR. * $p < 0.05$, *** $p < 0.001$,

**** $p < 0.0001$.

As one would predict, the spiking properties of neurons are directly correlated with the size

of their currents. The maximum size of voltage-gated sodium currents and potassium

currents measured in each cell (Max. I_K and Max. I_{NA}) was significantly dependent on the

particular iPSC line concerned (Fig 3E and G, $p < 0.0001$ in both cases, *ANOVA*). The

mean Max. I_K +/- SEM was 180.1 +/- 16.5, 176 +/- 10.8, 222.5 +/- 19.3 and 105.7 +/- 8.0 pA

for the C1a, C1b, P1a and P2b cell lines respectively. The mean Max. I_{NA} +/- SEM was -

166.5 +/- 18.6, -277.1 +/- 23.0, -268.0 +/- 30.5 and -108.0 +/- 14.2 pA for the C1a, C1b, P1a

and P2b cell lines. When pooled together, however, we noted that both the Max I_K and Max

I_{Na} were significantly smaller in the cell lines coming from SCA7 patients compared to the

control cell lines (**Fig 3F** and **H**, $p = 0.03$ and $p = 0.0005$, *Mann-Whitney Test*).

SCA7 iPSC-derived retinal photoreceptors show transcriptional

aberrations

After characterising the SCA7 iPSC-derived NPCs and neurons, whether we could produce

retinal photoreceptors from the same cell lines. For the retinal differentiation of iPSCs, the

Matrigel "sandwich" system facilitated a rapid self-organisation and differentiation of the

pluripotent stem cells into structures containing cells morphologically indicative of columnar

neuroepithelia. These structures lost their integrity from day 4-5, and cells spread into an

adherent monolayer. The gradual emergence of cells with a neuronal morphology was

observed from day 10 to day 30, particularly in areas of low confluence. Following

differentiation of patient and control cells into retinal cells, immunocytochemical analyses

were performed on cells at the end of the differentiation period (day 30), to determine

whether the cells expressed retinal cell markers. The cells were stained for either the

disease-causing protein ATXN7 (Fig 4A) or the retinal cell markers CRX and RCVRN (Fig

4B-C). No obvious differences in morphology were observed between SCA7 patient and

control-derived cells. The differentiated cells displayed varying expression levels of the

retinal genes *CRX, PAX6, RCVRN* and *OTX2* (Fig 4D-G).

Fig 4: iPSC-derived retinal cells exhibit aberrant transcriptional changes.

Immunocytochemical analysis of SCA7 (top panel) and control (bottom panel) retinal cells

showed positive staining for ATXN7 (A), cone-rod homeobox (CRX, B) and recoverin

(RCVRN, C) on day 30 of the differentiation experiment. Nuclei in blue. qPCR results

confirmed the expression of *CRX* (D), *PAX6* (E), *RCVRN* (F) and *OTX2* (G) in all five lines,

compared with the iPSC lines from which they were derived. All levels shown relative to

beta actin. (H) Expression of 23 selected genes in SCA7 patient iPSC-derived retinal cells

(n = 2) shown relative to unaffected control cells (n = 1). All levels shown relative to beta

actin and unaffected control cells. *$p < 0.05$.

To determine transcriptional changes in the retinal cells, we added retinal cell specific

genes to the gene panel used in the NPCs. The additions to the gene panel include:

arrestin 3 (*ARR3*); cone-rod homeobox (*CRX*); guanine nucleotide binding protein (G

protein); alpha transducing activity polypeptide 1 (*GNAT1*); Microphthalmia-associated

transcription factor (*MITF*); neural retina leucine zipper (*NRL*); orthodenticle homeobox 2

(*OTX2*); paired box 6 (*PAX6*); recoverin (*RCVRN*); rhodopsin (*RHO*) and retinal pigment

epithelium-specific protein 65kDa (*RPE65*). Like the NPC cells, both ATXN7 (p = 0.04) and

KAT2A (p = 0.02) were down regulated in the retinal cells. As with the NPCs, *ATXN7* CAG

repeat alleles were confirmed using an RT-PCR (S Fig 5B). In addition, we also noticed

expression changes in genes that appeared to be specific to the SCA7 photoreceptors. These

included *GRIA2*, encoding the glutamate receptor, ionotropic, AMPA2 (GluR2) (downregulated, p =

0.04); and three retinal-specific genes, *OTX2* (upregulated, p = 0.002), *RCVRN* (downregulated, p =

0.01) and *RPE65* (upregulated, p = 0.001). Lastly, and in contrast to the SCA7 NPCs, BEX1

expression appear to be upregulated in SCA7 photoreceptors (p = 0.0006). The opposing

in BEX1 transcription changes (up in photoreceptors and down in NPCs) may be indicative

of a possible differential response by the different cell types to the presence of mutant

ATXN7.

Discussion

This study describes the generation and characterisation of the first iPSCs from the South

African SCA7 patient cohort, through the transduction of patient dermal fibroblasts with

Sendai viral vectors (17, 18). Furthermore, we show that we are able to differentiate these

iPSCs from SCA7 into neurons and retinal photoreceptors. The SCA7 patient iPSCs

generated in this study were shown to be able to differentiate into a population of NPCs,

capable of self-renewal for up to 30 passages, and with morphological similarities to

neuroepithelial cells, as has been previously reported (20). The generation of sustainable

populations of neural precursors from SCA7 patients is significant, as these cells may act as

a starting point for the generation of numerous disease-affected neuronal subtypes, which

can be expanded for large-scale experiments at relatively low cost. Both patient- and

control- derived NPCs appeared capable of differentiating into neurons expressing βIII-

Tubulin and GABA, with comparable efficiencies. This corresponds with results from

previous studies of neurodegenerative conditions, including other polyQ diseases (25, 26,

33), which found no difference in differentiation potential between patient and control iPSCs.

In addition, no obvious difference in the ability to generate GABA-positive processes could

be detected between SCA7 and control iPSC-derived neurons.

Electrophysiological studies were carried out to establish whether there were any functional

differences in the intrinsic properties of patient and control cells. The majority of cells

recorded were capable of generating spiking activity including single and multiple action

potentials, an indication of neuronal maturity (32). Despite the presence of significant

differences in spiking responses between the four cell lines, we did not observe a reliable

trend between the control and patient derived neurons. However, we did find that the

differences in both the maximum I_K and I_{Na} between SCA7 patient derived cells and

controls. We do acknowledge a major limitation in these experiences being significant

variability across cell lines. This is likely due to unappreciated differences in culturing

conditions between cell lines, which may have masked our ability to detect a reliable

difference in spiking responses caused by the mutant ATXN7. Alternatively, it may be that

mutant ATXN7 does not affect the spiking properties of neurons at early stages of

development, suggesting that an extrinsic stressor of some kind might be required to elicit a

pathological phenotype. Indeed, similar functional analysis of neurons derived from patients

with SCA3, a related polyQ-repeat disorder, showed no difference between control and

patient cells until neurons were excited via bath application of glutamate (26).

Importantly, we observed differences in resting membrane potential and input resistance

between control and SCA7 patient derived neurons. Patient cells had more negative resting

membrane potentials and increased input resistance compared to control cells. The former

would serve to reduce the excitability of neurons containing mutant ATXN7 in response to

synaptic input. Reductions in Purkinje cell excitability have previously been observed in

animal models of SCA1 and SCA2 (34, 35) and could represent a common functional

endpoint in several polyQ disorders (36). Future work will involve exploring the underlying

mechanisms that might explain these differences in resting membrane potential and input

resistance. Importantly, it will be necessary to repeat this physiological functional analysis

on iPSC derived neurons of the Purkinje cell lineage in order to fully recapitulate the

cerebellar-specific elements of the disease using the recently published protocols (37, 38).

Retinal differentiation of iPSCs yielded a heterogeneous cell population after 30 days,

containing a large proportion of cells expressing the photoreceptor markers CRX and

RCVRN, as well as retinal cell genes *PAX6*, *OTX2*, *CRX* and *NRL* in the differentiating

cells. The cells expressed varying levels of the photoreceptor genes, *RCVRN* and *RHO*,

confirming that the retinal differentiation protocol was capable of producing cells expressing

markers of "mature" photoreceptors, albeit at relatively low levels of efficiency.

A PCR-based assay was used to confirm the size of the *ATXN7* CAG repeat alleles in

cDNA from patient- and control-derived fibroblasts, iPSCs, neural and retinal cells. Besides

serving as a "fingerprinting" assay, confirming the identity of the lines, these results offered

a semi-quantitative analysis of the expression levels of mutant and wildtype *ATXN7* alleles,

confirming that both alleles were expressed in all cell lines. Despite the inherent instability

of the CAG repeat within the *ATXN7* gene (39), automated fluorescent genotyping of the

PCR products indicated that no contractions or large-scale expansions had occurred during

either reprogramming or differentiation, consistent with previous reports of other iPSC-

derived models of polyQ disorders, including SCA3 and HD (25, 26).

Immunocytochemical analysis of ATXN7 expression in SCA7 patient neural and retinal cells

showed diffuse expression within the nucleus (neurons) or nucleus and cytoplasm (retinal

photoreceptors). No obvious aggregates were observed in either the iPSCs or the

differentiated cells (Fig 1E and Fig 4A). This strongly suggests that the differentiated cells

represent a population of cells at an early stage of development, rather than recapitulating

the age or disease stage of the patient from which the primary cells were derived. Previous

studies employing similar models for the study of neurodegenerative disease have raised

concerns regarding the relevance of modelling adolescent- and adult-onset diseases over

the short lifespan of cultured neurons (33, 40). These findings suggest that pathological

hallmarks of disease such as the formation of aggregates may take decades to manifest,

requiring the gradual accumulation of toxic proteins as a result of age-dependent

deficiencies in protein homeostasis. Although some studies suggest that aggregates may

be detected at earlier stages, the major determinants of aggregate formation remain the

length of the polyQ expansion, and the levels of expression of the polyQ-containing protein

(41, 42). Thus, a cell line derived from an individual expressing endogenous levels of a

moderately expanded ATXN7 protein may be less likely to demonstrate an observable

cellular phenotype. Alternatively, the aggregation of mutant protein may require prolonged

periods in culture, or an exogenous trigger, such as exposure to oxidative stress or

neurotoxins, or excitation-induced calcium influx (26).

The role of transcriptional dysregulation in the polyQ diseases has been extensively

documented, particularly in cell and animal models of SCA7 (5, 6, 23, 24, 43, 44). The

identification of gene expression changes, which precede the onset of symptoms, suggests

strongly that alterations in transcription may be among the earliest manifestations of

disease (45). Thus, it follows that gene expression changes may be used as a tool to

identify a disease-associated phenotype in cells representing early stages of development

(46).

In order to investigate gene expression changes in the SCA7 iPSCs and iPSC-derived

neurons generated here, a panel of candidate transcripts were selected, in which robust

changes had been previously demonstrated (4, 6, 24, 47-49). The iPSC-derived neurons

and retinal photoreceptors displayed changes in expression of these key transcripts,

suggesting that these cells may serve as useful models of neurodegenerative disease

progression and for the testing of potential therapies (Fig 1E and Fig 4H).

Of the three genes consistently downregulated across both differentiated cell types, two

(*ATXN7* and *KAT2A*) encode components of the STAGA transcriptional co-activator

complex.   Previous studies in SCA7 patient fibroblasts and mouse models have

demonstrated a disease-associated increase in *ATXN7* expression, mediated by non-

coding RNAs (47, 50). The contradictory decrease in *ATXN7* expression in SCA7 NPCs

and photoreceptors observed here could reflect the early developmental stage of the cells,

but further analysis of the regulatory pathways will be required in order to elucidate the

basis for this apparent decrease in the disease-causing protein in affected cell types.

KAT2A encodes the histone acetyltransferase GCN5. Although there are no established

links between *ATXN7* and the expression of *KAT2A*, numerous studies have identified a

functional interaction between the two proteins, which results in changes in STAGA activity

in *in vitro* and *in vivo* models of SCA7 (4, 51, 52). Loss of GCN5 expression has been

shown to result in increased retinal degeneration in SCA7 mice (51).

The interaction between ATXN7 and CRX has been hypothesised to be a key factor behind

the development of retinal degeneration in SCA7 patients (44). Therefore, the expression

of multiple known CRX targets, which were previously shown to be down-regulated in SCA7

mice, were included in the gene expression experiments. None of these target genes

(including *ARR3*, *GNAT1* or *RHO*) showed consistent changes in patient cells. However,

transcriptional changes in the expression of additional retinal genes, including *OTX2*

(involved in the determination of photoreceptor cell fate), *RCVRN* (expressed in

photoreceptors), and *RPE65* (expressed in retinal pigment epithelial cells), were noted in

the patient derived cells. A significant degree of heterogeneity was observed in the

differentiated retinal cells, both in terms of morphology, and gene/protein expression (Fig 4),

therefore additional investigation will be required on additional iPSC cell lines to determine

whether these differences can be attributed to experimental differences or pathogenic

mechanisms.

Downregulation of the HSP genes *HSP70* and *DNAJA1* was observed in SCA7 patient

NPCs. A decrease in levels of these two *HSP*s has been previously reported in both SCA7

mice, and human patient lymphoblasts (6, 24). Although this decrease in expression was

hypothesised in mice to represent an advanced stage of disease progression, the early

developmental stage recapitulated by our model suggests that decreases in certain HSP

genes may instead be an inherent defect, which could predispose certain populations of
cells to degeneration.

The generation of patient-specific, disease-relevant cell types is particularly important in
neurodegenerative diseases; as such cells provide a unique model in which to evaluate
disease pathogenesis without the complications associated with transgene overexpression
in cell or animal models. In addition, the use of cells containing the patient's own genetic
background offers the opportunity to investigate potential modifiers of disease onset and
progression (53, 54). Perhaps most importantly to the South African context, iPSC-derived
neurons provide the first opportunity to evaluate the efficacy of the allele-specific RNAi-
based therapy developed by Scholefield et al (12), in disease-affected cells.

One significant caveat of this study remains the small number of patients assessed - a
consequence of the rare nature of the condition - and the challenges associated with patient
recruitment in a developing world setting, in which many of those affected are unable to
access tertiary healthcare. Future studies will focus on recruitment, in order to extend these
investigations in a larger patient cohort. To control for the inherent genetic variability
associated with comparisons between unrelated patients, future work will also focus on the
generation of isogenic control lines, by means of CRISPR/Cas9-mediated genome editing.

Nevertheless, the SCA7 iPSCs generated here serve as a resource for differentiation into a
variety of disease-associated cell types, providing an ideal model in which to study
neurodegenerative diseases. The results of this study provide evidence of a disease
phenotype in iPSC-derived cells from the South African SCA7 patient cohort. We hope that
our data will contribute to the ongoing efforts to use iPSC cells to study the pathogenesis of

neurodegenerative disorders, which are needed in development of population-specific
therapies.

Acknowledgements

Funding for this work was provided by Ataxia UK, Commonwealth Scholarship Commission (UK),
John Fell OUP Fund, National Research Foundation (South Africa), National Research Foundation
(South Africa, Competitive Programme for Rated Researchers CPR201106240000019696), Medical
Research Council (South Africa), Harry Crossley Foundation, Deutscher Akademischer Austausch
Dienst, University of Cape Town Research Council, Wellcome Trust, Parkinson's Disease UK,
Medical Research Council UK, Blue Brain Project, and the James Martin 21[st] Century School.
Electrophysiology equipment was provided by the École Polytechnique Fédérale de Lausanne,
Switzerland. The authors thank Dr Mahito Nakanishi for kind provision of the Sendai virus vector,
Ms Jane Vowles for assistance with iPSC generation, and Ms Theresa Ruppelt and colleagues at
the South African National Health Laboratory Service for karyotype analyses. We honour the late Ms
Ingrid Baumgarten for her invaluable assistance over many years, particularly in establishing patient
fibroblast cultures.

References

1.   Orr HT, Zoghbi HY. Trinucleotide repeat disorders. Annu Rev Neurosci.
2007;30:575-621.

2.   Gouw L, Digre K, Harris C, Haines J, Ptacek L. Autosomal dominant cerebellar
ataxia with retinal degeneration Clinical, neuropathologic, and genetic analysis of a large
kindred. Neurology. 1994;44(8):1441-.

3.　Garden GA, La Spada AR. Molecular pathogenesis and cellular pathology of

spinocerebellar ataxia type 7 neurodegeneration. The Cerebellum. 2008;7(2):138-49.

4.　Palhan VB, Chen S, Peng G-H, Tjernberg A, Gamper AM, Fan Y, et al.

Polyglutamine-expanded ataxin-7 inhibits STAGA histone acetyltransferase activity to

produce retinal degeneration. Proceedings of the National Academy of Sciences of the

United States of America. 2005;102(24):8472-7.

5.　Abou-Sleymane G, Chalmel F, Helmlinger D, Lardenois A, Thibault C, Weber C, et

al. Polyglutamine expansion causes neurodegeneration by altering the neuronal

differentiation program. Human molecular genetics. 2006;15(5):691-703.

6.　Chou A-H, Chen C-Y, Chen S-Y, Chen W-J, Chen Y-L, Weng Y-S, et al.

Polyglutamine-expanded ataxin-7 causes cerebellar dysfunction by inducing transcriptional

dysregulation. Neurochemistry international. 2010;56(2):329-39.

7.　Ström A-L, Forsgren L, Holmberg M. A role for both wild-type and expanded ataxin-7

in transcriptional regulation. Neurobiology of disease. 2005;20(3):646-55.

8.　Ward JM, Stoyas CA, Switonski PM, Ichou F, Fan W, Collins B, et al. Metabolic and

organelle morphology defects in mice and human patients define spinocerebellar ataxia

type 7 as a mitochondrial disease. Cell reports. 2019;26(5):1189-202. e6.

9.　Smith D, Bryer A, Watson L, Greenberg L. Inherited polyglutamine spinocerebellar

ataxias in South Africa. SAMJ: South African Medical Journal. 2012;102(8):683-6.

10.   Greenberg J, Solomon G, Vorster A, Heckmann J, Bryer A. Origin of the SCA7 gene

mutation in South Africa: implications for molecular diagnostics. Clinical genetics.

2006;70(5):415-7.

11.   Smith DC, Atadzhanov M, Mwaba M, Greenberg LJ. Evidence for a common founder

effect amongst South African and Zambian individuals with Spinocerebellar ataxia type 7.

Journal of the neurological sciences. 2015;354(1):75-8.

12.   Scholefield J, Greenberg LJ, Weinberg MS, Arbuthnot PB, Abdelgany A, Wood MJ.

Design of RNAi hairpins for mutation-specific silencing of ataxin-7 and correction of a SCA7

phenotype. PloS one. 2009;4(9):e7232.

13.   Scholefield J, Watson L, Smith D, Greenberg J, Wood MJ. Allele-specific silencing of

mutant Ataxin-7 in SCA7 patient-derived fibroblasts. European Journal of Human Genetics.

2014;22(12):1369.

14.   Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic

and adult fibroblast cultures by defined factors. cell. 2006;126(4):663-76.

15.   Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction

of pluripotent stem cells from adult human fibroblasts by defined factors. cell.

2007;131(5):861-72.

16.   Freshney RI. Specialized cells. Culture of Animal Cells: A Manual of Basic Technique

and Specialized Applications, Sixth Edition. 2000:383-432.

17.   Nishimura K, Sano M, Ohtaka M, Furuta B, Umemura Y, Nakajima Y, et al.

Development   of   defective   and   persistent   Sendai   virus   vector   a   unique   gene

delivery/expression system ideal for cell reprogramming. Journal of Biological Chemistry.

2011;286(6):4760-71.

18. Nakanishi M, Otsu M. Development of Sendai virus vectors and their potential

applications in gene therapy and regenerative medicine. Current gene therapy.

2012;12(5):410-6.

19. Martí M, Mulero L, Pardo C, Morera C, Carrió M, Laricchia-Robbio L, et al.

Characterization of pluripotent stem cells. Nature protocols. 2013;8(2):223.

20. Li W, Sun W, Zhang Y, Wei W, Ambasudhan R, Xia P, et al. Rapid induction and

long-term self-renewal of primitive neural precursors from human embryonic stem cells by

small molecule inhibitors. Proceedings of the National Academy of Sciences.

2011;108(20):8299-304.

21. Taylor J, Grote SK, Xia J, Vandelft M, Graczyk J, Ellerby LM, et al. Ataxin-7 can

export from the nucleus via a conserved exportin-dependent signal. Journal of Biological

Chemistry. 2006;281(5):2730-9.

22. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE

guidelines: minimum information for publication of quantitative real-time PCR experiments.

Clinical chemistry. 2009;55(4):611-22.

23. Yoo S-Y, Pennesi ME, Weeber EJ, Xu B, Atkinson R, Chen S, et al. SCA7 knockin

mice model human SCA7 and reveal gradual accumulation of mutant ataxin-7 in neurons

and abnormalities in short-term plasticity. Neuron. 2003;37(3):383-401.

24.   Tsai H-F, Lin SJ, Li C, Hsieh M. Decreased expression of Hsp27 and Hsp70 in

transformed lymphoblastoid cells from patients with spinocerebellar ataxia type 7.

Biochemical and biophysical research communications. 2005;334(4):1279-86.

25.   Camnasio S, Carri AD, Lombardo A, Grad I, Mariotti C, Castucci A, et al. The first

reported generation of several induced pluripotent stem cell lines from homozygous and

heterozygous Huntington's disease patients demonstrates mutation related enhanced

lysosomal activity. Neurobiology of disease. 2012;46(1):41-51.

26.   Koch P, Breuer P, Peitz M, Jungverdorben J, Kesavan J, Poppe D, et al. Excitation-

induced ataxin-3 aggregation in neurons from patients with Machado–Joseph disease.

Nature. 2011;480(7378):543.

27.   Moore AR, Filipovic R, Mo Z, Rasband MN, Zecevic N, Antic SD. Electrical

excitability of early neurons in the human cerebral cortex during the second trimester of

gestation. Cerebral Cortex. 2008;19(8):1795-805.

28.   Perrier AL, Tabar V, Barberi T, Rubio ME, Bruses J, Topf N, et al. Derivation of

midbrain dopamine neurons from human embryonic stem cells. Proceedings of the National

Academy of Sciences of the United States of America. 2004;101(34):12543-8.

29.   Vazin T, Becker KG, Chen J, Spivak CE, Lupica CR, Zhang Y, et al. A novel

combination of factors, termed SPIE, which promotes dopaminergic neuron differentiation

from human embryonic stem cells. PloS one. 2009;4(8):e6606.

30.   Belinsky GS, Moore AR, Short SM, Rich MT, Antic SD. Physiological properties of

neurons derived from human embryonic stem cells using a dibutyryl cyclic AMP-based

protocol. Stem cells and development. 2011;20(10):1733-46.

31.   Belinsky GS, Rich MT, Sirois CL, Short SM, Pedrosa E, Lachman HM, et al. Patch-

clamp recordings and calcium imaging followed by single-cell PCR reveal the

developmental profile of 13 genes in iPSC-derived human neurons. Stem cell research.

2014;12(1):101-18.

32.   Moody WJ, Bosma MM. Ion channel development, spontaneous activity, and activity-

dependent development in nerve and muscle cells. Physiological reviews. 2005;85(3):883-

941.

33.   Soldner F, Hockemeyer D, Beard C, Gao Q, Bell GW, Cook EG, et al. Parkinson's

disease patient-derived induced pluripotent stem cells free of viral reprogramming factors.

Cell. 2009;136(5):964-77.

34.   Duvick L, Barnes J, Ebner B, Agrawal S, Andresen M, Lim J, et al. SCA1-like disease

in mice expressing wild-type ataxin-1 with a serine to aspartic acid replacement at residue

776. Neuron. 2010;67(6):929-35.

35.   Hansen ST, Meera P, Otis TS, Pulst SM. Changes in Purkinje cell firing and gene

expression precede behavioral pathology in a mouse model of SCA2. Human molecular

genetics. 2012;22(2):271-83.

36.   Chopra R, Shakkottai VG. The role for alterations in neuronal activity in the

pathogenesis of polyglutamine repeat disorders. Neurotherapeutics. 2014;11(4):751-63.

37.	Muguruma K. Self-organized cerebellar tissue from human pluripotent stem cells and
disease modeling with patient-derived iPSCs. The Cerebellum. 2018;17(1):37-41.

38.	Watson LM, Wong MM, Vowles J, Cowley SA, Becker EB. A simplified method for
generating Purkinje cells from human-induced pluripotent stem cells. The Cerebellum.
2018;17(4):419-27.

39.	Monckton DG, Cayuela ML, Gould FK, Brock GJ, De Silva R, Ashizawa T. Very large
(CAG) n DNA repeat expansions in the sperm of two spinocerebellar ataxia type 7 males.
Human molecular genetics. 1999;8(13):2473-8.

40.	Hartl FU, Bracher A, Hayer-Hartl M. Molecular chaperones in protein folding and
proteostasis. Nature. 2011;475(7356):324.

41.	Miller J, Arrasate M, Shaby BA, Mitra S, Masliah E, Finkbeiner S. Quantitative
relationships between huntingtin levels, polyglutamine length, inclusion body formation, and
neuronal death provide novel insight into huntington's disease molecular pathogenesis.
Journal of Neuroscience. 2010;30(31):10541-50.

42.	Arrasate M, Mitra S, Schweitzer ES, Segal MR, Finkbeiner S. Inclusion body
formation reduces levels of mutant huntingtin and the risk of neuronal death. nature.
2004;431(7010):805.

43.	Zijlstra M, Rujano M, Van Waarde M, Vis E, Brunt E, Kampinga H. Levels of DNAJB
family members (HSP40) correlate with disease onset in patients with spinocerebellar
ataxia type 3. European Journal of Neuroscience. 2010;32(5):760-70.

44.   La Spada AR, Fu Y-H, Sopher BL, Libby RT, Wang X, Li LY, et al. Polyglutamine-

expanded ataxin-7 antagonizes CRX function and induces cone-rod dystrophy in a mouse

model of SCA7. Neuron. 2001;31(6):913-27.

45.   Helmlinger D, Tora L, Devys D. Transcriptional alterations and chromatin remodeling

in polyglutamine diseases. TRENDS in Genetics. 2006;22(10):562-70.

46.   Feyeux M, Bourgois-Rocha F, Redfern A, Giles P, Lefort N, Aubert S, et al. Early

transcriptional changes linked to naturally occurring Huntington's disease mutations in

neural derivatives of human embryonic stem cells. Human molecular genetics.

2012;21(17):3883-95.

47.   Sopher BL, Ladd PD, Pineda VV, Libby RT, Sunkin SM, Hurley JB, et al. CTCF

regulates ataxin-7 expression through promotion of a convergently transcribed, antisense

noncoding RNA. Neuron. 2011;70(6):1071-84.

48.   Wang H-L, Chou A-H, Lin A-C, Chen S-Y, Weng Y-H, Yeh T-H. Polyglutamine-

expanded ataxin-7 upregulates Bax expression by activating p53 in cerebellar and inferior

olivary neurons. Experimental neurology. 2010;224(2):486-94.

49.   Luthi-Carter R, Strand AD, Hanson SA, Kooperberg C, Schilling G, La Spada AR, et

al. Polyglutamine and transcription: gene expression changes shared by DRPLA and

Huntington's disease mouse models reveal context-independent effects. Human molecular

genetics. 2002;11(17):1927-37.

50.   Tan JY, Vance KW, Varela MA, Sirey T, Watson LM, Curtis HJ, et al. Cross-talking

noncoding RNAs contribute to cell-specific neurodegeneration in SCA7. Nature Structural

and Molecular Biology. 2014;21(11):955.

51.   Chen YC, Gatchel JR, Lewis RW, Mao C-A, Grant PA, Zoghbi HY, et al. Gcn5 loss-

of-function accelerates cerebellar and retinal degeneration in a SCA7 mouse model. Human

molecular genetics. 2011;21(2):394-405.

52.   Helmlinger D, Hardy S, Eberlin A, Devys D, Tora L, editors. Both normal and

polyglutamine-expanded ataxin-7 are components of TFTC-type GCN5 histone

acetyltransferase-containing complexes. Biochemical Society Symposia; 2006: London;

Portland on behalf of The Biochemical Society; 1999.

53.   Bilen J, Bonini NM. Genome-wide screen for modifiers of ataxin-3 neurodegeneration

in Drosophila. PLoS genetics. 2007;3(10):e177.

54.   Marsh JL, Thompson LM. Drosophila in the study of neurodegenerative disease.

Neuron. 2006;52(1):169-78.

729 Supporting information

S1 Fig: Genogram of SCA7 and control patients. Genograms illustrating two families

from which the SCA7 patient samples were taken from. A related control, sibling of P2b,

was used. From the three patients (two SCA7 and one control), 4 iPSC clones where

derived (2 SCA7 lines and 2 Control lines).

S2 Fig: Representative karyogram from a iPSC line. Karyogram from iPSC clone P1

showing no gross abnormalities.

S3 Fig: iPSC characterisation. (A) Representative immunocytochemistry image showing

positive OCT4 staining (red, top panel) and TRA-1-60 staining (green, bottom panel) in

iPSCs. (B), Immunocytochemistry in newly infected fibroblasts (top panel) and iPSC

colonies (bottom panel) co-stained with primary antibodies against OCT4 (red) and the viral

nucleocapsid protein (green) showed effective silencing of the reprogramming Sendai virus.

DAPI staining shown in blue. Expression of pluripotency markers in iPSC lines, determined

by qPCR. All five iPSC lines (P1a, P2a, P2b, C1a, C1b) expressed OCT4 (C), SOX2 (D)

and NANOG (E), compared to low expression levels in the original donor fibroblasts (F), or

cells subjected to the retinal differentiation protocol for 10 days (D10, pooled data from lines

P2b and C1b). All levels shown relative to beta (β)-actin.

S4 Fig: Quantitative PCR results from SCA7 derived fibroblasts. There were no

significant transcriptional changes in SCA7 patient-derived fibroblasts compared to

unaffected control fibroblasts.

S5 Fig: Gel from semi-quantitative PCR of iPSC-derived NPCs and retinal cells. Gel

electrophoresis after qPCR on NPCs (A) and retinal cells (B**)**. Samples from all three

individuals showed a single band at approximately 355bp, corresponding with a wild-type

allele, whilst patient cell lines P1 and P2b showed an additional larger band corresponding

to a mutant allele. CAG repeat length in NPCs was evaluated at varying passages

(indicated by p13, p15 or p17).

S Table 1: Primary antibodies

S Table 2: Secondary antibodies

S Table 3: Primer sequences

Data access

All data presented in this manuscript can be accessed here:

https://www.dropbox.com/sh/2t5aydvclgd9tp3/AABIEZPqWIULiCyoaym-i4iaa?dl=0

A Passive   **B** Abortive spike   **C** Single spike   **D** Recurrent spikes

E

Cell fraction (%)

- C1a
- C1b
- P1
- P2b

****

Passive   Abortive   Single   Recurrent

F

Resting membrane potential (mV)

**

C1a   C1b   P1   P2b

G

Resting membrane potential (mV)

C   SCA7

***