1 **Title:** Natural variation in *couch potato* mediates rapid evolution of learning and reproduction in
2 natural populations of *Drosophila melanogaster*
3

4 **Authors:** Emily L. Behrman[1,2,3], Tadeusz J. Kawecki[2*] and Paul Schmidt[1*]
5
Author Affiliations:
1. Department of Biology, University of Pennsylvania, 433 S. University Ave. Philadelphia, PA
19104
2. Department of Ecology and Evolution, University of Lausanne, Biophore, CH-1015
Lausanne, Switzerland
3. Current address: Janelia Research Campus, Howard Hughes Medical Institute, 19700 Helix
Drive Ashburn, VA 20147
* Authors contributed equally
14
15 **Orcid IDs**:
16 Emily Behrman: 0000-0002-2472-9635
17 Tadeusz Kawecki: 0000-0002-9244-1991
18 Paul Schmidt: 0000-0002-8076-6705
19
20

21 **Running head:** Rapid learning evolution
22

23 **Corresponding Author:**
24 Emily L. Behrman
25 Email Address: Bemily@sas.upenn.edu
26 Mailing address: 19700 Helix Drive
27   Ashburn, VA 20147
28 Phone: 571.209.4000

30 **Keywords**
31 Rapid adaptation, learning, fecundity, trade-offs, *Drosophila melanogaster, couch potato*

Abstract

Seasonal oscillations in temperate environments between the different selection regimes

of winter and summer produce cyclical selection pressures that may drive rapid evolution of

diverse traits. We investigated the evolutionary dynamics of learning ability in natural

populations over this rapid seasonal timescale. Associative learning was tested in common

garden-raised *Drosophila melanogaster* collected from a natural population in the spring and fall

over three consecutive years. The spring flies learned consistently better than fall flies, revealing

seasonal evolution of improved learning performance in nature. Fecundity showed the opposite

seasonal pattern, suggesting a trade-off between learning and reproduction, which we confirmed

at the level of individual females. This trade-off is mediated at least in part by natural

polymorphism in the RNA binding protein *couch potato* (*cpo*), with a haplotype favored during

summer showing poorer learning performance and higher fecundity than a haplotype favored

over winter. Thus, seasonal environments can drive rapid cyclical evolution of learning

performance, but the evolutionary dynamics may be driven by pleiotropic effects of alleles

selected for other reasons.

Impact statement

Evolution is traditionally considered to be a very slow, gradual process, but recent studies show

that some organisms evolve rapidly for a variety of traits. However, there is still little known

about the rate at which behaviors evolve in the wild. Complex behaviors may evolve due to their

fitness benefits or because natural selection acts on a trait that is genetically correlated with that

behavior. Learning ability is an important behavior for many aspects of an organism's biology,

but may be costly and use resources that could be otherwise allocated to other important

functions, such as reproduction. We measure learning ability in fruit flies collected from a

natural population as it evolves across seasons. We find that the complex behavior of learning

ability evolves very fast the wild at the cost of reproduction: the spring populations have higher

learning ability but lay fewer eggs compared to the fall. We found that natural variants of the

couch potato help mediate the trade-off between reproduction and learning in natural

populations. This shows how evolution of complex traits can occur due to selection on genes that

affect multiple traits.

Introduction

It is becoming clear that many life history traits can evolve very rapidly over time scales

that were previously assumed to be constant (Kingsolver et al. 2001; Grant and Grant 2002;

Schmidt and Conde 2006; Carroll et al. 2007; Thompson 2013; Behrman et al. 2015, 2018), but

little is known about the rate at which behaviors evolve. Behaviors may be less prone to evolve

rapidly because behavior can be highly plastic and shaped by learning. Learning is important for

many aspects of animal biology including foraging, spatial orientation, predator avoidance,

aggression, social interactions, and sexual behavior (Yurkovic et al. 2006; Dukas 2008; Stensmyr

et al. 2012; Mansourian et al. 2016). Learning decouples behavioral phenotypes from genotypes

by allowing individuals to develop adaptive behavioral responses to changing environments.

Under a narrow set of circumstances learning may accelerate genetically-based evolutionary

change of behavior (Mayr 1974; Mery and Kawecki 2004b; Paenke et al. 2007), but under most

circumstances it is predicted to slow the rate at which behaviors evolve (Dukas 2004; Paenke et

al. 2007).

The ability to learn is itself a product of evolution, but we know next to nothing about

how rapidly its characteristics evolve in nature. Examples of genetic differences in learning

performance have been reported between closely related species (Odling-Smee et al. 2008;

Hoedjes and Smid 2014) and between conspecific populations (Croston et al. 2015; Froissart et

al. 2017), but the timescale over which these differences have evolved is unknown. Rapid

evolution of improved learning performance in laboratory selection experiments in rats, bees,

blowflies and *Drosophila* indicates that natural populations have copious standing genetic

variation for learning ability (Tryon 1940; Mcguire and Hirschth 1977; Brandes et al. 1988;

Mery and Kawecki 2002; Zwoinska et al. 2017). However, in those experiments learning was

directly under selection, whereas selection on learning in nature is indirect and mediated by the

fitness consequences of behaviors it modifies. An environmental change that results in an

increased need for learning in one behavior (e.g., foraging) might have no effect, or even

decrease, its benefits in a different context (e.g., predator avoidance), dampening changes in net

selection on learning.

The evolutionary dynamics of learning ability may be driven by fitness consequences of

the behavioral modifications it causes or by selection on genetically correlated traits. Correlated

selection due to trade-offs may arise because resources that might be invested in growth,

reproduction, maintenance or defense are diverted to develop and maintain the energetically

costly neural tissue required for learning and memory (Johnston 1982; Mery and Kawecki 2005;

Dukas 2008). The hypothesized reproductive cost of learning (Johnston 1982) is supported by

genetic trade-offs between learning and reproduction in the cabbage white butterfly, *Pieris*

rapae, (Snell-Rood et al. 2011) and an operant reproductive cost of learning in *Drosophila* (Mery

and Kawecki 2004). Thus, rapid evolutionary change of learning ability could be triggered by

environmental changes that alter selection on either learning ability or on correlated traits such as

fecundity.

Multivoltine species are a promising system to study rapid evolution because each

generation may experience significantly different conditions across seasonal time. The

alternating conditions between different selection regimes of winter and summer drive rapid

cyclical evolution of *Drosophila melanogaster* life history, stress resistance and immunity traits

with a hypothesized reproductive trade-off (Schmidt and Conde 2006; Behrman et al. 2015,

2018). Here we investigate if learning performance evolves rapidly over an annual timescale in a

natural population of *D. melanogaster* and if reproductive costs of learning affect the dynamics

in the wild. We predicted learning ability of wild *Drosophila* to vary with season as the

behaviors that involve learning change over seasonal time. Increased ability to learn may be

favored in the summer as many behavioral tasks associated with increased summer fitness are

modulated by learning in *Drosophila* laboratory studies: foraging, pathogen avoidance, and

sexual and social interactions (Dukas 1998, 2005; Sarin and Dukas 2009; Battesti et al. 2012;

Zrelec et al. 2013; Hollis and Kawecki 2014). However, it is possible that learning may be

involved behaviors that are important for overwintering survival (e.g., shelter identification,

pathogen avoidance). Alternatively, if selection acts on correlated traits, then seasonal variation

in learning ability could be driven by these indirect effects – particularly on fecundity in the

summer (Mery and Kawecki 2004) and stress tolerance in the winter (Mery and Kawecki 2005).

We assessed aversive and appetitive learning in common-garden-raised flies derived from

wild spring and fall field collections over three consecutive years. We found that learning

performance was consistently higher in the post-winter collections in the spring compared to the

post-summer collections in the fall. An inverse relationship between learning ability and

fecundity between seasonal collections and at an individual level indicated that a reproductive

cost of learning was involved in the seasonal dynamics. This pointed to a potential role of *couch*

potato (*cpo*), a pleiotropic RNA-binding protein highly expressed in the nervous system

including the mushroom body (Bellen et al. 1992a,b). Fecundity and reproductive diapause

differences in natural *cpo* variants correlate with differential *cpo* expression; the frequency of the

variants in wild *Drosophila* changes with latitude and season (Behrman et al. n.d.; Schmidt et al.

2008; Kolaczkowski et al. 2011; Fabian et al. 2012; Cogni et al. 2013; Bergland et al. 2014). We

found that learning performance and fecundity in flies carrying natural *cpo* haplotypes that are

more common in the spring versus the fall paralleled the seasonal pattern in the natural

population. These results suggest that rapid fluctuating evolution of learning ability in wild

Drosophila is at least in part driven by pleiotropic effects of *cpo* polymorphism, which mediates

a trade-off between reproduction and traits that promote overwinter survival.

Methods

***Drosophila* samples**

Seasonal populations derived from nature

Seasonal differences in learning were assessed by comparing outbred seasonal

populations reconstructed from isofemale lines, hereafter referred to as spring and fall

populations. Gravid females aspirated off of decaying fruit in the spring (June) and fall

(November) at Linvilla Orchards in Media, PA (39.9ºN, -75.4ºE), across three consecutive years

(2012-2014) were used to establish isofemale lines that were maintained in standard laboratory

conditions (25ºC, 12L:12D) on a four-week transfer cycle. After all collections were complete,

representative populations from each collection were re-constructed using 40 isofemale lines per

collection and were maintained in common garden culture for more than 10 non-overlapping

generations. We infer that differences in learning ability among the populations tested in the

standard laboratory environment are due to genetic differences among the populations.

cpo recombinant outbred populations

Three SNPs in *cpo* were used as markers for the temperate haplotype (cpo^{TTA}) that is

more common at high latitudes and in the spring versus the tropical haplotype (cpo^{CGT}) that is

more common at low latitudes and in the fall. (Behrman et al. n.d.): two intronic SNPs

(3R:13790130 and 3R:13791280) and one putative non-synonymous coding change

(3R:13793588, *D. melanogaster* reference genome v.5.39, (Cogni et al. 2013; Bergland et al.

2014). The non-synonymous coding change is validated using in situ hybridization (Bellen et al.

1992a) but the exon is not included in MODENCODE (Celniker et al. 2009). Learning

performance of the haplotypes was assessed using recombinant outbred populations (ROPs;

(Paaby et al. 2014; Behrman et al. 2018, n.d.)) that were each fixed for one haplotype with a

randomized genetic background. ROPs for cpo^{TTA} and cpo^{CGT} were constructed using nine

independent, homozygous *Drosophila* Genetic References Panel (MacKay et al. 2012) lines. Ten

gravid females from each line were permitted to lay eggs for 48h; after at least 10 non-

overlapping generations of recombination with the other liens containing the same haplotype

among the offspring, each ROP was fixed for either cpo^{TTA} or cpo^{CGT} in a heterogeneous, outbred

background.

cpo knockdown

We tested if *cpo* regulation changes learning ability because the *cpo* ROPs have different

levels of whole-body *cpo* expression (Behrman et al. n.d.) and other traits are correlated with

differential *cpo* expression (Schmidt et al. 2008). The UAS/GAL4 system was used to express

dsRNA for RNA interference (RNAi) to knockdown *cpo* expression. Two biological replicate

UAS constructs were created using different insertion sites paired with their respective insertion

site controls located on the second and third chromosomes, respectively, from the Transgenic

RNAi Project (TRiP): (BDSC-60388 with attP40 control BDSC-36304) and (BDSC-28360 with

attP2 control BDSC-36303). The hypothesis that lower *cpo* expression increases learning was

tested using lines that drive GAL4 expression in the mushroom, ellipsoid and fan shaped bodies,

subesophageal ganglion, antennal & optic lobes, protocerebrum & median bundle (BDSC-

30818). Three *cpo*-specific steroid-activated Gal4 geneswitch drivers were used knock down *cpo*

expression in different neuron combinations of adult flies: trachea-associated cells and subsets of

ventral nerve cord and sensory neurons (BDSC-40315), a subset of sensory neurons (BDSC-

40319) and sense organ support cells and subsets of ventral nerve cord, brain and sensory

neurons (BDSC-40334). BDSC-38461 was used as a negative control to express GAL4 in flight

muscles under control of the Actin 88F promotor. All crosses were made with the female

containing the GAL4 driver and male containing the UAS construct.

Learning

Aversive shock learning in the laboratory

Flies were conditioned to associate one of two odorants with an aversive mechanical

shock in an aversive olfactory learning assay (Mery and Kawecki 2005). Flies were reared in

common laboratory conditions (25˚C, 12L:12D) at standardized density of 100 eggs per vial,

sorted into groups by sex under light CO_2 anesthesia 24h prior to the learning assessment, and

then assayed at 3-5 days of age. Single-sex groups of 30 flies were conditioned to associate

either methylcyclohexanol (MCH, 800 uL/L) or 3-octanol (OCT, 600 uL/L) with a mechanical

shock (CS^+). Three cycles of conditioning were conducted, each consisting of the sequence: 30 s

exposure to one odor paired with mechanical shock (pulsating 1s every 5s), 60 s break of humid

air, 30 s exposure to other odor with no shock (CS^-), 60 s humid air. We tested odorant

preference of the conditioned flies after a 1 h retention interval by giving them 60 s to choose

between odorants in an elevator T-maze. The *cpo* haplotypes were also assessed for immediate

response (5 min retention) and long-term memory (24 h retention). Half of the fly groups were

conditioned to avoid each odorant to account for innate preference for either odor.

To exclude impairment in odor perception or innate preference we performed two

additional controls. Controls for absolute preference subjected flies to mechanical shock cycle

without odorant before presenting a choice between a single odorant and air in the T-maze.

Controls for relative preference tested innate odor preference in naïve flies by giving a choice

between the two odors in the T-maze.

Appetitive learning in a natural environment

Appetitive conditioning learning assays were adapted from previously developed food

substrate-based protocols (Mery and Kawecki 2002; Zrelec et al. 2013), but were implemented in

a natural setting in an experimental orchard over three replicate days. Flies were reared in

standard laboratory conditions at controlled density and 3d cohorts were sorted into groups of 50

by sex. Flies were marked by treatment using fluorescent powder according to learning treatment

and source population and kept for 12h on an agar substrate. The flies were then exposed to

either strawberry or apple food for 8h and had a 4h rest period on a fresh agar substrate before

being released into the outdoor testing phase. Four hundred flies from each treatment, season and

sex combination were released together into the same outdoor mesocosm, a 0.6x0.6x1.8m mesh

cage with plant bedding covering the ground. Each mesocosm was placed underneath a peach

tree inside a larger 8 m^3 cage. Eight pairs of strawberry and apple unidirectional traps were

dispersed around the mesocosm and flies were scored as learning if they selected the same food

type that they had been previously exposed to during the conditioning period. The powder

marking was an effective method of labeling the flies and all of the trapped flies showed traces

of the powder; this also indicates that no additional flies infiltrated the experiment from outside.

Fecundity

Fecundity of the reconstructed seasonal populations was measured by placing 25 virgin

females from each reconstructed seasonal population into culture bottles (Paaby et al. 2014) with

25 males from a standardized stock (BDSC-3605). Food plates were changed daily for ten days

to count the number of eggs. Average fecundity was calculated daily across 10 replicate bottles

per population.

Individual trade-off between learning and reproduction was assessed in the spring

reconstructed seasonal populations from 2013 and 2014. Virgins were collected, aged in vials of

standard food for 3 days, and subject to the aversive olfactory conditioning and test described

above. The flies were thus divided in "learners" (those that chose CS-) and "non-learners" (those

that chose CS+). From each replicate test, five "learner" and five "non-learner" females were

placed onto food with 5 males; all available flies were used in the few replicates with less than

five flies. The flies were housed on food supplemented with topical yeast to promote egg

production before the assay and on grape-agar substrate during the egg collection. Daily

fecundity was counted for 3d post learning assessment and the mean fecundity was calculated

based on the number of females in the vial.

Statistical analysis

All statistical analyses were performed using the R software (R Core Team, version R

3.2.2). To analyze learning performance, we modeled proportion of flies selecting the odor

against which they had been conditioned (CS^+) using generalized linear models with mixed

effects fit by maximum likelihood (package *lme4,* Bates et al. 2014). To test for the significance

of main effects and interactions, we used type II Wald chi-square test from the package *car* and

functional analysis of variance (ANOVA, Fox and Weisberg 2011). Flies that chose no odor and

remained in the center of the T-maze were excluded from analysis. For illustration purposes, the

proportion of flies learning was rescaled as $2 \times$ proportion $- 1$ to make it range from -1 to 1 with

0 indicating no learning.

For the seasonal populations, we used the following model:

Response = Season + Year + Direction of conditioning + Replicate

where Season, Year and Direction of conditioning are fixed effects and Replicate is a random

effect.

The effect of the natural *cpo* haplotypes was determined using the following model:

Response = Genotype + Sex + Direction of conditioning + Retention Interval + Replicate

where Genotype, Sex, Direction of conditioning and Retention Interval are fixed effects and

Replicate is a random effect.

To assess the RNAi knockouts, we used the following model:

Response = cpo*Gal4*Direction of conditioning + UAS + Replicate

where *cpo* (presence or absence), Gal4 (location), Sex and Direction of conditioning are fixed

effects and UAS driver and Replicate are random effects. Tukey's Honestly Significant

Difference test was used as a post-hoc evaluation of the effect of *cpo* in specific tissues using the

lsmeans package (Lenth 2016) in R.

Population measurement of fecundity was assessed using mixed model ANOVAs with

the fixed effects of season and year (seasonal populations) and genotype (*cpo* ROP) and the

random effect of replicate bottle. Individual fecundity of flies that learned compared to those that

did not learn was calculated using a paired T-test.

Results

Rapid evolution of learning and fecundity in a natural population

Learning ability in the wild *D. melanogaster* population evolved rapidly and repeatedly

across seasonal time. The spring collections learned better than the fall from the same year when

assessed using laboratory aversive conditioning (χ^2_1=4.37, p=0.036; Figure 1a). These behavioral

differences were not due to differences in ability to perceive odor, as there was no effect of

season in absolute preference between odorant and solvent in unconditioned flies (χ^2_1=0.24,

p=0.62 Figure 1b). The spring collections also tended to show higher appetitive learning in an

assay performed in outdoor mesocosms: spring flies were better at learning the association

between food and odor as they returned to the known high-quality food source at higher

frequency than the fall flies (χ^2_1= 2.56, p=0.11, Figure 1c). Taken together, the combined

aversive and appetitive conditioning results strengthen the evidence for a higher learning ability

of the spring flies ($\chi^2_4 = 11.0$, $p = 0.026$, Fisher's method for combining *p*-values).

The seasonal pattern of fecundity was opposite to that for learning ability at the

population level: the fall populations laid on average 30% more eggs per female per day than the

spring populations (Figure 1d; $F_{1,57} = 44.2$, p < 0.0001). A negative association between learning

and reproduction was also apparent at the individual level. Females that did not learn in the

aversive learning assay laid an average of six more eggs (25% more) per day compared to the

females that avoided the shock-associated odor (t_{14}= -2.84, p = 0.013, Figure 1e).

Natural variation in cpo sequence affects learning and fecundity

Recombinant outbred populations (ROPs) homozygous for the spring cpo^{TTA} haplotype

had higher aversive learning than the fall ROP; this pattern persisted across all retention intervals

between conditioning and testing ($\chi_1^2 = 34.88$, p = 3.5×10^{-9}; Figure 2a). In the absence of

conditioning, the spring ROPs showed a weaker avoidance of odors paired with air than the fall

with 1h between training and conditioning (χ_1^2=8.46, p=3.6×10^{-3}, Figure 2b), but not at 5m or

24h retention intervals. The *cpo* ROPs did not differ in the relative preference of naïve flies

choosing between the odors ($\chi_1^2 = 0.016$, p = 0.90, Figure 2c). The spring ROPs also had higher

appetitive associative learning when assessed in the natural mesocosms (χ_1^2=3.62, p=0.057,

Figure 2a). The spring cpo^{TTA} ROP was characterized by lower fecundity, with females laying

36% fewer eggs than females from the fall cpo^{CGT} ROP (Figure 2d, F_2 =8.23, p=0.01).

cpo expression in the peripheral nervous system affects learning

The two intronic SNPs in the *cpo* haplotype may regulate *cpo* expression as the flies with

the spring *cpo* haplotype have lower full body *cpo* expression than flies that carry the fall *cpo*

haplotype (Behrman et al. n.d.). Therefore, we tested if *cpo* expression mediates the differences

in learning and found a tissue-specific effect of *cpo* on learning with a significant interaction

between *cpo* expression and Gal4 driver tissue (χ^2_5=14.26 p=0.014, Figure 3). There was no

difference in learning when *cpo* was knocked down in the Actin 88F promotor control

(BSC38461), in the broad brain knockout (BDSC30818), in a subset of ventral nerve cord and

sensory neurons (BDSC-40315) or in the sense organ support cells and subsets of ventral nerve

cord, brain and sensory neurons (BDSC-40334). Knocking down *cpo* expression in a subset of

the sensory neurons (BDSC-40319) decreased learning (Tukey HSD: z=3.35 p=0.0008).

Discussion

Rapid evolution in natural populations

Learning and fecundity both evolve rapidly in a natural population of *D. melanogaster*

over the scale of approximately 10-15 generations from spring to fall and approximately 1-2

generations between fall and spring. The differences can be attributed to annual cyclical genetic

changes in natural populations because environmental effects are removed by rearing and testing

these populations in common laboratory conditions. The repeatability across years indicates that

this is not a result of genetic drift but instead a deterministic evolutionary process and genomic

data excludes gene flow through migration generating the seasonal cycles (Bergland et al. 2014).

Therefore, the rapid and repeatable seasonal changes in learning and reproduction are consistent

with seasonally fluctuating selection. Learning is thus not only a mechanism of plasticity that

allows organisms to respond rapidly to environmental change (Crombach and Hogeweg 2008);

we demonstrate that learning ability itself can evolve rapidly in nature.

Learning evolved rapidly across seasonal time with higher learning in the spring

compared to the fall. The results are counter to the prediction that learning is favored during the

summer when flies are active and use behaviors that involve learning when assessed in

laboratory. However, cognitive abilities may evolve to match the demands posed by an

organism's biological and physical environment (Healy and Jones 2002; Smid and Vet 2016).

Energetically challenging environments (e.g., cold winters or severe droughts) are hypothesized

to favor cognitive performance at the cost of other physiological systems receiving less resources

(Maille and Schradin 2016). For example, cache seed recovery success in several bird species

suggests that learning may be important for overwintering survival in harsh environments

(Bednekoff et al. 1997; Pravosudov and Clayton 2002; Olson et al. 2004). Although flies do not

cache food, it is possible that learning is also important for other aspects of *D. melanogaster*

overwintering survival in hash climates, such as the ability to find a suitable overwintering site.

Alternatively, rapid evolution of learning may be driven by pleiotropic effects of alleles that

fluctuate in frequency for other reasons such as a correlated trade-off with fecundity.

Trade-off between learning and fecundity

The rapid evolution of learning and fecundity are consistent with patterns of seasonal

evolution of other life history traits in natural populations of *D. melanogaster.* Previous studies

have demonstrated that spring populations collected after the winter bottleneck are more

vigorous with higher propensity for reproductive diapause (Schmidt and Conde 2006), greater

stress resistance (Behrman et al. 2015), higher post-infection survival (Behrman et al. 2018).

Here, we show that spring flies are also better at learning compared to fall flies collected from

the same location. Our finding of higher reproductive output in fall supports the hypothesis that

seasonal oscillations in traits and allele frequencies are caused by alternating selection for

robustness required for survival in the harsh winter conditions and for reproduction and

population growth during the summer expansion (Schmidt et al. 2005; Schmidt and Paaby 2008).

Our data demonstrate a pattern of cyclical seasonal selection with general robustness

favored in the winter and fecundity selected for during the summer. However, it remains unclear

which traits are directly being selected for, particularly given genetic correlations among traits

and complexity of genetic architecture. The negative correlation between learning and

reproduction is consistent with previous studies across a range of taxa (Galea et al. 1994; Mery

and Kawecki 2004; Snell-Rood et al. 2011). However, artificial selection in *D. melanogaster*

indicates independent trajectories of cognitive and reproductive aging, indicating some

differences in the genetic architecture of these traits with age (Zwoinska et al. 2017). It is

possible that rapid evolution of learning may be a product of the negative correlation between

learning and fecundity. The exponential population growth throughout the summer may select

for increased fecundity and therefore result in lower learning ability as a correlated response.

Natural variants in cpo affect learning

Selection on pleiotropic genes could result in the correlated trait changes in the

population across seasonal time. Variants of *cpo* show latitudinal clines and seasonal fluctuations

in frequency (Schmidt et al. 2008; Kolaczkowski et al. 2011; Fabian et al. 2012; Cogni et al.

2013; Bergland et al. 2014). *cpo* is known to be involved in many traits including activity,

dormancy, fecundity and lifespan (Behrman et al. n.d.; Bellen et al. 1992b; Schmidt et al. 2008),

but learning is a new phenotype for this pleiotropic gene. Learning performance and fecundity of

the recombinant outbred populations homozygous for the *cpo* haplotype variants matched the

seasonal pattern of those traits in the wild. The haplotype with SNP variants that are more

frequent in the spring showed higher learning and lower reproduction than the fall haplotype.

The flies containing the spring *cpo* haplotype showed a weaker avoidance of odors when paired

with air during the control. Previous work has shown that a knockdown of *cpo* through p-element

insertion also reduced odor avoidance (Sambandan et al. 2006). This leaves the possibility that

flies containing the spring *cpo* haplotypes have a reduced perception of odors; however, this

should have led to impaired rather than improved learning. Thus, the difference in learning

between the haplotypes is thus unlikely to be due to differences in ability to perceive odor.

Rather, it appears that natural variation in *cpo* haplotypes, or genetic variants linked to them,

serve as an integrator for sensing and responding to environmental changes. Lower full-body *cpo*

expression in the spring ROPs (Behrman et al. n.d.) may contribute to the higher learning ability

in these flies. We tested if *cpo* regulation affects learning ability using GAL4-UAS and found the

opposite pattern: decreasing *cpo* expression in the sensory neurons and ventral nerve cord

decreased learning ability. Our results indicate that *cpo* expression in the peripheral nervous

system is involved in learning. The contrary pattern in the ROPs may due to masking by other

tissues (e.g, ovaries) that express high levels of *cpo*. Altogether, these results imply that the

seasonally fluctuating evolution of learning ability is at least in part mediated by polymorphisms

in *cpo*.

Pleiotropy as a driving force in rapid evolution of learning

The effect of *cpo* on numerous fitness traits beside learning is consistent with evidence

that many learning genes have broad pleiotropic effects (Dubnau et al. 2002; Butcher et al.

2006). For example, a natural polymorphism in *D. melanogaster foraging* (*for*) gene is highly

pleiotropic: in addition to learning (Kaun et al. 2007; Mery et al. 2007) it affects many traits,

including foraging behavior (Fitzpatrick and Sokolowski 2004) and aggregation (Wang and

Sokolowski 2017).

415   The overall force of selection acting on pleiotropic polymorphisms in natural populations

reflects their aggregate impact on survival and reproduction mediated by the diverse ecologically

relevant traits they influence. It is unclear which traits are directly under selection for their

fitness benefits and which evolve as a byproduct of natural selection on correlated traits. The

learning differences between *cpo* haplotypes are likely to generate fitness differences that

contribute to seasonal adaptation in natural populations, either through alternative coadapted

strategies or are a non-adaptive mechanistic consequence of gene action.

Acknowledgements:

This work was supported by NSF GRF DGE-0822 (ELB), the Rosemary Grant Award from the

Society for the Study of Evolution (ELB), the Teece Dissertation Research Fellowship from the

University of Pennsylvania (ELB), the Peachey Environmental Fund from the Department of

Biology, University of Pennsylvania (ELB), and NIH R01GM100366 (PS), as well as a research

funding from the University of Lausanne to TJK.

Author contributions:

ELB, TJK and PS designed the experiment. ELB performed the experiment and did the analyses.

ELB, TJK and PS wrote the manuscript.

References:

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting Linear Mixed-Effects Models

using lme4. J. Stat. Softw. 67:1–48.

Battesti, M., C. Moreno, D. Joly, and F. Mery. 2012. Report Spread of Social Information and

Dynamics of Social Transmission within Drosophila Groups. Curr. Biol. 22:309–313.

Bednekoff, P. A., R. P. Balda, A. C. Kamil, and A. G. Hile. 1997. Long-term spatial memory in

four seed-caching corvid species. Anim. Behav 53:335–341.

Behrman, E. L., A. O. Bergland, D. A. Petrov, and P. S. Schmidt. n.d. Intragenic epistasis in

couch potato and its effect on climatic adaptation in natural populations in Drosophila

melanogaster.

Behrman, E. L., V. M. Howick, M. Kapun, F. Staubach, A. O. Bergland, D. A. Petrov, B. P.

Lazzaro, and P. S. Schmidt. 2018. Rapid seasonal evolution in innate immunity of wild

Drosophila melanogaster. Proc. R. Soc. B 285:20172599.

Behrman, E. L., S. S. Watson, K. R. O'Brien, M. S. Heschel, and P. S. Schmidt. 2015. Seasonal

variation in life history traits in two Drosophila species. J. Evol. Biol. 28:1691–1704.

Bellen, H. J., S. Kooyer, D. D'Evelyn, and J. Pearlman. 1992a. The Drosophila couch potato

protein is expressed in nuclei of peripheral neuronal precursors and shows homology to

RNA-binding proteins. Genes Dev. 6:2125–36.

Bellen, H. J., H. Vaessin, E. Bier, A. Kolodkin, D. D'Evelyn, S. Kooyer, and Yuh Nung Jan.

1992b. The Drosophila couch potato gene: An essential gene required for normal adult

behavior. Genetics 131:365–375.

Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, D. A. Petrov, K. R.

O'Brien, P. S. Schmidt, and D. A. Petrov. 2014. Genomic Evidence of Rapid and Stable

Adaptive Oscillations over Seasonal Time Scales in Drosophila. PLoS Genet. 10:e1004775.

Brandes, C., B. Frish, and R. Menzel. 1988. Time course of memory formation differs in

honeybee lines selected for good and poor learning. Anim. Behav. 36:981–985.

Butcher, L. M., J. K. Kennedy, and R. Plomin. 2006. Generalist genes and cognitive

neuroscience. Curr. Opin. Neurobiol. 16:145–151.

Carroll, S. P., A. P. Hendry, D. N. Reznick, and C. W. Fox. 2007. Evolution on ecological time-

scales. Funct. Ecol. 21:387–393.

Celniker, S. E., L. A. L. Dillon, M. B. Gerstein, K. C. Gunsalus, S. Henikoff, G. H. Karpen, M.

Kellis, E. C. Lai, J. D. Lieb, D. M. MacAlpine, G. Micklem, F. Piano, M. Snyder, L. Stein,

K. P. White, and R. H. Waterston. 2009. Unlocking the secrets of the genome. Nature

459:927–930.

Cogni, R., C. Kuczynski, S. Koury, E. Lavington, E. L. Behrman, K. R. O'Brien, P. S.

Schmidt, W. F. Eanes, K. R. O'Brien, P. S. Schmidt, and W. F. Eanes. 2013. The Intensity

of Selection Acting on the Couch Potato Gene-Spatial-Temporal Variation in a Diapause

Cline. Evolution (N. Y). 68:538–548.

Crombach, A., and P. Hogeweg. 2008. Evolution of evolvability in gene regulatory networks.

PLoS Comput. Biol. 4:e1000112.

Croston, R., C. L. Branch, D. Y. Kozlovsky, T. C. Roth, L. D. LaDage, C. A. Freas, and V. V

Pravosudov. 2015. Potential mechanisms driving population variation in spatial memory

and the hippocampus in food-caching chickadees. Pp. 354–371 *in* Integrative and

Comparative Biology.

Dubnau, J., A.-S. Chiang, and T. Tully. 2002. Neural substrates of memory: From synapse to

system. J. Neurobiol. 54:238–253.

Dukas, R. 2004. Evolutionary Biology of Animal Cognition. Annu. Rev. Ecol. Evol. Syst.

35:347–374.

Dukas, R. 2008. Evolutionary Biology of Insect Learning Learning: the acquisition of neuronal

representations of new information. Annu. Rev. Entomol 53:145–60.

Dukas, R. 1998. Evolutionary Ecology of Learning. Pp. 129–174 *in* Cognitive Ecology: The

Evolutionary Ecology of Information Processing and Decision Making.

Dukas, R. 2005. Learning affects mate choice in female fruit flies. Behav. Ecol. 16:800–804.

Fabian, D. K., M. Kapun, V. Nolte, R. Kofler, P. S. Schmidt, C. Schlötterer, and T. Flatt. 2012.

Genome-wide patterns of latitudinal differentiation among populations of Drosophila

melanogaster from North America. Mol. Ecol. 21:4748–4769.

Fitzpatrick, M. J., and M. B. Sokolowski. 2004. In Search of Food: Exploring the Evolutionary

Link Between cGMP-Dependent Protein Kinase (PKG) and Behaviour. Integr. Comp. Biol.

44:28–36.

Fox, J., and S. Weisberg. 2011. An {R} Companion to Applied Regression.

Froissart, L., M. Giurfa, S. Sauzet, and E. Desouhant. 2017. Cognitive adaptation in asexual and

sexual wasps living in contrasted environments. PLoS One 12.

Galea, L. A. M., M. Kavaliers, K. P. Ossenkopp, D. Innes, and E. L. Hargreaves. 1994. Sexually

dimorphic spatial learning varies seasonally in two populations of deer mice. Brain Res.

635:18–26.

Grant, P. R., and B. R. Grant. 2002. Unpredictable evolution in a 30-year study of Darwin's

finches. Science (80-.). 296:707–711.

Healy, S. D., and C. M. Jones. 2002. Animal learning and memory: an integration of cognition

and ecology. Zoology 105:321–327.

Hoedjes, K. M., and H. M. Smid. 2014. Natural variation in long-term memory formation among

Nasonia parasitic wasp species. Behav. Processes 105:40–45.

Hollis, B., and T. J. Kawecki. 2014. Male cognitive performance declines in the absence of

sexual selection. Proc. R. Soc. B Biol. Sci. 281:20132873.

Johnston, T. D. 1982. Selective costs and benefits in the evolution of learning. Adv. Study

Behav. 12:65–106.

Kaun, K. R., T. Hendel, B. Gerber, and M. B. Sokolowski. 2007. Natural variation in Drosophila

larval reward learning and memory due to a cGMP-dependent protein kinase. Learn. Mem.

14:342–349.

Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. E. Hill, A.

Hoang, P. Gibert, and P. Beerli. 2001. The strength of phenotypic selection in natural

populations. Am. Nat. 157:245–61.

Kolaczkowski, B., A. D. Kern, A. K. Holloway, and D. J. Begun. 2011. Genomic Differentiation

Between Temperate and Tropical Australian Populations of Drosophila melanogaster.

Genetics 187:245–260.

Lenth, R. V. 2016. Least-Squares Means: The R Package lsmeans. J. Stat. Softw. 69.

MacKay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, D. Zhu, S. Casillas, Y.

Han, M. M. Magwire, J. M. Cridland, M. F. Richardson, R. R. H. Anholt, M. Barrón, C.

Bess, K. P. Blankenburg, M. A. Carbone, D. Castellano, L. Chaboub, L. Duncan, Z. Harris,

M. Javaid, J. C. Jayaseelan, S. N. Jhangiani, K. W. Jordan, F. Lara, F. Lawrence, S. L. Lee,

P. Librado, R. S. Linheiro, R. F. Lyman, A. J. MacKey, M. Munidasa, D. M. Muzny, L.

Nazareth, I. Newsham, L. Perales, L. L. Pu, C. Qu, M. Ràmia, J. G. Reid, S. M. Rollmann,

J. Rozas, N. Saada, L. Turlapati, K. C. Worley, Y. Q. Wu, A. Yamamoto, Y. Zhu, C. M.

Bergman, K. R. Thornton, D. Mittelman, and R. A. Gibbs. 2012. The Drosophila

melanogaster Genetic Reference Panel. Nature 482:173–178.

Maille, A., and C. Schradin. 2016. Ecophysiology of cognition: How do environmentally

induced changes in physiology affect cognitive performance? Biol. Rev. 92:1101–1112.

Mansourian, S., J. Corcoran, A. Enjin, C. Löfstedt, M. Dacke, and M. C. Stensmyr. 2016. Fecal-

Derived Phenol Induces Egg-Laying Aversion in Drosophila. Curr. Biol. 26:2762–2769.

Mayr, E. 1974. Behavior Programs and Evolutionary Strategies: Natural selection sometimes

favors a genetically "closed" behavior program, sometimes an "open" one. Am. Sci.

62:650–659.

Mcguire, T. R., and J. Hirschth. 1977. Behavior-genetic analysis of Phormia regina:

Conditioning, reliable individual differences, and selection. 74:5193–5197.

Mery, F., A. T. Belay, A. K-C So, M. B. Sokolowski, T. J. Kawecki, and G. E. Robinson. 2007.

Natural polymorphism affecting learning and memory in Drosophila. Proc. Natl. Acad. Sci.

104:13051–13055.

Mery, F., and T. J. Kawecki. 2004. An operating cost of learning in Drosophila melanogaster.

Anim. Behav. 68:589–598.

Mery, F., and T. J. Kawecki. 2005. Cost of Long-Term Memory in Drosophila. Science (80-.).

308:1148.

Mery, F., and T. J. Kawecki. 2002. Experimental evolution of learning ability in fruit flies. Pnas

99:14274–14279.

Mery, F., and T. J. Kawecki. 2004b. The effect of learning on experimental evolution of

resocurce preference in Drosopihla melanogaster. Evolution (N. Y). 58:757–767.

Odling-Smee, L. C., J. W. Boughman, and V. A. Braithwaite. 2008. Sympatric species of

threespine stickleback differ in their performance in a spatial learning task. Behav. Ecol.

Sociobiol. 62:1935–1945.

Olson, D. J., A. C. Kamil, R. P. Balda, and P. J. Nims. 2004. Performance of Four Seed-Caching

Corvid Species in Operant Tests of Nonspatial and Spatial Memory. J. Comp. Psychol.

109:1–9.

Paaby, A. B., A. O. Bergland, E. L. Behrman, and P. S. Schmidt. 2014. A highly pleiotropic

amino acid polymorphism in the Drosophila insulin receptor contributes to life-history

adaptation. Evolution (N. Y). 68:3395–3409.

Paenke, I., B. Sendhoff, and T. J. Kawecki. 2007. Influence of Plasticity and Learning on

Evolution under Directional Selection. Am. Nat. 170:E47–E58.

Pravosudov, V. V., and N. S. Clayton. 2002. A test of the adaptive specialization hypothesis:

Population differences in caching, memory, and the hippocampus in black-capped

chickadees (Poecile atricapilla). Behav. Neurosci. 116:515–522.

Sambandan, D., A. Yamamoto, J.-J. Fanara, T. F. C. Mackay, and R. R. H. Anholt. 2006.

Dynamic genetic interactions determine odor-guided behavior in Drosophila melanogaster.

Genetics 174:1349–63.

Sarin, S., and R. Dukas. 2009. Social learning about egg-laying substrates in fruitflies. Proc. R.

Soc. B Biol. Sci. 276:4323–4328.

Schmidt, P. S., and D. R. Conde. 2006. Environmental heterogeneity and the maintenance of

genetic variation for reproductive diapause in Drosophila melanogaster. Evolution 60:1602–

11.

Schmidt, P. S., and A. B. Paaby. 2008. Reproductive diapause and life-history clines in North

American populations of Drosophila melanogaster. Evolution (N. Y). 62:1204–1215.

Schmidt, P. S., A. B. Paaby, and M. S. Heschel. 2005. Genetic variance for diapause expression

and associated life histories in Drosophila melanogaster. Evolution (N. Y). 59:2616–2625.

Schmidt, P. S., C.-T. C.-T. Zhu, J. Das, M. Batavia, L. Yang, and W. F. Eanes. 2008. An amino

acid polymorphism in the couch potato gene forms the basis for climatic adaptation in

Drosophila melanogaster. Proc. Natl. Acad. Sci. U. S. A. 105:16207–16211.

Smid, H. M., and L. E. Vet. 2016. The complexity of learning, memory and neural processes in

an evolutionary ecological context. Curr. Opin. Insect Sci. 15:61–69.

Snell-Rood, E. C., G. Davidowitz, and D. R. Papaj. 2011. Reproductive tradeoffs of learning in a

butterfly. Behav. Ecol. 22:291–302.

Stensmyr, M. C., H. K. M. Dweck, A. Farhan, I. Ibba, A. Strutz, L. Mukunda, J. Linz, V. Grabe,

K. Steck, S. Lavista-Llanos, D. Wicher, S. Sachse, M. Knaden, P. G. Becher, Y. Seki, and

B. S. Hansson. 2012. A Conserved Dedicated Olfactory Circuit for Detecting Harmful

Microbes in Drosophila. Cell 151:1345–1357.

Thompson, J. N. 2013. Relentless evolution. University of Chicago.

Tryon, R. C. 1940. Studies in individual differences in maze ability. VII. The specific

components of maze ability, and a general theory of psychological components. J. Comp.

Psychol. 30:283–335.

Wang, S., and M. B. Sokolowski. 2017. Aggressive behaviours, food deprivation and the

foraging gene. R. Soc. Open Sci. 4:170042.

Yurkovic, A., O. Wang, A. C. Basu, and E. A. Kravitz. 2006. Learning and memory associated

with aggression in Drosophila melanogaster. Proc. Natl. Acad. Sci. U. S. A. 103:17519–

17524.

Zrelec, V., M. Zini, S. Guarino, J. Mermoud, J. Oppliger, A. Valtat, V. Zeender, and T. J.

594  Kawecki. 2013. *Drosophila* rely on learning while foraging under semi-natural conditions.

595  Ecol. Evol. 3:4139–4148.

596 Zwoinska, M. K., A. A. Maklakov, T. J. Kawecki, and B. Hollis. 2017. Experimental evolution

597  of slowed cognitive aging in Drosophila melanogaster. Evolution (N. Y). 71:662–670.

Figures:

Figure 1. Rapid evolution of learning and reproduction in rapid evolution in natural populations

of *Drosophila melanogaster*. (a) Higher learning (mean +/-SE) in the spring populations

compared to the fall replicated across three years: 2012 (square), 2013 (circle), 2014 (triangle).

(b) No difference absolute preference in either population when unconditioned flies are given the

choice between one odor and solvent. (c) Spring populations return to positive conditioning food

at a higher rate than fall populations. (d) Lower reproduction in the spring populations compared

to the fall over ten days. (e) Individuals that do not learn in the aversive conditioning assay have

higher daily reproductive output than those that learn over three days. Difference between

individuals is shown in small outlined shapes and the mean difference +/- SE in the large, filled

shapes.

Figure 2. Natural variants in *couch potato* (*cpo*) involved in trade-off between learning and

reproduction. (a) Flies containing the spring (cpo^{TTA}) haplotype have higher learning (mean +/-

SE) than flies that contain the fall (cpo^{CGT}) haplotype across all retention intervals between

conditioning and testing using aversive shock, as well as in the appetitive conditioning of the

natural mesocosm. (b) Unconditioned flies containing the spring haplotype have a slightly higher

absolute preference for odor instead of solvent compared to the flies containing the fall

haplotype. (c) However, there is no relative preference for either of the experimental odors in

unconditioned flies. (d) Flies containing the fall haplotype had higher daily reproductive output

than flies containing the spring haplotype.

Figure 3. Effect of *couch potato* (*cpo*) expression on aversive shock learning. Learning scores

(mean +/- SE) for flies with normal *cpo* expression (filled) and *cpo* knockdown (outline). No

difference in learning when *cpo* was knocked down in the Actin 88F promotor control

(BSC38461), in the broad brain knockout (BDSC30818), in a subset of ventral nerve cord and

sensory neurons (BDSC-40315) or in the sense organ support cells and subsets of ventral nerve

cord, brain and sensory neurons (BDSC-40334). Knocking down *cpo* expression in a subset of

the sensory neurons (BDSC-40319) decreased learning.