Transcriptional deconvolution reveals consistent functional subtypes of pancreatic cancer epithelium and stroma

Jing He,*[1,2,8] H. Carlo Maurer,*[3,8] Sam R. Holmstrom,*[3,8] Tao Su,[8] Aqeel Ahmed,[8] Hanina Hibshoosh,[4,8] John A. Chabot,[5,8] Paul E. Oberstein,[6,8] Antonia R. Sepulveda,[4,8] Jeanine M. Genkinger,[7,8] Jiapeng Zhang,[9] Alina C. Iuga,[4,8] Andrea Califano,[1,2,8,12] Mukesh Bansal,[11,12] and Kenneth P. Olive[3,4,8,12,13]

[1] Department of Biomedical informatics
[2] Department of Systems Biology
[3] Department of Medicine, Division of Digestive and Liver Diseases
[4] Department of Pathology and Cell Biology
[5] Department of Surgery, Division of GI/Endocrine Surgery
[6] Department of Medicine, Division of Hematology and Oncology
[7] Department of Epidemiology, Mailman School of Public Health
[8] Herbert Irving Comprehensive Cancer Center, Columbia University Medical Center, New York, NY 10032
[9] Department of Computer Science and Engineering, University of California, San Diego
[10] Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York
[11] PsychoGenics Inc., Tarrytown, New York, USA.
[*] These authors contributed equally
[12] Correspondence should be addressed to K.P.O. (kenolive@columbia.edu), A.C. (califano@c2b2.columbia.edu), or M.B. (mukesh.bansal@psychogenics.com)
[13] Lead contact.

Summary

Bulk tumor tissues comprise intermixed populations of neoplastic cells and multiple stromal cell lineages. We used laser capture microdissection and RNA sequencing to disentangle the transcriptional programs active in the malignant epithelium and stroma of pancreatic ductal adenocarcinoma (PDA). This led to the development of a new algorithm (ADVOCATE) that accurately predicts the compartment fractions of bulk tumor samples and can computationally purify bulk gene expression data from PDA. We also present novel stromal subtypes, derived from 110 microdissected PDA stroma samples, that were enriched in extracellular matrix– and immune–associated processes. Finally, we applied ADVOCATE to systematically evaluate cross–compartment subtypes spanning four patient cohorts, revealing consistent functional classes and survival associations despite substantial compositional differences.

Keywords
Pancreatic Cancer, Pancreatic Ductal Adenocarcinoma, Gene Expression, Deconvolution, Laser Capture Microdissection, Subtypes, Stroma

Introduction

All carcinomas harbor both transformed malignant cells and non-transformed stromal cells, in varying proportions (Aran et al., 2015). Pancreatic ductal adenocarcinoma (PDA) is among the most stroma–rich cancers, with a complex inflammatory microenvironment that typically dominates the tumor parenchyma. Expected to be responsible for over 43,000 deaths per year in the United States, it is a common, aggressive malignancy that responds poorly to therapeutic intervention (Oberstein and Olive, 2013; Siegel et al., 2017). Within the stromal compartment of PDA, diverse fibroblast, myeloid, lymphoid, endothelial and other cell lineages contribute to both pro– and anti–tumor processes, including angiogenesis and epithelial differentiation (Rhim et al., 2014), tissue stiffness (Jacobetz et al., 2012; Provenzano et al., 2012), drug delivery (Olive et al., 2009), and local immunosuppression (Vonderheide and Bayne, 2013). These functions are orchestrated through a host of paracrine signals that pass between and within the epithelial and stromal compartments– communication that is quickly altered upon tissue disruption. Thus, efforts to parse transcriptional programs of PDA should take into account active processes in both compartments, ideally in an *in situ* context.

Despite extensive genomic characterization (Bailey et al., 2016; Biankin et al., 2012; Jones et al., 2008; Waddell et al., 2015; Witkiewicz et al., 2015), individual DNA mutations have thus far failed to provide confirmed prognostic or theranostic information for PDA. Indeed, only a small fraction of pancreatic tumors is predicted to harbor "druggable" genetic alterations (Bailey et al., 2016; Witkiewicz et al., 2015). As an alternative to genetic biomarkers, transcriptional classifiers for PDA have been explored using bulk tumor samples (Bailey et al., 2016; Collisson et al., 2011; Moffitt et al., 2015). While each study differs in the number of subtypes described, a shared message is that ductal pancreatic tumors include at least two groups distinguished by markers of epithelial differentiation state, with the more poorly–differentiated subtype (i.e. "Basal-like", "Squamous", or "Quasi-Mesenchymal") exhibiting reduced overall survival relative to well-differentiated subtypes (i.e. "Classical" or "Progenitor"). However, the contributions of stromal cells are handled differently in each instance, leading to some debate as to the merits of different proposed subtypes. To clarify this issue, we endeavored to directly profile gene expression from

purified neoplastic epithelium and associated stroma isolated from frozen human PDA samples.

Several techniques may be employed to isolate cellular subsets from bulk tissue including magnetic separation, fluorescence assisted cell sorting (FACS), and laser capture microdissection (LCM). The first two techniques rely on population–specific antibodies to separate a suspension of cells following disruption of the tumor. Unfortunately, the extremely fibrotic extracellular matrix of PDA necessitates prolonged enzymatic digestion to achieve a single-cell suspension, during which time transcriptional profiles are invariably altered. Moreover, PDA diffusely infiltrates the surrounding pancreatic parenchyma (Hruban, 2007) so that even tumor samples enriched by FACS for epithelial markers can include contributions from normal, atrophic, pre-neoplastic, or metaplastic epithelial cells. Laser capture microdissection (LCM) provides a powerful solution, allowing the isolation of relatively pure compartment–specific tissue samples based on morphological features, without disrupting the delicate interplay of intercellular communication. However, performing LCM while maintaining RNA quality suitable for sequencing is technically challenging, costly, and labor intensive. As a result, there are currently few examples of large collections of cancer gene expression profiles derived from LCM samples.

We present here expression profiles of laser capture microdissected malignant epithelium and matched reactive stroma for 66 human pancreatic ductal adenocarcinomas. These data informed the development of a novel computational algorithm for the **A**daptive **D**econ**V**olution **O**f **CA**ncer **T**issue **E**xpression (ADVOCATE), which we used to extend our analyses to external datasets. ADVOCATE may be used to perform two functions on bulk tumor expression profiles. First, it can infer the fractional contribution made by each compartment to the original bulk sample, while accounting for potential infiltration of uncharacterized tissue, as well as bias from different profiling platforms. Second, it can transform a bulk tissue transcriptome into separate "virtual" profiles for each sub–compartment. Though prior approaches exist for related applications (Abu-Alainin et al., 2016; Kuhn et al., 2011; Zhong et al., 2013), they make several assumptions, such as the linear mixing of compartments and the specificity of marker genes, leading to reduced performance relative to ADVOCATE. Furthermore, a crucial new feature is that ADVOCATE

can fully infer complete gene expression profiles for modeled subpopulations on a sample–by–sample basis, which may then be used by a variety of downstream analytical pipelines.

Given the multitude of cancer-related processes mediated by stromal elements, and the specific prominence of stromal infiltration in many tumor types including PDA, molecular subtyping of stromal signaling programs may offer a distinct and useful counterpoint to epithelial classification. A prior study utilized an elegant computational approach to indirectly infer stroma-specific subtypes, using a limited set of stroma–associated expression factors (Moffitt et al., 2015). In contrast, we provide a novel stromal classification derived from direct, genome-wide analysis of 110 experimentally purified stromal profiles, 66 of which were matched to the corresponding tumor epithelial compartment. Two prominent subtypes emerged from our analyses: one enriched for extracellular matrix associated pathways (ECM–rich) and the other for immune/cytokine associated pathways (Immune–rich), respectively. Application of this novel and comprehensive stroma–specific gene signature to ADVOCATE-based inference of virtual stroma profiles from bulk tumor samples led to consistent identification of the same functional subtypes across three large, independent PDA gene expression datasets; similar consistency was also observed using an existing epithelium–specific classification system. Notably, meta-analysis of all four cohorts revealed partial association of the ECM–rich stroma subtype with the Basal–like epithelial subtype. Finally, we demonstrated that incorporating molecular subtypes from both the epithelial and stromal compartments into a combined classification led to the identification of subtypes that are strongly associated with patient survival across multiple cohorts, thus producing a consistent, compartment-specific interpretation of datasets that could not previously be effectively reconciled, based on bulk gene expression profile data.

Results

Transcriptional profiling of pancreatic cancer epithelium and stroma

To study the separate transcriptional programs of intact pancreatic tumor epithelium and stroma, we optimized a robust protocol for maintaining RNA integrity during laser capture

microdissection of frozen tumor tissues, yielding total RNA suitable for library preparation and RNA sequencing. We first applied this LCM–RNA–Seq technique to 60 primary PDA specimens that were harvested and frozen intraoperatively by the Columbia University Tumor Bank in collaboration with the Columbia Pancreas Center (see Tables S1,2 for patient characteristics). For each tumor, we generated paired gene expression profiles from the malignant epithelium and nearby reactive stroma, as distinguished by cell morphology (Figure 1A). Extensive quality control metrics confirmed the high quality of resulting RNA libraries (Figures 1B,C and Figures S1A-D) (Adiconis et al., 2013; Shanker et al., 2015). Critically, samples from the two compartments separated spontaneously along the first component of a Principal Component Analysis (PCA) with virtually no overlap (Figure 1D), and were distinguished by expression of established marker genes for epithelial cells (KRT19, EPCAM, CDH1) versus markers of various stromal cell types, including leukocytes (PTPRC, CD4, CD163), endothelial cells (VWF, ENG, CDH5), and cancer associated fibroblasts (CAFs) (ACTA2, DCN, FAP) (Figure 1E). We observed that technical variance was substantially lower than biological variance (Figures S1E-F) and found that different malignant areas captured from a single tumor clustered closely, suggesting that the intra-tumoral transcriptional heterogeneity of that tumor was less than the inter-tumoral heterogeneity of PDA (Figures S1G-H). In summary, our analysis shows that LCM–RNA–Seq produces robust, genome–wide, compartment–specific gene expression profiles.

We next examined the most differentially expressed genes between the epithelium and the stroma (Table S3) and compared their expression to immunohistochemistry for the corresponding proteins in The Human Protein Atlas (HPA) pathology database (Pontén et al., 2008). We restricted our analysis to proteins for which the highest-quality antibodies were available (n= 321), based on established HPA criteria (Table S4). Of these, we evaluated the immunostaining patterns for the 50 genes whose LCM–RNA–Seq expression was most differentially expressed for each compartment, examining a minimum of six PDA samples per tested protein. This analysis yielded confirmatory staining patterns for 47 of 50 epithelial proteins and 36 of the 50 stromal proteins (Figures 1 F, G, Table S5). For example, Figures 1 H, I show two members of the galectin protein family, LGALS4 and LGALS1, with inverse staining patterns in the two compartments, consistent with our predictions. Critically, none of the proteins were found expressed in a pattern opposite that

predicted; most genes lacking supportive staining were simply not detected, perhaps due to post–translational regulation. Thus, through the use of LCM–RNA–Seq, we compiled a comprehensive repertoire of compartment–specific genes serving as a novel, tumor-specific resource for the pancreatic cancer field.

A framework to deconvolve compartment–specific gene expression profiles from bulk data

Multiple large–scale gene expression datasets for PDA have been reported (Bailey et al., 2016; Collisson et al., 2011; Moffitt et al., 2015), each one providing important contributions to our understanding of the disease. However, cross-comparative analysis of these datasets has been challenging, due to differences in expression profiling platforms, inclusion criteria, sample preparation, and other technical details. As a result, a consistent interpretation of the gene expression profile clusters emerging from these studies is still elusive, especially as it relates to stromal subtypes.

Availability of experimentally–purified, compartment–specific, paired expression profiles offers a unique opportunity to unify these resources by using transcriptional deconvolution to remove the variability introduced by the diverse tissue composition of the individual samples in each dataset. In contrast to many prior optimization techniques that were based on quadratic programming, we utilized a machine-learning approach to train a probabilistic algorithm (ADVOCATE), consisting of four steps, as outlined in Figure 2A: 1) we first fit a Gaussian-mixture model to the collection of normalized gene expression profile pairs to build a probability density model representing the expression of each gene in each compartment (i.e. epithelium and stroma). 2) Given a bulk gene expression profile, the optimal contribution by each compartment is then computed, compatible with the compartment- and gene-specific probability density models inferred from Step 1, thus predicting its epithelial- and stroma-specific contributions. 3) These contributions are then integrated across all genes to infer the most likely composition of the bulk sample given the underlying model from Step 1. 4) Finally, ADVOCATE combines the information from all three steps to infer epithelial and stroma–specific virtual gene expression profiles. The latter would hopefully recapitulate the results of more–costly and laborious LCM–RNA–Seq.

While we focus here on the epithelial and stromal compartments of PDA, the large number of genes available from gene expression profiles should pose no technical limitations on the number of compartments that can be modeled by the analysis, provided that distinct, sample-matched, compartment–specific expression profiles are available to train the algorithm. In addition, the *residuals* of these analyses (i.e., the component of the gene expression signature that cannot be explained by compartment-specific models) can be used to evaluate infiltration of additional unspecified cell lineages in the tumor, thus preventing them from affecting the inference of compartment-specific profiles.

Computational validation of ADVOCATE performance

We carried out a series of *in silico* and experimental assays to validate ADVOCATE's performance. First we computationally generated a set of pure epithelial and stromal gene expression profiles by randomly sampling from the gene expression probability density of each gene as captured by the LCM–RNA–Seq profiles of each compartment (see Methods). We then computationally mixed these profiles in varying proportions to create an equivalent set of synthetic bulk samples with known compartment fraction composition. We then trained ADVOCATE using compartment–specific synthetic samples and used it to predict the composition of the synthetic bulk samples. Error rates from this analysis were < 3% (Figure 2B).

Next we generated a more realistic set of "semi–synthetic" bulk samples by computationally mixing profiles from actual LCM–derived epithelial and stromal sample pairs, in varying proportions (Figure 2C). Again, ADVOCATE predicted the compartment mixture rate with very high accuracy (>90%) for the majority (~80%) of the samples. In a complementary analysis, we estimated compartment–specific LCM sample purity by leave–one–out cross-validation (LOOCV), whereby the composition of each individual LCM sample was estimated after training the model on the remaining 59 pairs. Figure 2D shows that ADVOCATE correctly predicted a very high (>90%) epithelial and stromal composition for the majority of compartment–specific LCM–derived samples (73%), as expected, with the residual error likely due to biological and technical variability in the LCM–derived sample profiles.

Interestingly, both analyses identified the same three outlier samples, presenting >20% error, see Figure 2C and 2D (orange lines/bars). Careful histopathological examination of these samples revealed that one (E17) was poorly differentiated and therefore may plausibly exhibit a more stroma–like signature (Figures S2A-B). A second (S10) had large areas of highly cellular stroma intermixed with fibrotic regions, which could plausibly lead to a more epithelial–like stroma signature (Figure S2C-E). No obvious pathological distinctions were apparent in the third sample (Figure S2F) suggesting either imprecise microdissection, an unusually high level of epithelial delamination into the stroma (Rhim et al., 2012), or poor quality/noise in the gene expression profile. These findings were consistent with the clustering of these three samples near the interface between epithelial and stromal samples by PCA (Figure 1D). When discarding these outlier samples, ADVOCATE's performance on the entire set further improved to 93.54%. Together, the results provide evidence of the robustness of ADVOCATE in predicting bulk tumor composition from synthetic data and its resilience when applied to more realistic experimental data. A power analysis performed using the LOOCV technique found that experimental error diminished only modestly as the size of the training set exceeded 20 samples, suggesting fairly modest sample requirements for training ADVOCATE on a new tumor type (Figure 2E).

Experimental validation of compartment fraction prediction with ADVOCATE

We assessed ADVOCATE's ability to predict the composition of samples whose compartment fraction had been experimentally assessed and compared its performance to that of four previously published algorithms: ESTIMATE (Yoshihara et al., 2013), deconRNAseq (Gong and Szustakowski, 2013), PSEA (Kuhn et al., 2011), and DSA (Zhong et al., 2013). We began by analyzing gene expression profiles from pure epithelial and stromal tissues isolated by LCM on PDA samples (Figures 3A, S3A-B). Among the five algorithms, ADVOCATE predicted the highest epithelial fraction in microdissected epithelial samples, and the second-lowest epithelial fraction in stromal samples; the deconRNAseq algorithm predicted low epithelial fractions in both the stromal and epithelial samples. We next used ADVOCATE to predict the compartment fractions of the CUMC bulk samples for

which nuclei counting had been performed (Figure S3C), as well as bulk PDA samples from TCGA, which are annotated with DNA-based purity estimates using the ABSOLUTE algorithm (Figure S3D). Among the five algorithms, ADVOCATE predicted the highest purity for the respective compartment in the LCM samples, and scored closest to the DNA-based reference standards for the CUMC and TCGA bulk PDA samples (nuclei count and ABSOLUTE prediction, respectively) (Figure S3E).

Carcinoma–derived cell lines are often used to provide a pure epithelial reference for gene expression studies. Thus, we used ADVOCATE to predict the compartment fractions of 40 PDA cell lines in the Cancer Cell Line Encyclopedia (CCLE) (Barretina et al., 2012). As expected, all PDA cell lines were scored as predominantly epithelial with a mean epithelial fraction of 94% (Figure S3F). However, nine lines were predicted to have >10% stromal fraction. This should not be interpreted to imply the presence of stromal cells in these lines. Rather, this result suggests that either the original tumor was poorly differentiated or that the cells underwent epithelial–to–mesenchymal transition (EMT) during their adaptation to tissue culture conditions, since stromal signatures are enriched in mesenchymal genes (Ross et al., 2000). Consistent with that interpretation, sarcoma and mesothelioma cell lines (n = 28) were predicted to be significantly more "stromal" than PDA cell lines (P = 2.30x10^{-25}, Kolmogorov-Smirnov (KS) test, Figure S3G). These findings highlight potential issues in the common practice of using epithelial tumor cell lines indiscriminately as a "pure" epithelial reference for purity assessments.

Finally, to directly test the ability of ADVOCATE to predict compartment fractions from bulk tissue, we performed RNA–Seq on bulk tissue from 15 PDA samples and compared compartment fraction predictions to histopathological assessments. Bulk tumor samples may also contain varying amounts of other tissue types (such as normal or atrophic pancreas, lymphoid aggregates, nerve plexus, and blood vessels) that are routinely found intermixed with frank carcinoma in pancreatic tumors. To account for these confounding factors, we incorporated a third "Residuals" compartment into this analysis, comprising genes that could not be modeled based on the probability density functions inferred from epithelial and stromal tissue, such as genes with low expression probability in both compartments (see Methods). This results in a 3-compartment analysis that can be used to

minimize the effect of unknown tissue infiltration on the estimate of epithelial/stromal composition and in the inference of the associated gene expression profiles.

We note that multiple studies have reported discrepancies between tumor purity estimates from pathology review and those from molecular analyses (Carter et al., 2012; Song et al., 2012; Yoshihara et al., 2013), either as a result of technical limitations or of sampling different areas of the tumor for the respective analyses. To minimize the latter effect, we assessed the histology of tissue sections immediately adjacent to those used for LCM-RNA-Seq. Two blind, independent histopathology assessments of tumor composition were performed on hematoxylin and eosin (H&E) stained frozen tissue sections. First, the areas of epithelium, stroma, and other tissue were estimated by a gastrointestinal pathologist. Second, individual nuclei were counted in the epithelial and stromal compartments for multiple representative tumor areas (Figure 3B). We found that tumor content was highly correlated between the two measures ($\rho = 0.77$), with the area assessment showing lower epithelial content than nuclei counting (Figure 3C). ADVOCATE's predictions tracked very well with nuclei count assessment, with a **m**ean **a**bsolute **p**ercent **e**rror (MAPE) of 15% (Figure 3D). However, it systematically overestimated tumor content, relative to area assessments (MAPE = 32%, Figure 3E), perhaps suggesting that nuclear counts better reflect gene expression contributions from distinct compartments as mRNA abundance is relatively constant on a cell-by-cell basis. Overall, ADVOCATE yielded the lowest MAPE of the five algorithms assessed (Figure S3E).

Experimental validation of virtual expression profiles generated from bulk tumors

The second function offered by ADVOCATE is to extract compartment–specific gene expression data from bulk samples (referred to as "virtual profiles" hereafter). To assess the accuracy of virtual profiles, we performed both bulk RNA-Seq and LCM–RNA–Seq on six tumor samples. We then determined the correlation between the LCM profiles and their bulk counterparts, before and after deconvolution (Figure 3F). Without deconvolution, there was only modest correlation with $\rho = 0.59$ and $\rho = 0.48$ for epithelium and stromal LCM comparisons, respectively. By contrast, following deconvolution with ADVOCATE, the correlation between experimental LCM and virtual profiles increased significantly to $\rho =$

0.71 for the epithelium and $\rho = 0.74$ for the stroma (P = 1.8x10^{-5} combined, two-sample Kolmogorov-Smirnov test). Critically, expression of lineage–specific markers in virtual epithelial and stromal samples closely tracked with the results of experimental microdissection (Figure 3G vs. Figure 1E). Indeed, both hierarchical clustering and PCA showed divergence of virtual epithelial and stromal profiles relative to the intermediate clustering of bulk profiles (Figures 3H, I). Taken together, these data demonstrate that ADVOCATE is effective in deconvolving compartment–specific expression profiles from bulk tissue.

Finally, we sought to assess the general suitability of ADVOCATE for deconvolving tumor subtypes based on published gene expression microarray data from laser-capture microdissected breast cancer samples (Oh et al., 2015) (Figures S4A, B). We trained ADVOCATE on a subset of these samples and then performed LOOCV analysis, finding the predicted fraction to be >80% in the majority of samples from either compartment (Figure S4C). We then applied this implementation to the remaining samples in the dataset, including relatively pure samples that clustered with the training set, as well as others likely to have intermediate compartment representation, based on PCA analysis (yellow points in Figure S4B). As expected, most of the samples from this test were predicted to have high purity, but the intermediate samples from PCA analysis emerged as the least pure ones. (Figure S4D). Finally, we used this implementation of ADVOCATE to predict the compartment fraction of an independent gene expression microarray dataset from LCM breast cancer samples (N=36, Agilent platform, GSE68744) and compared these results to purity predictions from ESTIMATE (Figure S4E). Critically, ADVOCATE successfully predicted quite pure fractions for the relevant compartment on these independent LCM samples. By contrast, ESTIMATE predicted lower purity in both compartments, particularly in epithelial samples. These results demonstrate both the suitability of ADVOCATE for use with different gene expression platforms, and the generalizability of the ADVOCATE algorithm across tumor types when provided appropriate training samples.

Compartment fraction analysis reveals distinct compositions of public PDA datasets

Three prior studies have presented expression–based classification systems, derived from large collections of bulk PDA profiles, that relate to biological properties or outcome (Bailey et al., 2016; Collisson et al., 2011; Moffitt et al., 2015). In each study, a subtype exhibiting molecular characteristics consistent with a poorly differentiated state (termed "Quasi-Mesenchymal", "Basal-like", or "Squamous", respectively) was shown to have poor overall survival, relative to a subtype exhibiting a signature reflective of pancreatic tissue origin ("Classical" or "Progenitor"), with varying numbers of additional groups also presented. Moffitt et. al. further parsed molecular subtypes from the tumor stroma using non-negative matrix factorization (NMF, a statistical approach to separating the constituent parts of an object. See Discussion.)(Lee and Seung, 1999). Each of these reports has contributed to our basic understanding of pancreatic cancer biology, but a systematic evaluation of compartment–specific contributions is needed to identify subtypes that are broadly applicable across biologically and technically heterogeneous PDA patient cohorts.

We used ADVOCATE to assess compartment fraction from gene expression profiles of three independent cohorts: (a) UNC Chapel Hill (UNC, n = 125), (b) the International Cancer Genome Consortium (ICGC, PACA-AU RNA-Seq dataset, n = 93), and (c) The Cancer Genome Atlas (TCGA, PAAD dataset, n = 137) (see Methods for inclusion criteria). We reasoned that there are three potential sources of relevant heterogeneity in these data: 1) differences in the epithelium/stroma ratio in areas of frank carcinoma; 2) variation in the representation of uncharacterized tissue (e.g., normal pancreas ductal epithelium, pancreatitis, lymph nodes, etc.) in the bulk sample; and 3) technical differences (e.g. expression platform, library preparation method, etc.). We further reasoned that the contributions of the latter two sources should be effectively captured by the "Residuals" component of the 3-compartment implementation of ADVOCATE, since they would lead to gene distributions distinct from those representing stromal and epithelial tissues in the LCM datasets used for algorithm training. Subtracting the Residuals component helps control technical and biological heterogeneity independent of true compartment fraction and produces more accurate compartment-specific inferences for both fractional composition and gene expression profiles. Using this approach, we found that the epithelial and stromal fractions varied significantly between the cohorts with 46%, 67% and 55% epithelium for the

ICGC, UNC and TCGA cohorts, respectively (p < 0.001, one-way ANOVA) (Figure 4A). This highlights critical differences in composition between tumor collections curated with different inclusion criteria or enrichment practices.

We next focused on the expression level and compartment–specificity of the genes used by each of the three published PDA classification systems (Figure 4B) (Bailey et al., 2016; Collisson et al., 2011; Moffitt et al., 2015). We noted that the Bailey classifier was developed using Ensembl gene annotation and that the Immunogenic subtype includes numerous recombined immunoglobulin genes that are not designated in the NCBI annotation used for the CUMC dataset (Tables S17-20). Therefore, to assess the compartment–specificity of Bailey classifier genes, we used a version of the CUMC dataset that was remapped using the Ensembl GRCh37 gene annotation. Notably, we observed that the genes used to define the Classical, Basal-like, and Progenitor subtypes were heavily weighted towards epithelium–specific expression. Conversely, those used to define the Activated, Normal and Immunogenic subtypes were weighted towards the stroma. The Quasi-Mesenchymal (QM) and Squamous gene sets were well–expressed and represented a mixture of epithelial and stromal identity, consistent with a more poorly differentiated state. Finally, the majority of genes that define the Exocrine and ADEX subtypes exhibited very low expression in the LCM-RNA-Seq datasets, suggesting that their expression in bulk tissue is derived from cell types largely absent from our microdissected samples. Together, these data provide insight into the cellular compartments that contribute to molecular gene signatures built from bulk tumor tissue samples.

Transcriptional deconvolution improves functional classification across cohorts
An important feature of a robust classification system is its capacity to identify sample subsets that are functionally similar across independent datasets. Variable cohort composition and other confounding factors can potentially lead to the co-stratification of functionally unrelated samples. For example, application of the epithelial signature from Moffitt et. al. (hereafter referred to as Moffitt–E) to bulk profiles from TCGA identifies two groups of samples, just as with the original UNC dataset on which Moffitt–E was trained. However, gene set variance analysis of the two TCGA groups finds intermixing of the functions associated with the Basal–like and Classical groups (Figures 4C). Having

established that compartment-specific fractional representation varies substantially between these cohorts, we wondered whether removing the expression contributions of stromal cells would aid classification efforts.

To address this challenge, we used ADVOCATE to generate virtual epithelial and stromal expression profiles from the bulk samples of each PDA cohort (producing new datasets vUNC, vTCGA, and vICGC). In each case, virtual profiles displayed clear expression of established cell–specific marker genes (Figure 4D and Figure S5A, B). Notably, bulk samples were distributed between the corresponding virtual epithelium and virtual stroma samples by hierarchical clustering (Figures S5 C–E).

To address whether deconvolution improved cross–cohort consistency of molecular classifiers, we analyzed the underline{v}irtual underline{e}pithelial profiles from each cohort (veUNC, veTCGA, and veICGC) using the Moffitt–E epithelial classification signature. Notably, deconvolution of TCGA data led to a realignment in functional associations, resulting in two groups that could be clearly recognized as Basal-like and Classical, based on enriched gene sets (Figure 4E). Similar functional associations were observed when the Moffitt–E classifier was applied to our experimentally–derived epithelial LCM–RNA–Seq (CUMC LCM–E) dataset (Figure 4F). We also noted that application of the Moffitt–E classifier to the veICGC dataset revealed excellent alignment with the pancreatic progenitor and squamous subtypes described by (Bailey et al., 2016) (SMC = 0.91) (Table S6). Together, these data indicate that removal of stromal expression data from bulk tumor datasets can improve the functional similarity of the groups identified by classification systems.

Identification of Immune-rich and ECM-rich subtypes of PDA stroma

The profuse stromal desmoplasia of pancreatic ductal adenocarcinoma is a defining feature of this malignancy. Previously, Moffitt and colleagues used NMF (Lee and Seung, 1999) on a large number of bulk PDA expression profiles to infer a PDA stromal classifier that identified two subtypes, designated "Activated" and "Normal" (Moffitt et al., 2015). These subtypes were interpreted to reflect the biology of cancer associated fibroblasts and was associated with clinical outcome. Building from this, we sought to capture expression

signals from the stroma as a whole, including the many myeloid, lymphoid, endothelial and other cell types that are commonly present in the PDA microenvironment. To this end, we expanded the stromal LCM–RNA–Seq cohort described above to include samples from a total of 110 unique patients. NMF with consensus clustering identified two prominent molecular subtypes among these samples. Clear functional identities were established for these subtypes using gene set variance analysis (GSVA), leading to their designations as: an "Immune–rich" group characterized by numerous immune and interleukin signals; and an "ECM–rich" group, characterized by numerous extracellular matrix–associated pathways (Figure 5A). We next extracted a gene signature distinguishing these two stromal subtypes, making use of the compartment specificity analysis described above to filter for stroma-specific genes (see Supplementary Methods and Supplementary Tables 22 – 24). Application of this signature to the virtual stroma profiles yielded two prominent clusters, as reflected across the UNC, ICGC, and TCGA cohorts (Figures 5B–D). Critically, in each cohort, the two clusters were again characterized by their enrichment for gene sets associated with ECM deposition or immune processes, indicating a robust and consistent performance of this new, stroma-specific "CUMC–S" signature.

Epithelial and stromal subtypes are partially linked and associated with survival differences

Having determined the epithelial and stromal subtypes of all CUMC, UNC, ICGC, and TCGA samples, we were able to assess the degree of variation in the subtype composition of these datasets. Notably, we found that both the epithelial and stromal composition of each tumor cohort varied considerably across datasets. Within the epithelium, the Basal–like group comprised 29%, 41%, and 27% of cases in the veUNC, veICGC, and veTCGA cohorts, respectively (Figures 5 E–G), and 36% of our epithelial LCM–RNA–Seq profiles (Figure S6A). Within the stroma, the ECM–rich subtype comprised 62%, 52%, and 31% of cases in the vsUNC, vsICGC, and vsTCGA cohorts, respectively (Figures 5 E–G), and 47% of our stromal LCM–RNA–Seq samples (Figure S6A). These observations serve to further highlight the significant heterogeneity between independent collections of pancreatic tumor specimens.

We next sought to assess the associations of epithelial and stromal subtypes with survival outcomes. Examining the epithelial samples, we found that removing stromal gene expression with ADVOCATE improved the survival association between Classical and Basal–like tumors in all three bulk datasets, with a particularly strong effect on TCGA outcomes (Figures 6 A–C) where 45% of the samples were re-classified after deconvolution. For the stromal subtypes, we observed at least a trend towards reduced survival among ECM-rich tumors in all three datasets (a finding made more apparent by deconvolution); however, this only reached significance in the ICGC cohort (Figures 6D–F). Together, these data indicate that (i) differences in tumor composition between different large-scale gene expression datasets can affect the predictive power of established classifier signatures for PDA, and (ii) transcriptional deconvolution can help overcome this hurdle, improving the reproducibility of outcome prediction.

The existence of numerous paracrine signaling pathways whose activity is affected by oncogenic mutations implies that stromal transcriptional programs should be heavily influenced by epithelial identity (Laklai et al., 2016). We examined this corollary by ascertaining the association of epithelial and stromal subtypes in our experimental LCM dataset as well as in those from the virtual UNC, ICGC, and TCGA datasets. We found that in the ICGC and TCGA cohorts, the ECM-rich stroma subtype was preferentially associated with the Basal-like epithelial subtype; the UNC and CUMC cohorts trended in this direction but did not reach significance. However, a meta-analysis of the 393 samples from all four datasets yielded an Odds Ratio of 2.7 for the association of Basal-like epithelium and ECM-rich stroma (Figure S6B, random effects model: OR 2.7 [1.33 – 5.53], $p < 0.001$), indicating a partial association between epithelial and stromal compartments.

The imperfect alignment of the epithelial and stromal subtypes offered the possibility that combination subtypes might vary in their survival associations as compared to either compartment alone (Figures 6 G–I). Indeed, consideration of combined epithelial and stromal combination subtypes affected the outcome prediction, particularly in the case of the UNC cohort where combination subtyping of deconvolved samples found a particularly poor outcome for Basal–like/ECM–rich tumors relative to Classical/Immune–rich tumors (HR = 3.76 for combined subtyping vs. 2.11 for epithelial subtyping alone (Figure 6I vs. 6C).

Together, these data highlight the relationship between Basal-like epithelium with ECM-rich stroma in pancreatic cancer and the strong association of this combination with poor overall survival.

Discussion

The traditional understanding of genetic mutations as drivers of tumor development has led to a focus on the malignant compartment that is exemplified by the term "tumor purity", which regards the stroma as mere contamination. However, with the understanding that stromal cells play critical roles in both promoting and restraining pancreatic tumor progression (Neesse et al., 2015), the consensus view of the stromal compartment has shifted to that of a critical partner – or foil – to the malignant epithelium. Indeed, in some contexts the stroma can even play a dominant role, as epitomized by the success of stroma–targeted immunotherapy in treating aggressive cancers such as metastatic melanoma and non–small cell lung cancer. In this light, we sought to study the interplay of PDA epithelium and stroma in their native state, separated by LCM from otherwise intact samples, but matched by patient so that the reciprocal signals active in each compartment might be examined.

A key outcome of this work is to unify our understanding of molecular subtypes in pancreatic ductal adenocarcinoma. To do this, we first examined the properties of subtypes resulting from existing classification schemes, making use of compartment–fraction estimation function of ADVOCATE as well as our annotation of the expression levels and compartment–specificity of each gene in our LCM-RNA-Seq dataset. We noted the substantial heterogeneity of compartment fraction between the UNC, ICGC, and TCGA cohorts, and detailed how certain proposed subtypes may have emerged from variations in tissue composition. Indeed, the removal of stromal expression signals from the Moffitt–E signature resulted in the reclassification of nearly half the samples and improved discernment of the functional processes associated with the Classical and Basal–like subtypes in each cohort.

Functional similarity in subtypes is an important concept for tumor classification. It is fully possible, even commonplace, to generate functionally distinct groups when applying the same classification signature to different sample cohorts. Variation in technical features such as library preparation method or expression platform, as well as biological heterogeneity can have an outsized impact on sample classification systems. Thus, in our effort to establish a novel classification system for PDA stroma, we placed the greatest emphasis on the reproducibility of molecular phenotypes across multiple cohorts. Following this process, we observed with great interest the emergence of two prominent molecular subtypes in the stroma with pronounced enrichment for two different aspects of stromal biology: ECM deposition and remodeling versus immune–related processes. The dichotomy between these two functional groups emphasizes the now well–described role that the inflammatory microenvironment plays in modulating the local immune response to PDA.

Examining both the epithelial and stromal subtypes together in combination across 393 pancreatic tumor specimens led to the finding that there is considerable heterogeneity in cross-compartmental dependencies across the 393 specimens examined, with some evidence of dependence between the global stromal transcriptional program and its epithelial counterpart. An expansive body of literature has accumulated describing myriad signaling interactions between the epithelium and stroma of pancreatic cancer (Bailey and Leach, 2012). Furthermore, mutation–driven epithelial signaling programs such as those induced by oncogenic K–ras have profound impacts on stromal cell biology (Pylayeva-Gupta et al., 2012; Ying et al., 2012). However, K–ras itself is mutated in 95% of PDA cases, so it cannot alone explain the differences between the prominent stromal subtypes. By examining all four tumor cohorts, we found a strong association between an ECM–rich stroma and Basal-like epithelium. The latter finding corroborates the concept that epithelial traits promoting dedifferentiation in PDA, such as the loss of SMAD4 expression, may in fact shape a more matricellular, and likely more rigid, stromal phenotype (Laklai et al., 2016). This association was less prominent in the UNC cohort and not detectable in our smaller group of LCM tumors. Combined evidence from all cohorts, however, generally supports the idea of a modest cross-compartment dependence. Further studies will be needed to better understand the dynamics of cross-compartment subtypes in PDA by taking into account additional variables such as mutation status or

environmental/epidemiological factors. To this end, we provide two novel tools for the PDA field: a compartment-specific gene signature that may discriminate between ECM-rich and Immune-rich stromas as well as the ADVOCATE framework which may help to assess the compartment composition of PDA bulk expression profiles and provide virtually purified expression for downstream analyses.

It is noteworthy that patient stratification according to both the epithelial and stromal combination subtype is strongly associated with patient outcome. Specifically, tumors with Basal–like epithelium and an ECM–rich stroma have a substantially worse overall survival compared to those with a Classical epithelium and Immune–rich stroma (HR = 3.76, 3.81, and 2.63 for UNC, ICGC, and TCGA, respectively). This effect size compares favorably to other known single variables in pancreatic cancer biology, including lymph node status (HR = 1.5), postoperative CA19-9 level (HR = 3.6) or the number of high penetrance driver genes (HR = 1.4)(Berger et al., 2008; Yachida et al., 2012) while also providing a biological context. Unfortunately, differences in the clinicopathological data reported for each cohort precluded a more sophisticated multivariate model. Nonetheless, we expect that this approach to subtyping will have immediate applications, for example, in interpreting the results of small-scale clinical trials where random inequalities of molecular subtypes could dramatically affect the expected survival between groups or relative to historical controls.

The generation of virtual compartment–specific expression profiles from bulk profiles is a unique tool in the computational biology field. Other algorithms often provide a means to infer compartment expression from a large group of samples based on linear modeling of putative marker gene expression (Gill et al., 2014; Kuhn et al., 2011), which depends on the availability of truly cell-type specific marker genes being expressed in an essentially binary manner. By contrast, ADVOCATE uses a machine learning approach to model the expression distribution for hundreds of genes discriminating between the training conditions (i.e. epithelium and stroma), thus achieving substantial redundancy in the prediction and generating results that are consistent with maximum likelihood approaches. This both informs its compartment fraction prediction function and enables the deconvolution function to be performed on an individual–sample basis (provided appropriate reference standards for the sample are available from which scaled normalized expression may be obtained).

The ADVOCATE framework may easily be expanded to include additional compartments or cell types, given suitable experimental data. However, it cannot generate virtual profiles for any compartments that have not been sampled; rather, it can only estimate the fraction of residual tissue yielding gene expression contributions that cannot be effectively modeled by the compartment-specific probability densify functions. In practice, this "Residuals" compartment helps address both issues of technical variability, such as those arising from different profiling platforms, as well as biological variability, such as the one resulting from infiltration of unknown cellular lineages. Thus, even when the Residual compartment fraction is high, this helps optimize the contribution of the compartments that are well modeled by the training LCM data and also optimally infer the corresponding gene expression profiles, thus minimizing the effect of confounding factors related to poorly represented tissue infiltration. The 3-compartment version of ADVOCATE is therefore useful as a noise-reduction tool, as also exemplified in Figure 4A. Considering the technical challenges of performing LCM–RNA–Seq and other enrichment methods, we expect this computational approach will reduce the need for experimental sample manipulation in the future. Moreover, while we presented implementations of ADVOCATE specific to pancreatic and breast cancer, we expect that the algorithm, which requires very limited sample sizes for training, can be generally applied to model subpopulations across the cancer field. This will provide an important framework for handling the cellular heterogeneity of cancer and further expanding the utility of large–scale gene expression profile collections.

To facilitate both understanding of the technical details of algorithm implementation and to allow the research community to make full use of this tool, we are making the source code available on Bioconductor in conjunction with publication of this manuscript.

Experimental procedures

The information provided here is a succinct summary of the experimental procedures. Extensive and detailed information is provided in supplementary information.

Samples studied

Information is provided from a total of 129 PDA patients who underwent surgery at the Columbia Pancreas Center. Of these, LCM–RNA–Seq data are presented for both the epithelium and stroma of 64 cases, for the stroma of an additional 59 cases, bulk RNA–Seq data are presented for 9 cases, and both LCM and bulk RNA–Seq are presented for 6 cases. Patients provided surgical informed consent which was approved by a local ethics committee (IRB # AAAB2667). Samples were frozen intraoperatively by the Columbia University Tumor Bank.

Laser capture microdissection and RNA sequencing

Cryosections of OCT–embedded tissue blocks were transferred to PEN membrane glass slides and stained with cresyl violet acetate. Adjacent sections were H&E stained for pathology review. Laser capture microdissection was performed on a PALM MicroBeam microscope (Zeiss), collecting at least 1000 cells per compartment. RNA was extracted and libraries prepared using the Ovation RNA-Seq System V2 kit (NuGEN). Libraries were sequenced to a depth of 30 million, 100bp, single-end reads.

Computational methods

The ADVOCATE framework is described in detail in supplementary materials. An R package for implementing ADVOCATE has been developed and will be made generally available in Bioconductor in conjunction with publication of this manuscript.

Accession numbers

Expression data were deposited into publically accessible databases with accession numbers to be provided prior to publication.

Author Contributions

K.P.O., A.C., and M.B. conceived the overall framework and supervised the experimental and methodological components of the analyses. H.C.M. and S.R.H. developed experimental methodologies and performed LCM–RNA–Seq. J.H. and M.B. developed the ADVOCATE algorithm and wrote related software. J.Z. provided guidance on deconvolution methods. J.H. and H.C.M. curated datasets. J.H., H.C.M., M.B., and K.P.O. analyzed and interpreted results. J.A.C., H.H., A.A, and T.S. contributed to tissue procurement and banking, and A.A. performed preliminary pathology assessments of banked samples. A.C.I. and A.R.S. marked regions of each sample to be laser capture microdissected on adjacent sections. A.C.I. annotated the histopathological features of each sample with further review of select cases by K.P.O. Regulatory oversight for the use of human samples/data was provided by H.H. and J.G. The manuscript was written by J.H., H.C.M., K.P.O., and A.C. with editing by M.B., A.C.I., S.R.H, and J.G. Figures were constructed by J.H., H.C.M., and K.P.O. Supervision and oversight were provided by K.P.O., A.C., and M.B.

Conflicts

A.C. is a founder and shareholder of DarwinHealth Inc. and a member of the Tempus Inc. SAB and shareholder. K.P.O. is a member of the SAB for Elstar Therapeutics. Columbia University is a shareholder of DarwinHealth Inc.

Acknowledgements

The authors would like to thank Federico M. Giorgi and Alexander Lachmann for advice on computational methods, and Richard Moffitt for valuable critique of the manuscript. This work was supported by the National Cancer Institute (NCI) Cancer Target Discovery and Development program (1U01CA168426 to A.C.), NCI Research Centers for Cancer Systems Biology Consortium (1U54CA209997 to A.C. and K.P.O.), NCI Outstanding Investigator Award (R35CA197745-02 to A.C.) and NCI Research Project Grant (R01CA157980 to K.P.O.). Financial support was also provided by the Columbia University Pancreas Center. H.C.M. received support from a Mildred Scheel Postdoctoral Fellowship (Deutsche Krebshilfe). P.E.O. received support from the NIH NCATS (KL2TR001874).

Figure Legends

Figure 1. Compartment–specific gene expression profiling of pancreatic tumors

(**A**) Images of Cresyl Violet stained human PDA frozen sections before and after laser capture microdissection of malignant epithelial and adjacent stromal cells. (**B**) RIN values for RNA samples derived from the indicated compartment (N = 60 each). (**C**) Number of genes and transcripts detected at > 1 FPKM in the samples from (B). (**D**) Principal component analysis of the profiles from (C). 60 pairs of epithelial and stromal LCM samples. Color graduation shows pairing of samples from the same tumor. Three samples discussed later are labeled. (**E**) Heatmap showing the expression of marker genes for the indicated cell types in each sample. (**F, G**) Validation of genes predicted as epithelium–specific (F) or stroma–specific (G) based on mRNA expression. Bar height and color shading reflect the certainty (t-statistic) of differential expression. The box color below each bar summarizes results of immunohistochemistry on PDA sections from the Human Protein Atlas (HPA). IHC staining pattern was categorized as strongly or weakly supportive of the predicted compartment (blue/red), indeterminate (grey), absent (white), or opposite the predicted pattern (black). (**H**) An example epithelium–specific gene, LGALS4, showed a protein staining pattern that was strongly consistent with its mRNA expression (at left). Blue and red arrows indicate PDA epithelium and stroma, respectively. (**I**) LGALS1 exhibited a highly stroma-specific expression pattern.

Figure 2. ADVOCATE framework and computational validation

(**A**) Outline of the ADVOCATE framework where LCM–RNA–Seq profiles are used to model genome–wide gene expression distributions for each gene across compartments. Bulk tissue profiles can be deconvolved using compartment–specific, gene–wise probabilities, yielding two products: i) predicted compartment fractions in the bulk tumor and ii) virtual gene expression profiles for the epithelium and stroma of that sample. Numbers indicate steps described in main text. (**B**). Compartment fraction analysis performed with ADVOCATE (green points) on a computationally generated (synthetic) set of epithelial and stromal profiles mixed together *in silico* in varying proportions. (**C**) Compartment fraction analysis performed with ADVOCATE (green/yellow points) on a semi-synthetic set of samples generated by mixing actual experimental pairs of LCM expression profiles together in varying proportions, for each of 60 tumors. The yellow lines highlight three outlier

samples. (**D**) Leave-one-out cross validation of compartment fraction using ADVOCATE where the epithelial (blue) and stromal (red) fractions, are predicted for each individual pair of LCM–RNA–Seq profiles after training the model using the remaining 59 pairs. The outliers from (C) are indicated as yellow bars. (**E**) Power analysis for training the ADVOCATE algorithm. Plot shows 95% (dark blue) and 85% (light blue) percentile of prediction errors relative to sample sizes based on LOOCV analysis. Errors decreases quickly as sample size increases, stabilize after ~ 20 samples, and reach a minimum at 60 samples.

Figure 3. Experimental validation of ADVOCATE

(**A**) Predicted compartment fractions for CUMC epithelial (left) and stromal (right) LCM–RNA–Seq profiles using ADVOCATE (red), ESTIMATE (blue), PSEA (green), DSA (purple) and deconRNAseq (orange). (**B**) Example of nuclei counting for the epithelial (teal dots) and stromal (orange dots) compartments. Nuclei per compartment were divided by the total to yield the compartmental fraction. (**C-E**) Comparison between tumor epithelium content estimated by (**C**) nuclei counting vs. area estimation, (**D**) nuclei counting vs. ADVOCATE and (**E**) area estimation vs. ADVOCATE for 15 tumors from the Columbia University Medical Center (CUMC) collection. MAPE denotes **M**ean **A**bsolute **P**ercentage **E**rror as a measure of prediction accuracy. (**F**) Correlation of normalized gene expression between experimental LCM–RNA–Seq profiles and: permuted bulk profiles (left, negative control); bulk expression profiles (middle); and virtual epithelial and stroma profiles purified from bulk using ADVOCATE (right). In each pair, blue indicates epithelium and red indicates stroma. (**G**) Heatmap showing expression of indicated marker genes in virtual epithelial and stromal profiles derived from the 15 bulk tumors described in (C). (**H**) Hierarchical clustering and (**I**) PCA of 15 bulk CUMC tumors and their virtual compartment–specific derivatives.

Figure 4. Analysis and classification of pancreatic tumor cohorts and classifiers

(**A**) Compartment fraction analysis of pancreatic tumors from the ICGC (blue), UNC (red), and TCGA (grey) cohorts. The 3-compartment implementation of ADVOCATE was used for fraction estimation to reduce noise from technical and non-tumor sources, and the epithelial and stromal estimate were then scaled to sum up to 1. (**B**) Analysis of genes comprising three published classifiers for PDA using LCM–RNA–Seq data. Upper panel depicts the

distribution of t-statistics for each of the indicated classifier gene lists as derived from a paired differential gene expression analysis between the epithelium and stroma. Positive values indicate stromal enrichment. Lower panel depicts the geometric mean expression in fragments per million mapped fragments (FPM) for the genes from each classifier gene list across all LCM–RNA–Seq samples. (**C**) Heatmap depicts the enrichment of each TCGA bulk sample for the indicated gene sets. Gene sets indicated functions associated with Basal-like (red) or Classical (blue) traits. (**D**) Heatmap depicts the enrichment of expression of indicated marker genes in deconvolved virtual epithelial and stromal profiles from the TCGA cohort. (**E**) Hierarchical clustering TCGA bulk tumors (black) and their virtual epithelial (blue) and stromal (red) derivatives. (**F,G**) Heatmaps of the top 30 differentially expressed genes between two groups that were obtained by clustering virtual epithelial TCGA (F) and CUMC LCM (G) profiles using the Moffitt–E classifier. GSVA scores for indicated gene sets are represented at the top. Tumors with Basal–like (red bar) or Classical (blue bar) traits, respectively, can be detected in both virtual and LCM tumor cohorts.

Figure 5. Stromal Subtyping of deconvolved external cohorts

(**A-D**) Heatmaps of the top 30 DEG between groups obtained by clustering stromal LCM–RNA–Seq samples from CUMC tumors (A), and virtual stromal (vs) profiles from the UNC (B), ICGC (C) and TCGA (D) cohorts, respectively. Clustering was based on the expression of a signature derived from stromal LCM profiles from 110 individual patients (CUMC-S classifier, see Supplementary Methods). Top section of heat-map depicts GSVA scores per sample for indicated gene sets. In each virtual stroma dataset, two groups were identified, one with features indicating elevated extracellular matrix deposition and remodeling ("ECM–rich", purple) and another enriched in various immune and interleukin pathways ("Immune-rich", green). (**E-G**) Multilayered donut plots showing (i) the alignment of epithelial with stromal subtypes for each tumor in each cohort and (ii) the proportion of each epithelial subtype. Separate pie charts summarize the proportion of stromal subtypes per cohort.

Figure 6. Combined epithelial and stromal subtypes associate with overall survival

(**A-C**) Kaplan-Meier (KM) survival analysis of patients with resected PDA from the indicated cohorts shows that the detection of a differential prognosis among the epithelial subtypes is enhanced by transcriptional deconvolution across all cohorts. Thick red and blue lines show the survival curves of tumor groups identified among deconvolved virtual epithelium profiles, while thinner light red and blue lines represent those groups detected by subtyping bulk expression profiles. Below each KM-plot, the hazard ratios (HR) of a Basal-Like tumor status from a Cox proportional hazards model (CPHM) and their 80% (blue), 90% (yellow) and 95% (orange) confidence interval are compared between virtual epithelial (ve) and bulk expression classification, respectively. (**D-F**) Kaplan-Meier (KM) survival analysis depicts overall survival relative to stromal subtype. Classifications derived from bulk tumor profiles are shown with thin lines while those derived from virtual stroma profiles are shown with thick lines. Below each KM-plot, HRs for an ECM-rich stromal status from a CPHM are compared between virtual stromal (vs) and bulk classification, respectively, as described above. Stromal subtypes are only associated with outcome in the ICGC cohort with ECM-rich tumors having a worse prognosis. (**G-I**) KM survival analysis of combined epithelial and stromal subtypes in the indicated cohorts. Red lines indicate Basal-like tumors with an ECM-rich stroma while blue lines indicate Classical tumors with an Immune-rich stroma. All other tumors are represented as a grey line. Pie charts summarize the proportion of each category per cohort. HRs from a CPHM demonstrate that the combined Basal-like/ECM-rich subtype is strongly associated with reduced survival in PDA patients in all three cohorts.

References

Abu-Alainin, W., Gana, T., Liloglou, T., Olayanju, A., Barrera, L. N., Ferguson, R., Campbell, F., Andrews, T., Goldring, C., Kitteringham, N.*, et al.* (2016). UHRF1 regulation of the Keap1-Nrf2 pathway in pancreatic cancer contributes to oncogenesis. The Journal of pathology *238*, 423-433.

Adiconis, X., Borges-Rivera, D., Satija, R., DeLuca, D. S., Busby, M. A., Berlin, A. M., Sivachenko, A., Thompson, D. A., Wysoker, A., Fennell, T.*, et al.* (2013). Comparative analysis of RNA sequencing methods for degraded or low-input samples. Nat Methods *10*, 623-629.

Aran, D., Sirota, M., and Butte, A. J. (2015). Systematic pan-cancer analysis of tumour purity. Nat Commun *6*, 8971.

Bailey, J. M., and Leach, S. D. (2012). Signaling pathways mediating epithelial-mesenchymal crosstalk in pancreatic cancer: Hedgehog, Notch and TGFbeta. In Pancreatic Cancer and Tumor Microenvironment, P.J. Grippo, and H.G. Munshi, eds. (Trivandrum (India)).

Bailey, P., Chang, D. K., Nones, K., Johns, A. L., Patch, A. M., Gingras, M. C., Miller, D. K., Christ, A. N., Bruxner, T. J., Quinn, M. C.*, et al.* (2016). Genomic analyses identify molecular subtypes of pancreatic cancer. Nature *531*, 47-52.

Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A. A., Kim, S., Wilson, C. J., Lehar, J., Kryukov, G. V., Sonkin, D.*, et al.* (2012). The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature *483*, 603-307.

Berger, A. C., Garcia, M., Jr., Hoffman, J. P., Regine, W. F., Abrams, R. A., Safran, H., Konski, A., Benson, A. B., 3rd, MacDonald, J., and Willett, C. G. (2008). Postresection CA 19-9 predicts overall survival in patients with pancreatic cancer treated with adjuvant chemoradiation: a prospective validation by RTOG 9704. J Clin Oncol *26*, 5918-5922.

Biankin, A. V., Waddell, N., Kassahn, K. S., Gingras, M.-C., Muthuswamy, L. B., Johns, A. L., Miller, D. K., Wilson, P. J., Patch, A.-M., Wu, J.*, et al.* (2012). Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. Nature *491*, 399-405.

Carter, S. L., Cibulskis, K., Helman, E., McKenna, A., Shen, H., Zack, T., Laird, P. W., Onofrio, R. C., Winckler, W., Weir, B. A.*, et al.* (2012). Absolute quantification of somatic DNA alterations in human cancer. Nat Biotechnol *30*, 413-421.

Collisson, E. A., Sadanandam, A., Olson, P., Gibb, W. J., Truitt, M., Gu, S., Cooc, J., Weinkle, J., Kim, G. E., Jakkula, L.*, et al.* (2011). Subtypes of pancreatic ductal adenocarcinoma and their differing responses to therapy. Nat Med *17*, 500-503.

Gill, B. J., Pisapia, D. J., Malone, H. R., Goldstein, H., Lei, L., Sonabend, A., Yun, J., Samanamud, J., Sims, J. S., Banu, M.*, et al.* (2014). MRI-localized biopsies reveal subtype-specific differences in molecular and cellular composition at the margins of glioblastoma. Proc Natl Acad Sci U S A *111*, 12550-12555.

Gong, T., and Szustakowski, J. D. (2013). DeconRNASeq: a statistical framework for deconvolution of heterogeneous tissue samples based on mRNA-Seq data. Bioinformatics *29*, 1083-1085.

Hruban, R. H. P., M.B.; Klimstra, D.S. (2007). Tumor of the Pancreas, Vol 4, (Washington DC: American Registry of Pathology).

Jacobetz, M. A., Chan, D. S., Neesse, A., Bapiro, T. E., Cook, N., Frese, K. K., Feig, C., Nakagawa, T., Caldwell, M. E., Zecchini, H. I.*, et al.* (2012). Hyaluronan impairs vascular function and drug delivery in a mouse model of pancreatic cancer. Gut.

Jones, S., Zhang, X., Parsons, D. W., Lin, J. C.-H., Leary, R. J., Angenendt, P., Mankoo, P., Carter, H., Kamiyama, H., Jimeno, A*., et al.* (2008). Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. Science *321*, 1801-1806.

Kuhn, A., Thu, D., Waldvogel, H. J., Faull, R. L., and Luthi-Carter, R. (2011). Population-specific expression analysis (PSEA) reveals molecular changes in diseased brain. Nature methods *8*, 945-947.

Laklai, H., Miroshnikova, Y. A., Pickup, M. W., Collisson, E. A., Kim, G. E., Barrett, A. S., Hill, R. C., Lakins, J. N., Schlaepfer, D. D., Mouw, J. K*., et al.* (2016). Genotype tunes pancreatic ductal adenocarcinoma tissue tension to induce matricellular fibrosis and tumor progression. Nat Med *22*, 497-505.

Lee, D. D., and Seung, H. S. (1999). Learning the parts of objects by non-negative matrix factorization. Nature *401*, 788-791.

Moffitt, R. A., Marayati, R., Flate, E. L., Volmar, K. E., Loeza, S. G. H., Hoadley, K. A., Rashid, N. U., Williams, L. A., Eaton, S. C., Chung, A. H., and Others (2015). Virtual microdissection identifies distinct tumor-and stroma-specific subtypes of pancreatic ductal adenocarcinoma. Nat Genet *47*, 1168-1178.

Neesse, A., Algul, H., Tuveson, D. A., and Gress, T. M. (2015). Stromal biology and therapy in pancreatic cancer: a changing paradigm. Gut *64*, 1476-1484.

Oberstein, P. E., and Olive, K. P. (2013). Pancreatic cancer: why is it so hard to treat? Therapeutic advances in gastroenterology *6*, 321-337.

Oh, E.-Y., Christensen, S. M., Ghanta, S., Jeong, J. C., Bucur, O., Glass, B., Montaser-Kouhsari, L., Knoblauch, N. W., Bertos, N., Saleh, S. M. I*., et al.* (2015). Extensive rewiring of epithelial-stromal co-expression networks in breast cancer. Genome Biology *16*, 128.

Olive, K. P., Jacobetz, M. A., Davidson, C. J., Gopinathan, A., McIntyre, D., Honess, D., Madhu, B., Goldgraben, M. A., Caldwell, M. E., Allard, D*., et al.* (2009). Inhibition of Hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. Science *324*, 1457-1461.

Pontén, F., Jirström, K., and Uhlen, M. (2008). The Human Protein Atlas-a tool for pathology. J Pathol *216*, 387-393.

Provenzano, P. P., Cuevas, C., Chang, A. E., Goel, V. K., Von Hoff, D. D., and Hingorani, S. R. (2012). Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma. Cancer Cell *21*, 418-429.

Pylayeva-Gupta, Y., Lee, K. E., Hajdu, C. H., Miller, G., and Bar-Sagi, D. (2012). Oncogenic Kras-Induced GM-CSF Production Promotes the Development of Pancreatic Neoplasia. Cancer Cell *21*, 836-847.

Rhim, A. D., Mirek, E. T., Aiello, N. M., Maitra, A., Bailey, J. M., McAllister, F., Reichert, M., Beatty, G. L., Rustgi, A. K., Vonderheide, R. H*., et al.* (2012). EMT and dissemination precede pancreatic tumor formation. Cell *148*, 349-361.

Rhim, A. D., Oberstein, P. E., Thomas, D. H., Mirek, E. T., Palermo, C. F., Sastra, S. A., Dekleva, E. N., Saunders, T., Becerra, C. P., Tattersall, I. W*., et al.* (2014). Stromal elements act to restrain, rather than support, pancreatic ductal adenocarcinoma. Cancer Cell *25*, 735-747.

Ross, D. T., Scherf, U., Eisen, M. B., Perou, C. M., Rees, C., Spellman, P., Iyer, V., Jeffrey, S. S., Van de Rijn, M., Waltham, M*., et al.* (2000). Systematic variation in gene expression patterns in human cancer cell lines. Nat Genet *24*, 227-235.

Shanker, S., Paulson, A., Edenberg, H. J., Peak, A., Perera, A., Alekseyev, Y. O., Beckloff, N., Bivens, N. J., Donnelly, R., Gillaspy, A. F*., et al.* (2015). Evaluation of commercially

available RNA amplification kits for RNA sequencing using very low input amounts of total RNA. J Biomol Tech *26*, 4-18.

Siegel, R. L., Miller, K. D., and Jemal, A. (2017). Cancer Statistics, 2017. CA Cancer J Clin *67*, 7-30.

Song, S., Nones, K., Miller, D., Harliwong, I., Kassahn, K. S., Pinese, M., Pajic, M., Gill, A. J., Johns, A. L., Anderson, M.*, et al.* (2012). qpure: A Tool to Estimate Tumor Cellularity from Genome-Wide Single-Nucleotide Polymorphism Profiles. PLoS One *7*, e45835.

Vonderheide, R. H., and Bayne, L. J. (2013). Inflammatory networks and immune surveillance of pancreatic carcinoma. Curr Opin Immunol *25*, 200-205.

Waddell, N., Pajic, M., Patch, A.-M., Chang, D. K., Kassahn, K. S., Bailey, P., Johns, A. L., Miller, D., Nones, K., Quek, K.*, et al.* (2015). Whole genomes redefine the mutational landscape of pancreatic cancer. Nature *518*, 495-501.

Witkiewicz, A. K., McMillan, E. A., Balaji, U., Baek, G., Lin, W. C., Mansour, J., Mollaee, M., Wagner, K. U., Koduru, P., Yopp, A.*, et al.* (2015). Whole-exome sequencing of pancreatic cancer defines genetic diversity and therapeutic targets. Nature communications *6*, 6744.

Yachida, S., White, C. M., Naito, Y., Zhong, Y., Brosnan, J. A., Macgregor-Das, A. M., Morgan, R. A., Saunders, T., Laheru, D. A., Herman, J. M.*, et al.* (2012). Clinical significance of the genetic landscape of pancreatic cancer and implications for identification of potential long-term survivors. Clin Cancer Res *18*, 6339-6347.

Ying, H., Kimmelman, A. C., Lyssiotis, C. A., Hua, S., Chu, G. C., Fletcher-Sananikone, E., Locasale, J. W., Son, J., Zhang, H., Coloff, J. L.*, et al.* (2012). Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. Cell *149*, 656-670.

Yoshihara, K., Shahmoradgoli, M., Martínez, E., Vegesna, R., Kim, H., Torres-Garcia, W., Treviño, V., Shen, H., Laird, P. W., Levine, D. A.*, et al.* (2013). Inferring tumour purity and stromal and immune cell admixture from expression data. Nat Commun *4*, 2612.

Zhong, Y., Wan, Y. W., Pang, K., Chow, L. M., and Liu, Z. (2013). Digital sorting of complex tissues for cell type-specific gene expression profiles. BMC bioinformatics *14*, 89.