1

Transposable elements are important contributors to standing variation in gene expression in *Capsella grandiflora*

Jasmina Uzunović[1*], Emily B. Josephs[2,3], John R. Stinchcombe[1], Stephen I. Wright[1]

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Canada

[2]Department of Evolution and Ecology, University of California, Davis, California, USA

[3]Center for Population Biology, University of California, Davis, California, USA.

* Corresponding author

E-mail: jasmina.uzunovic@mail.utoronto.ca

Abstract

Transposable elements (TEs) make up a significant portion of eukaryotic genomes, and thus are important drivers of genome evolution. However, the evolutionary forces controlling TE copy number and the extent to which TEs affect phenotypic variation on a genome-wide scale are still unclear. We characterised TE insertion polymorphism and its effects on gene expression in 124 whole genome sequences from a single population of *Capsella grandiflora*. The frequency of insertions was negatively correlated with distance to genes, as well as density of conserved non-coding elements, suggesting that the negative effects of TEs on gene regulation are important in limiting their abundance. Rare TE variants strongly influence gene expression variation, predominantly through downregulation. In contrast, rare single nucleotide polymorphisms (SNPs) contribute equally to up- and down-regulation, but have a weaker effect. Taken together, these results imply that TEs are a significant contributor to gene expression variation and can be more likely than rare SNPs to cause extreme changes in gene expression.

Author Summary

Transposable elements (TEs), mobile DNA elements with the ability to excise from the genome and reinsert in new locations, are important components of genomic diversity. Due to their abundance and mobility, TEs play an influential role in genomic evolution, often deleterious. Here we show that TEs in a population of the plant *Capsella grandiflora* are most deleterious when they insert in genic and regulatory regions. We find that TEs indeed are associated with unusual levels of gene expression, predominantly decreased expression. Furthermore, this effect is stronger than the association of single nucleotide polymorphisms with

44   gene expression variation, highlighting the importance of TE contribution to the maintenance of

45   expression variation.

Introduction

Transposable elements are one of the most abundant and pervasive types of selfish

genetic element, making up over 80% of the genome in some species [1]. They experience

selective pressure to increase their own numbers, and at least in some species their abundance is

thought to reflect a balance between their rate of increase through transposition and the action of

selection on the host due to their deleterious fitness effects [2,3]. Although the population

dynamics of TEs have been characterized in a handful of model systems, our understanding of

the main selective factors controlling TE abundance, and the relative importance of TEs vs. other

sources of genetic variation (such as SNPs) in maintaining phenotypic variation remains limited.

Population studies of TEs in the compact genomes of *Drosophila melanogaster*,

Arabidopsis thaliana, and *Caenorhabditis elegans* have shown that most euchromatic insertions

are at very low population frequency [4–8]. A bias towards rare insertions suggests that

pervasive purifying selection is acting against TEs in these species. There are two main

hypotheses for why TEs are subject to purifying selection [3,9]. First, ectopic recombination

between TEs on non-homologous chromosomes or distinct locations on the same chromosome

can lead to aberrant chromosomal rearrangements, which are costly for fitness [10,11].

Alternatively, insertions in or near genic regions can negatively affect gene function, regulation,

and expression [12].

Studies in *D. melanogaster* populations provide evidence of selection acting against both

ectopic recombination between TEs [4] and the deleterious effects of gene disruption [13,14]. In

A. thaliana, distance to genes seems to be more important than ectopic recombination in

mediating selection against new insertions, consistent with the gene disruption hypothesis

playing a primary role [6,15]. TE abundance is negatively correlated with gene density,

suggesting TEs are harmful when they insert near protein-coding genes. Similarly, in *C. elegans*

TEs are enriched in regions characterised by low gene density and high recombination rate,

suggesting proximity to genes is a more important factor in TE survival as opposed to ectopic

recombination [8]. However, both *A. thaliana* and *C. elegans* are highly self-fertilizing, and

elevated homozygosity may minimize the effects of ectopic recombination [15–17]. Thus, while

we have preliminary evidence about the relative importance of ectopic recombination and gene

disruption in the regulation of TE abundance, it is difficult to generalize about their relative

importance with so few studies, especially in outcrossing species.

Given the evidence for selective pressure against gene disruption in a number of systems

[6,14,18], TEs may be a significant source of genetic variation for fitness-related phenotypes,

and can have a major impact on the evolution of the rest of the genome, especially functional

elements such as genes or conserved regulatory regions. Active TEs initiate mutations with

diverse molecular effects as they move around the genome. Most of these are deleterious, such as

disrupting genes or gene regulatory elements [19], causing structural rearrangements, and

silencing or decreasing expression of nearby genes through the spread of host-induced epigenetic

silencing of TEs [6,7,14,20,21]. However, TEs can also contribute putatively adaptive mutations

such as gene duplications, new gene regulatory elements, and selective regulation of gene

expression under stressful conditions [22–24]. Despite our knowledge about the many specific

ways in which different TEs can impact genes, there are few studies examining the extent to

which TEs as a class contribute to genetic and phenotypic variation.

A genome-wide survey of eQTL loci from a single large population of the self-

incompatible outcrossing plant *Capsella grandiflora,* shows around a third of genes are

associated with local eQTLs (those present within 5 kb of the gene) [25]. The variants associated

with gene expression are rarer than expected under neutrality, and their frequency is negatively

correlated with the size of their effect on gene expression [25], consistent with purifying

selection playing a large role in genetic variation for gene expression in *C. grandiflora*.

However, Josephs and colleagues [25] looked solely at SNPs, which are only a subset of the total

genetic variation that exists, and this system provides an excellent opportunity to explore the

relative role of 'classical mutation' vs. the importance of transposable element activity in

maintaining genetic variation in a natural population.

In this study, we examined TE polymorphism and its effects on expression variation

using the same population studied by Josephs et al. We identify insertions across all individuals

and explore the relationship between TE population frequency, rates of crossing over, and

density of coding sites and non-coding conserved sequences. We test the effect of TEs on gene

expression variation using genome-wide expression data. We then performed the same tests on

rare SNPs to compare the relative role of the two types of genetic variants.

Results and discussion

We combined high-confidence TE calls from 124 *C. grandiflora* plants using the

RelocaTE2 pipeline [26]. In total, we found 95,740 TEs in the population with an average of

2468 insertions per individual and a median of 2675 insertions per individual (range of 497 to

3131). In a survey of 216 *A. thaliana* accessions, there were 23,095 unique (non-reference) TE

variants (with a mean of 1657 per accession) [7]. The higher abundance of TEs in the *C.*

grandiflora population could reflect higher TE activity in outcrossers vs selfers leading to more

low frequency insertions, since the *A. thaliana* study did not include insertions shared with the

reference genome) [27,28].

116   The frequency spectrum of TEs in the population shows a major skew towards low-

frequency insertions compared to 4-fold degenerate SNPs, with a strong enrichment of insertions

present in only one individual (Fig 1A). The trend of mostly low-frequency insertions is

consistent with observations from other organisms and indicates there is strong negative

selection acting against TEs [5–8,29,30].

Fig 1. Patterns of TE insertion frequency from population data. (A) Frequency spectrum of

TEs (red) and 4-fold SNPs (blue). (B) Relationship between TE frequency and distance to the

closest gene. The first point represents the average frequency of all TEs which are within a

protein-coding gene. Subsequent points average the frequency of all TEs in 500 bp windows

from the closest gene. The blue line shows a linear model fit through the data. r = 0.94, linear

coefficient has a p < 0.0001. C and D Relationship between average TE frequency and

proportion of (C) coding DNA and (D) conserved non-coding sequence (CNS) in 100 kb

windows across the genome. The blue curve shows a quadratic model fit. Spearman's ρ = -0.562

in (C) and -0.619 in (D).

Selection against gene disruption regulates TE numbers

There are two main hypotheses for why TEs are deleterious in the genome: (i) they can

cause ectopic recombination between two non-homologous insertions or (ii) their insertion can

disrupt gene function or expression. In the first case, TEs should be more deleterious in areas of

high recombination rate, under the assumption that ectopic recombination rate mirrors the

homologous crossover rate. We used estimates of recombination rate in bins of 100 kb across the

genome and compared to the average TE population frequency of all insertions present in that

bin (S5 Fig). There is no significant correlation between TE frequency and recombination rate

across the genome, suggesting ectopic recombination does not play a large role in outcrossing *C.*

grandiflora, consistent with results from the highly selfing *A. thaliana* [15]. Ectopic

recombination is expected to be of greater importance in outcrossing species due to their higher

heterozygosity; a TE is more likely to recombine non-homologously if it does not have a

homologous partner to pair with. However, there is no evidence for a larger role of ectopic

recombination in this outcrossing species than in a related selfing species. While finer-scale

estimates of recombination might reveal a significant effect and it is possible that ectopic

recombination rates are less correlated with rates of crossing-over than is often assumed, the

previously observed lack of an effect of recombination on TE abundance in *A. thaliana* [15]

cannot be attributed to its selfing mating system.

If TEs are deleterious because they disrupt gene function, then insertions should be most

deleterious, and consequently at lowest population frequency, when they are located in or near

genes. Plotting TE frequency against distance from the closest gene, in bins of 500 bp, reveals a

significant positive correlation (Fig 1B; $r = 0.94$), suggesting that TEs closer to genes experience

stronger purifying selection. This result is in line with a recent publication in the same species

[31].

Another way to test the gene disruption hypothesis is to plot the relationship between

average TE frequency and coding site density in 100 kb windows across the genome. In testing

this, we found a significant negative correlation, suggesting that TEs in regions of high gene

density are kept at low population frequency (Fig 1C; Spearman's ρ: -0.562). We also evaluated

TE frequency relationship to conserved non-coding sequences (CNSs), using CNSs conserved

across 9 Brassicaceae species, many of which show evidence of being regulatory elements [32].

Plotting TE frequency vs CNS density also leads to a strong negative correlation, implying TEs

face selective pressures due to negative effects of inserting into or near regulatory elements (Fig

1D; Spearman's ρ: -0.619). There is an obvious auto-correlation between coding site density and

CNS density as regulatory elements will be found close to genes. To check that the relationship

of CNS density and TE frequency cannot be fully explained by the correlation between genes

and CNSs, we built two quadratic regression models chosen because they showed the best fit to

the data. In the first model, TE frequency is predicted only by proportion of coding sites while in

the second, the proportion of CNSs is added as an additional predictor (the latter model is shown

in Table 1; adjusted $R^2 = 0.502$). Using an F-test on the two models showed the model with

coding sites and CNSs explains significantly more variation than the model with coding sites

alone ($p < 0.0001$). These results suggest that important TE effects on genes extend not only to

direct gene disruption and methylation-mediated gene silencing, but also to disruption of

regulatory sequence. The effect of TEs on CNSs could be due to direct disruption of enhancers,

methylation of regulatory sequence caused by TE silencing, or spatial disruption of gene

regulation caused by insertions between a gene and its regulatory region leading to aberrant

expression.

Table 1. Quadratic regression model of average TE frequency.

Model: Average TE frequency ~ proportion of coding sequences (CSs) + (proportion of coding sequences)^2 + proportion of CNSs + (proportion of CNSs)^2		
Coefficient	**Estimate**	**P-value**
Proportion of coding sequences	-0.100	< 0.0001
(Proportion of coding sequences)^2	0.126	< 0.0001
Proportion of CNSs	-0.236	< 0.0001
(Proportion of CNSs)^2	1.399	0.000381
Adjusted R^2	0.502	< 0.0001

TEs contribute significantly to gene expression variation

We have shown that TEs contribute to sequence variation, mostly as rare mutations that are under stronger purifying selection near genes and regulatory regions. Given this, we next investigated the extent to which TE insertions alter gene expression and contribute to standing genetic variation. One approach for testing rare variants is to ask whether individuals with extreme gene expression are often associated with TEs near their genes, or whether there is a greater burden of rare variants at the phenotypic extremes. Individual genes are sorted into expression rank bins, and then the total number of all local rare variants is plotted for all genes in all individuals. If rare variants cause a change in gene expression, either an increase or decrease, then this will result in a parabolic shape [33].

To facilitate later comparison with SNP data, we used 109 individuals which passed TE, expression and SNP quality filters. Using all TEs with a MAF < 3% and within 500 bp of a gene, we found that individuals with more TEs near a gene are more likely to have extreme expression values, and TEs appear to have a strong effect on decreasing gene expression, and a weaker effect of increasing expression at the extremes (Fig 2A; $R^2 = 0.450$). As a null control, we permuted mutation count and expression rank; as expected, permutations lead to no significant correlation, demonstrating the significance of our observed effect (S6 Fig). This general result suggests that TEs do contribute to expression variation and that TE silencing and gene disruption effects may be more common than TE-mediated upregulation. This result is in contrast with a recent analysis in *Arabidopsis thaliana*, where TE insertions driving increased expression appeared to be as prevalent as expression-decreasing effects ([6], but see [7]).

Fig 2. Relationship between (A) TE and (B) SNP count and gene expression rank bin for all rare variants (MAF < 3%) within 500 bp of a gene. C) Ratio of average number of rare variants in expression outliers relative to non-outliers for TEs and SNPs. 'Upper' refers to expression-increasing outliers, and 'lower' refers to expression-decreasing outliers. We ran 1000 permutations of the data to obtain normal distributions of the ratio statistic, and in all cases the observed statistic fell outside of the permuted distribution values.

While TEs appear to strongly downregulate gene expression, it is difficult from such an analysis to directly quantify the strength of the effect. We used a complementary statistical

approach by calculating the average number of TE insertions across genes in individuals that are

expression outliers relative to the average number in non-expression outliers. The ratio value for

TEs is 1.13, and expression outlier individuals have significantly more TEs than non-outlier

individuals (Fig 2C; 95% quantile confidence interval: 1.10-1.17 from 1000 bootstraps). When

considering only downregulated expression outliers, the ratio of TEs in outliers vs non-outliers is

a much higher value of 1.48, consistent with the pattern observed in the burden of rare variants

test (Fig 2C; 95% quantile confidence interval: 1.36-1.61 from 1000 bootstraps). These results

suggest that the standing genetic variation contributed by TEs has a significant effect on

disrupting gene expression, pushing individuals to extreme expression values for a trait which is

under stabilizing selection and that TE insertions are more likely to decrease expression than

increase it.

Breaking down the results by TE type reveals large differences in the direction and type

of effect that different TE types have on gene expression. Long terminal repeat (LTR) elements,

which are large retrotransposons, show the strongest effect on decreasing gene expression ($R^2 =$

0.471) while other TE types tested show significant but less pronounced effects (Fig 3, S7 Fig).

LTRs are the most abundant and the most methylated TE type in most plant genomes [34],

including *C. grandiflora*. Methylated TEs, which have been silenced by the host genome to

prevent their transcription and transposition, can have stronger effects on nearby gene

expression, either through the spread of methylation past the TE or by making the chromatin less

accessible to transcription factors and transcriptional machinery.

Fig 3. Ratio of average number of rare variants in expression outliers relative to non-outliers for different TE types. Black – all outliers; blue – expression-decreasing outliers; red – expression-increasing outliers.

Genic insertions cause the strongest expression decrease

If TEs affect gene expression by disrupting gene regulatory regions, then there should be a strong effect caused by TEs upstream of the gene. However, the strongest correlation with extreme gene expression occurred when considering TEs inserted into genes (S8 Fig; $R^2 =$ 0.524). Upstream TEs had a lower overall effect; interestingly they were only significantly over-enriched in increased expression outliers, with no significant effect in decreased expression outliers (Fig 4). This is somewhat similar to observations in *A. thaliana*, where upstream insertions showed a more symmetrical effect on expression than insertions inside or downstream of genes [7]. The upstream effect was only significant for TEs within 100 bp of the transcription start site (TSS) (Fig 4, S9 Fig), consistent with previous results suggesting that the greatest abundance of upstream conserved non-coding sites were 200 bp away from the TSS [32].

Similarly, upstream eQTLs were most abundant within 1kb from the TSS (closer intervals were

not tested) [25]. One possible explanation for the weaker and more symmetrical overall effect of

insertions upstream of genes is that a major effect of TE insertions into regulatory sequence is an

effect on expression noise, rather than a specific downregulation. Recent work on de novo

mutational effects in regulatory regions in yeast [35] suggest many more mutations have an

effect on the variance in gene expression than on the expression mean. TE effects could be

similar in regulatory regions, driving a pattern of purifying selection associated with increased

variance in average expression.

Fig 4. Ratio of average number of rare variants in expression outliers relative to non-

outliers for TEs upstream of, in, and downstream of genes. The up- and downstream

categories are non-overlapping. Black – all outliers; blue – expression-decreasing outliers; red –

expression-increasing outliers.

Intragenic TEs appear to cause the strongest shift in expression, so we further categorised

by the genic region of insertion. TEs in all genic regions were significantly over-enriched in low

expression gene outliers, with the strongest effect from insertions into coding regions (Fig 4, S10

Fig). In contrast, only TEs into the 5'UTR and coding sites were also significantly over-enriched

in high expression gene outliers. The 5'UTR contains regulatory sites, so TE insertions may be

contributing extra regulatory sites leading to increased transcription of the whole gene. TEs are

known to be a large source of promoter and regulatory regions which can be co-opted by the host

genome and contribute to the evolution of regulatory networks [36,37]. The large effect of TEs

in coding regions on decreasing gene expression could be due to faster decay of mRNAs with

large mutations, for example by the nonsense-mediated decay (NMD) pathway [38]. However,

the strong enrichment of TEs in low expression outliers when they insert into coding sites could

be driven by a technical artifact where reads with TEs are dropped due to poor mapping (see

Methods). When considering gene expression broken down into individual exons, there were no

significant differences in enrichment of TEs in exons upstream of the insertion, downstream of

the insertion and the exon containing the insertion (S4 Fig). Therefore there is not evidence for

technical artifacts which could lead to misinterpretation of results. Furthermore, TEs appear to

equally reduce expression of all exons, suggesting that TEs are not causing early transcription

breaks. The impact of TEs in introns could be driven by targeted silencing of insertions in introns

[39].

TE insertions show stronger enrichment in expression outliers than SNPs

Josephs and colleagues [25] previously showed that a third of genes had an eQTL in this

population of *C. grandiflora*, and that there is a significant negative correlation between eQTL

MAF and effect size, suggesting that purifying selection is mostly responsible for maintaining

genetic variation for gene expression. However, they tested SNPs with a MAF > 5%, and so the

contribution of rare variants to gene expression variation is still unclear.

We explored the effect of rare SNPs with a MAF < 3% on gene expression. Overall, these

variants are present in much larger number, and are equally likely to cause gene up- or down-

regulation (Fig 2B; R^2 = 0.517). This is in contrast with a recent study in maize, where SNP

variants upstream of genes tended to show greater evidence of downregulation [40]. When

comparing the ratio of variants in expression outliers vs non-outliers, SNPs had a smaller effect

than TEs (1.04 vs 1.13, respectively; Fig 2C). As a class of mutation, TEs are 9% more over-

represented in outliers than SNPs. In low expression outliers, there is a considerably greater gap

between TE and SNP over-representation of 43% (1.48 vs 1.05 ratio values, respectively; Fig

2C). Both tests of the burden of rare variants on gene expression are genome-wide, made by

summing across all genes and all individuals in the population. As such, the results highlight the

global impact that both TEs and SNPs have on gene expression variation. The outlier test used a

stringent cut-off of 4 standard deviations away from the mean to define outliers, so it may be that

TEs are on average more likely to cause a large change in expression and are over-enriched in

this test. SNPs, on the other hand, may be on average more likely to cause a small change in

expression and are more important in explaining small variation in expression. Therefore,

individual TEs may be strong candidates for having large effects on expression (large-effect

QTL), but considering their numbers SNPs are also likely important, with smaller individual

effects on average.

A major challenge when comparing SNPs to TEs is detection power. Calling non-

reference TEs with short-read data is difficult, especially in outcrossing species where many

insertions are heterozygous, leading to more false positives and false negatives than when calling

SNPs [41]. These power issues could lead to biases in our population frequency estimates. There

may be many more rare TEs in the population which the pipeline is not detecting, however, in

this case our results would be undervaluing the true effect sizes and are therefore conservative

estimates of the impact of rare TEs. The outlier expression test controls for this problem because

it calculates a ratio of TE numbers between two groups, both of which have equal detection

power, allowing for the comparison of TE and SNP effect size independent of the ability to call

these variants.

Conclusions

This study explored the relationship between TEs and the rest of the genome in a

population of *C. grandiflora*. We find that selection against disruption of gene function and

regulation plays an important role in limiting TE copy number, but we see no evidence for

selection against ectopic recombination. TE insertions in or near genes strongly downregulate

expression and to a smaller extent upregulate expression, with the effect differing by the type of

TE and the genic region into which it has transposed. In comparison, SNPs have an overall

weaker effect and are equally likely to up- and down-regulate expression. These results highlight

the important contribution of TEs to standing genetic and phenotypic variation.

Methods

Study organism

The self-incompatible, outcrossing plant *Capsella grandiflora*, is a good model for using

population genomic approaches to make inferences about natural selection due to its low

population structure and large effective population size ($N_e \approx 600,000$) [42–44]. The species is

a member of the Brassicaceae family and is native to Northern Greece and Southern Albania.

There is evidence of prevalent purifying and positive selection on SNPs in *C. grandiflora*

[43,45].

Genomic data and expression

The DNA and RNA data have been previously described [25,46]. Briefly, plants were

sampled from a single population located near Monodendri, Greece (population Cg-9 in [44]).

Seeds were collected and grown in the University of Toronto glasshouses and random crosses

were performed. The offspring of these crosses were grown in the University of Toronto growth

chambers, where leaf tissue was collected and used for DNA and RNA extraction. Whole

genome sequencing was done using 100 cycles of paired-end sequencing in a Hiseq 2000 with

Truseq libraries (Illumina), resulting in 100 bp paired-end reads. RNA was sequenced at the

Genome Quebec Innovation Centre in an Illumina HiSeq resulting in 100 bp paired-end or

single-ended reads. For consistency, one read was randomly chosen among the paired end reads

and they were subsequently treated as single ended. The reference genome used for mapping

reads is *Capsella rubella*, a close relative of *C. grandiflora*, which diverged approximately

200,000 years ago [28,42].

RNA reads were mapped to an exon-only reference generated from the *C. rubella*

reference genome [25,28] using Stampy 1.0.21 with default settings [47]. Expression level was

measured with the HTSeq.scripts.count feature from HTSeq [48] and normalized for sequencing

depth by dividing read counts by median read count of the entire sample. Genes with a median

expression level below five reads per sample before normalization were removed from the

analyses.

Calling TEs

360   We built a new TE library to use with the detection pipeline, with a focus on TEs that

show evidence of being recently active. Using pindel [49], we found large deletions (>150 bp) in

30 of the *C. grandiflora* short-read sequenced individuals relative to the *C. rubella* reference

genome. These represent candidate TE insertions that occured in *C. rubella* since they in the

reference genome at those locations but absent from those locations in the *C. grandiflora* sample.

The PASTEClassifier pipeline [50] allowed us to classify the sequences obtained from the *C.*

rubella reference into TE order based on the hierarchical TE classification system [51]. Any

sequences that remained unclassified were removed.

368   We used the RelocaTE2 [26] pipeline to detect both reference and non-reference TE

insertions in 124 *C. grandiflora* individuals from our population. Briefly, RelocaTE2 is based on

a combined split read and discordant read pairs mapping approach. All fastq files were

subsampled down to 70X coverage to avoid elevated TE calls in individuals with higher

coverage, and 23 individuals with below 70X coverage were removed. RelocaTE2 was run using

bowtie2 as the TE aligner, using the custom-built TE library described above, and the *C. rubella*

reference genome with default settings. We used bwa [52] and samtools [53] to align short reads

to the reference genome, and then passed these alignments to RelocaTE2 to genotype TE

insertions as heterozygous or homozygous. Heterozygous TE insertions will have reads aligning

to the reference over the insertion site because one of the homeologs does not have a TE.

To create a population level table of TE insertions and their genotype (homozygous

absent, heterozygous, or homozygous present), we merged all the insertions from the RelocaTE2

'high-confidence' labelled files, which were insertions supported by at least one junction read

from both sides. Any insertions called between individuals which were within 50 bp of one

another and were of the same TE classification type were considered as the same insertion.

While RelocaTE2 can genotype insertions as homozygous or heterozygous, there is some

evidence from the TE population frequency spectrum that these calls are not always accurate.

When using allele counts (0, 1, or 2 for absent, heterozygous, homozygous) to make a frequency

spectrum, there is a staggered pattern which goes away when examining the spectrum using

individual count (0 or 1 for presence or absence; see S1 Fig and Fig 1). This suggests that

homozygosity is being over-called. For all rare variant gene expression analyses (see below), we

used individual counts, effectively ignoring homozygous vs heterozygous calls, however results

were qualitatively the same when performed with allele counts (S2 Fig).

TE validation

Validating RelocateTE2's ability to find TEs is important, especially in outbred

heterozygous population data where only half of the reads will support a heterozygous TE

insertion. RelocaTE2 was previously tested on a rice strain of 20-fold genome coverage of 100

bp paired-end Illumina short reads aligned to the rice genome, and resulted in 93% specificity

(the percent of calls which were true positives) [26]. The *C. grandiflora* data should have a

comparable level of specificity because it has higher coverage than the rice data previously

tested. Nevertheless, we used an independent bioinformatics test to estimate the true-positive rate

(the proportion of new TEs called by RelocaTE2 that are actually TEs). We ran the SPAdes de

novo assembler [54] on all TE calls on scaffold 1 for five individuals, chosen for their high

coverage to increase the chance of success of the de novo assembler. The high coverage

individuals should not bias the true-positive rate estimate because all individuals were

subsampled to the same coverage (70X) for TE calling. We assembled all reads mapping to the

region 1000 bp upstream and downstream of the TE insertion call and then aligned the de novo

assembly contigs to the TE library using BLAST+ on default settings [55]. If we identified

significant hits to the TE library, we considered the TE call to be a true positive.

The average true positive value across the 5 individuals was 80% for TEs labelled as

high-confidence by RelocaTE2 (see table in Appendix for breakdown by individual). The true

positive value for non-high-confidence TEs was 60%. For subsequent analyses, we therefore

focused on the high-confidence calls. Analyses considering both high-confidence and low-

confidence calls led to similar results, but with lower significance, consistent with the greater

noise contributed by the lower-confidence calls (S3 Figure).

Gene and genome annotation

TE insertions into genes were categorised as exonic, intronic, 3' UTR or 5' UTR based

on the *C. rubella* genome annotation [28]. Similarly, 4-fold SNPs were identified for comparison

with TE site frequency spectrum. Conserved non-coding sites (CNSs) are determined through

alignment of 9 Brassicaceae species to the *C. rubella* genome as described in [32,45].

Recombination rate data comes from fitting a third-order polynomial crossover events from an

F2 genetic map derived from a cross between a *C. rubella* and *C. grandiflora* individual [28,56].

We ran quadratic multiple linear regression models predicting TE frequency from the

proportion of coding sites alone, or both coding sites and CNSs using the R lm function [57]. We

used the anova function to test whether the model with two predictor variables explained more

variation than the simpler model.

SNP dataset

Initial SNP calling and filtering was previously described [25,46]. From the 124

individuals used in TE detection, 15 were dropped from further analyses due to an excess of NA

SNP genotype calls (over 50,000 NA calls/individual). Any SNPs where more than 9 individuals

had an NA call were also dropped. The cutoffs were empirically determined as large outliers

when plotting histograms of NA calls per individual and NA calls per SNP. All other NA calls

were assumed to be homozygous for the reference allele; given the focus on rare variants, this

assumption is unlikely to have a strong effect on results.

Burden of rare variants test

The statistical test we used for determining the effect of rare variants (both TEs and

SNPs) on gene expression was first described by Zhao and colleagues [33]. Briefly, the test

estimates a genome-wide correlation between total rare variant count and gene expression

relative to the rest of the population, testing whether individuals with more extreme expression

values for a certain gene have more rare genetic variants located near that gene. Individuals are

sorted into expression rank bins and the number of rare variants located in and near a gene is

summed across all genes in the genome, for each expression rank bin. If there is a burden of rare

variants, and having more of them pushes individuals to the extremes of expression (relative to

the rest of the population), there will be a quadratic relationship between rare variant count and

expression rank. Note that each gene is assessed independently, such that at individual genes,

different individuals can end up in the tails of the distribution.

The rare variant burden plots were created using TEs and SNPs with a minor allele

frequency (MAF) less than 3% from all 109 individuals. For all plots, we used TEs and SNPs

which were within a gene or 500 bp upstream or downstream. We identified a small subset of

genes where some individuals showed extreme numbers of rare SNPs, possibly reflecting

alignment errors or paralogous sequences. Any genes where at least one individual had over 40

rare SNPs were removed (874 genes dropped in total).

For plots which considered only a single genic region (5'UTR, exon, intron, or 3'UTR),

there may be some bias due to genes where UTRs were not annotated. Note that given the

residual uncertainty of exact TE insertion locations (approx. 50 bp), this means that there will be

some noise associated with annotation of insertion position.

Expression outlier test

We developed an alternative method to test the effect of rare burdens on gene expression

which allows for a more direct comparison between the effect of TEs and SNPs. Again, we used

all variants with a MAF < 3% from the 109 individuals which passed both TE and SNP filtering

and which were located within a gene or 500 bp away.

The test involves determining individuals which are expression outliers, and comparing

their number of rare variants with those of non-outlier individuals, repeated for each gene.

Outliers were called as 4 standard deviations away from the mean, where mean and standard

deviation were calculated using the expression values of individuals between the 1st and 3rd

quartile. Estimating the mean and standard deviation from individuals between the 1st and 3rd

quartiles reduces bias in these estimates caused by large expression value outliers and non-

symmetrical distributions. As this method underestimates variation (relative to the total sample),

an extreme cut-off of 4 standard deviations (SD) was used to call outliers. When testing without

discrimination between upper and lower outliers, all genes were used except for 11, in which all

individuals were within 4 SD of the mean (total of 18874 genes). When distinguishing between

upper and lower outliers, we only considered genes where the mean was at least 4 SD above 0, to

ensure it was possible to have lower outliers. After removing genes where all individuals were

within 4 SD of the mean, there were 10290 genes tested for upper outliers and 14849 genes

tested for lower outliers.

For every gene, we computed the number of rare variants per individual to control for

differences in number of outliers and non-outliers among and between genes, and checked for a

significant difference by running a Wilcoxon signed rank test using the R wilcox.test function for

both TEs and SNPs [57]. To determine a per-variant effect size, we calculated the ratio of the

mean number of rare variants per outlier individual to the mean number of rare variants per non-

outlier individual across all genes. This ratio expresses how many more variants there are per

outlier vs non-outlier. Taking the difference between the TE ratio and SNP ratio gives a

quantitative measure of how much more likely the average TE is to cause an expression outlier

vs the average SNP.

We ran 1000 bootstraps of the ratio using the R boot() package in order to calculate 95%

quantile confidence intervals [58]. 1000 permutations were also run to build a null distribution,

where the outlier vs non-outlier label was switched with a 50% probability for every gene.

Expression outlier by exon

492   We performed a slightly altered version of the expression outlier test in order to control

493 for a possible technical artifact which could lead to the appearance of TE insertions in exons

494 decreasing gene expression. RNA reads that include a TE may be dropped in filtering due to

495 poor mapping, leading to an appearance of reduced overall transcript expression. To test this, we

496 divided total gene expression into individual exon expression. For each gene with a TE insertion

497 in at least one individual in the population, we created three categories: average expression of all

498 exons upstream of the TE insertion, expression of the exon with the TE insertion, and average

499 expression of all exons downstream of the TE insertion. If there was a technical artifact, and

500 RNA reads with TEs were dropped due to poor mapping, we would expect that only exons with

501 TEs had TEs associated with low expression outliers. Upstream and downstream exons wouldn't

502 show TE enrichment in low expression outliers, because their expression would not be affected

503 by poor mapping.

Acknowledgements

506   We thank Aneil Agrawal for key recommendations for statistical methods and Rob Ness

507 for important suggestions on data validation. We thank Young Wha Lee and Niroshini

508 Epitawalage for assistance collecting data. Thomas Bureau, Mathieu Blanchette, Daniel Schoen,

509 Paul Harrison, Alan Moses, Adrian Platts, and Eef Harmsen contributed to the design and

510 implementation of the Value-directed Evolutionary Genomics Initiative (VEGI).

References

513 1. Hurst GD, Werren JH. The role of selfish genetic elements in eukaryotic evolution. Nat Rev

514   Genet. 2001;2: 597–606.

2. Doolittle WF, Sapienza C. Selfish genes, the phenotype paradigm and genome evolution.

Nature. 1980;284: 601–603.

3. Charlesworth B, Charlesworth D. The population dynamics of transposable elements. Genet

Res. 1983;42: 1–27.

4. Petrov DA, Fiston-Lavier A-S, Lipatov M, Lenkov K, González J. Population genomics of

transposable elements in *Drosophila melanogaster*. Mol Biol Evol. 2011;28: 1633–1644.

5. Cridland JM, Macdonald SJ, Long AD, Thornton KR. Abundance and distribution of

transposable elements in two *Drosophila* QTL mapping resources. Mol Biol Evol. 2013;30:

2311–2327.

6. Quadrana L, Bortolini Silveira A, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA,

et al. The *Arabidopsis thaliana* mobilome and its impact at the species level. Elife. 2016;5:

6919–6931.

7. Stuart T, Eichten S, Cahn J, Karpievitch Y, Borevitz J, Lister R. Population scale mapping

of transposable element diversity reveals links to gene regulation and epigenomic variation.

Elife. 2016;5: e20777.

8. Laricchia KM, Zdraljevic S, Cook DE, Andersen EC. Natural variation in the distribution

and abundance of transposable elements across the *Caenorhabditis elegans* species. Mol

Biol Evol. 2017;34: 2187–2202.

9. Barrón MG, Fiston-Lavier A-S, Petrov DA, González J. Population genomics of

transposable elements in *Drosophila*. Annu Rev Genet. 2014;48: 561–581.

10. Langley CH, Montgomery E, Hudson R, Kaplan N, Charlesworth B. On the role of unequal

exchange in the containment of transposable element copy number. Genet Res. 1988;52:

223–235.

11. Charlesworth B, Lapid A, Canada D. The distribution of transposable elements within and

between chromosomes in a population of *Drosophila melanogaster*. II. Inferences on the

nature of selection against elements. Genet Res. 1992;60: 115–130.

12. Bennetzen JL, Wang H. The contributions of transposable elements to the structure,

function, and evolution of plant genomes. Annu Rev Plant Biol. 2014;65: 505–530.

13. Lee YCG. The role of piRNA-mediated epigenetic silencing in the population dynamics of

transposable elements in *Drosophila melanogaster*. PLoS Genet. 2015;11: e1005269.

14. Lee YCG, Karpen GH. Pervasive epigenetic effects of *Drosophila* euchromatic transposable

elements impact their evolution. Elife. 2017;6: e25762.

15. Wright SI, Agrawal N, Bureau TE. Effects of recombination rate and gene density on

transposable element distributions in *Arabidopsis thaliana*. Genome Res. 2003;13: 1897–

1903.

16. Charlesworth D, Charlesworth B. Transposable elements in inbreeding and outbreeding

populations. Genetics. 1995;140: 415–417.

17. Wright SI, Schoen DJ. Transposon dynamics and the breeding system. Genetica. 1999;107:

139–148.

18. Laricchia KM, Zdraljevic S, Cook DE, Andersen EC. Natural variation in the distribution

and abundance of transposable elements across the *Caenorhabditis elegans* species. Mol

Biol Evol. 2017;34: 2187–2202.

19. Cridland JM, Thornton KR, Long AD. Gene expression variation in *Drosophila*

melanogaster due to rare transposable element insertion alleles of large effect. Genetics.

2015;199: 85–93.

20. Hollister JD, Smith LM, Guo Y-L, Ott F, Weigel D, Gaut BS. Transposable elements and

small RNAs contribute to gene expression divergence between *Arabidopsis thaliana* and

Arabidopsis lyrata. Proc Natl Acad Sci U S A. 2011;108: 2322–2327.

21. Wang X, Weigel D, Smith LM. Transposon variants and their effects on gene expression in

Arabidopsis. PLoS Genet. 2013;9: e1003255.

22. González J, Lenkov K, Lipatov M, Macpherson JM, Petrov DA. High rate of recent

transposable element-induced adaptation in *Drosophila melanogaster*. PLoS Biol. 2008;6:

2109–2129.

23. Emera D, Casola C, Lynch VJ, Wildman DE, Agnew D, Wagner GP. Convergent evolution

of endometrial prolactin expression in primates, mice, and elephants through the

independent recruitment of transposable elements. Mol Biol Evol. 2012;29: 239–247.

24. Makarevitch I, Waters AJ, West PT, Stitzer M, Hirsch CN, Ross-Ibarra J, et al.

Transposable elements contribute to activation of maize genes in response to abiotic stress.

PLoS Genet. 2015;11: e1004915.

25. Josephs EB, Lee YW, Stinchcombe JR, Wright SI. Association mapping reveals the role of

purifying selection in the maintenance of genomic variation in gene expression. Proc Natl

Acad Sci U S A. 2015;112: 15390–15395.

26. Chen J, Wrightsman TR, Wessler SR, Stajich JE. RelocaTE2: a high resolution transposable

element insertion site mapping tool for population resequencing. PeerJ. 2017;5: e2942.

27. Dolgin ES, Charlesworth B, Cutter AD. Population frequencies of transposable elements in

selfing and outcrossing *Caenorhabditis* nematodes. Genet Res. 2008;90: 317–329.

28. Slotte T, Hazzouri KM, Ågren JA, Koenig D, Maumus F, Guo Y-L, et al. The *Capsella*

rubella genome and the genomic consequences of rapid mating system evolution. Nat

Genet. 2013;45: 831–835.

29. Tian Z, Zhao M, She M, Du J, Cannon SB, Liu X, et al. Genome-wide characterization of

nonreference transposons reveals evolutionary propensities of transposons in soybean. Plant

Cell. 2012;24: 4422–4436.

30. Novick PA, Smith JD, Floumanhaft M, Ray DA, Stéphane B. The evolution and diversity of

DNA transposons in the genome of the lizard *Anolis carolinensis.* Genome Biol Evol.

2011;3: 1–14.

31. Horvath R, Slotte T. The role of small RNA-based epigenetic silencing for purifying

selection on transposable elements in *Capsella grandiflora.* Genome Biol Evol. 2017;9:

2911–2920.

32. Haudry A, Platts AE, Vello E, Hoen DR, Leclercq M, Williamson RJ, et al. An atlas of over

90,000 conserved noncoding sequences provides insight into crucifer regulatory regions. Nat

Genet. 2013;45: 891–898.

33. Zhao J, Akinsanmi I, Arafat D, Cradick TJ, Lee CM, Banskota S, et al. A burden of rare

variants associated with extremes of gene expression in human peripheral blood. Am J Hum

Genet. 2016;98: 299–309.

34. Ahmed I, Sarazin A, Bowler C, Colot V, Quesneville H. Genome-wide evidence for local

DNA methylation spreading from small RNA-targeted sequences in *Arabidopsis.* Nucleic

Acids Res. 2011;39: 6919–6931.

35. Metzger BPH, Duveau F, Yuan DC, Tryban S, Yang B, Wittkopp PJ. Contrasting

Frequencies and Effects of cis- and trans-Regulatory Mutations Affecting Gene Expression.

Mol Biol Evol. 2016;33: 1131–1146.

36. Hirsch CD, Springer NM. Transposable element influences on gene expression in plants.

Biochimica Biophys Acta. 2016;1860: 157-165

37. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements: from

608   conflicts to benefits. Nat Rev Genet. 2017;18: 71–86.

609   38. Garneau NL, Wilusz J, Wilusz CJ. The highways and byways of mRNA decay. Nat Rev

610   Mol Cell Biol. 2007;8: 113–126.

611   39. Liu N, Lee CH, Swigut T, Grow E, Gu B, Bassik MC, et al. Selective silencing of

612   euchromatic L1s revealed by genome-wide screens for L1 regulators. Nature. 2018;553:

613   228–232.

614   40. Kremling KAG, Chen SY, Su MH, Lepak MK, Romay MC, Swarts, KL, et al.

615   Dysregulation of expression correlates with rare-allele burden and fitness loss in maize.

616   Nature. 2018;555: 520–523.

617   41. Ewing AD. Transposable element detection from whole genome sequence data. Mob DNA.

618   2015;6: 24.

619   42. Foxe JP, Slotte T, Stahl EA, Neuffer B, Hurka H, Wright SI. Recent speciation associated

620   with the evolution of selfing in *Capsella*. Proc Natl Acad Sci U S A. 2009;106: 5241–5245.

621   43. Slotte T, Foxe JP, Hazzouri KM, Wright SI. Genome-wide evidence for efficient positive

622   and purifying selection in *Capsella grandiflora*, a plant species with a large effective

623   population size. Mol Biol Evol. 2010;27: 1813–1821.

624   44. St. Onge KR, Källman T, Slotte T, Lascoux M, Palmé AE. Contrasting demographic history

625   and population structure in *Capsella rubella* and *Capsella grandiflora*, two closely related

626   species with different mating systems. Mol Ecol. 2011;20: 3306–3320.

627   45. Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, et al.

628   Evidence for widespread positive and negative selection in coding and conserved noncoding

629   regions of *Capsella grandiflora*. PLoS Genet. 2014;10: e1004622.

46. Josephs EB, Wright SI, Stinchcombe JR, Schoen DJ. The relationship between selection,

network connectivity, and regulatory variation within a population of *Capsella grandiflora*.

Genome Biol Evol. 2017;9: 1099–1109.

47. Lunter G, Goodson M. Stampy: a statistical algorithm for sensitive and fast mapping of

Illumina sequence reads. Genome Res. 2011;21: 936–939.

48. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput

sequencing data. Bioinformatics. 2015;31: 166–169.

49. Ye K, Schulz MH, Long Q, Apweiler R, Ning Z. Pindel: a pattern growth approach to detect

break points of large deletions and medium sized insertions from paired-end short reads.

Bioinformatics. 2009;25: 2865–2871.

50. Hoede C, Arnoux S, Moisset M, Chaumier T, Inizan O, Jamilloux V, et al. PASTEC: an

automatic transposable element classification tool. PLoS One. 2014;9: e91929.

51. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified

classification system for eukaryotic transposable elements. Nat Rev Genet. 2007;8: 973–

982.

52. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25: 1754–1760.

53. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

54. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a

new genome assembly algorithm and its applications to single-cell sequencing. J Comput

Biol. 2012;19: 455–477.

55. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10: 421.

56. Slotte T, Hazzouri KM, Stern D, Andolfatto P, Wright SI. Genetic architecture and adaptive

significance of the selfing syndrome in *Capsella*. Evolution. 2012;66: 1360–1374.

57.   R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. 2016.

58. Canty A, Ripley B. boot: Bootstrap R (S-Plus) Functions. R package version 1.3-20. 2017.

Supporting Information

S1 Fig. The TE population frequency spectrum made using allele counts.

S2 Fig. Relationship between TE count and gene expression rank bin for all rare TEs

within 500 bp of a gene. TE genotype was considered, where heterozygous TEs were counted

once and homozygous TEs were counted twice.

S3 Fig. Relationship between TE count and gene expression rank bin for all rare TEs,

including low-confidence calls, within 500 bp of a gene.

S4 Fig. Ratio of average number of rare variants in expression-decreasing outliers relative

to non-outliers for TEs. Expression is calculated separately for each exon. 'Up' refers to

average expression of all exons upstream of where the TE inserted. 'In' refers to the exon where

the TE inserted. 'Down' refers to average expression of all exons downstream of where the TE

inserted.

S5 Fig. Relationship between average TE frequency and average cross-over rate in 100 kb

bins across the genome (Pearson's r = 0.039).

S6 Fig. Relationship between permuted TE count and gene expression rank bin for all rare

TEs within 500 bp of a gene.

S7 Fig. Relationship between TE count and gene expression rank bin for different TE

types.

S8 Fig. Relationship between TE count and gene expression rank bin divided by TE

insertion location relative to gene. Upstream/downstream refers to 500 bp away from gene

annotation start/end, respectively.

S9 Fig. Ratio of average number of rare variants in expression outliers relative to non-

outliers for TEs inserted upstream of gene transcription start site. Upstream intervals are

non-overlapping. 'Upper' refers to expression-increasing outliers, and 'lower' refers to

expression-decreasing outliers.

S10 Fig. (A-D) Relationship between TE count and gene expression rank bin divided by

genic region of the insertion.

S1 Table. TE validation statistics showing the proportion of false positives, as determined

through an independent bioinformatics approach, for five tested individuals.

S1 File. All TE calls used in analyses across the population.