1

USP9X deubiquitylates DVL2 to regulate WNT pathway specification

Casey P. Nielsen[1], Kristin K. Jernigan[1], Nicole L. Diggins, Donna J. Webb and Jason A.

MacGurn [1][*]

[1] Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN, United

States

*Address correspondence to Jason A. MacGurn; email: jason.a.macgurn@vanderbilt.edu

Running title: Regulation of WNT pathway specification by USP9X

Current Character Count: 52,314 (with spaces).

Summary

The WNT signaling network is comprised of multiple receptors that relay various input

signals via distinct transduction pathways to execute multiple complex and context-specific

output processes. Integrity of the WNT signaling network relies on proper specification between

canonical and non-canonical pathways, which presents a regulatory challenge given that

several signal transducing elements are shared between pathways. Here, we report that

USP9X, a deubiquitylase, and WWP1, an E3 ubiquitin ligase, regulate a ubiquitin rheostat on

DVL2, a WNT signaling protein. Our findings indicate that USP9X-mediated deubiquitylation of

DVL2 is required for canonical WNT activation, while increased DVL2 ubiquitylation is

associated with localization to actin-rich projections and activation of the planar cell polarity

(PCP) pathway. We propose that a WWP1-USP9X axis regulates a ubiquitin rheostat on DVL2

that specifies its participation in either canonical WNT or WNT-PCP pathways. These findings

have important implications for therapeutic targeting of USP9X in human cancer.

Introduction

The canonical WNT β-catenin signaling pathway is involved in regulating many cellular processes such as cell fate determination during embryonic development, cell proliferation, and adult tissue homeostasis. Thus, it is not surprising that aberrant activation of the canonical Wnt pathway is known to occur in many types of cancer (MacDonald et al., 2009; Saito-Diaz et al., 2013). There are also several noncanonical WNT signaling pathways including the WNT-Planar Cell Polarity (WNT-PCP) pathway which controls cell migration and tissue polarity. Dysregulation of the WNT-PCP pathway has been linked to cancer invasion and metastasis (Katoh, 2005; Luga et al., 2012; Wang, 2009). While the canonical WNT β-catenin pathway and the noncanonical WNT-PCP pathway use divergent effector mechanisms to regulate distinct cellular functions, these pathways share membrane receptor components and the cytoplasmic WNT transducer protein dishevelled (DVL). Despite its key role in both pathways, the mechanisms dictating DVL participation in canonical or noncanonical WNT signaling are yet to be elucidated.

Initiation of the canonical WNT β-catenin pathway occurs when extracellular WNT ligand binds to the co-receptors Frizzled (FZD) and low-density lipoprotein receptor-related protein 5/6 (LRP5/6) leading to recruitment of DVL and AXIN to the WNT ligand receptor complex (MacDonald et al., 2009). This ultimately results in the inhibition of β-catenin ubiquitylation and degradation such that stabilized β-catenin can enter the nucleus to initiate a transcriptional program (MacDonald et al., 2009; Saito-Diaz et al., 2013). On the other hand, core WNT-PCP pathway components Van Gogh-Like 1 (VANGL1), FZD, Prickle (Pk), DVL, and others function to activate RHOA, c-Jun N-terminal kinase (JNK), and nemo-like kinase (NLK) signaling cascades in order to coordinate tissue polarity and cell motility through regulation of actin dynamics (Glinka et al., 2011).

Ubiquitylation is known to be involved in key regulatory steps of both the canonical WNT

and noncanonical WNT-PCP pathways. For example, ubiquitin-mediated regulation of

cytoplasmic β-catenin stability is well characterized (Marikawa and Elinson, 1998). In addition,

other steps of the WNT pathway upstream of β-catenin stabilization undergo regulation by the

ubiquitin system. Notably, several members of the NEDD4 family of E3 ubiquitin ligases

(SMURF1, ITCH, and NEDD4L) have been found to negatively regulate stability of WNT

pathway components. SMURF1 interacts with and ubiquitylates AXIN, inhibiting its interaction

with the WNT co-receptor LRP5/6 (Ding et al., 2013; Fei et al., 2014; Fei et al., 2013; Tanksley

et al., 2013; Wei et al., 2012). Both ITCH and NEDD4L promote degradation of DVL2 (Cadavid

et al., 2000; Ding et al., 2013; Fei et al., 2014; Fei et al., 2013; Wei et al., 2012). Additionally,

the deubiquitylase (DUB) USP34 was found to antagonize ubiquitylation of AXIN, promoting its

stabilization and function in the canonical WNT pathway (Lui et al., 2011). SMURF1 and

SMURF2 negatively regulate the WNT-PCP pathway by targeting WNT-PCP receptor

component Prickle1 for degradation (Narimatsu et al., 2009). Furthermore, DVL2 is also known

to undergo positive regulation by the ubiquitin system. For example, K63-linked ubiquitylation of

the N-terminal DAX domain, which is known to mediate dynamic polymerization of DVL2, has

been implicated as a positive regulator of DVL2 signal transduction (Schwarz-Romond et al.,

2007; Tauriello et al., 2010). These examples highlight the complex regulation of canonical and

non-canonical WNT pathways by ubiquitin conjugation and deconjugation machinery.

The NEDD4 family of E3 ubiquitin ligases is conserved from yeast to humans and has

been implicated in the ubiquitin-mediated endocytosis of many plasma membrane proteins

including surface receptors, ion channels, and nutrient transporters (David et al., 2013; He et al.,

2008; Hryciw et al., 2004; Kuratomi et al., 2005; Lee et al., 2011; Lin et al., 2011; MacGurn et

al., 2012). There are 9 NEDD4 family E3 ubiquitin ligases encoded in the human genome, each

with distinct physiological functions (Rotin and Kumar, 2009). While not all NEDD4 family E3

ubiquitin ligases have been characterized extensively, several have a strong preference for

catalyzing K63-linked polyubiquitin chains *in vitro (Ingham et al., 2004; Kim and Huibregtse,*

2009). Given the broad role of NEDD4 family E3 ubiquitin ligases in the regulation of membrane

trafficking and cell signaling, it is not surprising that they are often found to be dysregulated in

human diseases including cancer (Bernassola et al., 2008). NEDD4 family members have a

conserved domain structure including an N-terminal C2 domain that binds lipids, a C-terminal

HECT ubiquitin ligase domain, and 2-4 internal WW domains. The WW domains have a high

affinity for PY (Pro-Pro-x-Tyr) motifs and mediate interactions with substrates, adaptors, and

regulatory factors (Ingham et al., 2004). Here, we report that the NEDD4 family member WWP1

interacts with USP9X and DVL2, and we find that these associations are governed by

interactions between the WW domains of WWP1 and PY motifs present in both USP9X and

DVL2. Importantly, we find that these interactions control a ubiquitylation rheostat on DVL2 that

regulates both canonical WNT and non-canonical WNT-PCP activation in human breast cancer

cells. Thus, we find that this USP9X-DVL2-WWP1 axis is a key regulatory node in the WNT

pathway with the potential to influence both canonical and noncanonical WNT signaling in the

context of human cancer.

Results

The WW domains of WWP1 coordinate interactions with USP9X and DVL2

In an attempt to identify factors that regulate WWP1 function we used SILAC-based

quantitative proteomics to generate a WWP1 interaction profile in MDA-MB-231 cells (a triple

negative breast cancer cell line). This experiment identified interactions with various plasma

membrane (PM) and PM-associated proteins, factors associated with membrane trafficking,

signaling factors, and elements of the ubiquitin-proteasome system (**Table S1** and **FIG S1A**). To

validate results obtained by quantitative proteomics, we tested for co-purification of a subset of

putative WWP1 interacting proteins with FLAG-WWP1 stably expressed in MDA-MB-231 cells

(**FIG 1A**). This approach confirmed FLAG-WWP1 interactions with several putative interactors -

including USP9X, DVL2 and SPG20 – although some putative interactors such as ARRDC1 and

USP24 could not be confirmed by this approach. Considering the possibility that some putative

interacting proteins may artificially result from stable overexpression of FLAG-WWP1, we

performed native co-immunoprecipitation of WWP1 and identified interactions with USP9X and

DVL2 at endogenous expression levels (**FIG S1B**). Importantly, other studies have reported

interactions between USP9X and NEDD4 family E3 ubiquitin ligases – including SMURF1 (Xie

et al., 2013) and ITCH (Mouchantaf et al., 2006). Similarly, DVL2 was previously reported to

interact with other NEDD4 family E3 ubiquitin ligases – including WWP2 (Mund et al., 2015) and

NEDD4L (Ding et al., 2013; Wei et al., 2012). Notably, both USP9X and DVL2 contain multiple

PY motifs capable of binding the WW domains of NEDD4 family E3 ubiquitin ligases (**FIG 1B**).

Based on our findings and previous reports we decided to further investigate the basis of

physical interactions between WWP1, USP9X, and DVL2. Since USP9X and DVL2 both contain

multiple PY motifs we hypothesized that the WW domains of WWP1 might function to scaffold

these proteins into a complex. To test this, we generated WWP1 WW domain point mutant

variants designed to disrupt PY motif interactions (WxxP → FxxA) (Chen et al., 1997; Ermekova

et al., 1997; Gajewska et al., 2001) and analyzed DVL2 and USP9X interaction by co-

immunoprecipitation. We found that disruption of individual WW domains within WWP1 did not

diminish interaction with USP9X or DVL2 (**FIG S1C** and **S1D**). Indeed, we unexpectedly

observed that the *wwp1-ww2* mutant exhibited significantly greater interaction with DVL2 (**FIG**

S1C and **S1D**). Although individual WW domains were dispensable for interaction with DVL2

and USP9X, we found that a WWP1 variant with point mutations disrupting each WW domain

(*wwp1-4ww*) exhibited significantly decreased interaction with USP9X and complete loss of

interaction with DVL2 (**FIG 1C** and **1D**). Together, these findings indicate that *(i)* WW domain

scaffolding partially contributes to the interaction between WWP1 and USP9X, *(ii)* WW domain

interactions are required for WWP1 engagement with DVL2, and *(iii)* the WW domains of

WWP1 function redundantly for interaction with both DVL2 and USP9X.

To further probe the WW domain specificity underpinning WWP1 interactions with DVL2

and USP9X, we generated a panel of WWP1 variants containing only a single in-tact WW

domain (designated as *wwp1*-WW1, *wwp1*-WW2, *wwp1*-WW3, and *wwp1*-WW4). Importantly,

we found that any individual WW domain is sufficient to restore interaction with USP9X,

suggesting complete redundancy for this interaction (**FIG 1E**). In contrast, we found that only

WW1 and WW3 were sufficient to restore interaction with DVL2 (**FIG 1E** and **1F**), indicating that

WW2 and WW4 do not contribute to the interaction between WWP1 and DVL2. Furthermore,

the observation that *wwp1*-WW2 and *wwp1*-WW4 interact with USP9X but not DVL2 indicates

that the USP9X-WWP1 interaction occurs independent of DVL2 engagement. Overall, these

findings suggest that the WW domains of WWP1 scaffold interactions with both USP9X and

DVL2 in a manner that is redundant for USP9X and partially redundant for DVL2.

PY Motifs in USP9X and DVL2 Promote Interaction with WWP1

Based on the observation that WW domains of WWP1 are important for interacting with USP9X and DVL2, we hypothesized that PY motifs in USP9X and DVL2 are required to engage WWP1. DVL2 contains two PY motifs ($FPAY_{393}$ and $PPPY_{568}$) that flank the DEP domain (**FIG 1B**). To test if either of these PY motifs are required for DVL2 to bind WWP1, we generated DVL2 variants with point mutations at individual PY motifs ($FPAY_{393} \rightarrow FAAA_{393}$ and $PPPY_{568} \rightarrow PAPA_{568}$) and performed co-immunoprecipitation analysis using FLAG-DVL2 as bait. This analysis revealed that PY1 ($FPAY_{393}$) is dispensable for interaction with WWP1 while PY2 ($PPPY_{568}$) is required to interact with WWP1 (**FIG 1G**). Furthermore, these results indicate that disruption of the DVL2-WWP1 interaction does not affect DVL2 engagement with USP9X (**FIG 1G**). Taken together, our results indicate that the DVL2-WWP1 interaction is governed by binding of the PY2 motif of DVL2 to the WW1 or WW3 domains of WWP1.

We hypothesized that PY motifs in USP9X also contribute to its interaction with WWP1. USP9X contains four PY motifs ($QPQY_{409}$, $HPRY_{1017}$, $NPQY_{2437}$, and $APLY_{2515}$), with the last two occurring at the C-terminus of the protein (**FIG 1B**). To test if these C-terminal PY motifs are required for interaction with WWP1, we generated a USP9X variant deleted for the C-terminal portion of the protein ($USP9X_{1-2433}$, or *usp9x-Δpy3-4*). Importantly, deletion of the two C-terminal PY motifs resulted in complete loss of interaction with WWP1 (**FIG 1H**). However, point mutations in these C-terminal PY motifs ($NPQY_{2437} \rightarrow NAQA_{2437}$ (*py3*) and $APLY_{2515} \rightarrow AALA_{2515}$ (*py4*)) resulted in only partial loss of WWP1 interaction (**FIG S1E**). Notably, in each experiment loss of USP9X interaction with WWP1 did not affect its interaction with DVL2, indicating that the USP9X-DVL2 interaction occurs independently of WWP1. Thus, WWP1 engages both USP9X (partially) and DVL2 (primarily) via non-exclusive WW-PY interactions while DVL2 and USP9X exhibit the ability to interact independently of WWP1 (**FIG 1I**).

WWP1 and USP9X establish a ubiquitin rheostat on DVL2

DVL2 was previously reported to be a substrate of several NEDD4 family E3 ubiquitin

ligases including ITCH, NEDD4L, and NEDL1 (Ding et al., 2013; Miyazaki et al., 2004;

Narimatsu et al., 2009; Nethe et al., 2012; Wei et al., 2012) although both positive and negative

regulation of DVL2 by NEDD4 family members has been reported. However, WWP1 has not

previously been reported to regulate DVL2. Given the physical associations between WWP1

and DVL2 we hypothesized that DVL2 might be subject to ubiquitin conjugation by the E3

activity of WWP1. We found that recombinant WWP1 exhibited ubiquitin conjugation activity

toward HA-tagged DVL2 (purified from human STF-293 cell lysates) (**FIG 2A** and **FIG S2A-B**)

and this activity is lost in a lysine-deficient variant of DVL2 (*dvl2-K0*, **FIG S2C**). Importantly, we

also found that USP9X can reverse the ubiquitylation of DVL2 by WWP1 (**FIG 2B**), indicating

that DVL2 is a substrate for both WWP1 and USP9X. Strikingly, the K48-specific deubiquitylase

OTUB1 and the K63-specific deubiquitylase AMSH did not exhibit activity towards WWP1-

ubiquitylated DVL2 (**FIG 2B**), suggesting that WWP1-conjugated ubiquitin on DVL2 is unlikely to

contain K63- or K48- linked polymers.

Since WWP1 and USP9X both interact with and operate on DVL2, we hypothesized that

the coordination of WWP1 E3 ubiquitin ligase activity and USP9X deubiquitylase activity might

serve to establish a ubiquitin rheostat on DVL2. To test this, we incubated DVL2 with different

molar ratios of WWP1 and USP9X and analyzed the outcome. Importantly, we found that

different molar ratios of WWP1 and USP9X affected the extent of ubiquitylation on DVL2 (**FIG**

2C-D, **S2D** and **S2E**). These findings indicate that coordinated WWP1 and USP9X activities can

establish a ubiquitin rheostat on DVL2.

USP9X promotes canonical WNT activation

Given the established function of DVL2 as a key signal transducing protein in the WNT

pathway and given that NEDD4 family members have been reported to negatively regulate

WNT signaling (Tanksley et al., 2013) potentially via ubiquitylation of DVL2 (Ding et al., 2013),

we decided to test if USP9X regulates WNT signaling. We found that knockdown or knockout of

usp9x attenuated WNT activation as measured by a TopFLASH reporter assay (**FIG 3A** and

S3A) and by β–catenin stabilization (**FIG 3A**). The WNT signaling defect observed in *usp9x*

knockout cells was complemented by transient expression of wildtype FLAG-USP9X but not a

catalytic dead (C1556V, or CD) variant (**FIG 3B**), indicating that catalytic activity of USP9X is

required for canonical WNT activation. Furthermore, we found that treating cells with WP1130, a

small molecule known to inhibit the deubiquitylase activity of USP9X (Kapuria et al., 2010),

inhibited canonical WNT activation by WNT3a with an IC_{50} of 2.9μM (**FIG 3C**). This result is

consistent with our findings that USP9X is required for canonical WNT activation, however

WP1130 is known to target other deubiquitylases (USP24, USP5, USP14, and UCH37) (de Las

Pozas et al., 2018; Kapuria et al., 2010; Kushwaha et al., 2015; Peterson et al., 2015). Thus, we

cannot exclude the possibility that inhibition of multiple DUB activities drives loss of WNT

activation in the presence of WP1130. To further explore the possibility that USP9X activity

promotes WNT activation we transiently transfected expression vectors for FLAG-USP9X

(wildtype or catalytic dead) and measured WNT activation in the context of WNT3a stimulation.

Importantly, we found that transient expression of wildtype FLAG-USP9X, but not a catalytic-

dead variant, resulted in increased activation of the canonical WNT pathway (**FIG 3D**).

We hypothesized that loss of canonical WNT activation in the absence of USP9X may

be due to unchecked activity of WWP1 or other NEDD4 family E3 ubiquitin ligases. Importantly,

we found that the loss of canonical WNT activation in *usp9x* knockout cells was suppressed by

coordinate knockdown of WWP1 or NEDD4L but not WWP2 (**FIG 3E**) suggesting that hyper-

activation of multiple NEDD4 family members, including NEDD4L and WWP1, may attenuate

WNT signaling in the absence of USP9X. Furthermore, we found that transient overexpression

of wildtype WWP1, but not a catalytic dead variant (C890S), inhibited TopFLASH activation by

WNT3a (**FIG S3B** and **S3C**). Unexpectedly, we also found that expression of the *wwp1-ww2*

variant (harboring point mutations in WW2) inhibited canonical WNT activation to a greater

extent that wildtype WWP1 (**FIG S3C**) while expression of the *wwp1*-WW2 variant (where all

WW domains are disrupted except for WW2) exhibited hyperactivation of canonical WNT in the

presence of WNT3a (**FIG S3B**). Thus, ability of WWP1 to antagonize canonical WNT activation

correlates with its ability to bind DVL2 (**FIG 1E-F** and **S1C-D**). Taken together, these findings

indicate that USP9X promotes canonical WNT activation while WWP1 antagonizes canonical

WNT activation.

USP9X regulation of WNT requires DVL2 deubiquitylation

Given our finding that WWP1 and USP9X interact with and operate on DVL2, we

hypothesized that USP9X regulates canonical WNT activation by affecting the ubiquitylation

status of DVL2. To address the mechanism of canonical WNT regulation by USP9X, we first

tested if USP9X is required for β-catenin stabilization by LiCl – an inhibitor of GSK3. Importantly,

LiCl-mediated stabilization did not require USP9X (**FIG S4A**), indicating USP9X functions

upstream of the β-catenin destruction complex. Importantly, we found that knockdown or

knockout of USP9X resulted in decreased steady state abundance of DVL2 protein and mobility

shifts by SDS-PAGE consistent with hyper-ubiquitylation (**FIG 4A** and **S4B**). To identify post-

translational modifications on DVL2 regulated by USP9X we performed a SILAC-MS on FLAG-

DVL2 affinity purified from MDA-MB-231 cells and *usp9x* knockout equivalents. This analysis

resolved many phospho-peptides and ubiquitin remnant (diGly) peptides derived from DVL2

with four internal lysines exhibiting elevated ubiquitylation in the absence of USP9X (**FIG 4B**

and **FIG S4C-E**). Although ubiquitylation at N-terminal lysines of DVL2 has been reported

(Madrzak et al., 2015; Tauriello et al., 2010) the ubiquitin modification events described in this

study (K343, K428, K477 and K615) have not previously been reported.

To test if USP9X-mediated regulation of WNT signaling requires DVL2 ubiquitylation we

expressed DVL2 fused to the UL36 DUB domain (from human herpes virus) and found that this

(but not a catalytic dead UL36 fusion) suppressed the WNT activation defect in the absence of

USP9X (**FIG 4C**). These results indicate that USP9X regulates WNT signaling at (or in very

close proximity to) the level of DVL2. To explore the possibility that USP9X-mediated

deubiquitylation of DVL2 at specific sites is required for WNT activation we tested a panel of

DVL2 lysine mutants (K343R, K429R, K477R and K615R) but found expression of these

mutants was insufficient to suppress the WNT signaling defect observed in the absence of

USP9X (**FIG S4F**). We next considered the possibility that USP9X may regulate multiple

ubiquitylation events on DVL2, as suggested by SILAC-MS analysis (**FIG 4B**). Importantly, we

found that expression of a "lysine-less" DVL2 variant with all encoded lysine residues switched

to arginine (DVL2-K0) is sufficient to restore WNT activation in the absence of USP9X (**FIG 4D**).

Taken together, these results indicate that increased ubiquitylation of DVL2 is required for the

attenuation of WNT signaling observed in the absence of USP9X. Thus, the USP9X-mediated

deubiquitylation of DVL2 is critical for activation of canonical WNT signaling.

DVL2 ubiquitylation status regulates interactions with canonical and WNT-PCP factors

We hypothesized that the altered ubiquitylation status of DVL2 in the absence of USP9X

regulates its function as a signal transduction protein. To test this hypothesis, we first

characterized the DVL2 interaction network in MDA-MB-231 cells using SILAC-MS (**TABLE S2,**

TABLE S3, and **FIG 5A**). These experiments revealed two near-stoichiometric interactions with

the AP-2 clathrin adaptor complex and with the non-canonical WNT factors VANGL1 and cofilin

which are known to participate in the planar cell polarity (PCP) pathway (Luga et al., 2012).

Additionally, we identified several sub-stoichiometric interactions with NEDD4 family E3 Ub

ligases and proteasome subunits (**TABLE S2, TABLE S3,** and **FIG 5A**). Many of these

interactions had been previously reported (Angers et al., 2006; Yu et al., 2007). Several

interactions were confirmed by co-IP (**FIG 5B**) and co-localization by immunofluorescence

microscopy (**FIG S5A**). Consistent with previous reports (Yu et al., 2007; Yu et al., 2010) we

also found that knockdown of AP-2 complex subunits inhibited canonical WNT activation in a

TopFLASH assay (data not shown).

To quantify how DVL2 interactions are affected by ubiquitylation, we performed SILAC-MS

proteomic analysis comparing the interaction profile of catalytic active (light) or catalytic dead

(heavy) UL36-DVL2 fusions (also used in **FIG 4C**). This revealed that catalytic dead UL36-DVL2

fusion (which can be stably ubiquitylated) interacts more with proteasome subunits and

VANGL1/cofilin while the catalytic active UL36-DVL2 fusion (which is not stably ubiquitylated)

interacts more with the AP-2 complex (**FIG 5C**). Importantly, we noticed in this experiment a

small set of peptides derived from the DVL2-UL36 catalytic active fusion protein and absent

from the DVL2-UL36 catalytic dead fusion protein (**FIG S5B**). Interestingly, these outlier

peptides all correspond to tryptic events that can only occur in the absence of ubiquitylation (at

positions K428, K477 and K615) (**FIG S5C**) indicating that these sites are largely ubiquitin-

modified in the DVL2-UL36 catalytic dead fusion protein. We also used SILAC-MS to quantify

how DVL2 interactions are altered in *usp9x* knockout cells and found similar results: DVL2

interacted more with VANGL1 and cofilin in *usp9x* knockout cells (where DVL2 ubiquitylation is

elevated) but interacted more with AP-2 in the presence of USP9X (where DVL2 is less

ubiquitylated) (**FIG 5D**). These results indicate ubiquitylated DVL2 interacts more with

VANGL1/cofilin – components of the WNT-PCP pathway – while deubiquitylated DVL2 interacts

more with the AP-2 complex, which is critical for canonical WNT activation.

USP9X regulates cellular distribution of DVL2 and antagonizes WNT-PCP

Previous studies have reported that VANGL1 and cofilin localize to F-actin rich

projections in migrating MDA-MB-231 cells (Hatakeyama et al., 2014) and that DVL2 is required

for WNT5a-induced cell migration in these cells (Zhu et al., 2012). Based on our findings that

loss of USP9X results in increased ubiquitylation of DVL2 and increased interactions between

DVL2 and VANGL1/cofilin we hypothesized that loss of USP9X may result in re-localization of

DVL2 to actin-rich projections in MDA-MB-231 cells. To test this, we used immunofluorescence

microscopy to characterize FLAG-DVL2 distribution in MDA-MB-231 cells in the presence or

absence of USP9X. In MDA-MB-231 cells treated with control siRNAs, DVL2 exhibited a

punctate distribution throughout the cell body, reflective of previous descriptions of DVL2

localization (**FIG 6A, top panels**) (Schwarz-Romond et al., 2005) whereas MDA-MB-231

siUSP9X knockdown cells displayed a redistribution of DVL2 to actin-rich projections (**FIG 6A,**

middle panels). Quantification of the cellular distribution of DVL2 in the presence and absence

of USP9X reveals a significant increase in DVL2 in actin-rich projections upon loss of USP9X

(**FIG 6B**). Importantly, coordinate knockdown of WWP1 prevented this re-distribution of DVL2,

indicating that WWP1 activity is required for re-distribution of DVL2 to actin-rich projections in

the absence of USP9X (**FIG 6A, bottom panel** and **FIG 6B**). Thus, our data indicates that

USP9X promotes DVL2 localization to puncta that also contain AP-2 (**FIG S5A**), while loss of

USP9X activity results in WWP1-dependent re-distribution of DVL2 to actin-rich projections in

MDA-MB-231 breast cancer cells.

One feature of WNT5a-mediated PCP activation is Rho activation, which is DVL2-

dependent and required for induction of cell migration by the WNT-PCP pathway (Zhu et al.,

2012). Given our findings, we hypothesized that DVL2 re-distribution to actin-rich projections in

MDA-MB-231 cells lacking USP9X might result in Rho activation. Indeed, we found that MDA-

MB-231 cells normally exhibit about 35% of maximal Rho activation, while cells lacking USP9X

exhibit 50% maximal Rho activation (**FIG 6C-D**). Taken together, these findings indicate that

loss of USP9X in MDA-MB-231 cells results in re-distribution of DVL2 to actin-rich projections

and Rho activation, which is similar to what occurs during WNT5a-mediated activation of the

WNT-PCP pathway in these cells (Luga et al., 2012; Zhu et al., 2012).

USP9X regulates cell motility in a manner dependent on DVL2 ubiquitylation

6   VANGL1, cofilin, and DVL2 are known to coordinate WNT-PCP activation to regulate

actin dynamics and promote cell motility, migration and metastasis of MDA-MB-231 cells

(Katoh, 2005; Luga et al., 2012; Yang and Mlodzik, 2015; Zhu et al., 2012). Based on these

previous studies, we hypothesized that USP9X-mediated regulation of DVL2 may antagonize

WNT-PCP-mediated cellular motility. Strikingly, we found that MDA-MB-231 breast cancer cells

lacking USP9X exhibited a significant increase in motility compared to wildtype equivalents (**FIG**

6E-F). This increase in cell motility was complemented in cells transiently transfected with GFP-

USP9X (**FIG S6A**). Consistent with this observation, we also found that treatment of MDA-MB-

231 cells with WP1130 – a small molecule inhibitor of USP9X – also resulted in a significant

increase in the speed of cell motility (**FIG S6B**). Importantly, the increased cell speed observed

in *usp9x* knockout cells was dependent on VANGL1 (**FIG S6C**) indicating the increased motility

phenotype requires and in-tact WNT-PCP pathway. To test if the increase cell motility observed

in *usp9x* mutant cells was due to increased autocrine signaling, we treated cells with LGK974, a

porcupine inhibitor that blocks the secretion of many WNT ligands, and found this had no effect

on the speed of usp9x knockout MDA-MB-231 cells (**FIG S6D**). Taken together, these results

indicate that loss of USP9X leads to aberrant activation of PCP-mediated motility independent

of autocrine signaling.

23   We hypothesized that the elevated ubiquitylation of DVL2 might contribute to the

increased motility phenotype observed in usp9x knockout MDA-MB-231 cells. Consistent with

this hypothesis, we observed that knockdown of WWP1 suppressed the increased motility

phenotype of *usp9x* knockout cells (**FIG 7A**), although transient overexpression of WWP1 did

not alter speed of MDA-MB-231 cells (**FIG S7A**). Notably, MDA-MB-231 cells transiently

expressing GFP-DVL2 did not exhibit any change in cell motility compared to untransfected

control cells (**FIG S7B**), however expression of either wildtype or lysine-deficient (K0) GFP-

DVL2 fully suppressed the increased motility phenotype of *usp9x* knockout MDA-MB-231 cells

(**FIG S7C-D**). This result implicates DVL2 as a mediator of the cell motility phenotype observed

for *usp9x* knockout cells, but it does not inform as to the role of ubiquitylation since both

wildtype and lysine-deficient DVL2 fully suppress the *usp9x* knockout phenotype. To test the

role of ubiquitylation, we analyzed cell motility of MDA-MB-231 cells stably expressing DVL2-

UL36 (catalytic active and catalytic dead) DUB fusion proteins. Strikingly, we found that

knockdown of *USP9X* resulted in increased motility of control cells and cells expressing catalytic

dead DVL2-UL36 fusions, but not of cells expressing catalytic active DVL2-UL36 fusions (**FIG**

7B). This result reveals that the increased motility phenotype observed in the absence of

USP9X requires increased ubiquitylation of DVL2. Finally, we tested if DVL2 ubiquitylation is

required for WNT5a-induced cell motility in MDA-MB-231 cells. Interestingly, we found that

expression of catalytic active DVL2-UL36 resulted in a significant dampening of WNT5a-induced

cell motility compared to control cells expressing catalytic dead DVL2-UL36, although the

observed inhibition was partial (**FIG 7C**). These results indicate that USP9X is a critical

antagonist of PCP activation and cell motility in MDA-MB-231 cells and that this function is

dependent on the ubiquitylation of DVL2. Based on our findings, we propose that the balance of

USP9X and WWP1 activities toward DVL2 regulates WNT pathway specification, with USP9X-

mediated DVL2 deubiquitylation promoting canonical WNT signaling and WWP1-mediated

DVL2 ubiquitylation driving WNT-PCP activation and cell migration (**FIG 7D**).

Discussion

Dishevelled has long been known to play a key role in the transduction of WNT receptor signaling, yet how it is regulated to specify participation in either canonical or non-canonical relays is not well understood. Here, we report that the NEDD4 family E3 ubiquitin ligase WWP1 interacts with DVL2 and the deubiquitylase USP9X, and that these interactions govern a ubiquitylation rheostat on DVL2 critical for both canonical WNT and WNT-PCP activation. Specifically, our data indicates that *(i)* USP9X, WWP1 and DVL2 physically associate to establish a ubiquitin rheostat on DVL2, *(ii)* USP9X-mediated deubiquitylation of DVL2 is required for canonical WNT activation, and *(iii)* USP9X antagonizes activation of the WNT-PCP pathway (and thus cell motility) by limiting the accumulation of ubiquitylated DVL2. Based on the data, we propose that the USP9X-WWP1 axis biases DVL2 engagement with either the canonical WNT pathway or the noncanonical WNT-PCP pathway (**FIG 7D**).

A Novel Regulatory Axis for WNT Pathway Specification

The functional diversity of WNT signaling is a result of the complex network design that inter-weaves canonical and non-canonical elements to coordinate multiple specific signaling outputs. Thus, it is important to understand the network design features that couple canonical and non-canonical pathways and contribute to pathway specificity. The role of USP9X in WNT pathway specification draws parallels to diversin, an ankyrin repeat protein also implicated in WNT pathway specification. Knockout of diversin in mice resulted in defective WNT-PCP signaling but hyper-activated signaling through the canonical WNT pathway (Allache et al., 2015; Jones et al., 2014) – phenotypes that are opposite to *usp9x* knockout but suggest a similar mode of pathway coupling. Importantly, diversin has also been reported to be overexpressed in breast cancer cells and contributes to both proliferation and invasion (Yu et al., 2014). Shared use of certain signaling factors - like dishevelled and diversin - between canonical and non-canonical WNT pathways may facilitate mechanisms of pathway

1   specification based on inverse coupling – effectively operating as a "zero sum game". For

2   example, if a shared component like dishevelled or diversin is limiting for pathway activation

3   then engagement by one pathway may consequently prevent activation of the coupled pathway.

4   Thus, regulatory mechanisms that impact stability of shared factors or their interaction networks

5   have strong potential to contribute to pathway specification.

6   DVL2 has been reported to undergo both positive and negative regulation by the

ubiquitin system through the action of multiple different E3 ubiquitin ligases and deubiquitylating

enzymes. For example, both ITCH and NEDD4L have been shown to negatively regulate DVL2

stability by conjugating K48-linked polyubiquitin chains on DVL2, targeting it to the proteasome

for degradation (Ding et al., 2013; Liu et al., 2014; Marikawa and Elinson, 1998; Mukai et al.,

2010; Tauriello et al., 2010; Wei et al., 2012). On the other hand, K63-linked polyubiquitin

modifications of the DVL2 DIX domain were reported to enhance canonical WNT activation

(Tauriello et al., 2010) while other reports have suggested these modifications may antagonize

canonical WNT activation (Madrzak et al., 2015). The apparent regulation of DVL2 by multiple

members of the NEDD4 family could result from functional redundancy, but could also reflect

differential regulation based on expression in different contexts or different cell types. Our

finding that siRNA targeting of NEDD4L or WWP1 partially rescued WNT activation in the

absence of USP9X (**FIG 3E**) is consistent with a redundancy model, but it does not exclude the

possibility that USP9X antagonizes multiple exclusive ubiquitylation events on DVL2 mediated

by different E3 ubiquitin ligases. Indeed, our data indicates that functional regulation of DVL2 by

USP9X involves reversal of multiple distinct ubiquitylation events (**FIG 4B**) – possibly mediated

by multiple E3 ubiquitin ligases - since no single K→R mutation was capable of suppressing the

loss of WNT activation phenotype observed in the absence of USP9X (**FIG S4F**). Furthermore,

the *in vitro* analysis of USP9X activity presented here suggests that USP9X may be unique in its

ability to operate on WWP1-ubiquitylated DVL2, and that it may have activity toward atypical

ubiquitin polymers (**FIG 2B-D**). Additional biochemical and genetic studies will be required to fully elucidate how USP9X affects various distinct ubiquitylation events on DVL2 that may be mediated by different E3 ubiquitin ligases.

Previous studies have reported conflicting roles for USP9X with respect to canonical WNT activation. Specifically, USP9X has previously been reported to positively regulate the stability of β-catenin in glioma cells (Yang et al., 2016). However, USP9X was also reported to negatively regulate the stability of β-catenin in mouse neural progenitor cells (Premarathne et al., 2017). Our studies are consistent with a positive role for USP9X in canonical WNT pathway activation and our data indicate that USP9X functions upstream of the destruction complex (**FIG S4A**). Although our findings are consistent with a role for USP9X in the regulation of WNT at the level of DVL2 we cannot exclude the possibility of a context-dependent role for USP9X at other tiers of the WNT pathway. Furthermore, it is possible that expression of different NEDD4 family members, other USP9X-interactors, or substrate adaptors and targeting factors could modify USP9X function in a context-specific manner. Although previous studies have reported that Wnt5a-mediated activation of the WNT-PCP pathway in human breast cancer cells causes increased cell migration as a result of DVL2/Daam1-dependent activation of RhoA (Luga et al., 2012; Zhu et al., 2012) USP9X has not previously been linked to the regulation of these activities. Data presented in this study indicates that USP9X and WWP1 interact to regulate the ubiquitylation status of DVL2, which impacts DVL2 localization to actin-rich projections, Rho activation, and motility in breast cancer cells. These findings reveal a novel mode for regulation of WNT-PCP activation and motility of human breast cancer cells.

USP9X: a Complex Factor in Cancer

Given that USP9X has been implicated in the regulation of multiple receptor signaling and trafficking pathways, it is not surprising that in recent years USP9X dysregulation has been

linked to many different types of cancers. USP9X has been suggested to exhibit oncogenic

behavior in some cancer contexts including multiple myeloma, melanoma, lung cancer, and

breast cancer (Kushwaha et al., 2015; Li et al., 2017; Potu et al., 2017; Schwickart et al., 2010).

In contrast, USP9X has also been proposed to have tumor suppressor activity in other cancers

including pancreatic ductal adenocarcinoma (Pérez-Mancera et al., 2012) and colorectal

adenocarcinoma (Khan et al., 2018). Recently, USP9X has been shown to be required for the

attenuation of EGFR signaling (Savio et al., 2016) which is suggestive of a tumor suppressor

function. Taken together, these reports suggest a complex role for USP9X in cancer

progression that may be highly context-dependent. Our findings suggest that USP9X functions

together with WWP1 to regulate a ubiquitin rheostat on DVL2. The ubiquitylation status of DVL2

determines its engagement with either canonical or non-canonical WNT factors. Based on these

findings we hypothesize that in some cellular contexts increased USP9X activity may sensitize

cells to activation of the canonical WNT pathway (**FIG 3D**), which promotes proliferation and

tumor growth, while simultaneously antagonizing the non-canonical WNT-PCP pathway which is

involved in cell motility and metastasis (**FIG 6C-F**). These results indicate that dysregulation of

USP9X can function to promote canonical WNT-driven proliferation while simultaneously

guarding against migration and metastasis mediated by the WNT-PCP pathway. Based on

these findings, we propose that small molecule inhibitors of USP9X may be effective as

antagonists of WNT-based proliferation (**FIG 3C**) but they may also ultimately promote migration

and metastasis (**FIG S6B**).

Regulation of cell signaling pathways by DUB-E3 interactions

The ubiquitin proteasome system contributes to the tight spatial and temporal regulation

of protein trafficking, localization, and stability. Recently, several examples of interactions

between E3 ubiquitin ligases and DUBs have been reported, raising questions about the

regulatory significance and design logic of such interactions. In principle, DUB-E3 interactions

could facilitate a number of useful design features, including E3 antagonization, E3

deubiquitylation, or polyubiquitin chain editing on substrates. E3 antagonization has the

potential to result in a futile cycle, but coordinated regulation of ligase and DUB activities could

also establish a ubiquitin rheostat on a shared substrate. Such a rheostat has been reported to

occur in the context of ER quality control machinery, where DUB activities can sharpen the

distinction between folded and misfolded membrane proteins (Zhang et al., 2013). Another

striking example involves the DUB USP7, which was shown to interact with the MAGE-L2-

TRIM27 E3 ubiquitin ligase to regulate the ubiquitylation status of WASH, a shared substrate,

on the endosome (Wu et al., 2014). Interestingly, USP7 has been reported to interact with

multiple E3 ubiquitin ligases (Kim and Sixma, 2017) but the functional significance of many of

these interactions remain unclear. E3-DUB interactions have also been reported to protect E3

ubiquitin ligases from degradation, thus promoting E3 activity. For example, the DUB USP19

promotes cellular proliferation by stabilizing KPC1 (also known as RNF123), a ubiquitin ligase

for the cyclin-dependent kinase inhibitor p27Kip1 (also known as CDKN1B) (Lu et al., 2009). A

striking example of polyubiquitin chain editing occurs with the protein A20, a potent inhibitor of

the NFkB signaling pathway that contains both a DUB domain and an E3 ubiquitin ligase

domain (Wertz et al., 2004). The DUB domain of A20 removes a K63-linked polyubiquitin chain

from the Receptor Interacting Protein (RIP), a protein essential for NFkB signaling.

Subsequently, the E3 ubiquitin ligase domain of A20 conjugates a K48-linked polyubiquitin

chain to RIP targeting it for degradation by the proteasome - thus inhibiting the NFkB pathway.

Thus, there are numerous examples of E3-DUB interactions that establish ubiquitin rheostats on

substrates, protect and stabilize E3 ubiquitin ligases, and coordinate activities to achieve

polyubiquitin chain editing in signaling complexes.

    We have identified a novel E3-DUB interaction that functions to provide layered

regulation of the WNT pathway. Our results indicate that the activities of WWP1 and USP9X are

antagonistic and function to establish a ubiquitin rheostat on DVL2 that is a critical regulator of

WNT pathway specification. Functionally, USP9X antagonizes WWP1 activity towards DVL2

(**FIG 2C-D**), although our data cannot exclude the possibility that WWP1-USP9x interactions

function in a chain-editing capacity on polyubiquitinated DVL2. Ultimately, rigorous biochemical

analysis dissecting mechanisms that regulate the balance of different ubiquitylation and

deubiquitylation activities toward DVL2 will be critical to understand how its specification for

either canonical or noncanonical WNT signaling pathways is determined.

STAR Methods section

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Jason MacGurn (Jason.a.macgurn@vanderbilt.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell lines:

1. HEK239 cells

Source: American Type Culture Collection (ATCC), Catalog #: CRL-1573

Culture/growth conditions: Cultured in DMEM with 10% FBS and 1%

penicillin/streptomycin at 37ºC in 5% CO_2

Sex: Female

Authentication: Yes (ATCC authenticated)

2. MDA-MB-231 cells

Source: ATCC, Catalog #: HTB-26

Culture/growth conditions: Cultured in RPMI with 10% FBS and 1%

penicillin/streptomycin at 37ºC in 5% CO_2

Sex: Female

Authentication: Yes (ATCC authenticated)

3. HEK293 STF cells

Source: ATCC, Catalog #: HEK 293 STF (ATCC® CRL-3249™)

Culture/growth conditions: Cultured in RPMI with 10% FBS and 1%

penicillin/streptomycin at 37ºC in 5% CO_2

Sex: Female

Authentication: Yes (ATCC authenticated)

METHOD DETAILS

Assays for measuring WNT activation

Topflash reporter assays in HEK293 STF cells were performed as follows: HEK293 STF cells

were seeded on 24-well plates at ~50% confluency. Cells were treated with 100ng/mL purified

recombinant WNT3a (R&D Systems) and 100ng/mL purified recombinant R-spondin (R&D

systems) 24 hours prior to lysis with 1X Passive Lysis buffer (Promega). Luciferase activity was

measured using the Steady-Glo luciferase Assay (Promega). Luciferase signal was normalized

to viable cell number using Cell-Titer Glo Assay (Promega). For siRNA knockdown studies, cells

were treated for 72 hours prior to lysis, for inhibitor studies cells were treated for 24 hours prior

to lysis, and for overexpression studies cells were transfected with plasmid DNA for 48 hours

prior to lysis.

Topflash reporter assays in MDA-MB-231 cells were performed as follows: Cells were plated,

co-transfected with TOPflash and Renilla expression plasmids at 24 hours and lysed at 48

hours using the Dual-Glo Luciferase Assay system (Promega). Luciferase signal was

normalized to co-transfected Renilla expression. Data were normalized to a positive control (set

at 100% activation). Assays were performed in triplicate and repeated at least 3 times.

B-catenin stabilization assay: HEK293 STF cells were cultured in Dulbecco's modified Eagle's

Medium (DMEM) and activated with WNT3A ligand or LiCl for 24 hours. Then cells were lysed

and nuclear/cytoplasmic fractionation was performed as previously described (Thorne et al.,

2010).

__Transfections__

Plasmid and siRNA transfections were performed using Lipojet (SignaGen, SL100468) and

Lipofectamine LTX with PLUS reagent (Thermo Fisher, 15338100) according to manufacturer's

protocol.

__Immunoblots__

Whole-cell lysates were generated by lysing cells in Laemmli Sample Buffer (Bio-Rad) with 5%

βME and subsequently boiling samples at 95ºC for 5 minutes. Proteins were analyzed by SDS-

PAGE and immunoblotting using the LiCor Odyssey reagents and protocols. Fluorescence

signaling was detected using an Odyssey CLx Infrared Scanner (LiCor).

__Cell lines__

HEK293 STF cells and MDA-MB-231 cells were purchased from the American Type Culture

Collection (ATCC). HEK293 STF cells were cultured in DMEM with 10% FBS and 1%

penicillin/streptomycin at 37ºC in 5% CO2. MDA-MB-231 cells were cultured in DMEM with 10%

FBS and 1% penicillin/streptomycin at 37ºC in 5% CO2. MDA-MB-231 cells stably expressing

FLAG-WWP1, FLAG-DVL2, FLAG-DUB-DVL2, or FLAG-DUB*-DVL2 were generated using the

pQCXIP retroviral vector system. MDA-MB-231 *usp9x* knockout cells were generated using a

CRISPR/Cas-9 genome editing system.

__SILAC-based Quantitative Proteomic Analysis__

Quantitative mass spectrometry analysis by SILAC was performed on MDA-MB-231 cells stably

expressing the indicated FLAG-tagged substrates. These cells were cultured in the presence of

either heavy or light isotopes (lysine and arginine) for at least 6 doublings to ensure

incorporation of these isotopes into 100% of the proteome. Affinity purification was performed as

previously described (Lee et al., 2017). Eluted proteins were digested with 1µg Trypsin Gold

(Promega) at 37 F overnight. Digested peptides were cleaned-up on a Sep-Pak C18 column.

Purified peptides were then dried, reconstituted in 0.1% trifluoroacetic acid, and analyzed by

LC-MS/MS using an Orbitrap XL mass spectrometer. Database search and SILAC quantitation

were performed using MaxQuant software.

Co-Immunoprecipitation Studies

For coimmunoprecipitation assays, cells were washed with cold PBS and lysed with a

nondenaturing lysis buffer (NDLB): 50mM tris (pH 7.4), 300mM NaCl, 5mM EDTA, 1% Triton X-

100, cOmplete protease inhibitor cocktail (Roche), Phosphatase inhibitor cocktail (Roche), 1mM

PMSF, 1mM Phenanthroline, 10mM Iodoacetamide, and 20μM MG132. Lysates were diluted to

1mg/mL with NDLB and incubated with Sigma FLAG EZ-view beads for 2 hours with rotation at

4°C. Beads were washed 3 times with cold NDLB and once with cold PBS. Proteins were then

eluted from beads using sample buffer and samples were processed for SDS-PAGE followed by

immunoblotting.

Fluorescence Microscopy

MDA-MB-231 cells were seeded on coverslips at 50% confluency. At 24 hours after seeding,

cells were transfected with FLAG-Dvl2 using LipoJet Transfection Reagent (Signagen). 72

hours after seeding cells were fixed and permeabilized using 4% paraformaldehyde (PFA) and

immunofluorescence block buffer (PBS + 0.1% Triton-x + 10% FBS). Cells were then labeled

with primary antibodies for 1 hour followed by a 1 hour incubation with alexa-fluor conjugated

secondary antibodies. Coverslips were mounted on slides using ProLong AntiFade mountant

(Thermo). Microscopy images were acquired using a DeltaVision Elite system (GE Healthcare)

and processed using SoftWoRx software (GE Healthcare, Chicago, IL).

In Vitro Ubiquitin Conjugation and Deubiquitylation Assays

Due to the limited solubility of full-length DVL2 in solution, in all *in vitro* conjugation and

deconjugation assays HA-DVL2 was purified from human HEK293 cells using magnetic anti-HA

beads (Pierce #88836) and all conjugation/deconjugation reactions were performed on HA-

DVL2 bound to anti-HA beads. Lysis and purification were performed as follows: 24 hours after

seeding, HEK293 cells were transfected with HA-Dvl2 and 48 hours post-transfection cells were

lysed in NDLB and clarified by centrifugation. Lysates bound anti-HA magnetic beads overnight

at 4ºC with rotation. Beads were washed 3 times in NDLB and 1 time in cold 1X PBS before

being resuspended in conjugation buffer (40 mM Tris pH 7.5, 10 mM MgCl2, and 0.6 mM DTT)

for immediate use in conjugation/deconjugation reactions.

For the conjugation reactions, 56 nM GST-Ube1 (Boston Biochem, Cambridge, MA, E-300),

0.77 mM UBE2D3, 2.3 mM Ub, HA-DVL2, and 60 nM FLAG-WWP1 (Sigma, SRP0229) were

incubated in conjugation buffer (40 mM Tris pH 7.5, 10 mM MgCl2, and 0.6 mM DTT) with 1 mM

ATP, at 37°C. Reactions were initiated by the addition of FLAG-WWP1. Samples were removed

at indicated time-points, boiled in 2x Laemmli sample buffer for 10 min, and analyzed by blotting

for HA-DVL2 with DVL2 antibody (Cell Signaling Technology, #3216) using goat anti-rabbit

secondary (IRDye 800 CW, LI-COR Biosciences, Lincoln, NE). Quantification was performed

using ImageStudioLite software (LI-COR).

For deconjugation reactions, polyubiquitinated substrate (HA-DVL2) was generated using the

conjugation assay described above within one exception in that 0.1 mM ATP was used. The

conjugation reaction was stopped by incubation with 0.75 units Apyrase in a 90 µl reaction at

30°C for 1 hr. For the deconjugation reaction, 100 nM His6-USP9X (Boston Biochem E-552),

100nM OTUB1, or 100nM AMSH was added to conjugated HA-DVL2 as well as 10X DUB buffer

(0.5 M Tris pH 7.5, 500 mM NaCl, 50mM DTT) to 1X. Reactions were initiated by the addition of

His6-USP9X protein. The reaction was incubated at 37ºC. Samples were removed at indicated

time-points and boiled in 2x Laemmli sample buffer. HA-DVL2 levels were determined as

described in the conjugation assay.

For Rheostat experiments, HA-DVL2 substrate was resuspended in conjugation buffer with

1mM ATP and treated with 56 nM GST-Ube1 , 0.77 mM UBE2D3, 2.3 mM Ub, and indicated

concentrations of FLAG-WWP1 and His-USP9X for 60 minutes at 37°C. Samples were then

boiled in 2x Lammli sample buffer for 10 minutes and analyzed by blotting for HA-DVL2 with

DVL2 antibody (CST #3216). Line profile analysis was performed using ImageJ (NIH).

Migration Assays

Cell culture dishes were coated with 5 µg/ml rat tail ColI (Corning #354236) in PBS for 1 h at

37°C then cells were plated at low density and incubated at 37°C for 1 hour in cell culture

medium. Cells were kept at 37°C in SFM4MAb medium (HyClone, Logan, UT) supplemented

with 2% FBS at pH 7.4 during imaging using phase-contrast and fluorescence microscopy

techniques. Images were acquired at 5 min intervals for 6 h, using DeltaVision Elite system

software (GE Healthcare) and cell speed was quantified using SoftWoRx software (GE

Healthcare, Chicago, IL). Wind rose plots were generated by transposing x,y coordinates of cell

tracks to a common origin.

Rho Activation Assays

Rho activation assays were performed using the Rho Activation Assay Kit (Millipore Sigma #17-

924). MDA-MB-231 cells (WT and *usp9x* KO cells) were grown until confluent, washed in ice-

cold 1XPBS, lysed in ice-cold 1X Mg^{2+} Lysis/Wash buffer (MLB) (Millipore Sigma #20-168),

scraped and collected in microfuge tubes on ice, and incubated at 4°C for 15 minutes with

agitation. Lysates were then cleared of cellular debris by centrifugation at 14,000xg for 5

minutes at 4°C. Clarified lysate was then bound to Rho Assay Reagent (Millipore Sigma #14-

383) at 4°C for 45 minutes with gentle agitation. Agarose beads were then washed 3 times with

1X MLB and proteins were eluted in 2x Laemmli reducing sample buffer (BioRad #1610737) by

boiling at 95°C for 5 minutes. Samples were fractionated by SDS-PAGE and analyzed by

blotting using the Anti-Rho (-A, -B, -C) clone 55 antibody (Millipore Sigma #05-778).

GTPγS/GDP loading for positive and negative controls were prepared as described above with

an additional 30 minute incubation at 30°C with agitation of cell lysate in the presence of 10mM

EDTA and either 100uM GTPγS (Millipore Sigma #20-176) or 1mM GDP (Millipore Sigma #20-

177). This step was performed prior to binding the Rho Assay Reagent.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical Analysis

Detailed statistical analysis can be found in the accompanying Statistical Reporting Documents

for each figure. For each statistical analysis a Student's T-test was used to test for a statistically

significant difference between the means of the two variables of interest. The alpha value for

each experiment was set at 0.05 and a p-value was calculated using the Student's t-test

function in Microsoft Excel to determine statistical significance. Each figure with statistical

analysis represents n≥3 for all coIP and luciferase assay experiments, where n represents

biological replicates. For quantitation of all microscopy-based data an n≥25 was used where n

represents single cells. In all figures including statistics, error bars represent standard deviation

of the mean. Details of statistical analysis can also be found in figure legends.

1 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Monoclonal ANTI-FLAG® M2 antibody produced in mouse	Millipore	Product #: F1804
USP9X/Y Antibody (E-12)	Santa Cruz	Cat #: sc-365353
WWP1 monoclonal antibody (M01), clone 1A7	Abnova	Catalog #: H00011059-M01
Dvl2 Antibody #3216	Cell Signaling Technology	Product # 3216S
GAPDH (14C10) Rabbit mAb #2118	Cell signaling technology	Product # 2118S
USP24 (S-18) antibody	Cell Signaling Technology	Catalog #: sc-82080
Anti-beta Catenin antibody [E247]	Abcam	ab32572
Tubulin antibody	Vanderbilt Antibody and Protein Resource Core	N/A
NEDD4L Antibody #4013	Cell Signaling Technology	Product # 4013S
WWP2 (AIP2) Antibody (A-3)	Santa Cruz	Catalog #: sc-398090
HA antibody	Vanderbilt Antibody and Protein Resource Core	N/A
Alpha adaptin antibody (AP2A1) AC1-M11	Thermo Fisher	Cat #: MA3-061
Anti-AP2M1 antibody [EP2695Y]	Abcam	ab75995
Anti-VANGL1	Sigma	Product #: HPA025235
Cofilin (D3F9) XP Rabbit mAb	Cell Signaling Technology	D3F9
Anti-Rho (-A, -B, -C) clone 55	Millipore Sigma	Catalog # 05-778
Anti-HA magnetic beads	Pierce	Catalog # 88836
Bacterial and Virus Strains		
Biological Samples		
Chemicals, Peptides, and Recombinant Proteins		
WP1130	UBPBio	Cat #: F2120

MG-132	APExBIO	Cat #: A2585
Phenanthroline	Sigma-Aldrich	Cat #: P9375
Iodoacetamide	Sigma-Aldrich	Cat #: I1149
LGK-974	Selleck Chem	Catalog No.S7143
Recombinant Human/mouse Wnt5a	R&D System	Cat #: 645-WN
WWP1 Active human recombinant, expressed in baculovirus infected insect cells	Sigma-Aldrich	Prod #: SRP0229
USP9x-His6, isoform 2, human recombinant	Boston Biochem	Cat. # E-552
Recombinant OTUB1	UBPBio	Catalog #: H3000
AMSH, human recombinant	Boston Biochem	Cat. # E-548B
GST-Ubiquitin E1 Enzyme (UBE1), S.cer. recombinant	Boston Biochem	Cat. # E-300
Recombinant Human UbcH5c/UBE2D3	Boston Biochem	Cat. # E2627
Recombinant Mouse Wnt-3a	R&D Systems	Cat #: 1324-WN
Recombinant Mouse R-spondin	R&D Systems	Cat #: 3474-RS
Corning® Collagen I, Rat Tail	Corning	Product #: 354236
Heavy Lysine: L-Lysine-13C6,15N2 hydrochloride	Sigma-Aldrich	Product #: 608041
Heavy Arginine: L-Arginine-13C6,15N4 hydrochloride	Sigma-Aldrich	Product #: 608033
Critical Commercial Assays		
Steady Glo Luciferase Assay System	Promega	Catalog #: E2510
Dual Glo Luciferase Assay System	Promega	Catalog #: E2920
Rho Activation Assay Kit	Millipore Sigma	Catalog #: 17-294
Deposited Data		
Experimental Models: Cell Lines		
Human cells: MDA-MB-231 cells	ATCC	Product #: MDA-MB-231 (ATCC® HTB-26™)
Human cells: HEK293 cells	ATCC	Product #: 293 [HEK-293] (ATCC® CRL-1573™
Human cells: HEK 293 STF	ATCC	Product #: HEK 293 STF (ATCC® CRL-3249)

Human cells: MDA-MB-231 cells stably expressing FLAG-DVL2, FLAG-DUB-DVL2, or FLAG-DUB*-DVL2	This paper	N/A
Human cells: MDA-MB-231 *usp9x* KO cells +/- FLAG-DVL2	This paper	N/A
Experimental Models: Organisms/Strains		
Oligonucleotides		
Control siRNA: orKKJ1-s, nonsilencing/pGL2 sense, CGUACGCGGAAUACUUCGAUU	This paper	N/A
Usp9x siRNA #1 sequence: AGAAAUCGCUGGUAUAAAUU	This paper	N/A
Usp9x siRNA #2 sequence: ACACGAUGCUUUAGAAUUUU	This paper	N/A
Nedd4L siRNA #1 sequence: GCUAGACUGUGGAUUGAGUUU	This paper	N/A
Nedd4L siRNA #2 sequence: UGAGGAUCAUUUGUCCUACUU	This paper	N/A
WWP1 siRNA #1 sequence: GAAGUCAUCUGUAACUAAA**UU**	This paper	N/A
WWP1 siRNA #2 sequence: GCAGAGAAAUACUGUUUAU**UU**	This paper	N/A
WWP2 siRNA #1 sequence: GGGAGAAGAGACAGGACAA**UU**	This paper	N/A
WWP2 siRNA #2 sequence: CAGGAUGGGAGAUGAAAUA**UU**	This paper	N/A
VANGL1 Silencer Select siRNA	Thermo Fisher	Catalog #: 4392420
Recombinant DNA		
Plasmid: FLAG-Wwp1	This paper	N/A
Plasmid: FLAG-Wwp1-ww1 (W377F, P380A)	This paper	N/A
Plasmid: FLAG-Wwp1-ww2 (W409F, P412A)	This paper	N/A
Plasmid: FLAG-Wwp1-ww3 (W484F, P487A)	This paper	N/A
Plasmid: FLAG-Wwp1-ww4 (F524A, P527A)	This paper	N/A
Plasmid: FLAG-Wwp1-4ww (W377F, P380A, W409F, P412A, W484F, P487A, F524A, P527A)	This paper	N/A
Plasmid: FLAG-Wwp1-WW1 (W409F, P412A, W484F, P487A, F524A, P527A)	This paper	N/A
Plasmid: FLAG-Wwp1-WW2 (W377F, P380A, W484F, P487A, F524A, P527A)	This paper	N/A
Plasmid: FLAG-Wwp1-WW3 (W377F, P380A, W409F, P412A, F524A, P527A)	This paper	N/A

Plasmid: FLAG-Wwpw-WW4 (W377F, P380A, W409F, P412A, W484F, P487A)	This paper	N/A
Plasmid: FLAG-DVL2 PY1: $FPAY_{393} \rightarrow FAAA_{393}$	This paper	N/A
Plasmid: FLAG-DVL2 PY2: $PPPY_{568} \rightarrow PAPA_{568}$	This paper	N/A
Plasmid: FLAG-DVL2 2PY: $FPAY_{393} \rightarrow FAAA_{393}$ and $PPPY_{568} \rightarrow PAPA_{568}$	This paper	N/A
Plasmid: $USP9X_{1-2433}$, or *usp9x-Δpy3-4* $NPQY_{2437} \rightarrow NAQA_{2437}$ (*py3*) and $APLY_{2515} \rightarrow AALA_{2515}$ (*py4*)	This paper	N/A
Plasmid: UL36-DVL2	Stringer & Piper (2011) doi: 10.1083/jcb.201008121 Kettenhorn et al. (2005) Doi: 10.1016/j.molcel.2005.07.003	N/A
Plasmid: UL36 (C65A) catalytic dead-DVL2	Stringer & Piper (2011) doi:10.1083/jcb.201008121 Kettenhorn et al. (2005) Doi: 10.1016/j.molcel.2005.07.003	N/A
Plasmid: FLAG-DVL2-K0 (all lysines K→R)	This paper	N/A
Plasmid: FLAG-USP9X	This paper	N/A
Plasmid: FLAG-USP9X (C1556V) catalytic dead	This paper	N/A
Plasmid: FLAG-WWP1 (C890S) catalytic dead	This paper	N/A
Plasmid: GFP-DVL2	This paper	N/A
Plasmid: GFP-DVL2-K0 (all lysines K→R)	This paper	N/A
Plasmid: GFP-WWP1	This paper	N/A
Plasmid: GFP-USP9X	This paper	N/A
Software and Algorithms		
SoftWorx	GE	
MaxQuant	Max Planck Institute of Biochemistry	
ImageJ	NIH	
ImageStudioLite software	LICORE	
Other		

Nielsen et al.

Acknowledgements

We are very grateful to K. Rose for helpful advice regarding technical aspects and analysis of

quantitative proteomic data. We are also grateful to E. Lee, K. Saito-Diaz, V. Ng for reagents

and technical advice. We also thank T. Graham, S. M. Ahmed, S. Lee, and N. Hepowit for

critical reading of this manuscript. This research was supported by NIH grant R00 GM101077

(to J.A.M.) and GI SPORE grant in GI Cancer (P50 CA095103).

1 References

Allache, R., Wang, M., De Marco, P., Merello, E., Capra, V., and Kibar, Z. (2015). Genetic studies of
ANKRD6 as a molecular switch between Wnt signaling pathways in human neural tube defects. Birth
Defects Res A Clin Mol Teratol *103*, 20-26.
Angers, S., Thorpe, C.J., Biechele, T.L., Goldenberg, S.J., Zheng, N., MacCoss, M.J., and Moon, R.T. (2006).
The KLHL12-Cullin-3 ubiquitin ligase negatively regulates the Wnt-beta-catenin pathway by targeting
Dishevelled for degradation. Nat Cell Biol *8*, 348-357.
Bernassola, F., Karin, M., Ciechanover, A., and Melino, G. (2008). The HECT family of E3 ubiquitin ligases:
multiple players in cancer development. Cancer Cell *14*, 10-21.
Cadavid, A.L., Ginzel, A., and Fischer, J.A. (2000). The function of the Drosophila fat facets
deubiquitinating enzyme in limiting photoreceptor cell number is intimately associated with
endocytosis. Development *127*, 1727-1736.
Chen, H.I., Einbond, A., Kwak, S.J., Linn, H., Koepf, E., Peterson, S., Kelly, J.W., and Sudol, M. (1997).
Characterization of the WW domain of human yes-associated protein and its polyproline-containing
ligands. J Biol Chem *272*, 17070-17077.
David, D., Nair, S.A., and Pillai, M.R. (2013). Smurf E3 ubiquitin ligases at the cross roads of oncogenesis
and tumor suppression. Biochim Biophys Acta *1835*, 119-128.
de Las Pozas, A., Reiner, T., De Cesare, V., Trost, M., and Perez-Stable, C. (2018). Inhibiting Multiple
Deubiquitinases to Reduce Androgen Receptor Expression in Prostate Cancer Cells. Sci Rep *8*, 13146.
Ding, Y., Zhang, Y., Xu, C., Tao, Q.H., and Chen, Y.G. (2013). HECT domain-containing E3 ubiquitin ligase
NEDD4L negatively regulates Wnt signaling by targeting dishevelled for proteasomal degradation. J Biol
Chem *288*, 8289-8298.
Ermekova, K.S., Zambrano, N., Linn, H., Minopoli, G., Gertler, F., Russo, T., and Sudol, M. (1997). The
WW domain of neural protein FE65 interacts with proline-rich motifs in Mena, the mammalian homolog
of Drosophila enabled. J Biol Chem *272*, 32869-32877.
Fei, C., He, X., Xie, S., Miao, H., Zhou, Z., and Li, L. (2014). Smurf1-mediated axin ubiquitination requires
Smurf1 C2 domain and is cell cycle-dependent. J Biol Chem *289*, 14170-14177.
Fei, C., Li, Z., Li, C., Chen, Y., Chen, Z., He, X., Mao, L., Wang, X., Zeng, R., and Li, L. (2013). Smurf1-
mediated Lys29-linked nonproteolytic polyubiquitination of axin negatively regulates Wnt/β-catenin
signaling. Mol Cell Biol *33*, 4095-4105.
Gajewska, B., Kamińska, J., Jesionowska, A., Martin, N.C., Hopper, A.K., and Żołądek, T. (2001). WW
domains of Rsp5p define different functions: determination of roles in fluid phase and uracil permease
endocytosis in Saccharomyces cerevisiae. Genetics *157*, 91-101.
Glinka, A., Dolde, C., Kirsch, N., Huang, Y.L., Kazanskaya, O., Ingelfinger, D., Boutros, M., Cruciat, C.M.,
and Niehrs, C. (2011). LGR4 and LGR5 are R-spondin receptors mediating Wnt/β-catenin and Wnt/PCP
signalling. EMBO Rep *12*, 1055-1061.
Hatakeyama, J., Wald, J.H., Printsev, I., Ho, H.Y., and Carraway, K.L. (2014). Vangl1 and Vangl2: planar
cell polarity components with a developing role in cancer. Endocr Relat Cancer *21*, R345-356.
He, Y., Hryciw, D.H., Carroll, M.L., Myers, S.A., Whitbread, A.K., Kumar, S., Poronnik, P., and Hooper, J.D.
(2008). The ubiquitin-protein ligase Nedd4-2 differentially interacts with and regulates members of the
Tweety family of chloride ion channels. J Biol Chem *283*, 24000-24010.
Hryciw, D.H., Ekberg, J., Lee, A., Lensink, I.L., Kumar, S., Guggino, W.B., Cook, D.I., Pollock, C.A., and
Poronnik, P. (2004). Nedd4-2 functionally interacts with ClC-5: involvement in constitutive albumin
endocytosis in proximal tubule cells. J Biol Chem *279*, 54996-55007.
Ingham, R.J., Gish, G., and Pawson, T. (2004). The Nedd4 family of E3 ubiquitin ligases: functional
diversity within a common modular architecture. Oncogene *23*, 1972-1984.

Jones, C., Qian, D., Kim, S.M., Li, S., Ren, D., Knapp, L., Sprinzak, D., Avraham, K.B., Matsuzaki, F., Chi, F.*,*
et al. (2014). Ankrd6 is a mammalian functional homolog of Drosophila planar cell polarity gene diego
and regulates coordinated cellular orientation in the mouse inner ear. Dev Biol *395*, 62-72.
Kapuria, V., Peterson, L.F., Fang, D., Bornmann, W.G., Talpaz, M., and Donato, N.J. (2010).
Deubiquitinase inhibition by small-molecule WP1130 triggers aggresome formation and tumor cell
apoptosis. Cancer Res *70*, 9265-9276.
Katoh, M. (2005). WNT/PCP signaling pathway and human cancer (review). Oncol Rep *14*, 1583-1588.
Khan, O.M., Carvalho, J., Spencer-Dene, B., Mitter, R., Frith, D., Snijders, A.P., Wood, S.A., and Behrens,
A. (2018). The deubiquitinase USP9X regulates FBW7 stability and suppresses colorectal cancer. J Clin
Invest.
Kim, H.C., and Huibregtse, J.M. (2009). Polyubiquitination by HECT E3s and the determinants of chain
type specificity. Mol Cell Biol *29*, 3307-3318.
Kim, R.Q., and Sixma, T.K. (2017). Regulation of USP7: A High Incidence of E3 Complexes. J Mol Biol *429*,
3395-3408.
Kuratomi, G., Komuro, A., Goto, K., Shinozaki, M., Miyazawa, K., Miyazono, K., and Imamura, T. (2005).
NEDD4-2 (neural precursor cell expressed, developmentally down-regulated 4-2) negatively regulates
TGF-beta (transforming growth factor-beta) signalling by inducing ubiquitin-mediated degradation of
Smad2 and TGF-beta type I receptor. Biochem J *386*, 461-470.
Kushwaha, D., O'Leary, C., Cron, K.R., Deraska, P., Zhu, K., D'Andrea, A.D., and Kozono, D. (2015). USP9X
inhibition promotes radiation-induced apoptosis in non-small cell lung cancer cells expressing mid-to-
high MCL1. Cancer Biol Ther *16*, 392-401.
Lee, Y.S., Park, J.S., Kim, J.H., Jung, S.M., Lee, J.Y., Kim, S.J., and Park, S.H. (2011). Smad6-specific
recruitment of Smurf E3 ligases mediates TGF-β1-induced degradation of MyD88 in TLR4 signalling. Nat
Commun *2*, 460.
Li, X., Song, N., Liu, L., Liu, X., Ding, X., Song, X., Yang, S., Shan, L., Zhou, X., Su, D.*, et al.* (2017). USP9X
regulates centrosome duplication and promotes breast carcinogenesis. Nat Commun *8*, 14866.
Lin, A., Hou, Q., Jarzylo, L., Amato, S., Gilbert, J., Shang, F., and Man, H.Y. (2011). Nedd4-mediated AMPA
receptor ubiquitination regulates receptor turnover and trafficking. J Neurochem *119*, 27-39.
Liu, C.C., Kanekiyo, T., Roth, B., and Bu, G. (2014). Tyrosine-based signal mediates LRP6 receptor
endocytosis and desensitization of Wnt/β-catenin pathway signaling. J Biol Chem *289*, 27562-27570.
Lu, Y., Adegoke, O.A., Nepveu, A., Nakayama, K.I., Bedard, N., Cheng, D., Peng, J., and Wing, S.S. (2009).
USP19 deubiquitinating enzyme supports cell proliferation by stabilizing KPC1, a ubiquitin ligase for
p27Kip1. Mol Cell Biol *29*, 547-558.
Luga, V., Zhang, L., Viloria-Petit, A.M., Ogunjimi, A.A., Inanlou, M.R., Chiu, E., Buchanan, M., Hosein, A.N.,
Basik, M., and Wrana, J.L. (2012). Exosomes mediate stromal mobilization of autocrine Wnt-PCP
signaling in breast cancer cell migration. Cell *151*, 1542-1556.
Lui, T.T., Lacroix, C., Ahmed, S.M., Goldenberg, S.J., Leach, C.A., Daulat, A.M., and Angers, S. (2011). The
ubiquitin-specific protease USP34 regulates axin stability and Wnt/β-catenin signaling. Mol Cell Biol *31*,
2053-2065.
MacDonald, B.T., Tamai, K., and He, X. (2009). Wnt/beta-catenin signaling: components, mechanisms,
and diseases. Dev Cell *17*, 9-26.
MacGurn, J.A., Hsu, P.C., and Emr, S.D. (2012). Ubiquitin and membrane protein turnover: from cradle to
grave. Annu Rev Biochem *81*, 231-259.
Madrzak, J., Fiedler, M., Johnson, C.M., Ewan, R., Knebel, A., Bienz, M., and Chin, J.W. (2015).
Ubiquitination of the Dishevelled DIX domain blocks its head-to-tail polymerization. Nat Commun *6*,
6718.

Marikawa, Y., and Elinson, R.P. (1998). beta-TrCP is a negative regulator of Wnt/beta-catenin signaling
pathway and dorsal axis formation in Xenopus embryos. Mech Dev *77*, 75-80.
Miyazaki, K., Fujita, T., Ozaki, T., Kato, C., Kurose, Y., Sakamoto, M., Kato, S., Goto, T., Itoyama, Y., Aoki,
M*., et al.* (2004). NEDL1, a novel ubiquitin-protein isopeptide ligase for dishevelled-1, targets mutant
superoxide dismutase-1. J Biol Chem *279*, 11327-11335.
Mouchantaf, R., Azakir, B.A., McPherson, P.S., Millard, S.M., Wood, S.A., and Angers, A. (2006). The
ubiquitin ligase itch is auto-ubiquitylated in vivo and in vitro but is protected from degradation by
interacting with the deubiquitylating enzyme FAM/USP9X. J Biol Chem *281*, 38738-38747.
Mukai, A., Yamamoto-Hino, M., Awano, W., Watanabe, W., Komada, M., and Goto, S. (2010). Balanced
ubiquitylation and deubiquitylation of Frizzled regulate cellular responsiveness to Wg/Wnt. EMBO J *29*,
2114-2125.
Mund, T., Graeb, M., Mieszczanek, J., Gammons, M., Pelham, H.R., and Bienz, M. (2015). Disinhibition of
the HECT E3 ubiquitin ligase WWP2 by polymerized Dishevelled. Open Biol *5*, 150185.
Narimatsu, M., Bose, R., Pye, M., Zhang, L., Miller, B., Ching, P., Sakuma, R., Luga, V., Roncari, L.,
Attisano, L*., et al.* (2009). Regulation of planar cell polarity by Smurf ubiquitin ligases. Cell *137*, 295-307.
Nethe, M., de Kreuk, B.J., Tauriello, D.V., Anthony, E.C., Snoek, B., Stumpel, T., Salinas, P.C., Maurice,
M.M., Geerts, D., Deelder, A.M*., et al.* (2012). Rac1 acts in conjunction with Nedd4 and dishevelled-1 to
promote maturation of cell-cell contacts. J Cell Sci *125*, 3430-3442.
Peterson, L.F., Sun, H., Liu, Y., Potu, H., Kandarpa, M., Ermann, M., Courtney, S.M., Young, M.,
Showalter, H.D., Sun, D*., et al.* (2015). Targeting deubiquitinase activity with a novel small-molecule
inhibitor as therapy for B-cell malignancies. Blood *125*, 3588-3597.
Potu, H., Peterson, L.F., Kandarpa, M., Pal, A., Sun, H., Durham, A., Harms, P.W., Hollenhorst, P.C.,
Eskiocak, U., Talpaz, M*., et al.* (2017). Usp9x regulates Ets-1 ubiquitination and stability to control NRAS
expression and tumorigenicity in melanoma. Nat Commun *8*, 14449.
Premarathne, S., Murtaza, M., Matigian, N., Jolly, L.A., and Wood, S.A. (2017). Loss of Usp9x disrupts cell
adhesion, and components of the Wnt and Notch signaling pathways in neural progenitors. Sci Rep *7*,
8109.
Pérez-Mancera, P.A., Rust, A.G., van der Weyden, L., Kristiansen, G., Li, A., Sarver, A.L., Silverstein, K.A.,
Grützmann, R., Aust, D., Rümmele, P*., et al.* (2012). The deubiquitinase USP9X suppresses pancreatic
ductal adenocarcinoma. Nature *486*, 266-270.
Rotin, D., and Kumar, S. (2009). Physiological functions of the HECT family of ubiquitin ligases. Nat Rev
Mol Cell Biol *10*, 398-409.
Saito-Diaz, K., Chen, T.W., Wang, X., Thorne, C.A., Wallace, H.A., Page-McCaw, A., and Lee, E. (2013). The
way Wnt works: components and mechanism. Growth Factors *31*, 1-31.
Savio, M.G., Wollscheid, N., Cavallaro, E., Algisi, V., Di Fiore, P.P., Sigismund, S., Maspero, E., and Polo, S.
(2016). USP9X Controls EGFR Fate by Deubiquitinating the Endocytic Adaptor Eps15. Curr Biol *26*, 173-
183.
Schwarz-Romond, T., Fiedler, M., Shibata, N., Butler, P.J., Kikuchi, A., Higuchi, Y., and Bienz, M. (2007).
The DIX domain of Dishevelled confers Wnt signaling by dynamic polymerization. Nat Struct Mol Biol *14*,
484-492.
Schwarz-Romond, T., Merrifield, C., Nichols, B.J., and Bienz, M. (2005). The Wnt signalling effector
Dishevelled forms dynamic protein assemblies rather than stable associations with cytoplasmic vesicles.
J Cell Sci *118*, 5269-5277.
Schwickart, M., Huang, X., Lill, J.R., Liu, J., Ferrando, R., French, D.M., Maecker, H., O'Rourke, K., Bazan,
F., Eastham-Anderson, J*., et al.* (2010). Deubiquitinase USP9X stabilizes MCL1 and promotes tumour cell
survival. Nature *463*, 103-107.

Tanksley, J.P., Chen, X., and Coffey, R.J. (2013). NEDD4L is downregulated in colorectal cancer and
inhibits canonical WNT signaling. PLoS One *8*, e81514.
Tauriello, D.V., Haegebarth, A., Kuper, I., Edelmann, M.J., Henraat, M., Canninga-van Dijk, M.R., Kessler,
B.M., Clevers, H., and Maurice, M.M. (2010). Loss of the tumor suppressor CYLD enhances Wnt/beta-
catenin signaling through K63-linked ubiquitination of Dvl. Mol Cell *37*, 607-619.
Wang, Y. (2009). Wnt/Planar cell polarity signaling: a new paradigm for cancer therapy. Mol Cancer Ther
8, 2103-2109.
Wei, W., Li, M., Wang, J., Nie, F., and Li, L. (2012). The E3 ubiquitin ligase ITCH negatively regulates
canonical Wnt signaling by targeting dishevelled protein. Mol Cell Biol *32*, 3903-3912.
Wertz, I.E., O'Rourke, K.M., Zhou, H., Eby, M., Aravind, L., Seshagiri, S., Wu, P., Wiesmann, C., Baker, R.,
Boone, D.L.*, et al.* (2004). De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF-
kappaB signalling. Nature *430*, 694-699.
Wu, X., Fukushima, H., North, B.J., Nagaoka, Y., Nagashima, K., Deng, F., Okabe, K., Inuzuka, H., and Wei,
W. (2014). SCFβ-TRCP regulates osteoclastogenesis via promoting CYLD ubiquitination. Oncotarget *5*,
4211-4221.
Xie, Y., Avello, M., Schirle, M., McWhinnie, E., Feng, Y., Bric-Furlong, E., Wilson, C., Nathans, R., Zhang, J.,
Kirschner, M.W.*, et al.* (2013). Deubiquitinase FAM/USP9X interacts with the E3 ubiquitin ligase SMURF1
protein and protects it from ligase activity-dependent self-degradation. J Biol Chem *288*, 2976-2985.
Yang, B., Zhang, S., Wang, Z., Yang, C., Ouyang, W., Zhou, F., Zhou, Y., and Xie, C. (2016). Deubiquitinase
USP9X deubiquitinates β-catenin and promotes high grade glioma cell growth. Oncotarget *7*, 79515-
79525.
Yang, Y., and Mlodzik, M. (2015). Wnt-Frizzled/planar cell polarity signaling: cellular orientation by facing
the wind (Wnt). Annu Rev Cell Dev Biol *31*, 623-646.
Yu, A., Rual, J.F., Tamai, K., Harada, Y., Vidal, M., He, X., and Kirchhausen, T. (2007). Association of
Dishevelled with the clathrin AP-2 adaptor is required for Frizzled endocytosis and planar cell polarity
signaling. Dev Cell *12*, 129-141.
Yu, A., Xing, Y., Harrison, S.C., and Kirchhausen, T. (2010). Structural analysis of the interaction between
Dishevelled2 and clathrin AP-2 adaptor, a critical step in noncanonical Wnt signaling. Structure *18*, 1311-
1320.
Yu, X., Wang, M., Dong, Q., and Jin, F. (2014). Diversin is overexpressed in breast cancer and accelerates
cell proliferation and invasion. PLoS One *9*, e98591.
Zhang, Z.R., Bonifacino, J.S., and Hegde, R.S. (2013). Deubiquitinases sharpen substrate discrimination
during membrane protein degradation from the ER. Cell *154*, 609-622.
Zhu, Y., Tian, Y., Du, J., Hu, Z., Yang, L., Liu, J., and Gu, L. (2012). Dvl2-dependent activation of Daam1 and
RhoA regulates Wnt5a-induced breast cancer cell migration. PLoS One *7*, e37823.

Figure 1. WWP1 interacts with USP9X and DVL2 in MDA-MB-231 human breast cancer cells. (**A**) Co-immunoprecipitation analysis testing hits from SILAC-MS analysis in (FIG S1A). FLAG-WWP1 was affinity purified from MDA-MB-231 cell lysates. Input and immunoprecipitated (IP) fractions were blotted for the indicated species. (**B**) Schematic of known domains in WWP1, DVL2, and USP9X. (**C**) Analysis of co-purification of DVL2 and USP9X with WWP1 variants. The indicated WWP1 variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. The *wwp1-4ww* mutant indicates that all four WW domains have been mutated to disrupt the ability to bind PY motifs. (**D**) Quantification of USP9X co-purification with FLAG-WWP1 variants (normalized to WT) over multiple experiments (n=3). Red asterisk indicates a significant difference based on Student's t-test analysis. (**E**) Analysis of co-purification of DVL2 and USP9X with WWP1 variants. The indicated WWP1 variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. Each mutant variant of WWP1 contains a single in-tact WW domain (indicated), while the remaining three are mutated to disrupt PY motif binding. (**F**) Quantification of DVL2 co-purification with FLAG-WWP1 variants (normalized to WT) over multiple experiments (n=3). (**G**) Analysis of co-purification of WWP1 and USP9X with DVL2 variants. The indicated mutant DVL2 variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. Mutant variants of DVL2 included point mutations disrupting the ability of its PY motifs to interact with WW domains. (**H**) Analysis of co-purification of WWP1 and DVL2 with USP9X variants. The indicated USP9X variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. Mutant variants of USP9X included a small C-terminal truncation which results in deletion of both C-terminal PY motifs. (**I**) A schematic model for how USP9x and DVL2 both interact with WWP1 via PY motif interactions. The data also suggest an interaction between USP9X and DVL2 that is independent of WWP1 (dotted red line).

A

B

C

D

Nielsen et al.

Figure 2 WWP1 and USP9X operate on DVL2 to establish a ubiquitylation rheostat. **(A)** *In vitro* ubiquitylation reactions were performed using recombinant purified FLAG-WWP1 (60nM) and HA-DVL2 purified from cultured cells. E3 conjugation reactions were allowed to proceed for 60 minutes before the conjugation reaction was terminated. **(B)** Deconjugation assays were performed using HA-DVL2 ubiquitylated *in vitro* by WWP1 (as generated in **(A)**). Equivalent molar ratios of USP9X, OTUB1 and AMSH (100nM) were added and each reaction was allowed to proceed for 60 minutes before the deconjugation reaction was terminated and HA-DVL2 was resolved by SDS-PAGE and immunoblot (bottom panel). For each sample, a ratio of modified:unmodified HA-DVL2 was measured using ImageJ. The modified:unmodified HA-DVL2 ratio was averaged over multiple replicate experiments (n=3). The red asterisk indicates a statistically significant difference (p<0.05) compared to DVL2 that has been modified by WWP1 and subsequently treated with OTUB1 or AMSH. **(C)** Representative immunoblot of *in vitro* ubiquitylation/deubiquitylation reactions (replicate experiments shown in **FIG S2D-E**). Experiments were performed using indicated combinations of purified FLAG-WWP1 and 6xHIS-USP9X with HA-DVL2 purified from cultured cells (see **FIG S2A-B**). Reactions were allowed to proceed for 60 minutes before being terminated and HA-DVL2 was resolved by SDS-PAGE and immunoblot. **(D)** Line density plots from the HA-DVL2 immunoblot data in **(C)** illustrate differences in extent of ubiquitylation in each sample. Replicate experiments are shown in **FIG S2D-E**.

Nielsen et al.

Figure 3. USP9X promotes canonical WNT activation. **(A)** Analysis of ligand-stimulated WNT activation in MDA-MB-231 cells (blue bars) and *usp9x* knockout equivalents (gray bars). TopFLASH luciferase assays were used to measure WNT activation (top panel) and immunoblotting was performed to assess stabilization of nuclear β-catenin. * indicates WNT activation in *usp9x* knockout cells is significantly decreased (p<0.05) compared to sibling MDA-MB-231 cells. **(B)** MDA-MB-231 cells (wildtype and *usp9x* knockout) were stimulated with WNT3a ligand and R-spondin and WNT activation was measured by TopFLASH luciferase assays. Prior to activation, cells were transiently transfected with empty vector (EV) or vector expression USP9X (wildtype (WT) or catalytic dead (CD; C1566V)). **(C)** Ligand-stimulated WNT activation was measured in the presence of indicated concentrations of WP1130. Based on the data an IC50 was estimated. **(D)** MDA-MB-231 cells were transfected with empty vector or plasmids for CMV-driven expression of wildtype (WT) or catalytic dead USP9X. **(E)** Ligand-stimulated WNT activation was measured using TopFLASH luciferase assays for MDA-MB-231 cells and *usp9x* knockout equivalents. Cells were transfected with control siRNA or siRNA targeting knockdown of NEDD4L, WWP1 or WWP2.

Figure 4. USP9X promotes canonical WNT activation via deubiquitylation of DVL2. **(A)** Immunoblotting analysis of FLAG-DVL2 mobility by SDS-PAGE from lysates of MDA-MB-231 cells or *usp9x* knockout equivalents. A low exposure α-FLAG blot is shown to illustrate differences in DVL2 abundance, while the higher exposure α-FLAG is shown to illustrate the higher MW species evident in *usp9X* knockout cells. **(B)** SILAC-MS profiling of post-translational modification sites (phosphorylation and ubiquitylation) on DVL2. In this experiment, FLAG-DVL2 purified from MDA-MB-231 cells (heavy) was compared to *usp9x* knockout equivalents (light). A schematic illustrating the domain structure of DVL2 is shown at the top with asterisks to indicate the position of detected ubiquitylation events. **(C)** Analysis of ligand-stimulated WNT activation was measured using TopFLASH reporter assays in STF-293 cells transfected with the indicated expression vectors. UL36 is the DUB domain fused to DVL2 in these experiments. **(D)** Following siRNA knockdown of USP9X, ligand-stimulated WNT activation was measured using TopFLASH reporter assays in STF-293 cells transfected with the indicated HA-DVL2 expression vectors. DVL2-K0 is a variant where all encoded Lys residues are substituted with Arg residues. * represents a statistically significant difference when compared to indicated sample (p<0.05). n.s. represents no significant difference relative to indicated comparison.

Nielsen et al.

Figure 5. DVL2 ubiquitylation state specifies association with canonical WNT or WNT-PCP pathway factors. **(A)** SILAC-MS was used to resolve the DVL2 interaction network in MDA-MB-231 human breast cancer cells. Near-stoichiometric interactions (solid lines) were detected with the AP-2 clathrin adaptor complex and VANGL1-cofilin, which comprise a subcomplex of the WNT-PCP pathway. Sub-stoichiometric interactions (dotted lines) were identified for NEDD4 family E3 ubiquitin ligases and proteasome subunits. **(B)** Interactions detected by SILAC-MS were confirmed by coimmunoprecipitation. **(C and D)** SILAC-MS analysis was performed to measure how DVL2 interactions are impacted by catalytic active vs. catalytic dead DUB fusion **(C)** or by loss of *USP9X* **(D)**. Scatter plots depict individual H:L measurements (LOG$_2$ transformed) for each peptide resolved for the indicated proteins. The mean value and standard deviation for each protein is indicated. For the experiment shown in **(C)**, several peptides with extremely low H:L ratios are not depicted here but are shown in **FIG S5B**. These outlier peptides correspond to sites of ubiquitin modification (**FIG S5C**) shown in **FIG 4B**. Differences between DVL2 (bait) and all interacting proteins (AP2 complex, VANGL1, CFL1, and proteasome subunits) are statistically significant (p<0.005).

Nielsen et al.

Figure 6. USP9X regulates DVL2 localization and antagonizes WNT-PCP activation. **(A)** Immunofluorescence imaging of FLAG-DVL2 (green) and actin (phalloidin, blue) in MDA-MB-231 human breast cancer cells transfected with control siRNA (top panels), siRNA targeting USP9X (middle panels), or a combination of siRNA targeting USP9X and siRNA targeting WWP1 (bottom panels). **(B)** Quantification of DVL2 cellular distribution over a population of MDA-MB-231 cells (n > 100). **(C)** WT MDA-MB-231 and *usp9x* KO variant cell lysates were assayed for active Rho by pulldown assay. Addition of GDP or non-hydrolyzable GTP (GTPγS) to cell lysates served as negative and positive controls, respectively. **(D)** Quantification of Rho activation shown in **FIG 6C** (n=3) was performed by measuring the amount of active Rho detected in parent and *usp9x* knockout cells and normalizing to the GTPγS (maximal activation) control. The red asterisk indicates a significant difference when compared to WT cells (p<0.05) **(E)** MDA-MB-231 cells were plated at low density and cell migration (µm/hour) speed was measured. Red asterisk indicates a significant difference between the tested populations (n>30). **(F)** Rose plots showing trajectories of individual MDA-MB-231 cells (left) and *usp9x* knockout equivalents (right).

Nielsen et al.

Figure 7. USP9X-mediated regulation of cell motility requires DVL2 ubiquitylation. **(A)** Migration speed of MDA-MB-231 *usp9x* KO cells treated with either control siRNA or siRNA targeting WWP1 was measured after plating at low density for the indicated conditions. Red asterisk indicates a significant difference compared to control siRNA treated cells (p<0.005). **(B)** MDA-MB-231 cell lines stably expressing DVL2-DUB fusions (catalytic active and catalytic dead) were treated with control siRNA or siRNA targeting USP9X and cell speed was measured for 25 cells. **(C)** MDA-MB-231 cell lines stably expressing DVL2-DUB fusions (catalytic active and catalytic dead) underwent mock or WNT5a treatment and cell speed was measured for 25 cells. The red asterisk indicates a statistically significant difference (p<0.005) compared to MDA-MB-231 cells stably expressing the catalytic dead DUB-DVL2 fusion treated with WNT5a. **(D)** Model for DVL2 regulation by the USP9X-WWP1 axis to establish a ubiquitin rheostat on DVL2 that influences interaction preference with either the clathrin adaptor AP-2 complex, which is important for canonical WNT signaling, or the VANGL1-cofilin complex, which is important for the WNT-PCP pathway. Thus, the USP9X-WWP1 axis regulates DVL2 ubiquitylation to determine WNT pathway specification.

Nielsen et al.

Supplemental Figure S1. Related to Figure 1. USP9X, WWP1, and DVL2 interactions are scaffolded by WW-PY interactions **(A)** Schematic summarizing putative WWP1 interacting proteins identified from SILAC-MS. Complete data set for this experiment is reported in Table S1. **(B)** Endogenous WWP1 was immunoprecipitated from MDA-MB-231 cells. Input (10%) and immunoprecipitates were resolved by SDS-PAGE and immunoblotted for the indicated species. **(C)** Analysis of co-purification of DVL2 and USP9X with WWP1 variants. The indicated WWP1 variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. Mutant variants of WWP1 included point mutations disrupting the ability of individual WW domains to bind PY motifs. **(D)** Quantification of DVL2 co-purification with FLAG-WWP1 variants (normalized to WT) over multiple experiments (n=3). **(E)** Analysis of co-purification of WWP1 and DVL2 with USP9X mutant variants. The indicated mutant USP9X variants were FLAG affinity purified, and quantitative immunoblots were performed to assess co-purification of interacting factors. *py3* and *py4* mutations indicate point mutations in PY3 and PY4, respectively, while *2py* indicates both C-terminal PY motifs were mutated. *C1566V* is mutated at the catalytic cysteine residue of USP9X.

Supplemental Figure S2. Related to Figure 2. Analysis of WWP1 and USP9x activity toward DVL2 *in vitro*.
(A) Purity of recombinant FLAG-WWP1 and 6XHIS-USP9X used in *in vitro* conjugation/deconjugation reactions (FIG 2) shown by SYPRO Ruby gel staining. **(B)** Purity of HA-DVL2 (purified from cultured cells) used in *in vitro* conjugation/deconjugation reactions (FIG 2) shown by SYPRO Ruby gel staining. **(C)** *In vitro* ubiquitylation reactions were performed using recombinant purified FLAG-WWP1 (60nM) and HA-DVL2 (wildtype and K0, a variant where all lysine residues are mutated to arginine) purified from cultured cells. E3 conjugation reactions were allowed to proceed for 60 minutes before the conjugation reaction was terminated. **(D-E, left panels)** Biological replicates of *in vitro* ubiquitylation rheostat reaction shown in FIG 2C-D, experiments were performed using indicated combinations of purified FLAG-WWP1 and 6XHIS-USP9X with HA-DVL2 purified from cultured cells. Reactions were allowed to proceed for 60 minutes before reaction was terminated and HA-DVL2 was resolved by SDS-PAGE and immunoblot. (**D-E, right panels**) Quantification of HA-DVL2 ubiquitylation under indicated conditions using ImageJ line profile analysis of immunoblot.

Nielsen et al.

Supplemental Figure S3. Related to Figure 3. USP9X and WWP1 regulate canonical WNT activation. **(A)** Analysis of ligand-stimulated WNT activation in the presence of control siRNA or siRNAs targeting USP9X knockdown. TopFLASH luciferase assays were used to measure WNT activation (top panel) and immunoblotting was performed to assess knockdown efficiency (bottom panel). The asterisk indicates significant difference compared to control siRNA treatment ($p < 0.05$). **(B)** TopFLASH activation of STF-293 cells transfected with empty vector or plasmids for CMV-driven expression of wildtype (WT) FLAG-WWP1, *wwp1-C890S* (a catalytic dead variant) or *wwp1*-WW2 (a variant with only WW2 intact) (see **FIG 1E** and **1F**). The asterisk indicates a statistically significant difference compared to the empty vector control ($p < 0.05$) while n.s. indicates no significant difference compared to the empty vector control. **(C)** TopFLASH activation of STF-293 cells transfected with empty vector or plasmids for CMV-driven expression of wildtype (WT) FLAG-WWP1 or a *ww2* mutant variant (see **FIG S1C** and **S1D**). Single asterisk indicates statistically significant difference compared to empty vector control ($p < 0.05$), while double asterisk indicates statistically significant difference compared to both empty vector control and wildtype WWP1 overexpression ($p < 0.005$).

Nielsen et al.

Nielsen et al.

Supplemental Figure S4. Related to Figure 4. USP9X regulates DVL2 ubiquitylation state. **(A)** Analysis of LiCl stimulation of WNT activation following transfection with control siRNA or siRNAs targeting USP9X knockdown. Immunoblotting was performed to assess knockdown efficiency and stabilization of nuclear β–catenin. **(B)** Immunoblotting analysis of DVL2 mobility by SDS-PAGE following transfection with control siRNA or siRNAs targeting USP9X knockdown. **(C-E)** Raw data for SILAC quantification (corresponding to **FIG 4B**) of individual ubiquitylation events on DVL2 including diGly modification of K343 **(C)**, K428 **(D)** and K484 **(E)**. MS2 spectra for each diGly peptide is shown in the left panel. To the right of each MS2 spectra, chromatograms are shown for heavy and light unmodified (middle) and diGly-modified (right) peptides, illustrating the quantification of each peptide. In each case, the diGly-modified form of the peptide is significantly elevated in the light sample, which is derived from the *usp9x* mutant sample (see **FIG 4B**). **(F)** Functional analysis of DLV2 lysine mutants. STF-293 reporter cells were transiently transfected with the indicated FLAG-DVL2 plasmids and siRNAs and analyzed for WNT activation using the TopFLASH luciferase assay.

B

SILAC Quantification
H: FLAG-DUBcat.dead-Dvl2
L: FLAG-DUBactive-Dvl2

C

Outlier Peptides

Peptide	H:L ratio (normalized)
$_{416}$GLSVHTDMASVTK$_{428}$	0.014791
$_{416}$GLSVHTDMASVTK$_{428}$	0.017268
$_{429}$AMAAPESGLEVR$_{440}$	0.015015
$_{477}$KYASGLLK$_{484}$	0.023585
$_{616}$SGSGSESEPSSR$_{627}$	0.04083

Supplemental Figure S5. Related to Figure 5. DVL2 ubiquitylation regulates interactions. **(A)** Immunofluorescence microscopy analysis was performed on MDA-MB-231 breast cancer cells expressing FLAG-DVL2. **(B)** Expanded graph of SILAC-MS analysis in **FIG 5C** showing outlier DVL2 peptides. **(C)** Description of outlier DVL2 peptides shown in **(B)** with corresponding normalized H:L ratio.

Supplemental Figure S6. Related to Figure 6. USP9X antagonizes cell motility. **(A)** The indicated MDA-MB-231 cell lines were transiently transfected with a GFP-USP9X expression vector and cell migration speed was measured for untransfected and transfected (GFP-positive) cells (n=25). The red asterisk represents a statistically significant difference compared to untransfected *usp9x* KO cells. **(B)** Migration speed of MDA-MB-231 cells was measured after treatment with DMSO control or WP1130 (300nM). There is a statistically significant difference when comparing cell speed after WP1130 treatment to DMSO-treated control ($p<0.005$). **(C)** Migration speed of MDA-MB-231 cells was measured after treating with control siRNA or siRNA targeting VANGL1. There was a statistically significant difference ($p<0.05$) in cell speed with VANGL1 knockdown when compared to control siRNA treatment in the indicated cell lines. Bottom panel: Corresponding immunoblots confirming VANGL1 knockdown with siRNA treatment. **(D)** Migration speed of indicated MDA-MB-231 cells after treatment with DMSO control or LGK974 (1nM). There was no significant difference with LGK974 treatment when compared to DMSO control in indicated cell types.

Nielsen et al.

Supplemental Figure 7. Related to Figure 7. DVL2 expression suppresses the motility phenotype of *usp9x* knockout cells. **(A)** The indicated MDA-MB-231 cell lines were transiently transfected with a GFP-WWP1 expression vector and cell migration speed was measured for untransfected and transfected (GFP-positive) cells (n=25). There was no significant difference relative to untransfected cells. **(B-C)** The indicated MDA-MB-231 cell lines were transiently transfected with a GFP-DVL2 expression vector and cell migration speed was measured for untransfected and transfected (GFP-positive) cells (n=25). Red asterisk indicates a statistically significant difference (p<0.005) when compared to untransfected control in the indicated cell line (n=25). **(D)** The indicated MDA-MB-231 cell lines were transiently transfected with a GFP-DVL2-K0 (a construct in which all lysines have been mutated to arginines) expression vector and cell migration speed was measured for untransfected and transfected (GFP-positive) cells (n=25). Red asterisk indicates a significant difference (p<0.005) when compared to untransfected control in the indicated cell line (n=25).

Table S1. WWP1 Interaction Profile using SILAC-MS

category	protein	known function / description	pept. count	H:L ratio	stoich. (norm.)
Bait	WWP1	NEDD4 family E3 ubiquitin ligase (bait)	289	36.01	1.00
membrane traffic	SCAMP3	endosomal recycling	18	37.43	0.16
	ARF4	polarized exocytosis	8	7.13	0.14
	RAB14	regulation of Golgi-to-endosome transport	5	9.22	0.07
	ARF5	endocytosis of integrins	3	12.40	0.05
	SEC23IP	regulates ER exit	9	5.47	0.03
plasma membrane / membrane associated	CCDC85C	function unknown	21	17.47	0.16
	SPG20	function unknown	33	47.71	0.16
	AMOTL2	regulation of cell junctions; Hippo signaling	22	38.39	0.09
	AMOTL1	regulation of cell junctions; Hippo signaling	21	33.19	0.07
	RAPGEF6	function unknown	31	28.00	0.06
	MC1R	melanocortin receptor	3	26.71	0.03
	AXL	receptor tyrosine kinase	8	5.18	0.03
	ERBB2IP	binds and stabilizes ERBB2 receptor	12	9.36	0.03
	FNDC3B	function unknown	10	5.19	0.03
	SLC2A1	facilitative glucose transporter	3	6.67	0.02
	LRP10	putative receptor; function unknown	4	12.62	0.02
	EGFR	epidermal growth factor receptor	6	8.66	0.02
	ADAM9	cell surface metalloproteinase	4	6.04	0.02
	NPC1	intracellular cholesterol transporter	4	8.53	0.01
signaling	WBP2	Hippo pathway and hormone signaling	26	17.07	0.32
	PDLIM7	bone formation and BMP signaling pathways	30	22.18	0.21
	PTPN14	protein tyrosine phosphatase	75	22.11	0.20
	TMEM55B	phosphatidylinositol 4,5-bisphosphate 4-phosphatase	4	6.14	0.05
	DVL2	signal transducer, WNT pathway	9	25.67	0.04
	PTPN21	protein tyrosine phosphatase	4	10.24	0.01
ubiquitin proteasome system	USP9X	deubiquitylase	94	25.37	0.12
	FAM175B	member of the BRISC complex; deubiquitylase	9	13.22	0.07
	TXNIP	thioredoxin interacting protein; may function as E3 adaptor	8	8.86	0.06
	BRCC3	member of the BRISC complex; deubiquitylase	5	21.72	0.05
	ARRDC1	thioredoxin interacting protein; may function as E3 adaptor	6	20.14	0.04
	C19ORF62	member of the BRISC complex; deubiquitylase	3	9.28	0.03
	USP24	deubiquitylase	20	23.66	0.02

Light: MDA-MB-231 cells stably expressing pQCXIP (empty vector)
Heavy: MDA-MB-231 cells stably expressing FLAG-WWP1
Selection Criteria: (1) number of peptides quantified ≥ 3; (2) H:L ratio > 5

Table S2. DVL2 Interaction Profile using SILAC-MS

category	protein	known function / description	pept. count	H:L ratio	stoich. (norm.)
Bait	DVL2	WNT pathway signaling protein	29	28.70	1.00
AAA ATPase	ATAD3A	mitochondria-localized AAA ATPase of unknown function	18	8.58	0.72
	ATAD3B		14	**	0.55
endocytosis	AP2M1	AP2 clathrin adpator complex	10	19.79	0.58
	AP2B1		19	8.83	0.51
	AP2A1		18	37.89	0.47
	AP2S1		2	10.99	0.36
	AP2A2		8	**	0.22
	EPS15L1	endocytic adaptor proteins; sort ubiquitylated cargo for endocytosis	3	**	0.09
	EPS15		2	**	0.06
signaling	VANGL1	WNT-PCP signaling protein	5	**	0.24
	DVL3	WNT pathway signaling protein	5	**	0.18
	DVL1		4	**	0.15
	CK1delta	kinase; regulates canonical WNT activation	2	**	0.12
	WDR26	signal transduction protein	2	13.00	0.08
	RANBP9	signal transduction protein	2	**	0.07
ubiquitin proteasome system	NEDD4L	NEDD4 family E3 ubiquitin ligase	4	22.38	0.10
	PSD11	proteasome subunits	2	5.58	0.12
	PSA7		2	36.59	0.20
cytoskeleton	Cofilin-1	regulator of actin dynamics	3	5.70	0.46

Light: MDA-MB-231 cells stably expressing pQCXIP (empty vector)

Heavy: MDA-MB-231 cells stably expressing FLAG-DVL2

Selection Criteria: (1) number of peptides quantified ≥ 2; (2) H:L ratio > 5

** indicates signal in heavy channel but no signal in light channel

Table S3. DVL2 Interaction Profile in WNT-activated cells using SILAC-MS

category	Protein	known function / description	pept. count	H:L ratio	stoich. (norm.)
bait	DVL2	WNT pathway signaling protein	43	107.20	1.00
AAA ATPase	ATAD3B	mitochondria-localized AAA ATPase of	15	**	0.40
	ATAD3A	unknown function	25	62.37	0.67
endocytosis and trafficking	AP2B1	AP2 clathrin adaptor complex	49	52.27	0.90
	AP2M1		17	43.41	0.67
	AP2A1		38	60.82	0.67
	AP2S1		5	71.38	0.60
	AP2A2		30	27.94	0.55
	CAVIN1	caveolae formation and organization	10	18.36	0.44
	CAVIN2		8	**	0.32
	SQSTM1	autophagy receptor	10	10.15	0.39
	EPS15L1	endocytic adaptor proteins; sort ubiquitylated cargo for endocytosis	7	6.31	0.14
	EPS15		2	**	0.04
signaling	VANGL1	WNT-PCP signaling protein	13	**	0.42
	CK1 delta	casein kinase isoforms; regulate canonical WNT activation	17	**	0.70
	CKII beta		7	**	0.56
	CK1 epsilon		12	**	0.49
	CKII alpha		9	**	0.39
	PLK1	polo-like kinase-1	16	**	0.45
	DVL3	WNT pathway signaling protein	15	**	0.36
	RANBP9	signal transduction protein	15	**	0.35
	WDR26	signal transduction protein	13	74.44	0.34
ubiquitin proteasome system	NEDD4L	NEDD4 family E3 ubiquitin ligase	29	35.38	0.51
	WWP2		12	**	0.24
	WWP1		3	**	0.06
	PSA7	proteasome subunits	2	**	0.14
	PRS10		3	**	0.13
	PSA1		2	**	0.13
	PSD11		2	**	0.08
transcription	FOXK2	transcriptional regulators	18	**	0.47
	FOXK1		13	**	0.30
protein folding	BAG3	co-factor for HSP70	14	**	0.42

Light: MDA-MB-231 cells stably expressing pQCXIP (empty vector) stimulated with 200ng/mL mouse-Wnt3a and 100ng/mL mouse-Rspondin

Heavy: MDA-MB-231 cells stably expressing FLAG-DVL2 stimulated with 200ng/mL mouse-Wnt3a and 100ng/mL mouse-Rspondin

Selection Criteria: (1) number of peptides quantified ≥ 2; (2) H:L ratio > 5
** indicates signal in heavy channel but no signal in light channel