Changes to the microbiome of the stomach and intestine of Threespine Sticklebacks

associated with infection by the cestode *Schistocephalus solidus*.

Megan A Hahn[1], Nolwenn M Dheilly[1*]

[1]School of Marine and Atmospheric Sciences, Stony Brook University, 100 Nicolls Road, Stony

Brook NY, 11774

*Corresponding author

E-mail: nolwenn.dheilly@stonybrook.edu

Abstract

Helminth parasites are extremely diverse, prevalent, and infective of virtually all phyla. However, until recently, parasitic helminth infections have been studied as interactions between the host and the parasite, leaving out key players; microbes. Microbes play key roles in the biology of all living organisms, and this concept is challenging our understanding of host and parasite interactions. Because the microbiome controls many aspects of the physiology and behavior of their host, the discovery that many helminths can disrupt their host microbiome, either deliberately or as a side effect of infection, is of great significance. However, this phenomenon has been exclusively studied in helminth parasites that inhabit the gut, and hence share the same environment as their host microbiome. In this study we investigate the composition of the microbiome of threespine sticklebacks infected by the cestode *Schistocephalus solidus* who grows in the sterile body cavity of its intermediate host. Using a combination of qPCR and 16S sequencing approaches we show that *S. solidus* infection is associated with a significant increase in bacterial load and changes in diversity and composition of the microbiome of bpoth the stomach and intestine of sticklebacks.. We hypothesize that *S. solidus* interacts with host microbes either directly during the first 24 hours of infection, or indirectly, through modulation of the host immune system. The role of the stickleback microbiome in the outcome of *S. solidus* infection remains to be investigated.

Author summary

The classical paradigm assumes that symptoms associated with a parasite infection are due to the parasite itself. However, parasites can alter their host microbiome, meaning that microbes could be the ones responsible for associated diseases, and be the target for new therapeutic strategies. Here, we investigate the association between infection by *S. solidus* and the diversity and

composition of the host gut microbiome. Our study reveals for the first time that a cestode

parasite, that grows within the body cavity of its host, could be responsible for changes in the

microbiome composition of its host. This suggests that the interplay between parasite and host

associated microbes may be due to indirect mechanism, such as modulation of the host immune

response. Functional studies are now necessary to determine the mechanisms behind parasite-

associated alterations of their host microbiomes.

Introduction

The growing recognition of the fundamental role of microbes in animals and plant biology is

changing our understanding of life. The holobiont is defined as the combination of host, and all its

associated microbes [1]. We are only starting to comprehend the potential of applying these

concepts to host-parasite interactions to decipher the role of host-microbe, parasite-microbe, and

host-parasite-microbe interactions in the ecology and evolution of parasites [2]. Microbes provide

crucial services to the host, including disease resistance. For instance, there is considerable

evidence of the defensive role of bacterial symbionts in insects [3-6]. We are now able to take

advantage of this property and resistant mosquitoes carrying the endosymbiotic bacterium

Wolbachia are being engineered and released in natural populations to reduce the spread of

diseases such as malaria, dengue, and Zika [7-12]. In vertebrates too, there is growing evidence

that the microbiome participates in resistance mechanisms against diseases. For example, the

disruption of the host microbiome at early life-stages, or just before infection can increase the

susceptibility to diseases [13-15] and transplantation of the microbiome of resistant individuals

into susceptible individuals can be sufficient to provide protection [16].

Given the various direct and indirect roles of microbes in host defense, chances are that parasites

have gained the ability to take advantage of microbes to increase their virulence. Indeed,

evolutionary theories predict that parasite counter-defense strategies are shaped by the host defense

mechanisms that they encounter at the time of infection, which explains the major role of immune

evasion, and of manipulation of immune pathways in parasite virulence mechanisms [17].

Alteration of the host microbiome has been associated with parasitic helminth infection in various

species including pigs, hamsters, cattle, goats, and sheep [18-24]. In rats, infection by the cestode

Hymenolepis diminuta results in a shift from *Bacilli* to *Clostridia* species [25]. The liver fluke

Opisthorchis viverrini, associated with hepatobiliary diseases and cholangiocarcinoma, modifies

its host intestinal microbiome and promotes *Helicobacter pylori* infection in the liver [26-28].

Similar effects were observed when studying natural populations. For instance, infection by soil

transmitted nematodes *Trichuris spp.* in human individuals in Malaysia is correlated with an

increased microbial diversity in the gut and a differential abundance of a subset of OTUs, with

enrichment in the family *Paraprevotellaceae* [24]. Kreisinger et al.[29] showed that alterations of

the gut microbiome composition are species specific; wild mice (*Apodemus flavicollis*) naturally

infected by three different helminths (the nematodes *Heligmosomoides polygyrus* and *Syphacia*

spp., and the cestode *Hymenolepis spp.*) had significantly different microbiomes. However, all

these studies focused on parasites' alteration of the microbiome of the tissue within which the

parasite resides. Here, we characterized the stomach and intestine microbiomes of Threespine

sticklebacks, *Gasterosteus aculeatus*, parasitized or not with the cestode parasite *Schistocephalus*

solidus. Within 24 hours after ingestion of an infected copepod, *S. solidus* passes through the

intestine wall and reaches the body cavity of the fish where it matures into an adult worm, infective

for birds [30]. Thus, *S. Solidus* directly interacts with the gut microbiome of the threespine

stickleback for a very brief period. In the body cavity *S. solidus* may, however, impact the host

microbiome indirectly by modulating the expression of immune response genes [31-33].

Results

Threespine sticklebacks were collected live from Cheney Lake in Anchorage Alaska and shipped

back to Stony Brook University. Tissue collection was conducted after a three week acclimation

period in order to allow *S. solidus* to transit into the body cavity and grow to a conspicuous size if

the fish had only recently been infected. Threespine sticklebacks were then starved for 12 hours,

euthanized in tricaine methane sulphonate solution (MS222) and dissected in sterile conditions.

Stomach (St) and Intestine (In) collected from six non-infected fish (HGa) and six fish parasitized

by *S. solidus* (PGa) were collected and processed for analysis of their microbiome composition.

Swab samples of the body cavity of two non-infected and two parasitized individuals were also

collected and used as negative controls.

Real-time qPCR of 16S rRNA gene

We used a quantitative PCR (qPCR) approach with universal primers targeting the bacteria 16S

rRNA gene as a proxy to investigate the relationship between bacteria abundance in the fish host

and *S. solidus* infection, (Fig 1, S1 Fig). Indeed, variations in 16S abundance among samples may

be due to changes in bacterial load or changes in bacterial community composition.

Microbial diversity and composition

The bacterial community composition and diversity was determined by sequencing of the bacterial

16S rRNA gene V4 hypervariable region. Sequencing yielded an average of 43,645 (±SE 16,027)

high quality paired-end reads for *G. aculeatus* tissue samples. Sequencing depth of control samples

was higher and yielded an average of 83,875 (±SE 10,080) high quality reads. We resolved

Amplicon Sequence Variants (ASVs) to avoid applying the arbitrary 97% threshold that defines

molecular OTUs but has limited biological meaning [34]. The most abundant 100 ASVs in *G.*

aculeatus represented a cumulative average of 85.1 % (±SE 1.1) of the paired-end reads in *G.*

aculeatus tissue samples whereas it represented 7.8 % (±SE 0.4) reads in control samples.

Conversely, the top 100 ASVs in control samples represented a cumulative average of 84.8% (±SE

1.4) of the paired-end reads in control samples, and only 4.4% (±SE 1.8) of the paired-end reads

in *G. aculeatus* tissue samples. ASVs abundant in control samples were extremely similar to

common bacterial contaminants of the MoBio kit used for bacterial DNA extraction and were

removed from the analysis [35]. ASVs that were removed as contaminants can be found in S1

Table.

Alpha and beta diversity values were calculated on an ASV table rarified to 10,000 reads.

Comparisons between non-infected and parasitized individuals yielded significant differences in

alpha diversity values for Faith's Phylogenetic Diversity ($p < 0.05$, Table 1, Fig 2). Significant

differences in Chao1 diversity were observed between stomach and intestine samples ($p < 0.05$,

Table 1, Fig 2).

Table 1. Results of two factor ANOVA on alpha diversity metrics of bacterial communities

for all sample groupings including intestine (In), stomach (St), non-infected sticklebacks (H

Ga), parasitized sticklebacks (P Ga) at the 10,000 rarefaction level.

Comparisons		Evenness		Faith PD		Shannon		Chao1	
Group 1	Group 2	F	p-value	F	p-value	F	p-value	F	p-value
H Ga	P Ga	1.040	0.319	8.562	0.008*	0.086	0.772	1.199	0.287
Ga St	Ga In	1.152	0.295	0.753	0.395	3.470	0.077	6.614	0.018*

| **Interaction** | 0.123 | 0.730 | 1.175 | 0.291 | 0.545 | 0.470 | 1.759 | 0.199 |

* indicates a significant p-value for the given comparison

Principal coordinate analysis on weighted and unweighted UniFrac distances showed significant

clustering of non-infected and parasitized tissue samples along PC1 (unpaired t-test p<0.05, Fig 2,

S3 Fig and S3 and S4 Tables) and of intestine and stomach samples (Weighted UniFrac; PC2,

paired t-test p<0.05, S5 and S6 Fig, S2 Table, Unweighted UniFrac; PC4 and PC17 paired t-test

p<0.05, S3 Table). Community dissimilarity metrics (S5 Table) further confirmed that non-

infected and parasitized individuals host significantly different microbiomes (PERMANOVA;

weighted UniFrac P<0.05; unweighted UniFrac P<0.005) with significant differences in dispersion

(PERMDISP; weighted UniFrac p<0.05, unweighted UniFrac p<0.05). These results indicate that

the microbiome of parasitized individuals is overall less variable than the microbiome of non-

infected individuals. Together with the reduced phylogenetic diversity, it indicates a significant

reduction in abundance of variants of bacteria, rather than overall changes in species composition,

No significant differences were observed between stomach and intestine samples using these

metrics.

Taxa distribution and differential abundance analysis

We used DESeq2 to identify bacteria phylotypes differentially abundant among samples. The

proportion of ASVs differentially abundant between non-infected and parasitized *G. aculeatus*,

was similar to the proportion of ASVs differentially abundant between intestines and stomachs of

G. aculeatus, or differentially abundant between non-infected and parasitized individuals in

stomach and intestine (Table 2, S6 and S7 tables).

Table 2. Number of differentially abundant ASVs/genus identified by DESeq2 (Adj p value

<0.05).

Comparison between		Total	More in A	More in B
A	**B**			
Healthy (H)	**Parasitized (P)**	44 / 1	22 / 1	22 / 0
Intestine (In)	**Stomach (St)**	43 /4	8 / 1	35 / 3
Infection Status:Tissue Interaction		48 / 3		

Stickleback intestine and stomach microbiome communities were largely dominated by few ASVs

of *Clostridiales*, *Rickettsiales, Burkholderiales* and *Desulfovibrionales* (Fig 3). The cumulative

median relative abundance of the five most abundant ASVs in the threespine stickleback gut

microbiome communities represented 56% of the reads, while the next five most abundant ASVs

represented 20% of the reads. On few occasions, individual fish had high abundance of ASVs that

only occurred in low abundance in other samples. For example, individual Ga12 (Fig 3) had high

abundance of *Gammaproteobacteria* of the family *Shewanella* with a cumulative relative

abundance in the stomach and intestine of 31.4% and 41.7%. Yet, the cumulative mean relative

abundance across all other fish is 9.2% and 8.7% for the stomach and intestine, respectively.

Despite the very subtle differences in overall diversity between stomach and intestine samples,

significant differences in relative abundance of 43 ASVs were observed, but only 3 genera of low

abundance bacteria showed significant differences. Seventeen ASVs were found significantly

more abundant in intestine than stomach and 26 ASVs were found significantly more abundant in

stomach than intestine (Table 2). In particular, stomachs were enriched in various

Gammaproteobacteria of the order *Alteromonadales,* in *Alphaproteobacteria* of the order

Rickettsiales and *Sphingomonadales*, and a *Bacilli* of the order *Lactobacillales.* Intestines were

enriched in *Clostridia*, of the order *Clostridiales,* and in *Cytophaga* of the order *cytophagale.* Of

interest, different sequence variants of *Alphaproteobacteria* of the order *Rhizobiales,*

Betaproteobacteria of the order *Burkholderiales*, and *Spartobacteria* of the order

Chtoniobacteriales were enriched in the intestine and stomach of sticklebacks.

Similarly, only three low abundance bacteria were found differentially abundant between non-

infected and parasitized individuals (Table 2, S6 table) suggesting that different variants of

bacteria from the same genus are found in non-infected and parasitized individuals. Thirty-nine of

the ASVs were less abundant in parasitized individuals, with a majority of *Alphaproteobacteria* of

the order *Rickettsiales*, *Betaproteobacteria* of the order *Burkholderiales*, and

Gammaproteobacteria of the orders *Alteromonadales* and *Oceanospirillales*. Twenty-two ASVs

were more abundant in parasitized individuals including *Clostridia* of the order *Clostridiales*, and

Bacilli of the order *Bacillales, Alphapoteobacteria* of the order *Rhizobiales* and

Sphingomonadales, Gammaproteobacteria of the order *Aeromonadales* and *Spirochaetes* of the

order *Borreliales*.

Discussion

In this study, we analyzed the effect of infection by *S. solidus* on the composition of the stomach,

(which is often omitted from gut microbial composition analyses) , and intestine microbiome of

its threespine stickleback host. Both RT-qPCR analyses used to quantify 16S rRNA gene

abundance and 16S rRNA sequencing used to characterize microbiome diversity and composition

revealed significant differences in the microbiome of both the stomach and intestine of the fish,

even though the parasite was not in contact with these microbes anymore. Helminths have been

found to impact the gut microbiome of humans and livestock [18-20, 22, 23, 26]. This modification

of the host microbiome has been suggested to participate in strategies to impact the host immune

system and establish infection [36]. However, these studies focused on intestinal parasites. In this

study, we found that parasite infection is associated with an increase in abundance of 16S rRNA

genes, but a reduction in phylogenetic diversity associated with a reduction of beta diversity and a

small shift in bacterial microbiome composition. The correlation between our observation of

changes in fish gut microbiome composition and infection status could result from two non-

mutually exclusive hypotheses: non-infected individuals may carry defensive bacteria that reduce

the chance of successful infection by *S. solidus*, or *S. solidus* infection may result in an alteration

of the host microbiome. Our study identifies a number of ASVs significantly differentially

abundant in the stomach and/or intestine of non-infected and parasitized individuals that could be

targeted in future studies to test these hypotheses using laboratory controlled experimental

infections.

Plerocercoids of *S. solidus* were not in contact with the host gut microbiome at the time of sample

collection. Therefore, two hypotheses could be explored going forward to determine how *S. solidus*

infection might impact its host microbiome. It is possible that *S. solidus* directly interacts with

stomach and intestine microbes during the first 24 hours of infection, before it transits into the

body cavity of the fish [30]. The host gut microbiome can have important effects on host fitness

and health, including immune and metabolic function [37, 38]. Thus, early alteration of the host

gut microbiome at the onset of infection could provide long-term advantages to the parasite.

Alternatively, changes in the fish microbiome composition could result from cross-talks between

the microbiome and the immune system. Indeed, *S. solidus* alters different pathways of its host

immune system at different time points over the course of an infection [39], that could indirectly

impact the fish ability to regulate its microbiome composition. Also, a comparison of germ-free

and conventional threespine sticklebacks revealed variations in the recruitment of granulocytes,

and in expression of immune response genes [40, 41](including IL-22) that play key roles in host

protection and facilitation of helminth establishments[42, 43]. Further studies are therefore

necessary to determine at which time point *S. solidus* infection results in changes in the stickleback

microbiome.

It should be noted here that this study focused on field-caught individuals. Given the very high

prevalence of *S. solidus* in Cheney lake [44], it is very likely that all non-infected individuals had

been exposed to *S. solidus*. As such, parasite exposure, if not parasite infection, probably represents

gut homeostasis in natural conditions. It means that more important differences can be expected

from future laboratory control experiments focusing on the differences in the microbiome

composition of healthy non-exposed sticklebacks, and both exposed but non-infected and

successfully infected sticklebacks. Because high parasite prevalence is highly common in field

samples of any species, and because this study, and many others have found a significant

correlation between parasitism and changes in microbiome composition, we would like to

conclude by emphasizing that the presence of macro-parasites in field caught individuals should

always be assessed before conducting a microbiome study in natural conditions.

Methods

Sample collection

In September 2015, threespine stickleback (*Gasterosteus aculeatus*) specimens were collected live

from Cheney Lake in Anchorage Alaska (61° 12' 17" N, 149° 45' 33" W). Fish were caught using

un-baited minnow traps placed along the north shoreline of the lake between 0.25 and 2 m deep,

and at least 2 m apart. As stickleback have relatively short lifespans and it is easy to distinguish

young of the year fish from fish that are 1+ or 2+ years old, we took fish that ranged from 3-4 cm

in length to ensure that we sampled from the same cohort [45]. Fish were shipped to Stony Brook

University, kept in 20-gallon tanks of 6% seawater at 5°C at the Flax Pond Marine Lab, and fed

mysis shrimp twice per day.

Following an acclimation period of 3 weeks, and 12 hours of starvation, fish were anaesthetized

and euthanized in tricaine methane sulphonate solution (MS222), decapitated, and dissected in

sterile conditions. Prior to dissections all surfaces were cleaned with 80% ethanol. Between each

subsequent fish dissection, tools and dissecting plate were cleaned with 80% ethanol and betadine

to remove potential contaminants. Before dissection, fish were brushed with betadine to prevent

contamination of the body cavity with microbes found on the surface of the fish. For dissection,

an incision was made along the lateral line of the fish body, around the bony pelvis. The cut

extended from the pectoral fins to just anterior of the anus to avoid cutting the intestine. After the

initial incision, the bony pelvis was pulled away, opening up the body cavity. The sex of the fish

was assessed by visual inspection of gonads at the time of dissection and then confirmed using

PCR with sex specific primers as described in (REF). The presence of *S. solidus* was recorded, and

stomach and intestines were flash frozen in liquid nitrogen for future DNA extraction. The number

of parasites per infected fish is provided in table 3. Swab samples of the body cavity of two non-

infected and two parasitized individuals were also collected. All samples were then stored at -80°C

until use.

Table 3: Sex of threespine sticklebacks used in this study and number of *S. solidus* parasite

found within parasitized individuals.

Non-infected Fish	Sex	Parasitized Fish	Sex	# of parasites
Ga1		Ga7		3
Ga2		Ga8		4
Ga3		Ga9		4
Ga4		Ga10		1
Ga5		Ga11		1
Ga6		Ga12		2

* M: Male; F: Female

Ethics statement

Stickleback collection followed guidelines for scientific fish collection by the State of Alaska

Department of Fish and Game in accordance with Fish sampling permit #SF2015-263 and #P17-

025 and fish transport permits #15A-0084 and 17A-0024 provided to NMD. Fish were maintained

at Stony Brook University under the License to collect or possess #1949 provided by the New

York State Department of Environmental Conservation to NMD. Fish experiments were conduct

following protocols described in Institutional Animal Care and Use Committee (IACUC) #237429

and # 815164 to Michael Bell and NMD, respectively. Fish euthanasia was conducted using

MS222 and decapitation before parasite and tissue sampling. All experiments were performed in

accordance with relevant guidelines and regulations in the Public Health Service Policy (PHS) on

Humane Care and Use of Laboratory Animals.

DNA extraction

Tissues from six non-infected fish (HGa) and six infected fish (PGa) were collected for DNA

extraction. DNA purification from stomach (St) and intestine (In) tissues was performed using a

MoBIO Powerlyzer Powersoil Kit (Cat#12855) with the following modifications to the

manufacturer's protocol. First, all tissue samples were placed in tough tubes with 3mm glass beads.

To each tube, we added 750μL of the bead beating solution supplied with the extraction kit.

Samples underwent three rounds of 20 sec bead beating cycles and were kept on ice between each

round. The homogenate was transferred to the MoBIO supplied 0.1mm bead tube and incubated

with 60μL of the provided C1 solution for 10 min at 60°C. After vortexing, centrifugation, and

filtration with solution C5 according to manufacturer's protocol, 75μL of solution C6 was added

to the spin filter in a clean 2mL tube and incubated at room temperature for 5 min before

centrifugation at 10,000g for 30 sec. This final step was repeated to obtain a final DNA extract of

150μL.

Real-time qPCR

Bacteria from all samples were quantified by qPCR using the QuantStudio 6 Flex RT PCR System

(Fisher Scientific). Two 16S primers were used (Forward: TCCTACGGGAGGCAGCAGT,

Reverse: GACTACCAGGGTATCTAATCCTGTT [46] and

Forward: GTGSTGCAYGGYTGTCGTCA, Reverse: ACGTCRTCCMCACCTTCCTC [47]) and

compared to an *E. coli* standard for which cell count was obtained using a hemocytometer (Stony

Brook, NY). The qPCR mixture (20μL) was composed of 10μL of SYBR green master mix (CAT#

4309155), 4.8μL Molecular biology grade water (CAT#46000), 0.6μL forward primer, 0.6uL

reverse primer, and 4μL DNA.

16S rDNA amplification and Illumina MiSeq sequencing

We sequenced the stomach and intestine microbiota from six non-infected and six parasitized

individuals. Swabs of the body cavity, and negative controls from the DNA extraction were also

sequenced. The 16S rDNA V4-V5 hypervariable region was amplified with *E. coli* 515f and 806r

primers (with barcode on the forward primer) as specified by the Earth Microbiome Project [48-

51]. We used a 28 cycle PCR (5 cycle used on PCR products) using the HotStarTaq Plus Master

Mix Kit (Qiagen, USA) under the following conditions: 94°C for 3 minutes, followed by 28 cycles

of 94°C for 30 seconds, 53°C for 40 seconds and 72°C for 1 minute, after which a final elongation

step at 72°C for 5 minutes was performed. After amplification, PCR products are checked in 2%

agarose gel to determine the success of amplification and the relative intensity of bands. Multiple

samples are pooled together (e.g., 100 samples) in equal proportions based on their molecular

weight and DNA concentrations. Pooled samples are purified using calibrated Ampure XP beads.

Then the pooled and purified PCR product is used to prepare illumina DNA library. Sequencing

was performed at MR DNA to obtain 250bp paired end reads (www.mrdnalab.com, Shallowater,

TX, USA) on a MiSeq following the manufacturer's guidelines. Sequence data were initially

processed using MR DNA analysis pipeline (MR DNA, Shallowater, TX, USA). In summary,

sequences were joined, depleted of barcodes then sequences <150bp removed, sequences with

ambiguous base calls removed.

Microbiome composition analysis

The 16S rDNA gene Illumina reads were further processed using methods implemented in

QIIME2, Phyloseq [52, 53] and R as detailed in the supplementary information (Supp. File 1).

Briefly, samples were demultiplexed and quality filtered. ASVs (amplicon sequence variants) were

picked against the Greengene database using the DADA2 pipeline. . The final ASV table was then

rarefied to a depth of 3000 sequences per sample for comparisons between the microbiomes of *S.*

solidus and stickleback tissue samples, and 10,000 sequences for comparisons among stickleback

tissue samples. Rarefaction curves showed that at these depths of sampling, we were able to sample

a large portion of the ASV diversity present while still retaining all samples (Supp File 2).

Diversity measures and statistical analyses

Analyses were performed using QIIME2 and the Phyloseq and DESeq2 packages in R [53, 54].

Each of the following diversity metrics and statistical analyses were carried out at the 10,000

rarefaction levels to perform comparisons (i) among all sample categories (HGaIn, PGaIn, HGaSt,

PGaSt), (ii) between tissue (all In vs all St), and (iii) depending on infection status (all HGa versus

all PGa). Pielou's Evenness,   Shannon, Simpson, and Faith's Phylogenic Diversity

(PD_Whole_tree, with Greegene taxonomy tree) alpha diversity metrics were used to estimate

within species diversity. Alpha diversity was compared between sample types using pairwise

Kruskal-Wallis H-tests. Beta diversity was calculated with both weighted and unweighted UniFrac

metrics. PERMANOVA at 10,000 permutations was carried out to identify significant differences

between categories of samples. We tested for homogeneity of multivariate dispersion using

PERMDISP. Principal coordinate analysis (PCoA) was performed to look for patterns in

unconstrained multivariate space and a combination of paired and unpaired pairwise t-tests were

used to test if axes significantly discriminate groups of samples. DESeq2 was used on raw data to

examine differences in individual microbe abundance.

Data Accessibility

323   All data is being made available through the NCBI database under BioSample accessions

324   BioSample accessions SAMN08800337 to SAMN08800360, SAMN08805227, SAMN08805228,

325   SAMN08805230 and SAMN08805232.

Acknowledgements

We would like to acknowledge the expertise and assistance of Dr. Michael Bell in fish collection

and dissection techniques.

References

1.   Margulis L. Symbiogenesis and Symbionticism1991. 1-14 p.

2.   Dheilly NM. Holobiont–Holobiont interactions: redefining host–parasite interactions. PLoS Pathog. 2014;10(7):e1004093.

3.   Oliver KM, Russell JA, Moran NA, Hunter MS. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. Proc Natl Acad Sci. 2003;100(4):1803-7.

4.   Teixeira L, Ferreira Á, Ashburner M. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. PLoS Biol. 2008;6(12):e1000002.

5.   Xie J, Butler S, Sanchez G, Mateos M. Male killing *Spiroplasma* protects *Drosophila melanogaster* against two parasitoid wasps. J Hered. 2014;112(4):399.

6.   Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL. *Wolbachia* infections are virulent and inhibit the human malaria parasite *Plasmodium falciparum* in *Anopheles gambiae*. PLoS Pathog. 2011;7(5):e1002043.

7.   Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, et al. The genome sequence of *Drosophila melanogaster*. Science. 2000;287(5461):2185-95. Epub 2000/03/25. PubMed PMID: 10731132.

8.   Aliota MT, Peinado SA, Velez ID, Osorio JE. The wMel strain of *Wolbachia* reduces transmission of Zika virus by *Aedes aegypti*. Sci Rep. 2016;6:srep28792.

9.   Dutra HLC, Rocha MN, Dias FBS, Mansur SB, Caragata EP, Moreira LA. *Wolbachia* blocks currently circulating Zika virus isolates in Brazilian *Aedes aegypti* mosquitoes. Cell Host Microbe. 2016;19(6):771-4.

10.   Pan X, Thiem S, Xi Z. Wolbachia-Mediated Immunity Induction in Mosquito Vectors. Arthropod Vector: Controller of Disease Transmission, Volume 1: Elsevier; 2017. p. 35-58.

11.   Carrington LB, Tran BCN, Le NTH, Luong TTH, Nguyen TT, Nguyen PT, et al. Field-and clinically derived estimates of *Wolbachia*-mediated blocking of dengue virus transmission potential in *Aedes aegypti* mosquitoes. Proc Natl Acad Sci USA. 2018;115(2):361-6.

12.   Wang S, Dos-Santos AL, Huang W, Liu KC, Oshaghi MA, Wei G, et al. Driving mosquito refractoriness to *Plasmodium falciparum* with engineered symbiotic bacteria. Science. 2017;357(6358):1399-402.

13.   Knutie SA, Wilkinson CL, Kohl KD, Rohr JR. Early-life disruption of amphibian microbiota decreases later-life resistance to parasites. Nature communications. 2017;8(1):86.

14.   Theriot CM, Koenigsknecht MJ, Carlson Jr PE, Hatton GE, Nelson AM, Li B, et al. Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase susceptibility to *Clostridium difficile* infection. Nature communications. 2014;5:3114.

15.   Schuijt TJ, Lankelma JM, Scicluna BP, e Melo FdS, Roelofs JJ, de Boer JD, et al. The gut microbiota plays a protective role in the host defence against pneumococcal pneumonia. Gut. 2015:gutjnl-2015-309728.

16.   Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K, et al. Wild
mouse gut microbiome protects laboratory mice against lethal influenza virus infection and
colorectal cancer. Am Assoc Immnol; 2017. p. 5.
17.   Schmid-Hempel P. Immune defence, parasite evasion strategies and their relevance for
'macroscopic phenomena'such as virulence. Philos Trans R Soc Lond B Biol Sci.
2009;364(1513):85-98.
18.   Wu S, Li RW, Li W, Beshah E, Dawson HD, Urban Jr JF. Worm burden-dependent
disruption of the porcine colon microbiota by *Trichuris suis* infection. PloS ONE.
2012;7(4):e35470.
19.   Li RW, Wu S, Li W, Huang Y, Gasbarre LC. Metagenome plasticity of the bovine
abomasal microbiota in immune animals in response to *Ostertagia ostertagi* infection. PloS ONE.
2011;6(9):e24417.
20.   Nicholls C. Endoscopy, physiology and bacterial flora of sheep infected with abomasal
nematodes. 1987.
21.   Li RW, Wu S, Li W, Navarro K, Couch RD, Hill D, et al. Alterations in the porcine colon
microbiota induced by the gastrointestinal nematode *Trichuris suis*. Infect Immun.
2012;80(6):2150-7.
22.   Li RW, Li W, Sun J, Yu P, Baldwin RL, Urban JF. The effect of helminth infection on the
microbial composition and structure of the caprine abomasal microbiome. Sci Rep. 2016;6.
23.   Cantacessi C, Giacomin P, Croese J, Zakrzewski M, Sotillo J, McCann L, et al. Impact of
experimental hookworm infection on the human gut microbiota. J Infect Dis. 2014;210(9):1431-
4.
24.   Lee SC, San Tang M, Lim YA, Choy SH, Kurtz ZD, Cox LM, et al. Helminth colonization
is associated with increased diversity of the gut microbiota. PLoS Negl Trop Dis. 2014;8(5):e2880.
25.   McKenney EA, Williamson L, Yoder AD, Rawls JF, Bilbo SD, Parker W. Alteration of
the rat cecal microbiome during colonization with the helminth *Hymenolepis diminuta*. Gut Micro.
2015;6(3):182-93. doi: 10.1080/19490976.2015.1047128. PubMed PMID: PMC4615828.
26.   Plieskatt JL, Deenonpoe R, Mulvenna JP, Krause L, Sripa B, Bethony JM, et al. Infection
with the carcinogenic liver fluke *Opisthorchis viverrini* modifies intestinal and biliary microbiome.
FASEB J. 2013;27(11):4572-84.
27.   Chng KR, Chan SH, Ng AHQ, Li C, Jusakul A, Bertrand D, et al. Tissue microbiome
profiling identifies an enrichment of specific enteric bacteria In *Opisthorchis viverrini* associated
cholangiocarcinoma. EBioMedicine. 2016.
28.   Itthitaetrakool U, Pinlaor P, Pinlaor S, Chomvarin C, Dangtakot R, Chaidee A, et al.
Chronic *Opisthorchis viverrini* Infection Changes the Liver Microbiome and Promotes
Helicobacter Growth. PloS ONE. 2016;11(11):e0165798.
29.   Kreisinger J, Bastien G, Hauffe HC, Marchesi J, Perkins SE. Interactions between multiple
helminths and the gut microbiota in wild rodents. Phil Trans R Soc Biol.
2015;370(1675):20140295.
30.   Hammerschmidt K, Kurtz J. *Schistocephalus solidus*: Establishment of tapeworms in
sticklebacks–fast food or fast lane? Exp Parasitol. 2007;116(2):142-9.
31.   Kurtz J, Wegner KM, Kalbe M, Reusch TB, Schaschl H, Hasselquist D, et al. MHC genes
and oxidative stress in sticklebacks: an immuno-ecological approach. Philos Trans R Soc Lond B
Biol Sci. 2006;273(1592):1407-14.

32.   Kurtz J, Kalbe M, Aeschlimann PB, Häberli MA, Wegner KM, Reusch TB, et al. Major
histocompatibility complex diversity influences parasite resistance and innate immunity in
sticklebacks. Philos Trans R Soc Lond B Biol Sci. 2004;271(1535):197-204.
33.   Hammerschmidt K, Kurtz J. Evolutionary implications of the adaptation to different
immune systems in a parasite with a complex life cycle. Philos Trans R Soc Lond B Biol Sci.
2005;272(1580):2511-8.
34.   Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. ISME. 2017;11(12):2639.
35.   Glassing A, Dowd SE, Galandiuk S, Davis B, Chiodini RJ. Inherent bacterial DNA
contamination of extraction and sequencing reagents may affect interpretation of microbiota in
low bacterial biomass samples. Gut pathog. 2016;8(1):24.
36.   Dheilly NM, Poulin R, Thomas F. Biological warfare: Microorganisms as drivers of host–
parasite   interactions.   Infect   Genet   Evol.   2015;34:251-9.   doi:
dx.doi.org/10.1016/j.meegid.2015.05.027.
37.   Round JL, Mazmanian SK. The gut microbiota shapes intestinal immune responses during
health and disease. Nat Rev Immunol. 2009;9(5):313-23.
38.   Devaraj S, Hemarajata P, Versalovic J. The human gut microbiome and body metabolism:
implications for obesity and diabetes. Clin Chem. 2013;59(4):617-28.
39.   Scharsack JP, Koch K, Hammerschmidt K. Who is in control of the stickleback immune
system: interactions between *Schistocephalus solidus* and its specific vertebrate host. Proc R Soc
Lond [Biol]. 2007;274(1629):3151-8.
40.   Milligan-Myhre K, Small CM, Mittge EK, Agarwal M, Currey M, Cresko WA, et al. Innate
immune responses to gut microbiota differ between threespine stickleback populations. Disease
Models and Mechanisms. 2015:dmm. 021881.
41.   Small CM, Milligan-Myhre K, Bassham S, Guillemin K, Cresko WA. Host genotype and
microbiota contribute asymmetrically to transcriptional variation in the threespine stickleback gut.
Genome biology and evolution. 2017;9(3):00-.
42.   L Makepeace B, Martin C, D Turner J, Specht S. Granulocytes in helminth infection-who
is calling the shots? Current medicinal chemistry. 2012;19(10):1567-86.
43.   Turner J-E, Stockinger B, Helmby H. IL-22 mediates goblet cell hyperplasia and worm
expulsion in intestinal helminth infection. PLoS pathogens. 2013;9(10):e1003698.
44.   Bell MA, Heins DC, Wund MA, von Hippel FA, Massengill R, Dunker K, et al.
Reintroduction of threespine stickleback into Cheney and Scout Lakes, Alaska. Evol Ecol Res.
2016;17(2):157-78.
45.   Heins D, Eidam D, Baker J. Timing of Infections in the Threespine Stickleback
(Gasterosteus aculeatus) by Schistocephalus solidus in Alaska. The Journal of parasitology.
2016;102(2):286-9.
46.   Larsen N, Vogensen FK, van den Berg FW, Nielsen DS, Andreasen AS, Pedersen BK, et
al. Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. PloS one.
2010;5(2):e9085.
47.   Horz H, Vianna M, Gomes B, Conrads G. Evaluation of universal probes and primer sets
for assessing total bacterial load in clinical samples: general implications and practical use in
endodontic antimicrobial therapy. J Clin Microbiol. 2005;43(10):5332-7.
48.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.
ISME. 2012;6(8):1621-4.

49.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et
al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc
Natl Acad Sci USA. 2011;108(Supplement 1):4516-22.
50.   Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, et al.
Improved bacterial 16S rRNA gene (V4 and V4-5) and fungal internal transcribed spacer marker
gene primers for microbial community surveys. mSystems. 2016;1(1):e00009-15.
51.   Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA
primers for marine microbiomes with mock communities, time series and global field samples.
Environ microbiol. 2015.
52.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: high-
resolution sample inference from Illumina amplicon data. Nature methods. 2016;13(7):581-3.
53.   McMurdie PJ, Holmes S. Phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PloS ONE. 2013;8(4):e61217.
54.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al.
QIIME allows analysis of high-throughput community sequencing data. Nature methods.
2010;7(5):335-6.

Supporting information

474 **S1 File: Document contains S1 to S7 Figs with corresponding legends**

475 **S2 File: Contains S1 to S9 tables with corresponding legends**

Figures

480 **Fig 1 Bacterial load as estimated using qPCR of 16S gene**. Bacterial load in the stomach (St)

481 and intestine (In) of non-infected *G. aculeatus* (HGa), and parasitized *G. aculeatus* (PGa).

Fig 2. Comparison of alpha diversity metrics and principal coordinates analyses on beta-diversity metrics. A. alpha diversity values in the stomach (St) and intestine (In) of non-infected *G. aculeatus* (HGa), and parasitized *G. aculeatus* (PGa). B. Principal coordinates analysis based on Unweighted Unifrac distance at 10,000 rarefaction level showing differences between HGa and PGa.

490 **Fig 3**. **Relative abundance of the 25 most abundant ASVs in the intestine (A and C) and**

491 **stomach (B and D) of non-infected (A and B, samples Ga1 to 6) and parasitized (C and D,**

492 **samples Ga 7 to 12) Threespine Sticklebacks.** The legend indicates the family (f_) and genus

493 (g_) names.

494