Title: Sexual reproduction in the fungal foliar pathogen *Zymoseptoria tritici* is driven by

antagonistic density-dependence mechanisms

Running head: Density dependence in *Zymoseptoria tritici* sexuality

Authors: Frédéric Suffert *, Ghislain Delestre, Sandrine Gélisse [1]

UMR BIOGER, INRA, AgroParisTech, Université Paris-Saclay, 78850 Thiverval-Grignon,

France.

* corresponding author ; ORCID 0000-0001-6969-3878 ; email: frederic.suffert@inra.fr

[1] The authors are listed in descending order of the importance of their contributions.

Acknowledgments: We thank Nathalie Retout (INRA BIOGER, France) for technical

assistance and Dr. Frédéric Hamelin (Agrocampus Ouest, France) for preliminary discussions

on ecological modeling aspects. We thank Julie Sappa for her help correcting our English. We

thank Dr. Alexey Mikaberidze and Prof. Bruce McDonald (ETH Zürich, Switzerland) for

providing the ST99CH-3D7 and ST99CH-1A5 strains of *Z. tritici* and seeds of the wheat cv.

Runal. This study was supported by a grant from the European Union Horizon Framework

2020 Program (Grant Agreement no. 634179, EMPHASIS project) covering the 2015-2019

period.

Abstract

This study provides empirical evidence for antagonistic density-dependence mechanisms driving sexual reproduction in the wheat fungal pathogen *Zymoseptoria tritici*. Biparental crosses with 12 increasing inoculum concentrations, in controlled conditions, showed that sexual reproduction in *Z. tritici* was impacted by an Allee effect due to mate limitation and a competition with asexual multiplication for resource allocation. We found that asexual multiplication was itself affected by competition for host resources between the two parental isolates, as illustrated by the decrease in asexual fruiting body density and sporulation intensity observed with increasing lesion density. Consistent with these results for controlled co-inoculation, we found that the intensity of sexual reproduction varied with both cropping period and the vertical position of the host tissues in the field. An optimal lesion density maximizing offspring (ascospores) number was established, and its eco-evolutionary consequences are considered here. Two ecological mechanisms may be involved: competition for resources between the two modes of reproduction (decrease in the host resources available for sexual reproduction due to their prior use in asexual multiplication), and competitive disequilibrium between the two parental isolates, due to differential interaction dynamics with the host, for example, leading to an imbalance between mating types. Conceptual models based on these results suggest that sexual reproduction plays a key role in the evolution of pathogenicity traits, including virulence and aggressiveness. Ecological knowledge about the determinants of sexual reproduction in *Z. tritici* may, therefore, open up new perspectives for the management of other fungal foliar pathogens with dual modes of reproduction.

Keywords

43 Allocation resource, asexual multiplication, competition, fungal pathogen, plant disease

44 epidemiology, sexual reproduction

Introduction

An ability to reproduce is one of the most fundamental features of life. It has been suggested

that the benefits of sexual reproduction include the purging of deleterious mutations from the

genome and the production of a recombinant progeny well-adapted to a changing

environment. Density-dependent mechanisms are common in microbial ecology and play a

crucial role in sexual reproduction. The modeling of density-dependent selection would shed

light on the ability of populations to persist despite "sex"-related conflicts, but simple models

seem unable to predict the diversity of the responses observed in nature [1]. A few empirical

studies have focused on the role of sex in influencing fungal population and community

structures [2, 3].

Several density-dependent mechanisms may operate simultaneously in a population, and it is

important to understand how they act, both independently and together. Positive (facilitation)

density-dependent mechanisms, such as demographic Allee effects (increasing relationship

between overall individual fitness and population density), linked to mate-finding, for

example [4], are frequently taken into account in population dynamics studies. Negative

(restriction) density-dependent mechanisms, such as competition for resources in cases of

high population density, have also been widely described. The effects of competition for

resources on individual fitness at low and high densities are a key, complex question in

population and evolutionary ecology. For instance, Neiman et al. [5] detected a strong

positive effect of high population density on reproductive output in the presence of adequate

food supplies in *Potamopyrgus antipodarum* (freshwater snail), but they also identified food

limitation as the primary source of negative density dependence. Generally, in situations of

high population density, negative density dependence often erases the demographic Allee

effect through resource competition.

The impact of density-dependent mechanisms on sexual reproduction has mostly been studied

in animal ecology, where encounters between male and female animals result from active

behavior influenced by sexual dimorphism or mating advantage [1]. For instance, Stelzer [6]

focused on two different mechanisms of density-dependent population regulation — resource

exploitation and sexual reproduction — in populations of the rotifer *Brachionus calyciflorus*

displaying different investments in sex: populations reproduce clonally at low densities, with

the proportion of sexual individuals increasing at higher densities. These two mechanisms

have been shown to apply to several facultatively sexual animals, but similar antagonistic

density-dependent mechanisms may also operate in fungal microorganisms with dual modes

of reproduction.

80   *   *   *

Competitive interactions and their effect on fitness during reproduction, particularly for

asexual reproduction, have been investigated in the emblematic fungal plant pathogens

Puccinia graminis f. sp. *tritici*, the causal agent of wheat stem rust [7], and *Phytophthora*

infestans, the causal agent of potato late blight [8]. The findings obtained were consistent with

the ecological principle that resource availability decreases over time, with increasing host

tissue mortality, but also during multiple infections, in which different parasite strains share

the host resources. Clément et al. [8] focused on the consequences of multiple infections on

asexual reproduction, and dissociated the effects of single-site and double-site infections and

those of competition between identical and different isolates during double-site infections.

Allee effects due to mate limitation have been reported to affect the dynamics of some fungal

pathogens at large spatial scales, such as *Hymenoscyphus fraxineus*, the causal agent of ash

dieback [9], and *Mycosphaerella fijiensis*, the causal agent of black Sigatoka disease in

banana plants [10]. Sexual spores for dispersal are produced only when compatible mating

types come into direct contact. However, mate limitation at the plant scale was not confirmed

by experimental studies, probably because there are still many gaps in our knowledge

concerning the ways in which sexual reproduction occurs in host tissues and its determinants.

However, this effect forms the theoretical basis of demogenetic models combining sexual and

asexual reproduction, in which the impact of mate-finding limits the speed of pathogen spread

[11, 12].

Thus, on the one hand, pathogen population density can affect competition between

individuals for limited host resources at a given reproductive stage (e.g. asexual reproduction)

or the competition between asexual and sexual stages. On the other hand, it may also affect

the probability of finding a mate and, thus, the intensity of sexual reproduction. Antagonistic

interactions between negative and positive density-dependent mechanisms have never been

studied experimentally in the reproduction of fungal plant pathogens. They could have been

investigated by Clément et al. [8] for *P. infestans* co-infections with isolates of mating types

A1 and A2 [13], but sexual reproduction was deliberately prevented by inoculation with

incompatible isolates only.

109   *   *   *

Sexual reproduction in *Zymoseptoria tritici*, the causal agent of Septoria leaf blotch of wheat,

requires a physical encounter between two compatible strains (Mat1-1 and Mat1-2; [14, 15]).

It generally occurs on senescent tissues (leaf and stem debris), with pseudothecia (sexual

fruiting bodies producing ascospores) appearing 46 to 76 days after the initial infection in

field conditions [16]. The very active sexual cycle of *Z. tritici* has crucial consequences for

the epidemiology of Septoria leaf blotch, because it determines the amount of primary

inoculum (wind-dispersed ascospores produced on wheat debris) available for the initiation of

subsequent epidemics [17]. Sexual reproduction also plays an essential role in maintaining the

diversity of local pathogen populations and determining their rate of adaptation to selection

pressures, such as those exerted by resistant hosts or fungicide treatments. Given the high

population diversity, whatever the spatial scale considered [18], a wheat plant should be

considered to be subject to "multiple infections". The sex ratio is consequently close to 1 at

the local scale (Morais et al., unpubl.) and the imbalance effects sometimes reported for other

organisms are negligible.

Avirulent *Z. tritici* strains were recently shown to cross with virulent strains on wheat

cultivars carrying the corresponding resistance gene [19]. This key finding calls into question

current views about the determinants of sexual reproduction, including, in particular, the

common belief that individuals unable to infect wheat tissues cannot reproduce sexually.

However, the proportion of crossing events resulting from the "exclusive paternal

parenthood" model was not quantified at the leaf scale in the natural context of highly diverse

populations. The vast majority of crossing events probably occur when leaf lesions caused by

two compatible strains either coalesce or are located very close together. In practice, it

remains unclear how close together infections must be for effective mating to occur. Metcalfe

[20] found internal hyphae spanning mean distances of no more than 11 mm in a susceptible

cultivar just before the production of pycnidia, suggesting that infection foci located more

than 20 mm apart may be isolated. The number of offspring (i.e. the number of ascospores)

should be proportional to the number of crossing events, and correlated with disease severity,

which is a good proxy for lesion density. This has been shown at field scale: more severe

epidemics are associated with higher levels of ascospore production during epidemics in the

following year [21, 22]. Moreover, the fitness (aggressiveness) of the *Z. tritici* parental

isolates and a time lag between the inoculations with the two parental strains can affect sexual

reproduction intensity [23].

The density-dependent mechanisms driving the interaction between *Z. tritici* and wheat during

the asexual stage remain unclear. In particular, it remains unknown how the fungus exploits

the resources of it host to obtain nutrients for growth, how it invades the apoplast during the

long asymptomatic infection phase, and how the necrotrophic phase triggered [24].

Objective and strategy

We investigated the density dependence of both asexual and sexual reproductive mechanisms

in *Z. tritici*, a relevant fungal model organism for such eco-epidemiological investigations.

What impact does lesion density in host tissues have on sexual reproduction at the leaf scale?

Could we generate empirical data for the establishment of a density-dependence relationship?

Would this relationship be monotonic or more complex, due to competition for resources

between the two modes of reproduction? This study addressed these questions. We first

analyzed epidemiological datasets collected in field conditions, to confirm our initial

assumptions. We then assessed the intensities of asexual and sexual reproduction after five *Z.*

tritici biparental crosses on adult wheat plants over a period of three years.

Materials & methods

Preliminary analysis of field epidemiological data

Field epidemiological data were analyzed to assess the impact of disease severity (proxy of

asexual multiplication intensity) and the vertical position of the host tissues on the intensity of

subsequent sexual reproduction, during the epidemic period and then during the interepidemic

period. We reanalyzed a dataset obtained in Danish field conditions by Eriksen & Munk [16],

together with the data obtained in our three-year experiment in France.

Epidemic period – The relationship between the proportion of pseudothecia among the total

fruiting bodies (pycnidia and pseudothecia) on wheat leaves during a growing season and

mean disease severity (expressed as the percentage of the leaf area covered by pycnidia and

pseudothecia) were determined with the experimental data for the field experiment performed

by Eriksen & Munk [16] (Table 1).

Interepidemic period – In a field plot sown with a monoculture of wheat cv. Soissons from

2007 to 2016 at the Grignon experimental station [22], standing stubble was left over an area

of 20 m^2 during the fall of 2015-2016, 2016-2017 and 2017-2018. Stubble samples were

collected three or four times each year, from September to January. Stems and leaves were

partitioned from lower to upper layers, according to their vertical position (0-5 cm, 5-10 cm,

10-15 cm, 15-25 cm, 25-35 cm, 35-50 cm, 50-65 cm). The dry tissues of each layer of wheat

stubble were weighed, and ascospore production, as a proxy of sexual reproduction intensity,

was quantified as described by Suffert et al. [17].

Fungal material

Five crosses were performed, with eight *Z. tritici* isolates (FS0802 × FS1006, FS0813 ×

FS0732, FS0808 × FS0806, FS1022 × FS1008 [25]) collected in 2010 from wheat cv.

Soissons in France, and two isolates (3D7 × 1A5 [26]) collected in 1999 from wheat cv.

Galaxie and Lono (Table 1). The compatibility of the five pairs of isolates was determined by

PCR amplification of the two mating-type idiomorphs [15]. The effective ability of the eight

French isolates to reproduce sexually on adult plants was first checked in semicontrolled

conditions [23]. Subcultures of each isolate stored at -80°C were grown for five days in Petri

dishes containing PDA (potato dextrose agar, 39 g L^{-1}) at 18°C in the dark. For each isolate,

an initial conidial suspension was prepared with a Malassez counting chamber. Serial

dilutions (1:5) of each initial monoparental suspension were prepared, from 10^7 conidia mL^{-1}

to 50 conidia mL^{-1}. Twelve biparental suspensions were then prepared by mixing 25 mL each

of the appropriate monoparental suspensions and adding two drops of surfactant (Tween 20;

Sigma, France).

Plant material

Seeds of the wheat cv. Soissons or Runal (moderately susceptible to Septoria tritici blotch)

were sown on December 18 2014, December 14 2015, and December 15 2016 (Table 1) in

Jiffy peat pots, which were then kept in greenhouse conditions for two weeks. Seedlings were

vernalized in a growth chamber for 8 weeks at 8°C, with a 10-h light/14-h dark photoperiod.

They were then returned to the greenhouse and left to acclimate for one week before

transplantation into individual pots filled with 1 liter of commercial compost (Klasmann Peat

Substrat 4; Klasmann France SARL, France). We added 4 g of slow-release fertilizer

(Osmocote Exact 16-11-11N-P-K 3MgO Te) to each pot. The plants were also watered with

Hydrokani C2 fertilizer (Hydro Agri Spécialités, France), diluted 1:100 and poured into the

saucers under the pots. Plants were sprayed once with spiroxamine (Flexity SC at 300 g L^{-1};

BASF, France) for the specific prevention of powdery mildew (*Blumeria graminis* f.sp.

tritici), no later than six weeks before inoculation. During the growth period, the plants were

illuminated with natural daylight supplemented with 400 W sodium vapor lamps between

6.00 a.m. and 9.00 p.m. The air temperature was kept below 20°C during the 15-h light period

and above 12°C during the 9-h dark period. Plants were thinned to three stems per pot during

the growth period.

Inoculation procedure

Co-inoculations with the five biparental suspensions were carried out after the wheat heads had fully emerged (one in 2015, on April 30; two in 2016, on May 3; two in 2017, on April 25; Table 1). In 2017, two complete sets of wheat plants cv. Soissons were also inoculated with each monoparental suspension (FS1022 and FS1008). The suspensions were applied with an atomizer (Ecospray, VWR, France), on three adult plants (nine stems), as described by Suffert et al. [23]. Plants were turned during the 10-second spraying event, to ensure even coverage with inoculum. Infection was promoted and cross-contamination prevented by enclosing each trio of plants inoculated with the same suspension in a sealed transparent polyethylene bag humidified with distilled water for 72 h. Wheat plants were then kept in a greenhouse for about 13 weeks (from April 30 to July 24 2015, from 3 May to July 28 2016, from 25 April to August 2 2017; Figure 1a). Air temperature was kept above 18°C during the spring, and reached 30°C several times during summer.

Assessment of fitness traits during the asexual stage

Disease severity on leaves was assessed by eye five weeks after inoculation (on June 2 2015, June 6 2016 and May 27 2017), as the percentage of the leaf area displaying necrosis (mean for the F1, F2, and F3 leaves for each stem; 1, 2, 3 and 5%, thereafter by increments of 5% up to 100%; Figure 2). In 2016, the density of pycnidia (number of pycnidia divided by the necrotic area) and the proportion of pycnidia exuding a cirrhus (Figure 2b-c) were also estimated for plants inoculated with each of the two biparental suspensions, FS0813 × FS0732 and FS0808 × FS0806 (Table 1), from the necrotic area in a 6-cm leaf section from four randomly selected F1 leaves per inoculum concentration.

Promotion of sexual reproduction and ascosporogenesis

Co-inoculated plants were placed outdoors during the summer and fall (from July 24 2015 to October 9 2015, from July 28 2016 to November 14 2016, and from July 28 2017 to December 4 2017) to induce the maturation of pseudothecia (Figure 1d). Each trio of co-inoculated plants was tied up with stiff wire, with the stems and leaves clamped so as to prevent contact with other trios of plants. On October 14 2015, November 16 2016, and December 6 2017, dry leaves and stems from each trio of plants were cut into 2-cm fragments and allowed to dry, in separate open boxes, in a laboratory atmosphere, at 18°C for one week. Ascospore release was assessed immediately after drying, to check the maturity of the pseudothecia. The pseudothecia were considered mature in 2015 and 2017. By contrast, in 2016, the plant debris was placed outdoors again until January 16 2017, to promote pseudothecium maturation.

Assessment of the intensity of sexual reproduction

For each inoculum concentration, the debris was weighed, soaked in water for 20 min and spread on dry filter paper in a box (24 × 36 cm), the lid of which was left half open to decrease the relative humidity of the air around the debris. Eight Petri dishes (90 mm in diameter) containing PDA medium were then placed upside down 1 cm above the debris. The boxes were placed in the dark, at 18°C for 19 h. The Petri dishes were then closed and incubated in the same conditions. Each discharge event was replicated four times (from November 2015 to January 2016, from January 2016 to February 2017, and from December 2017 to January 2018).

The ascospores released onto the Petri dishes germinated after 24 h. Six days later, yeast-like

colonies resembling cream-colored convoluted heaps were observed. The colonies were

counted under a microscope four and seven days after ascospore discharge. It was assumed

that each colony resulted from the germination of a single ascospore, and that clusters of

colonies appeared just above mature pseudothecia from which several ascospores had been

discharged. A cumulative ascospore discharge index (ΣADI), defined as the total number of

ascospores discharged per gram of wheat debris [17, 23], was calculated for each discharge

event, as follows:

$$\Sigma ADI = \sum_{e=1}^{n} \frac{1}{8} \cdot \frac{x.y}{\pi.r^2} \cdot \frac{NBcol}{DW}$$ eq. 1

where NBcol is the total number of colonies in the eight Petri dishes, DW is the total dry

weight of debris spread out in the box, r is the radius of a Petri dish (4.5 cm), x and y are the

width and length of the box (24 cm and 36 cm, respectively), and n is the number of discharge

events ($n = 4$).

Assessment of yield components

Yield components were assessed in 2016 and 2017 (Table 1). At maturity, the spikes were cut

and hand-threshed to separate the kernels from the chaff. Kernels were then counted and

weighed to estimate the grain weight per spike (GWS) and thousand-kernel weight (TKW) for

each cross and inoculum concentration.

Data analysis

A quadratic model (eq. 2) was used to describe empirically the relationship between sexual

reproduction intensity (SRI) and asexual multiplication intensity (AMI).

$$SRI = a.\,AMI^2 + b.\,AMI + c \qquad\qquad \text{eq. 2}$$

where *a, b* and *c* are parameters.

A second model integrating two types of density-dependent constraints (eq. 3) was used. It

was based on a logistic function f(α,β,γ,AMI) (eq. 4) describing a positive density-dependent

mechanism and coupled with a function g(μ,σ,ρ,AMI) (eq. 5) describing a negative density-

dependent mechanism.

$$SRI = f(\alpha, \beta, \gamma, AMI) \times g(\mu, \sigma, \rho, AMI) \qquad\qquad \text{eq. 3}$$

$$f(\alpha, \beta, \gamma, AMI) = \alpha.\,e^{\left(-\beta.e^{-\gamma.IAM}\right)} \qquad\qquad \text{eq. 4}$$

$$g(\mu, \sigma, \rho, AMI) = \mu - (\mu - \sigma)\left(e^{\left(-\ln\mu.e^{(-\rho.IAM)}\right)} - 0.01\right) \qquad\qquad \text{eq. 5}$$

where α, β, γ, μ, σ, and ρ are parameters.

Both models were adjusted with the data generated by the five crosses, with the estimation of

SRI by determination of the cumulative ascospore discharge index (ΣADI) and of AMI with

the percentage of the total area displaying necrosis.

Results

Sexual reproduction in field conditions varies according to the cropping period and the

vertical position of the infected host tissues in the canopy

The epidemiological data obtained by Eriksen & Munk [16] in field conditions showed that

the proportion of pseudothecia (proxy for the intensity of sexual reproduction) present on

wheat leaves varies with disease severity (proxy for asexual multiplication intensity; Figure 3)

over the cropping season. Our reanalysis of these data made it possible to limit bias due to

fruiting dynamics (the latent period is much longer for pseudothecia than for pycnidia). The

proportion of pseudothecia was maximal for severities of 50 to 80%. This proportion

remained stable or even decreased at severities above 80%.

Monitoring of the dynamics of sexual reproduction on standing stubble over three

interepidemic periods revealed a vertical structuring of mean ascospore production on

senescent wheat tissues (Online Resource 1). This vertical structuration changed over time.

Ascospore numbers peaked in November (2015, 2016) or December (2017). Ascospore

production generally increased with distance from the ground, up to a height of 25 cm (Figure

4). This layer corresponds to the part of the stubble not usually exported at harvest. The 25-35

cm layer made the largest contribution to sexual reproduction. Above this height, ascospore

production follows the opposite trend, gradually decreasing between 35 and 65 cm. The

lowest layer of standing stubble, which includes the crown and what remains of the first

leaves to emerge before tillering, generated less than 5% of the total number of ascospores.

The general interaction between *Z. tritici* and wheat is density-dependent

The interaction between the pathogen and the host plant is clearly density-dependent.

Regardless of the year in which the cross was performed (2015, 2016 or 2017), the biparental

suspension used and the host cultivar (Soissons or Runal), inoculum concentration (total

number of *Z. tritici* conidia in the biparental suspension per mL) was positively related to the

estimated mean disease severity (percentage of necrotic area) for the three uppermost leaves

of the adult wheat plants (Figure 5). Soissons was particularly susceptible to the isolates with

which it was co-inoculated (mean sporulation area of 60% for an inoculum concentration of 5

$\times$ 10^5 conidia.mL^{-1}), whereas Runal was more resistant (35%). For both varieties, disease

severity (80%) was maximal at an inoculum concentration of 10^7 conidia.mL^{-1}.

Mean severity on the three uppermost leaves of adult wheat plants cv. Soissons and Runal

was inversely correlated with total grain weight per spike (GWS), and thousand kernel weight

(TKW): late (inoculation after flowering) but intense attacks (sporulation over 80% of the

area) decreased grain yield by up to 20-25% (Figure 6). *Z. tritici* attacks had a strong impact

on wheat yield, demonstrating that the interaction between the pathogen and the host plant

was density-dependent, i.e. a function of pathogen inoculum pressure.

Competition for resources affects asexual multiplication in *Z. tritici*

We observed robust, negative relationships between mean disease severity and the density of

pycnidia and between mean disease severity and the percentage of exuding pycnidia in 2016

(Figure 7): more intense *Z. tritici* attacks were associated with a lower density of pycnidia,

and a lower percentage of exuding pycnidia. For both FS813 $\times$ FS732 and FS808 $\times$ FS806

crosses, severe attacks (mean sporulation area of 80%) halved the density of pycnidia (150 vs.

300 pycnidia.cm^{-2}) and the percentage of exuding pycnidia (45 vs. 90%). Thus, the asexual

development of *Z. tritici* is affected by competition for resource allocation.

A comparison of the disease severity reached after inoculation with a biparental suspension

and after inoculation with a suspension containing only one isolate (with an equivalent

number of conidia to the mixture) highlighted competitive disequilibrium between the two

parental isolates. This finding provides a second line of evidence that the asexual

development of *Z. tritici* is driven by density-dependent processes. Soissons appeared to be

more susceptible to FS1008 than to FS1022, regardless of the concentration of the inoculum:

in practice, FS1008 was more aggressive than FS1022 (e.g. disease severity with FS1008 was

twice that with FS1022 for inoculum concentrations of 10^4 to 10^6 conidia.mL^{-1}; Figure 8). At

inoculum concentrations of 50 to 10^5 conidia.mL^{-1}, FS1008 induced a less severe attack (<

20% on average) if the other, less aggressive isolate (FS1022) was present at the same

concentration. This difference was significant provided that the number of FS1022 conidia

was the same. It disappeared at the highest inoculum concentrations (> 10^5 conidia.mL^{-1},

corresponding to a disease severity of 35-40%).

Competition with asexual multiplication for resource allocation affects sexual

reproduction in *Z. tritici*

Ascospores were collected after each of the five crosses in 2015, 2016 and 2017. The number

of ascospores was largest for the FS802 × FS1006 cross in 2015 and lowest for the FS808 ×

FS806 cross in 2016 (Online Resource 2; Figure 9). Moreover, in 2016, the first ascospores

were collected later in the season (January), due to relatively the dry weather from August to

September 2016 (65 mm of rainfall, versus the mean of 170 mm generally observed this

period). A similar effect of year was also observed with the ascomycetes *Leptosphaeria*

maculans (Marie-Hélène Balesdent, INRA BIOGER, pers. comm.) and *Septoria linicola*

(Annette Penaud, Terres Inovia, pers. comm.) in the same experimental area.

For all crosses, the number of ascospores discharged increased with inoculum concentration.

The smallest numbers of ascospores were collected for the lowest inoculum concentrations

(50 or 10^2 conidia.mL^{-1}). The number of ascospores discharged at the highest inoculum level

(10^7 conidia.mL^{-1}) was however systematically below the maximum number. For crosses on

Soissons, this maximum was reached for inoculum concentrations of 5×10^4 conidia.mL^{-1}

(FS802 × FS1006 and FS813 × FS732) to 10^6 conidia.mL^{-1} (FS813 × FS732 and FS1022 ×

FS1008). It was reached for an inoculum concentration of $5 × 10^6$ conidia.mL^{-1} for the cross

performed on Runal (3D7 × 1A5), on which disease severity was significantly lower than on

Soissons for inoculum levels of $5 × 10^2$ to $5 × 10^6$ conidia.mL^{-1} (Figure 5). This pattern was

even more marked for the relationship between effective disease severity (proxy for asexual

multiplication intensity) and the normalized number of ascospores (proxy for sexual

reproduction intensity; Figure 10). The quadratic model and a model integrating two density-

dependent constraints converged for three crosses (FS802 × FS1006, FS808 × FS806, FS1022

× FS1008), with the intensity of sexual reproduction increasing to a peak for disease severities

of 35 to 50% and then decreasing thereafter. This pattern was particularly marked for the

FS802 × FS1006 and FS808 × FS806 crosses. The initial ascending part of the curves

illustrates an Allee effect (difficulty finding mates at low pathogen densities), whereas the

second, descending part of the curve highlights competition for resource allocation between

the two modes of reproduction (fewer host resources available for sexual reproduction due

their previous mobilization for asexual multiplication).

Discussion

In field conditions, the different layers of wheat plants, from the soil to the head, contribute

differently to *Z. tritici* reproduction. Mean *Z. tritici* ascospore production was maximal

between 25 and 35 cm above the ground. This significant impact of vertical positioning is

complementary to the findings of Pfender & Wootke [27], who established that the

pseudothecia of *Pyrenophora tritici-repentis* survived well in standing stubble and upper

mulch layers, but not in straw or mulch located in the soil or directly on its surface. At least

two antagonistic mechanisms are likely to account for the increase and subsequent decrease in

sexual reproduction intensity with increasing height of the canopy layer above the ground.

Higher positions in the canopy are associated with a lower density of lesions. The probability

of crosses occurring, therefore, also decreases with height. Conversely, higher positions in the

canopy are occupied by younger wheat tissues, on which infections are likely to be more

recent, with a lower probability of ascosporogenesis having had time to occur. Lower down in

the canopy, the wheat tissues are older and the pseudothecia are already empty or degraded.

These empirical observations confirmed our initial assumption of antagonistic density-

dependent mechanisms controlling *Z. tritici* sexual reproduction in natural conditions.

402   * * *

Traits relating to asexual multiplication *in planta,* such as the density and exudation capacity

of pycnidia, are affected by competition between the two parental isolates for resources. This

result is consistent with the lower level of pycnidial coverage observed after inoculation with

mixtures of isolates than after inoculation with single isolates [28, 29]. The competitive

disequilibrium between the two parental isolates may also account for the density dependence

of asexual development in *Z. tritici*. This finding is consistent with previous results obtained

in the same experimental conditions in a previous study [23], in which an interval of two to

three weeks between inoculation with the first parental isolate and inoculation with the second

was found to be detrimental to ascosporogenesis. In this case, the host tissues were likely to

be colonized by the first isolate, leaving fewer host resources available for the second isolate,

consistent with the establishment of competition during the asexual stage of the disease cycle.

Interactions of this type were studied by Clement et al. [8], who identified two basic response

patterns, consisting of an increase or decrease in reproductive fitness in multiple infections of

potato leaves with *P. infestans,* depending on pathogen genotype. In the case of *Z. tritici,* the

activation of host defenses by the less virulent or less aggressive isolate may limit the

subsequent development of a second more aggressive isolate at low inoculum pressure.

Conversely, at high inoculum pressure, the host sites for infection are likely to be saturated,

regardless of the aggressiveness of the isolates causing the infection. The less aggressive

isolate (i.e. to which the host is less susceptible) may trigger defense responses that ultimately

prevent the more aggressive isolate from developing to the extent that it would have if used

alone for inoculation. The highest inoculum pressure was so high that this mechanism was not

expressed.

* * *

Sexual reproduction is driven by at least two density-dependent mechanisms with opposite

impacts: a positive Allee effect due to mate limitation, and with a negative effect due to

competition with asexual multiplication for resources. We provide here the first experimental

evidence that sexual reproduction in a fungal foliar pathogen is driven by antagonistic

density-dependent mechanisms.

Positive density dependence of sexual reproduction - Higher densities of *Z. tritici* lesions on

leaves and stems are associated with a higher probability of encounter and a higher intensity

of sexual reproduction. Mate limitation has been identified as the most common mechanism

underlying Allee effects [4]. This experimental result, obtained at the plant scale in semi-

controlled conditions, is supported by the positive correlations previously established in

natural conditions at both the field and landscape scales: ascospore production is generally

greatest after the most severe epidemics [21, 22].

Negative density dependence of sexual reproduction - Strong Septoria tritici blotch attacks (i.e

intense asexual multiplication) limit the resources available for sexual reproduction, resulting

in a detrimental effect. We showed that competition between the two parental isolates during

asexual multiplication (accentuated, for example, by differences in aggressiveness or a time

interval between the two inoculations) is plausible and may lead to an imbalance between

mating types. However, this explanation appears to be less relevant in natural conditions,

given the high diversity of the pathogen population even at the leaf scale (sex ratio close to 1).

As illustrated by the negative impact on fruiting and sporulating capacity, competition also

limits the potential development of parental isolates in a similar way: host resources are

overexploited during the asexual phase, to the detriment of sexual reproduction.

The optimal equilibrium between asexual and sexual reproduction in semi-controlled

conditions corresponds to a disease severity of 30 to 45%. This optimum severity is lower

than that observed in the field on the leaves of the lower plant layers, on which severity may

reach very high levels (80-100%) and which senesce faster than the leaves of the upper layers.

Eriksen & Munk [16] reported that pseudothecia form only at disease severities exceeding

38%, following the coalescence of lesions, and that the highest proportion of pseudothecia

occurs on leaves with a disease severity of 50 to 80%. Furthermore, the model we fitted to

these experimental data revealed no negative density-dependent effect, perhaps because the

epidemiological survey stopped earlier (August) than in our own experiment conditions,

suggesting that the pseudothecia may not have been mature in the upper plant layers.

458   *   *   *

As a final step in this study, we developed an overall synthetic representation of the

relationship between the investments of *Z. tritici* in the two modes of reproduction at the host-

tissue scale, taking into account the ecological processes discussed above. In Figure 11, we

propose a theoretical relationship between asexual multiplication intensity (disease severity)

and sexual reproduction intensity (ability to produce offspring). The number of fruitful

encounters between parental isolates initially increases with increasing disease severity (1).

During this first stage, competition between isolates may lead to mating-type disequilibrium,

with an impact on sexual reproduction (2). At disease severities of more than 30-45%, a high-

density pathogen population becomes disadvantageous, due to competition for host resources

between the asexual and sexual modes of reproduction (3). This model is complementary to

the findings of Suffert et al. [23], formalized by the title "Fashionably late partners have more

fruitful encounters". We can now extend the metaphor by providing *Z. tritici* with additional

advice for even more fruitful encounters: "Do not live as a hermit but avoid crowded places!"

This view was already put forward by Kokko & Rankin [1], in their study entitled "Lonely

hearts or sex in the city?"

Is an optimal disease severity of 30-45% an appropriate ecological equilibrium determining

the evolutionary endpoints of selection in *Z. tritici*? This theoretical question needs to be

addressed in an evolutionary ecology perspective, and we can now use empirical data to try to

answer it. Indeed, a successful parasitic life depends on optimal exploitation of the host to

satisfy key functions directly involved in reproductive fitness. Theoretically, both weakly

aggressive isolates causing only a few small lesions, and very aggressive isolates causing

large numbers of large lesions, have a lower probability of transmitting a part of their genetic

background by sexual reproduction than isolates with intermediate levels of aggressiveness.

This view is complementary to the findings of Kema et al. [19], which explained the

maintenance of avirulent individuals in the pathogen population despite the widespread

deployment of resistance genes (e.g. AvrStb6 despite the extensive presence of Stb6 in the

wheat lines used in breeding programs worldwide [30]; Thierry Marcel, INRA BIOGER,

com. pers.). Sexual reproduction therefore plays a key role in both the evolution of virulence

and aggressiveness. We can hypothesize that sexual reproduction is less intense on very

susceptible wheat varieties than on less susceptible varieties, but this counter-intuitive

hypothesis requires testing in future experiments.

A previous study revealed an absence of functional trade-offs between the two modes of

reproduction in *Z. tritici*: no adaptive compromise was established between pathogenicity

(asexual multiplication on adult wheat plants) and transmission (intensity of sexual

reproduction on wheat debris [23]), in analyses of pathogen life traits at the individual level.

An overall epidemiological trade-off was, however, established between intra- and

interannual scales over a larger spatiotemporal scale (field and surrounding area), probably

driven by the consequences of sexual reproduction for local population dynamics relating to

selection and counter-selection [22]. According to the findings reported here, the potential

antagonism between the asexual and sexual modes of reproduction and its epidemiological

consequences appear to be more complex, due to a dependence on scale, from host tissues up

to agricultural landscape scale. Sexual reproduction is clearly a key process in the eco-

epidemiology of Septoria tritici blotch. An understanding of its determinants may open up

new perspectives for the management of other foliar fungal pathogens with dual modes of

reproduction.

References

1. Kokko H, Rankin DJ (2010) Lonely hearts or sex in the city? Density-dependent effects in

mating systems. Philosophical Transactions in the Royal Society B 361:319-334.

2. Lee SC, Ni M, Li W, Shertz C, and Heitman J (2010) The evolution of sex: a perspective

from the fungal kingdom. Microbiology and Molecular Biology Reviews 74:298-340.

3. Lively CM (2009) The maintenance of sex: host-parasite coevolution with density-

dependent virulence. Journal of Evolutionary Biology 22:2086-2093.

4. Gascoigne J, Berec L, Gregory S, Courchamp F (2009) Dangerously few liaisons: a review

of mate-finding Allee effects. Population Ecology 51:355-372.

5. Neiman M, Warren D, Rasmussen B, Zhang S (2013) Complex consequences of increased

density for reproductive output in an invasive freshwater snail. Evolutionary Ecology

27:1117-1127.

6. Stelzer CP (2012) Population regulation in sexual and asexual rotifers: an eco-evolutionary

feedback to population size? Functional Ecology 26:180-188.

7. Newton MR, Kinkel LL, Leonard KJ (1997) Competition and density-dependent fitness in

a plant parasitic fungus. Ecology 78:1774-1784.

8. Clément JAJ, Magalon H, Glais I, Jacquot E, Andrivon D (2012) To be or not to be

solitary: *Phytophthora infestans*' dilemma for optimizing its reproductive fitness in

multiple infections. PLoS ONE 7: e37838.

9. Gross A, Holdenrieder O, Pautasso M, Queloz V, Sieber TN (2014) *Hymenoscyphus*

pseudoalbidus, the causal agent of European ash dieback. Molecular Plant Pathology

15:5-21.

10. Halkett F, Coste D, Platero GG, Zapater MF, Abadie C, Carlier J (2010) Genetic

discontinuities and disequilibria in recently established populations of the plant

pathogenic fungus *Mycosphaerella fijiensis*. Molecular Ecology 19:3909-3923.

11. Hamelin FM, Castella F, Doli V, Marçais B, Ravigné V, Lewis MA (2016) Mate finding,

sexual spore production, and the spread of fungal plant parasites. Bulletin of

Mathematical Biology 78:695-712.

12. Ravigné V, Lemesle V, Walter A, Mailleret L, Hamelin L (2017) Mate limitation in

fungal plant parasites can lead to cyclic epidemics in perennial host populations.

Bulletin of Mathematical Biology 79:430-447.

13. Hamed BH, Gisi U (2013) Generation of pathogenic F1 progeny from crosses
of *Phytophthora infestans* isolates differing in ploidy. Plant Pathology 62:708-718.

14. Kema GHJ, Verstappen ECP, Todorova M, Waalwijk C (1996) Successful crosses and
molecular tetrad and progeny analyses demonstrate heterothallism in *Mycosphaerella*
graminicola. Current Genetics 30:251-258.

15. Waalwijk CO, Mendes O, Verstappen EC, De Waard MA, Kema GH (2002) Isolation and
characterization of the mating-type idiomorphs from the wheat septoria leaf blotch
fungus *Mycosphaerella graminicola*. Fungal Genetics and Biology 35:277-286.

16. Eriksen L, Munk L (2003) The occurrence of *Mycosphaerella graminicola* and its
anamorph Septoria tritici in winter wheat during the growing season. European Journal
of Plant Pathology 109:253-259.

17. Suffert F, Sache I, Lannou C (2011) Early stages of septoria tritici blotch epidemics of
winter wheat: Build-up, overseasoning, and release of primary inoculum. Plant
Pathology 60: 166-177.

18. Linde CC, Zhan J, McDonald BA (2002) Population structure of *Mycosphaerella*
graminicola: from lesions to continents. Phytopathology 92:946-955.

19. Kema GHJ, Mirzadi Gohari A, Aouini L, Gibriel HAY, Ware SB, van den Bosch F,
Manning-Smith R, Alonso-Chavez V, Helps J, Ben M'Barek S, Mehrabi R, Diaz-
Trujillo C, Zamani E, Schouten HJ, van der Lee TAJ, Waalwijk C, de Waard MA, de
Wit PJGM, Verstappen ECP, Thomma BPHJ, Meijer HJG, Seidl MF (2018) Stress and
sexual reproduction affect the dynamics of the wheat pathogen effector AvrStb6 and
strobilurin resistance. Nature Genetics, in press.

20. Metcalfe RJ (1998) Selection for resistance to demethylation inhibitor fungicides in

Mycosphaerella graminicola on wheat. PhD, The University of Reading.

21. Cowger C, McDonald BA, Mundt CC (2002) Frequency of sexual reproduction by

Mycosphaerella graminicola on partially resistant wheat cultivars. Phytopathology

92:1175-1181.

22. Suffert F, Goyeau H, Sache I, Carpentier F, Gélisse S, Morais D, Delestre G (2018)

Epidemiological trade-off between intra- and interannual scales in the evolution of

aggressiveness in a local plant pathogen population. Evolutionary Applications, in press

23. Suffert F, Delestre G, Carpentier F, Walker AS, Gazeau G, Gélisse S, Duplaix C (2016)

Fashionably late partners have more fruitful encounters: impact of the timing of co-

infection and pathogenicity on sexual reproduction in *Zymoseptoria tritici*. Fungal

Genetics and Biology 92:40-49.

24. Sánchez-Vallet A, McDonald MC, Solomon PS, McDonald BA (2015) Is *Zymoseptoria*

tritici a hemibiotroph? Fungal Genetics and Biology 79: 29-32.

25. Suffert F, Ravigné V, Sache I (2015) Seasonal changes drive short-term selection for

fitness traits in the wheat pathogen *Zymoseptoria tritici*. Applied and Environmental

Microbiology 81:6367-6379.

26. Zhan J, Kema GHJ, Waalwijk C, McDonald BA (2002) Distribution of mating type alleles

in the wheat pathogen *Mycosphaerella graminicola* over spatial scales from lesions to

continents. Fungal Genetics and Biology 36:128-136.

27. Pfender WF, Wootke SL (1988) Microbial communities of *Pyrenophora*-infested wheat

straw as examined by multivariate analysis. Microbial Ecology 15:95-113.

580 28. Zelikovitch N, Eyal Z (1991) Reduction in pycnidial coverage after inoculation of wheat

581   with mixtures of isolates of *Septoria tritici*. Plant Disease 75:907-910.

582 29. Eyal Z (1992) The response of field-inoculated wheat cultivars to mixtures of Septoria

583   tritici isolates. Euphytica 61:25-35.

584 30. Saintenac C, Lee W-S, Cambon F, Rudd JJ, King RC, Marande W, Powers SJ, Bergès H,

585   Phillips AL, Uauy C, Hammond-Kosack KE, Langin T, Kanyuka K (2018) Wheat

586   receptor-kinase-like protein Stb6 controls gene-for-gene resistance to fungal pathogen

587   *Zymoseptoria tritici*. Nature Genetics, in press.

Table 1. List of fitness traits and yield components assessed during the asexual and sexual stages for the five *Zymoseptoria tritici* crosses on adult wheat plants.

Year	Parental isolate	Mating type	Host cv.	Fitness traits			Yield components	
				Asexual stage		Sexual stage		
				Necrotic area	Density and exudation capacity of pycnidia	Ascospore production	GWS [1]	TKW [2]
2015	INRA09-FS0802 INRA09-FS1006	Mat1-1 Mat1-2	Soissons	×		×		
2016	INRA09-FS0813 INRA09-FS0732	Mat1-1 Mat1-2	Soissons	×	×	×	×	×
2016	INRA09-FS0808 INRA09-FS0806	Mat1-1 Mat1-2	Soissons	×	×	×	×	×
2017	INRA09-FS1022 INRA09-FS1008	Mat1-1 Mat1-2	Soissons	× [3]		×	× [3]	× [3]
2017	ST99CH-3D7 ST99CH-1A5	Mat1-1 Mat1-2	Runal	×		×	×	×

[1] grain weight per spike

[2] thousand-kernel weight

[3] also assessed on two sets of plants inoculated separately with each monoparental suspension

Fig.1 (a) Adult wheat plants cv. Soissons (three replicates per inoculum concentration) five
weeks after their inoculation in the greenhouse with 12 biparental suspensions of
Zymoseptoria tritici isolates (FS808 and FS806), at concentrations of 50 to 10^7 conidia.mL^{-1}.
Plants are ranked from the lowest (left) to the highest concentration (right). (b-c) Sporulating
area on the flag leaf, characterized by a high density of pycnidia, with high (b) and low (c)
exudation capacities. (d) Trios of plants stored outdoors to induce sexual reproduction and
ascosporogenesis.

602
603

Fig.2 Visual rating scale used to quantify the intensity of *Zymoseptoria tritici* asexual multiplication (disease severity assessed as a percentage of necrotic area) on the three uppermost leaves (F1, F2 and F3) of adult wheat plants. The images show the 2-7 cm section of the flag leaves over their point of insertion on the stem.

Fig.3 Relationship between the proportion of pseudothecia among the total fruiting bodies on
wheat leaves in field conditions and the mean severity of Septoria tritici blotch expressed as a
percentage of the leaf area covered by pycnidia or pseudothecia. A model (bold line)
integrating different density-dependent constraints (see eq. 3 in Material & Methods) was
fitted to some of the experimental data (◆) obtained by Eriksen et al. [16] (Table 1) in a two-
year field survey in Denmark. Data for leaves L8 to L11 (basal leaf layers) were excluded (×)
because no pseudothecia were found whatever the level of disease severity, probably due to
their advanced state of decomposition; data for leaves L5 to L7 (collected before mid-June),
L1 (flag leaf), and L2 (collected before August), were also excluded (+), because sexual
reproduction was probably incomplete (29-53 days between the appearance of the first
pycnidia and the appearance of the first pseudothecia [16]).

Fig.4 Impact of the vertical position of wheat tissues in standing stubble left in the field in the
fall (stems and leaves, from lower to upper layers: 0-5 cm, 5-10 cm, 10-15 cm, 15-25 cm, 25-
35 cm, 35-50 cm, 50-65 cm) on the intensity of *Zymoseptoria tritici* sexual reproduction
expressed as the number of ascospores collected per gram of debris (logarithmic scale) on
three or four dates (from September to January) over three interepidemic periods (2015-2016,
2016-2017, 2017-2018). The mean contribution of each stubble layer to sexual reproduction is
expressed as a percentage of the total number of ascospores collected during each
interepidemic period.

Fig.5 Relationship between *Zymoseptoria tritici* inoculum concentration (total number of
conidia per mL) and mean intensity of asexual multiplication (disease severity assessed as a
% of necrotic area) estimated on the three uppermost leaves of adult wheat plants. Inoculation
with biparental suspensions of isolates FS802 × FS1006 (2015), FS813 × FS732 (2016),
FS808 × FS806 (2016), and FS1022 × FS1008 (2017) were performed on cv. Soissons;
Inoculations with biparental suspensions of isolates 3D7 × 1A5 (2017) were performed on cv.
Runal.

Fig.6 Relationship between the mean intensity of asexual multiplication in *Zymoseptoria tritici* (disease severity assessed as a % of necrotic area) on the three uppermost leaves of adult wheat plants cv. Soissons (black symbols) or Runal (white symbol), and (a, c, e) total grain weight per spike (GWS) or (b, d, f) thousand-kernel weight (TKW). Disease severity was assessed five weeks after inoculation with monoparental or biparental suspensions of isolates FS806, FS808, FS732, and FS813 in 2016, and FS1022, FS1008, 1A5, and 3D7 in 2017. Vertical bars represent the standard deviation.

651
652

 Fig.7 Relationship between the mean severity of Septoria tritici blotch (percentage of necrotic
area) and (a) density of pycnidia (number of pycnidia per cm^2) or (b) percentage of pycnidia
displaying exudation. Assessments were performed five weeks after the inoculation of adult
wheat plants cv. Soissons with biparental suspensions of *Zymoseptoria tritici* isolates FS813 ×
FS732 (2016) and FS808 × FS806 (2016). Vertical bars represent the standard deviation.

659
660

Fig.8 Relationship between inoculum concentration of *Zymoseptoria tritici* FS1008 or
FS1022 isolate (total number of conidia per mL) in suspension alone or mixed with a second
isolate present in the same amount and the mean severity of Septoria tritici blotch (percentage
of necrotic area) estimated on the three uppermost leaves of adult wheat plants cv. Soissons.
The number indicates the difference (in %) between the severity of the disease induced by
FS1008 (the most aggressive isolate) inoculated alone and the severity of the disease induced
by the FS1008 × FS1022 mixture (for the same number of FS1008 conidia); a negative value
indicates that disease severity was lower after inoculation with a mixture.

Fig.9 Impact of inoculum concentration (total number of conidia per mL) on the intensity of
sexual reproduction in *Zymoseptoria tritici* (number of ascospores per g of wheat debris).
Inoculation with biparental suspensions of isolates (a) FS802 × FS1006 (2015), (b) FS813 ×
FS732 (2016), (c) FS808 × FS806 (2016), and (d) FS1022 × FS1008 (2017) were performed
on cv. Soissons, and with (e) 3D7 × 1A5 (2017) on cv. Runal. Each bar corresponds to the
cumulative number of ascospores collected per gram of wheat debris over four independent
discharge events (rep1 to rep4). The complete dataset is given in Table S1.

Fig.10 Empirical relationship between asexual multiplication intensity in *Zymoseptoria tritici*
(disease severity assessed as a percentage of necrotic area) and sexual reproduction intensity
(normalized number of ascospores for each cross expressed in %). A quadratic model (dotted
lines; see eq. 2 in Material & Methods) and a model integrating two density-dependent
constraints (bold lines; see eq. 3) were adjusted on the data for three crosses (FS802 × FS1006
in red, FS808 × FS806 in orange, FS1022 × FS1008 in yellow). Convergence was not
obtained for the other two crosses (white points).

686
687

Fig.11 Theoretical relationship between asexual multiplication intensity in *Zymoseptoria*
tritici (disease severity assessed as a percentage of necrotic area) and sexual reproduction
intensity (normalized number of ascospores produced per g of wheat tissue), assuming that
the conidia of two compatible parental isolates were deposited in equal proportions on leaves.
Grey areas represent the decrease in sexual reproduction intensity due to: (1) an Allee effect
(difficulty finding the opposite mating type at low population densities); (2) induced
resistance (activation of host defense mechanisms by one of the parental isolates, leading to a
mating-type disequilibrium); (3) competition for host resource allocation between asexual
multiplication and sexual reproduction. Sexual reproduction intensity is expected to be
optimal for disease severities of 30 to 45%.

FS802 × FS1006 on wheat cv. Soissons (2015)

Inoculum concentration [1]	Disease severity [2]	Mean number of ascospores collected per gram of debris					
Date	05-02-2015	11-12-2015	11-19-2015	12-03-2015	01-21-2016	-	Total
50	0.8 ± 0.3	103.2	72.5	7.9	26.6		210.2
10^2	2.5 ± 0.6	0.3	7.5	91.2	0.3		99.2
5×10^2	3.0 ± 0.5	13.5	105.6	71.5	3.7		194.3
10^3	4.8 ± 1.3	66.4	132.9	51.2	11.4		261.9
5×10^3	14.2 ± 2.2	265.0	184.6	518.7	60.9		1029.1
10^4	17.5 ± 3.5	159.2	543.2	236.3	24.6		963.3
5×10^4	29.2 ± 3.4	976.8	656.1	504.4	35.5		2172.7
10^5	38.3 ± 4.7	316.2	999.9	529.3	90.8		1936.1
5×10^5	61.7 ± 3.1	76.3	1225.2	433.8	150.0		1885.2
10^6	65.0 ± 4.6	98.7	1238.5	353.6	49.9		1740.7
5×10^6	79.2 ± 4.2	10.1	781.8	747.6	69.3		1608.8
10^7	77.5 ± 4.9	395.0	660.1	120.6	3.1		1178.7

703 [1] total number of conidia per mL
704 [2] percentage of necrotic area ± standard deviation

FS813 × FS732 on wheat cv. Soissons (2016)

Inoculum concentration [1]	Disease severity [2]	Mean number of ascospores collected per gram of debris						
Date	05-06-2016	11-17-2016	01-19-2017	02-15-2017	03-01-2017	03-15-2017	Total
50	1.1 ± 0.4	0.0	0.0	2.0	0.0	0.0	2.0
10^2	1.0 ± 0.5	0.0	0.0	0.0	4.5	0.0	4.5
5×10^2	0.9 ± 0.3	0.0	0.0	0.0	0.0	0.0	0.0
10^3	1.6 ± 0.4	0.0	0.0	0.0	0.0	0.0	0.0
5×10^3	9.3 ± 2.0	0.0	0.0	9.6	0.2	2.9	12.7
10^4	17.1 ± 2.6	0.0	0.0	10.6	10.0	21.2	41.8
5×10^4	33.3 ± 2.1	0.0	0.0	80.6	47.8	98.5	227.0
10^5	41.3 ± 3.8	0.0	0.0	138.8	266.9	409.0	814.7
5×10^5	57.2 ± 3.7	0.0	14.4	154.7	172.4	134.8	476.3
10^6	67.8 ± 1.7	0.0	0.7	467.9	135.4	755.9	1360.0
5×10^6	75.6 ± 1.5	0.0	0.0	72.2	108.1	189.6	369.9
10^7	78.3 ± 2.4	0.0	10.6	338.2	230.3	694.5	1273.7

705   [1] total number of conidia per mL

706   [2] percentage of necrotic area ± standard deviation

FS808 × FS806 on wheat cv. Soissons (2016)

Inoculum concentration [1]	Disease severity [2]	Mean number of ascospores collected per gram of debris						
Date	05-06-2016	11-17-2016	01-19-2017	02-15-2017	03-01-2017	03-15-2017	Total
50	1.6 ± 0.3	0.0	0.0	0.2	2.1	0.0	2.3
10^2	1.4 ± 0.3	0.0	0.0	0.0	0.0	0.0	0.0
5×10^2	2.8 ± 0.7	0.0	0.0	9.0	7.6	21.9	38.5
10^3	5.9 ± 1.5	0.0	4.1	2.6	21.5	36.9	65.0
5×10^3	7.3 ± 1.5	0.0	0.0	8.1	0.0	11.1	19.2
10^4	9.8 ± 2.6	0.0	0.0	6.9	0.0	4.2	11.2
5×10^4	28.9 ± 2.6	0.0	22.2	321.2	83.9	78.0	505.3
10^5	43.9 ± 3.1	0.0	42.6	147.7	42.3	247.6	480.2
5×10^5	60.0 ± 2.9	0.0	22.7	76.5	3.1	81.8	184.0
10^6	65.0 ± 3.3	0.0	0.0	56.7	60.2	32.8	149.7
5×10^6	75.6 ± 1.6	0.0	0.6	119.2	28.3	175.4	323.5
10^7	74.4 ± 2.2	0.0	8.2	13.6	25.5	4.0	51.2

707 [1] total number of conidia per mL

708 [2] percentage of necrotic area ± standard deviation

FS1022 × FS1008 on wheat cv. Soissons (2017)

Inoculum concentration [1]	Disease severity [2]	Mean number of ascospores collected per gram of debris						
Date	05-27-2017	11-09-2017	12-07-2017	01-04-2017	01-01-2018	01-24-2018	Total
50	3.2 ± 0.2	0.0	18.3	1.7	12.7	16.1	48.8
10^2	3.3 ± 0.2	0.0	38.8	64.6	75.6	16.8	195.7
5×10^2	5.7 ± 0.5	0.0	0.9	22.0	35.0	34.8	92.8
10^3	6.4 ± 0.7	0.0	12.5	79.2	96.6	104.8	293.1
5×10^3	5.9 ± 0.5	0.0	61.2	45.5	51.4	90.9	249.0
10^4	6.9 ± 1.2	0.0	2.2	0.4	54.2	77.0	133.8
5×10^4	22.2 ± 2.2	0.0	2.8	53.6	11.6	84.9	152.8
10^5	25.0 ± 2.4	0.0	0.1	129.5	176.1	145.6	451.3
5×10^5	56.9 ± 3.6	0.0	26.2	15.5	7.3	117.0	166.0
10^6	75.6 ± 5.0	0.0	60.6	381.1	254.4	247.5	943.6
5×10^6	94.4 ± 0.4	0.0	18.4	3.2	16.3	153.5	191.4
10^7	98.3	0.0	97.3	136.4	92.9	194.3	521.0

709   [1] total number of conidia per mL

710   [2] percentage of necrotic area ± standard deviation

3D7 × 1A5 on wheat cv. Runal (2017)

Inoculum concentration [1]	Disease severity [2]	Number of ascospores collected per gram of debris						
Date	05-27-2017	11-09-2017	12-07-2017	01-04-2017	01-01-2018	01-24-2018	Total
50	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10^2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5×10^2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10^3	0.0	0.0	0.0	0.0	0.0	2.2	2.2
5×10^3	0.3 ± 0.1	0.0	0.0	1.4	0.0	0.2	1.6
10^4	2.1 ± 0.3	0.0	97.6	117.2	98.0	150.2	463.0
5×10^4	4.1 ± 0.2	0.0	15.5	29.5	51.4	124.8	221.2
10^5	7.6 ± 0.3	0.0	19.2	81.0	79.5	106.2	286.0
5×10^5	21.7 ± 0.7	0.0	58.4	95.0	185.0	473.8	812.1
10^6	34.7 ± 2.8	0.0	0.0	217.4	85.8	420.5	723.7
5×10^6	80.6 ± 0.4	0.0	386.2	87.5	200.2	1181.4	1855.3
10^7	88.3 ± 1.8	0.0	73.2	19.6	4.2	201.5	298.5

[1] total number of conidia per mL

[2] percentage of necrotic area ± standard deviation

Online Resource 2. (a) Spatiotemporal dynamics of *Zymoseptoria tritici* sexual reproduction
in standing stubble left in the field during the fall, over three interepidemic periods (2015-
2016, 2016-2017, 2017-2018). (b, c, d) Impact of the vertical position of wheat tissues (stems
and leaves, from lower to upper layers: 0-5 cm, 5-10 cm, 10-15 cm, 15-25 cm, 25-35 cm, 35-
50 cm, 50-65 cm) on the number of ascospores collected per gram of debris (logarithmic
scale) on three or four sampling dates (from September to January) over the three
interepidemic periods. The dotted lines indicate the standard error.

722