The cellular NMD pathway restricts Zika virus infection and is targeted by the viral

capsid protein

Krystal A. Fontaine[a], Kristoffer E. Leon[a,b], Mir M. Khalid[a], Sakshi Tomar[a], David Jimenez-

Morales[a,c]*, Mariah Dunlap[a], Julia A. Kaye[a], Priya S. Shah[c]*, Steve Finkbeiner[a,d], Nevan

J. Krogan[a,c] and Melanie Ott[a,e]#

[a]Gladstone Institutes, San Francisco, CA

[b]Medical Scientist Training Program, University of California, San Francisco, CA

[c]Quantitative Biology Institute, Department of Cellular and Molecular Pharmacology,

University of California, San Francisco, CA

[d]Department of Neurology and Physiology, University of California, San Francisco, CA,

[e]Department of Medicine, University of California, San Francisco, CA

Running title: Zika virus capsid protein targets the NMD pathway

#Address correspondence to Melanie Ott, melanie.ott@gladstone.ucsf.edu.

*Present address: Priya S. Shah, Departments of Chemical Engineering and Microbiology

and Molecular Genetics, University of California, Davis, CA.

David Jimenez-Morales, PhD, Department of Medicine, Division of Cardiovascular

Medicine, Stanford University.

K.A.F and K.E.L contributed equally to this work.

Abstract word count: 198 | Text word count: 4,426

Abstract

Zika virus (ZIKV) infection of neural progenitor cells (NPCs) *in utero* is associated with

neurological disorders, such as microcephaly, but a detailed molecular understanding of

ZIKV-induced pathogenesis is lacking. Here we show that *in vitro* ZIKV infection of human

cells, including NPCs, causes disruption of the nonsense-mediated mRNA decay (NMD)

pathway. NMD is a cellular mRNA surveillance mechanism that is required for normal

brain size in mice. Using affinity purification-mass spectrometry, we identified multiple

cellular NMD factors that bind to the viral capsid protein, including the central NMD

regulator up-frameshift protein 1 (UPF1). Endogenous UPF1 interacted with the ZIKV

capsid protein in co-immunoprecipitation experiments, and capsid expression post-

transcriptionally downregulated UPF1 protein levels, a process that we confirmed occurs

during ZIKV infection. Cellular fractionation studies show that the ZIKV capsid protein

specifically targets nuclear UPF1 for degradation via the proteasome. A further decrease

in UPF1 levels by RNAi significantly enhanced ZIKV infection in NPC cultures, consistent

with a model in which NMD restricts ZIKV infection in the fetal brain. We propose that

ZIKV, via the capsid protein, has evolved a strategy to lower UPF1 levels and dampen

antiviral activities of NMD, which in turn contributes to neuropathology *in vivo.*

Importance

Zika virus (ZIKV) is a significant global health threat, as infection has been linked to

serious neurological complications, including microcephaly. Using a human stem cell-

derived neural progenitor model system, we find that a critical cellular quality control

process called the nonsense-mediated mRNA decay (NMD) pathway is disrupted during

ZIKV infection. Importantly, disruption of the NMD pathway is a known cause of

microcephaly and other neurological disorders. We further identify an interaction between

the capsid protein of ZIKV and up-frameshift protein 1 (UPF1), the master regulator of

NMD, and show that ZIKV capsid targets UPF1 for degradation. Together, these results

offer a new mechanism for how ZIKV infection can cause neuropathology in the

developing brain.

Introduction

ZIKV is a mosquito-borne RNA virus that belongs to the *Flaviviridae* family. First

isolated in Uganda in 1947, ZIKV remained relatively obscure for decades following its

discovery because infection was associated with only mild disease. However, more

severe clinical manifestations, including microcephaly, have been observed during the

recent spread of ZIKV through the Americas (1). ZIKV infection induces cell cycle arrest

and apoptosis in neural progenitor cells (NPCs) in *in vitro* studies and *in vivo* mouse

models, with the latter resulting in cortical thinning and microcephaly (2-6). While it is now

established that ZIKV infection during pregnancy is a causative agent of microcephaly

(7), the molecular mechanisms underlying ZIKV-induced neuropathogenesis remain

largely unknown.

Similar to other flaviviruses, ZIKV contains a single-stranded, positive-sense RNA

genome of ~11 kb in size. The genome encodes a single polyprotein that is post-

translationally processed by both host and viral proteases to produce 3 structural and 7

nonstructural proteins (8, 9). The flavivirus capsid, which is the first protein encoded in

the genome, is a major structural element required for the encapsidation of the RNA

genome during virion assembly (10). While flavivirus replication is known to occur in the

cytoplasm, a significant portion of the viral capsid protein localizes to the nucleus during

infection (10, 11). Although the role of nuclear capsid during infection is less clear, several

functions have been suggested. The capsid protein from dengue virus, a close relative of

ZIKV, binds to core histones and inhibits nucleosome formation, thus implicating the

protein in altering host gene expression (12). Furthermore, several flavivirus capsid

proteins, including ZIKV capsid, localize to the nucleolus, with many interacting with

nucleolar proteins to promote viral particle production (13-16).

The nonsense-mediated mRNA decay (NMD) pathway was initially discovered as

a highly conserved quality control system that destroys transcripts containing premature

termination codons (PTCs)(17). Following splicing of pre-mRNAs, a multi-subunit protein

complex called the exon-junction complex (EJC) is deposited onto mRNAs near the sites

of exon-exon junctions. If a PTC is found ~50–55 nucleotides upstream of an EJC, the

mRNA will be subjected to NMD-mediated degradation initiated by the recruitment of the

RNA helicase up-frameshift protein 1 (UPF1). UPF1 plays a central role in the NMD

pathway by linking the translation termination event to the assembly of a surveillance

complex, resulting in NMD activation (18). Interestingly, microcephaly has been

associated with genetic mutations that result in the impairment of the NMD pathway.

While knockout of Upf1 and other NMD factors is embryonic lethal in mice (19), mice

haploinsufficient for the EJC components Magoh, Rbm8a, and Eif4a3 exhibit aberrant

neurogenesis and microcephaly (20-22).

In addition to PTC-containing transcripts, it is now known that the NMD pathway

recognizes a broader range of RNA substrates. Notably, the NMD controls the "normal"

expression of ~10% of the cellular transcriptome and is regarded as a post-transcriptional

mechanism of gene regulation (23). Furthermore, the NMD pathway also regulates viral

infections. While it was first reported that UPF1 promotes the infectivity of HIV-1 progeny

virions (24), replication of several human RNA viruses, including human T-cell

lymphotropic virus type 1 (HTLV-1), Semliki Forest virus and Sindbis virus, is enhanced

following UPF1 knockdown, implicating UPF1 and the NMD pathway, either directly or

indirectly, in the host antiviral response (25-28). As ZIKV infection and NMD impairment

both promote microcephaly development, and we previously described disruption of the

NMD pathway in cells infected with a related flavivirus, the hepatitis C virus (HCV) (29),

we hypothesized that ZIKV infection manipulates the cellular NMD pathway, a process

contributing to ZIKV-induced neuropathology.

Results

The NMD pathway is impaired during ZIKV infection. To determine if ZIKV infection

affects NMD, we infected human hepatoma cells (Huh7) and human induced pluripotent

stem cell (iPSC)-derived NPCs with ZIKV for 48 h. We isolated total RNA from infected

cells and measured mRNA levels of three canonical NMD substrates: asparagine

synthetase (ASNS), cysteinyl-tRNA synthetase (CARS), and SR protein SC35 (29).

ASNS, CARS, and SC35 transcripts were significantly elevated in Huh7 cells and NPCs

following infection with Asian lineage ZIKV strain P6-740 (Fig. 1a). Levels of NMD

substrates were also elevated in Huh7 cells infected with the contemporary ZIKV clinical

isolate PRVABC59 (Puerto Rico, 2015) (Fig. 1a). We found that the ZIKV-induced

increase in NMD transcripts did not reflect a global increase in transcription, as mRNA

levels of housekeeping genes, including glyceraldehyde 3-phosphate dehydrogenase

(GAPDH), were not altered in infected cells (Fig. 1a). Together, these results indicate that

the NMD pathway is impaired in ZIKV-infected cells.

NMD substrates are regulated through the ATP-dependent RNA helicase activity

of UPF1, the central regulator of NMD (18). To determine if ZIKV infection broadly affects

NMD, we utilized two publicly available RNA sequencing (RNA-Seq) datasets to compare

genome-wide transcriptional alterations found during ZIKV infection (6) to those found

following UPF1 knockdown (30). As shown in Figure 1b, there is a significant overlap in

upregulated genes between these two datasets. Interestingly, several of the overlapping

genes are canonical NMD substrates (31-35) involved in cell cycle arrest and induction

of apoptosis, two conditions linked to ZIKV-associated neuropathology (5). These genes

include DNA damage-inducible transcript 3 (DDIT3) (36) and growth arrest and DNA

damage-inducible protein 45 alpha and beta (GADD45A and GADD45B, respectively)

(37). Via quantitative real-time RT-PCR, we confirmed that transcripts of each were

upregulated following infection of Huh7 cells with ZIKV PRVABC59, while the mRNA

levels of the housekeeping genes GAPDH, hypoxanthine phosphoribosyltransferase 1

(HPRT1), and lactate dehydrogenase A (LDHA) were not elevated (Fig. 1c). Combined,

these data show that ZIKV infection is associated with dysregulated expression of NMD

substrates relevant to ZIKV-mediated neuropathogenesis.

ZIKV capsid interacts with the NMD pathway. We previously showed that the core

protein of HCV, as well as the capsid protein of the related flaviviruses dengue virus and

West Nile virus, interact with the host protein within bgcn homolog (WIBG/PYM1), an EJC

disassembly factor associated with NMD (29, 38). To examine potential interactions

between ZIKV and the NMD pathway, we separately analyzed data generated from an

affinity purification-mass spectrometry (AP-MS) screen to specifically query whether the

capsid protein of ZIKV interacts with NMD-associated host factors (Shah et al.,

submitted). ZIKV-host protein-protein interaction (PPI) maps were generated in HEK293T

cells using ZIKV proteins from the Ugandan 1947 strain MR 766 or the French Polynesian

2013 strain H/PF/2013 as bait proteins.

From this analysis, we found that ZIKV capsid proteins interacted with several

components of the NMD pathway, including multiple members of the EJC complex, as

well as UPF1 and UPF3B, the latter an NMD effector that stimulates UPF1 helicase

activity (Fig. 2a) (39). Importantly, the NMD host factors that interacted with each of the

two different capsid proteins greatly overlapped, revealing that the interaction between

capsid and the NMD pathway is conserved across the Asian and African lineages of ZIKV

(Fig. 2a). Next, we validated the binding of ZIKV capsid to select NMD host factors by co-

immunoprecipitating FLAG-tagged capsid protein with endogenous UPF3B or UPF1 in

HEK293T cells. Both UPF3B and UPF1 proteins co-immunoprecipitated with ZIKV

capsid, thus confirming the AP-MS results (Fig. 2b,c, respectively). UPF1 interacted with

the viral capsid protein independently from its RNA-binding and ATPase/helicase

capacities, as ZIKV capsid co-immunoprecipitated with UPF1 mutants deficient in these

functions following overexpression (Fig 2d).

ZIKV capsid degrades UPF1, the master regulator of NMD. Surprisingly, we

consistently observed a decrease in UPF1, but not UPF3B, protein levels in the input

lysate of ZIKV capsid-transfected cells, pointing to a specific perturbation of endogenous

UPF1 expression by ZIKV capsid (Fig. 2c). To confirm that UPF1 protein levels are

dysregulated during ZIKV infection, we performed western blot analysis of infected Huh7

cells and NPCs. Cellular UPF1 protein levels were consistently downregulated by ~50%

in ZIKV-infected Huh7 cells, whereas a ~25% reduction was observed in ZIKV-infected

NPCs (Fig. 2e), mirroring the difference in infection efficiencies achieved in these two cell

systems. UPF1 transcript levels were not decreased in ZIKV-infected cells or following

ZIKV capsid overexpression, indicating that UPF1 is post-transcriptionally downregulated

during ZIKV infection (Supplemental Fig. 1a,b, respectively).

Because ZIKV capsid and UPF1 both localize to the nucleus and the cytoplasm

(13, 40) we performed fractionation studies in ZIKV capsid-transfected HEK293T cells to

determine if UPF1 is downregulated within a specific cellular compartment. Capsid

expression markedly decreased nuclear UPF1 levels, whereas cytoplasmic levels were

unchanged (Fig. 3a). We next examined a potential role for the autophagic and

proteasomal pathways, both of which are known to mediate nuclear protein degradation,

in ZIKV capsid-induced UPF1 downregulation. As shown in Supplemental Figure 2,

nuclear UPF1 levels in ZIKV capsid-transfected cells were not rescued by inhibition of

cellular autophagy via bafilomycin A1 treatment. However, nuclear UPF1 levels were

restored in a dose-dependent manner following treatment with the proteasome inhibitor

bortezomib (Fig. 3b), indicating enhanced proteasomal degradation of nuclear UPF1 in

the presence of ZIKV capsid. Although ZIKV capsid colocalized with endogenous UPF1

in the cytoplasm of transfected Huh7-Lunet cells (Mander's colocalization coefficient of

~57%), we detected very little colocalization within the nucleus (~7%) (Supplemental

Figure 3). We hypothesized that this was due to specific degradation of UPF1 by ZIKV

capsid in the nucleus. Indeed, when cells were treated with bortezemib, the fraction of

nuclear UPF1 colocalizing with ZIKV capsid increased ~8-fold, while the fraction of

nuclear capsid interacting with UPF1 remained unchanged (Figure 3c). These results

demonstrate that ZIKV capsid interacts with UPF1 both in the cytoplasm and nucleus, but

specifically targets nuclear UPF1 for proteasomal degradation.

UPF1 is a restriction factor of ZIKV. To test the effect of lowered UPF1 levels on ZIKV

infection, we further decreased UPF1 expression prior to ZIKV infection by transfecting

NPCs with either non-targeting siRNA or a pool of UPF1-specific siRNAs. We then

infected the transfected cells with ZIKV and measured viral RNA levels, as well as

infectious titers, 48 h post-infection (hpi). UPF1 knockdown was successful in siRNA-

treated cells, as confirmed by western blot analysis (Fig. 4a). The depletion of UPF1 in

NPCs prior to infection resulted in a significant increase in both ZIKV RNA levels and

infectious virus production (Fig. 4b,c, respectively), indicating that expression of UPF1

restricts ZIKV infection at or before the RNA replication stage. To differentiate between

these two stages, we analyzed double-stranded RNA (dsRNA) intermediates

representing presumed viral RNA replication centers in infected NPCs (Fig. 4d)(41).

Using confocal microscopy and 3D reconstruction analyses, we observed no significant

difference in the number and size of dsRNA foci per cell when comparing ZIKV-infected,

UPF1-depleted NPCs to ZIKV-infected cultures expressing UPF1 (Fig. 4e,f). Instead, we

found a significant increase in the number of infected cells in NPC cultures when UPF1

was depleted, indicating that UPF1 regulates permissivity of NPCs to ZIKV infection at an

early stage prior to viral RNA replication (Fig. 4g).

Discussion

In summary, we identified the NMD pathway as a restriction mechanism for ZIKV

infection in human NPCs. NMD was partially inactivated in ZIKV-infected NPCs through

expression of the viral capsid protein and the resulting degradation of host nuclear UPF1.

As further weakening NMD by depleting UPF1 resulted in a marked increase in the

number of infected cells, we propose a model in which an evolutionary "arms race"

between cellular NMD and ZIKV determines whether a cell is successfully infected

(Fig.4h).

Downregulation of UPF1 by ZIKV capsid is not complete, and is likely limited by

the damaging effects of NMD disruption, as illustrated by the upregulation of genes

involved in cell growth arrest and apoptosis. Indeed, knockout of UPF1 and other

members of the NMD pathway is embryonic lethal in mice (19). However, mice

haploinsufficient for NMD factors upstream of UPF1, including Magoh, Rbm8a, and

Eif4a3, develop microcephaly (20-22). Thus, the reduction in nuclear UPF1 we observe

in ZIKV-infected NPCs could contribute to the microcephaly phenotype caused by ZIKV

infection in the fetal brain. While fetal and adult NPCs appear to be transcriptionally

distinct (42), it has been shown that adult NPCs are also permissive to ZIKV infection

(43). As the NMD pathway is a ubiquitous cellular surveillance mechanism, it is likely that

ZIKV capsid targets UPF1 for degradation in any cell type that is susceptible to ZIKV

infection. Accordingly, we have found that UPF1 is degraded following infection of both

NPCs and hepatic Huh7 cells.

Why ZIKV capsid specifically downregulates nuclear UPF1, and how nuclear UPF1

contributes to ZIKV restriction remain unanswered. Several studies suggest that NMD is

associated with the nucleus, although this issue remains controversial. Multiple

transcripts, such as those encoding T cell receptor beta, triosephosphate isomerase and

mouse major urinary protein, have been shown to be specifically degraded in purified

nuclei or reduced in nuclear fractions (44). These data support the model that selectively

depleting nuclear UPF1 levels disrupts NMD function in ZIKV-infected cells. In addition,

UPF1 is involved in several other processes within the nucleus, including nuclear-

associated RNA metabolism, cell cycle progression, and DNA replication (40). Therefore,

by targeting nuclear UPF1, ZIKV could disrupt these processes and program target cells

for viral replication. Notably, viral RNA replication is thought to occur solely within the

cytoplasmic compartment (45, 46). Using confocal microscopy and 3D reconstruction, we

did not detect dsRNA foci localized within the nuclei of ZIKV-infected cells (data not

shown), supporting our finding that UPF1 does not restrict viral RNA replication. While

our results suggest a role for nuclear UPF1 in ZIKV restriction, it is possible that UPF1

also serves as a restriction factor of ZIKV within the cytoplasm. Previously, it was shown

that UPF1 suppresses alphavirus replication by degrading the incoming viral RNA

following uncoating in the cytosol (28). Thus, ZIKV may possess an additional mechanism

to prevent cytoplasmic UPF1 from targeting its incoming RNA genome for destruction.

Our data reveal that nuclear UPF1 is degraded by ZIKV capsid in a proteasome-

dependent manner. While the nuclear proteasome has not been specifically linked to

microcephaly, it plays critical roles in the regulation of chromatin structure, gene

expression, DNA repair, and protein quality control (47). Thus, the co-opting of the nuclear

proteasome by ZIKV capsid to degrade UPF1 could disrupt its normal proteasomal

activity and further contribute to the cytopathic effects associated with ZIKV infection.

Furthermore, given that the capsid protein of the closely related dengue virus can

translocate across cell membranes (48), it is possible that capsid released from apoptotic,

ZIKV-infected cells can enter neighboring, uninfected cells to degrade UPF1, thus

increasing permissivity of bystander cells to ZIKV infection. Studies are ongoing to

determine the precise molecular mechanism of ZIKV capsid-mediated UPF1 degradation

and how UPF1 depletion enhances ZIKV replication, directly or indirectly. Ultimately,

these data may help inform new therapeutic approaches, as reinforcement of the antiviral

properties of the NMD pathway is expected to enhance resistance of NPCs to ZIKV

infection and to promote normal neurodevelopment in infected fetuses.

Materials and Methods

Viruses and cells. Two Asian lineage strains of ZIKV, P6-740 (ATCC VR-1845) and

PRVABC59 (ATCC VR-1843), were used for all experiments. ZIKV stocks were

propagated in Vero cells (ATCC) and titers were determined by plaque assays on Vero

cells. Huh7 cells (ATCC), Huh7-Lunet cells (Ralf Bartenschlager, Heidelberg University)

and Vero cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) with

10% fetal bovine serum (FBS), 2 mM L-glutamine, 100 U/mL penicillin, and 100 µg/mL

streptomycin. HEK293T cells (ATCC) were maintained in DMEM/H21 medium

supplemented with 10% FBS, 100 U/mL penicillin, 100 µg/mL streptomycin, and 1 mM

sodium pyruvate or DMEM with 10% FBS, 2 mM L-glutamine, 100 U/mL penicillin, and

100 µg/mL streptomycin. Human iPSC-derived NPCs were generated and maintained

as described previously (49). All of the human fibroblast cell lines used to generate iPSCs

came from the Coriell Institute for Medical Research and Yale Stem Cell Center. The

iPSCs used in these studies were the CTRL2493nXX, CS2518nXX, and Cs71iCTR-

20nXX lines. CTRL2493nXX was derived from the parental fibroblast line ND31845 that

was biopsied from a healthy female at 71 years of age. CS2518nXX was derived from the

parental fibroblast line ND30625 that was biopsied from a healthy male at 76 years of

age. CS71iCTR-20nXX was derived from the parental fibroblast line ND29971 that was

biopsied from a female at 61 years of age. For virus infections, NPCs plated on Matrigel-

coated (Corning) multi-well plates or Huh7 cells were infected with ZIKV at a multiplicity

of infection (MOI) of 0.1 or 1 for 2 h at 37°C. Infected cells were harvested at 48 hpi for

all analyses.

Affinity purification, mass spectrometry, and AP-MS scoring. The ZIKV capsid open

reading frames (ORFs) from the Ugandan 1947 strain MR 766 or the French Polynesian

2013 strain H/PF/2013 were cloned into pCDNA4_TO with a C-terminal 2xStrep II affinity

tag for expression in human cells. The viral capsid proteins (three biological replicates),

as well as GFP (two biological replicates) and empty vector (ten biological replicates) as

negative controls, were expressed in HEK293T cells and affinity purifications were

performed as previously described (50). Briefly, clarified lysates were incubated with

Strep-Tactin Superflow (IBA) overnight at 4°C. Proteins were eluted with 50 mM Tris pH

7.5, 150 mM NaCl, 1 mM EDTA containing 2.5 mM Desthiobiotin (IBA) for 30 min at 4°C.

Lysates and affinity purified eluates were analyzed by western blot and silver stain PAGE

to confirm expression and purification. Purified protein eluates were digested with trypsin

for LC-MS/MS analysis. Samples were denatured and reduced in 2M urea, 10 mM

NH_4HCO_3, 2 mM DTT for 30 min at 60°C, then alkylated with 2 mM iodoacetamide for 45

min at room temperature. Trypsin (Promega) was added at a 1:100 enzyme:substrate

ratio and digested overnight at 37°C. Following digestion, samples were concentrated

using C18 ZipTips (Millipore) according to the manufacturer's specifications. Peptides

were resuspended in 15 µL of 4% formic acid and 3% ACN, and 1-2 µL of sample was

loaded onto a 75 µm ID column packed with 25 cm of Reprosil C18 1.9 µm, 120Å particles

(Dr. Maisch GmbH). Peptides were eluted into a Q-Exactive Plus (Thermo Fisher) mass

spectrometer by gradient elution delivered by an Easy1200 nLC system (Thermo Fisher).

The gradient was from 4.5% to 32% acetonitrile over 53 min. All MS spectra were

collected with oribitrap detection, while the 20 most abundant ions were fragmented by

higher energy collisional dissociation (HCD) and detected in the orbitrap. All data was

searched against the SwissProt Human protein sequences, combined with ZIKV

sequences and GFP. Peptide and protein identification searches, as well as label-free

quantitation, were performed using the MaxQuant data analysis algorithm and all peptide

and protein identifications were filtered to a 1% false-discovery rate (51, 52). SAINTq (53)

was used to calculate the probability of bait-prey interactions for both Ugandan ZIKV

capsid and French Polynesian ZIKV capsid against the negative controls, including GFP

and empty vector, with protein intensities as input values. We applied a combined

threshold of probability of interaction (AvgP) greater than 0.90 and Bayesian False

Discovery Rate of less than 0.05.

Quantitative real-time reverse transcription-PCR (qRT-PCR). Total cellular RNA was

isolated from Huh7 cells and NPCs using the RNeasy Mini Kit (Qiagen). cDNA was

synthesized with oligo(dT)$_{18}$ (ThermoFisher Scientific) primers, random hexamer (Life

Technologies) primers, and AMV reverse transcriptase (Promega). The cDNA was then

used in SYBR Green PCR Master Mix (ThermoFisher Scientific) according to

manufacturer's instructions and analyzed by qPCR (Bio-Rad ABI 7900). The primers used

for ASNS, CARS, SC35 1.7, GAPDH, HPRT1, LDHA, and 18S rRNA have been

described previously (29). The additional primers used were ZIKV PRVABC59 forward

primer 5'- GAG ACG AGA TGC GGT ACA GG -3', ZIKV PRVABC59 reverse primer 5'-

CGA CCG TCA GTT GAA CTC CA -3', UPF1 forward primer 5'- CTG CAA CGG ACG

TGG AAA TAC -3', UPF1 reverse primer 5'- ACA GCC GCA GTT GTA GCA C -3', DDIT3

forward primer 5'- CTG CTT CTC TGG CTT GGC TG -3', DDIT3 reverse primer 5'- GCT

CTG GGA GGT GCT TGT GA -3', GADD45A forward primer 5'- GAG CTC CTG CTC

TTG GAG AC -3', GADD45A reverse primer 5'- GCA GGA TCC TTC CAT TGA GA -3',

GADD45B forward primer 5'- TGA CAA CGA CAT CAA CAT C -3', and GADD45B reverse

primer 5'- GTG ACC AGA GAC AAT GCA G -3'. Relative levels of each transcript were

normalized by the delta threshold cycle method to the abundance of 18S rRNA or

GAPDH, with mock-infected cells or vector-transfected cells set to 1.

Western blot analysis. Cells were lysed in RIPA lysis buffer (50 mM Tris-HCl, pH 8, 150

mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, supplemented with Halt™

protease inhibitor cocktail (ThermoFisher Scientific)) to obtain whole cell lysates or lysed

using the NE-PER nuclear and cytoplasmic extraction kit (ThermoFisher Scientific) to

obtain cytoplasmic and nuclear fractions. Proteins were separated by SDS-PAGE and

transferred to nitrocellulose membranes (Bio-Rad). Blots were incubated with the

indicated primary antibody: anti-UPF3B (ab134566, Abcam), anti-UPF1 (12040, Cell

Signaling Technology, Inc.), anti-ZIKV Capsid (C) (GTX133304, GeneTex), anti-Flag

(F7425, Sigma-Aldrich), anti-β-actin (A5316, Sigma-Aldrich), anti-ZIKV Envelope (E)

(GTX133314, GeneTex), anti-SP1 (sc-14027, Santa Cruz Biotechnology), anti-GAPDH

(5174, Cell Signaling Technology, Inc.), anti-Myc tag (ab9106, Abcam), anti-Strep tag

(ab18422, Abcam), and anti-p62 (ab56416, Abcam). Proteins were visualized by

chemiluminescent detection with ECL and ECL Hyperfilm (Amersham). Differences in

band intensity were quantified by densitometry using ImageJ.

Immunoprecipitations. Cells were lysed in either RIPA lysis buffer or IP lysis buffer

(150mM NaCl, 50mM Tris pH 7.4, 1mM EDTA, 0.5% NP-40 substitute, supplemented

with Halt™ protease inhibitor cocktail (ThermoFisher Scientific) at 4°C and passed

through a G23 needle. Clarified lysates were immunoprecipitated with Flag M2 agarose

(Sigma), anti-Myc tag (ab9106, Abcam), or normal rabbit IgG (sc-2027, Santa Cruz

Biotechnology) overnight, washed in lysis buffer, and resuspended in Laemmli buffer for

SDS-PAGE. Western blot analysis of immunoprecipitated proteins was performed as

described above.

Immunofluorescence. Transfected Huh7-Lunet cells or infected NPCs were collected at

48 h and plated onto 22 × 22 mm #1.5 coverslips. Cells were fixed in 4%

paraformaldehyde, permeabilized with 0.1% Triton X-100, and blocked in 3% bovine

serum albumin. Cells were then immunostained with the indicated antibodies: anti-Strep

Tag (Abcam, ab184224), anti-UPF1 (Abcam, ab109363), human anti-DENV mAb 1.6D

(Sharon Isern and Scott Michael, Florida Gulf Coast University), which recognizes the

ZIKV envelope protein, anti-dsRNA mAb J2 (SCICONS), and the appropriate fluorophore-

conjugated secondary antibodies. Coverslips were mounted onto glass slides using

Vectashield® Mounting Medium with DAPI (Vector Laboratories) and analyzed by

fluorescence microscopy (Zeiss Axio Observer ZI) or confocal microscopy (Zeiss LSM

880). For acquiring high-resolution images, cells were imaged on the Zeiss LSM 880 with

Airyscan using a 20x/0.8 or 63x/1.4 M27 oil immersion objective. A total of 15-20 (20x

objective) or 60-80 (63x objective) Z-slices were acquired every 0.88 μm or 0.15 μm,

respectively. The resulting Z-stack was reconstructed and rendered in 3D using Imaris

software (Bitplane). Viral dsRNA foci were reconstructed via the Imaris spot detection

function, which provided an analysis of total number and mean volume of foci within a

cell, for images acquired using the 20x objective. Strep-tagged ZIKV capsid, UPF1, and

dsRNA channels acquired using the 63x objective were reconstructed using the Imaris

surfaces package. The Imaris colocalization function was used to determine overlap of

fluorescence. Thresholding for background fluorescence was determined by the Imaris

automatic thresholding tool that utilizes the Costes approach (54). The thresholded

Mander's correlation coefficient (MCC) measures the fraction of voxels with fluorescence

positive for one channel that also contain fluorescence from another channel. The MCC

is typically more appropriate for analysis of three-dimensional colocalization (55).

Statistical analysis. Statistical differences between groups were analyzed using either

a two-tailed unpaired Student's *t*-test or a two-tailed ratio paired Student's *t*-test, as stated

in the figure legends. Hypergeometrical tests were used to calculate the probability of an

overlap in gene dysregulation between ZIKV-infected NPCs and UPF1-depleted cells and

to calculate the probability of ZIKV capsid bait-prey interactions. Data are represented as

mean ± s.e.m. Statistical significance was defined as *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$,

and ****$P \leq 0.0001$.

Acknowledgements. The authors would like to thank all members of the Ott laboratory,

as well as Roman Camarda, Marius Walter, and Anna Maurer for helpful discussions and

advice throughout the preparation of this manuscript. We thank Chia-Lin Tsou, the

Gladstone Stem Cell Core, and Meredith Calvert from the Gladstone Microscopy Core for

technical assistance, and Ralf Bartenschlager (Heidelberg University), Lynne Maquat

(University of Rochester), and Sharon Isern and Scott Michael (Florida Gulf Coast

University) for reagents. We are grateful to Veronica Fonseca for administrative support,

John Carroll and Teresa Roberts for graphical design, and to Kathryn Claiborn, Eric

Martens, and Gary Howard for editorial assistance. This work was supported by

NIH/NIAID F32AI112262 to P.S.S., NIH/NINDS R01 NS101996-01 to S.F., NIH/NIAID

U19AI1186101 to N.J.K., DOD/DARPA HR0011-11-C-0094 (PROPHECY) to N.J.K.,

NIH/NIAID R01 AI097552 to M.O., and the James B. Pendleton Charitable Trust.

References

1. Fauci AS, Morens DM. 2016. Zika Virus in the Americas--Yet Another Arbovirus Threat.

N Engl J Med 374:601-604. 10.1056/NEJMp1600297.

2. Cugola FR, Fernandes IR, Russo FB, Freitas BC, Dias JLM, Guimarães KP, Benazzato

C, Almeida N, Pignatari GC, Romero S, Polonio CM, Cunha I, Freitas CL, Brandão WN,

Rossato C, Andrade DG, Faria DdP, Garcez AT, Buchpigel CA, Braconi CT, Mendes E,

Sall AA, Zanotto, Paolo M. de A., Peron JPS, Muotri AR, Beltrão-Braga PCB. 2016. The

Brazilian Zika virus strain causes birth defects in experimental models. Nature 534:267-

271. 10.1038/nature18296.

3. Shao Q, Herrlinger S, Yang S, Lai F, Moore JM, Brindley MA, Chen J. 2016. Zika virus

infection disrupts neurovascular development and results in postnatal microcephaly with

brain damage. Development 143:4127-4136. 10.1242/dev.143768.

4. Souza BSF, Sampaio GLA, Pereira CS, Campos GS, Sardi SI, Freitas LAR, Figueira

CP, Paredes BD, Nonaka CKV, Azevedo CM, Rocha VPC, Bandeira AC, Mendez-Otero

R, Dos Santos RR, Soares MBP. 2016. Zika virus infection induces mitosis abnormalities

and apoptotic cell death of human neural progenitor cells. Scientific reports 6:39775.

10.1038/srep39775.

5. Li C, Xu D, Xu Z, Ye Q, Hong S, Jiang Y, Liu X, Zhang N, Shi L, Qin C. 2016. Zika

Virus Disrupts Neural Progenitor Development and Leads to Microcephaly in Mice. Cell

Stem Cell 19:120-126. 10.1016/j.stem.2016.04.017.

6. Tang H, Hammack C, Ogden S, Wen Z, Qian X, Li Y, Yao B, Shin J, Zhang F, Lee E,

Christian K, Didier R, Jin P, Song H, Ming G. 2016. Zika Virus Infects Human Cortical

Neural Progenitors and Attenuates Their Growth. Cell Stem Cell 18:587-590.

10.1016/j.stem.2016.02.016.

7. Sonja A Rasmussen, Denise J Jamieson, Margaret A Honein, Lyle R Petersen. 2016.

Zika Virus and Birth Defects--Reviewing the Evidence for Causality. The New England

journal of medicine 374:1981-1987. 10.1056/NEJMsr1604338.

8. Harris E, Holden KL, Edgil D, Polacek C, Clyde K. 2006. Molecular biology of

flaviviruses. Novartis Found Symp 277:253.

9. Lindenbach BD, Rice CM. 2003. Molecular biology of flaviviruses. Adv Virus Res 59:23-

61.

10. Oliveira ERA, Mohana-Borges R, de Alencastro RB, Horta BAC. 2017. The flavivirus

capsid protein: Structure, function and perspectives towards drug design. Virus Res

227:115-123. 10.1016/j.virusres.2016.10.005.

11. den Boon JA, Diaz A, Ahlquist P. 2010. Cytoplasmic Viral Replication Complexes.

Cell Host Microbe 8:77-85. 10.1016/j.chom.2010.06.010.

12. Colpitts TM, Barthel S, Wang P, Fikrig E. 2011. Dengue virus capsid protein binds

core histones and inhibits nucleosome formation in human liver cells. PLoS ONE

6:e24365. 10.1371/journal.pone.0024365.

13. Lukasz P Slomnicki, Dong-Hoon Chung, Austin Parker, Taylor Hermann, Nolan L

Boyd, Michal Hetman. 2017. Ribosomal stress and Tp53-mediated neuronal apoptosis in

response to capsid protein of the Zika virus. Scientific Reports (Nature Publisher Group)

7:1-15. 10.1038/s41598-017-16952-8.

14. Tsuda Y, Mori Y, Abe T, Yamashita T, Okamoto T, Ichimura T, Moriishi K, Matsuura

Y. 2006. Nucleolar protein B23 interacts with Japanese encephalitis virus core protein

and participates in viral replication. Microbiol Immunol 50:225-234.

15. Xu Z, Hobman TC. 2012. The helicase activity of DDX56 is required for its role in

assembly of infectious West Nile virus particles. Virology 433:226-235.

10.1016/j.virol.2012.08.011.

16. Rawlinson SM, Moseley GW. 2015. The nucleolar interface of RNA viruses. Cell

Microbiol 17:1108-1120. 10.1111/cmi.12465.

17. Peccarelli M, Kebaara BW. 2014. Regulation of natural mRNAs by the nonsense-

mediated mRNA decay pathway. Eukaryotic cell 13:1126-1135. 10.1128/EC.00090-14.

18. Hug N, Longman D, Cáceres JF. 2016. Mechanism and regulation of the nonsense-

mediated decay pathway. Nucleic Acids Res 44:1483-1495. 10.1093/nar/gkw010.

19. Han X, Wei Y, Wang H, Wang F, Ju Z, Li T. 2018. Nonsense-mediated mRNA decay:

a 'nonsense' pathway makes sense in stem cell biology. Nucleic acids research 46:1038-

1051. 10.1093/nar/gkx1272.

20. Silver DL, Watkins-Chow DE, Schreck KC, Pierfelice TJ, Larson DM, Burnetti AJ, Liaw

H, Myung K, Walsh CA, Gaiano N, Pavan WJ. 2010. The exon junction complex

component Magoh controls brain size by regulating neural stem cell division. Nat

Neurosci 13:551-558. 10.1038/nn.2527.

21. Mao H, Pilaz L, McMahon JJ, Golzio C, Wu D, Shi L, Katsanis N, Silver DL. 2015.

Rbm8a haploinsufficiency disrupts embryonic cortical development resulting in

microcephaly. J Neurosci 35:7003-7018. 10.1523/JNEUROSCI.0018-15.2015.

22. Mao H, McMahon JJ, Tsai Y, Wang Z, Silver DL. 2016. Haploinsufficiency for Core

Exon Junction Complex Components Disrupts Embryonic Neurogenesis and Causes

p53-Mediated Microcephaly. PLoS Genetics 12:e1006282.

10.1371/journal.pgen.1006282.

23. Kurosaki T, Maquat LE. 2016. Nonsense-mediated mRNA decay in humans at a

glance. J Cell Sci 129:461-467. 10.1242/jcs.181008.

24. Serquiña AKP, Das SR, Popova E, Ojelabi OA, Roy CK, Göttlinger HG. 2013. UPF1

is crucial for the infectivity of human immunodeficiency virus type 1 progeny virions. J

Virol 87:8853-8861. 10.1128/JVI.00925-13.

25. Jerome M Molleston, Sara Cherry. 2017. Attacked from All Sides: RNA Decay in

Antiviral Defense. Viruses 9:2. 10.3390/v9010002.

26. Balistreri G, Bognanni C, Mühlemann O. 2017. Virus Escape and Manipulation of

Cellular Nonsense-Mediated mRNA Decay. Viruses 9:. 10.3390/v9010024.

27. Rigby RE, Rehwinkel J. 2015. RNA degradation in antiviral immunity and

autoimmunity. Trends in Immunology 36:179-188. 10.1016/j.it.2015.02.001.

28. Balistreri G, Horvath P, Schweingruber C, Zünd D, McInerney G, Merits A,

Mühlemann O, Azzalin C, Helenius A. 2014. The host nonsense-mediated mRNA decay

pathway restricts Mammalian RNA virus replication. Cell Host Microbe 16:403-411.

10.1016/j.chom.2014.08.007.

29. Ramage HR, Kumar GR, Verschueren E, Johnson JR, Von Dollen J, Johnson T,

Newton B, Shah P, Horner J, Krogan NJ, Ott M. 2015. A combined proteomics/genomics

approach links hepatitis C virus infection with nonsense-mediated mRNA decay. Mol Cell

57:329-340. 10.1016/j.molcel.2014.12.028.

30. Tani H, Imamachi N, Salam KA, Mizutani R, Ijiri K, Irie T, Yada T, Suzuki Y, Akimitsu

N. 2012. Identification of hundreds of novel UPF1 target transcripts by direct

determination of whole transcriptome stability. RNA Biology 9:1370-1379.

10.4161/rna.22360.

31. Weischenfeldt J, Damgaard I, Bryder D, Theilgaard-Mönch K, Thoren LA, Nielsen FC,

Jacobsen SEW, Nerlov C, Porse BT. 2008. NMD is essential for hematopoietic stem and

progenitor cells and for eliminating by-products of programmed DNA rearrangements.

Genes Dev 22:1381-1396. 10.1101/gad.468808.

32. Li T, Shi Y, Wang P, Guachalla LM, Sun B, Joerss T, Chen Y, Groth M, Krueger A,

Platzer M, Yang Y, Rudolph KL, Wang Z. 2015. Smg6/Est1 licenses embryonic stem cell

differentiation via nonsense-mediated mRNA decay. EMBO J 34:1630-1647.

10.15252/embj.201489947.

33. Brazão TF, Demmers J, van IJcken W, Strouboulis J, Fornerod M, Romão L, Grosveld

FG. 2012. A new function of ROD1 in nonsense-mediated mRNA decay. FEBS Lett

586:1101-1110. 10.1016/j.febslet.2012.03.015.

34. Chan W, Huang L, Gudikote JP, Chang Y, Imam JS, MacLean JA, Wilkinson MF.

2007. An alternative branch of the nonsense-mediated decay pathway. EMBO J 26:1820-

1830. 10.1038/sj.emboj.7601628.

35. Nelson JO, Moore KA, Chapin A, Hollien J, Metzstein MM. 2016. Degradation of

Gadd45 mRNA by nonsense-mediated decay is essential for viability. Elife 5:.

10.7554/eLife.12876.

36. Jauhiainen A, Thomsen C, Strömbom L, Grundevik P, Andersson C, Danielsson A,

Andersson MK, Nerman O, Rörkvist L, Ståhlberg A, Åman P. 2012. Distinct cytoplasmic

and nuclear functions of the stress induced protein DDIT3/CHOP/GADD153. PLoS ONE

7:e33208. 10.1371/journal.pone.0033208.

37. Salvador JM, Brown-Clay JD, Fornace AJ. 2013. Gadd45 in stress signaling, cell cycle

control, and apoptosis. Adv Exp Med Biol 793:1-19. 10.1007/978-1-4614-8289-5_1.

38. Gehring NH, Lamprinaki S, Kulozik AE, Hentze MW. 2009. Disassembly of exon

junction complexes by PYM. Cell 137:536-548. 10.1016/j.cell.2009.02.042.

39. Chamieh H, Ballut L, Bonneau F, Le Hir H. 2008. NMD factors UPF2 and UPF3 bridge

UPF1 to the exon junction complex and stimulate its RNA helicase activity. Nat Struct Mol

Biol 15:85-93. 10.1038/nsmb1330.

40. Varsally W, Brogna S. 2012. UPF1 involvement in nuclear functions. Biochem Soc

Trans 40:778-783. 10.1042/BST20120052.

41. Klema VJ, Padmanabhan R, Choi KH. 2015. Flaviviral Replication Complex:

Coordination between RNA Synthesis and 5′-RNA Capping. Viruses 7:4640-4656.

10.3390/v7082837.

42. Maisel M, Herr A, Milosevic J, Hermann A, Habisch H, Schwarz S, Kirsch M,

Antoniadis G, Brenner R, Hallmeyer-Elgner S, Lerche H, Schwarz J, Storch A. 2007.

Transcription profiling of adult and fetal human neuroprogenitors identifies divergent

paths to maintain the neuroprogenitor cell state. Stem Cells 25:1231-1240.

10.1634/stemcells.2006-0617.

43. Li H, Saucedo-Cuevas L, Regla-Nava J, Chai G, Sheets N, Tang W, Terskikh A,

Shresta S, Gleeson J. 2016. Zika Virus Infects Neural Progenitors in the Adult Mouse

Brain and Alters Proliferation. Cell Stem Cell 19:593-598. 10.1016/j.stem.2016.08.005.

44. Nickless A, Bailis JM, You Z. 2017. Control of gene expression through the nonsense-

mediated RNA decay pathway. Cell & Bioscience 7:1-12. 10.1186/s13578-017-0153-7.

45. Cortese M, Goellner S, Acosta EG, Neufeldt CJ, Oleksiuk O, Lampe M, Haselmann

U, Funaya C, Schieber N, Ronchi P, Schorb M, Pruunsild P, Schwab Y, Chatel-Chaix L,

Ruggieri A, Bartenschlager R. 2017. Ultrastructural Characterization of Zika Virus

Replication Factories. Cell Rep 18:2113-2123. 10.1016/j.celrep.2017.02.014.

46. Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, Schwarz

MC, Sánchez-Seco MP, Evans MJ, Best SM, García-Sastre A. 2016. Zika Virus Targets

Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host Microbe 19:882-890.

10.1016/j.chom.2016.05.009.

47. von Mikecz A. 2006. The nuclear ubiquitin-proteasome system. J Cell Sci 119:1977-

1984. 10.1242/jcs.03008.

48. Freire JM, Veiga AS, Conceição TM, Kowalczyk W, Mohana-Borges R, Andreu D,

Santos NC, Da Poian AT, Castanho, Miguel A. R. B. 2013. Intracellular nucleic acid

delivery by the supercharged dengue virus capsid protein. PLoS ONE 8:e81450.

10.1371/journal.pone.0081450.

49. HD iPSC Consortium. 2017. Developmental alterations in Huntington's disease neural

cells and pharmacological rescue in cells and mice. Nat Neurosci 20:648-660.

10.1038/nn.4532.

50. Jäger S, Gulbahce N, Cimermancic P, Kane J, He N, Chou S, D'Orso I, Fernandes J,

Jang G, Frankel AD, Alber T, Zhou Q, Krogan NJ. 2011. Purification and characterization

of HIV-human protein complexes. Methods 53:13-19. 10.1016/j.ymeth.2010.08.007.

51. Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate proteome-

wide label-free quantification by delayed normalization and maximal peptide ratio

extraction, termed MaxLFQ. Mol Cell Proteomics 13:2513-2526.

10.1074/mcp.M113.031591.

52. Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.

Nat Biotechnol 26:1367-1372. 10.1038/nbt.1511.

53. Teo G, Koh H, Fermin D, Lambert J, Knight JDR, Gingras A, Choi H. 2016. SAINTq:

Scoring protein-protein interactions in affinity purification - mass spectrometry

experiments with fragment or peptide intensity data. Proteomics 16:2238-2245.

10.1002/pmic.201500499.

54. Costes SV, Daelemans D, Cho EH, Dobbin Z, Pavlakis G, Lockett S. 2004. Automatic

and quantitative measurement of protein-protein colocalization in live cells. Biophys J

86:3993-4003. 10.1529/biophysj.103.038422.

55. Dunn KW, Kamocka MM, McDonald JH. 2011. A practical guide to evaluating

colocalization in biological microscopy. Am J Physiol Cell Physiol 300:C742.

10.1152/ajpcell.00462.2010.

56. Paul LM, Carlin ER, Jenkins MM, Tan AL, Barcellona CM, Nicholson CO, Michael SF,

Isern S. 2016. Dengue virus antibodies enhance Zika virus infection. Clinical &

Translational Immunology 5:n/a. 10.1038/cti.2016.72.

Figure Legends

Figure 1. ZIKV infection disrupts the NMD pathway.

(a) Transcript levels of NMD substrates and housekeeping genes from Huh7 cells or

NPCs mock-infected or infected with ZIKV strain P6-740 or the contemporary clinical

isolate PRVABC59. Cells were infected at a multiplicity of infection (MOI) of 0.1 or 1 and

harvested at 48 hpi. Data are represented as mean ± s.e.m. *P* values were calculated by

unpaired Student's *t*-test. *$P \leq 0.05$; **$P \leq 0.01$; ns, not significant. n= 3 independent

experiments.

(b) Venn diagram showing overlap of significantly upregulated genes associated with

ZIKV infection of NPCs and UPF1 knockdown in HeLa cells. RNA-Seq analyses of mock-

infected or ZIKV-infected NPCs harvested at 56 hpi and control siRNA-treated or UPF1

siRNA-treated HeLa TO cells harvested at 72 h post-transfection (hpt). The GeneProf

hypergeometric probability calculator was then used to generate a hypergeometric *P*

value. ****$P \leq 0.0001$.

(c) Transcript levels of housekeeping genes and select genes involved in cell cycle growth

arrest and apoptosis that were identified in (b). Huh7 cells were mock-infected or infected

with ZIKV PRVABC59 at an MOI of 0.1 or 1 and harvested at 48 hpi. Data are represented

as mean ± s.e.m. *P* values were calculated by unpaired Student's *t*-test. *$P \leq 0.05$; **$P \leq$

0.01; ***$P \leq 0.001$; ns, not significant. n= 3 independent experiments.

Figure 2. The capsid protein of ZIKV interacts with the NMD pathway.

(a) Ugandan ZIKV capsid (Ug Cap, MR 766) and French Polynesian ZIKV capsid (Fp Cap, H/PF/2013) PPI maps that show significant enrichment for host NMD-associated factors (purple), as identified by AP-MS (SAINTq probability score > 0.9 and FDR < 0.05). Ten interactions between Fp Cap and host NMD factors (hypergeometrical test, P value = 7.16 x 10^{-10}) and eight interactions between Ug Cap and host NMD factors (P value = 3.45 x 10^{-7}) were identified.

(b) Co-immunoprecipitation (co-IP) and western blot analysis of HEK293T cells transfected with vector or Flag-tagged ZIKV capsid (H/PF/2013, Asian lineage) and harvested at 48 hpt to immunoprecipitate endogenous UPF3B. The upper band detected in the IP Capsid blot represents a non-specific artifact.

(c) Co-IP and western blot analysis of HEK293T cells transfected with vector or Flag-tagged ZIKV capsid and harvested at 48 hpt to immunoprecipitate endogenous UPF1.

(d) Myc-tag co-IP and western blot analysis of HEK293T cells transfected with Strep-tagged ZIKV capsid and Myc-UPF1 (WT), Myc-UPF1-C126S (RNA-binding mutant), or Myc-UPF1-G495R/G497E (ATPase/helicase mutant) and harvested at 48 hpt to immunoprecipitate ZIKV capsid.

(e) Western blot analysis of UPF1 levels in mock-infected and ZIKV-infected (PRVABC59, MOI of 1) Huh7 cells or mock-infected and ZIKV-infected (P6-740, MOI of 1) NPCs harvested at 48 hpi, with β-actin and ZIKV envelope (ZIKV E) or ZIKV capsid (ZIKV C) protein serving as loading and infection controls, respectively. Densitometric analyses were performed using ImageJ to quantify relative band intensities. Data are represented

as mean ± s.e.m. *P* values were calculated by unpaired Student's *t*-test. **P ≤ 0.01; ***P

≤ 0.001. n= 3 independent experiments.

Figure 3. ZIKV capsid degrades UPF1, the master regulator of NMD, via a

proteasome-dependent mechanism.

(a) Western blot analysis of UPF1 levels in subcellular fractionated HEK293T cells

transfected with vector or Flag-tagged ZIKV capsid (H/PF/2013, Asian lineage) for 48 h.

GAPDH was used as a cytoplasmic marker and SP1 as a nuclear marker to ensure

optimal fractionation. Densitometric analyses were performed using ImageJ to quantify

relative band intensities. Data are represented as mean ± s.e.m. *P* values were calculated

by unpaired Student's *t*-test. **P ≤ 0.01; ns, not significant. n= 3 independent experiments.

(b) Western blot analysis of nuclear UPF1 levels in fractionated HEK293T cells

transfected with vector or Flag-tagged ZIKV capsid for 48 h. Cells were treated with

DMSO or increasing concentrations of the proteasome inhibitor bortezomib for 24 h

before harvest. Densitometric analyses were performed using ImageJ to quantify relative

band intensities. Data are represented as mean ± s.e.m. *P* values were calculated by

one-way ANOVA with multiple comparisons. *P ≤ 0.05; ns, not significant. n= 3

independent experiments.

(c) Representative 3D confocal microscopy images of the nuclei of Huh7-Lunet cells

transfected with Strep-tagged ZIKV capsid. Cells were treated at 24 hpt with DMSO or 10

nM bortezomib and processed for immunostaining at 48 hpt with antibodies against Strep-

tag (turquoise) and endogenous UPF1 (purple). DAPI (blue) was used to stain and define

the nuclei. Each channel was reconstructed digitally for visualization of the 3D

colocalization. The thresholded Mander's correlation coefficients were determined and *P*

value was calculated by unpaired Student's *t*-test. **P ≤ 0.01. n = 8 cells per condition.

Scale bar represents 3 µm.

Figure 4. UPF1 knockdown enhances the permissivity of NPCs to ZIKV infection.

(a) Western blot analysis of UPF1 levels in NPCs transfected with non-targeting siRNA

(siNT) or a pool of UPF1-specific siRNAs (siUPF1) at 96 hpt. Densitometric analyses were

performed using ImageJ to quantify relative band intensities. Data are represented as

mean ± s.e.m. *P* value was calculated by unpaired Student's *t*-test. **P ≤ 0.01. n= 3

independent experiments using one NPC line.

(b) ZIKV RNA levels in siNT-treated or siUPF1-treated NPCs infected with ZIKV strain

PRVABC59 at an MOI of 0.1 or 1 and harvested at 48 hpi. Data are represented as mean

± s.e.m. *P* value was calculated by two-tailed ratio paired Student's *t*-test. ***P ≤ 0.001.

n= 3 independent experiments using one NPC line.

(c) Released infectious virus from siNT-treated or siUPF1-treated, ZIKV-infected (MOI of

1) NPCs harvested at 48 hpi. Data are represented as mean ± s.e.m. *P* value was

calculated by unpaired Student's *t*-test. **P ≤ 0.01. n= 3 independent experiments using

one NPC line.

(d) Representative confocal microscopy images of a ZIKV-infected, siNT-treated NPC or

a ZIKV-infected, siUPF1-treated NPC with the nuclei stained with DAPI (blue) and ZIKV

dsRNA foci stained with the anti-dsRNA mAb J2 (teal). 3D image rendering and

reconstructed dsRNA foci were produced using the Imaris spot detection function. Scale

bar represents 2 μm.

(e) Number of dsRNA foci were averaged for each cell. Data are represented as mean ±

s.e.m. P value was calculated by two-tailed ratio paired Student's t-test. ns, not significant.

n = 3 independent experiments using two NPC lines, with 3-10 cells analyzed per

condition for each experiment.

(f) Measurement of dsRNA foci volume were averaged for each cell. Data are represented

as mean ± s.e.m. P value was calculated by two-tailed ratio paired Student's t-test. ns,

not significant. n = 3 independent experiments using two NPC lines, with 3-10 cells

analyzed per condition for each experiment.

(g) Infection rates of siNT-treated or siUPF1-treated, ZIKV-infected (MOI of 1) NPCs

measured at 48 hpi. Fixed cells were subjected to the anti-DENV mAb 1.6D, which also

recognizes the ZIKV envelope protein (56). Data are represented as mean ± s.e.m. P

value was calculated by two-tailed ratio paired Student's t-test. **P ≤ 0.01. n = 3

independent experiments using two NPC lines.

(h) Model of the interaction between the capsid protein of ZIKV and UPF1 of the NMD

pathway.

Supplemental Figure 1. UPF1 is not transcriptionally downregulated during ZIKV

infection or following ZIKV capsid overexpression.

(a) UPF1 transcript levels from Huh7 cells mock-infected or infected with ZIKV strain

PRVABC59 at an MOI of 0.1 or 1 and harvested at 48 hpi. Data are represented as mean

± s.e.m. *P* values were calculated by unpaired Student's *t*-test. ns, not significant. n= 3

independent experiments.

(b) UPF1 transcript levels from HEK293T cells transfected with vector or Strep-tagged

ZIKV capsid (H/PF/2013, Asian lineage) and harvested at 48 hpt. Data are represented

as mean ± s.e.m. *P* values were calculated by unpaired Student's *t*-test. ns, not

significant. n= 3 independent experiments.

Supplemental Figure 2. ZIKV capsid-induced UPF1 degradation is not dependent

on autophagy.

Western blot analysis of nuclear UPF1 levels in fractionated HEK293T cells transfected

with vector or Flag-tagged ZIKV capsid for 48 h. Cells were treated with DMSO or the

autophagy inhibitor bafilomycin A1 (Baf) (10 nM) for 24 h before harvest. Levels of p62,

which is degraded by autophagy, were monitored to confirm autophagic inhibition

following bafilomycin A1 treatment.

Supplemental Figure 3. ZIKV capsid colocalizes with endogenous UPF1.

Representative 3D confocal microscopy images of Huh7-Lunet cells transfected with vector or

Strep-tagged ZIKV capsid. Cells were processed for immunostaining at 48 hpt and probed with

antibodies against Strep-tag (turquoise) and endogenous UPF1 (purple). DAPI (blue) was used

to stain the nuclei. Each channel was reconstructed digitally for visualization of the 3D

colocalization. The thresholded Mander's correlation coefficient for ZIKV capsid was 0.57 (n =

17), indicating that approximately 57% of the voxels positive for capsid fluorescence were also

positive for UPF1 fluorescence. Scale bar represents 5 μm.

a

UPF1

β-Actin

b

c

d dsRNA

siNT

siUPF1

e

f

g

h Zika

Capsid

UPF1

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3