Leaps and bounds: geographical and ecological distance constrained the colonisation of the

Afrotemperate by *Erica*

Pirie, M.D. [1, 2, 8], Kandziora, M. [1,4], Nürk, N.M. [5], Le Maitre, N.C. [2], Mugrabi de Kuppler, A. [6], Gehrke, B. [1],

Oliver, E.G.H. [7], and Bellstedt, D.U. [2]

Author affiliations

[1]Institut für Organismische und Molekulare Evolutionsbiologie, Johannes Gutenberg-Universität,

Anselm-Franz-von-Bentzelweg 9a, 55099 Mainz, Germany

[2]Department of Biochemistry, University of Stellenbosch, Private Bag X1, Matieland 7602, South

Africa

[3]University Museum of Bergen, University of Bergen, Bergen, Norway

[4]Life and Environmental Sciences, School of Natural Sciences, University of California, Merced, USA

[5]Institute of Plant Systematics, Bayreuth Centre of Ecology and Environmental Research (BayCEER),

University of Bayreuth, Universitätsstraße 30, 95447 Bayreuth, Germany

[6]INRES Pflanzenzüchtung, Rheinische Friedrich-Wilhelms-Universität Bonn, Katzenburgweg 5, 53115

Bonn, Germany

[7]Department of Botany and Zoology, University of Stellenbosch, Private Bag X1, Matieland 7602,

South Africa

[8]Author for correspondence: pirie@uni-mainz.de; phone: +49 6131 3922928

Abstract

The coincidence of long distance dispersal and biome shift is assumed to be the result of a

multifaceted interplay between geographical distance and ecological suitability of source and sink

areas. Here, we test the influence of these factors on the dispersal history of the flowering plant

genus *Erica* (Ericaceae) across the Afrotemperate. We quantify similarity of *Erica* climate niches per

biogeographic area using direct observations of species, and test various colonisation scenarios while

estimating ancestral areas for the *Erica* clade using parametric biogeographic model testing. We infer

that the overall dispersal history of *Erica* across the Afrotemperate is the result of infrequent

colonisation limited by geographic proximity and niche similarity. However, the Drakensberg

Mountains represent a colonisation sink, rather than acting as a "stepping stone" between more

distant and ecologically dissimilar Cape and Tropical African regions. Strikingly, the most dramatic

examples of species radiations in *Erica* were the result of single unique dispersals over longer

distances between ecologically dissimilar areas, contradicting the rule of phylogenetic biome

conservatism. These results highlight the importance of rare biome shifts, in which a unique dispersal

event fuels evolutionary radiation.

Keywords: Afrotemperate; historical biogeography; phylogenetic biome conservatism; Cape Floristic

Region; climatic niche shift; *Erica*; evolution; Madagascar; model testing

Introduction

The current day distributions of many plant groups are the result of long distance dispersal (LDD)

(Muñoz et al., 2004; Alsos et al., 2007; Kadereit & Baldwin, 2012; Ruhfel et al., 2016; Jordano, 2017).

Such events are thought to be rare (Nathan, 2006; but see Viana & al., 2015), but rarer still might be

plant dispersals across long distances between different biomes (Crisp et al., 2009). The coincidence

of intercontinental dispersal and biome shift, such as inferred in *Lupinus* (Drummond et al., 2012),

Bartsia (Uribe-Convers & Tank, 2015), and *Hypericum* (Nürk, Michling & Linder, 2018), is assumed to

be the result of a multifaceted interplay between geographical distance and ecological suitability of

source and sink areas (Donoghue & Edwards, 2014). Here, we test the influence of these factors on

the biogeographic history of the flowering plant genus *Erica* (Ericaceae).

The more than 800 *Erica* species across Europe and Africa provide an excellent example with which

to test the impact of geographical and ecological distance on biogeographic history. Just 21 of the

species are found in Central and Western Europe, Macaronesia, the Mediterranean and the Middle

East. This species-poor assemblage nevertheless most likely represents the ancestral area of the

clade (McGuire & Kron, 2005; Mugrabi de Kuppler et al., 2015; Kowalski & Fagúndez, 2017) where

the oldest lineages began to diversify c. 30 Ma (Pirie et al., 2016). From around 15 Ma, a single

lineage dispersed across different biomes of the Afrotemperate (sensu White, 1981): today 23

species are known from the high mountains of Tropical Africa; 51 in Southern Africa's Drakensberg

Mountains; c. 41 in Madagascar and the Mascarene islands; and c. 690 in the Cape Floristic Region of

South Africa (Oliver, 2012; Pirie et al., 2016). Present day habitats of *Erica* species tend to be low

nutrient and fire prone (Ojeda, 1998), but hence still differ markedly in ecology, from the

Mediterranean climates of southern Europe and the Cape to colder climes of northern Europe and

the non-seasonal temperate habitats of the high mountains in Tropical Africa. These habitats are also

separated by considerable geographic distances, isolated by expanses of inhospitable ecosystems

and/or ocean. Nonetheless, similar distribution patterns across Europe and Africa are observed in

different plant groups (e.g. Gizaw et al., 2016; Mairal, Sanmartín & Pellissier, 2017).

Organisms adapted to different habitats respond differently to changing environmental conditions

(Mairal, Sanmartín & Pellissier, 2017; Chala et al., 2017). Distribution patterns of arid-adapted plant

groups, for which suitable habitats in Africa have been more contiguously distributed (Bellstedt et al.,

2012), might thus be best described by biogeographic scenarios emphasising vicariance processes,

such as for example the "Rand Flora" representing plant lineages that show similar disjunct

distributions around the continental margins of Africa (Sanmartín et al., 2010; Pokorny et al., 2015),

or the "African arid corridor" hypothesis that seeks to explain disjunct distributions between the

Horn of Africa and arid south-western Africa (Verdcourt, 1969; White, 1983). Models that invoke

concerted patterns of LDD might instead apply to plants adapted to habitats that remained largely

isolated over time (Knox & Palmer, 1998; Galbany-Casals et al., 2014; Nürk et al., 2015; Míguez et al.,

2017). Examples include the shared arid adapted elements of Macronesia and adjacent North-West

Africa and Mediterranean (Kim et al., 2008; Fernández-Palacios et al., 2011; García-Aloy et al., 2017),

and the more mesic temperate or tropical alpine habitats of the "sky islands" of East Africa, in which,

for example, multiple lineages originated from northern temperate environments (Gehrke & Linder,

2009; Gizaw et al., 2013, 2016).

A more specific biogeographic scenario, inferred from Cape clades with distributions very similar to

that of *Erica,* involves dispersal north from the Cape to the East African mountains via the

Drakensberg ("Cape to Cairo"; Galley & al., 2007). McGuire & Kron (2005) proposed a different

scenario for *Erica* instead: southerly stepping stone dispersal through the African high mountains to

the Cape. Both scenarios, however, imply that dispersal is more frequent between adjacent

areas/over shorter distances. Short distance or stepping stone dispersal may indeed be more

probable than LDD (Nathan, 2006), and distance alone could conceivably be more important than

directionality (Linder et al., 2013). On the other hand, the probabilities of LDDs are hard to model

(Higgins & Richardson, 1999; Nathan, 2006), in part because (observable) LDD events also involve

successful establishment in more or less distinct environments (Donoghue & Edwards, 2014). Thus

geographic distance and ecological suitability might individually constrain the biogeographic history

of plants, or the interplay between both factors may be decisive (Donoghue, 2008; Carvajal-Endara et

al., 2017), so much so that clades with similar ecological tolerances and origin might show

convergence to similar distribution patterns (Merckx et al., 2015; Gizaw et al., 2016; Mairal,

Sanmartín & Pellissier, 2017).

In this paper, we ask whether and to what extent geographic proximity or climatic niche similarity

constrained the colonisation of the Afrotemperate by *Erica*. Until recent work (Pirie, Oliver &

Bellstedt, 2011; Pirie et al., 2016), too little was known of the phylogenetic relationships of the 97%

of *Erica* species outside Europe to be able to address such questions. Specifically, we test six

biogeographic models, as illustrated in Fig. 1: Three that test the influence of geographic distance,

climatic niche similarity, and the combination of both; and three differing stepping stone models that

each imply geographical distance effects promoting dispersal predominantly between adjacent

areas: northerly "Cape to Cairo", "Southerly stepping stone" and a model that invokes elements of

both, the "Drakenberg melting pot" hypothesis.

Materials and Methods

Phylogenetic hypothesis: Analyses were based on phylogenetic trees (Pirie et al., 2016) which

represent c. 60% of the c. 800 species of *Erica* from across their geographic range and DNA

sequences from multiple plastid markers (*trnT-trnL* and *trnL-trnF-ndhJ* spacer sequences for all taxa,

with exemplar sampling of *trnL* intron, *atpI-atpH* spacer, *trnK-matK* intron and *matK* gene, *psbM-trnH*

spacer, *rbcL* gene, *rpl16* intron, *trnL-rpl32* spacer sequences) and nuclear ribosomal (nrDNA) internal

transcribed spacer (ITS; for all taxa). For the biogeographic analyses here, we adopt the phylogenetic

hypothesis of Pirie et al. (2016), the best tree inferred under Maximum Likelihood (ML) using RAxML

(Stamatakis, 2006), based on the combined data and 597 taxa and rate smoothed using RELTIME

(Tamura et al., 2012) with a single secondary calibration point derived from a wider fossil calibrated

analysis of Ericaceae (Schwery et al., 2015). Pirie & al. (2016) identified a "Cape clade" that included

all but one of the sampled species of *Erica* found in the CFR. The single exception was *E.*

pauciovulata, which was placed within a polytomy including the Cape clade and other Afrotemperate

lineages. This may, however, be artefactual due to sequence anomalies in the *trnL-trnF-ndhJ* spacer

region of *E. pauciovulata*. Preliminary results based on additional sampling including nrDNA ETS (Pirie

et al. in prep.) confirm the monophyly of Cape clade including *E. pauciovulata*, and we therefore

exclude this taxon from biogeographic analyses to avoid inferring an independent colonisation of the

CFR as a result of its uncertain position.

Defining the pure-distance and the niche-based models: Five biogeographic areas of the *Erica*

distribution were defined following Pirie & al. (2016): Europe (including northern Africa); Tropical

Africa (TA); Madagascar; Drakensberg; Cape. We obtained occurrence data for *Erica* species from our

own collections, and from PRECIS (representing mostly southern African collections, held by the

South African National Biodiversity Institute; http://newposa.sanbi.org/) and GBIF

(https://www.gbif.org/) databases. We curated the species occurrence data by removing obviously

erroneous locality data, duplicated records, and records with imprecise occurrence data (coordinates

with ≤ 3 decimal places, many of which represented centroids of quarter degree squares which were

originally represented in PRECIS), but did not further consider the source of or information on the

precision of the geographical coordinates, because these are most often not stated in the database-

derived occurrence records. This resulted in 6818 individual occurrences representing the species in

the phylogenetic trees (Appendix 1). Based on these individual occurrences we estimated the

geographic range of the species in the five biogeographic areas (resulting in what we term 'area

ranges', i.e. the distributional range of all of the species in a certain biogeographic area) We assessed

the area ranges in a biogeographic area placing a buffer of one minutes of arc in radius (ca. 11 km)

and 50 m elevation around the individual species occurrences (Europe 4667, Tropical Africa 42,

Madagascar 70, Drakensberg 58, and Cape 1981 occurrences; Appendix 2) conservatively aiming at a

representative approximation of spatial extent (Nakazato, Warren & Moyle, 2010; Anacker & Strauss,

2014) and ecological conditions of the species' distribution in a respective biogeographic area. These

area ranges, which include up to several thousands of spatial points, were used in the subsequent

analyses to calculate geographical and ecological distances between biogeographic areas.

To incorporate a measure of geographic proximity among areas in a solely distance-based

biogeographic model (the 'geographic distance' model; Fig. 1), we calculated the overall minimum

geographic pairwise distances between the area ranges according to Meeus (1999) in WGS84

projection using the raster 2.3-33 package (Hijmans, 2015) in R (R Development Core Team, 2013).

We converted geographic distances into dispersal rate multipliers (0-1, whereby the largest distance

has the smallest dispersal probability), while comparing the effect of scaling the distances linearly

(applying a linear model with intercept of 1 and a slope of -1.52^{-07}) and exponentially (-0.25, -1 and -

2).

To incorporate in a niche-based biogeographic model, the 'niche similarity' model (Fig. 1), a measure

of climatic similarity between the biogeographic areas we built a multidimensional environmental

model representing the full space of all available climates in the global study area (i.e. most of

Europe and entire Africa, represented by >0.5 million spatially independently sampled point

locations; Appendix 2) using principal component analysis (PCA) in R's ade4 1.6-2 (Dray & Dufour,

2007). We then calculated the pairwise climate similarity between the area ranges (as defined by the

species occurrence data; see above) as Schoener's D (Schoener, 1968; ranging from 0 = no similarity

to 1 = identical) while correcting for the entire climatic variance in the global study area and between

the biogeographic areas (Broennimann et al., 2012) by kernel density smoothers using ecospat 2.1.1

(Broennimann et al., 2014). Finally, the biogeographic model for the niche similarity hypothesis was

defined using the pairwise Schoener's D values for the combined PCA axes 1 and 2 directly as

dispersal rate multipliers between areas (for details see protocol in Appendix 2).

Finally, to consider both geographical and environmental distances in a joint model, also accounting

for a negative correlation between both geographic and environmental distances (Kendall's R = -

0.64), we used two rate multiplier matrices, representing both climatic niche and physical distance

(converted into probabilities; see above), as input.

Biogeographic model testing and ancestral area reconstruction: We used BioGeoBEARS (Matzke,

2013) for parametric model testing, whilst aware of the debate surrounding these models and their

comparison (Ree & Sanmartín, 2018; see Results and Discussion). The above defined biogeographic

models (Fig. 1) were parameterized using different dispersal rate multipliers (see below and

Appendix 3) and compared to null models that do not incorporate any constraints. As input data we

used the rate-smoothed ML phylogeny reduced to one tip per sampled species (Pirie & al., 2016; the

"best tree"), a file delimiting the distributional range of species, and a file indicating

connectivity/distance between the different areas of the *Erica* distribution (varying for the different

biogeographic models; Fig. 1, Appendix 3). Model fit of the different nested and non-nested models

was tested using the Akaike Information Criterion (AIC) and the delta AIC (Burnham & Anderson,

2002). For model testing we additionally used nine trees from the RAxML bootstrap analyses of Pirie

& al. (2016) of the same dataset (rate-smoothed using the ape package in R; Paradis & al., 2004).

These trees were selected to represent the possible resolutions of phylogenetic uncertainty between

the geographically restricted major clades (Appendix 4) but were otherwise chosen randomly with

respect to topologies and branch lengths. All hypotheses were implemented with combinations of

dispersal-extinction-cladogenesis (DEC; Ree et al., 2005; Ree & Smith, 2008), Bayarea-like or DIVA-

like models, with or without allowing long distance dispersal (the "+J" model of Matzke, 2014). We

focus on DEC and DEC+J models because these generally fit the data better than Bayarea-like or

DIVA-like models.

Prior to comparing the different biogeographic hypotheses, we tested whether an unconstrained

model fitted the data better than a) restricting the maximum number of areas at nodes to two;

and/or b) implementing an adjacent area matrix (Appendix 3; Results). The Southerly stepping stone,

Cape to Cairo, and Drakensberg melting pot hypotheses were then run under a range of different

dispersal multipliers (0.00, 0.01, 0.05, 0.075, 0.1, 0.25 and 0.5; and for the DEC+J model also on the

nine bootstrap trees with dispersal multipliers of 0.01, 0.1, 0.25 and 0.5) to test whether these

arbitrary values influenced the results. For niche- and distance-based biogeographic models

Schoener's *D* values and geographic distances scaled to probabilities (see above) were used directly

as dispersal multipliers (Appendix 3). We also assessed the impact on model fit (given the best fitting

model) of a number of different values for the parameter "w", which is an exponent for the dispersal

multipliers (which otherwise was fixed to "1"; Appendix 3); and coding of *E. arborea* as European

(following Mugrabi de Kuppler et al., 2015), rather than as widespread between Europe and Tropical
Africa. Further details and example files for the BioGeoBEARS analyses are presented in Appendix 3.

Estimating dispersal rates: For the best models under both DEC+J and DEC, given the best tree, we
estimated the number and type of biogeographic events across the clade using Biogeographical
Stochastic Modelling (BSM) as implemented in BioGeoBEARS (Matzke, 2014). BSM simulates histories
of the times and locations of dispersal events. Frequencies were estimated by taking the mean and
standard deviation of event counts from 50 BSMs. We also compared the results to that of simple
parsimony optimisation using Mesquite v3.31 (Maddison & Maddison, 2006), under the assumption
that LDD events are simply rare (Pirie et al., 2012). We incorporated phylogenetic uncertainty by
summarising the results over the complete sample of 252 RAxML bootstrap trees adapted from Pirie
& al. (2016), and coding *E. arborea* either as widespread between Europe and Tropical Africa or
European (Appendix 5).

214
Results

Niche similarity model: The environmental space that represents all climates available in the study
area – most of Europe and all of Africa – and that was used to approximate the climatic similarity
between biogeographic areas (area ranges) calculated as hypervolume corrected Schoener's *D*
accounting for climatic variation in the study area and between the area ranges, explained >88% of
the climate variation on the first two PCA axes. Despite the range of differing conditions within areas,
e.g. with rainfall seasonality differing according to elevation, variation between area ranges was
considerable (the distribution and the median values for Schoener's *D* per PCA axis pairwise for the
areas are presented in Appendix 6, and for the combined axes 1 and 2 in Appendix 7). According to
the latter, the Cape and Drakensberg areas are climatically most similar (*D*: 0.71) and Europe and
Madagascar are most different (*D*: 0.21). Most similar to the European are the Cape and Drakensberg
climates (both *D*: 0.35), and the Tropical Africa climate (*D*: 0.27; Fig. 2C).

Biogeographic model testing: Assuming that AIC values of the differing models can be compared (but
see Ree and Sanmartín, 2018), DEC/DEC+J models generally fit the data better than Bayarea-like or
DIVA-like models and DEC+J models generally fit the data better than equivalent DEC ones (Appendix
8). Under DEC+J, models including an adjacent area matrix fitted the data better than those without
constraint to dispersal. We additionally fixed the maximum number of ancestral areas to two,
increasing the speed of the analyses without negatively impacting model fit. Under DEC, models with
maximum areas at nodes restricted to two fitted the data better than those without constraint to
ancestral ranges. Under both DEC+J and DEC, geographic distance fitted the data better when

translated linearly into dispersal rate probabilities (0-1) than when scaled exponentially (Appendix 8);

we therefore focus on models using the probabilities, referring to them simply as "geographical

distance". The DEC+J results in general do not show the flaws as reported by Ree and Sanmartín

(2018). For example, the values for range expansion (parameter d) were similar and low (0.0030 and

0.0027 per Ma respectively; Appendix 9). Under DEC+J, cladogenetic dispersal (parameter j) was

0.0024 per node, i.e. lower than d (particularly given an average branch length across the *Erica*

phylogeny of 1.78 Ma, variance of 11.67) and much lower than the maximum permitted value (3).

Under DEC+J given the best tree, the "Drakensberg melting pot", "geographic distance", and

"Southerly stepping stone" models revealed the best fit (lowest AIC with deltaAIC ≥2); under DEC the

Drakensberg melting pot model alone scored best, but with AIC 141 compared to AIC 131 for DEC+J

(Appendix 8). Adopting DEC+J as the generally better fitting and biologically more realistic model (see

Discussion), we assessed the results given phylogenetic uncertainty represented by selected

bootstrap trees. Based on the bootstrap trees, the combined niche-geographic distance hypothesis

was often among the best fitting models (deltaAIC <2 given eight of nine trees), scoring better than

pure distance (deltaAIC <2 for five trees), or niche similarity (deltaAIC <2 for four trees) alone. The

"Cape to Cairo" model generally fitted better than most other biogeographic scenarios (deltaAIC <2

for eight of nine trees, compared to Drakensberg melting pot (deltaAIC <2 for two of nine trees) and

southerly stepping stone (not amongst the best fitting models); Table 1; Appendix 8).

Ancestral area reconstruction: Overall, we infer a colonisation path of *Erica* from Europe to the Cape

via an initial migration to Tropical Africa, under DEC+J and irrespective of best fitting model or

phylogenetic uncertainty. When *E. arborea* is treated as widespread between Europe and Tropical

Africa, the common ancestor of the African/Madagascan clade is inferred to have been similarly

widespread. When *E. arborea* is treated as ancestrally European, dispersal from Europe to Tropical

Africa is inferred without a transitional widespread distribution. Under DEC, the colonisation path to

the Cape is also via an initial migration to Tropical Africa, then a widespread distribution between

Tropical Africa and the Cape, followed by an extinction in Tropical Africa. Whether *E. arborea* is

treated as widespread between Europe and Tropical Africa or not, the common ancestor of the

African/Madagascan clade is inferred to have been similarly widespread between Europe and

tropical Africa. Ancestral area reconstructions given the best tree under the best fitting models (as

well as under a model without range or dispersal constraints for comparison; in each case under both

DEC+J and DEC) are presented in Appendix 13. Overall, ancestral areas inferred under parsimony

were consistent with those inferred under parametric models (more so with those under DEC+J,

given that widespread distributions are not incorporated into standard character optimisation), with

the numbers and directions of shifts unaffected by phylogenetic uncertainty.

The vast majority of biogeographic events inferred using BSM under both DEC+J and DEC were

within-area speciation (97.15 % and 96.26% respectively; Appendix 9). Under DEC+J, few range

expansion events were inferred between Europe and Tropical Africa and between Tropical Africa and

the Drakensberg region, with most between Cape and the Drakensberg regions (Appendix 10).

Dispersal rates between area ranges inferred under BSM are summarised in Fig. 2C. A single founder

event (parameter j) was inferred from Tropical Africa to the Cape region, with fewer events between

the Drakensberg and Tropical Africa and between Tropical Africa and Madagascar. Overall, most

founder events took place from Tropical Africa (1.96 [standard deviation of 0.47] events averaged

across 50 BSM; Appendix 11). In addition to the most commonly inferred range expansions given

DEC+J, under DEC additional range expansions were inferred from Tropical Africa to Madagascar and

from Tropical Africa to the Cape (Appendix 10). With each range expansion under DEC, the

corresponding ancestral distribution was widespread. Under both DEC+J and DEC dispersal rates

between Tropical Africa and the Drakensberg were roughly symmetrical, as opposed to those

between the Cape and the Drakensberg or between Europe and Tropical Africa which were

asymmetrical (Fig. 2; Appendix 12).

284
285
Discussion

In this study, we modelled shifts between biomes and dispersals over larger distances in the

evolution of *Erica*, in order to test six hypotheses for the origins of Afrotemperate plant groups (Fig.

1). Three models concerned general factors considered of importance in limiting plant dispersal:

geographical distance, similarity of realised climatic niches, and a combination of geographical and

ecological proximity. The remaining three models described specific colonisation hypotheses of the

Afrotemperate, in each case proposing a stepwise shift in distributions between adjacent areas.

These models differed in the area of origin and in the direction of dispersal: northerly dispersal from

the Cape ("Cape to Cairo"), versus southerly dispersal from Europe ("Southerly stepping stone"), or a

combination of both (termed here "Drakensberg melting-pot").

Of the stepping-stone-dispersal models, "Cape to Cairo" and/or "Drakensberg melting-pot" fit the

data better than "Southerly stepping stone" for all but the best tree, but relative fit of the models

was somewhat sensitive to phylogenetic uncertainty (Table 1). By contrast, the positions of areas

relative to one another, and the similarities in their realised climatic niche, were consistently

prominent in our results. Of the distance models, the combination of geographical and ecological

distance fit the data well, and given the phylogenetic uncertainty, often better than either of these

factors individually or the stepping stone models. The generally better fit of the combined geographic

and realised niche model affirms the concerted importance of both factors in shaping distributional

patterns of plants (Donoghue 2008; Donoghue and Edwards, 2014). Of the nine range expansion

events that we inferred (DEC+J, best tree, best model), seven respectively were between adjacent

areas or between areas with similar environmental conditions (where "similar" is arbitrarily defined

as a pairwise Schoener's D > 0.5; Fig. 2). Overall, this represents striking evidence for geographical

and ecological distance constraining past and present distributions of *Erica* species, similar to that

inferred for other Mediterranean climate plant groups (Skeels & Cardillo, 2017). Irrespective of

model fit, the sequence of dispersal events that we inferred from ancestral area reconstructions,

based on both the set of best fitting parametric models and a parsimonious interpretation of the

infrequent dispersal events (Fig. 2), does resemble a "Drakensberg melting-pot" scenario. The

Drakensberg acted as a sink for dispersals from the adjacent Cape and Tropical African regions, but

not as a stepping stone (or indeed a "springboard"; Galley & al., 2007).

Cape lineages found in the Drakensberg have not dispersed to Tropical Africa, and neither have

Tropical Africa lineages found in the Drakensberg dispersed further to the Cape. This is unexpected,

not only because of the low distances and high niche similarities involved, but also because of the

equivalent events inferred in other similarly distributed plant groups (Galley et al., 2007). Striking in a

different way are three unique events: the single dispersals from Europe to Tropical Africa, out of

Tropical Africa to the Cape, and out of Tropical Africa to Madagascar, which were each over much

longer distances. The dispersals to Tropical Africa and to Madagascar both might have involved shifts

in realised niches (indicated by low Schoener's D values of 0.298 and 0.274 respectively); that to the

Cape, borderline so (Schoener's D of 0.560; Fig. 2). Notably, the dispersals to tropical Africa and to

the Cape coincided with clear increases in diversification rate (Pirie et al., 2016).

Potential explanations for these apparent exceptions to the general importance of geographical and

ecological distance might be found in the context of the changing climates and geology of the African

continent during the timeframe of the *Erica* radiation. The summer-arid climate of the present day

Cape has been linked to the establishment of the cold Benguela current off the south-west African

coast in the mid Miocene 14-10 Ma (Marlow et al., 2000; Zachos et al., 2001). Evidence from pollen

deposited in nearby marine sediments shows an accumulation of typical Cape lineages since roughly

the same time, including Ericaceae (Dupont et al., 2011), supported by further evidence from recent

dated phylogenies both for the ages of clades in the Cape (e.g. Verboom & al., 2009; Hoffmann & al.,

2015) and the origins of fire adapted lineages (Bytebier et al., 2011). The gradual change from a more

tropical to a mesic flora and initiation of a regular fire regime in south-western Africa might be

ecological changes important for the establishment of *Erica* in the Cape. Whilst the mountains of the

Western Cape, home to much of the *Erica*-dominated fynbos vegetation, long predate Miocene

climatic changes, the origins of the Drakensberg and Tropical African high mountains, *Erica's* area of

first establishments in Africa, are more recent, with uplift in these regions creating montane habitats
from the Miocene onwards (McCarthy & Rubidge, 2005).

Thus, shifting climates and mountain building created an archipelago of temperate islands across
sub-Saharan Africa that were available for colonisation by plants able to tolerate the novel
conditions. These included *Erica* species, which had begun to diversify c. 30 Ma in the Northern
Hemisphere (Pirie et al., 2016), and which as a clade are characterised by drought resistant leaves
(Van Wilgen, Higgins & Bellstedt, 1990) as well as adaptations to post-fire regeneration (Ojeda,
1998). However, our analyses of the realised climatic niches of *Erica* species in their different biomes
suggests that despite these pre-existing drought and fire adaptations, colonisation of new areas by
Erica involved further adaptation to rather different climatic conditions, as inferred for tropical alpine
Hypericum in South America (Nürk, Michling & Linder, 2018) and hypothesised for tropical high
alpine plants in general (Gehrke, 2018). In this context, biome shifts and increased diversification
rates may be linked: the open field presented by these newly formed, isolated, temperate habitats
may have facilitated both the chance establishment of suboptimally adapted plants and their
subsequent *in situ* shift into new adaptive zones, promoting accelerated diversification.

Neither differences in ecological nor geographic distance present an obvious explanation for why
dispersal to the Drakensberg was not followed by further independent colonisations, particularly of
the Cape. One possibility could be that within the Drakensberg, Cape and Tropical African elements
occupy somewhat differing niches: the former, such as the widespread Cape-Drakensberg species *E.*
cerinthoides and *E. caffra* predominantly at lower elevations, the latter at higher elevations under
conditions differing more to those in the Cape. Another could be niche pre-emption (Silvertown,
2004), whereby the single colonisation and rapid species radiation of Cape *Erica* prevented further
colonisation by similar competitors.

Widespread species such as *E. cerinthoides* and *E. caffra*, found in the Cape and Drakensberg, and *E.*
arborea, found in Europe and Tropical Africa (Désamoré & al., 2011; Gizaw & al., 2013), are
exceptional in *Erica*. Almost all extant species are restricted to just one of the areas as defined here
(including species such as *E. silvatica* and *E. benguelensis* that are widespread across disjunct areas
within Tropical Africa), and species radiations leading to most of the present day diversity of *Erica*
were within single areas (Pirie & al., 2016). This suggests that species distributions were restricted
throughout the evolution of the *Erica* African/Madagascan clade, and that the areas remained
isolated during this period (i.e. the last c. 15 Ma; Pirie & al., 2016). We would also argue that it lends
credibility to results obtained under DEC+J, in which some range shifts were treated as cladogenetic
dispersal events (instead of by inferring seemingly implausible widespread distributions), despite
arguable drawbacks in the implementation of that model (Ree & Sanmartín, 2018).

Nevertheless, the extent and position of suitable habitats across the Afrotemperate shifted

considerably during this timeframe, and the implicit assumption of our analyses, that they can be

treated as consistent during the *Erica* diversification, is a considerable simplification. This may not

impact the results of a broader scale analysis such as the one we present here, but might be

important in the context of diversifications within regions, such as those within the Cape (Linder,

2003; Dupont et al., 2011; Schnitzler et al., 2012; Hoffmann, Verboom & Cotterill, 2015) or

Drakensberg (Bentley, Verboom & Bergh, 2014). To assess the impact of climatic changes on the

dramatic radiation of Cape *Erica*, for example, it would be important to translate realised niches into

past distributions to model the shifting extents and interconnectedness of populations through time

(cf. Mairal & al., 2017).

Conclusions: The overall picture to be gleaned from the colonisation history of *Erica* across the

Afrotemperate is one of infrequent dispersal limited by geographic distance and ecological similarity.

Lack of dispersals where they might be expected – in the case of *Erica*, the Drakensberg acting as a

sink, rather than stepping stone to wider dispersal – can point to biological and historical

idiosyncrasies of particular lineages. Our results also show the importance of single unique events

that can run counter to general trends. In *Erica*, three particularly long distance dispersals, two

potentially with shifts in the realised niche, were followed by species radiations – most notably in the

Cape – that dominate the narrative of the group as a whole. Our results serve to further emphasise

the importance of such rare events, in which unique biome shifts fuel dramatic evolutionary

radiations.

Acknowledgements: We thank J. Fagúndez, A. Hitchcock, R. Turner, M. Muasya, C. Stirton, R. Clark, B.

Bytebier, M. Pimentel, F. Ojeda, C. Merry, and many others for providing samples and Cape Nature

and South Africa National Parks for assistance with permits. We also gratefully acknowledge the

computing time granted on the supercomputer Mogon at Johannes Gutenberg University Mainz

(www.hpc.uni-mainz.de), and F. Michling for providing R code. Invaluable constructive comments on

the previous preprint version of this paper was provided by Andrea Meseguer, Simon Joly, Florian

Boucher, and two anonymous reviewers. Funding was provided by the South African National

Research Foundation (NRF; DUB and MDP); a postdoctoral fellowship from the Claude Leon

Foundation (MDP); DFG (PI1169/1-1, PI1169/1-2, and PI1169/2-1 to MDP); and the Ministerium für

Klimaschutz, Umwelt, Landwirtschaft, Natur- und Verbraucherschutz des Landes Nordrhein-

Westfalen, the Faculty of Agriculture Lehr- und Forschungsschwerpunkt „Umweltverträgliche und

Standortgerechte Landwirtschaft", Bonn University; and the Landgard foundation (AMK). Any

opinion, finding and conclusion or recommendation expressed in this material is that of the authors
and the NRF does not accept liability in this regard.

Author contributions': DUB, MDP & EGHO: conceived the research; NCM, AM, MDP, BG, EGHO &
DUB: generated data; MDP, MK & NMN: designed analytical approach; MK: performed BioGeoBEARS
analyses; NMN: performed GIS-based environmental analyses; MDP: led the writing (to which all
authors contributed).

References

Alsos IG., Eidesen PB., Ehrich D., Skrede I., Westergaard K., Jacobsen GH., Landvik JY., Taberlet P.,
Brochmann C. 2007. Frequent long-distance plant colonization in the changing Arctic. *Science*
316:1606–9. DOI: 10.1126/science.1139178.
Anacker BL., Strauss SY. 2014. The geography and ecology of plant speciation: range overlap and
niche divergence in sister species. *Proceedings of the Royal Society B: Biological Sciences*
281:20132980–20132980. DOI: 10.1098/rspb.2013.2980.
Bellstedt DU., Galley C., Pirie MD., Linder HP. 2012. The Migration of the Palaeotropical Arid Flora:
Zygophylloideae as an Example. *Systematic Botany* 37:951–959. DOI:
10.1600/036364412X656608.
Bentley J., Verboom G., Bergh NG. 2014. Erosive processes after tectonic uplift stimulate vicariant
and adaptive speciation: evolution in an Afrotemperate-endemic paper daisy genus. *BMC*
Evolutionary Biology 14:27. DOI: 10.1186/1471-2148-14-27.
Broennimann O., Fitzpatrick MC., Pearman PB., Petitpierre B., Pellissier L., Yoccoz NG., Thuiller W.,
Fortin MJ., Randin C., Zimmermann NE., Graham CH., Guisan A. 2012. Measuring ecological
niche overlap from occurrence and spatial environmental data. *Global Ecology and*
Biogeography 21:481–497. DOI: 10.1111/j.1466-8238.2011.00698.x.
Broennimann O., Petitpierre B., Randin C., Engler R., Breiner F., D'Amen M., Pellissier L., Pottier J., Pio
D., Mateo RG. 2014. ecospat: Spatial ecology miscellaneous methods. *R package version* 1.
Burnham K., Anderson D. 2002. *Model Selection and Multimodel Inference*. New York, Berlin,
Heidelberg: Springer Science & Business Media.
Bytebier B., Antonelli A., Bellstedt DU., Linder HP. 2011. Estimating the age of fire in the Cape flora of
South Africa from an orchid phylogeny. *Proceedings of the Royal Society B: Biological Sciences*
278:188–195. DOI: 10.1098/rspb.2010.1035.
Carvajal-Endara S., Hendry AP., Emery NC., Davies TJ. 2017. Habitat filtering not dispersal limitation

shapes oceanic island floras: species assembly of the Galápagos archipelago. *Ecology Letters*

20:495–504. DOI: 10.1111/ele.12753.

Chala D., Zimmermann NE., Brochmann C., Bakkestuen V. 2017. Migration corridors for alpine plants

among the 'sky islands' of eastern Africa: do they, or did they exist? *Alpine Botany* 127:133–

144. DOI: 10.1007/s00035-017-0184-z.

Crisp MD., Arroyo MTK., Cook LG., Gandolfo MA., Jordan GJ., McGlone MS., Weston PH., Westoby

M., Wilf P., Linder HP. 2009. Phylogenetic biome conservatism on a global scale. *Nature*

458:754–756. DOI: 10.1038/nature07764.

Désamoré A., Laenen B., Devos N., Popp M., González-Mancebo JM., Carine MA., Vanderpoorten A.

2011. Out of Africa: north-westwards Pleistocene expansions of the heather *Erica arborea*.

Journal of Biogeography 38:164–176. DOI: 10.1111/j.1365-2699.2010.02387.x.

Donoghue MJ. 2008. Colloquium paper: a phylogenetic perspective on the distribution of plant

diversity. *Proceedings of the National Academy of Sciences of the United States of America*

105:11549–55. DOI: 10.1073/pnas.0801962105.

Donoghue MJ., Edwards EJ. 2014. Biome Shifts and Niche Evolution in Plants. *Annual Review of*

Ecology, Evolution, and Systematics 45:547–572. DOI: 10.1146/annurev-ecolsys-120213-

091905.

Dray S., Dufour AB. 2007. The ade4 Package: Implementing the Duality Diagram for Ecologists.

Journal of Statistical Software 22:1–20. DOI: 10.1.1.177.8850.

Drummond CS., Eastwood RJ., Miotto STS., Hughes CE. 2012. Multiple Continental Radiations and

Correlates of Diversification in Lupinus (Leguminosae): Testing for Key Innovation with

Incomplete Taxon Sampling. *Systematic Biology* 61:443–460. DOI: 10.1093/sysbio/syr126.

Dupont LM., Linder HP., Rommerskirchen F., Schefuß E. 2011. Climate-driven rampant speciation of

the Cape flora. *Journal of Biogeography* 38:1059–1068. DOI: 10.1111/j.1365-

2699.2011.02476.x.

Fernández-Palacios JM., De Nascimento L., Otto R., Delgado JD., García-Del-Rey E., Arévalo JR.,

Whittaker RJ. 2011. A reconstruction of Palaeo-Macaronesia, with particular reference to the

long-term biogeography of the Atlantic island laurel forests. *Journal of Biogeography* 38:226–

246. DOI: 10.1111/j.1365-2699.2010.02427.x.

Galbany-Casals M., Unwin M., Garcia-Jacas N., Smissen RD., Susanna A., Bayer RJ. 2014. Phylogenetic

relationships in Helichrysum (Compositae: Gnaphalieae) and related genera: Incongruence

between nuclear and plastid phylogenies, biogeographic and morphological patterns, and

implications for generic delimitation. *Taxon* 63:608–624. DOI: 10.12705/633.8.

Galley CC., Bytebier B., Bellstedt DU., Linder HP. 2007. The Cape element in the Afrotemperate flora:

from Cape to Cairo? *Proceedings of the Royal Society B: Biological Sciences* 274:535–543. DOI:

10.1098/rspb.2006.0046.

García-Aloy S., Vitales D., Roquet C., Sanmartín I., Vargas P., Molero J., Kamau P., Aldasoro JJ.,

Alarcón M. 2017. North-west Africa as a source and refuge area of plant biodiversity: a case

study on Campanula kremeri and Campanula occidentalis. *Journal of Biogeography* 44:2057–

2068. DOI: 10.1111/jbi.12997.

Gehrke B. 2018. Staying cool: preadaptation to temperate climates required for colonising tropical

alpine-like environments. *PhytoKeys* in press:111. DOI: 10.3897/phytokeys.96.13353.

Gehrke B., Linder HP. 2009. The scramble for Africa: pan-temperate elements on the African high

mountains. *Proceedings of the Royal Society B: Biological Sciences* 276:2657–2665. DOI:

10.1098/rspb.2009.0334.

Gizaw A., Brochmann C., Nemomissa S., Wondimu T., Masao CA., Tusiime FM., Abdi AA., Oxelman B.,

Popp M., Dimitrov D. 2016. Colonization and diversification in the African 'sky islands': insights

from fossil-calibrated molecular dating of *Lychnis* (Caryophyllaceae). *New Phytologist* 211:719–

734. DOI: 10.1111/nph.13937.

Gizaw A., Kebede M., Nemomissa S., Ehrich D., Bekele B., Mirré V., Popp M., Brochmann C. 2013.

Phylogeography of the heathers *Erica arborea* and *E. trimera* in the afro-alpine 'sky islands'

inferred from AFLPs and plastid DNA sequences. *Flora - Morphology, Distribution, Functional

Ecology of Plants* 208:453–463. DOI: 10.1016/j.flora.2013.07.007.

Higgins SI., Richardson DM. 1999. Predicting Plant Migration Rates in a Changing World: The Role of

Long-Distance Dispersal. *The American Naturalist* 153:464–475. DOI: 10.1086/303193.

Hijmans RJ. 2015. raster: Geographic data analysis and modeling. [R package].

Hoffmann V., Verboom GA., Cotterill FPD. 2015. Dated Plant Phylogenies Resolve Neogene Climate

and Landscape Evolution in the Cape Floristic Region. *PLOS ONE* 10:e0137847. DOI:

10.1371/journal.pone.0137847.

Jordano P. 2017. What is long-distance dispersal? And a taxonomy of dispersal events. *Journal of

Ecology* 105:75–84. DOI: 10.1111/1365-2745.12690.

Kadereit JW., Baldwin BG. 2012. Western Eurasian-western North American disjunct plant taxa: The

dry-adapted ends of formerly widespread North temperate mesic lineages-and examples of

long-distance dispersal. *Taxon* 61:3–17.

Kim SC., McGowen MR., Lubinsky P., Barber JC., Mort ME., Santos-Guerra A. 2008. Timing and tempo

of early and successive adaptive radiations in Macaronesia. *PLoS ONE* 3:1–7. DOI:

10.1371/journal.pone.0002139.

Knox EB., Palmer JD. 1998. Chloroplast DNA evidence on the origin and radiation of the giant lobelias

in eastern Africa. *Systematic Botany* 23:109–149.

Kowalski R., Fagúndez J. 2017. *Maiella miocaenica* gen. et sp. nov., a New Heather Genus (Ericeae,

Ericaceae) from the Central European Miocene. *International Journal of Plant Sciences* 178:411–
420. DOI: 10.1086/691461.

Linder HP. 2003. The radiation of the Cape flora, southern Africa. *Biological Reviews* 78:597–638.
DOI: Doi 10.1017/S1464793103006171.

Linder HP., Antonelli A., Humphreys AM., Pirie MD., Wüest RO. 2013. What determines
biogeographical ranges? Historical wanderings and ecological constraints in the danthonioid
grasses. *Journal of Biogeography* 40:821–834. DOI: 10.1111/jbi.12070.

Maddison WP., Maddison DR. 2006. Mesquite: a modular system for evolutionary analysis.

Mairal M., Sanmartín I., Pellissier L. 2017. Lineage-specific climatic niche drives the tempo of
vicariance in the Rand Flora. *Journal of Biogeography* 44:911–923. DOI: 10.1111/jbi.12930.

Marlow JR., Lange CB., Wefer G., Rosell-Mele A. 2000. Upwelling intensification as part of the
Pliocene-Pleistocene climate transition. *Science* 290:2288. DOI: 10.1126/science.290.5500.2288.

Matzke NJ. 2013. Probabilistic Historical Biogeography: New Models for Founder-Event Speciation,
Imperfect Detection, and Fossils Allow Improved Accuracy and Model- Testing. University of
California, Berkeley.

Matzke NJ. 2014. Model Selection in Historical Biogeography Reveals that Founder-Event Speciation
Is a Crucial Process in Island Clades. *Systematic Biology* 63:951–970. DOI:
10.1093/sysbio/syu056.

McCarthy T., Rubidge B. 2005. *The story of earth & life: a southern African perspective on a 4.6*
billion-year journey. Cape Town: Struik Publishers.

McGuire AF., Kron KA. 2005. Phylogenetic relationships of European and African ericas. *International*
Journal of Plant Sciences 166:311–318. DOI: 10.1086/427478.

Meeus J. 1999. *Astronomical algorithms*. Willman-Bell.

Merckx VSFT., Hendriks KP., Beentjes KK., Mennes CB., Becking LE., Peijnenburg KTCA., Afendy A.,
Arumugam N., De Boer H., Biun A., Buang MM., Chen PP., Chung AYC., Dow R., Feijen FAA.,
Feijen H., Soest CF Van., Geml J., Geurts R., Gravendeel B., Hovenkamp P., Imbun P., Ipor I.,
Janssens SB., Jocqué M., Kappes H., Khoo E., Koomen P., Lens F., Majapun RJ., Morgado LN.,
Neupane S., Nieser N., Pereira JT., Rahman H., Sabran S., Sawang A., Schwallier RM., Shim PS.,
Smit H., Sol N., Spait M., Stech M., Stokvis F., Sugau JB., Suleiman M., Sumail S., Thomas DC.,
Van Tol J., Tuh FYY., Yahya BE., Nais J., Repin R., Lakim M., Schilthuizen M. 2015. Evolution of
endemism on a young tropical mountain. *Nature* 524:347–350. DOI: 10.1038/nature14949.

Míguez M., Gehrke B., Maguilla E., Jiménez-Mejías P., Martín-Bravo S. 2017. Carex sect.
Rhynchocystis (Cyperaceae): a Miocene subtropical relict in the Western Palaearctic showing a
dispersal-derived Rand Flora pattern. *Journal of Biogeography* 44:2211–2224. DOI:
10.1111/jbi.13027.

Mugrabi de Kuppler AL., Fagúndez J., Bellstedt DU., Oliver EGH., Léon J., Pirie MD. 2015. Testing
reticulate versus coalescent origins of *Erica lusitanica* using a species phylogeny of the northern
heathers (Ericeae, Ericaceae). *Molecular Phylogenetics and Evolution* 88:121–131. DOI:
10.1016/j.ympev.2015.04.005.

Muñoz J., Felicísimo AM., Cabezas F., Burgaz AR., Martínez ILB-4319. 2004. Wind as a long-distance
dispersal vehicle in the Southern Hemisphere. *Science* 304:1144–1147.

Nakazato T., Warren DL., Moyle LC. 2010. Ecological and geographic modes of species divergence in
wild tomatoes. *American Journal of Botany* 97:680–693. DOI: 10.3732/ajb.0900216.

Nathan R. 2006. Long-Distance Dispersal of Plants. *Science* 313:786–788. DOI:
10.1126/science.1124975.

Nürk NM., Michling F., Linder HP. 2018. Are the radiations of temperate lineages in tropical alpine
ecosystems pre-adapted? *Global Ecology and Biogeography* 27:334–345. DOI:
10.1111/geb.12699.

Nürk NM., Uribe-Convers S., Gehrke B., Tank DC., Blattner FR. 2015. Oligocene niche shift, Miocene
diversification - cold tolerance and accelerated speciation rates in the St. John's Worts
(Hypericum, Hypericaceae). *BMC evolutionary biology* 15:80. DOI: 10.1186/s12862-015-0359-4.

Ojeda F. 1998. Biogeography of seeder and resprouter Erica species in the Cape Floristic Region—
Where are the resprouters? *Biological Journal of the Linnean Society* 63:331–347. DOI:
10.1111/j.1095-8312.1998.tb01521.x.

Oliver EGH. 2012. Ericaceae. In: Manning JC, Goldblatt P eds. *Plants of the Greater Cape Floristic
Region, Volume 1: The Core Cape Flora*. Strelitzia. Pretoria: South African National Biodiversity
Institute (SANBI Publishing), 482—511.

Paradis E., Claude J., Strimmer K. 2004. APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics 20:289–290. DOI: 10.1093/bioinformatics/btg412.

Pirie MD., Humphreys AM., Antonelli A., Galley C., Linder HP. 2012. Model uncertainty in ancestral
area reconstruction: A parsimonious solution? *Taxon* 61:652–664. DOI: 10.5167/uzh-64515.

Pirie MD., Oliver EGH., Bellstedt DU. 2011. A densely sampled ITS phylogeny of the Cape flagship
genus Erica L. suggests numerous shifts in floral macro-morphology. *Molecular Phylogenetics
and Evolution* 61:593–601. DOI: 10.1016/j.ympev.2011.06.007.

Pirie MD., Oliver EGH., Mugrabi de Kuppler A., Gehrke B., Le Maitre NC., Kandziora M., Bellstedt DU.
2016. The biodiversity hotspot as evolutionary hot-bed: spectacular radiation of Erica in the
Cape Floristic Region. *BMC Evolutionary Biology* 16:190. DOI: 10.1186/s12862-016-0764-3.

Pokorny L., Riina R., Mairal M., Meseguer AS., Culshaw V., Cendoya J., Serrano M., Carbajal R., Ortiz
S., Heuertz M., Sanmartín I. 2015. Living on the edge: timing of Rand Flora disjunctions
congruent with ongoing aridification in Africa. *Frontiers in Genetics* 6:154. DOI:

10.3389/fgene.2015.00154.

R Development Core Team. 2013. R: A language and environment for statistical computing.

Ree RH., Moore BR., Webb CO., Donoghue MJ. 2005. A likelihood framework for inferring the

581   evolution of geographic range on phylogenetic trees. *Evolution; international journal of organic*

582   *evolution* 59:2299–2311. DOI: 10.1554/05-172.1.

583   Ree RH., Sanmartín I. 2018. Conceptual and statistical problems with the DEC+J model of founder-

584   event speciation and its comparison with DEC via model selection. *Journal of Biogeography*.

585   DOI: 10.1111/jbi.13173.

586   Ree RH., Smith SA. 2008. Maximum likelihood inference of geographic range evolution by dispersal,

587   local extinction, and cladogenesis. *Systematic biology* 57:4–14. DOI:

588   10.1080/10635150701883881.

589   Ruhfel BR., Bove CP., Philbrick CT., Davis CC. 2016. Dispersal largely explains the gondwanan

590   distribution of the ancient tropical clusioid plant clade. *American Journal of Botany* 103:1117–

591   1128. DOI: 10.3732/ajb.1500537.

592   Sanmartín I., Anderson CL., Alarcon M., Ronquist F., Aldasoro JJ. 2010. Bayesian island biogeography

593   in a continental setting: the Rand Flora case. *Biology Letters* 6:703–707. DOI:

594   10.1098/rsbl.2010.0095.

595   Schnitzler J., Graham CH., Dormann CF., Schiffers K., Peter Linder H. 2012. Climatic niche evolution

596   and species diversification in the Cape flora, South Africa. *Journal of Biogeography* 39:2201–

597   2211. DOI: 10.1111/jbi.12028.

598   Schoener TW. 1968. The Anolis Lizards of Bimini: Resource Partitioning in a Complex Fauna. *Ecology*

599   49:704–726. DOI: 10.2307/1935534.

600   Schwery O., Onstein RE., Bouchenak-Khelladi Y., Xing Y., Carter RJ., Linder HP. 2015. As old as the

601   mountains: The radiations of the Ericaceae. *New Phytologist* 207:355–367. DOI:

602   10.1111/nph.13234.

603   Silvertown J. 2004. The Ghost of Competition Past in the Phylogeny of Island Endemic Plants. *J. Ecol.*

604   92:168–173. DOI: 10.1111/j.1365-2745.2004.00853.x.

605   Skeels A., Cardillo M. 2017. Environmental niche conservatism explains the accumulation of species

606   richness in Mediterranean-hotspot plant genera. *Evolution*:582–594. DOI: 10.1111/evo.13179.

607   Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

608   thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690. DOI:

609   10.1093/bioinformatics/btl446.

610   Tamura K., Battistuzzi FU., Billing-Ross P., Murillo O., Filipski A., Kumar S. 2012. Estimating divergence

611   times in large molecular phylogenies. *Proceedings of the National Academy of Sciences of the*

612   *United States of America* 109:19333–19338. DOI: 10.1073/pnas.1213199109.

Uribe-Convers S., Tank DC. 2015. Shifts in diversification rates linked to biogeographic movement
into new areas: An example of a recent radiation in the Andes. *American Journal of Botany*
102:1854–1869. DOI: 10.3732/ajb.1500229.

Verboom GA., Archibald JK., Bakker FT., Bellstedt DU., Conrad F., Dreyer LL., Forest FF., Galley CC.,
Goldblatt P., Henning JF., Mummenhoff K., Linder HP., Muasya AM., Oberlander KC., Savolainen
V., Snijman DA., Niet T van der T van der., Nowell TL. 2009. Origin and diversification of the
Greater Cape flora: Ancient species repository, hot-bed of recent radiation, or both? *Molecular*
Phylogenetics and Evolution 51:44–53. DOI: 10.1016/j.ympev.2008.01.037.

Verdcourt B. 1969. The arid corridor between the North-East and South-West areas of Africa. In: van
Zinderen Bakker EM ed. *Paleoecology of Africa*. Cape Town: Balkema, 140–144.

Viana DS., Gangoso L., Bouten W., Figuerola J. 2015. Overseas seed dispersal by migratory birds.
Proceedings of the Royal Society B: Biological Sciences 283:20152406. DOI:
dx.doi.org/10.1098/rspb.2015.2406.

White F. 1981. The history of the Afromontane archipelago and the scientific need for its
conservation. *African Journal of Ecology* 19:33–54. DOI: 10.1111/j.1365-2028.1981.tb00651.x.

White F. 1983. The vegetation of Africa: a descriptive memoir to accompany the
UNESCO/AETFAT/UNSO vegetation map of Africa. *Natural Resources Research* 20:1–356.

Van Wilgen BW., Higgins KB., Bellstedt DU. 1990. The role of vegetation structure and fuel chemistry
in excluding fire from forest patches in the fire-prone fynbos shrublands of South Africa. *The*
Journal of Ecology 78:210–222. DOI: 10.2307/2261046.

Zachos J., Pagani M., Sloan L., Thomas E., Billups K. 2001. Trends, rhythms, and aberrations in global
climate 65 Ma to present. *Science* 292:686–693. DOI: 10.1126/science.1059412.

Figures

Fig. 1: Biogeographic hypotheses. The pure geographic distance model; niche similarity, implying

colonisation of areas with the most similar climatic niche (Donoghue 2008); these together

constituting a combined geographic and niche similarity model; stepping stone: stepwise southerly

colonisation of the Afrotemperate from Europe (following McGuire and Kron, 2005); Cape to Cairo:

stepwise northerly colonisation of the Afrotemperate from the Cape (following Galley et al., 2007);

necessarily preceded by LDD from Europe; and Drakensberg melting pot: the Drakensberg colonised

by both southerly and northerly stepwise dispersal.

Fig. 2: Biogeographic scenario. A) Inferred dispersal scenario between the biogeographic areas

(colour coded "area ranges" as assessed by the buffered species occurrence data) depicted on the

global study area. **B**) The phylogeny of *Erica* with a representation of ancestral areas derived from a

single BSM analysis using the best tree and model (under DEC+J). **C**) Climate similarity between area

ranges given as hypervolume corrected pairwise climate similarity (Schoener's D; with $D \geq 0.5$ in dark

grey, and $D < 0.5$ in light grey; Table S8) with superimposed black arrows scaled to the dispersal rates

per Ma inferred between areas (Table S13). Ma, million years.

Table 1: Best fitting biogeographic models given the best tree (DEC+J and DEC) and nine selected

bootstrap trees (DEC+J). Dispersal multipliers are indicated where relevant, as are the Log likelihood

(LnL), Akaike Information Criterion (AIC), and overall deltaAIC scores for models. Models with

deltaAIC of 0 are indicated in bold type. DMP=Drakensberg mekting-pot; Dist=Distance;

SSS=Southerly stepping-stone; CtoC=Cape to Cairo

Tree	Model	Dispersal multiplier	LnL	AIC	deltaAIC
Best (DEC+J)	**DMP**	**0.5**	**-62.5**	**131**	**0**
	DMP	**0.1**	**-62.5**	**131**	**0**
	DMP	**0.25**	**-62.5**	**131**	**0**
	DMP	0.75	-62.6	131.2	0.2
	Dist	-	-62.8	131.6	0.58
	SSS	0.5	-63.1	132.3	1.3
	SSS	0.25	-63.3	132.7	1.7
Best (DEC)	**DMP**	**0.75**	**-68.6**	**141.2**	**0**
	DMP	0.5	-68.7	141.3	0.1
BS 0_0	**Niche+Dist**	**-**	**-61.2**	**128.4**	**0**
	CtoC	0.25	-61.7	129.4	1
	CtoC	0.1	-62.1	130.1	1.7
	Dist	-	-62.2	130.4	2
0_1	**Niche+Dist**	**-**	**-65.6**	**137.1**	**0**
	CtoC	0.1	-65.8	137.6	0.5
	CtoC	0.25	-66.1	138.3	1.2
0_2	**Niche+Dist**	**-**	**-60.2**	**126.3**	**0**
	CtoC	0.25	-60.4	126.7	0.4
	CtoC	0.1	-60.4	126.9	0.6
	Dist	-	-61	127.9	1.6
1_0	**CtoC**	**0.1**	**-58.7**	**123.5**	**0**
	CtoC	0.25	-59.5	124.9	1.4
1_1	**Niche+Dist**	**-**	**-61.7**	**129.4**	**0**
	Niche	-	-62.6	131.2	1.8
1_2	**Niche+Dist**	**-**	**-55.9**	**117.9**	**0**
	CtoC	0.25	-56.5	119	1.1
	CtoC	0.1	-56.9	119.7	1.8
	Dist	-	-56.9	119.7	1.8
2_0	**Niche+Dist**	**-**	**-62.3**	**130.6**	**0**
	CtoC	0.25	-62.6	131.2	0.6
	Dist	-	-62.8	131.5	0.9

	CtoC	0.1	-62.9	131.8	1.2
	CtoC	0.5	-63.1	132.3	1.7
	Niche	-	-63.2	132.4	1.8
2_1	**Dist**	**-**	**-56.5**	**119.1**	**0**
	DMP	0.1	-56.6	119.2	0.1
	DMP	0.25	-56.6	119.2	0.1
	DMP	0.5	-57	120	0.9
	CtoC	0.5	-57.2	120.4	1.3
	Niche+Dist	-	-57.2	120.4	1.3
	CtoC	0.25	-57.5	121	1.9
	Niche	-	-57.6	121.1	2
2_2	**Dist**	**-**	**-65.3**	**135.6**	**0**
	Niche+Dist	-	-65.2	136.4	0.8
	DMP	0.25	-65.5	137	1.4
	CtoC	0.5	-65.5	137	1.4
	DMP	0.1	-65.5	137.1	1.5
	CtoC	0.25	-65.6	137.1	1.5
	Niche	-	-65.8	137.5	1.9
	DMP	0.5	-65.8	137.6	2

Appendices

Appendix 1: Methods: occurrence data

Appendix 2: Methods: Global environmental space, area ranges, and climate similarity analysis

Appendix 3: Methods: Biogeographic models; example files for BioGeoBEARS analyses

Appendix 4: Methods: Selected bootstrap trees used to represent phylogenetic uncertainty
between geographically restricted major clades

Appendix 5: Methods: Mesquite file used for parsimony ancestral state reconstruction including
RAXML bootstrap trees

Appendix 6: Results: pairwise climate similarity (Schoener's *D*) between biogeographic areas per PC
axis

Appendix 7: Results: Pairwise climate similarity (Schoener's *D*) between biogeographic areas for
combined PC axes

Appendix 8: Results of the different models under DEC+J and DEC (generally the better models
compared to DIVA-like and BAYAREA-like-models)

Appendix 9: Results: Summary of event counts from 50 biogeographical stochastic mappings under
the best inferred model using the best tree

Appendix 10: Results: Number of range-expansion dispersal events (mean and standard deviation
of all observed "d" dispersals) averaged across 50 biogeographical stochastic mappings under the
best inferred model using the best tree.

Appendix 11: Results: Number of cladogenetic dispersal events (mean and standard deviation of all
observed jump 'j' dispersals) averaged from 50 biogeographical stochastic mappings under the best
inferred model using the best tree.

Appendix 12: Results: Number of all dispersal events (mean and standard deviation of all observed
anagenetic 'a', 'd' dispersals, PLUS cladogenetic founder/jump dispersal) averaged from 50
biogeographical stochastic mappings under the best inferred model using the best tree.

Appendix 13: Results: Ancestral area reconstructions inferred using BioGeoBEARS given the best
tree under the best fitting model given A: DEC+J; B: DEC; and without range or dispersal constraint:
C: DEC+J; D: DEC. For each model the single most probable state is shown first (boxes with areas at
nodes) followed by the relative probability of each state represented with pie charts at nodes.
Areas are represented by colours: Dark blue for Europe (E); green for Tropical Africa (T); yellow for

Madagascar (M); light blue for Drakensberg (D); red for Cape (C); and further colours for

widespread distributions as indicated in the legends.

Longer distance /
dissimilar niche
(dispersal multiplier
[DM]: 0.0-0.4)

Intermediate distance /
niche similarity
(DM: 0.4-0.7)

Short distance /
 similar niche (DM: 0.7-1.0)

Geographic distance

Niche similarity

Cape to Cairo

Southerly stepping stone

Drakensberg melting pot

A

B

C

Europe

Tropical Africa

Cape

Madagascar

Drakensberg

60   50   40   30   20   10   0 Ma

2000 km

0°