Discovery of biased orientation of human DNA motif sequences

affecting enhancer-promoter interactions and transcription of genes

Naoki Osato[1][*]

[1]Department of Bioinformatic Engineering, Graduate School of Information Science

and Technology, Osaka University, Osaka 565-0871, Japan

[*]Corresponding author

E-mail address: naokiosato11@gmail.com, nosato@ist.osaka-u.ac.jp

Abstract

Chromatin interactions have important roles for enhancer-promoter interactions

(EPI) and regulating the transcription of genes. CTCF and cohesin proteins are located

at the anchors of chromatin interactions, forming their loop structures. CTCF has

insulator function limiting the activity of enhancers into the loops. DNA binding

sequences of CTCF indicate their orientation bias at chromatin interaction anchors –

forward-reverse (FR) orientation is frequently observed. DNA binding sequences of

CTCF were found in open chromatin regions at about 40% - 80% of chromatin

interaction anchors in Hi-C and in situ Hi-C experimental data. It has been reported that

long range of chromatin interactions tends to include less CTCF at their anchors. It is

still unclear what proteins are associated with chromatin interactions.

To find DNA binding motif sequences of transcription factors (TF) such as CTCF

affecting the interaction between enhancers and promoters of transcriptional target

genes and their expression, here, I predicted human transcriptional target genes of TF

bound in open chromatin regions in enhancers and promoters in monocytes and other

cell types using experimental data in public database. Transcriptional target genes were

predicted based on enhancer-promoter association (EPA). EPA was shortened at the

genomic locations of FR orientation of DNA binding motifs of TF, which were

supposed to be at chromatin interaction anchors.

The expression level of the target genes predicted based on EPA was compared

with target genes predicted from only promoters. Total 351 biased orientation of DNA

motifs (192 forward-reverse (FR) and 179 reverse-forward (RF) orientation, the reverse

complement sequences of some DNA motifs are also registered in databases, so the total

number is smaller than the number of FR and RF) affected the expression level of

putative transcriptional target genes significantly in monocytes of four people in

common. Moreover, EPI predicted using FR or RF orientation of some DNA motifs

were overlapped with chromatin interaction data (Hi-C) more than the other EPA (Total

62 biased orientation of DNA motifs, 41 FR and 24 RF showed this result).

Keywords: transcriptional target genes, gene expression, transcription factors, enhancer,

enhancer-promoter interactions, chromatin interactions, CTCF, cohesion, open

chromatin regions

Background

Chromatin interactions have important roles for enhancer-promoter interactions

(EPI) and regulating the transcription of genes. CTCF and cohesin proteins are located

at the anchors of chromatin interactions, forming their loop structures. CTCF has

insulator function limiting the activity of enhancers into the loops (Fig. 1A). DNA

binding sequences of CTCF indicate their orientation bias at chromatin interaction

anchors – forward-reverse (FR) orientation is frequently observed (de Wit et al. 2015;

Guo et al. 2015). About 40% - 80% of chromatin interaction anchors of Hi-C and in situ

Hi-C experiments include DNA binding motif sequences of CTCF. However, it has been

reported that long range of chromatin interactions tends to include less CTCF at their

anchors (Jeong et al. 2017). Other DNA binding proteins such as ZNF143, YY1, and

SMARCA4 (BRG1) are found to be associated with chromatin interactions and EPI

(Bailey et al. 2015; Barutcu et al. 2016; Weintraub et al. 2017). CTCF, cohesin, ZNF143,

YY1 and SMARCA4 have other biological functions as well as chromatin interactions

and enhancer-promoter interactions. The DNA binding motif sequences of the

transcription factors are found in open chromatin regions near transcriptional start sites

(TSS) as well as chromatin interaction anchors.

DNA binding motif sequence of ZNF143 was enriched at both chromatin

interaction anchors. ZNF143's correlation with the CTCF-cohesin cluster relies on its

weakest binding sites, found primarily at distal regulatory elements defined by the

'CTCF-rich' chromatin state. The strongest ZNF143-binding sites map to promoters

bound by RNA polymerase II (POL2) and other promoter-associated factors, such as the

TATA-binding protein (TBP) and the TBP-associated protein, together forming a

'promoter' cluster (Bailey et al. 2015).

DNA binding motif sequence of YY1 does not seem to be enriched at both

chromatin interaction anchors (Z-score < 2), whereas DNA binding motif sequence of

ZNF143 is significantly enriched (Z-score > 7; Bailey et al. 2015 Figure 2a). In the

analysis of YY1, to identify a protein factor that might contribute to EPI, (Ji et al. 2015)

performed chromatin immune precipitation with mass spectrometry (ChIP-MS), using

antibodies directed toward histones with modifications characteristic of enhancer and

promoter chromatin (H3K27ac and H3K4me3, respectively). Of 26 transcription factors

that occupy both enhancers and promoters, four are essential based on a CRISPR

cell-essentiality screen and two (CTCF, YY1) are expressed in >90% of tissues

examined (Weintraub et al. 2017). These analyses started from the analysis of histone

modifications of an enhancer mark rather than chromatin interactions. Other protein

factors associated with chromatin interactions may be found from other researches.

As computational approaches, machine-learning analyses to predict chromatin

interactions have been proposed (Schreiber et al. 2017; Zhang et al. 2017). However,

they were not intended to find DNA motif sequences of transcription factors affecting

chromatin interactions, EPI, and the expression level of transcriptional target genes,

which were examined in this study.

DNA binding proteins involved in chromatin interactions are supposed to affect

the transcription of genes in the loop formed by chromatin interactions. In my previous

analyses, the expression level of human putative transcriptional target genes was

affected, according to the criteria of enhancer-promoter association (EPA) (Fig. 1B;

(Osato 2018)). EPI predicted based on EPA shortened at the genomic locations of

forward-reverse orientation of CTCF binding sites affected the expression level of

putative transcriptional target genes the most, compared with the expression level of

transcriptional target genes predicted from only promoters (Fig. 1B). The expression

level tended to be increased in monocytes and CD4$^+$ T cells, implying that enhancers

activated the transcription of genes, and decreased in ES and iPS cells, implying that

enhancers repressed the transcription of genes. These analyses suggested that enhancers

affected the transcription of genes significantly, when EPI were predicted properly.

Other DNA binding proteins involved in chromatin interactions, as well as CTCF,

would locate at chromatin interaction anchors with a pair of biased orientation of DNA

binding motif sequences, affecting the expression level of putative transcriptional target

genes in the loop formed by the chromatin interactions. As experimental issues of the

analyses of chromatin interactions, chromatin interaction data are changed, according to

experimental techniques, depth of DNA sequencing, and even replication sets of the

same cell type. Chromatin interaction data may not be saturated enough to cover all

chromatin interactions. Supposing these characters of DNA binding proteins associated

with chromatin interactions and avoiding experimental issues of the analyses of

chromatin interactions, here I searched for DNA motif sequences of transcription factors

and DNA repeat sequences, affecting EPI and the expression level of putative

transcriptional target genes in monocytes of four people and other cell types without

using chromatin interaction data. Then, putative EPI were compared with chromatin

interaction data.

Results

Search for biased orientation of DNA motif sequences

Transcription factor binding sites (TFBS) were predicted using open chromatin

regions and DNA motif sequences of transcription factors collected from various

databases and journal papers (see Methods). Transcriptional target genes were predicted

using TFBS in promoters and enhancer-promoter association (EPA) shortened at the

genomic locations of DNA binding motif sequences of transcription factors such as

CTCF and cohesins (RAD21 and SMC3). To find DNA motif sequences of transcription

factors, other than CTCF and cohesin, and DNA repeat sequences affecting the

expression level of putative transcriptional target genes, transcriptional target genes of

each transcription factor were predicted based on EPA shortened at the genomic

locations of DNA motif sequences, and the expression level of the putative

transcriptional target genes was compared with those predicted from promoters using

Mann Whiteney U test (p-value < 0.05). The number of transcription factors showing a

significant difference of expression level of putative transcriptional target genes was

counted. To examine whether the orientation of DNA motif sequences shortening EPA

affected the number of transcription factors showing a significant difference of

expression level of putative transcriptional target genes, the number of the transcription

factors was compared among forward-reverse, reverse-forward, and any orientation of

DNA motif sequences shortening EPA, using chi-square test (p-value < 0.05). To avoid

missing DNA motif sequences showing a relatively weak statistical significance by

multiple testing collection, the above analyses were conducted using monocytes of four

people independently, and DNA motif sequences found in monocytes of four people in

common were selected. Total 351 of biased (192 forward-reverse and 179

reverse-forward) orientation of DNA binding motif sequences of transcription factors

have been found in monocytes of four people in common, whereas only one any

orientation (i.e. without considering orientation) of DNA binding motif sequences have

been found in monocytes of four people in common (Fig. 2; Table 1; Supplemental

Table S1). Forward-reverse orientation of DNA motif sequences included CTCF,

cohesin (RAD21 and SMC3), ZNF143 and YY1, which are associated with chromatin

interactions and enhancer-promoter interactions (EPI). SMARCA4 (BRG1) is

associated with topologically associated domain (TAD), which is higher-order

chromatin organization. The DNA binding motif sequences of SMARCA4 was not

registered in the databases used in this study. Forward-reverse orientation of DNA motif

sequences included SMRC2, a member of the same SWI/SNF family of proteins as

SMARCA4. These results suggested that the expression level of putative transcription

target genes of some transcription factors would be different, depending on the genomic

locations (enhancers or promoters) of DNA binding motif sequences of the transcription

factors in monocytes of four people.

To examine whether the biased orientation of DNA binding motif sequences of

transcription factors was observed in other cell types, the same analyses were conducted

in other cell types. However, for other cell types, experimental data of one sample were

available in ENCODE database, so the analyses of DNA motif sequences were

performed by comparing with the result in monocytes of four people. Among the biased

orientation of DNA binding motif sequences found in monocytes, 79, 76, 91, 61, and 59

DNA binding motif sequences were also observed in CD4$^+$ T cells, H1-hESC, iPS,

Huvec and MCF-7 respectively, including CTCF and cohesin (RAD21 and SMC3)

(Table 2; Supplemental Table S2). The scores of DNA binding motif sequences were the

highest in monocytes, and the other cell types showed lower scores. The results of the

analysis of DNA motif sequences in CD20$^+$ B cells and macrophages did not include

CTCF and cohesin, because these analyses can be utilized in cells where the expression

level of putative transcriptional target genes of each transcription factor show a

significant difference between promoters and EPA shortened at the genomic locations of

DNA motif sequences. Some experimental data of a cell did not show a significant

difference between promoters and EPA (Osato 2018).

Instead of DNA binding motif sequences of transcription factors, DNA repeat

sequences were also examined. The expression level of transcriptional target genes

predicted based on EPA shortened at the genomic locations of DNA repeat sequences

was compared with the expression level of transcriptional target genes predicted from

promoters. Three reverse-forward orientation of DNA repeat sequences showed a

significant difference of expression level of putative transcriptional target genes in

monocytes of four people in common (Table 3). Among them, LTR16C DNA repeat

sequence was observed in iPS and H1-hESC.

Co-location of biased orientation of DNA motif sequences

To examine association of the 351 biased (forward-reverse and reverse-forward)

orientation of DNA binding motif sequences of transcription factors. Co-location of the

DNA binding motif sequences in open chromatin regions in monocytes was analyzed.

The number of open chromatin regions with the same pairs of DNA binding motif

sequences was counted, and when the pairs of DNA binding motif sequences were

enriched with statistical significance (chi-square test, p-value $< 1.0 \times 10^{-10}$), they were

listed (Table 4; Supplemental Table S3). Open chromatin regions overlapped with

histone modification of an enhancer mark (H3K27ac) (total 26,095 regions) showed a

larger number of enriched pairs of DNA motifs than all open chromatin regions (Table

4; Supplemental Table S3). H3K27ac is known to be enriched at chromatin interaction

anchors (Phanstiel et al. 2017). As already known, CTCF was found with cohesin such

as RAD21 and SMC3 (Table 4). Top 30 pairs of FR and RF orientations of DNA motifs

co-occupied in the same open chromatin regions were shown (Table 4). Total number of

pairs of DNA motifs was 601, consisting of 155 kinds of DNA motifs, when the pair of

DNA motifs were observed in more than 80% of the number of open chromatin regions

with the DNA motifs.

Comparison with chromatin interaction data

To examine whether the biased orientation of DNA motif sequences is associated

with chromatin interactions, enhancer-promoter interactions (EPI) predicted based on

enhancer-promoter associations (EPA) were compared with chromatin interaction data

(Hi-C). Due to the resolution of Hi-C experimental data used in this study (50kb), EPI

were adjusted to 50kb resolution. EPI were predicted based on three types of EPA: (i)

EPA without being shortened by DNA motif sequences, (ii) EPA shortened at the

genomic locations of DNA motif sequences of transcription factors such as CTCF and

cohesin (RAD21, SMC3) without considering their orientation, and (iii) EPA shortened

at the genomic locations of forward-reverse or reverse-forward orientation of DNA

motif sequences of transcription factors. EPA (iii) showed the highest ratio of EPI

overlapped with chromatin interactions (Hi-C) using DNA binding motif sequences of

CTCF and cohesin (RAD21 and SMC3), compared with the other two types of EPA

(Figure 3). Total 62 biased orientation (41 FR and 24 RF) of DNA motif sequences

including CTCF and cohesin showed a higher ratio of EPI overlapped with Hi-C than

the other types of EPA in monocytes (Table 5). Chromatin interaction data were

obtained using different samples from DNase-seq, open chromatin regions, so

individual differences seemed to be large from the results of this analysis. Since, for

some DNA motif sequences of transcription factors, the number of EPI overlapped with

chromatin interactions was small, if higher resolution of chromatin interaction data

(such as Hi-ChIP, in situ DNase Hi-C, and in situ Hi-C data, or a tool to improve the

resolution such as HiCPlus) is available, the number of EPI overlapped with chromatin

interactions would be increased and the difference of the numbers among three types of

EPA might be larger and more significant than the current results (Rao et al. 2014;

Ramani et al. 2016; Mumbach et al. 2017; Zhang et al. 2018).

Discussion

To find DNA motif sequences of transcription factors and DNA repeat sequences

affecting human putative transcriptional target genes, DNA motif sequences were

searched from open chromatin regions of monocytes of four people. Total 351 biased

(192 forward-reverse and 179 reverse-forward) orientation of DNA motif sequences

were found in monocytes of four people in common, whereas only one any orientation

(i.e. without considering orientation) of DNA motif sequence was found to affect

putative transcriptional target genes, suggesting that EPA shortened at the genomic

locations of forward-reverse or reverse-forward orientation of DNA motif sequences of

transcription factors is an important character for EPI and transcription of genes.

When DNA motif sequences were searched from monocytes of one person, a

larger number of biased orientation of DNA motif sequences affecting human putative

transcriptional target genes has been found. When the number of people, from which

experimental data were obtained, was increased, the number of DNA motif sequences

found in all people in common decreased and in some cases, known transcription

factors involved in chromatin interactions such as CTCF and cohesin (RAD21 and

SMC3) were not identified by statistical tests. This would be caused by individual

difference of the same cell type, low quality of experimental data, and experimental

errors. Moreover, though forward-reverse orientation of DNA binding motif sequences

of CTCF and cohesin is frequently observed at chromatin interaction anchors, the

percentage of forward-reverse orientation is not 100, and other orientations of the DNA

binding motif sequences are also observed. Though DNA binding motif sequences of

CTCF and cohesins are found in various open chromatin regions, DNA binding motif

sequences of some transcription factors would be observed less frequently in open

chromatin regions. The analyses of cells of a number of people would avoid missing

relatively weak statistical significance of DNA motif sequences in cells of each person

by multiple testing correction of thousands of statistical tests. A DNA motif sequence

was found with p-value < 0.05 in cells of one person and the DNA motif sequence

found in the same cell type of four people in common would have p-value < $(0.05)^4$ =

0.00000625. Actually, DNA motif sequences with p-value slightly less than 0.05 in

monocytes of one person were observed in monocytes of four people in common.

EPI were compared with chromatin interactions (Hi-C) in monocytes. EPA

shortened at the genomic locations of DNA binding motif sequences of CTCF and

cohesin (RAD21 and SMC3) showed a significant difference of the ratios of EPI

overlapped with chromatin interactions, according to three types of EPA (FR, any, and

others). Using open chromatin regions overlapped with ChIP-seq experimental data of

histone modification of an enhancer mark (H3K27ac), the ratio of EPI not overlapped

with Hi-C was reduced and the difference of the ratio of EPI overlapped with chromatin

interactions among the three types of EPA became larger (data not shown). (Phanstiel et

al. 2017) also reported that there was an especially strong enrichment for loops with

H3K27 acetylation peaks at both ends (Fisher's Exact Test, p = 1.4 $\times$ 10^{-27}). However,

the total number of EPI overlapped with chromatin interactions was also reduced using

H3K27ac peaks, so more chromatin interaction data would be needed to obtain reliable

results in this study. As an issue of experimental data, chromatin interaction data were

obtained using different samples from DNase-seq, open chromatin regions, so

individual differences were not reflected to chromatin interaction data. Moreover, the

resolution of chromatin interaction data used in this study is about 50kb, thus the

number of chromatin interactions was relatively small (72,284 at 50kb resolution with

cutoff score of CHiCAGO tool > 1). EPA shortened at the genomic locations of DNA

binding motif sequences of transcription factors found in various open chromatin

regions such as CTCF and cohesin (RAD21 and SMC3) tend to be overlapped with a

larger number of chromatin interactions than DNA binding motif sequences of

transcription factors less frequently observed in open chromatin regions. Therefore, to

examine the difference of the numbers of EPI overlapped with chromatin interactions,

according to three types of EPA (FR, any, and others), the number of chromatin

interactions should be large enough. If the resolution of chromatin interaction data

would be increased by using other experimental data such as Hi-ChIP, in situ DNase

Hi-C, and in situ Hi-C data, or a tool to improve the resolution of chromatin interaction

data such as HiCPlus, more DNA binding motif sequences of transcription factors less

frequently observed in open chromatin regions would be examined (Rao et al. 2014;

Ramani et al. 2016; Mumbach et al. 2017; Zhang et al. 2018).

Methods

Search for biased orientation of DNA motif sequences

To examine transcriptional regulatory target genes, bed files of hg38 of Blueprint

DNase-seq data for Monocytes CD14$^+$ (EGAD00001002286) were obtained from

Blueprint web site (http://dcc.blueprint-epigenome.eu/#/home), and the bed files of

hg38 were converted into those of hg19 using Batch Coordinate Conversion (liftOver)

web site (https://genome.ucsc.edu/util.html). Bed files of hg19 of ENCODE

CD4$^+$_Naive_Wb11970640 (GSM1014537; UCSC Accession: wgEncodeEH003156),

H1-hESC (GSM816632; UCSC Accession: wgEncodeEH000556), iPSC (GSM816642;

UCSC Accession: wgEncodeEH001110), HUVEC (GSM1014528; UCSC Accession:

wgEncodeEH002460),   and   MCF-7   (GSM816627;   UCSC   Accession:

wgEncodeEH000579)   were   obtained   from   the   ENCODE   websites

(http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDgf/;

http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDnase/).

To identify transcription factor binding sites (TFBS) from the DNase-seq data,

TRANSFAC (2013.2), JASPAR (2010), UniPROBE, BEEML-PBM, high-throughput

SELEX, Human Protein-DNA Interactome, and transcription factor binding sequences

of ENCODE ChIP-seq data were used (Wingender et al. 1996; Newburger and Bulyk

2009; Portales-Casamar et al. 2010; Xie et al. 2010; Zhao and Stormo 2011; Jolma et al.

2013; Kheradpour and Kellis 2014). Position weight matrices of transcription factor

binding sequences were transformed into TRANSFAC matrices and then into MEME

matrices using in-house scripts and transfac2meme in MEME suite (Bailey et al. 2009).

Transcription factor binding sequences of transcription factors derived from vertebrates

were used for further analyses. Searches were conducted for transcription factor binding

sequences from each narrow peak using FIMO with p-value threshold of 10^{-5} (Grant et

al. 2011). Transcription factors corresponding to transcription factor binding sequences

were searched computationally by comparing their names and gene symbols of HGNC

(HUGO Gene Nomenclature Committee) -approved gene nomenclature and 31,848

UCSC   known   canonical   transcripts

(http://hgdownload.cse.ucsc.edu/goldenpath/hg19/database/knownCanonical.txt.gz),   as

transcription factor binding sequences were not linked to transcript IDs such as UCSC,

RefSeq, and Ensembl transcripts.

Target genes of a transcription factor were assigned when its TFBS was found in

DNase-seq narrow peaks in promoter or extended regions for enhancer-promoter

association of genes (EPA). Promoter and extended regions were defined as follows:

promoter regions were those that were within distances of ±5 kb from transcriptional

start sites (TSS). Promoter and extended regions were defined as per the following

association rule, which is the same as that defined in Figure 3A of a previous study

(McLean et al. 2010): the single nearest gene association rule, which extends the

regulatory domain to the midpoint between the TSS of the gene and that of the nearest

gene upstream and downstream without the limitation of extension length. Extended

regions for EPA were shortened at the genomic locations of DNA binding sites of

transcription factors that were the closest to a transcriptional start site, and

transcriptional target genes were predicted from the shortened enhancer regions using

TFBS. Furthermore, promoter and extended regions for EPA were shortened at the

genomic locations of forward–reverse orientation of DNA binding sites of transcription

factors. When forward or reverse orientation of DNA binding sites were continuously

located in genome sequences several times, the most external forward–reverse

orientation of DNA binding sites were selected. The genomic positions of genes were

identified using 'knownGene.txt.gz' file in UCSC bioinformatics sites (Karolchik et al.

2014). The file 'knownCanonical.txt.gz' was also utilized for choosing representative

transcripts among various alternate forms for assigning promoter and extended regions

for EPA. From the list of transcription factor binding sequences and transcriptional

target genes, redundant transcription factor binding sequences were removed by

comparing the target genes of a transcription factor binding sequence and its

corresponding transcription factor; if identical, one of the transcription factor binding

sequences was used. When the number of transcriptional target genes predicted from a

transcription factor binding sequence was less than five, the transcription factor binding

sequence was omitted.

For gene expression data, RNA-seq reads mapped onto human hg19 genome

sequences were obtained, including ENCODE long RNA-seq reads with poly-A of

H1-hESC, iPSC, HUVEC, IMR90, and MCF-7 (GSE26284, GSM958733,

GSM2344099, GSM2344100, GSM958734, and GSM765388), and UCSF-UBC human

reference epigenome mapping project RNA-seq reads with poly-A of naive CD4$^+$ T

cells (GSM669617). Two replicates were present for H1-hESC, iPSC, HUVEC, and

MCF-7, and a single one for CD4$^+$ T cells. RPKMs of the RNA-seq data were

calculated using RSeQC (Wang et al. 2012). For monocytes, Blueprint RNA-seq RPKM

data (GSE58310, GSE58310_GeneExpression.csv.gz, Monocytes_Day0_RPMI) was

used (Saeed et al. 2014). Based on RPKM, UCSC transcripts with expression levels

among top 30% of all the transcripts were selected in each cell type.

The expression level of transcriptional target genes predicted based on EPA

shortened at the genomic locations of DNA motif sequences of transcription factors or

DNA repeat sequences was compared with the expression level of transcriptional target

genes predicted from promoter. For each DNA motif sequence shortening EPA,

transcriptional target genes were predicted using about 3,000 − 5,000 DNA binding

motif sequences of transcription factors, and the expression level of putative

transcriptional target genes of each transcription factor was compared between EPA and

only promoter using Mann-Whitney test (p-value < 0.05). The number of transcription

factors showing a significant difference of expression level of putative transcriptional

target genes between EPA and promoter was compared among forward-reverse,

reverse-forward, and any orientation of DNA motif sequences shortening EPA using

chi-square test (p-value < 0.05). When DNA motif sequences of transcription factors or

DNA repeat sequences shortening EPA showed a significant difference of expression

level of putative transcriptional target genes in forward-reverse, reverse-forward, or any

orientation in monocytes of four people in common, the DNA motif sequences were

listed.

Though forward-reverse orientation of DNA binding motif sequences of CTCF

and cohesin is frequently observed at chromatin interaction anchors, the percentage of

forward-reverse orientation is not 100, and other orientations of the DNA binding motif

sequences are also observed. Though DNA binding motif sequences of CTCF and

cohesin are found in various open chromatin regions, DNA binding motif sequences of

some transcription factors would be observed less frequently in open chromatin regions.

The analyses of cells of a number of people would avoid missing relatively weak

statistical significance of DNA motif sequences in cells of each person by multiple

testing correction of thousands of statistical tests. A DNA motif sequence was found

with p-value < 0.05 in cells of one person and the DNA motif sequence found in the

same cell type of four people in common would have p-value < 0.05^4 = 0.00000625.

Co-location of biased orientation of DNA motif sequences

Co-location of biased orientation of DNA binding motif sequences of

transcription factors was examined. The number of open chromatin regions with the

same pairs of DNA binding motif sequences was counted, and when the pairs of DNA

binding motif sequences were enriched with statistical significance (chi-square test,

p-value < 1.0 × 10^{-10}), they were listed.

Comparison with chromatin interaction data

For chromatin interactions, 'PCHiC_peak_matrix_cutoff0.txt.gz' file was

downloaded from 'Promoter Capture Hi-C in 17 human primary blood cell types' web

site (https://osf.io/u8tzp/files/), and chromatin interactions for Monocytes with scores of

CHiCAGO tool > 1 were extracted from the file (Javierre et al. 2016).

EPI were predicted using three types of EPA in monocytes: (i) EPA shortened at

the genomic locations of forward-reverse or reverse-forward orientation of DNA motif

sequences of transcription factors, (ii) EPA shortened at the genomic locations of any

orientation (i.e. without considering orientation) of DNA motif sequences of

transcription factors, and (iii) EPA without being shortened by DNA motif sequences.

EPI predicted using the three types of EPA in common were removed. First, EPI

predicted based on EPA (i) were compared with chromatin interactions (Hi-C). The

resolution of chromatin interaction data used in this study was 50kb, so EPI were

adjusted to 50kb, before their comparison. The number and ratio of EPI overlapped with

chromatin interactions were counted. Second, EPI were predicted based on EPA (ii), and

EPI predicted based on EPA (i) were removed from the EPI. The number and ratio of

EPI overlapped with chromatin interactions were counted. Third, EPI were predicted

based on EPA (iii), and EPI predicted based on EPA (i) and (ii) were removed from the

EPI. The number and ratio of EPI overlapped with chromatin interactions were counted.

The number and ratio of the EPI compared two times between EPA (i) and (iii), and

EPA (i) and (ii) (binomial distribution, p-value < 0.025 for each test).

Acknowledgements

The supercomputing resource was provided by Human Genome Center of the Institute

of Medical Science at the University of Tokyo. Computations were partially performed

on the NIG supercomputer at ROIS National Institute of Genetics. Publication charges

for this article were funded by JSPS KAKENHI Grant Number 16K00387. This

research was partially supported by the Platform Project for Supporting in Drug

Discovery and Life Science Research（Platform for Dynamic Approaches to Living

System）from Japan Agency for Medical Research and Development (AMED). This

research was partially supported by Development of Fundamental Technologies for

Diagnosis and Therapy Based upon Epigenome Analysis from Japan Agency for

Medical Research and Development (AMED). This work was partially supported by

JST CREST Grant Number JPMJCR15G1, Japan.

423
424

425　**Figures**

Figure 1. Chromatin interactions and enhancer-promoter association. (A) Forward–

reverse orientation of CTCF-binding sites is frequently found in chromatin interaction

anchors. CTCF can block the interaction between enhancers and promoters limiting the

activity of enhancers to certain functional domains (de Wit et al. 2015; Guo et al. 2015).

(B) Computationally-defined regulatory domains for enhancer-promoter association

(McLean et al. 2010). The *single nearest gene* association rule extends the regulatory

domain to the midpoint between this gene's TSS and the nearest gene's TSS both

upstream and downstream. Enhancer-promoter association was shortened at the

genomic locations of forward-reverse orientation of DNA binding motif sequences of

transcription factors (e.g. CTCF in the figure).

Figure 2. Biased-orientation of DNA motif sequences of transcription factors. Total 351

of biased orientation of DNA binding motif sequences of transcription factors have been

found to affect the expression level of putative transcriptional target genes in monocytes

of four people in common, whereas only one any orientation (i.e. without considering

orientation) of DNA binding motif sequences have been found to affect the expression

level.

Figure 3. Comparison of DNA binding motif sequences of transcription factors with

chromatin interactions. Total 62 biased orientation (41 FR and 24 RF) of DNA motif

sequences including CTCF and cohesin showed a higher ratio of EPI overlapped with

Hi-C than the other types of EPA in monocytes. FR: forward-reverse orientation of

DNA motifs. any: any orientation of DNA motifs. others: enhancer-promoter

association not shortened at the genomic locations of DNA motifs. * p-value < 10^{-5}.

Tables

Table 1. Top 90 biased orientation of DNA binding motif sequences of transcription

factors in monocytes. TF: DNA binding motif sequences of transcription factors. Score:

$-\log_{10}$ (p-value).

Forward-reverse orientation

TF	Score	TF	Score	TF	Score
SRF	75.75	YLL054C	34.01	NR2F2	25.16
IRF7	71.35	ZIC1	33.85	PLAGL1	24.95
MINI20	61.98	SMC3	33.67	REST	24.83
CTCF	61.90	ESR2	32.37	GAL4P	24.48
E2F	59.63	TP63	32.28	ZNF143	23.97
STAT1	59.24	ZNF836	31.96	HLH2_HLH3	23.85
ZNF317	56.27	STA5A	31.54	BM9	23.63
CEBPG	55.76	MYB	31.23	YB1	23.29
ZNF623	55.01	AT3G16280	31.22	PRDM15	22.93
ZNF93	54.53	E2F8	30.92	ZNF92	22.52
HMBOX1	53.29	P53	30.86	CPRF1	22.42
SMAD2_SMAD3_SMAD4	50.96	STA5B	30.24	ZNF214	22.41
FOXN2	50.18	STAT3	30.15	PTF1A	22.39
CLAMP	49.82	AP1	30.05	SMRC2	22.16
RDR1	46.44	AT3G04100	29.77	ZNF695	22.08
ZNF646	44.25	CEBPZ	29.73	ZNF28	22.03
TFE3	42.76	RAD21	28.98	ZNF316	21.96
CMYB	42.28	GATA2	28.82	TATA	21.90
KR	41.45	SMAD2	28.81	NF1B	21.75
STAT4	40.84	KLF6	28.08	PHX2B	21.12
ZNF716	40.69	HXB1	27.93	ATHB1	20.96
KLF14	39.94	GLI1	27.39	DEC	20.92
DREB2E	39.93	GCMA	27.03	ZNF709	20.88
FOXA1	38.43	ZNF676	26.96	YY1	20.87
USF	38.04	E47	26.62	EIF5A2	20.85
HME1	37.31	EHF	26.62	ZN148	20.75
ZN219	35.67	TCF12	26.59	PAZ5	20.35
D	35.16	RSRFC4	26.53	PO6F1	20.24
IKZF1	34.09	HIC1	25.64	FOXO4	19.97
YBR033W	34.01	EGR2	25.23	EBF	19.56

463 Reverse-forward orientation

TF	Score	TF	Score	TF	Score
RFX5	102.60	GABP1_GABP2	36.27	WT1	29.37
TCF1	75.65	RDS2	35.16	E74A	29.16
SRF	65.83	TCFAP2B	34.85	CREM	28.95
ZNF28	64.19	TBP	34.77	ZNF316	28.64
TCF2	62.76	ZNF670	34.63	HAND1_E47	28.37
ZNF225	62.48	ZBRK1	34.60	ZNF449	28.26
CEBPB	59.99	ZNF687	34.12	ZNF233	27.97
NFY	59.26	NFKB2	33.35	SOX2	26.77
ZNF286B	55.03	RFX3	33.04	RXRA	26.37
CTCF	53.46	NRF3	32.99	MYF6	26.15
PAX5	52.87	VND	32.75	MAFK	26.12
STAT1	52.41	RNF96	32.57	CEBPZ	26.03
ISL1	50.03	SP3	32.52	PO2F2	25.89
ETV7	47.38	FLI1	32.45	MADS10	25.62
KLF16	46.75	RBAK	32.22	CETS2	25.35
SP1	46.33	NR2F1	32.10	HSF1	25.06
SMAD3	45.57	NKX3	31.85	GCN4P	24.53
STAT5B	45.02	ZIC2	31.68	CRZ1	24.46
TEAD1	43.30	PARP	31.49	AHR	23.86
ZFP579	43.25	ZNF195	31.36	CRZ1P	23.80
ZBTB2	42.58	BDP1	31.14	PAX6	23.78
ZNF343	41.31	SOX9	31.03	TOE3	23.67
DOBOX5	39.35	SP8	30.79	IRF4	22.77
NANOG	39.14	EMBP1B	30.62	RAP1	22.54
LEU3	38.92	ZNF121	30.34	ETV3	22.20
NRF2	38.83	ZFP202	30.31	FOXP2	21.97
OBOX2	38.30	CD59	30.15	TATA	21.93
ZNF652	38.14	RXRB	29.87	CDX2	21.75
ZNF30	37.89	STAT5A	29.61	NFAC1	21.54
MYOG	36.64	SOX11	29.39	SP4	21.44

467 **Table 2.** Top 30 forward-reverse orientation of DNA binding motif sequences of

468 transcription factors in MCF-7 and iPS. TF: DNA binding motif sequences of

469 transcription factors. Score: $-\log_{10}$ (p-value).

470 MCF-7   iPS

TF	Score	TF	Score
CTCF	23.81	ZIC3	19.62
RAD21	16.51	ZNF521	19.24
SMC3	15.93	TFE3	16.93
E2F	10.41	ZNF317	16.74
CAC	9.82	NKX25	16.54
HOXA7	8.59	HES1	15.64
STA5B	7.60	ZBTB24	14.51
REST	7.55	AP1	13.99
SIRT6	7.35	SMC3	13.83
KLF3	6.82	RAD21	13.24
YY1	5.59	YLL054C	12.80
BM9	5.44	ZNF836	11.89
DREB2E	5.29	TBX5	11.72
PLAL1	5.27	CTCF	11.54
ZN219	5.19	STAT1	11.49
STAT1	4.73	RUNX2	10.58
SNT2	4.71	WT1	10.57
ETS	3.46	ZIC1	9.91
ZBTB6	3.42	MAX	8.85
PACC	3.39	KLF3	8.60
XRCC4	3.23	KLF14	8.29
DBP	2.83	ZNF143	8.11
SETB1	2.82	E2F5	7.96
RREB1	2.80	SIRT6	7.86
E2F5	2.68	HME1	7.79
CLAMP	2.52	YY1	7.31
ZNF143	2.50	STA5B	7.03
BTEB2	2.47	MAZ	6.69
YKL222C	2.40	ZN148	6.50
AP1	2.28	E47	6.03

474 **Table 3.** Biased orientation of DNA repeat sequences in monocytes. Score: $-\log_{10}$

475 (*p*-value).

DNA repeat seq.	Score
AluSg	7.52
L1ME4b	8.83
LTR16C	2.94

Table 4. Top 30 of co-locations of biased orientation of DNA binding motif sequences

of transcription factors in monocytes. Co-locations of DNA motif sequences of CTCF

with other biased orientation of DNA motif sequences were shown in the separate table.

Motif 1,2: DNA binding motif sequences of transcription factors. # overlap: the number

of open chromatin regions including both Motif 1 and Motif 2. # motif 1: the number of

open chromatin regions including Motif 1. # motif 2: the number of open chromatin

regions including Motif 2. # others: the number of open chromatin regions not including

Motif 1 or Motif 2. Open chromatin regions overlapped with histone modification

(H3K27ac) were used (Total 26,095 regions).

Motif 1	Motif 2	# both	# motif 1	# motif 2	# others	*p*-value
CTCF	SMAD2	4527	781	4478	16309	0
CTCF	RREB1	4379	929	4757	16030	0
CTCF	ZN263	4357	951	4875	15912	0
CTCF	SP1	4321	987	4052	16735	0
CTCF	RAD21	3804	1504	225	20562	0
CTCF	KLF6	3695	852	3800	17748	0
CTCF	SMC3	1259	178	1977	22681	0
CTCF	RXRA	1220	259	2559	22057	0

Motif 1	Motif 2	# both	# motif 1	# motif 2	# others	*p*-value
SMAD2	ZN263	7427	1578	1805	15285	0
SMRC2	ZN263	7354	1192	1893	15656	0
RREB1	SP1	7208	1957	1165	15765	0
SIRT6	ZN263	7126	1154	2121	15694	0
MAZ	SP1	7052	909	1321	16813	0
MAZ	ZN263	7022	999	2225	15849	0
SP1	ZN263	7009	1364	2223	15499	0
SMAD2	SP1	6925	2080	1448	15642	0
MAZ	RREB1	6920	967	2245	15963	0
SIRT6	SMAD2	6766	1514	2239	15576	0
SIRT6	SMRC2	6668	1612	1878	15937	0
PLAL1	SMAD2	6637	1074	2368	16016	0
MAZ	SMAD2	6627	1334	2378	15756	0
RARG	SMAD2	6602	1012	2403	16078	0
PLAL1	ZN263	6571	1140	2661	15723	0
EGR2	RREB1	6565	787	2600	16143	0
KLF6	RREB1	6563	932	2573	16027	0
MAZ	SMRC2	6562	1459	1984	16090	0
RREB1	ZN263	6551	1607	2681	15256	0
KLF6	SP1	6541	954	1832	16768	0
PLAL1	RREB1	6496	1215	2640	15744	0
EGR2	SP1	6492	860	1881	16862	0
KLF6	SMAD2	6491	1004	2514	16086	0
RARG	ZN263	6390	1224	2842	15639	0
PLAL1	SP1	6318	1393	2055	16329	0
KLF6	ZN263	6304	1191	2928	15672	0
EGR2	SMAD2	6270	1082	2735	16008	0
EGR2	ZN263	6253	1099	2979	15764	0
EGR2	MAZ	6219	1133	1742	17001	0
RARG	RREB1	6219	1395	2917	15564	0

492   **Table 5.** Comparison of DNA binding motif sequences of transcription factors with

493   chromatin interaction data. TF: DNA binding motif sequences of transcription factors.

494   Score: $-\log_{10}$ (*p*-value).

495   Forward-reverse   Reverse-forward orientation

TF	Score 1	Score 2
AP1	28.64	3.75
AT3G04100	24.51	7.07
BTEB2	8.64	2.69
CTCF	6.09	6.80
E2F	19.43	3.24
E2F5	10.58	2.02
EGR2	4.28	10.11
ETS	7.66	5.74
FOSB	7.98	1.99
FOXO4	7.98	5.84
GCM1	2.51	3.47
GCR	6.39	4.50
IRF7	16.36	2.03
KLF14	18.52	8.08
KLF3	6.42	6.43
MAZ	1.82	4.55
MIG2	42.64	6.22
PARG	5.38	4.21
PITX3	10.31	5.62
PLAL1	11.50	5.42
PO6F1	31.28	7.60
PTF1A	8.91	4.50
PU1	3.86	1.71
RAD21	9.28	5.16
RREB1	7.47	6.40
RUNX2	2.97	1.68
SETB1	1.63	2.08
SMAD2	4.10	3.48
SMC3	10.24	6.21
SP1	20.98	4.64
TCFAP2A	5.10	1.63
TF3B	3.21	5.75
TF3CB	20.26	4.66
TFE3	94.90	3.64
YY1	13.88	7.04
ZBTB6	15.08	3.93
ZIC1	5.37	1.63
ZN148	14.51	4.47
ZN219	15.44	7.99
ZNF28	11.88	3.36
ZNF460	14.14	1.63

TF	Score 1	Score 2
ASCL2	6.23	3.80
ATF3	17.21	2.23
BC11A	1.65	4.07
BDP1	23.49	6.85
CTCF	12.32	11.74
E2F2	23.33	1.95
EBF1	14.83	7.11
GABP1_GABP2	6.43	2.80
IRF3	2.38	1.98
IRF8	4.30	4.62
NRF3	17.95	1.78
PAX5	48.74	2.74
POLR3A	41.85	9.98
RAD21	1.79	2.26
RNF96	7.69	1.93
RPN4	21.32	4.48
RXRA	29.89	11.87
SMAD4	41.27	2.44
SP2	2.09	11.15
SP4	12.74	1.84
SP8	20.03	2.17
TCFAP2A	8.90	3.53
TCFAP2E	5.72	1.84
ZFP202	23.15	4.85

References

Bailey SD, Zhang X, Desai K, Aid M, Corradin O, Cowper-Sal Lari R, Akhtar-Zaidi B,
Scacheri PC, Haibe-Kains B, Lupien M. 2015. ZNF143 provides sequence
specificity to secure chromatin interactions at gene promoters. *Nat Commun* **2**:
6186.

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble
WS. 2009. MEME SUITE: tools for motif discovery and searching. *Nucleic*
acids research **37**: W202-208.

Barutcu AR, Lajoie BR, Fritz AJ, McCord RP, Nickerson JA, van Wijnen AJ, Lian JB,
Stein JL, Dekker J, Stein GS et al. 2016. SMARCA4 regulates gene expression
and higher-order chromatin structure in proliferating mammary epithelial cells.
Genome research **26**: 1188-1201.

de Wit E, Vos ES, Holwerda SJ, Valdes-Quezada C, Verstegen MJ, Teunissen H,
Splinter E, Wijchers PJ, Krijger PH, de Laat W. 2015. CTCF Binding Polarity
Determines Chromatin Looping. *Molecular cell* **60**: 676-684.

Grant CE, Bailey TL, Noble WS. 2011. FIMO: scanning for occurrences of a given
motif. *Bioinformatics (Oxford, England)* **27**: 1017-1018.

Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y et al.
2015. CRISPR Inversion of CTCF Sites Alters Genome Topology and
Enhancer/Promoter Function. *Cell* **162**: 900-910.

Javierre BM, Burren OS, Wilder SP, Kreuzhuber R, Hill SM, Sewitz S, Cairns J,
Wingett SW, Varnai C, Thiecke MJ et al. 2016. Lineage-Specific Genome
Architecture Links Enhancers and Non-coding Disease Variants to Target Gene
Promoters. *Cell* **167**: 1369-1384 e1319.

Jeong M, Huang X, Zhang X, Su J, Shamim M, Bochkov I, Reyes J, Jung H, Heikamp
E, Presser Aiden A et al. 2017. A Cell Type-Specific Class of Chromatin Loops
Anchored at Large DNA Methylation Nadirs. *bioRxiv*.

Ji X, Dadon DB, Abraham BJ, Lee TI, Jaenisch R, Bradner JE, Young RA. 2015.
Chromatin proteomic profiling reveals novel proteins associated with
histone-marked genomic regions. *Proc Natl Acad Sci U S A* **112**: 3841-3846.

Jolma A, Yan J, Whitington T, Toivonen J, Nitta KR, Rastas P, Morgunova E, Enge M,
Taipale M, Wei G et al. 2013. DNA-binding specificities of human transcription
factors. *Cell* **152**: 327-339.

Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, Dreszer TR,
Fujita PA, Guruvadoo L, Haeussler M et al. 2014. The UCSC Genome Browser
database: 2014 update. *Nucleic acids research* **42**: D764-770.

Kheradpour P, Kellis M. 2014. Systematic discovery and characterization of regulatory
motifs in ENCODE TF binding experiments. *Nucleic acids research* **42**:

2976-2987.

McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM,
Bejerano G. 2010. GREAT improves functional interpretation of cis-regulatory
regions. *Nature biotechnology* **28**: 495-501.

Mumbach MR, Satpathy AT, Boyle EA, Dai C, Gowen BG, Cho SW, Nguyen ML,
Rubin AJ, Granja JM, Kazane KR et al. 2017. Enhancer connectome in primary
human cells identifies target genes of disease-associated DNA elements. *Nature*
genetics **49**: 1602-1612.

Newburger DE, Bulyk ML. 2009. UniPROBE: an online database of protein binding
microarray data on protein-DNA interactions. *Nucleic acids research* **37**:
D77-82.

Osato N. 2018. Characteristics of functional enrichment and gene expression level of
human putative transcriptional target genes. *BMC Genomics* **19**: 957.

Phanstiel DH, Van Bortle K, Spacek D, Hess GT, Shamim MS, Machol I, Love MI,
Aiden EL, Bassik MC, Snyder MP. 2017. Static and Dynamic DNA Loops form
AP-1-Bound Activation Hubs during Macrophage Development. *Molecular cell*
67: 1037-1048.e1036.

Portales-Casamar E, Thongjuea S, Kwon AT, Arenillas D, Zhao X, Valen E, Yusuf D,
Lenhard B, Wasserman WW, Sandelin A. 2010. JASPAR 2010: the greatly
expanded open-access database of transcription factor binding profiles. *Nucleic*
acids research **38**: D105-110.

Ramani V, Cusanovich DA, Hause RJ, Ma W, Qiu R, Deng X, Blau CA, Disteche CM,
Noble WS, Shendure J et al. 2016. Mapping 3D genome architecture through in
situ DNase Hi-C. *Nat Protoc* **11**: 2104-2121.

Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn
AL, Machol I, Omer AD, Lander ES et al. 2014. A 3D map of the human
genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**:
1665-1680.

Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajanirefah A, Matarese F, Cheng SC,
Ratter J, Berentsen K, van der Ent MA et al. 2014. Epigenetic programming of
monocyte-to-macrophage differentiation and trained innate immunity. *Science*
(New York, NY) **345**: 1251086.

Schreiber J, Libbrecht M, Bilmes J, Noble W. 2017. Nucleotide sequence and DNaseI

sensitivity are predictive of 3D chromatin architecture. *bioRxiv*.

Wang L, Wang S, Li W. 2012. RSeQC: quality control of RNA-seq experiments.
Bioinformatics (Oxford, England) **28**: 2184-2185.

Weintraub AS, Li CH, Zamudio AV, Sigova AA, Hannett NM, Day DS, Abraham BJ,
Cohen MA, Nabet B, Buckley DL et al. 2017. YY1 Is a Structural Regulator of
Enhancer-Promoter Loops. *Cell* **171**: 1573-1588 e1528.

Wingender E, Dietze P, Karas H, Knuppel R. 1996. TRANSFAC: a database on
transcription factors and their DNA binding sites. *Nucleic acids research* **24**:
238-241.

Xie Z, Hu S, Blackshaw S, Zhu H, Qian J. 2010. hPDI: a database of experimental
human protein-DNA interactions. *Bioinformatics (Oxford, England)* **26**:
287-289.

Zhang H, Li F, Jia Y, Xu B, Zhang Y, Li X, Zhang Z. 2017. Characteristic arrangement
of nucleosomes is predictive of chromatin interactions at kilobase resolution.
Nucleic acids research **45**: 12739-12751.

Zhang Y, An L, Xu J, Zhang B, Zheng WJ, Hu M, Tang J, Yue F. 2018. Enhancing Hi-C
data resolution with deep convolutional neural network HiCPlus. *Nat Commun*
9: 750.

Zhao Y, Stormo GD. 2011. Quantitative analysis demonstrates most transcription factors
require only simple models of specificity. *Nature biotechnology* **29**: 480-483.