Discovery of directional chromatin-associated regulatory motifs

affecting human gene transcription

Naoki Osato[1*]

[1]Department of Bioinformatic Engineering, Graduate School of Information Science

and Technology, Osaka University, Osaka 565-0871, Japan

[*]Corresponding author

E-mail address: naokiosato11@gmail.com

Abstract

Background: Chromatin interactions are essential in enhancer-promoter interactions (EPIs) and transcriptional regulation. CTCF and cohesin proteins located at chromatin interaction anchors and other DNA-binding proteins such as YY1, ZNF143, and SMARCA4 are involved in chromatin interactions. However, there is still no good overall understanding of proteins associated with chromatin interactions and insulator functions.

Results: Here, I describe a systematic and comprehensive approach for discovering DNA-binding motifs of transcription factors (TFs) that affect EPIs and gene expression. This analysis identified 96 biased orientations [64 forward-reverse (FR) and 52 reverse-forward (RF)] of motifs that significantly affected the expression level of putative transcriptional target genes in monocytes, T cells, HMEC, and NPC and included CTCF, cohesin (RAD21 and SMC3), YY1, and ZNF143; some TFs have more than one motif in databases; thus, the total number is smaller than the sum of FRs and RFs. KLF4, ERG, RFX, RFX2, HIF1, SP1, STAT3, and AP1 were associated with chromatin interactions. Many other TFs were also known to have chromatin-associated functions. The predicted biased orientations of motifs were compared with chromatin interaction data. Correlations in expression level of nearby genes separated by the motif sites were then examined among 53 tissues.

Conclusion: One hundred FR and RF orientations associated with chromatin

interactions and functions were discovered. Most TFs showed weak directional biases at chromatin interaction anchors and were difficult to identify using enrichment analysis of motifs. These findings contribute to the understanding of chromatin-associated motifs involved in transcriptional regulation, chromatin interactions/regulation, and histone modifications.

Keywords: transcriptional regulation, transcription factors, enhancer-promoter interactions, chromatin interactions, CTCF, open chromatin regions, histone modifications

Background

Chromatin interactions play essential roles in enhancer-promoter interactions (EPIs) and transcriptional regulation. CTCF and cohesin proteins are located at the chromatin interaction anchors and facilitate the formation of loop structures. CTCF functions as an insulator by limiting the activity of enhancers in the loops (Fig. 1a) [1, 2]. The DNA-binding sequence of CTCF reveals its orientation bias at chromatin interaction anchors in terms of the forward-reverse (FR) orientation [1, 3, 4]. Other DNA-binding proteins such as ZNF143, YY1, and SMARCA4 (BRG1) are involved in chromatin interactions and EPIs [5-7]. ZNF143 is known to bind more often to promoters than distal regions, thereby establishing looping with the distal element bound by CTCF [5]. The DNA-binding motif of ZNF143 is enriched at chromatin

interaction anchors (Z-score > 7), whereas the DNA-binding motif of YY1 is not

enriched (Z-score < 2; [5] Figure 2a). CTCF, cohesin, ZNF143, YY1, and SMARCA4

are involved in various biological functions other than chromatin interaction. These

transcription factors (TFs) bind to open chromatin regions near transcriptional start sites

(TSS) as well, which may reduce the enrichment of the motifs at chromatin interaction

anchors.

Computational methods to predict chromatin interactions have been developed [8,

9]. However, they were not intended to find DNA motifs of TFs involved in chromatin

interactions. Xie et al. [10] proposed that the next challenge will be to develop

systematic methods to discern the specific functions of these motifs in a genome-wide

fashion.

DNA-binding proteins involved in chromatin interactions are considered to affect

the transcription of genes within the loops formed by chromatin interactions. To confirm

this hypothesis, I previously found that the expression level of human putative

transcriptional target genes of a TF was significantly changed according to

enhancer-promoter association (EPA) rules [11]. A regulatory domain for EPA was

limited within the FR orientation of CTCF-binding sites, and the transcriptional target

genes of each TF bound to enhancers and promoters were predicted using the EPA

domain (Fig. 1b and c) [12]. The expression levels predicted using the EPA domain

differed significantly from the expression levels predicted using promoters alone (or

closed chromatin regions, where enhancers would be silenced, and nearby genes would

express at low level [13]). The expression levels tended to increase in monocytes and

CD4$^+$ T cells, implying that TFs serve as activators of target genes. The expression

levels tended to decrease in embryonic stem (ES) and induced pluripotent stem (iPS)

cells, implying that TFs serve as repressors of target genes. The expression level of

putative transcriptional target genes of a TF that binds enhancers is changed

significantly when EPIs are predicted properly using the insulator sites of CTCF in

different cell types. This observation suggests that the difference in the expression level

reflects the accuracy of prediction of transcriptional target genes using a given EPA

domain. Using this approach, I planned to search for other DNA-binding proteins

involved in chromatin interactions, which affect the expression level of putative

transcriptional target genes through their insulator function.

Experimental analyses of chromatin interactions in molecular biology have

technical limitations. Chromatin interaction data are altered according to experimental

techniques, the number of DNA sequencing reads, and even replication sets of the same

cell type. Additionally, chromatin interaction data may not be abundant enough to

contain all chromatin interactions and may include experimental noise such as ligation

errors. To compensate the technical limitations of experimental analyses, a

computational search was conducted for DNA motifs of TFs that affect EPIs and the

expression level of putative transcriptional target genes in monocytes, T cells, human

mammary epithelial cells (HMEC), and neural progenitor cells (NPC) without using

chromatin interaction data. Then, I compared putative EPIs and transcription factor

binding sites (TFBSs) with chromatin interaction and gene expression data.

Results

Search for biased orientations of DNA motifs

TFBSs were predicted using open chromatin regions and DNA motifs of TFs collected from various databases and journals (see Methods). Transcriptional target genes were predicted based on TFBSs in promoters and enhancers. An EPA domain was limited within the TFBSs with insulator function, such as CTCF and cohesin (RAD21 and SMC3). To find motifs of TFs with insulator function other than CTCF and cohesin, an EPA domain was limited within the TFBSs of a putative insulator TF (Fig. 1c). Transcriptional target genes of each general TF bound in (i) distal open chromatin regions, which include enhancers, and (ii) closed chromatin regions in promoters were predicted using the EPA domain in order to estimate the enhancer activity of each general TF. The ratio of median expression level of target genes of each general TF between (i) and (ii) was calculated. The distribution of the ratios of all general TFs was compared among forward-reverse (FR), reverse-forward (RF) orientations, and non-oriented (i.e., without considering orientation) groups of TFBSs using a two-sided Mann-Whitney test (p-value $< 10^{-7}$). Putative insulator TFs with significant differences in the distributions of the ratios of expression level among FR, RF, and non-oriented were enumerated (Fig. 2a). TFBSs overlapping H3K27ac ChIP-seq peaks showed a significant difference in the expression level of their putative target genes between (i) and (ii), suggesting that the TFBSs were associated with enhancer (and promoter) signals of H3K27ac (Fig. 2a).

To confirm the predictions of TFBSs, the distributions of TFBSs were examined

in the human genome. TFBSs involved in chromatin interactions were identified near

TSS (Fig. 2b). ZNF143 is known to bind to the promoter of genes, thereby establishing

looping with the distal element bound by CTCF [5]. YY1 is a multi-functional

transcriptional regulator and is involved in transcriptional activation, repression, and

initiation [14, 15]. The distributions of TFBSs seemed to be consistent with the features

of the TFs. Most TFBSs were identified in intergenic and intron regions of the human

genome (Fig. 2c).

Several hundreds of FR and RF orientations were found in monocytes, T cells,

HMEC, and NPC, which included DNA motifs with weak biased orientations (Fig. 3).

The number of TFs was likely to be changed, according to the number of DNase-seq

reads, since the number of the reads in monocytes was larger than those of the other

three cell types. When DNase-seq reads increase, more TFBSs may be predicted to

some extent. Notably, 96 biased (64 FR and 52 RF) orientations of DNA-binding motifs

of TFs were discovered and found to be common between the four cell types, whereas

non-oriented DNA-binding motifs of TFs were not detected (Fig. 3; Table 1; Additional

file 1: Table S1). The FR orientation of DNA motifs included CTCF, cohesin (RAD21

and SMC3), YY1, and ZNF143, which are involved in chromatin interactions and EPIs.

Some TFs have both FR and RF orientations of DNA motifs; thus, the total number 96

is smaller than the sum of FRs and RFs. Palindromic DNA motifs of TFs do not have

FR and RF distinction; thus, they do not show a directional bias.

Among the 96 TFs, KLF4 [16], ERG [17, 18], RFX [19], RFX2 [20], HIF1 [21],

SP1 [22], STAT3 [23], and AP1 [24] were found to be associated with chromatin

interactions. An important example is KLF4, which is associated with three-dimensional

enhancer rewiring and transcriptional changes during the reprogramming of mouse

embryonic fibroblasts to pluripotent stem cells [16]. ATF3 [25], ZBTB7A [26], NRF1 [6,

27], TFAP2 [28], Myc [29, 30], the ETS family, ELK3 [31], ELK4 [32], RUNX3 [33],

USF1 [34, 35], AHR/ARNT [36], SP2 [37], EGR2 [38-40], ZBTB33 [41], NF-Y [42,

43], MITF [44, 45], NONO [46], FLI1 [47], E2F [48-50], BCL [51], CXXC1 [52],

E2F1 [53], EGR1 [54], REST [55, 56], RFX [57], SIN3A [58-60], AP1 [61], and STAT3

and STAT5 [62] were associated with chromatin remodeling, chromatin accessibility,

pioneer factor, and/or histone modification. TFBSs of the TFs other than those involved

in chromatin interactions would affect gene expression by chromatin-associated

functions. Huang et al. [34] have mentioned that the insulator element at the 5' end of

the chicken beta-globin locus acts as a barrier, thus protecting transgenes against

silencing effects of adjacent heterochromatin. The transcription factor USF1 binds

within the insulator at a site important in adjacent nucleosomes for generating histone

modifications associated with active chromatin and, by inference, with barrier function.

The biased orientations of TFBSs might be located near those of known TFs

involved in chromatin interactions such as CTCF, RAD21, SMC3, YY1, and ZNF143;

thus, they showed significant differences in the expression level of putative

transcriptional target genes among FR, RF, and non-orientation. However, among 447

FR and 496 RF orientations of DNA motifs in monocytes, the TFBSs of only 16 (4%)

FR and 28 (6%) RF orientations were located within 1 kb of $\geq$ 20% of the TFBSs of

either of the known TFs and their associated TFs such as CTCFL and YY2 (Additional

file 1: Table S2).

To examine whether the biased orientations of TFBSs were associated with

CTCF-binding sites, an EPA domain was limited within the TFBSs of biased

orientations of a TF in the 96 TFs found to be common among the four cell types. Then,

transcriptional target genes of each general TF were predicted from (i) open chromatin

regions in the EPA domain that did not include CTCF-binding sites upstream and

downstream of a gene, and (ii) closed chromatin regions in promoters. The ratio of

expression level between (i) and (ii) was calculated. The biased orientations of CTCF

consist of 19 DNA-binding motifs in monocytes. When upstream and downstream of

genes with the TFBSs of each DNA-binding motif sequence of CTCF were eliminated,

93, 92, 93, and 84 TFs out of 95, except for CTCF, showed a significant difference in

the distribution of the ratios of expression level (p-value $< 10^{-7}$) in monocytes, CD4$^+$ T

cells, HMEC, and NPC, respectively (Additional file 1: Table S3). This suggests that the

biased orientations of TFBSs are not significantly associated with CTCF binding-sites.

Comparison with chromatin interaction data

To examine whether the biased orientations of DNA motifs were associated with

chromatin interactions, EPIs predicted using an EPA domain were compared with

chromatin interaction data, using three replications (B2T1, B2T2, and B3T1) of

H3K27ac HiChIP chromatin interaction data in CD4$^+$ T cells separately [63]. The

resolutions of HiChIP chromatin interaction data and EPIs were adjusted to 5 kilobases

(kb). The number of unique HiChIP cis (i.e., in the same chromatin) chromatin

interactions was 19,926,360 at 5-kb resolution in B2T1 replication of T cells; 666,149 at

5-kb resolution with ≥1,000 counts for each interaction; and 78,209 at 5-kb resolution

with ≥6,000 counts for each interaction (Additional file 1: Table S4). EPIs were

predicted using three types of EPA domains: (i) EPA domain limited within the FR or

RF orientation of TFBSs of a TF with putative insulator function such as CTCF, (ii)

EPA domain limited within the TFBSs of a TF without considering their orientation, and

(iii) EPA domain which was not limited within TFBSs. Despite the limited resolution of

the chromatin interaction data, a total of 201 biased orientations [133 of 273 (49%) FR

and 90 of 211 (43%) RF; some TFs have both FR and RF motifs; thus, the total number

is smaller than the sum of FRs and RFs] of DNA motifs in T cells included CTCF,

cohesin (RAD21 and SMC3), ZNF143, and YY1 in three replications (B2T1, B2T2, and

B3T1). For the DNA motifs, a significantly higher ratio of EPIs overlapped with

HiChIP chromatin interactions with ≥1,000 counts for each interaction in EPA domain

(i) than in EPA domain (ii) and (iii) in T cells (Table 2). For a total of 390 biased

orientations [261 of 273 (96%) FR and 199 of 211 (94%) RF] of DNA motifs in T cells,

a significantly higher ratio of EPIs overlapped with the chromatin interactions in EPA

domain (i) than EPA domain (iii) (Table 2; Additional file 1: Table S4 and S5). The

difference between EPIs predicted using EPA domain (i) and (ii) seemed to be difficult

to distinguish using the chromatin interaction data and a statistical test in some cases.

However, for the difference between EPIs predicted using EPA domain (i) and (iii), a

larger number of biased orientations of DNA motifs were correlated with chromatin

interaction data. Most (95%) of the biased orientations of DNA motifs were correlated

with chromatin interactions when comparing EPIs predicted using EPA domain (i) and

(iii) with HiChIP chromatin interactions. Chromatin interaction data were obtained from

different samples of DNase-seq open chromatin regions; thus, individual differences

may exist. Individual differences of chromatin interactions were larger than those of

open chromatin regions [63]. Additionally, the analyses for Th17, Treg, and GM12878

were shown in Additional file 2.

In situ Hi-C data in HMEC, which are available in a 4D nucleome data portal,

were compared with putative EPIs. The number of unique in situ Hi-C cis chromatin

interactions was 121,873,642 at 3-kb resolution. For a total of 222 biased orientations

[155 of 310 (50%) FR and 108 of 269 (40%) RF] of DNA motifs, which included CTCF,

cohesin (RAD21 and SMC3), ZNF143, and YY1, a significantly higher ratio of EPIs

overlapped with in situ Hi-C chromatin interactions (≥ 1 count for each interaction, i.e.,

all interactions) in EPA domain (i) than in EPA domain (ii) and (iii) (Additional file 1:

Table S7). When comparing EPIs predicted using only EPA domain (i) and (iii) with the

chromatin interactions, for a total of 406 biased orientations [268 of 310 (86%) FR and

230 of 269 (86%) RF] of DNA motifs, a significantly higher ratio of EPIs overlapped

with the chromatin interactions in EPA domain (i) than EPA domain (iii) (Additional file

1: Table S7). There were six replications of in situ Hi-C experimental data of HMEC,

but other replications contained much smaller numbers of chromatin interactions. The

HiChIP chromatin interaction data in T cells were expected to be enriched with EPIs,

since they were produced with H3K27ac antibody to capture chromatin interactions. In

situ Hi-C data consist of all types of chromatin interactions and thus contain a smaller

number of EPIs than HiChIP data. Chromatin interactions with a small number of

counts may be artifacts caused by ligation errors during experiments [64, 65]. Therefore,

other replications of in situ Hi-C data with a smaller number of chromatin interactions

will not be available for this analysis.

Promoter capture Hi-C data of NPC were downloaded and analyzed by the same

procedure as that for T cells and HMEC [66]. The number of unique promoter capture

Hi-C cis chromatin interactions was 9,436,689 at 5-kb resolution. For a total of 109

biased orientations [61 of 205 (28%) FR and 60 of 241 (25%) RF] of DNA motifs,

which included CTCF, cohesin (RAD21 and SMC3), ZNF143, and YY1, a significantly

higher ratio of EPIs overlapped with promoter capture Hi-C chromatin interactions (≥ 1

count for each interaction, i.e. all interactions) in EPA domain (i) than in EPA domain

(ii) and (iii) (Additional file 1: Table S8). When comparing EPIs predicted using only

EPA domain (i) and (iii) with the chromatin interactions, for a total of 186 biased

orientations [99 of 205 (48%) FR and 103 of 241 (43%) RF] of DNA motifs, a

significantly higher ratio of EPIs overlapped with the chromatin interactions in EPA

domain (i) than EPA domain (iii) (Additional file 1: Table S8).

All three types of chromatin interaction data (HiChIP, in situ Hi-C, and promoter

capture Hi-C) indicated that EPIs predicted using EPA domain (i) that was limited

within TFBSs of known TFs involved in chromatin interactions (CTCF, RAD21, SMC3,

ZNF143 and YY1) overlapped with more chromatin interactions than those using EPA

domain (ii) and (iii) in six cell types (T cells, Th17, Treg, GM12878, HMEC, and NPC)

(the analyses for Th17, Treg, and GM12878 were shown in Additional file 2). In

monocytes, Hi-C data was available, but it was not analyzed due to the low resolution of

chromatin interaction data (50 kb).

The biased orientations of TFBSs of known TFs involved in chromatin

interactions such as CTCF, RAD21, SMC3, YY1, and ZNF143 might overlap with the

biased orientations of TFBSs of other TFs; thus, they showed a significantly higher ratio

of EPIs overlapped with chromatin interactions in EPA domain (i) than EPA domain (ii)

and (iii). However, among 273 FR and 211 RF orientations of DNA motifs of TFs not

including the known TFs in T cells, the TFBSs of only 15 (6%) FR and 9 (4%) RF

orientations of DNA motifs of TFs overlapped with $\geq$ 60% of the TFBSs of biased

orientations of DNA motifs of either of the known TFs at 5-kb resolution.

Moreover, to investigate the enhancer activity of EPIs, the distribution of

expression level of putative target genes of EPIs was compared between EPIs that

overlapped with HiChIP chromatin interactions and EPIs that did not overlap with them.

Though the target genes of EPIs were selected from top 4,000 transcripts (genes) in

terms of the expression level of all transcripts (genes) excluding transcripts not

expressed in T, Th17, Treg, and GM12878 cells separately, target genes of EPIs that

overlapped with chromatin interactions showed a significantly higher expression level

than non-overlapped EPIs, suggesting that EPIs that overlapped with chromatin

interactions activated the expression of target genes in the four cell types. For almost all

(99%-100%) FR and RF orientations of DNA motifs, a significantly higher expression

level of putative target genes was observed for EPIs that overlapped with chromatin

interactions than for EPIs that did not overlap in the four cell types (Additional file 1:

Table S5 and S6). DNA motifs showing a significantly lower expression level were not

observed in this analysis. HiChIP data were produced using H3K27ac antibody; thus,

chromatin interactions acting as repressors would not be identified. However, for in situ

Hi-C and promoter capture Hi-C data in HMEC and NPC, the significant difference in

the expression level of putative target genes of EPIs was observed in a small percentage

(less than 1%) of biased orientations of DNA motifs.

Recently, high resolution chromatin interaction data became publicly available

[67]. The biased orientations of TFBSs of TFs were examined using Micro-C chromatin

interaction data in human fibroblasts. The resolution of the data was adjusted to 200 bp.

TFBSs of CTCF, RAD21, SMC3, and SMRC2 showed a significant directional bias at

the pairs of chromatin interaction anchors (p-value $< 10^{-50}$). SMARCA4 is known to be

involved in higher-order chromatin structures such as the topologically associating

domain (TAD) [7]. DNA motifs of SMARCA4 did not exist in the databases of DNA

motifs of TFs used in this analysis. SMRC2 was not identified in this analysis because it

may be involved in higher-order chromatin structures that do not affect EPI and gene

expression directly by its insulator function. ATF3 and ZBTB7A in Table 1 were found

among the top 32 biased orientations of TFBSs of TFs at the pairs of chromatin

interaction anchors with p-value $< 10^{-50}$ (Additional file 1: Table S9). Deletion of an

ATF3-binding site affected the expression of a target gene, which was far apart from the

site [25]. ZBTB7A is involved in the changes in chromatin accessibility and the

regulation of gene expression [26]. TFBSs of YY1 and ZNF143 also indicated a weak

directional bias at the chromatin interaction anchors (Additional file 1: Table S9). In this

analysis, biased orientations of TFBSs of TFs were investigated based on the difference

of distribution of the expression level of putative transcriptional target genes. YY1 and

ZNF143 were found to have significant directional biases, thereby suggesting that the

analysis in this study could find TFs that did not have a strong directional bias based on

the number of TFBSs at chromatin interaction anchors. The other TFs, except for CTCF,

RAD21, SMC3, ATF3, and ZBTB7A in Table 1, showed weak directional biases at the

chromatin interaction anchors (Additional file 1: Table S9).

Correlation of expression level of gene pairs separated by TFBS

To investigate whether the biased orientations of TFBSs affected gene expression,

the expression level of protein-coding genes, that are most closely located upstream and

downstream of any biased orientations of TFBSs in the human genome, was compared

in monocytes. The expression levels of closely located genes including divergent gene

pairs tend to be correlated among tissues [68, 69]. When a biased orientation of TFBS

serves as an insulator (e.g. CTCF), the correlation of gene expression level would be

reduced. Among 96 biased (64 FR and 52 RF) orientations of TFBSs found in common

in monocytes, T cells, HMEC, and NPC, TFs with ≥ 50 genomic locations of the

TFBSs were selected after the elimination of ubiquitously expressed genes among 53

tissues and TFBSs near the genes with a coefficient of variance (CV) < 90. After this

filtering, 43 of 60 (72%) FR and 40 of 50 (80%) RF orientations showed significantly

lower correlations of expression levels of gene pairs separated by the TFBSs of a TF

than the correlations of all pairs of neighboring genes with intergene distance <1

megabase (Mb) (Mann-Whitney test, p-value < 0.05) (Table 3). The FR and RF

orientations of TFs included known TFs involved in chromatin interactions such as

CTCF, RAD21, SMC3, YY1, and ZNF143. The top 20 FR and RF orientations of DNA

motifs were selected in ascending order of the median correlation coefficient of

expression levels of gene pairs separated by the TFBSs of a TF. The 20 TFs included

known TFs involved in chromatin interactions such as CTCF, RAD21, and SMC3 (Fig.

4) (Mann-Whitney test, p-value < 0.01). Instead of using all TFBSs of a TF, when

TFBSs of a TF were limited to sites used in the analysis of biased orientations of DNA

motifs in an EPA domain, 33 of 55 (60%) FR and 36 of 47 (77%) RF orientations

showed significantly lower correlations of expression levels of gene pairs separated by

the TFBSs (Additional file 1: Table S10). With intergene distance < 20 kb, 21 of 62

(34%) FR and 8 of 52 (15%) RF orientations showed significantly lower correlations of

expression levels of gene pairs separated by the TFBSs of a TF than the correlations of

all pairs of neighboring genes (Additional file 1: Table S10).

A large number of TFBSs of a TF tend to show significantly lower correlations of

gene expression (median numbers of TFBSs of a TF showing significantly lower

correlations were 656 and 479 for FR and RF, respectively, and median numbers of

TFBSs of a TF not showing significantly lower correlations were 122 and 115 for FR

and RF, respectively. The Mann-Whitney test p-value = 0 and < 10^{-10} for FR and RF,

respectively. This suggests that if the criteria to examine the correlations of expression

levels of gene pairs separated by TFBSs are more stringent, and the number of TFBSs is

reduced, it would be more difficult to use a statistical test to compare the correlations of

expression levels of gene pairs separated by the reduced TFBSs, particularly for TFs

with a small number of TFBSs.

Among 91 biased orientations of DNA motifs of TFs found in common among

the four cell types, after excluding the five known TFs involved in chromatin

interactions such as CTCF, RAD21, SMC3, YY1, and ZNF143, TFBSs of 88 TFs were

located at ≥ 1 genomic regions between gene pairs not including TFBSs of the known

TFs and at ≥ 1 genomic regions between low correlation (< 0.13) of expression level of

gene pairs (Table 3). TFBSs of 52 TFs were placed at ≥ 10 genomic regions, though

some proportion of TFBSs of the known TFs might not be associated with insulator

function.

Co-location of motifs with biased orientations

To examine the association of directional DNA-binding motifs, co-location of

motifs within 200 bp (approximately the minimum length of an open chromatin region

of 146 bp) of each other was analyzed in monocytes, T cells, HMEC, and NPC. The

same pairs of motifs upstream and downstream of genes in EPA domain were

enumerated, and the pairs of motifs found in ≥ 100 genomic regions were listed (Table

4; Additional file 1: Table S11). A majority of the pairs of motifs overlapped with more

than 1 base, but the same tendency was found in the analysis of motifs of all TFs

(Additional file 1: Table S11). CTCF partially overlapped with cohesin RAD21 and

SMC3 (Table 4). The top 20 pairs of FR and RF orientations of motifs overlapped or

co-localized have been shown (Table 4). Unique pairs of overlapping or co-localized

motifs were 404 FR and 662 RF, consisting of 117 FR and 218 RF of unique DNA

motifs in monocytes. Only 10 overlapping or co-localized motifs were found to be

common among the four cell types, including pairs of CTCF, RAD21, SMC3, and YY1,

thereby implying that most overlapping and co-localized motifs serve in a cell-specific

manner (Additional file 1: Table S11). Paired DNA-binding motifs of TFs with biased

orientations tend to overlap, and a relatively small number of motifs are co-localized

upstream and downstream of genes. Overlapping motifs act as TFBSs for several TFs

either simultaneously or competitively [70-77]. CTCF and cohesin function

cooperatively with another TF and a protein such as KLF4 and BRD2 [78, 79]. KLF4

recruits cohesin to the *Oct4* enhancer region. BRD2 supports boundary (insulator)

activity of CTCF. The analysis of 457 ChIP-seq data sets from 119 human TFs showed

that secondary motifs were detected in addition to the canonical motifs of the TFs,

indicating tethered binding and co-binding between multiple TFs [80]. Wang et al. [80]

observed significant position and orientation preferences between many co-binding TFs.

Overlapping motifs found in this study may be biologically functional rather than

artifact. Though repeat sequences in the human genome were masked, more than a

hundred pairs of motifs overlapped with hundreds of upstream and downstream gene

regions.

Discussion

A systematic method to find DNA-binding motifs of TFs that affect EPIs and

gene expression without using chromatin interaction data is presented in this study.

DNA motifs of TFs such as CTCF and cohesin (RAD21 and SMC3) were searched in

the open chromatin regions of monocytes, T cells, HMEC, and NPC. A total of 96

biased (64 FR and 52 RF) orientations of DNA motifs were discovered and found to be

common among the four cell types. Remarkably, non-oriented (i.e., without considering

orientation) motifs was not found to affect the expression level of putative

transcriptional target genes, suggesting that an EPA domain limited within the FR or RF

orientation of TFBSs was a more accurate prediction of EPIs. This was supported by the

comparisons with chromatin interaction and gene expression data. FF and RR

orientations of motifs were not examined in this analysis since CTCF has FR or RF

orientation of their TFBSs [3]. Among the 96 biased orientations of DNA motifs studied,

14 were reported to be associated with chromatin interactions, and many other TFs have

chromatin-associated functions. The enrichment analysis of biased orientations of

TFBSs of the 96 TFs for pairs of chromatin interaction anchors indicated a weak

statistical significance, except for CTCF, RAD21, SMC3, ATB3, and ZBTB7A,

suggesting that the TFs were difficult to find using DNA motifs of TFs alone [5].

Without considering the difference of orientations among cell types, 175 biased

orientations were found in common among the four cell types. These numbers (96 and

175) are the numbers of unique TFs with biased orientations of DNA motifs and not the

numbers of unique DNA motifs. Possible causes of the discrepancy are as follows: (1) A

TF or an isoform of the same TF may bind to different DNA-binding motifs, according

to cell types and/or in the same cell type. About 50% or more alternative splicing

isoforms are differently expressed among tissues, thereby indicating that most

alternative splicing is subject to tissue-specific regulation [81-83]; (2) The same TF has

several DNA-binding motifs, and in some cases, one of the motifs is almost the same as

the reverse complement motif of another motif of the same TF in a database; (3) I

previously found that a complex of TFs would bind to a slightly different DNA-binding

motif by combining DNA-binding motifs from the complex in *C. elegans* [84]. In other

reports, the difference of binding specificity of TF pairs was investigated [85].

Cofactors also alter the binding specificity of TFs [70-72]. These increase the variety

and number of (biased orientations of) DNA-binding motifs, which are potentially

involved in different biological functions among cell types.

To predict TFBSs in genomic DNA, several tools and methods were explored and

then protein interaction quantitation (PIQ) was used [27]. PIQ predicts TFBSs by

reducing noise in DNase-seq data, and the comparison of the prediction with ChIP-seq

data has revealed that PIQ shows a better performance than other tools and digital

genome footprinting (DNase-DGF). The prediction of TFBSs using PIQ in this study

increased the number of biased orientations of motifs, thereby reducing the number of

non-oriented motifs. At first, the number of predicted motifs was expected to increase

relative to using consensus DNA-binding sequences of TFs and open chromatin regions

available in the ENCODE database, which seemed to be difficult to find all TFBSs in

the genome. However, the number of TFBSs was almost the same or slightly decreased

compared with the analysis of consensus sequences, suggesting that PIQ increased

true-positive predictions of TFBSs and reduced false-positive predictions.

When forming a homodimer or heterodimer with another TF, a TF may bind to

genomic DNA with a specific orientation of its TFBSs. The approach in this study

would be able to identify TFs forming a heterodimer. If the TFBS of only the mate to a

pair of TFs was found in an EPA domain, the EPA domain was limited within one side

of the TFBS, and transcriptional target genes were predicted using the EPA domain

limited within the side. In this analysis, the mate to both heterodimers and homodimers

of TFs can be tested to examine the effect on the expression level of transcriptional

target genes predicted using the EPA domain limited within one side. Biased

orientations of DNA-binding motifs may also be found in forward-forward or

reverse-reverse orientation.

TFs that are multi-functional transcriptional regulators such as YY1 may not

indicate a strong directional bias of their TFBSs. Though some proportion of TFBSs of

a TF involved in chromatin interactions and insulator functions is found in a biased

orientation in the genome, the other TFBSs of the TF may not form a dimer with

another TF and may not be present in a specific orientation in the genome. Thus, the

overall proportion of biased orientations of TFBSs of a TF may be reduced such that a

statistical test cannot detect it.

The approach presented in this study was tested to search for repeat sequences

that serve as insulator sites and affect the expression level of putative transcriptional

target genes of TFs. A small number of repeat sequences were identified and included

MIR, which act as insulators [86].

CTCF and cohesin-binding sites are frequently mutated in cancer [87]. Some

biased orientations of DNA motifs involved in chromatin interactions may be associated

with diseases including cancer.

The analyses conducted here suggest that biased orientations of DNA motifs of

discovered TFs affect EPIs and the expression level of putative transcriptional target

genes of general TFs bound in enhancers and promoters. This gives insight into the

nature of TFs and contributes to the understanding of functions and molecular

mechanisms of chromatin interactions, regulation of chromatin, histone modifications,

transcriptional regulation, and transcriptional cascades.

Methods

Search for biased orientations of DNA motifs

To identify transcription factor binding sites (TFBSs) from open and closed

chromatin regions, TRANSFAC (2019.2), JASPAR (2018), UniPROBE (2018),

high-throughput SELEX, transcription factor binding sequences of ENCODE ChIP-seq

data, and HOCOMOCO version 9 and 11 were used to predict insulator sites [88-94].

TRANSFAC (2019.2) was used to analyze enhancer-promoter interactions because

these data were sufficient to identify biased orientations of DNA motifs of insulator TFs

with less computational time, thereby reducing the number of non-oriented DNA motifs

of TFs. To reduce false-positive predictions of TFBSs in open chromatin regions, the

DNA motif search was performed by using the protein interaction quantitation (PIQ)

tool with 12,249 position frequency matrices converted from DNA motifs of vertebrate

TFs in the above-mentioned databases and DNase-seq data from GEO and ENCODE

database (GSM1024791 CD14$^+$ monocytes; GSM665812 CD4$^+$ T cell; GSM736634

Human mammary epithelial cell, HMEC; GSM878615 Neural progenitor cell, NPC;

ENCFF263GMV, ENCFF886XEV Th17; GSM1024741 Treg; GSM736620 GM12878)

[27]. To find open chromatin regions associated with enhancer and promoter activity,

narrow peaks of ChIP-seq experimental data of H3K27ac were predicted using MACS2

callpeak (GEO: GSM773004) [95]. To predict TFBSs of TFs from closed chromatin

regions, another method was employed. Position weight matrices of vertebrate

transcription factor binding sequences were converted into DNA consensus sequences,

which consist of an alphabet of 15 characters (the four bases A, C, G, T, the six two-fold

degenerate IUPAC codes R=[AG], Y=[CT], K=[GT], M=[AC], S=[GC], W=[AT], the

four three-fold degenerate IUPAC codes B=[CGT], D=[AGT], H=[ACT], V=[ACG],

and the four-fold degenerate character N=[ATGC]) using a convert matrix in the RSAT

website [96]. Transcription factor binding sequences were searched from genomic

regions other than narrow peaks of DNase-seq data in repeat-masked hg19 genome

sequences using the Match tool in TRANSFAC with a similarity score cutoff of 1 after

transforming the DNA consensus sequences of TFs into position frequency matrixes for

the Match search [97]. Repeat-masked hg19 genome sequences were downloaded from

the UCSC genome browser (http://genome.ucsc.edu/,

http://hgdownload.soe.ucsc.edu/goldenPath/hg19/bigZips/hg19.fa.masked.gz). TFs

corresponding to transcription factor binding sequences were searched computationally

and manually by comparing their names and gene symbols of HUGO Gene

Nomenclature Committee-approved gene nomenclature and 31,848 UCSC canonical

transcripts

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/database/knownCanonical.txt.gz)

because transcription factor binding sequences were not linked to transcript IDs such as

UCSC, RefSeq, and Ensembl transcripts.

Target genes of a TF were assigned when its TFBS was found in the promoter or

extended regions of a gene's EPA. Promoter and extended regions were defined as

follows: promoter regions were those that were within a distance of ±5 kb from

transcriptional start sites (TSS). Promoter and extended regions were defined as per the

following association rule, which is the same as that defined in Figure 3A of a previous

study [12]: the single nearest gene association rule, which extends the regulatory

domain to the midpoint between the TSS of the gene and that of the nearest gene

upstream and downstream without the limitation of extension length. The limitation of

extension length was ＜1 Mb in this study. Extended regions for an EPA were limited

within the TFBSs of a putative insulator TF closest to a TSS, and transcriptional target

genes were predicted using the limited EPA domain and TFBSs of a general TF in open

chromatin regions. Promoter and extended regions of an EPA were limited within the

forward–reverse (FR) orientation of TFBSs of a putative insulator TF. When the

forward or reverse orientation of TFBSs was located in the genome several times, the

most internal (i.e., closest to a TSS) forward–reverse orientation of the TFBSs was

selected. The genomic positions of genes were identified using the 'knownGene.txt.gz'

file in UCSC bioinformatics sites [98]. The file 'knownCanonical.txt.gz' was also

utilized to select representative transcripts among various alternate forms for assigning

promoter and extended regions of an EPA. From the list of TFBSs and transcriptional

target genes, redundant DNA motifs of TFs were removed by comparing the putative

target genes of a TFBS and its corresponding TF; if identical, one of the DNA motifs of

TF was chosen. When the number of transcriptional target genes predicted based on a

TFBS was <5, the DNA motif of TF was omitted.

For gene expression data, RNA-seq reads mapped onto the human hg19 genome

sequence were obtained from UCSF-UBC human reference epigenome mapping project

RNA-seq reads with poly-A of naive $CD4^+$ T cells (GEO: GSM669617). For monocytes,

Blueprint RNA-seq FPKM data

('C0010KB1.transcript_quantification.rsem_grape2_crg.GRCh38.20150622.results')

were downloaded from Blueprint DCC portal

(http://dcc.blueprint-epigenome.eu/#/files). RNA-seq reads with the poly-A of human

mammary epithelial cells (HMEC) in the Encyclopedia of DNA Elements at UCSC

(http://genome.ucsc.edu/encode/,

'wgEncodeCshlLongRnaSeqHmecCellPapAlnRep1.bam' file), H1-derived NPC (GEO:

GSM915326, ENCODE: ENCFF529SIO), Th17 (GSM2859479, NCBI SRA:

SRR6298326), Treg (GSM2859476, SRR6298323), and GM12878 (GSE78551,

ENCFF297QCE) were analyzed. RNA-seq reads were aligned in repeat-masked hg19

genome sequences using Burrows-Wheeler Aligner with default parameters [99].

FPKMs of the RNA-seq data were calculated by RSeQC [100]. Based on

log2-transformed FPKM, UCSC transcripts were arranged in a descending order of

expression level, and the top 4,000 transcripts were selected in each cell type.

The expression level of transcriptional target genes predicted using an EPA

domain limited within the TFBSs of a TF was compared with the expression level of

transcriptional target genes predicted using closed chromatin regions in promoters. For

each DNA motif limiting the EPA domain, transcriptional target genes were predicted

based on DNA-binding motifs of vertebrate TFs in the TRANSFAC database, and the

ratio of expression level of putative transcriptional target genes of each TF was

calculated between the EPA domain and closed chromatin regions in the promoters. The

distribution of the ratios of all TFs was compared among forward-reverse (FR),

reverse-forward (RF), and non-orientation (i.e., without considering orientation) of

TFBSs limiting an EPA domain using a two-sided Mann-Whitney test, (p-value $< 10^{-7}$).

Other parameters for the analysis were set as follows: the number of transcriptional

target genes of a TF was ≥ 50, the number of general TFs (not insulator TFs) to predict

transcriptional target genes was ≥ 50, and the number of TFBSs of a TF (insulator TF)

shortening an EPA was ≥ 100.

Comparison with chromatin interaction data

For comparison of EPIs with chromatin interactions (HiChIP) in CD4[+] T, Th17,

Treg, and GM12878 cells,

'GSM2705049_Naive_HiChIP_H3K27ac_B2T1_allValidPairs.txt',

'GSM2705050_Naive_HiChIP_H3K27ac_B2T2_allValidPairs.txt',

'GSM2705051_Naive_HiChIP_H3K27ac_B3T1_allValidPairs.txt',

'GSM2705054_Th17_HiChIP_H3K27ac_B1T2_allValidPairs.txt',

'GSM2705055_Th17_HiChIP_H3K27ac_B2T1_allValidPairs.txt',

'GSM2705056_Th17_HiChIP_H3K27ac_B3T1_allValidPairs.txt',

'GSM2705059_Treg_HiChIP_H3K27ac_B3T1_allValidPairs.txt',

'GSM2705041_GM_HiChIP_H3K27ac_B1_allValidPairs.txt',   and

'GSM2705042_GM_HiChIP_H3K27ac_B2_allValidPairs.txt' files were downloaded

from the Gene Expression Omnibus (GEO) database. The resolutions of chromatin

interactions and EPIs were adjusted to 5 kb before comparison. Chromatin interactions

with $\geq 6,000$ and $\geq 1,000$ counts for each interaction were analyzed in the comparison

in $CD4^+$ T cells. According to the unique numbers of chromatin interactions in

replications in Th17, Treg, and GM12878 cells, chromatin interactions with $\geq 1,000$

(B1T2 in Th17), $\geq 2,000$ (B2T1 and B3T1 in Th17), $\geq 6,000$ (B3T1 in Treg), and

$\geq 4,000$ (B1 and B2 in GM12878) counts for each interaction were analyzed in the

comparison.

EPIs were predicted on the basis of the following three types of EPA domains in

monocytes: (i) EPA domain limited within the FR or RF orientation of TFBSs of a TF,

(ii) EPA domain limited within the non-oriented (i.e., without considering orientation)

TFBSs of a TF, and (iii) EPA domain without being limited within a TFBS. EPIs

predicted using the three types of EPA domains in common were removed. EPIs

predicted using EPA domain (i) were removed from EPIs predicted using EPA domain

(ii). EPIs predicted using EPA domain (i) and (ii) were removed from EPIs predicted

using EPA domain (iii). The resolution of HiChIP chromatin interaction data was 1-5

kb; thus, EPIs were adjusted to 5 kb before comparison. The number and ratio of EPIs

overlapped with chromatin interactions were compared two times between EPI (i) and

(iii), and EPI (i) and (ii) (binomial distribution, p-value < 0.025 for each test, two-sided,

95% confidence interval).

For comparison of EPIs with chromatin interactions (in situ Hi-C) in HMEC, the

genomic positions of chromatin interactions in the 'HMEC_4DNFI97O9SAZ.pairs.gz'

file in 4D nucleome data portal (http://data.4dnucleome.org) were converted into the

hg19   version   of   human   genome   using   the   liftOver   tool

(http://hgdownload.soe.ucsc.edu/downloads.html#liftover).   The   resolutions   of

chromatin interaction data and EPIs were adjusted to 3 kb before comparison. All

chromatin interactions were used in the comparison. For comparison of EPIs with

chromatin interactions (promoter capture Hi-C) in NPC, the resolutions of chromatin

interaction data in the 'GSE86189_npc.po.all.txt.bz2' file (GEO: GSE86189) and EPIs

were adjusted to 5 kb before comparison. All chromatin interactions were used in the

comparison.

Putative target genes for the comparison of EPIs and chromatin interactions were

selected from top 4,000 transcripts in terms of the expression level. The expression level

of putative target genes of EPIs overlapped with chromatin interactions was compared

with EPIs not overlapped with them. When the same transcriptional target gene of a TF

in an enhancer was predicted using both an EPI overlapping a chromatin interaction and

a non-overlapping EPI, the target gene was eliminated. The distribution of expression

level of putative target genes was compared using a two-sided Mann-Whitney test

(p-value < 0.05).

Micro-C chromatin interaction data in human fibroblasts (4D nucleome data

portal ID: 4DNFICOEXGPJ) was converted into the hg19 version of the human genome

using the liftOver tool and was adjusted to 200-bp resolution. The same genomic

locations of paired chromatin interaction anchors were merged.

Correlation of expression level of gene pairs separated by TFBS

To examine the correlation of expression level of genes pairs separated by TFBSs

of a TF, gene expression data among 53 tissues were obtained from the

'GTEx_Analysis_2016-01-15_v7_RNASeQCv1.1.8_gene_median_tpm.gct.gz' file in

the GTEx database (https://gtexportal.org/home/). The correlation of log2-transformed

gene expression levels was analyzed based on RefSeq protein-coding transcripts

(RefSeq ID with 'NM_'). The same genomic locations of RefSeq transcripts were

removed, and TFBSs overlapped with RefSeq transcripts were also eliminated. Genes

closest to a TFBS and another gene were analyzed using BEDOPS closest-features

[101]. The Mann-Whitney test was performed using R.

Abbreviations

TF: Transcription factors; TFBS: Transcription factor binding sites; TSS:

Transcriptional start sites; EPA domain: Promoter and extended regions for

enhancer-promoter association; EPI: Enhancer-promoter interaction; FF:

Forward-forward; FR: Forward-reverse; RF: Reverse-forward; RR: Reverse-reverse;

ChIP-seq: ChIP-sequencing, chromatin immunoprecipitation followed by massively

parallel DNA sequencing; DNase-seq: DNase I hypersensitive sites sequencing;

RNA-seq: RNA sequencing; ENCODE: Encyclopedia of DNA elements;

Acknowledgements

Supercomputing resources were provided by the Human Genome Center of the Institute

of Medical Science at the University of Tokyo. This work was partially supported by

JSPS KAKENHI (Grant Number 16K00387) and JST CREST (Grant Number

JPMJCR15G1), Japan.

Figures

Fig. 1 Biased orientations of DNA motifs of TFs affecting enhancer-promoter interactions. **a** Forward–reverse orientation of CTCF-binding sites is frequently found in chromatin interaction anchors. CTCF can block the interaction between enhancers and promoters limiting the activity of enhancers to certain functional domains [1-4]. **b** Computationally-defined regulatory domains for enhancer-promoter association (EPA)

[12]. The *single nearest gene* association rule extends the regulatory domain to the

midpoint between this gene's TSS and the nearest gene's TSS both upstream and

downstream. The EPA domain was limited within the forward-reverse orientation of

TFBSs of a TF in this study (e.g. CTCF in the figure). **c** Transcriptional target genes of

each general TF bound in (i) distal open chromatin regions, which include enhancers,

and (ii) closed chromatin regions in promoters were predicted using an EPA domain. An

EPA domain was limited according to forward-reverse (FR), reverse-forward (RF)

orientations, and non-oriented TFBSs of a putative insulator TF. Each dot in scatter

plots represents the median expression level (FPKM) of target genes of a TF. The ratio

of median expression level between (i) (y-axis) and (ii) (x-axis) was calculated to

compare the distribution of the ratios of all general TFs among the three types of EPA

domains.

Fig. 2 Identification of TFBSs in open chromatin regions and properties of TFBSs. **a**
Comparison of expression level of putative transcriptional target genes. Each dot
represents the median expression level (FPKM) of target genes predicted based on

TFBSs of a TF in closed chromatin regions in promoters (x-axis) and an EPA domain

limited within the FR orientation of CTCF-binding sites (y-axis) in T cells (upper left

graph). The distribution of expression level was changed according to four criteria of

EPA domains (four graphs) [11]. Red dots indicate the median expression level of target

genes is significantly higher in an EPA domain than that in closed chromatin regions,

and blue dots indicate the median expression level of target genes is significantly lower

in an EPA domain than that in closed chromatin regions (Mann-Whitney test). The

expression level of target genes predicted using an EPA domain tend to be higher than

those predicted using closed chromatin regions, thereby implying that TFs serve as

activators of target genes. The ratio of median expression level of target genes between

an EPA domain and closed chromatin regions according to the four types of EPA

domains is shown in terms of probability density and cumulative distribution (middle

right two graphs). TFBSs overlapping H3K27ac ChIP-seq peaks were associated with a

significant difference of expression level of target genes, and TFBSs not overlapping

H3K27ac peaks were not associated with the difference (lower two graphs). This

suggests that the significant difference of expression level is associated with the

enhancer (and promoter) signals of H3K27ac. **b** Distribution of TFBSs of TFs near TSS.

ZNF143 and YY1 tended to be located near TSS, and CTCF tended to be observed

more in distant regions than those around TSS in monocytes. **c** Distribution of TFBSs of

TFs in the human genome in monocytes. Most of TFBSs were found in intergenic and

intron regions. YY1 is known to be at the TSS in the edge of exons of genes and is

involved in transcriptional activation, repression, and initiation.

Fig. 3 Biased orientations of TFBSs of TFs affect the transcription of genes by the

putative insulator function of the TFs. Ninety-six (sixty-four FR and fifty-two RF)

biased orientations of TFBSs of TFs showed significant differences in the distributions

of the ratios of expression level among FR, RF, and non-orientation in monocytes, T

cells, HMEC, and NPC, whereas non-oriented TFBSs of TFs did not show significant

differences in expression level.

Fig. 4 Genes separated by predicted TFBSs of biased orientations of TFs are less

correlated in gene expression. The top 20 TFs are selected according to ascending order

of the median correlation coefficient. The correlation coefficient between neighboring

gene pairs is shown in terms of probability density **a** and cumulative distribution **b**. The

brown line (All) represents the correlation between all neighboring genes; gray shading

in **a** and red line (random) in **b** represents correlation between randomly chosen gene

pairs; and the purple line (no_CTCF) represents the correlation between gene pairs not

separated by predicted CTCF-binding sites.

Tables

Table 1 Biased orientations of DNA-binding motifs of TFs in common between monocytes, T cells, HMEC, and NPC. TF: DNA-binding motifs of transcription factors.

Forward-reverse orientation

TF	P-value	TF	P-value	TF	P-value
AHR	0	MYC	0	**ZNF143**	0
ARNT	0	MYCN	0	ZNF740	0
ARNTL	7.96E-08	MZF1	0	ZNF770	6.51E-08
ATF2	3.09E-14	NFY	1.72E-11	ZNF84	0
ATF3	2.22E-15	NFYB	0		
BCL	0	NONO	0		
BHLHE41	5.55E-14	NRF1	1.63E-11		
CREB3L1	0	OVOL2	0		
CTCF	0	PLAG1	0		
CXXC1	0	PLAGL1	0		
E2F	0	**RAD21**	0		
E2F1	0	REST	0		
EGR1	0	RFX1	0		
EGR3	0	SIN3A	0		
EGR4	0	SIX5	0		
ELF1	0	**SMC3**	0		
ELF2	5.74E-13	SP2	1.04E-11		
ELK3	3.33E-15	TFAP2A	0		
ELK4	6.19E-11	TFAP2B	2.22E-16		
ERG	0	TFAP2C	0		
ETS	0	TP73	0		
ETS1	0	USF1	0		
ETV7	2.31E-08	XBP1	0		
HEY2	0	**YY1**	0		
HIC1	0	YY2	0		
HINFP	6.46E-14	ZBTB33	0		
INSM1	2.22E-16	ZBTB7A	0		
KLF4	0	ZFP62	0		
KLF7	9.07E-09	ZFX	0		
MAX	0	ZIC1	0		

Reverse-forward orientation

TF	P-value	TF	P-value
AP1	0	RFX2	0
ARNT	1.26E-11	RFX5	0
ATF3	0	RUNX3	0
BHLHE40	8.88E-16	SP1	0
CREB1	0	SP2	0
CTCF	0	SP4	0
E2F	2.32E-13	SPDEF	0
E2F3	0	SREBF2	0
EFNA2	6.61E-11	STAT	0
EGR1	0	TATA	0
EGR2	0	TFAP2B	2.00E-15
ETS1	0	TFAP2C	0
FLI1	1.05E-13	TFCP2L1	0
FLI1:FIGLA	0	USF1	7.55E-10
FOS	0	YBX1	0
GMEB2	0	**YY1**	0
HIF1A	0	YY2	0
KLF4	0	ZBTB33	0
MITF	0	ZBTB7B	0
MYB	0	ZNF100	6.22E-14
MYC	0	**ZNF143**	0
MYCN	0	ZNF432	0
MZF1	0		
NFKB1	0		
NFYA	2.31E-13		
NR3C1	0		
NRF1	0		
OSR2	0		
PBX3	0		
PLAG1	2.78E-14		

Table 2 Comparison of enhancer-promoter interactions (EPIs) with chromatin interactions in T cells. The upper tables show the number of biased orientations of DNA motifs, where a significantly higher ratio of EPIs that were predicted using an EPA domain (i) overlapped with HiChIP chromatin interactions than EPA domain (ii) and (iii) in T cells. The lower tables show that among 64 FR and 52 RF orientations of DNA

motifs found in common in monocytes, T cells, HMEC, and NPC, 44 (69%) FR and 28

(54%) RF were matched with the analysis of HiChIP chromatin interactions according

to three types of EPA domains. TF: DNA-binding motifs of a transcription factor. Score:

$-\log_{10}$ (p-value). Score 1: Comparison of EPA domain (i) (an EPA domain limited

within the FR or RF orientation of TFBSs) with EPA domain (iii) (an EPA domain not

limited). Score 2: Comparison of EPA domain (i) with EPA domain (ii) (an EPA domain

limited within the non-oriented TFBSs).

1. Comparison of EPI with HiChIP data among three EPA (i), (ii), and (iii)

Replication of HiChIP data	Total no. (FR + RF) of DNA motifs	No. of FR DNA motifs	No. of RF DNA motifs
B2T1	226	153	101
B2T2	221	149	98
B3T1	237	160	109
DNA motifs found in common among replications			
B2T1 and B2T2	212	141	95
B2T1 and B3T1	212	143	93
B2T2 and B3T1	208	138	93
B2T1, B2T2, and B3T1	201	133	90

2. Comparison of EPI with HiChIP data between two EPA (i) and (iii)

Replication of HiChIP data	Total no. (FR + RF) of DNA motifs	No. of FR DNA motifs	No. of RF DNA motifs
B2T1	395	262	205
B2T2	398	264	204
B3T1	394	263	203
DNA motifs found in common among replications			
B2T1 and B2T2	394	262	202
B2T1 and B3T1	391	261	202
B2T2 and B3T1	392	262	200
B2T1, B2T2, and B3T1	390	261	199

721 FR and RF orientations of DNA motifs found to be common in the four cell types.

722 Forward-reverse (FR)   Reverse-forward (RF)

TF	Score 1	Score 2
AHR	38.81	13.41
ARNTL	155.78	1.96
ATF3	323.31	35.22
BCL	153.72	3.52
CREB3L1	323.31	27.96
CTCF	323.31	3.50
CXXC1	69.92	3.30
E2F	122.12	21.87
E2F1	43.22	2.85
EGR1	79.09	3.79
EGR3	85.25	34.18
ELF1	203.38	3.86
ELF2	227.62	7.83
ELK4	117.83	4.81
ERG	199.81	25.70
ETS	196.80	1.63
ETS1	308.00	3.92
HIC1	223.55	4.56
INSM1	193.85	15.87
KLF4	209.71	9.38
KLF7	229.95	11.96
MAX	323.31	8.35
MYC	84.79	24.93
MYCN	165.16	25.87
MZF1	81.83	2.29
NFY	323.31	3.42
NONO	323.31	9.24
NRF1	45.13	30.75
OVOL2	323.31	31.61
PLAG1	177.90	11.02

TF	Score 1	Score 2
RAD21	258.21	4.81
RFX1	159.78	17.05
SMC3	222.09	1.76
SP2	94.11	9.54
TFAP2A	25.86	20.26
TFAP2B	90.03	6.10
TFAP2C	169.60	8.55
TP73	198.45	7.75
USF1	194.06	3.56
XBP1	323.31	44.58
YY1	62.59	5.30
ZBTB7A	52.89	30.58
ZFX	13.33	14.55
ZNF143	323.31	35.38

TF	Score 1	Score 2
ARNT	323.31	14.12
ATF3	323.31	14.64
BHLHE40	323.31	12.42
CTCF	240.61	4.56
E2F	78.16	7.07
E2F3	65.67	8.85
EFNA2	255.01	10.95
GMEB2	323.31	7.81
KLF4	213.35	9.49
MYB	323.31	17.68
MYC	323.31	12.23
MZF1	200.49	22.73
NFYA	323.31	11.75
NR3C1	43.16	8.31
OSR2	111.77	14.65
SP2	294.86	2.59
SP4	110.94	19.63
SPDEF	185.85	14.35
SREBF2	297.27	1.62
TATA	2.16	4.02
TFAP2B	29.46	12.62
TFAP2C	112.10	51.21
YY1	323.31	9.92
YY2	47.49	2.61
ZBTB33	90.29	7.78
ZBTB7B	43.90	23.59
ZNF100	27.67	4.75
ZNF143	63.03	9.69

Table 3 Correlation of the expression level of gene pairs separated by TFBSs of a TF.

Pairs of genes separated by 64 FR and 52 RF orientations of TFBSs of TFs found to be

common between monocytes, T cells, HMEC, and NPC were analyzed to find

significantly lower correlation of their expression level compared to the correlation of

all pairs of nearby genes among 53 tissues. Score: $-\log_{10}$ (p-value) of the

Mann-Whitney test. Median correlation: median correlation coefficient of expression

level of nearby gene pairs separated by TFBSs. The median correlation coefficient of

expression level of all pairs of nearby genes was 0.17. # TFBS: number of TFBSs of a

TF separating gene pairs with expression data. All TFBSs of a TF were used, except for

TFBSs separating pairs of ubiquitously expressed genes with coefficient of variance <

90.

Forward-reverse (FR)   Reverse-forward (RF)

TF	Score	Median correlation	# TFBS	TF	Score	Median correlation	# TFBS
AHR	1.80	0.12	520	AP1	3.87	0.11	744
ARNT	1.35	0.09	141	ARNT	2.95	0.10	453
ATF2	1.40	0.11	166	ATF3	4.34	0.10	665
ATF3	3.61	0.11	795	BHLHE40	2.49	0.07	155
BHLHE41	2.02	0.07	118	CREB1	1.79	0.11	303
CREB3L1	2.88	0.05	127	**CTCF**	5.96	0.09	721
CTCF	6.02	0.09	837	E2F	3.95	0.10	663
E2F	3.73	0.10	444	E2F3	4.05	0.09	616
E2F1	3.79	0.11	868	EFNA2	3.51	0.10	543
EGR1	2.97	0.10	391	EGR1	2.58	0.10	428
ELF1	4.86	0.10	714	EGR2	1.39	0.13	747
ELF2	3.88	0.11	781	ETS1	3.75	0.11	667
ELK3	1.86	0.12	547	FLI1:FIGLA	5.41	0.08	428
ELK4	2.44	0.11	663	FOS	3.51	0.11	777
ETS	3.34	0.11	523	GMEB2	1.36	0.09	310
ETS1	4.27	0.11	687	HIF1A	2.34	0.11	448
ETV7	3.68	0.11	662	KLF4	2.21	0.12	744
HIC1	1.65	0.11	380	MITF	3.14	0.10	419
HINFP	1.33	0.09	258	MYC	4.85	0.10	704
KLF4	2.00	0.13	718	MYCN	2.79	0.10	507
MYC	3.86	0.11	762	NFKB1	1.59	0.11	342
MYCN	2.36	0.11	583	NFYA	2.15	0.10	229
MZF1	1.74	0.10	261	NRF1	2.01	0.12	615
NFYB	2.97	0.10	264	OSR2	1.99	0.12	301
NRF1	2.21	0.11	640	PBX3	2.20	0.11	253
PLAG1	2.93	0.10	503	RFX2	2.12	0.09	168
PLAGL1	2.55	0.10	414	RFX5	3.03	0.05	75
RAD21	6.08	0.09	852	RUNX3	2.01	0.11	241
REST	4.35	0.11	920	SP1	2.73	0.11	725
RFX1	2.38	0.08	81	SP2	2.15	0.12	1090
SIN3A	3.31	0.11	813	SP4	2.33	0.11	479
SMC3	6.18	0.09	625	SPDEF	5.29	0.07	449
SP2	2.30	0.12	738	TFAP2B	3.42	0.11	855
TFAP2A	4.17	0.11	767	TFAP2C	2.05	0.11	576
TFAP2C	2.13	0.11	470	TFCP2L1	1.45	0.13	362
USF1	4.24	0.10	628	YBX1	1.34	0.13	160
XBP1	6.65	0.07	379	**YY1**	2.23	0.01	50
ZBTB33	4.90	0.10	701	ZBTB33	3.44	0.11	886
ZBTB7A	2.82	0.10	485	ZBTB7B	2.27	0.13	892
ZFP62	2.95	0.11	565	**ZNF143**	2.26	0.07	81
ZFX	2.69	0.11	581				
ZIC1	4.47	0.11	829				
ZNF770	3.38	0.08	271				

Table 4 Top 20 of the co-localized and overlapping biased orientations of DNA-binding

motifs of TFs in monocytes. Co-localized and overlapping DNA motifs of CTCF with

another biased orientation of DNA motifs are shown in separate tables. Motif 1 and

Motif 2: DNA-binding motifs of TFs. No.: the number of genomic locations with both

Motif 1 and Motif 2 upstream and downstream of genes in an EPA domain (< 1 Mb

distance from TSS of genes). The total number of genes was 30,523.

Forward-reverse (FR)

Motif 1	Motif 2	No.	Mean of distances bet. DNA sites (nt)	Median of distances bet. DNA sites (nt)	Mean of distances from TSS (nt)	Median of distances from TSS (nt)	Interquartile range of distance from TSS (nt)
FOS	JUN	3332	0.00	0	43202.95	18085	5524 - 48583
CTCF	**SMC3**	2876	0.01	0	34606.92	13716	3073 - 39471
FOSL2	JUN	2807	0.00	0	40389.84	17365	5424 - 45251
FOSB	FOSL2	2644	0.00	0	44302.93	17867	5633 - 47209
EGR1	EGR2	2397	0.00	0	14578.21	1251	247 - 11473
CTCF	**RAD21**	2352	0.00	0	33492.68	12913	2824 - 37672
RAD21	**SMC3**	2343	0.02	0	33572.80	14302	3856 - 37159
ATF3	FOSL2	2322	0.00	0	42784.04	17841	5366 - 48236
FOSL2	JUNB	2305	0.02	0	42884.36	17867	5706 - 46515
ETS2	ETSLIKE	2205	0.00	0	37660.66	15765	4146 - 41462
FOS	JUNB	2151	0.00	0	38403.20	16367	4849 - 44736
IRF	IRF4	1962	0.00	0	40794.41	15872	3969 - 45946
FOSL2	SMARCC1	1945	0.00	0	42429.69	17369	5094 - 46586
EGR1	WT1	1941	0.16	0	12485.23	427	93 - 5394
FOSB	JUN	1808	0.02	0	36648.74	16705	5156 - 40309
ATF3	JUN	1713	0.02	0	38101.24	17119	5294 - 43559
JUN	TFAP2A	1628	0.00	0	42262.97	18017	5089 - 47094
FOSL2	TFAP2A	1628	0.00	0	36606.38	16201	4949 - 41948
BACH2	SMARCC1	1462	0.16	0	38673.02	14778	4060 - 41262
ELF1	ETS1	1459	0.00	0	37056.72	14346	3370 - 40339

751   Reverse-forward (RF)

Motif 1	Motif 2	No.	Mean of distances bet. DNA sites (nt)	Median of distances bet. DNA sites (nt)	Mean of distances from TSS (nt)	Median of distances from TSS (nt)	Interquartile range of distance from TSS (nt)
EGR1	EGR3	3012	0.00	0	17005.00	709	149 - 10393
NR1I2	NR1I3	2932	0.00	0	49622.84	20548	5924 - 58436
ZIC1	ZIC3	2871	0.07	0	21511.64	3092	434 - 19128
EGR	WT1	2612	0.04	0	9949.38	396	102 - 4266
FOS	FOS:JUN	2512	0.01	0	37200.80	15466	4438 - 42373
EGR	EGR3	2293	0.17	0	11634.92	532	135 - 6044
MAFG	NFE2L1:MAFG	2272	0.03	0	34641.56	16917	5284 - 42097
KLF10	SP1	2233	0.00	0	9778.94	297	73 - 3077
EGR	EGR2	2160	0.16	0	10502.76	490	129 - 5227
KLF3	SP3	2036	0.00	0	6974.78	246	63 - 1447
KLF1	SP3	1771	0.00	0	7748.42	264	66 - 1829
KLF10	SP3	1734	0.00	0	7275.48	222	58 - 1320
KLF7	SP4	1639	0.00	0	12521.62	309	58 - 4894
KLF1	SP1	1574	0.05	0	6780.46	245	61 - 1438
AP1	JUNB	1572	0.05	0	34158.07	15451	4746 - 40315
EGR1	WT1	1543	0.31	0	12742.28	444	111 - 5553
EGR1	SP1:SP3	1393	0.08	0	7789.54	307	91 - 1886
FOS:JUN	JUNB	1386	0.14	0	33012.80	14245	3900 - 37988
SP1	SP3	1372	0.00	0	8845.31	293	77 - 1816
EHF	ELF1	1366	0.02	0	41217.39	13357	2842 - 43178

Co-localized and overlapping DNA motifs of CTCF with other DNA motifs.

FR orientation of CTCF

Motif 1	Motif 2	No.	Mean of distances bet. DNA sites (nt)	Median of distances bet. DNA sites (nt)	Mean of distances from TSS (nt)	Median of distances from TSS (nt)	Interquartile range of distance from TSS (nt)
CTCF	SMC3	2876	0.01	0	34606.92	13716	3073 - 39471
CTCF	RAD21	2352	0.00	0	33492.68	12913	2824 - 37672
CTCF	CTCFL	1151	0.00	0	30252.04	11606	2229 - 34539
CTCF	ZBTB7A	470	0.27	0	23227.96	7065	745 - 23496
CTCF	E2F	123	0.81	0	7913.36	222	61 - 1997
CTCF	WT1	120	0.00	0	3543.72	114	44 - 375
CTCF	MYC	110	2.30	0	31289.45	11948	2887 - 33972

RF orientation of CTCF

Motif 1	Motif 2	No.	Mean of distances bet. DNA sites (nt)	Median of distances bet. DNA sites (nt)	Mean of distances from TSS (nt)	Median of distances from TSS (nt)	Interquartile range of distance from TSS (nt)
CTCF	RXRA	1362	0.00	0	29981.33	10937	2100 - 31407
CTCF	CTCFL	429	0.00	0	25171.22	9834	1859 - 24741
CTCF	RAD21	337	0.00	0	38829.91	17973	6117 - 45749
CTCF	ZIC1	126	1.40	0	35480.65	10286	2316 - 29380
CTCF	ZIC3	108	0.00	0	30219.32	10904	4463 - 31037

References

1. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES, Aiden EL: **A 3D map of the**

human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 2014, **159:**1665-1680.

2. Shlyueva D, Stampfel G, Stark A: **Transcriptional enhancers: from properties to genome-wide predictions.** *Nat Rev Genet* 2014, **15:**272-286.

3. Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y, et al: **CRISPR Inversion of CTCF Sites Alters Genome Topology and Enhancer/Promoter Function.** *Cell* 2015, **162:**900-910.

4. de Wit E, Vos ES, Holwerda SJ, Valdes-Quezada C, Verstegen MJ, Teunissen H, Splinter E, Wijchers PJ, Krijger PH, de Laat W: **CTCF Binding Polarity Determines Chromatin Looping.** *Mol Cell* 2015, **60:**676-684.

5. Bailey SD, Zhang X, Desai K, Aid M, Corradin O, Cowper-Sal Lari R, Akhtar-Zaidi B, Scacheri PC, Haibe-Kains B, Lupien M: **ZNF143 provides sequence specificity to secure chromatin interactions at gene promoters.** *Nat Commun* 2015, **2:**6186.

6. Weintraub AS, Li CH, Zamudio AV, Sigova AA, Hannett NM, Day DS, Abraham BJ, Cohen MA, Nabet B, Buckley DL, et al: **YY1 Is a Structural Regulator of Enhancer-Promoter Loops.** *Cell* 2017, **171:**1573-1588 e1528.

7. Barutcu AR, Lajoie BR, Fritz AJ, McCord RP, Nickerson JA, van Wijnen AJ, Lian JB, Stein JL, Dekker J, Stein GS, Imbalzano AN: **SMARCA4 regulates gene expression and higher-order chromatin structure in proliferating mammary epithelial cells.** *Genome Res* 2016, **26:**1188-1201.

8. Schreiber J, Libbrecht M, Bilmes J, Noble W: **Nucleotide sequence and DNaseI sensitivity are predictive of 3D chromatin architecture.** *bioRxiv* 2017.

9. Zhang H, Li F, Jia Y, Xu B, Zhang Y, Li X, Zhang Z: **Characteristic arrangement of nucleosomes is predictive of chromatin interactions at kilobase resolution.** *Nucleic Acids Res* 2017, **45:**12739-12751.

10. Xie X, Lu J, Kulbokas EJ, Golub TR, Mootha V, Lindblad-Toh K, Lander ES, Kellis M: **Systematic discovery of regulatory motifs in human promoters and 3' UTRs by comparison of several mammals.** *Nature* 2005, **434:**338-345.

11. Osato N: **Characteristics of functional enrichment and gene expression level of human putative transcriptional target genes.** *BMC Genomics* 2018, **19:**957.

12.   McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM, Bejerano G: **GREAT improves functional interpretation of cis-regulatory regions.** *Nat Biotechnol* 2010, **28:**495-501.

13.   Arnold CD, Gerlach D, Stelzer C, Boryn LM, Rath M, Stark A: **Genome-wide quantitative enhancer activity maps identified by STARR-seq.** *Science* 2013, **339:**1074-1077.

14.   Xi H, Yu Y, Fu Y, Foley J, Halees A, Weng Z: **Analysis of overrepresented motifs in human core promoters reveals dual regulatory roles of YY1.** *Genome Res* 2007, **17:**798-806.

15.   Shrivastava A, Calame K: **An analysis of genes regulated by the multi-functional transcriptional regulator Yin Yang-1.** *Nucleic Acids Res* 1994, **22:**5151-5155.

16.   Di Giammartino DC, Kloetgen A, Polyzos A, Liu Y, Kim D, Murphy D, Abuhashem A, Cavaliere P, Aronson B, Shah V, et al: **KLF4 is involved in the organization and regulation of pluripotency-associated three-dimensional enhancer networks.** *Nat Cell Biol* 2019, **21:**1179-1190.

17.   Li F, Yuan Q, Di W, Xia X, Liu Z, Mao N, Li L, Li C, He J, Li Y, et al: **ERG orchestrates chromatin interactions to drive prostate cell fate reprogramming.** *J Clin Invest* 2020.

18.   Zhang Z, Chng KR, Lingadahalli S, Chen Z, Liu MH, Do HH, Cai S, Rinaldi N, Poh HM, Li G, et al: **An AR-ERG transcriptional signature defined by long-range chromatin interactomes in prostate cancer cells.** *Genome Res* 2019, **29:**223-235.

19.   Majumder P, Gomez JA, Chadwick BP, Boss JM: **The insulator factor CTCF controls MHC class II gene expression and is required for the formation of long-distance chromatin interactions.** *J Exp Med* 2008, **205:**785-798.

20.   Quigley IK, Kintner C: **Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops to Enable Multiciliated Cell Gene Expression.** *PLoS Genet* 2017, **13:**e1006538.

21.   Mimura I, Nangaku M, Kanki Y, Tsutsumi S, Inoue T, Kohro T, Yamamoto S, Fujita T, Shimamura T, Suehiro J, et al: **Dynamic change of chromatin conformation in response to hypoxia enhances the expression of GLUT3 (SLC2A3) by cooperative interaction of hypoxia-inducible factor 1 and KDM3A.** *Mol Cell Biol* 2012, **32:**3018-3032.

22.   Deshane J, Kim J, Bolisetty S, Hock TD, Hill-Kapturczak N, Agarwal A: **Sp1**
regulates chromatin looping between an intronic enhancer and distal
promoter of the human heme oxygenase-1 gene in renal cells. *J Biol Chem*
2010, **285:**16476-16486.

23.   Park JH, Choi Y, Song MJ, Park K, Lee JJ, Kim HP: **Dynamic Long-Range**
Chromatin Interaction Controls Expression of IL-21 in CD4+ T Cells. *J*
Immunol 2016, **196:**4378-4389.

24.   D'Ippolito AM, McDowell IC, Barrera A, Hong LK, Leichter SM, Bartelt LC,
Vockley CM, Majoros WH, Safi A, Song L, et al: **Pre-established Chromatin**
Interactions Mediate the Genomic Response to Glucocorticoids. *Cell Syst*
2018, **7:**146-160 e147.

25.   Kim DE, Procopio MG, Ghosh S, Jo SH, Goruppi S, Magliozzi F, Bordignon P,
Neel V, Angelino P, Dotto GP: **Convergent roles of ATF3 and CSL in**
chromatin control of cancer-associated fibroblast activation. *J Exp Med*
2017, **214:**2349-2368.

26.   Ramos Pittol JM, Oruba A, Mittler G, Saccani S, van Essen D: **Zbtb7a is a**
transducer for the control of promoter accessibility by NF-kappa B and
multiple other transcription factors. *PLoS Biol* 2018, **16:**e2004526.

27.   Sherwood RI, Hashimoto T, O'Donnell CW, Lewis S, Barkal AA, van Hoff JP,
Karun V, Jaakkola T, Gifford DK: **Discovery of directional and nondirectional**
pioneer transcription factors by modeling DNase profile magnitude and
shape. *Nat Biotechnol* 2014, **32:**171-178.

28.   Rothstein M, Simoes-Costa M: **Heterodimerization of TFAP2 pioneer factors**
drives epigenomic remodeling during neural crest specification. *Genome Res*
2020, **30:**35-48.

29.   Knoepfler PS, Zhang XY, Cheng PF, Gafken PR, McMahon SB, Eisenman RN:
Myc influences global chromatin structure. *EMBO J* 2006, **25:**2723-2734.

30.   Sabo A, Amati B: **Genome recognition by MYC.** *Cold Spring Harb Perspect*
Med 2014, **4**.

31.   Yang H, Schramek D, Adam RC, Keyes BE, Wang P, Zheng D, Fuchs E: **ETS**
family transcriptional regulators drive chromatin dynamics and
malignancy in squamous cell carcinomas. *Elife* 2015, **4:**e10870.

32.   Li L, Bhatia R: **The controversial role of Sirtuins in tumorigenesis - SIRT7**

joins the debate. *Cell Res* 2013, **23:**10-12.

33. Lee JW, Kim DM, Jang JW, Park TG, Song SH, Lee YS, Chi XZ, Park IY, Hyun
JW, Ito Y, Bae SC: **RUNX3 regulates cell cycle-dependent chromatin**
dynamics by functioning as a pioneer factor of the restriction-point. *Nat*
Commun 2019, **10:**1897.

34. Huang S, Li X, Yusufzai TM, Qiu Y, Felsenfeld G: **USF1 recruits histone**
modification complexes and is critical for maintenance of a chromatin
barrier. *Mol Cell Biol* 2007, **27:**7991-8002.

35. Li Y, Schulz VP, Deng C, Li G, Shen Y, Tusi BK, Ma G, Stees J, Qiu Y, Steiner
LA, et al: **Setd1a and NURF mediate chromatin dynamics and gene**
regulation during erythroid lineage commitment and differentiation.
Nucleic Acids Res 2016, **44:**7173-7188.

36. Swanson HI: **DNA binding and protein interactions of the AHR/ARNT**
heterodimer that facilitate gene activation. *Chem Biol Interact* 2002,
141:63-76.

37. Volkel S, Stielow B, Finkernagel F, Stiewe T, Nist A, Suske G: **Zinc finger**
independent genome-wide binding of Sp2 potentiates recruitment of
histone-fold protein Nf-y distinguishing it from Sp1 and Sp3. *PLoS Genet*
2015, **11:**e1005102.

38. Frob F, Wegner M: **The role of chromatin remodeling complexes in Schwann**
cell development. *Glia* 2020, **68:**1596-1603.

39. Wust HM, Wegener A, Frob F, Hartwig AC, Wegwitz F, Kari V, Schimmel M,
Tamm ER, Johnsen SA, Wegner M, Sock E: **Egr2-guided histone H2B**
monoubiquitination is required for peripheral nervous system myelination.
Nucleic Acids Res 2020.

40. Mager GM, Ward RM, Srinivasan R, Jang SW, Wrabetz L, Svaren J: **Active**
gene repression by the Egr2.NAB complex during peripheral nerve
myelination. *J Biol Chem* 2008, **283:**18187-18197.

41. Blattler A, Yao L, Wang Y, Ye Z, Jin VX, Farnham PJ: **ZBTB33 binds**
unmethylated regions of the genome associated with actively expressed
genes. *Epigenetics Chromatin* 2013, **6:**13.

42. Oldfield AJ, Yang P, Conway AE, Cinghu S, Freudenberg JM, Yellaboina S,
Jothi R: **Histone-fold domain protein NF-Y promotes chromatin accessibility**

for cell type-specific master transcription factors. *Mol Cell* 2014,
55:708-722.

43.   Nardini M, Gnesutta N, Donati G, Gatta R, Forni C, Fossati A, Vonrhein C,
Moras D, Romier C, Bolognesi M, Mantovani R: **Sequence-specific**
transcription factor NF-Y displays histone-like DNA binding and H2B-like
ubiquitination. *Cell* 2013, **152:**132-143.

44.   Laurette P, Strub T, Koludrovic D, Keime C, Le Gras S, Seberg H, Van Otterloo
E, Imrichova H, Siddaway R, Aerts S, et al: **Transcription factor MITF and**
remodeller BRG1 define chromatin organisation at regulatory elements in
melanoma cells. *Elife* 2015, **4**.

45.   Levy C, Fisher DE: **Dual roles of lineage restricted transcription factors: the**
case of MITF in melanocytes. *Transcription* 2011, **2:**19-22.

46.   Ma C, Karwacki-Neisius V, Tang H, Li W, Shi Z, Hu H, Xu W, Wang Z, Kong L,
Lv R, et al: **Nono, a Bivalent Domain Factor, Regulates Erk Signaling and**
Mouse Embryonic Stem Cell Pluripotency. *Cell Rep* 2016, **17:**997-1007.

47.   Riggi N, Knoechel B, Gillespie SM, Rheinbay E, Boulay G, Suva ML, Rossetti
NE, Boonseng WE, Oksuz O, Cook EB, et al: **EWS-FLI1 utilizes divergent**
chromatin remodeling mechanisms to directly activate or repress enhancer
elements in Ewing sarcoma. *Cancer Cell* 2014, **26:**668-681.

48.   Blais A, Dynlacht BD: **E2F-associated chromatin modifiers and cell cycle**
control. *Curr Opin Cell Biol* 2007, **19:**658-662.

49.   Taubert S, Gorrini C, Frank SR, Parisi T, Fuchs M, Chan HM, Livingston DM,
Amati B: **E2F-dependent histone acetylation and recruitment of the Tip60**
acetyltransferase complex to chromatin in late G1. *Mol Cell Biol* 2004,
24:4546-4556.

50.   Subtil-Rodriguez A, Vazquez-Chavez E, Ceballos-Chavez M,
Rodriguez-Paredes M, Martin-Subero JI, Esteller M, Reyes JC: **The chromatin**
remodeller CHD8 is required for E2F-dependent transcription activation of
S-phase genes. *Nucleic Acids Res* 2014, **42:**2185-2196.

51.   Choi J, Jeon S, Choi S, Park K, Seong RH: **The SWI/SNF chromatin**
remodeling complex regulates germinal center formation by repressing
Blimp-1 expression. *Proc Natl Acad Sci U S A* 2015, **112:**E718-727.

52.   Smith E, Shilatifard A: **The chromatin signaling pathway: diverse**

mechanisms of recruitment of histone-modifying enzymes and varied
biological outcomes. *Mol Cell* 2010, **40:**689-701.

53.   Magri L, Swiss VA, Jablonska B, Lei L, Pedre X, Walsh M, Zhang W, Gallo V,
Canoll P, Casaccia P: **E2F1 coregulates cell cycle genes and chromatin**
components during the transition of oligodendrocyte progenitors from
proliferation to differentiation. *J Neurosci* 2014, **34:**1481-1493.

54.   Sun Z, Xu X, He J, Murray A, Sun MA, Wei X, Wang X, McCoig E, Xie E,
Jiang X, et al: **EGR1 recruits TET1 to shape the brain methylome during**
development and upon neuronal activity. *Nat Commun* 2019, **10:**3892.

55.   Ooi L, Belyaev ND, Miyake K, Wood IC, Buckley NJ: **BRG1 chromatin**
remodeling activity is required for efficient chromatin binding by repressor
element 1-silencing transcription factor (REST) and facilitates
REST-mediated repression. *J Biol Chem* 2006, **281:**38974-38980.

56.   Cunliffe VT: **Eloquent silence: developmental functions of Class I histone**
deacetylases. *Curr Opin Genet Dev* 2008, **18:**404-410.

57.   Xu Y, Sengupta PK, Seto E, Smith BD: **Regulatory factor for X-box family**
proteins differentially interact with histone deacetylases to repress collagen
alpha2(I) gene (COL1A2) expression. *J Biol Chem* 2006, **281:**9260-9270.

58.   Saunders A, Huang X, Fidalgo M, Reimer MH, Jr., Faiola F, Ding J,
Sanchez-Priego C, Guallar D, Saenz C, Li D, Wang J: **The SIN3A/HDAC**
Corepressor Complex Functionally Cooperates with NANOG to Promote
Pluripotency. *Cell Rep* 2017, **18:**1713-1726.

59.   Salas-Armenteros I, Perez-Calero C, Bayona-Feliu A, Tumini E, Luna R,
Aguilera A: **Human THO-Sin3A interaction reveals new mechanisms to**
prevent R-loops that cause genome instability. *EMBO J* 2017, **36:**3532-3547.

60.   Adams GE, Chandru A, Cowley SM: **Co-repressor, co-activator and general**
transcription factor: the many faces of the Sin3 histone deacetylase (HDAC)
complex. *Biochem J* 2018, **475:**3921-3932.

61.   Biddie SC, John S, Sabo PJ, Thurman RE, Johnson TA, Schiltz RL, Miranda TB,
Sung MH, Trump S, Lightman SL, et al: **Transcription factor AP1 potentiates**
chromatin accessibility and glucocorticoid receptor binding. *Mol Cell* 2011,
43:145-155.

62.   Orlova A, Wagner C, de Araujo ED, Bajusz D, Neubauer HA, Herling M,

Gunning PT, Keseru GM, Moriggl R: **Direct Targeting Options for STAT3**
and STAT5 in Cancer. *Cancers (Basel)* 2019, **11**.

63.   Mumbach MR, Satpathy AT, Boyle EA, Dai C, Gowen BG, Cho SW, Nguyen
ML, Rubin AJ, Granja JM, Kazane KR, et al: **Enhancer connectome in**
primary human cells identifies target genes of disease-associated DNA
elements. *Nat Genet* 2017, **49:**1602-1612.

64.   Yardimci GG, Ozadam H, Sauria MEG, Ursu O, Yan KK, Yang T, Chakraborty
A, Kaul A, Lajoie BR, Song F, et al: **Measuring the reproducibility and**
quality of Hi-C data. *Genome Biol* 2019, **20:**57.

65.   Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T,
Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO, et al: **Comprehensive**
mapping of long-range interactions reveals folding principles of the human
genome. *Science* 2009, **326:**289-293.

66.   Jung I, Schmitt A, Diao Y, Lee AJ, Liu T, Yang D, Tan C, Eom J, Chan M, Chee
S, et al: **A compendium of promoter-centered long-range chromatin**
interactions in the human genome. *Nat Genet* 2019, **51:**1442-1449.

67.   Krietenstein N, Abraham S, Venev SV, Abdennur N, Gibcus J, Hsieh TS, Parsi
KM, Yang L, Maehr R, Mirny LA, et al: **Ultrastructural Details of**
Mammalian Chromosome Architecture. *Mol Cell* 2020, **78:**554-565 e557.

68.   Purmann A, Toedling J, Schueler M, Carninci P, Lehrach H, Hayashizaki Y,
Huber W, Sperling S: **Genomic organization of transcriptomes in mammals:**
Coregulation and cofunctionality. *Genomics* 2007, **89:**580-587.

69.   Xie X, Mikkelsen TS, Gnirke A, Lindblad-Toh K, Kellis M, Lander ES:
Systematic discovery of regulatory motifs in conserved regions of the
human genome, including thousands of CTCF insulator sites. *Proc Natl*
Acad Sci U S A 2007, **104:**7145-7150.

70.   Ansari AZ, Peterson-Kaufman KJ: **A partner evokes latent differences**
between Hox proteins. *Cell* 2011, **147:**1220-1221.

71.   Slattery M, Riley T, Liu P, Abe N, Gomez-Alcala P, Dror I, Zhou T, Rohs R,
Honig B, Bussemaker HJ, Mann RS: **Cofactor binding evokes latent**
differences in DNA binding specificity between Hox proteins. *Cell* 2011,
147:1270-1282.

72.   Merabet S, Mann RS: **To Be Specific or Not: The Critical Relationship**

Between Hox And TALE Proteins. *Trends Genet* 2016, **32:**334-347.

73.   Yoon SO, Chikaraishi DM: **Tissue-specific transcription of the rat tyrosine**
hydroxylase gene requires synergy between an AP-1 motif and an
overlapping E box-containing dyad. *Neuron* 1992, **9:**55-67.

74.   Ackerman SL, Minden AG, Williams GT, Bobonis C, Yeung CY: **Functional**
significance of an overlapping consensus binding motif for Sp1 and Zif268
in the murine adenosine deaminase gene promoter. *Proc Natl Acad Sci U S A*
1991, **88:**7523-7527.

75.   Chatterjee R, Zhao J, He X, Shlyakhtenko A, Mann I, Waterfall JJ, Meltzer P,
Sathyanarayana BK, FitzGerald PC, Vinson C: **Overlapping ETS and CRE**
Motifs ((G/C)CGGAAGTGACGTCA) preferentially bound by GABPalpha
and CREB proteins. *G3 (Bethesda)* 2012, **2:**1243-1256.

76.   He X, Syed KS, Tillo D, Mann I, Weirauch MT, Vinson C: **GABPalpha**
Binding to Overlapping ETS and CRE DNA Motifs Is Enhanced by
CREB1: Custom DNA Microarrays. *G3 (Bethesda)* 2015, **5:**1909-1918.

77.   Kin K, Chen X, Gonzalez-Garay M, Fakhouri WD: **The effect of non-coding**
DNA variations on P53 and cMYC competitive inhibition at cis-overlapping
motifs. *Hum Mol Genet* 2016, **25:**1517-1527.

78.   Wei Z, Gao F, Kim S, Yang H, Lyu J, An W, Wang K, Lu W: **Klf4 organizes**
long-range chromosomal interactions with the oct4 locus in reprogramming
and pluripotency. *Cell Stem Cell* 2013, **13:**36-47.

79.   Hsu SC, Gilgenast TG, Bartman CR, Edwards CR, Stonestrom AJ, Huang P,
Emerson DJ, Evans P, Werner MT, Keller CA, et al: **The BET Protein BRD2**
Cooperates with CTCF to Enforce Transcriptional and Architectural
Boundaries. *Mol Cell* 2017, **66:**102-116 e107.

80.   Wang J, Zhuang J, Iyer S, Lin X, Whitfield TW, Greven MC, Pierce BG, Dong
X, Kundaje A, Cheng Y, et al: **Sequence features and chromatin structure**
around the genomic regions bound by 119 human transcription factors.
Genome Res 2012, **22:**1798-1812.

81.   Wang ET, Sandberg R, Luo S, Khrebtukova I, Zhang L, Mayr C, Kingsmore SF,
Schroth GP, Burge CB: **Alternative isoform regulation in human tissue**
transcriptomes. *Nature* 2008, **456:**470-476.

82.   Chen M, Manley JL: **Mechanisms of alternative splicing regulation: insights**

from molecular and genomics approaches. *Nat Rev Mol Cell Biol* 2009,
10:741-754.

83. Das D, Clark TA, Schweitzer A, Yamamoto M, Marr H, Arribere J, Minovitsky S,
Poliakov A, Dubchak I, Blume JE, Conboy JG: **A correlation with exon**
expression approach to identify cis-regulatory elements for tissue-specific
alternative splicing. *Nucleic Acids Res* 2007, **35:**4845-4857.

84. Tabuchi TM, Deplancke B, Osato N, Zhu LJ, Barrasa MI, Harrison MM, Horvitz
HR, Walhout AJ, Hagstrom KA: **Chromosome-biased binding and gene**
regulation by the Caenorhabditis elegans DRM complex. *PLoS Genet* 2011,
7:e1002074.

85. Jolma A, Yin Y, Nitta KR, Dave K, Popov A, Taipale M, Enge M, Kivioja T,
Morgunova E, Taipale J: **DNA-dependent formation of transcription factor**
pairs alters their binding specificity. *Nature* 2015, **527:**384-388.

86. Wang J, Vicente-Garcia C, Seruggia D, Molto E, Fernandez-Minan A, Neto A,
Lee E, Gomez-Skarmeta JL, Montoliu L, Lunyak VV, Jordan IK: **MIR**
retrotransposon sequences provide insulators to the human genome. *Proc*
Natl Acad Sci U S A 2015, **112:**E4428-4437.

87. Katainen R, Dave K, Pitkanen E, Palin K, Kivioja T, Valimaki N, Gylfe AE,
Ristolainen H, Hanninen UA, Cajuso T, et al: **CTCF/cohesin-binding sites are**
frequently mutated in cancer. *Nat Genet* 2015, **47:**818-821.

88. Wingender E, Dietze P, Karas H, Knuppel R: **TRANSFAC: a database on**
transcription factors and their DNA binding sites. *Nucleic Acids Res* 1996,
24:238-241.

89. Portales-Casamar E, Thongjuea S, Kwon AT, Arenillas D, Zhao X, Valen E,
Yusuf D, Lenhard B, Wasserman WW, Sandelin A: **JASPAR 2010: the greatly**
expanded open-access database of transcription factor binding profiles.
Nucleic Acids Res 2010, **38:**D105-110.

90. Newburger DE, Bulyk ML: **UniPROBE: an online database of protein**
binding microarray data on protein-DNA interactions. *Nucleic Acids Res*
2009, **37:**D77-82.

91. Zhao Y, Stormo GD: **Quantitative analysis demonstrates most transcription**
factors require only simple models of specificity. *Nat Biotechnol* 2011,
29:480-483.

92. Jolma A, Yan J, Whitington T, Toivonen J, Nitta KR, Rastas P, Morgunova E,
Enge M, Taipale M, Wei G, et al: **DNA-binding specificities of human**
transcription factors. *Cell* 2013, **152:**327-339.

93. Kheradpour P, Kellis M: **Systematic discovery and characterization of**
regulatory motifs in ENCODE TF binding experiments. *Nucleic Acids Res*
2014, **42:**2976-2987.

94. Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD,
Rumynskiy EI, Medvedeva YA, Magana-Mora A, Bajic VB, Papatsenko DA, et
al: **HOCOMOCO: towards a complete collection of transcription factor**
binding models for human and mouse via large-scale ChIP-Seq analysis.
Nucleic Acids Res 2018, **46:**D252-D259.

95. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum
C, Myers RM, Brown M, Li W, Liu XS: **Model-based analysis of ChIP-Seq**
(MACS). *Genome Biol* 2008, **9:**R137.

96. van Helden J: **Regulatory sequence analysis tools.** *Nucleic Acids Res* 2003,
31:3593-3596.

97. Kel AE, Gossling E, Reuter I, Cheremushkin E, Kel-Margoulis OV, Wingender
E: **MATCH: A tool for searching transcription factor binding sites in DNA**
sequences. *Nucleic Acids Res* 2003, **31:**3576-3579.

98. Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, Dreszer
TR, Fujita PA, Guruvadoo L, Haeussler M, et al: **The UCSC Genome Browser**
database: 2014 update. *Nucleic Acids Res* 2014, **42:**D764-770.

99. Li H, Durbin R: **Fast and accurate short read alignment with**
Burrows-Wheeler transform. *Bioinformatics* 2009, **25:**1754-1760.

100. Wang L, Wang S, Li W: **RSeQC: quality control of RNA-seq experiments.**
Bioinformatics 2012, **28:**2184-2185.

101. Neph S, Kuehn MS, Reynolds AP, Haugen E, Thurman RE, Johnson AK, Rynes
E, Maurano MT, Vierstra J, Thomas S, et al: **BEDOPS: high-performance**
genomic feature operations. *Bioinformatics* 2012, **28:**1919-1920.

1092