Opposing roles of endosomal innate immunity proteins IFITM3 and TLR7 in human metapneumovirus infection

Temet M. McMichael[1,3], Yu Zhang[2,a], Adam D. Kenney[1,3], Lizhi Zhang[1,3], Mijia Lu[2,3], Mahesh Chemudupati[1,3], Jianrong Li[2,3], and Jacob S. Yount[1,3]

[1]Department of Microbial Infection and Immunity, [2]Department of Veterinary Biosciences, and [3]Infectious Diseases Institute, The Ohio State University, Columbus, OH

[a]Current affiliation: University of Pittsburgh, Pittsburgh, PA

Correspondence: J. Yount, PhD, Department of Microbial Infection and Immunity, the Ohio State University, 460 W 12th Ave, Biomedical Research Tower 790, Columbus, OH 43210 (yount.37@osu.edu).

ABSTRACT

Human metapneumovirus (hMPV) utilizes a bifurcated cellular entry strategy, fusing either with the plasma membrane or, after endocytosis, with the endosome membrane. Whether cellular factors restrict or enhance either entry pathway is largely unknown. We found that the interferon-induced transmembrane protein 3 (IFITM3) inhibits hMPV infection to an extent similar to endocytosis-inhibiting drugs, and an IFITM3 variant that accumulates at the plasma membrane in addition to its endosome localization provided increased virus restriction. Mechanistically, IFITM3 blocks hMPV F protein-mediated membrane fusion, and inhibition of infection was reversed by the membrane destabilizing drug amphotericin B. Conversely, we unexpectedly found that infection by some hMPV strains is enhanced by Toll-like receptor 7 (TLR7), an endosomal protein, suggesting that cellular entry via endocytosis may be particularly advantageous for hMPV despite eventual restriction of this pathway upon induction of IFITM3. Overall, our results identify IFITM3 and TLR7 as endosomal factors differentially regulating hMPV infection.

INTRODUCTION

Human metapneumovirus (hMPV) is a member of the *Pneumoviridae* family, a recent

classification distinguishing it from related *Paramyxoviridae* family members[1]. Since its

discovery in 2001[2], hMPV has been increasingly recognized as a significant cause of

respiratory infections in infants, children, and the elderly[3-6]. This virus infects over 85%

of the population by age ten[7], and is one of the most common causes of respiratory tract

infections in infants[3, 5]. The frequency of detection of hMPV in elderly patients with

respiratory infections approaches that of influenza virus[4]. However, unlike influenza

virus, there are currently no licensed vaccines or drugs targeting hMPV. While vaccination

strategies have been proposed for eliciting an adaptive immune response against hMPV[8,

9], roles for innate immune restriction factors in controlling the virus remain to be

identified.

The early classification of hMPV as a paramyxovirus suggested that it would exhibit

pH-independent fusion at the plasma membrane similarly to other prototypical

paramyxoviruses, such as Sendai virus (SeV)[10]. However, early reports indicated that

acidic pH enhanced fusion mediated by the F proteins of certain hMPV strains[11, 12].

Follow up studies utilizing single virus particle tracking and fusion assays showed that a

portion of hMPV fuses at the plasma membrane as is typical of paramyxoviruses, but also

that a significant fraction of the virus is endocytosed and fuses with endosomal membranes

in a manner characteristic of low pH-dependent viruses, such as influenza A virus (IAV)[13-

15]. These data suggested that hMPV utilizes a bifurcated cellular entry strategy with

characteristics typified by both SeV and IAV. From this, we reasoned that hMPV may be

susceptible to inhibition by interferon-induced transmembrane protein 3 (IFITM3), which

is a cellular antiviral factor that inhibits virus-endosome membrane fusion reactions[16-21].

IFITM3 broadly restricts endocytosed virus infections by blocking their fusion, thus preventing cytosolic entry of virus genomes[16]. Though the exact mechanism of fusion inhibition is not known, IFITM3 has been proposed to alter membrane properties including the structural rigidity and local curvature of endosome membranes[18, 22]. Our recent work demonstrated that a palmitoylated amphipathic helix within IFITM3 is required for its inhibition of virus protein-mediated membrane fusion, and may stabilize an intermediate of the membrane fusion process[23, 24]. Viruses such as IAV that enter cells primarily through endocytosis are strongly inhibited by IFITM3 while viruses such as SeV that fuse at the cell surface are poorly inhibited[25, 26]. IFITM3 KO mice and humans with deleterious IFITM3 gene polymorphisms experience severe influenza virus infections, confirming the importance of this protein in antiviral defense *in vivo*[27-30]. Effects of IFITM3 on hMPV have not yet been adequately investigated.

In the present study, we show that IFITM3 is able to inhibit hMPV infection, thus identifying the first known cellular restriction factor for this important respiratory pathogen. We observed partial inhibition of hMPV infection by endocytosis inhibitors and partial restriction by IFITM3, both of which are consistent with the dual entry mechanism proposed for this virus. We demonstrate that IFITM3 can block membrane fusion mediated by the hMPV F protein and, importantly, that manipulating the levels, localization, or activity of IFITM3 in cells can have significant effects on hMPV infection. Further, we found that an additional endosomal protein, Toll-like receptor 7 (TLR7), unexpectedly facilitates hMPV infection of cells. Taken together, these results suggest that

the endosomal entry route may be particularly beneficial for hMPV in cell types that express TLR7 despite eventual restriction of this pathway upon induction of IFITM3.

METHODS

Cell culture, transfections, and transductions.

HEK293T, A549, Vero E6, and MEF cells were grown in DMEM supplemented with 10% Equafetal FBS (Atlas biologicals). LLC-MK2 cells were grown in OPTI-MEM reduced serum media + Glutamax supplement (Thermo Fisher) with 2% FBS (Atlas biologicals). HAP1 WT and IFITM3 KO cells (Horizon Discovery) were grown in IMDM with 10% Equafetal FBS. Macrophage cell lines were grown in 1640 RPMI with 10% Equafetal FBS. All cells were grown at $37^{o}C$ with 5% CO_2 in a humidified incubator. HEK293T cells transduced with human TLR7 or vector control were purchased from InvivoGen. NEDD4 WT and KO MEFs were generated by Dr. Hiroshi Kawabe (Max Planck Institute) and were kindly provided to us by Dr. Matthew Pratt (University of Southern California)[31, 32]. Macrophage cell lines were generated by Dr. Douglas Golenbock (University of Massachusetts) and obtained through the NIH-sponsored BEI Resources. Cells were transfected with indicated plasmids using LipoJet transfection reagent (Signagen Laboratories) following the manufacturer's instructions. For generation of stable cell lines, myc-IFITMs were inserted in the pLenti-puro vector and VSV G pseudotyped viruses were generated as described previously[33].

Treatment with IFN, siRNA, endocytosis inhibitors, and amphotericin B.

Where indicated, cells were treated with human IFN-β (obtained through BEI resources and utilized at a 1:100 concentration) for 24 h to induce IFITM3 expression. Concurrent with IFN-β stimulation, IFITM3 knockdown was achieved using Dharmacon ON-TARGET

Plus Smart Pool human IFITM3-targeting (L-014116) or non-targeting control (D-001810-10-20) with Lipofectamine RNAiMAX transfection reagent (Invitrogen) according to the manufacturer's protocol. For experiments involving inhibitors or drugs, cells were treated with chlorpromazine or genistein at concentrations of 10μg/mL or 200μM, respectively, or with 2.7 μM amphotericin B.

Virus propagation, infection, and flow cytometry.

Influenza A virus A/PR/8/34 (H1N1) was propagated in 10 day old embryonated chicken eggs (Charles River) for 48 h at 37C and titered on MDCK cells. SeV expressing GFP (SeV-GFP) was generated by Dr. Dominique Garcin (University de Geneve) and provided to us by Dr. Mark Peeples (The Research Institute at Nationwide Children's Hospital). This virus was propagated in 10 day old embryonated chicken eggs at 37^{o}C for 40 h and titered on Vero E6 cells. hMPV and hMPV-GFP were generated by a reverse genetics system based on the NL/1/00 strain, a prototypic A1 strain, utilizing a previously described system[34]. The hMPV 02-202 (B1) strain was kindly provided to us by Dr. John Williams (University of Pittsburgh). The hMPV (A1), hMPV-GFP (A1), and hMPV (B1) strains were propagated in Vero E6 cells, concentrated by ultracentrifugation through a 20% sucrose cushion, and titered on LLC-MK2 cells. Cells were infected with influenza PR8 for 6 h or 24 h at a multiplicity of infection of 1 or 2.5. For flow cytometry quantification of infection, fixed and permeabilized influenza-infected cells were stained with anti-H1N1 influenza NP (obtained from BEI resources) followed by anti-mouse Alexa-647 secondary (Life Technologies). Cells were infected with hMPV at an MOI of 0.2 or 2 for 20 hours and stained using an anti-hMPV antibody (Millipore, MAB80138) at a 1:1000 dilution. Secondary staining was done using anti-mouse Alexa-488 antibody (Life Technologies), at a 1:1000 dilution. All

antibodies were diluted in 0.1 % Triton X-100 in PBS. Cells infected with GFP viruses were infected for 20 hours, isolated, fixed, and analyzed directly. Flow cytometry was performed on a FACSCanto II flow cytometer (BD Biosciences), and analyzed using FlowJo software.

Western blotting and antibodies.

For Western blotting, cells were lysed in buffer containing 0.1 mM triethanolamine, 150 mM NaCl, and 1% SDS at pH 7.4 supplemented with EDTA-free Protease Inhibitor Cocktail (Roche) and Benzonase Nuclease (Sigma) for 5 min at RT. Primary antibodies for IFITM3 (Proteintech group), HA (HA.11, Biolegend), GAPDH (Invitrogen), Actin (Abcam), Tubulin (Antibody Direct), MYC (Developmental Studies Hybridoma Bank at the University of Iowa, deposited by Dr. J. Michael Bishop, catalog no. 9E 10), and NEDD4 (Millipore) were diluted at 1:1000 in 0.1% Tween 20 in PBS. Secondary antibodies, anti-mouse IgG HRP (Millipore) or anti-rabbit IgG HRP (Cell Signaling), were diluted 1:10000 in 0.1% Tween 20 in PBS.

Cell-cell fusion assay.

A trypsin-independent hMPV F protein was cloned previously by mutating the RQSR motif of the hMPV NL/1/00 strain and used herein[35]. Cell-cell fusion assays were performed as outlined previously[24, 36]. To promote fusion, cells were pulsed with freshly prepared DMEM containing 25 mM MES at pH 5.0 for 2 min. Cells were then washed with PBS and incubated in standard media for 8 hours at 37°C, after which, they were washed once with PBS, and prepared for measurement of luciferase activity using the Promega Dual-Luciferase Reporter Assay System. All readings were recorder using a Promega GloMax-96 Microplate Luminometer.

RESULTS

Overexpression of IFITM3 inhibits hMPV infection.

To examine effects of IFITM3 on hMPV strain NL/1/00 (A1) infection, we compared susceptibility of HEK293T cells after stable transduction with empty vector or myc-tagged human IFITM3 (Figure 1A). Importantly, HEK293T cells do not express detectable IFITM3 at baseline and are thus a useful model for assessing virus infection upon introduction of IFITM3[37]. Percent hMPV infection of the IFITM3-expressing cells as measured by flow cytometry was decreased by approximately half compared to vector control (Figure 1B). As a positive control, we tested susceptibility of the cells to IAV and saw that, consistent with previous reports, infection was reduced almost completely in IFITM3-expressing cells (Figure 1A)[16, 24]. As a negative control, and as observed previously, we found that SeV was largely insensitive to expression of IFITM3 (Figure 1A)[25, 32]. These results are consistent with the endocytic entry of IAV, plasma membrane entry of SeV, and dual entry of hMPV. To further confirm the role of endocytosis in hMPV infection of HEK293T cells, we examined effects of two broad inhibitors of endocytosis, chlorpromazine and genistein. These inhibitors caused a nearly complete elimination of infection of cells by IAV while blocking hMPV infection by roughly 50%, correlating in magnitude to the extent of inhibition by IFITM3 for both viruses (Figure 1B,C). We additionally found that overexpression of IFITM1 partially inhibited hMPV infection of HEK293T cells, and that IFITM2 modestly inhibited infection, though both proteins strongly inhibited IAV infection (Supplementary Figure 1). These results suggested that IFITMs 1-3 are capable of inhibiting hMPV when overexpressed. However, given that IFITMs 1 and 2 do not compensate for loss of IFITM3 in viral infections *in vivo*[27-30, 38, 39], and given that our

early studies examining IFITM3 KO cells suggested an important role for IFITM3 in restricting hMPV, we focused our subsequent experiments on IFITM3.

Endogenous cellular IFITM3 limits hMPV infection.

We next examined the role of endogenously expressed IFITM3 in hMPV infection using IFITM3 KO HAP1 cells generated using CRISPR/CAS9 technology (Figure 2A). Infection of IFITM3 KO cells with hMPV was significantly increased both with and without interferon treatment as compared to WT cells (Figure 2A, B). Likewise, human A549 cells, a lung epithelial cell line, exhibited a significant increase in susceptibility to hMPV when IFITM3 was knocked down using siRNAs in both mock treated and IFN treated cells (Figure 2C and D). These results demonstrate that endogenous IFITM3 restricts hMPV infection and that other IFN-induced effectors cannot fully compensate for loss of IFITM3.

Amphotericin B reverses protective effects of IFITM3.

Amphotericin B is a membrane destabilizing antifungal drug that has been reported to negate the protective effects of IFITM3 against IAV[22]. We sought to confirm these results and to determine whether amphotericin B could similarly reverse IFITM3 restriction of hMPV. We pretreated IFITM3-expressing stable HEK293T cells or vector control cells with amphotericin B, and subsequently measured infection with IAV or hMPV. We observed potent restriction of IAV by IFITM3 that was completely ablated by amphotericin B (Figure 3A). Similarly to previous experiments, we observed that hMPV infection was decreased by roughly 50% in IFITM3-expressing cells, and that this was also completely reversed when cells were pretreated with amphotericin B (Figure 1B, 3B). These results may suggest that IAV and hMPV are inhibited by IFITM3 via similar mechanisms, and also that amphotericin

B may be clinically detrimental for hMPV-infected patients as has been previously suggested for IAV infections[22].

Mutation of the IFITM3 endocytic trafficking motif enhances hMPV restriction.

IFITM3 possesses a four amino acid YxxΦ endocytosis signal at residues 20-23 (YEML in IFITM3) that mediates its trafficking to endosomes and lysosomes from the plasma membrane[19, 20, 26]. A polymorphism in the human IFITM3 gene that is linked to severe influenza virus infections has been proposed to disrupt this motif by altering RNA splicing[27]. Multiple laboratories have demonstrated that mutation of the key residue, Y20, within this motif results in accumulation of IFITM3 at the plasma membrane and in a diminished ability to inhibit influenza virus when expressed at low levels[19, 20, 26]. When strongly expressed, IFITM3-Y20A can be visualized at both the plasma membrane and at endosomes, possibly owing to passive endocytosis of the protein[20, 26, 40]. Given the dual localization of IFITM3-Y20A at the plasma membrane and at endosomes, and the proposed dual entry mechanism of hMPV, we sought to examine the effect of the IFITM3-Y20A variant on hMPV infection. We thus generated a stable HEK293T cell line expressing myc-tagged IFITM3-Y20A and confirmed that its expression was similar to stable cells expressing WT IFITM3 (Figure 4A, B). Infection of these cells with hMPV again showed that IFITM3 significantly decreased infection, and remarkably, that IFITM3-Y20A expression resulted in a near-complete resistance of the cells to infection (Figure 4C). These results demonstrate that the IFITM3-Y20A mutant, which mimics the localization of a naturally occurring IFITM3 variant[19, 27, 41], has an enhanced ability to inhibit hMPV, correlating with its expanded pattern of localization.

Expression of IFITM3 restricts cell-cell fusion mediated by the hMPV F protein.

IFITM3 prevents formation of fusion pores between IAV and host membranes despite lipid mixing occurring between the virus and cell membranes[16, 17]. Thus, we reasoned that restriction of hMPV by IFITM3 likely also involves inhibition of fusion. Previous work by other groups has demonstrated that the hMPV fusion glycoprotein (F protein) is sufficient for virus membrane fusion and that it promotes cell-cell fusion (syncytia formation) when expressed in cells[11, 12]. Previous studies have also utilized cell-cell fusion assays to examine the ability of IFITMs to block fusion mediated by viral fusogens[18, 24]. Here, we used a characterized cell-cell fusion assay in which cells expressing the hMPV F protein were mixed with target cells expressing or lacking IFITMs[24]. Each population of cells was also transfected with distinct plasmids that produce luciferase only when they are present in the same cell, i.e., only when cell-cell fusion has occurred, such that fusion can be quantified by measurement of luciferase activity (Figure 5A). We first confirmed that hMPV F is capable of inducing fusion between HEK293T cell populations (Figure 5B). Importantly, fusion was only observed when hMPV F was present. Previous work by others has demonstrated that low pH is not required, but can increase fusion mediated by some hMPV F proteins[11, 12]. Indeed, pulsing of our mixed cell population with media at pH 5 resulted in enhanced luciferase production indicative of increased cell-cell fusion (Figure 5B). Thus, we utilized a pH 5 pulse in subsequent experiments to achieve optimal fusion. To examine effects of IFITM3 on hMPV F-mediated fusion, we transfected IFITM3, IFITM3-Y20A, or vector control into target cells. Consistent with past reports in which IFITM3 inhibited cell-cell fusion mediated by numerous viral fusion proteins[18, 24], WT IFITM3 inhibited F-mediated fusion (Figure 5B). The Y20A mutant, consistent with its increased

accumulation at the plasma membrane[19, 20, 26, 41], further decreased fusion (Figure 5B). These data demonstrate that IFITM3 is able to block membrane fusion mediated by the hMPV F protein.

Knockout of NEDD4 reduces hMPV infection.

We previously identified NEDD4 as the primary E3 ubiquitin ligase responsible for ubiquitination and turnover of steady-state IFITM3[32]. Depletion of NEDD4 results in reduced ubiquitination of IFITM3 and increased IFITM3 levels even in the absence of infection or IFN stimulation. As such, NEDD4 KO cells are more resistant to IAV infection dependent on their enhanced IFITM3 levels[32]. Using WT or NEDD4 KO mouse embryonic fibroblasts, we confirmed increased levels of basal IFITM3 in the KO versus WT cells (Figure 6A). Upon infection with hMPV, we observed that NEDD4 KO cells were significantly more resistant to hMPV infection than WT cells (Figure 6B). These results further support a role for IFITM3 as a cellular restriction factor able to inhibit hMPV infection and also identify NEDD4 as a novel target for inhibiting hMPV infections.

Toll-like receptor 7 promotes hMPV infection

It is unclear why hMPV has evolved to utilize an IFITM3-restricted entry pathway through endosomes when it is also capable of fusing at the plasma membrane[14]. Although the acidic pH of endosomes may enhance virus fusion, it is not required (Figure 5B)[11, 12, 14]. Other endosomal factors may also promote infection prior to the induction of IFITM3 by IFNs. For example, in a screen of more than 300 potential antiviral restriction factors, overexpression of Toll-like receptor 7 (TLR7), an endosome-localized TLR that is usally associated with an antiviral response, increased infection with an hMPV A2 strain to 131%

when normalized to infection of control cells[42]. To corroborate this result, we examined

infection susceptibility of HEK293T cells to the hMPV A1 strain used in our IFITM3 studies

when stably expressing TLR7 or vector control. Cells expressing TLR7 were infected at a

significantly higher rate, independently confirming the published large-scale screen (Figure

7A,B). To determine whether endogenous TLR7 affects hMPV infection, we measured the

infection susceptibility of WT and TLR7 KO murine macrophage cell lines. Consistent with

our overexpression results, the percent infection of TLR7 KO cells was significantly

decreased as compared to WT cells (Figure 7C). Infections of cells lacking TLR3, another

endosomal protein, or TLR4, which localizes to the cell surface, were not significantly

different from WT, providing additional specificity controls (Figure 7C). We next examined

whether hMPV strain 02-202 (B1) was affected by IFITM3 and TLR7 expression in

HEK293T cells. We found that, like the A1 strain, the B1 strain was susceptible to IFITM3

restriction (Supplementary Figure 2). However, we observed no effect of TLR7 on the B1

strain, distinguishing it from the A1 strain used in our studies and the A2 strain used

previously[42] (Figure 7, Supplementary Figure 2).   These results demonstrate that TLR7

facilitates infection by certain strains of hMPV, and along with results on IFITM3, highlight

the complex interactions between hMPV and cellular endosomes.

DISCUSSION

Our work has identified IFITM3 as the first confirmed restriction factor able to limit hMPV

infection of cells. Infection by hMPV was decreased by overexpression of IFITM3 (Figure

1A,B) and was increased by knockout or knockdown of endogenous IFITM3 (Figure 2).

Cellular pathways that regulate IFITM3 levels may thus represent new targets for limiting

hMPV infections. Indeed, NEDD4 KO cells, which accumulate high levels of IFITM3 due to

inefficient IFITM3 ubiquitination at steady state[32] were more resistant to hMPV infection than WT cells (Figure 6). Our data indicate that IFITM3 is able to restrict hMPV of the A1 and B1 lineages (Figure 1 and 2, and Supplementary Figure 2), and the previously mentioned large-scale overexpression screen of potential antiviral factors also showed partial inhibition of an A2 virus by IFITM3[42]. These findings indicate that IFITM3 broadly restricts multiple hMPV lineages.

Past work studying IAV and its inhibition by IFITM3 showed that the drug amphotericin B, which destabilizes lipid membranes, neutralizes the antiviral effects of IFITM3[22]. Our examination of hMPV infection of cells expressing IFITM3 showed that amphotericin B similarly negates the activity of IFITM3 against this additional virus (Figure 3B). Importantly, amphotericin B is a commonly used antifungal agent, and our new work suggests that its clinical use may enhance susceptibility to not only IAV, but also hMPV. This work also suggests that IFITM3 inhibits hMPV infection via effects on membranes. Using cell-cell fusion assays, we found that indeed, IFITM3 is able to inhibit membrane fusion mediated by the hMPV F protein (Figure 5). This work is in line with the previously established ability of IFITM3 to block fusion pore formation in IAV infection and cell-cell fusion mediated by a multitude of viral fusion proteins[16-18, 24].

TLR7 is generally considered an antiviral molecule since it detects viral single-stranded RNA and signals for the production of type I IFNs[43]. TLR7 is expressed on macrophages, conventional dendritic cells, and plasmacytoid dendritic cells[43], and our work suggests that its presence makes these cell types more susceptible to infection with certain hMPV strains. Staining of primary human lung airway epithelial cells with anti-TLR7 antibodies also indicated that TLR7 is expressed on these cells[44], which are

additional relevant targets of hMPV infection. How hMPV utilizes TLR7 to facilitate cellular infection is not yet mechanistically understood, though the increase in the percentage of cells infected when TLR7 is expressed may suggest that TLR7 is a cell entry factor (Figure 7). Remarkably, the ability of hMPV to coopt TLR7 is complemented by the reported inhibition of TLR7 signaling by the hMPV M2-2 protein[45]. Thus, hMPV has evolved to utilize TLR7 while also inhibiting its antiviral function. Detection of hMPV infection by retinoic acid inducible gene I (RIG-I), a cytosolic sensor of viral replication products, is similarly inhibited by the P or G proteins of some hMPV strains[46, 47]. Likewise, IFN signaling activity is also decreased in hMPV-infected cells via inhibition of Signal Transducer and Activator of Transcription (STAT) protein phosphorylation[48, 49]. Thus, hMPV utilizes several mechanisms to evade the type I IFN response while benefitting from the presence of TLR7.

Both IFITM3 and TLR7 are primarily endosomal proteins, though they have opposite effects on hMPV infection. In most cell types, IFITM3 levels are low prior to infection and its induction by IFNs[32]. Consistent with a role for IFN-induced proteins, such as IFITM3, in controlling hMPV infection *in vivo*, type I IFN receptor KO mice show increased hMPV titers in the lungs at early timepoints post infection[50]. It will be interesting to examine pathogenesis of hMPV infection in IFITM3 and TLR7 KO mice in the future, and to determine whether TLR7-blocking antibodies or nucleic acids can impact hMPV infections. Likewise, our work suggests that polymorphisms in the human IFITM3 gene that have been linked to severe IAV infections should also be examined for effects on hMPV infections[27-30]. Overall, we have revealed that two endosomal proteins, IFITM3 and TLR7, differentially regulate vulnerability of cells to hMPV infection. These findings

support the proposed role for endocytosis in hMPV infection, and have identified new factors that may be manipulated or targeted to prevent or treat infection by this important human pathogen.

Notes

Acknowledgments. We thank Dr. John Williams for providing us with the hMPV B1 strain used in this work.

Author Contributions. T.M.M., J.L., and J.S.Y. designed experiments and analyzed data. T.M.M., Y.Z., A.D.K., L.Z., M.L., and M.C. performed experiments. T.M.M. and J.S.Y. wrote the manuscript with editorial contributions from all authors.

Financial Support. This research was supported by NIH grant AI130110 (to J.S.Y.). T.M.M. was supported by a Gilliam Fellowship for Advanced Study from the Howard Hughes Medical Institute. A.D.K. was supported by The Ohio State University Systems and Integrative Biology Training Program funded by NIH grant GM068412.

Conflicts of Interest. There are no conflicts of interest regarding the publication of this manuscript. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Rima B, Collins P, Easton A, et al. ICTV Virus Taxonomy Profile: Pneumoviridae. J Gen Virol **2017**; 98:2912-3.

2. van den Hoogen BG, de Jong JC, Groen J, et al. A newly discovered human pneumovirus isolated from young children with respiratory tract disease. Nat Med **2001**; 7:719-24.

3. Williams JV, Harris PA, Tollefson SJ, et al. Human metapneumovirus and lower respiratory tract disease in otherwise healthy infants and children. N Engl J Med **2004**; 350:443-50.

4. Widmer K, Zhu Y, Williams JV, Griffin MR, Edwards KM, Talbot HK. Rates of hospitalizations for respiratory syncytial virus, human metapneumovirus, and influenza virus in older adults. J Infect Dis **2012**; 206:56-62.

5. Edwards KM, Zhu Y, Griffin MR, et al. Burden of human metapneumovirus infection in young children. N Engl J Med **2013**; 368:633-43.

6. Yan XL, Li YN, Tang YJ, et al. Clinical characteristics and viral load of respiratory syncytial virus and human metapneumovirus in children hospitaled for acute lower respiratory tract infection. J Med Virol **2017**; 89:589-97.

7. Sastre P, Ruiz T, Schildgen O, Schildgen V, Vela C, Rueda P. Seroprevalence of human respiratory syncytial virus and human metapneumovirus in healthy population analyzed by recombinant fusion protein-based enzyme linked immunosorbent assay. Virol J **2012**; 9:130.

8. Sun J, Wei Y, Rauf A, et al. Methyltransferase-defective avian metapneumovirus vaccines provide complete protection against challenge with the homologous Colorado strain and the heterologous Minnesota strain. J Virol **2014**; 88:12348-63.

9. Zhang Y, Wei Y, Zhang X, Cai H, Niewiesk S, Li J. Rational design of human metapneumovirus live attenuated vaccine candidates by inhibiting viral mRNA cap methyltransferase. J Virol **2014**; 88:11411-29.

10. Hoekstra D, Klappe K, de Boer T, Wilschut J. Characterization of the fusogenic properties of Sendai virus: kinetics of fusion with erythrocyte membranes. Biochemistry **1985**; 24:4739-45.

11. Schowalter RM, Chang A, Robach JG, Buchholz UJ, Dutch RE. Low-pH Triggering of Human Metapneumovirus Fusion: Essential Residues and Importance in Entry. Journal of Virology **2009**; 83:1511-22.

12. Herfst S, Mas V, Ver LS, et al. Low-pH-induced membrane fusion mediated by human metapneumovirus F protein is a rare, strain-dependent phenomenon. Journal of Virology **2008**; 82:8891-5.

13. Yang H, He H, Tan B, Liu EM, Zhao XD, Zhao Y. Human metapneumovirus uses endocytosis pathway for host cell entry. Mol Cell Probe **2016**; 30:231-7.

14. Cox RG, Mainou BA, Johnson M, et al. Human Metapneumovirus Is Capable of Entering Cells by Fusion with Endosomal Membranes. Plos Pathogens **2015**; 11.

15. Edinger TO, Pohl MO, Stertz S. Entry of influenza A virus: host factors and antiviral targets. Journal of General Virology **2014**; 95:263-77.

16. Feeley EM, Sims JS, John SP, et al. IFITM3 inhibits influenza A virus infection by preventing cytosolic entry. PLoS Pathog **2011**; 7:e1002337.

17. Desai TM, Marin M, Chin CR, Savidis G, Brass AL, Melikyan GB. IFITM3 restricts influenza A virus entry by blocking the formation of fusion pores following virus-endosome hemifusion. PLoS Pathog **2014**; 10:e1004048.

18. Li K, Markosyan RM, Zheng YM, et al. IFITM proteins restrict viral membrane hemifusion. PLoS Pathog **2013**; 9:e1003124.

19. Jia R, Xu F, Qian J, et al. Identification of an endocytic signal essential for the antiviral action of IFITM3. Cell Microbiol **2014**; 16:1080-93.

20. Chesarino NM, McMichael TM, Hach JC, Yount JS. Phosphorylation of the antiviral protein interferon-inducible transmembrane protein 3 (IFITM3) dually regulates its endocytosis and ubiquitination. J Biol Chem **2014**; 289:11986-92.

21. Chesarino NM, McMichael TM, Yount JS. Regulation of the trafficking and antiviral activity of IFITM3 by post-translational modifications. Future Microbiol **2014**; 9:1151-63.

22. Lin TY, Chin CR, Everitt AR, et al. Amphotericin B increases influenza A virus infection by preventing IFITM3-mediated restriction. Cell Rep **2013**; 5:895-908.

23. Yount JS, Moltedo B, Yang YY, et al. Palmitoylome profiling reveals S-palmitoylation-dependent antiviral activity of IFITM3. Nat Chem Biol **2010**; 6:610-4.

24. Chesarino NM, Compton AA, McMichael TM, et al. IFITM3 requires an amphipathic helix for antiviral activity. EMBO Rep **2017**; 18:1740-51.

25. Hach JC, McMichael T, Chesarino NM, Yount JS. Palmitoylation on conserved and nonconserved cysteines of murine IFITM1 regulates its stability and anti-influenza A virus activity. J Virol **2013**; 87:9923-7.

26. Compton AA, Roy N, Porrot F, et al. Natural mutations in IFITM3 modulate post-translational regulation and toggle antiviral specificity. EMBO Rep **2016**; 17:1657-71.

27. Everitt AR, Clare S, Pertel T, et al. IFITM3 restricts the morbidity and mortality associated with influenza. Nature **2012**; 484:519-23.

28. Wang Z, Zhang A, Wan Y, et al. Early hypercytokinemia is associated with interferon-induced transmembrane protein-3 dysfunction and predictive of fatal H7N9 infection. Proc Natl Acad Sci U S A **2014**; 111:769-74.

29. Pan Y, Yang P, Dong T, et al. IFITM3 Rs12252-C Variant Increases Potential Risk for Severe Influenza Virus Infection in Chinese Population. Front Cell Infect Microbiol **2017**; 7:294.

30. Allen EK, Randolph AG, Bhangale T, et al. SNP-mediated disruption of CTCF binding at the IFITM3 promoter is associated with risk of severe influenza in humans. Nat Med **2017**; 23:975-83.

31. Fouladkou F, Landry T, Kawabe H, et al. The ubiquitin ligase Nedd4-1 is dispensable for the regulation of PTEN stability and localization. P Natl Acad Sci USA **2008**; 105:8585-90.

32. Chesarino NM, McMichael TM, Yount JS. E3 Ubiquitin Ligase NEDD4 Promotes Influenza Virus Infection by Decreasing Levels of the Antiviral Protein IFITM3. PLoS Pathog **2015**; 11:e1005095.

33. Wang F, St Gelais C, de Silva S, et al. Phosphorylation of mouse SAMHD1 regulates its restriction of human immunodeficiency virus type 1 infection, but not murine leukemia virus infection. Virology **2016**; 487:273-84.

34. Zhang Y, Sun J, Wei Y, Li J. A Reverse Genetics Approach for the Design of Methyltransferase-Defective Live Attenuated Avian Metapneumovirus Vaccines. Methods Mol Biol **2016**; 1404:103-21.

35. Zhang Y, Wei Y, Li J, Li J. Development and optimization of a direct plaque assay for human and avian metapneumoviruses. J Virol Methods **2012**; 185:61-8.

36. Battles MB, Langedijk JP, Furmanova-Hollenstein P, et al. Molecular mechanism of respiratory syncytial virus fusion inhibitors. Nat Chem Biol **2016**; 12:87-93.

37. Melvin WJ, McMichael TM, Chesarino NM, Hach JC, Yount JS. IFITMs from Mycobacteria Confer Resistance to Influenza Virus When Expressed in Human Cells. Viruses **2015**; 7:3035-52.

38. Gorman MJ, Poddar S, Farzan M, Diamond MS. The Interferon-Stimulated Gene Ifitm3 Restricts West Nile Virus Infection and Pathogenesis. J Virol **2016**; 90:8212-25.

39. Poddar S, Hyde JL, Gorman MJ, Farzan M, Diamond MS. The Interferon-Stimulated Gene IFITM3 Restricts Infection and Pathogenesis of Arthritogenic and Encephalitic Alphaviruses. J Virol **2016**; 90:8780-94.

40. Peng T, Hang HC. Site-Specific Bioorthogonal Labeling for Fluorescence Imaging of Intracellular Proteins in Living Cells. J Am Chem Soc **2016**; 138:14423-33.

41. Jia R, Pan Q, Ding S, et al. The N-terminal region of IFITM3 modulates its antiviral activity by regulating IFITM3 cellular localization. J Virol **2012**; 86:13697-707.

42. Schoggins JW, MacDuff DA, Imanaka N, et al. Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. Nature **2014**; 505:691-5.

43. Petes C, Odoardi N, Gee K. The Toll for Trafficking: Toll-Like Receptor 7 Delivery to the Endosome. Front Immunol **2017**; 8:1075.

44. Ioannidis I, Ye F, McNally B, Willette M, Flano E. Toll-like receptor expression and induction of type I and type III interferons in primary airway epithelial cells. J Virol **2013**; 87:3261-70.

45. Kitagawa Y, Sakai M, Funayama M, Itoh M, Gotoh B. Human Metapneumovirus M2-2 Protein Acts as a Negative Regulator of Alpha Interferon Production by Plasmacytoid Dendritic Cells. Journal of Virology **2017**; 91.

46. Goutagny N, Jiang ZZ, Tian J, et al. Cell Type-Specific Recognition of Human Metapneumoviruses (HMPVs) by Retinoic Acid-Inducible Gene I (RIG-I) and TLR7 and Viral Interference of RIG-I Ligand Recognition by HMPV-B1 Phosphoprotein. Journal of Immunology **2010**; 184:1168-79.

47. Bao XY, Liu TS, Shan YC, Li K, Garofalo RP, Casola A. Human metapneumovirus glycoprotein G inhibits innate immune responses. Plos Pathogens **2008**; 4.

48. Hastings AK, Amato KR, Wen SC, Peterson LS, Williams JV. Human metapneumovirus small hydrophobic (SH) protein downregulates type I IFN pathway signaling by affecting STAT1 expression and phosphorylation. Virology **2016**; 494:248-56.

49. Ren JP, Kolli D, Liu TS, et al. Human Metapneumovirus Inhibits IFN-beta Signaling by Downregulating Jak1 and Tyk2 Cellular Levels. Plos One **2011**; 6.

50. Hastings AK, Erickson JJ, Schuster JE, et al. Role of type I interferon signaling in human metapneumovirus pathogenesis and control of viral replication. J Virol **2015**; 89:4405-20.

Figure 1: Endocytosis inhibitors and overexpressed IFITM3 restrict hMPV infection. A) Anti-myc and anti-Actin Western blotting of HEK293T cells stably transduced with vector control or myc-IFITM3. **B)** HEK293T cells stably transduced with vector control or myc-IFITM3 (IFITM3) were infected with IAV, hMPV, or SeV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three or more experiments with error bars representing standard deviation. **C)** HEK293T cells were treated for 1 h with chlorpromazine (Chlor) or genistein (Gen), or were mock treated, followed by infection with IAV or hMPV in the presence or absence of the inhibitors as indicated for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are from an experiment representative of three experiments with error bars representing standard deviation of triplicate samples. **B,C)** Asterisks indicate $p < 0.001$ and NS indicates not significant by Student's t-test.

Figure 2: Endogenous IFITM3 restricts hMPV infection. A,B) WT and IFITM3 KO HAP1 cells were mock treated or treated with IFN-β overnight. **A)** Anti-IFITM3 and anti-GAPDH Western blotting. **B)** Cells were infected with hMPV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three or more experiments with error bars representing standard deviation. **C,D)** A549 cells were mock treated or treated with IFN-β overnight. Cells were simultaneously transfected with siRNA targeting IFITM3 (siIFITM3) or non-targeting control siRNA (siControl) as indicated. **C)** Anti-IFITM3 and anti-GAPDH Western blotting. **D)** Cells were infected with hMPV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of triplicate samples from a representative experiment from more than 3 similar experiments with error bars representing standard deviation. **B,D)** Asterisks indicate $p<0.001$ by Student's t-test.

Figure 3: Amphotericin B reverses IFITM3 inhibition of hMPV infection. A,B) HEK293T cells stably transduced with vector control or myc-IFITM3 (IFITM3) were treated for 1 h with amphotericin B or were mock treated, followed by infection with IAV **(A)** or hMPV **(B)** in the presence or absence of amphotericin B as indicated for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three experiments with error bars representing standard deviation. Asterisks indicate p<0.001 as compared to all other samples on the respective graphs by Student's t-test.

Figure 4: Mutation of the IFITM3 endocytosis motif enhances restriction of hMPV infection. A) Anti-myc and anti-GAPDH Western blotting of HEK293T cells stably transduced with vector control, myc-IFITM3, or myc-IFITM3-Y20A. **B)** Cells as in **(A)** were analyzed by flow cytometry after staining for IFITM3. **C)** Cells as in **(A,B)** were infected with hMPV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three or more experiments with error bars representing standard deviation. Asterisk indicates p<0.001 and double asterisks represent p<0.0001 by Student's t-test.

Figure 5: IFITM3 can inhibit membrane fusion mediated by the hMPV F Protein. A) Scheme for cell-cell fusion assay. **B)** HEK293T cells were transfected with pFR-Luc plus plasmid expressing hMPV F protein. A second set of cells were transfected with pBD-NFκB. Mixed cell populations were treated with media at pH 5 or with standard media for 2 min. After 8 h, cell lysates were analyzed for luciferase activity indicative of cell-cell fusion. Bar graphs depict averages of triplicate samples representative of three experiments with error bars representing standard deviation. C) HEK293T cells were transfected with pFR-Luc plus plasmid expressing hMPV F protein. A second set of cells were transfected with pBD-NFκB plus plasmid expressing myc-IFITM3 (IFITM3), myc-IFITM3-Y20A (IFITM3-Y20A) or vector control. Mixed cell populations were treated with media at pH 5 for 2 min. After 8 h, cell lysates were analyzed for luciferase activity indicative of cell-cell fusion. Bar graphs depict averages of triplicate samples representative of three or more experiments with error bars representing standard deviation. **B,C)** Asterisks indicate $p< 0.001$ and double asterisks indicate $p<0.0001$ by Student's t-test.

Figure 6: NEDD4 KO cells are partially resistant to hMPV infection. A) Anti-IFITM3, anti-Tubulin, and anti-NEDD4 Western blotting for WT or NEDD4 KO MEFs. **B)** Cells as in **(A)** were infected with hMPV at two MOIs for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are an average of triplicate samples from an experiment representative of four similar experiments. Error bars represent standard deviation. Asterisks indicate p<0.0001 by Student's t-test.

Figure 7: TLR7 enhances hMPV infection. A) Anti-HA and anti-GAPDH Western blotting of HEK293T cells stably transduced with vector control or TLR7-HA. **B)** HEK293T cells stably transduced with vector control or TLR7-HA were infected with hMPV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three experiments with error bars representing standard deviation. Asterisk indicates $p<0.001$ by Student's t-test. **C)** WT, TLR3 KO, TLR4 KO, or TLR7 KO murine macrophages were infected with hMPV for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of three experiments with error bars representing standard deviation. Asterisk indicates $p<0.001$ compared to all other samples in the graph by Student's t-test.

Supplementary Figure 1: Overexpressed IFITMs restrict hMPV infection. A)
Anti-myc and anti-GAPDH Western blotting of HEK293T cells stably transduced
with vector control, myc-IFITM1, myc-IFITM2, or myc-IFITM3. **B,C)** HEK293T cells
stably transduced with vector control or myc-IFITMs were infected with IAV **(B)** or
hMPV **(C)** for 24 h. Percent infection was determined by flow cytometry after
staining with virus-specific antibodies using non-infected cells to set gates. Results
shown are averages of two or more experiments with error bars representing
standard deviation. Asterisks indicate $p<0.001$ as compared to vector control by
Student's t-test. Some data from Main Text Figure 1 is re-plotted here for
comparison between IFITM3 and other IFITMs.

Supplementary Figure 2: Strain-specific effects of IFITM3 and TLR7 on hMPV infection. A) HEK293T cells stably transduced with vector control or myc-IFITM3 (IFITM3) were infected with hMPV A1 or B1 strains for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of triplicate samples from an experiment representative of more than three experiments with error bars representing standard deviation. **B)** HEK293T cells stably transduced with vector control TLR7-HA (TLR7) were infected with hMPV A1 or B1 strains for 24 h. Percent infection was determined by flow cytometry after staining with virus-specific antibodies using non-infected cells to set gates. Results shown are averages of triplicate samples from an experiment representative of more than three experiments with error bars representing standard deviation. **A,B)** Asterisks represent p<0.001 and NS indicates not significant by Student's t-test.