Intracellular signaling by the *comRS* system in *Streptococcus mutans* genetic competence

Simon A.M. Underhill[1], Robert C. Shields[2], Justin R. Kaspar[2], Momin Haider[1], Robert A. Burne[2] and Stephen J. Hagen[*1]

[1]Department of Physics, University of Florida, 2001 Museum Road, Gainesville, Florida, USA 32611

[2] Department of Oral Biology, University of Florida, 1395 Center Drive, Gainesville, Florida, USA 32610

*Corresponding author

Tel. 352 392 4716

 Email: sjhagen@ufl.edu.

Abstract

Entry into genetic competence in streptococci is controlled by ComX, an alternative sigma factor for genes that enable the import of exogenous DNA. In *Streptococcus mutans* the immediate activator of *comX* is the ComRS quorum system. ComS is the precursor of XIP, a seven-residue peptide that is imported into the cell and interacts with the cytosolic receptor ComR to form a transcriptional activator for both *comX* and *comS*. Although intercellular quorum signaling by ComRS has been demonstrated, observations of bimodal expression of *comX* suggest that *comRS* may also function as an intracellular feedback loop, activating *comX* without export or detection of extracellular XIP. Here we use microfluidic and single cell methods to test whether ComRS induction of *comX* requires extracellular XIP or ComS. We find that individual *comS*-overexpressing cells activate their own *comX,* independent of the rate at which their growth medium is replaced. However, in the absence of lysis they do not activate *comS*-deficient mutants growing in co-culture. We also find that induction of *comR* and *comS* genes introduced into *E. coli* cells leads to activation of a *comX* reporter. Therefore ComRS control of *comX* does not require either the import or extracellular accumulation of ComS or XIP, or specific processing of ComS to XIP. We also find that endogenously and exogenously produced ComS/XIP have inequivalent effects on *comX* activation. These data are fully consistent with intracellular positive feedback in *comS* transcription as the origin of bimodal *comX* expression in *S. mutans*.

Importance

The ComRS system can function as a quorum sensing trigger for genetic competence in *S. mutans.* The signal peptide XIP, which is derived from the precursor ComS, enters the cell and interacts with the Rgg-type cytosolic receptor ComR to activate *comX,* which encodes the alternative sigma factor for the late competence genes. Previous studies have demonstrated intercellular signaling via ComRS, although release of the ComS or XIP peptide to the extracellular medium appears to require lysis of the producing cells. Here we test the complementary hypothesis that ComRS can drive *comX* through a purely intracellular mechanism that does not depend on extracellular accumulation or import of ComS or XIP. By combining single-cell, coculture and microfluidic approaches we demonstrate that endogenously produced ComS can allow ComRS to activate *comX* without requiring processing, export or import. These data provide insight into intracellular mechanisms that generate noise and heterogeneity in *S.mutans* competence.

Introduction

Streptococcus mutans inhabits human oral biofilms and is a primary etiological

agent of dental caries (1). Many of the behaviors that facilitate the growth, competition,

stress tolerance, and virulence of *S. mutans* are linked to the regulation of genetic

competence, a transient physiological state during which the organism can internalize

DNA from its environment (2-7). The competence pathway of *S. mutans* is complex, as

it receives input from extracellular peptide signals and environmental cues as well as

regulatory feedback (8-14). Consequently, several elements of the mechanism and

dynamics of the pathway are not well understood.

S. mutans initiates entry into the competent state by increasing the transcription

of the *comX* gene (sometimes referred to as *sigX*), which encodes an alternative sigma

factor that is required for the expression of approximately 30 late competence genes

(15,16). Expression of *comX* can be induced by the peptides CSP (competence

stimulating peptide) and XIP (*sigX*-inducing peptide), and the efficacy of these peptides

is strongly influenced by environmental conditions. CSP is derived by cleavage of a 21-

residue peptide from ComC and exported through an ATP-binding cassette transporter.

It is further processed to the active 18-residue peptide by the SepM protease (17).

Extracellular CSP is detected by the two-component signal transduction system

ComDE, with the phosphorylated response regulator ComE activating genes for

bacteriocin synthesis and immunity. ComE does not directly activate *comX*, but affects

comX indirectly, via a pathway that is not completely understood (18).

The immediate regulator of *comX* in *S. mutans* and in streptococci of the

salivarius, bovis and pyogenic groups is the ComRS system. ComR is an Rgg-like

cytosolic transcriptional regulator, and the type II ComS of *S. mutans* is a 17-residue

peptide (19). The C-terminus of ComS contains the 7-residue small hydrophobic peptide

XIP. Extracellular XIP is imported by the Opp permease and interacts with ComR to

form a transcriptional activator for both *comX* and *comS* (19, 20). Notably, the *S.*

mutans competence pathway contains at least two positive feedback loops, as

XIP/ComR activates *comS* and ComX activates *comE* expression (19, 20).

An intriguing property of *S. mutans* competence is that although exogenous CSP

and XIP can both activate *comX* and induce transformability, they do so under different

environmental conditions and they elicit qualitatively different behaviors in the

expression of *comX* (11). Exogenous CSP elicits a bimodal response in which less than

half of the population activates *comX*, whereas exogenous XIP elicits a unimodal

response in which all cells in the population activate *comX*. Further, CSP activates

comX only in complex media containing small peptides; this activation requires that cells

carry an intact *comS* but it does not require the Opp permease (11). By contrast,

exogenous XIP activates *comX* only in defined media lacking small peptides (11, 21);

this activation requires that cells carry the *opp* gene but does not require *comS* (19).

Therefore, although exogenous XIP can activate *comX* in a strain lacking *comS*, the

bimodal *comX* response to CSP requires an intact *comS* gene.

The observation that competence in several streptococcal species is directly

stimulated by an extracellular ComS-derived peptide suggests that ComRS constitutes

a novel type of Gram-positive quorum signaling system, in which the ComS-derived

signal XIP is processed and secreted, accumulates in the extracellular medium, and is

then reimported. This interpretation in *S. mutans* is supported by several experimental

observations. First, cells that carry *opp* take up exogenous XIP (in defined medium) and

activate *comX* with high efficiency (19, 21). Second, exogenous synthetic XIP is

dramatically more effective in stimulating transformability than is exogenous full-length

ComS (19). Third, filtrates of *S. mutans* cultures grown to $OD_{550} = 0.4$ in defined

medium were able to stimulate a P*comX* reporter strain (21). Moreover, LC-MS/MS

analysis of supernatants of *S. mutans* cultures grown to high density (22, 23) detected

µM quantities of XIP. Fourth, a transposon mutagenesis screen in *S. pyogenes*

identified the widely conserved *pptAB* ABC transporter as a possible exporter of short

hydrophobic peptides of the ComS type (24), raising the possibility that *S. mutans* may

also possess dedicated mechanisms for processing and export of ComS/XIP.

However such mechanisms have not yet been identified in *S. mutans*. There is

evidence that the Eep membrane protease facilitates the processing of *S. thermophilus*

ComS (25), but Eep did not affect processing of ComS of *S. mutans* (22). Further,

although *S. mutans* appears to encode a gene product with a fairly high degree of

homology to PptAB of *S. pyogenes*, deletion of the apparent *pptAB* genes had only a

weak effect on competence induction in mid-exponential phase cultures of *S. mutans*

(24). Consequently, the processing of ComS to XIP remains uncharacterized. The

import of XIP presents an additional puzzle for ComRS quorum signaling because the

permease Opp is required for XIP to activate *comX,* but is not required for activation by

CSP.

We recently demonstrated that XIP can function as a diffusible, intercellular

signal (26) in *S. mutans*. A ComS producing strain was able to induce *comX* response

in co-cultured cells lacking *comS*, in the absence of physical contact. However

intercellular signaling was greatly impaired by deletion of the gene *atlA* encoding the

primary autolysin. Therefore, lysis and not active transport appears to be the primary

route by which *S. mutans* externalizes the diffusible signal.

Here we use a combination of microfluidic and coculture methods to test the

complementary hypothesis that ComRS can also activate *comX* by an intracellular

mechanism, through endogenous ComS production, and that this mechanism does not

specifically require processing of ComS/XIP or its extracellular accumulation.

Results

An intact copy of *comS* alters the *comX* response to exogenous XIP

A previous study found that even when a *comS* deletion (Δ*comS*) and wild type

background were supplied with high concentrations of synthetic XIP, the transformation

efficiency of the Δ*comS* strain was only half as great as that of the wild type (19).

Fig. 1A-1B shows a similar experiment performed with individual cells carrying a

P*comX-gfp* plasmid-borne reporter. *S. mutans* UA159 (wild type) and Δ*comS* genetic

backgrounds were imaged while adhered within a microfluidic chamber and supplied

with a constant flow of defined medium (FMC) containing synthetic XIP. Although both

strains respond to exogenous XIP, the Δ*comS* strain consistently showed roughly 1.5-

fold lower P*comX* activity than the wild type, even at saturating XIP concentrations. The

threshold for *comX* response occurred at a roughly 2-fold lower XIP concentration in

UA159 than in the Δ*comS* strain (Fig. 1B). Therefore, the deletion of *comS* elevated the

threshold for a response to extracellular XIP and reduced the overall response at

saturation.

The deletion of *comS* also affected cell-to-cell variability (noise) in *comX*

expression*.* Figs. 1C and 1D show that the histograms of reporter fluorescence differ in

UA159 (wild type) and Δ*comS* cells. UA159 showed a generally broader (noisier) *comX*

response than did Δ*comS.* We quantified this difference by fitting the histograms to a

gamma distribution $\Gamma(n \mid a,b)$, a two-parameter probability distribution that can be used

to model cell-to-cell variability in *n,* the copy number for a bacterial protein (27). In a

simple physical model, the parameter *a* of the gamma distribution is related to the

number of mRNAs produced during the cell division time, while *b* is related to the

number of protein copies produced per mRNA transcript (28). As shown in Figs. 1E and

1F, the UA159 background has a roughly 2-fold higher value for parameter *a*

(transcription rate), while parameter *b* (translation) is similar for the two strains. As this

difference persists even at XIP concentrations exceeding 1 µM, these data show that

deletion of *comS* significantly affects *comX* expression, even when excess extracellular

XIP is provided.

Fluid replacement did not alter induction of *comX* in a *comS* overexpressing

strain

To test whether *comX* activation requires cells to import extracellular XIP from

their environment, we examined the effect of fluid flow rate in cells that overexpressed

comS from the strong P23 promoter on the plasmid pIB184 (26, 29) while adhered in a

microfluidic flow chamber. Cells carrying the 184*comS* overexpression plasmid produce

significantly higher levels of *comS* mRNA (Fig. S1) and activate *comX* in defined

medium lacking exogenous CSP or XIP (26). We anticipated that if cells were

immobilized and supplied with a continuous flow of fresh medium, high flow rates would

remove XIP (or ComS) that was released, leading to diminished *comX* activity. In order

to prevent XIP import via Opp, cells were supplied with a flow of complex medium (BHI)

lacking XIP or CSP. We loaded 184*comS* P*comX-rfp* cells, which carry both the *comS*

overexpression plasmid and a plasmid-borne P*comX-rfp* reporter, into four different

microfluidic flow chambers. Chambers were supplied with fresh complex medium

flowing at rates between 0.02 ml h^{-1} and 1 ml h^{-1}. These flow rates were sufficient to

completely replace the growth medium within each chamber on time intervals ranging

from 6 seconds to 10 minutes. We also studied (I) a P*comX-rfp* reporter in a UA159

background (negative control) and (II) a ComS-overproducing strain lacking a start

codon (ATG point-mutated to AAG) on its chromosomal *comS* (184*comS* P*comX-rfp*

Δ*comS*).

Figs. 2A shows that P*comX* was not activated in the UA159 background (leftmost

column). In contrast, the strain overexpressing *comS* and harboring an intact

chromosomal copy of *comS* showed a highly heterogeneous response, indicating that a

subpopulation of these cells strongly activated *comX* in the flowing complex medium.

Further, the rate of fluid flow had no effect on their *comX* expression. Fig. 2B shows the

fluorescence of the individual cells for which the signal exceeded the maximum P*comX*

activity (roughly 100 fluorescence units) seen in the UA159 negative control. Rather

than declining at high flow rates where the medium was rapidly replaced, the median

comX activity in the ComS-overproducing cells actually showed a very slight increase,

smaller than the cell-to-cell variability. These data show that overexpression of *comS*

can allow *S. mutans* to activate *comX*, even in complex medium which normally does

not permit activation of *comX* transcription by exogenously supplied XIP. The finding

that this activation is unaffected by rapid replacement of the medium implies that the

comX response in the *comS* overexpressing strain was not due to accumulation of

extracellular XIP (or ComS). Curiously however, this *comX* response did require an

intact chromosomal *comS*: Figs. 2A and 2B show that very few Δ*comS* cells activated

comX, even though they harbored the *comS* overexpression plasmid. Together with

Fig. 1, these data show that the chromosomal *comS* plays a role in *comX* activation that

is not fully complemented either by saturating concentrations of exogenous XIP or by

endogenous overproduction of ComS.

The presence of *comS* in the chromosome affects *comS* mRNA levels

We used RT-qPCR to measure *comS*, *comR* and *comX* transcript copy numbers

in mid-exponential phase cultures of the ComS overexpression, *comS* deletion, and

UA159 strains (Fig. S1). Transcription of *comS* (Fig. S1A) was significantly increased in

cells treated with exogenously added XIP in defined medium, compared to controls

lacking XIP or growing in complex medium. In addition, cells overexpressing *comS* in a

wild-type genetic background (184*comS*) and growing in complex medium had *comS*

transcript levels that were significantly higher than a *comS* overexpressing strain that

lacked the chromosomal copy of *comS* (184*comS* Δ*comS*). Levels of *comR* transcripts

showed no change across strains or conditions (Fig. S1C).

Population density of *comS* overexpressing cells does not determine the *comX*

response

To further test whether *comS*-overexpressing cells release extracellular XIP, we

measured the effect of *comS*-overexpressing (sender) populations on *comX* activation

in *comS*-deficient (receiver) cells growing in coculture. We mixed sender (184*comS*

P*comX-rfp*) and receiver (P*comX-gfp* Δ*comS*) cultures in different ratios and loaded

them into microfluidic chambers containing static medium without exogenous XIP. We

anticipated that, if senders released XIP or ComS into the extracellular medium, both

senders (RFP reporter) and receivers (GFP reporter) would respond by activating

comX, and that the average activation would increase with the ratio of senders to

receivers. We used defined medium (FMC) to facilitate import of ComS/XIP if present.

We analyzed the green and red fluorescence of the co-cultures to generate

histograms of individual cell fluorescence that reveal both the receiver (green) and

sender (red) *comX* response, shown in Figs. 3A and 3B respectively. Representative

microscopy images are shown in Figs. 3C-3H. In a control chamber containing only

receiver cells exposed to 50 nM XIP, the GFP fluorescence was enhanced, but RFP

fluorescence remained at the baseline level (Fig. 3D and second column in Figs. 3A,

3B). Similarly, a control chamber containing only sender cells showed enhanced RFP

fluorescence, but only baseline GFP fluorescence (Fig. 3H and rightmost column in

Figs. 3A, 3B). As expected, the median RFP fluorescence of the co-cultures

(Fig. 3B, 3F-3H) increased at high sender/receiver ratios. The median RFP

fluorescence of cells in the co-culture increased approximately in proportion to the

density of senders, as expected if each sender activated its own *comX*. RFP

fluorescence was constant over a period of four hours (Fig. S2).

However, when exogenous XIP was not provided, the GFP fluorescence of co-

cultures showed no dependence on the sender/receiver ratio. Over the 0-100% range of

co-culture ratios shown in Fig. 3A, the GFP fluorescence histograms did not exceed the

baseline level of the negative control (Δ*comS* strain alone) in the third column of Fig.

3A. This GFP response did not change appreciably over a period of four hours (Fig.

S2). Senders did not activate *comX* in the Δ*comS* receivers at any mixing ratio.

These data show that although overexpression of *comS* stimulates *comX* within

individual cells, over the 4 h time scale of this experiment this activation did not cause

extracellular, diffusible XIP to accumulate to levels sufficient to activate nearby Δ*comS*

cells.

Growth phase-dependent release of XIP

We previously showed that intercellular signaling by *S. mutans* ComRS can

occur late in growth, but is impeded by deletion of the *atlA* gene, which encodes a major

autolysin (26, 30). Loss of AtlA inhibits cell lysis, which appears to occur primarily in

stationary phase. We therefore tested whether competence signaling from sender

(*comS* overexpressing) to receiver (Δ*comS*) cells can be observed in the later phases of

growth. We prepared sender/receiver co-cultures in different ratios in defined medium

(to favor import of extracellular XIP if present). Because low pH suppresses the *comX*

response, we adjusted the pH during the experiment to ensure that cells would remain

responsive to *comX*-activating signals if present (9, 13). Every 2 h, the pH of the

cultures was adjusted to 7.0 by addition of 2 N NaOH, the OD_{600} was recorded, and an

aliquot of the culture was collected for fluorescence imaging of *comX* promoter activity.

The GFP fluorescence histograms of Fig. 4A show that *comX* expression in the Δ*comS*

strain increased very slightly at 12 h in comparison to expression at 2 h or 8 h. This

increase was more pronounced at higher ratios of sender to receiver cells, consistent

with some release of XIP from lysing senders late in the growth phase. The histograms

of Fig. 4B, like those of Fig. 3B, show a strong RFP response (consistent with

autoactivation of senders) with a slightly stronger response at earlier times.

The median GFP and RFP signals in the above histograms do not shift

dramatically with either time or co-culture ratio. However, the histograms in Fig. 4A

suggest moderate, density-dependent increases in receiver (green) fluorescence at

12 h. Figs. 4C and 4D highlight these changes by showing the value of the 99^{th}

percentile of red and green fluorescence respectively in the cultures, versus optical

density. In Fig. 4C, the GFP fluorescence of the most active receivers increases slightly

at higher OD_{600} values, a change that is slightly more pronounced at higher

sender:receiver ratios. By contrast, Fig. 4D shows no strong trend in the RFP

fluorescence of the 99^{th} percentile of senders, versus OD_{600}. None of the co-cultures

exhibited as strong a GFP response as the positive control (receiver + 50 nM synthetic

XIP, Fig. 4C), indicating that even 12 h growth did not lead to an extracellular XIP

accumulation as large as 50 nM. Overall these data are consistent with robust self-

activation of the *comS* overexpressing strain, accompanied by a modest release of XIP

(or ComS) to the extracellular medium during the late stages of the growth curve.

Binding *in vitro* of ComR to the *comS* and *comX* promoters

The observation of *comX* activation in the overexpressing (sender) strain, without

significant accumulation of XIP in the medium, implies that activation of *comX* does not

require export and reimport of XIP or ComS, if the chromosomal copy of *comS* is intact.

To test whether endogenously produced ComS, acting intracellularly, is sufficient to

activate *comX*, we tested whether unprocessed ComS could enable the binding of

ComR to the *comS* and *comX* promoter regions *in vitro*. Fig. 5A shows a fluorescence

polarization assay with purified recombinant ComR, synthetic XIP or ComS, and a

fluorescently labeled DNA oligomer corresponding to the *S. mutans comX* promoter

region containing the ComR binding site. The assay was performed in the presence and

absence of excess (10 µM) ComS or XIP. Because of the excess of peptide (10 µM)

relative to fluorescent DNA probe (1 nM), the probe polarization depends primarily on

the concentration of ComR added. Fig. 5A shows that in the absence of XIP or ComS,

ComR caused a weak rise in the fluorescence polarization of the DNA oligomer,

indicating relatively weak affinity, as observed for ComR of other streptococci (20).

However, in the presence of ComS or XIP the binding isotherm saturated at lower

ComR concentrations, indicating formation of a complex with higher affinity for the

comX promoter. Histidine tagging of ComR was found to reduce this affinity, as shown

in Fig. S3.

Fig. 5B shows a competition assay in which unlabeled ('cold') P*comX* and

P*comS* DNA oligomers having the same stem-loop structure as the labeled probe were

titrated into samples containing 1.5 µM ComR, 10 µM ComS or XIP, and the labeled

DNA (1 nM). The systematic decrease in polarization is consistent with competition for

ComR. The unlabeled P*comS* and P*comX* probes, which differ by 3 bases, appear to

have identical affinity for ComR.

Fig. 5A confirms a ComR-dependent interaction between ComS and the DNA

probe, although this interaction is weaker than that of XIP. It also shows that, at

saturating concentrations of ComR, ComS elicits only half of the total fluorescence

polarization that results from an equivalent concentration of XIP. This difference may

suggest that ComS and XIP induce a qualitatively different interaction between the

ComR/peptide complexes and the DNA probe. The solid curves in Fig. 5A and 5B are

calculated from a two-step, cooperative binding model (*Methods*) in which the

dissociation constants for the ComR/peptide complex (k_1) and the complex/promoter

(k_2), as well as the order of multimerization n of the ComR/peptide complex, are

variables. Although the data clearly indicate that both ComS and XIP interact with

ComR to bind the DNA probe, they do not permit a precise determination of the ComS

and XIP interaction parameters. As discussed in *Methods*, the data are consistent with a

range of parameter values, corresponding roughly to micromolar k_1 and nanomolar k_2

for both ComS and XIP, and similar cooperativity n for both peptides. The curves in

Fig. 5 show the model with a roughly two fold difference in the ComR dissociation

constants for ComS (k_1 = 3.2 µM, n = 2.5) and XIP (k_1 = 7.3 µM, n = 2.4).

ComRS can drive expression from the *comX* promoter in *E. coli*

As discussed above, the OppA permease, which imports extracellular XIP, is not required for CSP-driven stimulation of ComRS and *comX*. In addition, the export of XIP is accomplished by the non-specific mechanism of lysis. To test whether the ComRS system requires dedicated mechanisms for processing of ComS to XIP, we constructed a dual plasmid system that placed IPTG-inducible *comS* and *comR* into *E. coli* cells carrying a P*comX-gfp* reporter. Fig. 6 shows the GFP activity of *E. coli* BL21(DE3) cells possessing the pDL278 P*comX-gfp* fusion and a pACYC-Duet1 that was either (*a*) empty plasmid or else carried inducible (*b*) *comS,* (*c*) *comR*, or (*d*) *comR* + *comS* (Fig. 6A). Fig. 6B shows GFP activity of the control strains (a-c) in response to added IPTG. Following induction by IPTG, the *comS* and empty vector controls show only baseline GFP expression (Fig. 6B). However, *comR* alone was able to elicit some expression from the *comX* reporter, consistent with the weak DNA binding seen at high ComR concentrations in Fig. 5. When both *comR* and *comS* were present (Fig. 6C), the median *comX* expression was roughly 2-fold greater than with ComR alone at 1 mM IPTG, and roughly 3-fold higher at 300 µM IPTG. These data show that endogenously produced, unprocessed ComS is sufficient to enhance ComR-induced expression from the *comX* promoter.

Data are consistent with an intracellular feedback loop in *comS* transcription

The reduced *comX* response in *comS* deletion strains in Fig. 1A and Fig. 2A shows that an intact *comS* has a stimulatory effect on *comX* expression that is not entirely matched either by exogenous XIP or by overexpression of ComS from a

plasmid. One explanation for this effect is transcriptional positive feedback, in which

endogenously produced ComS retained within the cell interacts with ComR to stimulate

transcription from *comS* and *comX* (Fig. 7A). Import of extracellular XIP would be

expected to stimulate such feedback in cells carrying an intact *comS*, potentially

accounting for the stronger *comX* response to XIP that is observed in the wild type

versus Δ*comS* background.

We constructed a mathematical model of this mechanism as described in the

Methods and the Supplemental Information. The model addresses conditions in a

defined medium where extracellular XIP can be imported to the cell, and early growth

where lysis-driven release of ComS or XIP is not significant. Under these conditions the

model assumes that (I) extracellular XIP is imported to the cell but (II) neither ComS nor

XIP is exported. Further (III) both the ComS and XIP complexes of ComR are able to

elicit transcription of *comS* and *comX*, although not necessarily with equal transcription

rates or degree of multimerization. Assumption (III) is motivated by data indicating that

ComR binding does not absolutely require processing of ComS to XIP, by differences in

the saturating fluorescence polarization achieved with ComS and XIP in Fig. 5A, and by

the different saturating *comX* response (to XIP) of the Δ*comS* and UA159 backgrounds

in Fig. 1A. In parametrizing the model we allow a different order of multimerization in the

ComR/peptide complex that binds DNA: $n = 1$ (ComS) or $n = 2$ (XIP) (see *Discussion*).

In addition the maximal rate of transcription (ComX production) is permitted to depend

on whether a ComR/XIP or a ComR/ComS complex is bound to the promoter.

Fig. 7B shows the results of solving the steady state dynamical equations of the

model with parameter values that best reproduce the data (*Methods*). If only XIP (but

not full-length ComS) interacts with ComR to activate *comX*, the model gives identical

comX activation in both the Δ*comS* and wild-type backgrounds, in contradiction of

Fig. 1A. However if both the ComR/XIP and ComR/ComS complexes are permitted to

induce transcription, although with different maximal rates, the model reproduces the

different *comX* expression in the Δ*comS* and wild-type backgrounds. Parameter values

(Table S2) were found to preserve the relative orders of magnitude of the dissociation

constants of the transcriptional activators found in Fig. 5, with the best fit giving a

transcription rate that is between 4-fold and 200-fold greater for the ComS-ComR bound

promoter than for XIP-ComR activation of the gene.

Discussion

The ComRS system found in mutans, salivarius, pyogenic and bovis streptococci

has been described alternatively as a quorum sensing (19) or timing (25) mechanism

that directly controls *comX*, the master regulator of genetic competence. The ComS-

derived peptide XIP is readily imported by *S. mutans* in defined growth medium, where

it induces transformability with high efficiency. Key evidence supporting an intercellular

signaling role for XIP includes the detection by LC-MS/MS spectroscopy of XIP in

supernates of *S. mutans* that were grown to high cell densities (22, 23). In addition,

filtrates of *S. mutans* cultures grown to high density induced P*comX* activity in reporter

strains (21), indicating the presence of an active competence signal in the extracellular

medium. A recent co-culture study verified that XIP is freely diffusible in aqueous media

and that ComS-overexpressing senders are able to activate *comX* in nearby Δ*comS*

receiver mutants, with no cell-cell contact being required (26). However, deletion of *atlA,*

which encodes a surface-localized protein associated with envelope biogenesis and

autolysis (30), suppressed this intercellular signaling (26). Taken together these data

indicate that the *S. mutans* ComRS system can provide intercellular competence

signaling when autolysis releases sufficient concentrations of ComS or XIP.

We have previously argued that the bimodal expression of *comX* under

stimulation by CSP is a signature that *comX* is controlled by an intracellular

transcriptional feedback system: endogenously produced ComS (or its product XIP)

accumulates within some cells and then interacts with ComR to activate *comX.* Such

feedback mechanisms are often a cause of bimodality in bacterial gene expression,

including in other competence pathways (31, 32). The lack of an established

mechanism for processing and export of ComS, the sensitivity of *comX* bimodality to

endogenous production (via chromosomal *comS)* of XIP, and the fact that XIP import

via Opp is not required for the bimodal behavior, together suggest that purely

intracellular signaling in *comRS* may be possible, at least under some environmental

conditions. Such a mechanism would potentially allow individual *S. mutans* within a

population to differ in competence behavior, as observed experimentally (14).

The present data support this model by providing several lines of evidence that

the chromosomal *comS* plays a role in ComRS activation of *comX*, regardless of other

XIP or ComS that may be supplied. First, although the response of *comX* is different in

complex medium supplemented with CSP than in defined medium supplemented with

XIP, in both cases the behavior of *comX* is affected by the presence of the intact

chromosomal *comS* under the control of its cognate promoter. In complex medium,

comS is required in order for CSP to elicit any *comX* response; in defined media the

deletion of *comS* reduces the *comX* response (both average and variance) to

exogenous XIP, and raises the threshold XIP concentration for a response. Further,

even if it harbors a *comS* overexpression plasmid, a *comS* deletion strain expresses

comX much more weakly than does a *comS*-overexpressing strain that retains its

chromosomal *comS*. These data show that the cell's own native regulation of *comS*

affects its activation of *comX*, independent of whether it overproduces ComS from a

plasmid or imports exogenous XIP via Opp.

Our data also show that even in complex growth medium, which is known to

inhibit the uptake of extracellular XIP, ComS-overexpressing (sender) cells activate their

own *comX.* This autoactivation is unaffected by very rapid exchange (by flow) of the

medium, strongly suggesting that *comX* activation in these cells does not require

accumulation of extracellular XIP. However, this autoactivation requires a native *comS*

in addition to the overexpression plasmid. In addition, even in the ComS overexpression

strain, *comS* mRNA levels are significantly higher when the native *comS* is present, but

comR mRNA levels are unaffected (Fig. S1). These behaviors are consistent with

positive feedback where imported XIP or plasmid-generated ComS enable internal

amplification of *comS* expression.

Finally, the data show that *comS*-overexpressing cells fail to stimulate Δ*comS*

cells (receivers) in co-cultures in defined medium conditions, which are favorable for

import of XIP. As the Δ*comS* receivers do respond to exogenously added XIP, these

data indicate that the overexpressing cells activate their own *comX* without releasing

significant XIP to the medium. The weak intercellular signaling that is observed in co-

cultures grown to late growth phases is consistent with eventual lysis of sender cells,

possibly linked to autoactivation of the lytic pathway driven by ComX and ComDE.

The finding that diffusive signaling by ComS or XIP between *S. mutans* cells is

inefficient or lacks spatial range is consistent with the conclusion reached by Gardan *et*

al. using *S. thermophilus* (25)*.* Those authors found that the type I ComS peptide of *S.*

thermophilus was not secreted at detectable levels in a strain that produced it naturally,

although an overproducing strain did generate detectable ComS in the medium. They

argued that ComS does not diffuse through or accumulate in the medium, although it

may be able to signal between cells that are in physical contact. This proximity model

for ComS resembles a "self-sensing" quorum system (33) in which the secreted signal is

retained at elevated concentrations in the immediate surroundings of the cell, possibly

associated with the cell surface, so that the cell responds somewhat more strongly to its

own secreted signal than to that of the rest of the population.

Our observations suggest more strongly that intercellular quorum signaling

through ComS or XIP is not essential to ComRS control of *comX* in *S. mutans,* under

the conditions examined*.* A more essential component is the dynamics of the cell's own

comS transcription. A plausible mechanism for the bimodal response of *S. mutans*

comX to CSP stimulation is therefore that CSP stimulates the bistable *comRS* feedback

loop by facilitating, through an indirect mechanism, the positive feedback. It would be

sufficient for example to inhibit degradation of endogenous ComS, which depending on

basal transcription levels would trigger *comX* activation in at least some cells, leading to

the bimodal distribution of *comX* activity (11). Notably our data show that

overexpression of *comS* also leads to heterogeneous *comX* activity, suggesting that it

plays a role similar to exogenous CSP by facilitating *comS* autofeedback.

As it is unknown in *S. mutans* whether ComS is processed to XIP inside the cell,

either XIP or ComS could potentially act as the intracellular feedback signal. Although

S. mutans competence was shown to be unresponsive to exogenous full-length ComS

(19), this finding may reflect either selectivity by ComR or simply inefficient import of full

length ComS by Opp. ComS is significantly larger (17 residues) than peptides that are

typically transported by ABC transporters. Shanker *et. al.* (34) found that *S. mutans*

ComR is unresponsive to the ComS peptides produced by other streptococcal species,

although an eight residue XIP (ComS$_{10\text{-}17}$) did interact effectively with ComR to bind the

comS and *comX* promoters (20). Our fluorescence polarization data confirm that both

ComS and XIP can interact with ComR to bind the *comX* and *comS* promoter regions.

They also suggest that ComS and XIP may form ComR complexes of different degrees

of multimerization, a difference that could have interesting consequences for the

nonlinear dynamics of feedback regulation. Our mathematical model for transcriptional

autofeedback in the *comRS* system incorporates the data by assuming that

endogenously produced ComS is not released to the environment, although

extracellular XIP is imported and supplements the endogenous ComS in interacting with

ComR. Unless *E. coli* is equipped to process ComS to XIP, or perhaps that ComS

cleaves autocatalytically, our finding that *E. coli* carrying inducible *comR* and *comS* can

activate a *comX* reporter plasmid is further evidence that processing of ComS to XIP is

not absolutely required.

Other studies show some precedent for such regulation. Structural studies in *S.*

pyogenes have shown that some intracellular Rgg receptor proteins can bind

pheromones that differ in length and sequence (35). Crystallographic structures of

homologous ComR proteins (34, 36) show the SHP binding pocket of the ComR C-

terminus to fall in the tetratricopeptide repeat domain that is responsible for

multimerization, while the N-terminus helix-turn-helix structure binds DNA after an

induced structural rearrangement. The location of the SHP binding pocket could allow

the longer ComS to hinder multimerization when bound, resulting in a monomer binding

to its target, while XIP does not. Fig. S3 shows preliminary evidence that the ComS N-

terminus affects ComR binding in *S. mutans*. Neither ComS nor XIP could induce DNA

binding by an N-terminally 6x histidine tagged ComR, whereas XIP (but not ComS)

caused DNA binding activity in a C-terminally tagged ComR. These data indicate that

steric effects around the SHP binding pocket may influence DNA binding affinities.

Positive feedback occurs in many quorum sensing systems as the accumulation

of the chemical signal in the extracellular environment stimulates the cell to produce

additional signal or its cognate receptor. For example, in *Vibrio fischeri* the C8

homoserine lactone autoinducer stimulates expression of *ainS*, which encodes the

autoinducer synthase (37). In *Vibrio cholerae*, the CAI-1 signal stimulates production of

its receptor CqsS (38). In these cases, the extracellular signal concentration, which is

the positive feedback signal, is sensed by large numbers of cells, and so the population

responds homogenously. However, if an individual cell responds preferentially to its own

signal production, then the feedback signal is specific to the individual cell and the

behavior is qualitatively different. Individual feedback can convert a graded (or

unimodal) population response to a switched or bimodal response (39). Depending on

parameters such as the rate of signal production, noise levels or the cell density, the

response of the cells may then span a range from strongly social or quorum behavior to

purely autocrine or self-sensing (33) behavior in which cells respond independently and

the population becomes heterogeneous (40). Synthetic biology has exploited this

phenomenon in several bacterial quorum sensing systems to amplify the cell's

sensitivity to an exogenous signal. This can lower the quorum circuit's threshold

sensitivity to the signal, and it can also enhance the amplitude of the cell's full response

to that signal. Figs. 1A and 1B suggest that the chromosomal *comS* in *S. mutans*

roughly doubles the amplitude of *comX* response and lowers the XIP sensitivity

threshold roughly two-fold. This amplification is comparable to what was accomplished

in engineered, synthetic systems (41, 42).

As a result the ComRS system may have two modes of function in *S. mutans*. At

low population densities, during early growth, ComRS operates through intracellular

feedback, leading to population bimodality in *comX* expression. Here, only a small

subpopulation of cells activate the late competence genes. However, in later growth

phases or in mature biofilms, stress mechanisms that drive autolysis allow the release

of XIP, providing a diffusible signal that is detected by other cells and amplified through

the internal feedback mechanism to elicit a strong competence response. In this sense,

XIP may serve to broadcast localized stress conditions, stimulating *S. mutans* to

scavenge DNA resources opportunistically from nearby lysing cells (43, 44).

Materials and methods

Strains and growth conditions

S. mutans wild-type strain UA159 and mutant reporting/gene deletion strains

from glycerol freezer stock were grown in BBL BHI (Becton, Dickinson and co.) at 37°C

in 5% CO_2 overnight. Antibiotics were used at the following concentrations where

resistance is indicated in Table 1: erythromycin (10 µg ml^{-1}), kanamycin (1 mg ml^{-1}),

spectinomycin (1 mg ml^{-1}). For experiments in defined medium, strains were washed

twice by centrifugation, removal of supernatant and re-suspension in the defined

medium FMC (45). These were then diluted 20-fold into fresh FMC and allowed to grow

in the same incubator conditions until at optical density at 600 nm (OD_{600}) of 0.1.

Synthetic XIP (sequence GLDWWSL) was synthesized and purified to 98% purity by

NeoBioSci (Cambridge, MA).

E. coli strains were grown in LB at 37 °C shaking in an aerobic incubator

overnight. Antibiotics were used at the following concentrations where resistance is

indicated: ampicillin (10 µg ml^{-1}), chloramphenicol (34 µg ml^{-1}). For recombinant ComR

expression, the next day the overnight cultures were diluted 100-fold into LB containing

ampicillin at the indicated concentration and grown under the same incubator conditions

as overnight. For recombinant *comRS* expression inducing P*comX-gfp,* cells were

diluted 100-fold into LB containing chloramphenicol and spectinomycin and grown

shaking at 37 °C to an OD_{600} of 0.1, at which point IPTG was added and the cells grown

shaking at 30 °C until imaging.

556 **Table 1:** list of strains and plasmids used.

Strain or plasmid	Characteristics[*]	Source or reference
S. mutans strains		
P_comX-gfp_ (plasmid)	UA159 harboring P_comX_ GFP promoter fusion on plasmid pDL278.	(11)
P_comX-rfp_	UA159 harboring P_comX_ dsRed RFP promoter fusion on plasmid pDL278.	(26)
P_comX-gfp_ Δ_comS_	UA159 _comS_ gene replaced with a non-polar erythromycin resistance cassette. Harboring P_comX_ GFP promoter fusion on plasmid pDL278.	(11)
184_comS_ P_comX-rfp_	UA159 harboring pIB184_comS_ and P_comX_ dsRed RFP promoter fusion.	(26)
184_comS_ P_comX-rfp_ Δ_comS_	UA159 harboring pIB184_comS_ and P_comX_ dsRed RFP promoter fusion. _comS_ disrupted by	This study.

	point mutation in start codon (ATG to AAG).	
E. coli strains		
BL21(DE3)	Used for recombinant protein expression	New England Biolabs, MA
10-beta	Used for propagating plasmids during cloning	New England Biolabs, MA
SAMU215	BL21(DE3) carrying P*comX-gfp* on pDL27	This study
SAMU216	SAMU215 carrying pACYC-Duet1 empty vector	This study
SAMU217	SAMU215 carrying pACYC-Duet1 plasmid with *comR* on multiple cloning site (MCS) I	This study
SAMU218	SAMU215 carrying pACYC-Duet1 plasmid with *comS* on MCS II	This study
SAMU219	SAMU215 carrying pACYC-Duet1 plasmid with *comR* on MCS I and *comS* on MCS II	This study

Plasmids		
pIB184	Shuttle expression plasmid with the P23 constitutive promoter, Em^R	(29)
pDL278	*E. coli – Streptococcus* shuttle vector, Sp^R	(46)
pET45b(+)his-*comR*$_{UA159}$	pET45b(+) derivative containing the translational fusion P$_{T7lac}$-*6xhis-comR*$_{UA159}$, Ap^r	This study
pACYC-Duet 1	T7-*lac* inducible expression vector for co-expression of two proteins, Cm^R.	Millipore-Sigma, MA

*Em = erythromycin, Sp = spectinomycin, Ap = ampicillin, Cm = chloramphenicol.

Construction of *comS* point mutant

The start codon of the *comS* gene was mutated from ATG to AAG. The mutation

was introduced directly into the chromosome by site-directed mutagenesis using a PCR

product generated by overlap extension PCR (47). Potential mutants were screened

using mismatch amplification mutation analysis (MAMA) PCR (48), as previously

described (49, 50). The point mutation was confirmed by PCR and sequencing to

ensure that no further mutations were introduced into the *comS* gene and its flanking

regions.

Microfluidic experiments

Microfluidic experiments were performed using a seven-channel PDMS-cast

mixing array device, with fluorescence imaging and single-cell image analysis as

described previously (11, 51, 52). Cells carrying a P*comX-gfp* plasmid-based reporter

were grown to OD_{600} 0.1 from dilution in FMC and sonicated briefly using a Fisher

Scientific FB120 sonic dismembrator probe to split large chains. Sonicated cells were

then loaded into the device through a syringe capped with a 5 µm filter to remove any

remaining aggregations. FMC containing 1 mg ml^{-1} spectinomycin and a XIP gradient

produced from three inlets containing different concentrations of XIP (0 nM, 600 nM and

6 µM XIP inlets) passed through a mixing matrix was pumped through the cell chambers

at a steady rate of 0.08 ml h^{-1} to create a constant, different XIP concentration in each

cell chamber.

Gamma distributions (a two-parameter probability distribution describing the

amount of protein produced in sequential transcription and translation steps) were fit to

the single cell fluorescence distributions using Matlab to fit protein production to

theoretical description (28). The fit was applied to cells fluorescing above an arbitrary

cutoff of 40 units (around background) in order to prevent turned-off cells from skewing

the distribution. Parameters were rounded to three significant figures and reported in

Table S1.

Flow rate dependence experiment

In order to measure the flow rate dependence of XIP signaling we loaded cells

into a commercial six-channel microfluidic slide (IBIDI μ-slide VI, IBIDI GmbH). The six

channels (*a-f*) contained respectively (*a*) a red fluorescent protein (dsRed) *comX*

reporting strain (P*comX-rfp*) control channel flowing fresh BHI at 0.1 ml h^{-1}; (*b*)-(*e*) four

channels containing a *comS* overexpression strain 184*comS* P*comX-rfp* (*comS* on

plasmid pIB184 under the strong constitutive P23 promoter) with BHI at different flow

rates ranging from 0.02 ml h^{-1} up to 1ml h^{-1}, and (*f*) a 184*comS* P*comX-rfp* strain with a

point mutation disrupting the chromosomal *comS* gene (Δ*comS*) under flow at 0.1 ml h^{-}

^{1}. After 2 hours the plain BHI supplied was replaced with BHI supplemented with 50 μg

ml^{-1} chloramphenicol in order to halt further translation and allow any RFP in the cells to

fold. This was supplied at 0.1 ml h^{-1} flow rate for all channels. Four hours (the

maturation time of our RFP) after chloramphenicol addition final fluorescence images of

the cultures were taken. Due to the bimodal *comX* activation in BHI, a fluorescence

cutoff was set as the maximum RFP fluorescence observed in the PcomX-*rfp* negative

control. Cells exhibiting RFP fluorescence above this level were collected in an array

and the size of this sample as a percentage of the population as well as the median of

the above-cutoff fluorescence reported.

Channel co-culture experiment

We loaded co-cultures of a P*comX-gfp* Δ*comS* (responders) with the *comS*

overexpressing strain 184*comS* P*comX-rfp* (senders) into two commercial microfluidic

slides (IBIDI μ-slide VI) using static (not flowing) FMC medium and varying ratios of

comS overproducers:Δ*comS* responders (percentage by volume of OD_{600} 0.1 cultures

vortexed together). P*comX-rfp*, P*comX-gfp* Δ*comS,* P*comX-rfp* + 50 nM XIP and

P*comX-gfp* Δ*comS* + 50 nM XIP were used as controls. The end ports of the channels

were sealed with mineral oil to prevent drying of the medium in the channels. Images

were taken as in the microfluidic experiments and analysis performed similarly. In the

case of controls XIP was added to planktonic culture and the tube vortexed before

pipetting into the slide. Because the population was heterogeneous in both fluorescent

reporter type and *comX* expression, a fluorescence threshold was defined as the

maximum RFP fluorescence observed in the P*comX-rfp* negative control as previously.

The median of the RFP fluorescence observed above this cutoff in other samples was

used as a measure of how strongly the red cells were activating *comX* as a function of

their number density.

OD dependence of co-culture response

For tests of growth-phase dependence of signaling, co-cultures similar to those in

microfluidic channel slides were prepared. Overnight cultures were washed and diluted

40x into fresh FMC containing erythromycin (10 µg ml^{-1}) and spectinomycin (1 mg ml^{-1}).

Once grown to OD_{600} 0.05, these were mixed in ratios varying from to 0% *comS*

overexpressers to 100% overexpressers, defined by volume of *comS* overproducers

added divided by the volume of the Δ*comS* culture added. Low initial cell densities were

used to ensure that early, mid and late growth phases were probed for XIP release.

Every two hours the OD_{600} of the culture and its pH were measured. The pH was

corrected back to 7.0 using 2N sodium hydroxide if it had deviated below 6.5, in order to

measure reaction to any XIP released at late times into the culture. RFP and GFP

fluorescence were measured by pipetting a small amount of the culture onto a glass

coverslip and analyzing single cells. 99[th] percentile GFP fluorescence was then used to

determine if XIP was being released to the *comS* mutants in an OD_{600}-dependent

manner.

RT-qPCR measurement of *comS, comR* and *comX* transcripts

S. mutans cells were diluted 20-fold into BHI (P*comX-gfp* WT/BHI, 184*comS*

P*comX-rfp*, 184comS P*comX-rfp* ∆comS samples) or FMC (P*comX-gfp* WT/FMC ± XIP,

P*comX-gfp* ∆*comS* ± XIP samples, pIB184 / WT, pIB184ComS / WT). Where added,

XIP was supplied at OD_{600} = 0.1. Cells were harvested at OD_{600} = 0.5 by centrifugation

and re-suspended in RNA protectant buffer for 10 minutes. Samples were then

centrifuged, the supernatant removed and the pellets frozen at -80 °C. RNA extraction

was performed using the Qiagen RNEasy mini kit (Qiagen, USA). RNA sample

concentration and purity were measured using a Thermo Scientific NanoDrop One

Microvolume UV-Vis Spectrophotometer (Thermo Scientific, USA). 1 µg of RNA was

then reverse transcribed to cDNA using the Bio-Rad iScript reverse transcription kit with

random primers (Bio-Rad, USA). The qPCR was performed on a Bio-Rad CFX96 Real-

Time System using Bio-Rad Sso Advanced Universal SYBR Green Supermix with a 50-

fold dilution of the cDNA and 500 nM gene-specific primers. Sequences used for the

primers are given in Table S3 (supplemental information). A standard curve across 8

orders of magnitude of transcript copies (from 10^8 to 10^1) was used to determine

transcript count for each gene. For each sample the *comX*, *comR* and *comS* transcript

counts were then normalized by the 16S rRNA count for the same sample. Fig. S1

shows the median of this ratio, with error bar lengths given by the range from second

lowest to second highest ratio obtained.

Fluorescence polarization

ComR protein was obtained by cloning the *comR* gene into the 6x-His tagged

site on pET-45b(+) vector in *E. coli* 10-beta using standard PCR cloning methods. His-

ComR was then expressed in *E. coli* BL21(DE3) by induction with 1mM IPTG at mid-

exponential phase in LB. After 4 hours cells were lysed using lysozyme in B-PER lysis

buffer (ThermoFisher). Protein was then purified from clarified lysate using Ni-NTA

agarose affinity chromatography and the histidine tag cleaved using enterokinase max

at 4°C (EKMax, Invitrogen). The resulting protein solution was dialyzed into PBS pH 7.4

for experimental use. Native ComR concentration was measured by the Pierce BCA

assay (Thermo Scientific) and purity of the cleaved form verified by SDS-PAGE run

against an uncleaved sample.

Fluorescence polarization assays were performed in a 96-well plate with black

bottom and black sides in a Biotek Synergy 2 plate reader (Biotek Instruments inc.) in

the polarization mode. A 5' Bodipy FL-X labeled self-annealing stem-loop DNA strand

with sequence corresponding to PcomX (sequence 5'-BODIPY FL-X -

ATGGGACATTTATGTCCTGTCCCCCACAGGACATAAATGTCCCAT - 3'), synthesized

by ThermoFisher) was used as the binding aptamer and a filter set with excitation 485

nm, emission 528 nm was used for fluorescence excitation. 1 nM labeled DNA probe

was added to a reaction buffer previously described (53) supplemented with 1 mM

EDTA and 0.05 mg ml^{-1} salmon DNA. ComR was titrated in concentration in this buffer

alone, in the presence of 10 µM XIP or in the presence of 10 µM *comS*. The reactions

were incubated at 37°C for 20 minutes before reading. Synthetic ComS (sequence

MFSILTSILMGLDWWSL) for fluorescence polarization was synthesized and purified to

60% purity by Biomatik (Wilmington, DE).

Competing unlabeled probe assays were performed with 1.5 µM ComR in the

same buffer containing 1 nM P*comX* fluorescent DNA and 10 µM SHP (either ComS or

XIP). An unlabeled probe corresponding to either the P*comS* (sequence 5' - ACG

GGACATAAATGTCCTGTCCCCCACAGGACATTTATGTCCCGT - 3'), synthesized by

Thermo Fisher) or the above P*comX* probe was titrated into this solution and the

decreasing polarization plotted. Reactions were again incubated at 37°C for 20 minutes

before polarization readings taken. In all FP experiments reading was performed three

times on the same plate to estimate instrument error. The average polarization was

used for plotting and analysis. Details and parameters of the two-step binding model for

the FP data are given in the supporting information.

ComRS expression in *E. coli*

A recombinant *comRS* system was produced in *E. coli* by inserting *comR* into

multiple cloning site (MCS) I in the pACYC-Duet 1 vector and *comS* into MCS II by

standard cloning methods. BL21(DE3) cells containing P*comX-gfp* on pDL278 were

transformed with the resulting pACYC-Duet 1 *comRS* plasmid. Controls consisting of an

empty pACYC-Duet 1 and the vector containing only *comR* at MCS I and only *comS* at

MCS II were produced in the same way. Cells were diluted 100-fold from overnight into

LB containing chloramphenicol and spectinomycin and grown shaking at 37 °C until

they reached OD$_{600}$ 0.1. At this point IPTG was added in varying amounts to induce

expression of the T7-*lac* controlled recombinant system. Following addition of IPTG the

cells were grown with shaking at 30 °C and GFP fluorescence was imaged after 3 h.

Mathematical model of *comRS* control of *comX*

Deterministic modeling of *comX* activation by *comRS* was performed by least

squares fitting a chemical equilibrium model to the microfluidic data from experiments

for each of the wild type background P*comX* GFP strain and the Δ*comS* cells. Details of

the model and the robustness analysis are in the supporting information, with parameter

values given in Table S2.

Acknowledgments

This work was supported by 1R01 DE023339 from the National Institute of Dental and

Craniofacial Research. The authors thank Minjun Son, Chris Browngardt, Natalie

Maricic, Lin Zeng, Hey Min Kim and Sang-Joon Ahn for helpful discussions, provision of

mutant strains and advice on experimental procedures.

References

1. **Loesche, WJ.** 1986. Role of *Streptococcus mutans* in human dental decay.

Microbiol. Rev. **50:**353-380.

2. **Senadheera, DB, Cordova, M, Ayala, EA, Chavez de Paz, LE, Singh, K, Downey,**

JS, Svensater, G, Goodman, SD, Cvitkovitch, DG. 2012. Regulation of bacteriocin

production and cell death by the VicRK signaling system in *Streptococcus mutans*. J.

Bacteriol. **194:**1307-1316.

3. **Ahn, SJ, Lemos, JA, Burne, RA.** 2005. Role of HtrA in growth and competence of

Streptococcus mutans UA159. J. Bacteriol. **187:**3028-3038.

4. **Senadheera, MD, Lee, AW, Hung, DC, Spatafora, GA, Goodman, SD,**

Cvitkovitch, DG. 2007. The *Streptococcus mutans vicX* gene product modulates *gtfB/C*

expression, biofilm formation, genetic competence, and oxidative stress tolerance. J.

Bacteriol. **189:**1451-1458.

5. **Tremblay, YDN, Lo, H, Li, Y, Halperin, SA, Lee, SF.** 2009. Expression of the

Streptococcus mutans essential two-component regulatory system VicRK is pH and

growth-phase dependent and controlled by the LiaFSR three-component regulatory

system. Microbiology. **155:**2856-2865.

6. **Ahn, SJ, Wen, ZT, Burne, RA.** 2006. Multilevel control of competence development

and stress tolerance in *Streptococcus mutans* UA159. Infect. Immun. **74:**1631-1642.

7. **Qi, F, Merritt, J, Lux, R, Shi, W.** 2004. Inactivation of the *ciaH* Gene in

Streptococcus mutans diminishes mutacin production and competence development,

alters sucrose-dependent biofilm formation, and reduces stress tolerance. Infect.

Immun. **72:**4895-4899.

8. **Son, M, Ghoreishi, D, Ahn, SJ, Burne, RA, Hagen, SJ.** 2015. Sharply Tuned pH

Response of Genetic Competence Regulation in *Streptococcus mutans*: a Microfluidic

Study of the Environmental Sensitivity of *comX*. Appl. Environ. Microbiol. **81:**5622-5631.

9. **Guo, Q, Ahn, SJ, Kaspar, J, Zhou, X, Burne, RA.** 2014. Growth phase and pH

influence peptide signaling for competence development in *Streptococcus mutans*. J.

Bacteriol. **196:**227-236.

10. **Moye, ZD, Son, M, Rosa-Alberty, AE, Zeng, L, Ahn, SJ, Hagen, SJ, Burne, RA.**

2016. Effects of Carbohydrate Source on Genetic Competence in *Streptococcus*

mutans. Appl. Environ. Microbiol. **82:**4821-4834.

11. **Son, M, Ahn, S, Guo, Q, Burne, RA, Hagen, SJ.** 2012. Microfluidic study of

competence regulation in *Streptococcus mutans*: environmental inputs modulate

bimodal and unimodal expression of *comX*. Mol. Microbiol. **86:**258-272.

12. **Reck, M, Tomasch, J, Wagner-Döbler, I.** 2015. The Alternative Sigma Factor SigX

Controls Bacteriocin Synthesis and Competence, the Two Quorum Sensing Regulated

Traits in *Streptococcus mutans*. PLOS Genetics. **11:**e1005353.

13. **Son, M, Shields, RC, Ahn, S, Burne, RA, Hagen, SJ.** 2015. Bidirectional signaling

in the competence regulatory pathway of *Streptococcus mutans*. FEMS Microbiol. Lett.

362.

14. **Stephen J Hagen and,Minjun Son.** 2017. Origins of heterogeneity in

Streptococcus mutans competence: interpreting an environment-sensitive signaling

pathway. Physical Biology. **14:**015001.

15. **Khan, R, Rukke, HV, Høvik, H, Åmdal, HA, Chen, T, Morrison, DA, Petersen,**

FC. 2016. Comprehensive Transcriptome Profiles of *Streptococcus mutans* UA159 Map

Core Streptococcal Competence Genes. mSystems. **1 (2)**.

16. **Fontaine, L, Wahl, A, Fléchard, M, Mignolet, J, Hols, P.** 2015. Regulation of

competence for natural transformation in streptococci. Infection, Genetics and

Evolution. **33:**343-360.

17. **Hossain, MS, Biswas, I.** 2012. An extracelluar protease, SepM, generates

functional competence-stimulating peptide in *Streptococcus mutans* UA159. J.

Bacteriol. **194:**5886-5896.

18. **Perry, JA, Jones, MB, Peterson, SN, Cvitkovitch, DG, Lévesque, CM.** 2009.

Peptide alarmone signalling triggers an auto-active bacteriocin necessary for genetic

competence. Mol. Microbiol. **72:**905-917.

19. **Mashburn-Warren, L, Morrison, DA, Federle, MJ.** 2010. A novel double-

tryptophan peptide pheromone controls competence in *Streptococcus* spp. via an Rgg

regulator. Mol. Microbiol. **78:**589-606.

20. **Fontaine, L, Goffin, P, Dubout, H, Delplace, B, Baulard, A, Lecat-Guillet, N,**

Chambellon, E, Gardan, R, Hols, P. 2013. Mechanism of competence activation by

the ComRS signalling system in streptococci. Mol. Microbiol. **87:**1113-1132.

21. **Desai, K, Mashburn-Warren, L, Federle, MJ, Morrison, DA.** 2012. Development

of competence for genetic transformation of *Streptococcus mutans* in a chemically

defined medium. J. Bacteriol. **194:**3774-3780.

22. **Khan, R, Rukke, HV, Ricomini Filho, AP, Fimland, G, Arntzen, MO, Thiede, B,**

Petersen, FC. 2012. Extracellular identification of a processed type II ComR/ComS

pheromone of *Streptococcus mutans*. J. Bacteriol. **194:**3781-3788.

23. **Wenderska, IB, Lukenda, N, Cordova, M, Magarvey, N, Cvitkovitch, DG,**

Senadheera, DB. 2012. A novel function for the competence inducing peptide, XIP, as

a cell death effector of *Streptococcus mutans*. FEMS Microbiol. Lett. **336:**104-112.

24. **Chang, JC, Federle, MJ.** 2016. PptAB Exports Rgg Quorum-Sensing Peptides in

Streptococcus. Plos One. **11:**e0168461.

25. **Gardan, R, Besset, C, Gitton, C, Guillot, A, Fontaine, L, Hols, P, Monnet, V.**

2013. Extracellular Life Cycle of ComS, the Competence-Stimulating Peptide of

Streptococcus thermophilus. Journal of Bacteriology. **195:**1845-1855.

26. **Kaspar, J, Underhill, SAM, Shields, RC, Reyes, A, Rosenzweig, S, Hagen, SJ,**

Burne, RA. 2017. Intercellular communication via the *comX*-Inducing Peptide (XIP) of

Streptococcus mutans. J. Bacteriol. .

27. **Taniguchi, Y, Choi, PJ, Li, G, Chen, H, Babu, M, Hearn, J, Emili, A, Xie, XS.**

2010. Quantifying *E. coli* Proteome and Transcriptome with Single-Molecule Sensitivity

in Single Cells. Science. **329:**533.

28. **Friedman, N, Cai, L, Xie, XS.** 2006. Linking stochastic dynamics to population

distribution: an analytical framework of gene expression. Phys. Rev. Lett. **97:**168302.

29. **Biswas, I, Jha, JK, Fromm, N.** 2008. Shuttle expression plasmids for genetic

studies in *Streptococcus mutans*. Microbiology. **154:**2275-2282.

30. **Ahn, SJ, Burne, RA.** 2006. The *atlA* operon of *Streptococcus mutans*: role in

autolysin maturation and cell surface biogenesis. J. Bacteriol. **188:**6877-6888.

31. **Dubnau, D, Losick, R.** 2006. Bistability in bacteria. Mol. Microbiol. **61:**564-572.

32. **Maamar, H, Dubnau, D.** 2005. Bistability in the *Bacillus subtilis* K-state

(competence) system requires a positive feedback loop. Mol. Microbiol. **56:**615-624.

33. **Bareia, T, Pollak, S, Eldar, A.** 2017. Self-sensing in *Bacillus subtilis* quorum-

sensing systems. Nature Microbiology.

34. **Shanker, E, Morrison, DA, Talagas, A, Nessler, S, Federle, MJ, Prehna, G.**

2016. Pheromone Recognition and Selectivity by ComR Proteins among *Streptococcus*

Species. PLOS Pathogens. **12:**e1005979.

35. **Aggarwal, C, Jimenez, JC, Nanavati, D, Federle, MJ.** 2014. Multiple length

peptide-pheromone variants produced by *Streptococcus pyogenes* directly bind Rgg

proteins to confer transcriptional regulation. J. Biol. Chem. **289:**22427-22436.

36. **Talagas, A, Fontaine, L, Ledesma-Garca, L, Mignolet, J, Li de, IS, Lazar, N,**

Aumont-Nicaise, M, Federle, MJ, Prehna, G, Hols, P, Nessler, S. 2016. Structural

Insights into Streptococcal Competence Regulation by the Cell-to-Cell Communication

System ComRS. PLOS Pathogens. **12:**e1005980.

37. **Lupp, C, Ruby, EG.** 2004. *Vibrio fischeri* LuxS and AinS: comparative study of two

signal synthases. J. Bacteriol. **186:**3873-3881.

38. **Hurley, A, Bassler, BL.** 2017. Asymmetric regulation of quorum-sensing receptors

drives autoinducer-specific gene expression programs in *Vibrio cholerae*. PLOS

Genetics. **13:**e1006826.

39. **Becskei, A, Séraphin, B, Serrano, L.** 2001. Positive feedback in eukaryotic gene

networks: cell differentiation by graded to binary response conversion. Embo J.

20:2528.

40. **Youk, H, Lim, WA.** 2014. Secreting and sensing the same molecule allows cells to

achieve versatile social behaviors. Science. **343:**1242782.

41. **Sayut, DJ, Niu, Y, Sun, L.** 2006. Construction and Engineering of Positive

Feedback Loops. ACS Chem. Biol. **1:**692-696.

42. **Nistala, GJ, Wu, K, Rao, CV, Bhalerao, KD.** 2010. A modular positive feedback-

based gene amplifier. J. Biol. Eng. **4:**4-1611-4-4.

43. **Leung, V, Dufour, D, Lévesque, C,M.** 2015. Death and survival in *Streptococcus*

mutans: differing outcomes of a quorum-sensing signaling peptide. Frontiers in

Microbiology. **6:**1176.

44. **Perry, JA, Cvitkovitch, DG, Lévesque, CM.** 2009. Cell death in *Streptococcus*

mutans biofilms: a link between CSP and extracellular DNA. FEMS Microbiol. Lett.

299:261-266.

45. **Terleckyj, B, Willett, NP, Shockman, GD.** 1975. Growth of several cariogenic

strains of oral streptococci in a chemically defined medium. Infect. Immun. **11:**649-655.

46. **LeBlanc, DJ, Lee, LN, Abu-Al-Jaibat, A.** 1992. Molecular, genetic, and functional

analysis of the basic replicon of pVA380-1, a plasmid of oral streptococcal origin.

Plasmid. **28:**130-145.

47. **Ho, SN, Hunt, HD, Horton, RM, Pullen, JK, Pease, LR.** 1989. Site-directed

mutagenesis by overlap extension using the polymerase chain reaction. Gene. **77:**51-

59.

48. **Cha, RS, Zarbl, H, Keohavong, P, Thilly, WG.** 1992. Mismatch amplification

mutation assay (MAMA): application to the *c-H-ras* gene. PCR Methods Appl. **2:**14-20.

49. **Zeng, L, Das, S, Burne, RA.** 2010. Utilization of lactose and galactose by

Streptococcus mutans: transport, toxicity, and carbon catabolite repression. J. Bacteriol.

192:2434-2444.

50. **Ahn, SJ, Kaspar, J, Kim, JN, Seaton, K, Burne, RA.** 2014. Discovery of novel

peptides regulating competence development in *Streptococcus mutans*. J. Bacteriol.

196:3735-3745.

51. **Jeon, NL, Dertinger, SKW, Chiu, DT, Choi, IS, Stroock, AD, Whitesides, GM.**

2000. Generation of Solution and Surface Gradients Using Microfluidic Systems.

Langmuir. **16:**8311-8316.

52. **Kwak, IH, Son, M, Hagen, SJ.** 2012. Analysis of gene expression levels in

individual bacterial cells without image segmentation. Biochem. Biophys. Res.

Commun. **421:**425-430.

53. **Aggarwal, C, Jimenez, JC, Lee, H, Chlipala, GE, Ratia, K, Federle, MJ.** 2015.

Identification of Quorum-Sensing Inhibitors Disrupting Signaling between Rgg and Short

Hydrophobic Peptides in Streptococci. MBio. **6:**e00393-15.

54. **Lemme, A, Grobe, L, Reck, M, Tomasch, J, Wagner-Dobler, I.** 2011.

Subpopulation-specific transcriptome analysis of competence-stimulating-peptide-

induced *Streptococcus mutans*. J. Bacteriol. **193:**1863-1877.

Figure legends

Fig. 1: *comS* deletion is not fully complemented by synthetic XIP

(A) Comparison of P*comX-gfp* activity in *S. mutans* cells of the UA159 (wild type) background (magenta) and Δ*comS* mutant background (cyan). The median GFP fluorescence is shown for cells that were supplied with continuous flow of exogenous synthetic XIP in microfluidic chambers. Data are shown at 30 minutes (circles, dash-dot lines) and 90 minutes (squares, dashed lines) of flow. The smooth curves are spline fits to the data. (B) Median GFP levels in the two strains at the 90 minute time point of the flow experiment. Also shown are the histograms of the individual cell P*comX-gfp* reporter activity, versus exogenous XIP concentration, for (C) the UA159 background and (D) the Δ*comS* background. Solid black curves in (C) and (D) show the best fit gamma probability distribution for each histogram. A cutoff of 40 units of P*comX-gfp* fluorescence has been applied to exclude background autofluorescence. (E) Parameter *a* of the (two parameter) gamma probability distribution, obtained from fits in (C)-(D), reflecting the ratio of transcription rate to protein degradation rate, and (F) the distribution parameter *b*, reflecting the ratio of translation rate to the rate of mRNA degradation. In (E)-(F) cyan indicates the Δ*comS* mutant, and magenta indicates the UA159 background.

Fig. 2: Activation of ComX in a ComS-overexpressing strain is independent of rate of medium replacement

P*comX-rfp* reporter activity is shown in cells growing in microfluidic chambers supplied with continuously flowing fresh complex medium (BHI). (A) Histograms of individual cell

PcomX-rfp reporter fluorescence: (Leftmost column) wild-type background (negative

control) with flow at 0.1 ml h^{-1}; (Second column) ComS-overexpressing 184comS

ΔcomS background at 0.1 ml h^{-1}; (Columns 3-5) ComS-overexpressing (184comS)

background at 0.02 ml h^{-1}, 0.1 ml h^{-1}, 0.5 ml h^{-1} and 1 ml h^{-1}. (At a flow rate of 1 ml h^{-1}

the medium in each flow chamber is replaced every 6 s.) (B) RFP fluorescence of cells

that exceeded the wild type (negative control) red fluorescence in column 1 of (A). The

black bar indicates the median of data in each channel: (Leftmost column) 184comS

ΔcomS background; (Columns 2-5) ComS-overexpressing (184comS) background. All

RFP measurements were made 4 hours after addition of chloramphenicol to the

cultures.

Fig. 3: ComS overexpresser does not induce *comX* of Δ*comS* strain in coculture

Histograms of (A) GFP and (B) RFP fluorescence of individual cells in cocultures of

sender (184comS PcomX-rfp) and receiver (PcomX-gfp ΔcomS) strains. Strains were

grown to equal optical density, mixed in varying proportion, and then incubated in

microfluidic chambers containing stationary defined medium. Blue lines indicate

population medians. Panels (A) and (B) show fluorescence of: (Leftmost column)

UA159 background strain containing PcomX rfp reporter, without added XIP (negative

control); (Second column) PcomX-gfp ΔcomS (receiver) with added 50 nM XIP (positive

control); (Columns 3-11) Cocultures of sender and receiver, with columns labeled by

percentage by volume of 184comS (sender) culture in the initial preparation of the co-

culture. (C-H) Phase contrast images of cocultures, overlaid with red and green

fluorescence images: (C) PcomX-rfp reporter in UA159 background, with no added XIP

(negative control); (D) P*comX-gfp* Δ*comS* cells with 50 nM added XIP (positive control);

(E) P*comX-gfp* Δ*comS* (receiver) alone, with 0% sender; (F)-(H) cocultures containing

30%, 80%, and 100% sender respectively.

Fig. 4: Evidence for release of XIP in cocultures late in growth

Histograms of (A) GFP and (B) RFP fluorescence of individual cells in cocultures of

receiver (P*comX-gfp* Δ*comS*) and sender (P*comX-rfp* 184*comS*) strains, following

different incubation periods. Labels at top indicate the volume fraction comprised by the

sender strain in the preparation of the coculture. Histogram colors indicate incubation

times: 2 h (cyan), 8 h (magenta), 12 h (green). The lower panels show the 99[th]

percentile of the individual cell GFP (C) and RFP (D) fluorescence observed in the

culture, versus the cultureOD$_{600}$. Exogenous XIP was not added, except in the positive

(sender-only) control sample indicated by the inverted triangles in (C) and (D).

Fig. 5: ComS and XIP interact with ComR to bind the *comX* promoter

Fluorescence polarization assay testing interaction of ComS and XIP peptides with

ComR and the P*comX* transcriptional activation site. The DNA probe is labeled with a

Bodipy FL-X fluorophore. (A) Titration of ComR into a solution containing 10 μM of XIP

(blue) or full length ComS (green), labeled DNA probe (1 nM), and 0.05 mg ml^{-1} salmon

DNA. Negative control (black) contains no ComS or XIP peptide. (B) Competition assay

in which unlabeled promoter sequence DNA was titrated into a solution containing

ComR (1.5 μM), fluorescent DNA probe (1 nM) and peptide (either ComS or XIP, 10

μM). Unlabeled P*comX* DNA was used with XIP (blue) and ComS (green). An unlabeled

P*comS* probe was also tested for its ability to compete with the fluorescent P*comX*

probe in the presence of XIP (red) and ComS (gold). Solid curves indicate binding and

competition behavior predicted by the two step model described in *Methods*, in which

peptide (ComS or XIP) first forms a multimeric complex with ComR (k_1, n), and a single

copy of this complex binds to the (labeled or unlabeled) P*comX* DNA (k_2). For ComS

binding/competition (green), the curves represent $k_1 = 3.2$ µM, $k_2 = 2.2$ nM, $n = 2.5$. For

XIP binding/competition (blue), the curves represent $k_1 = 7.3$ µM, $k_2 = 33$ nM, $n = 2.4$.

Fig. 6: Activation of P*comX* by inducible *comRS* in *E. coli*

Activity of a P*comX-gfp* reporter within individual *E.coli* cells measured by fluorescence

microscopy. Strains (A) harbored a two-plasmid system that included the P*comX-gfp*

reporter as well as different combinations of *comR* and *comS* under control of the T7-*lac*

promoter**.** (B) GFP fluorescence of controls containing the pACYC-Duet1 empty vector,

comR alone or *comS* alone after 3 h of growth with shaking, following addition of 0, 300

µM and 1 mM IPTG. (C) GFP fluorescence of the strain possessing both *comR* and

comS after 3 h of growth with shaking in varying IPTG concentrations. The blue bar

represents the median cell fluorescence.

Fig. 7: proposed model for *comRS* feedback regulation of *comX*.

Model for *comS*-feedback enhanced activation of P*comX-gfp* reporter by exogenous

synthetic XIP. (A) Illustration of the feedback model and the role of *comS* and

extracellular XIP. ComS and XIP both interact with ComR to activate transcription of

comS and *comX.* At low concentrations of extracellular XIP, *comS* expression is very

low and *comX* is not expressed. At high concentrations of extracellular XIP, XIP is

imported efficiently by Opp and interacts with ComR to drive expression of both *comS*

and *comR*. Endogenously produced ComS is not readily exported in the absence of

lysis, and so intracellular accumulation of ComS drives elevated *comS* and *comX*

expression. Consequently *comX* expression at any given XIP concentration is boosted

by *comS* feedback. Cells lacking native *comS* can respond to synthetic XIP but cannot

activate *comX* to the same level as wild type. The figure describes behavior in defined

medium; in complex medium extracellular XIP is not imported (11). (B) Comparison of

model simulation with data. Red circles indicate median P*comX-gfp* fluorescence of the

UA159 background strain supplied with synthetic XIP in microfluidic flow; blue circles

indicate the median P*comX-gfp* fluorescence of the Δ*comS* background. Solid curves

represent calculated values from a fit in which 11 parameters were fit to the microfluidic

data, as described in *Methods.* The model relates the predicted ComX concentration to

the median GFP fluorescence by an offset and scale factor.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

B