Automated classification of synaptic vesicles in electron tomograms of *C. elegans* using machine learning

Kristin Verena Kaltdorf[1,2,3], Maria Theiss[2,3], Sebastian Markert[1], Mei Zhen[4], Thomas Dandekar[2,&], Christian Stigloher[1,&], Philip Kollmannsberger[3,&]

[&] TD, CS and PK are Joint Senior Authors.

[1]Imaging Core Facility, Biocenter, University of Würzburg, 97074 Würzburg, Germany

[2]Department of Bioinformatics, Biocenter, University of Würzburg, 97074 Würzburg, Germany

[3]Center for Computational and Theoretical Biology, Hubland North, building 32, University of Würzburg, 97074 Würzburg, Germany

[4]Department of Molecular Genetics, University of Toronto, Toronto, M5G 1X5, Canada

Correspondence to: Philip.Kollmannsberger@uni-wuerzburg.de

1.   Abstract

Synaptic vesicles (SVs) are a key component of neuronal signaling and fulfil different roles depending on their composition. In electron micrograms of neurites, two types of vesicles can be distinguished by morphological criteria, the classical "clear core" vesicles (CCV) and the typically larger "dense core" vesicles (DCV), with differences in electron density due to their diverse cargos. Compared to CCVs, the precise function of DCVs is less defined. DCVs are known to store neuropeptides, which function as neuronal messengers and modulators [1]. In *C. elegans*, they play a role in locomotion, dauer formation, egg-laying, and mechano- and chemosensation [2]. Another type of DCVs, also referred to as granulated vesicles, are known to transport Bassoon, Piccolo and further constituents of the presynaptic density in the center of the active zone (AZ), and therefore are important for synaptogenesis [3].

To better understand the role of different types of SVs, we present here a new automated approach to classify vesicles and to quantify their properties from electron tomograms. We combine machine learning with an extension of our previously developed vesicle segmentation workflow, the ImageJ macro 3D ART VeSElecT, to reliably distinguish CCVs and DCVs using image-based features. We apply this method to electron tomograms of the *C. elegans* NMJs, and find an increased fraction of DCVs as well as a higher mean distance between DCVs and AZs in dauer larvae when compared to adult hermaphrodites. Our machine learning approach could be applied to study properties of different synaptic vesicle pools in electron tomograms of diverse model organisms.

44 2.   Author summary

Vesicles are important components of the cell, and synaptic vesicles are central for

neuronal signaling. Two types of synaptic vesicles can be distinguished by electron

microscopy: the classical "clear core" vesicles (CCVs) and the typically larger "dense

core" vesicles (DCVs). The distinct appearance of vesicles is caused by their different

cargos.

To distinguish both vesicle types automatically and rapidly in 3D electron

micrographs, we present here a new automated approach to classify vesicles and

quantify their properties from electron tomograms. We combine machine learning with

an extension of our previously developed vesicle segmentation workflow, an ImageJ

macro, to reliably distinguish CCVs and DCVs using specific features in the images.

The approach allows to compare vesicle changes and was compared and validated

by microscopy experts on hand curated data-sets.

An application example compares vesicles of *C. elegans* dauer larvae and adult

hermaphrodites. The technique can be transferred to more extensive comparisons in

both stages as well as to other neurobiology questions regarding synaptic vesicles.

3.   Introduction

Caenorhabditis elegans is a well-studied model organism. Its small nervous system,

consisting of 302 neurons [4,5] allows for studies of the connectome in its entirety [5–

7]. A focus of research is on *C. elegans* young adult hermaphrodites, which makes

them a reference system for comparison with other developmental states in the life

cycle. The most remarkable of all *C. elegans* developmental states is the dauer larva.

When the L1 stage larvae sense adverse environmental conditions such as

heat, overpopulation and starvation, they activate an alternative state, diapause,

instead of continuing the reproductive life cycle [8,9]. Dauer larvae are characterized

by their ability to survive several months without any food uptake [10]. Although

metabolic activity is reduced [10], dauer larvae are able to move quickly, if necessary,

in particular to escape adverse environmental stimuli and in order to reach and

colonize new habitats. Improvement of environmental conditions leads to dauers to

exit diapause, resume development into pre-adult L4 larvae, and re-enter the

reproductive life cycle [9]. The mechanisms behind this strikingly fast developmental

transition, and the role of the dauer nervous system in sensing and responding to

environmental cues, remain largely enigmatic.

Neuronal signaling plays a critical role in regulating the dauer stage.

Neuronally released insulin-like peptides [11–14] and morphogens such as TGF-beta

[15,16] regulate the activation of dauer diapause. These observations suggest

potential changes of the nature of neuronal signaling in dauer larvae.

Neuropeptides are stored and released through dense core vesicles (DCVs)

[1,17,18]. Recent studies, particularly the electron microscopic reconstruction of *C.*

elegans prepared by high-pressure freezing, reveal that almost all synapses,

including the neuromuscular junctions (NMJs), contain two principal types of vesicles,

a small fraction of DCVs and the much more abundant clear core vesicles (CCVs)

[19–21]. CCVs are characterized by their clear inner core in electron micrographs,

which stands in strong contrast to the stained vesicle membrane. CCVs typically

contain neurotransmitters such as acetylcholine and GABA in the *C. elegans* NMJs

that are released into the synaptic cleft after fusion of the CCV with the presynaptic

cell membrane at the release sites at active zones [2]. In contrast to CCVs, the role of

DCVs is much more enigmatic. DCVs are characterized by their eponymous dark,

electron dense core that typically contains neuropeptides, small protein molecules

with about 3 - 100 amino acids [1] or other protein cargos such as active zone

proteins [3]. DCVs have a bigger average size of about 60 - 80 nm in comparison to

CCVs, which have a much smaller diameter [19], that can vary in different organisms

or developmental states [22,23]. Whereas CCVs cluster around the active zone,

DCVs tend to locate further away from the release site, typically at the rim of the SV

pool as described for *C. elegans* NMJs [19].

DCV exocytosis is considered to medicate slow synaptic and non-synaptic

signaling in neurons [1]. It typically occurs independently of the active zone and only

at high rates of neuronal activity [1,2]. In *C. elegans*, neuropeptides participate in a

wide variety of behaviors such as egg-laying, locomotion, dauer formation,

mechano- and chemosensation depending on the expression pattern of the peptides

[2]. Despite the known critical involvement of neural peptides in dauer development,

whether the dauer states exhibit differences in DCV distribution in the nervous

system remains unknown.

We used high-resolution 3D transmission electron microscopy (electron

tomography) to systematically investigate differences of DCV and CCV in NMJs of

C. elegans adult hermaphrodites and dauer larvae. Series of 2D images were taken

from different viewing angles by tilting the sample, and reconstructed *in silico* to

obtain one 3D image stack with near isotropic resolution of typically 4 – 5 nm,

determined by measuring the "unit membrane" thickness [19,25–28]. Quantification of

SV in 3D electron tomograms requires reconstruction of the vesicle pool of a sufficient

number of tomograms, which is very time-consuming and prone to mistakes when

performed manually.

To circumvent these shortcomings we previously developed *3D ART*

VeSElecT, a workflow for automated reconstruction of vesicles [22]: This ImageJ/Fiji

tool allows for automated reconstruction of synaptic vesicle pools in electron

tomograms, but is not able to differentiate between different types of vesicles. To

solve this issue, we here present a python-based machine learning approach that

trains vesicle classifiers, and a straightforward ImageJ/Fiji tool that applies a trained

classifier to previously detected vesicles.

We show that our technique can detect differences between the DCV and CCV

ratio in two developmental states, adult hermaphrodites and dauer larvae. The

application of these image analysis and machine learning tools for electron

tomograms is in principle useful to study differences in synaptic vesicle pools of many

other model organisms.

4. Results

3.1 Manual classification shows three distinct vesicle types

To compare vesicle properties in two different developmental states of *C. elegans*, dauer larvae and adult hermaphrodites, we used high-pressure freezing and subsequent freeze substitution to immobilize and fixate animals in a near-to-native state. Using electron tomography on ~250 nm cross-sections of the worm, we obtained high resolution 3D images that allow for investigation of the ultrastructure of the nematodes NMJs. We used this information to compare synaptic vesicle pool characteristics, aiming for an automated approach for vesicle classification.

Upon visual examination of electron microscopic datasets of the NMJs of *C. elegans* dauer samples, we noted an increased fraction of DCV in dauer larvae compared to adult hermaphrodites. To systematically quantify this, vesicles were automatically extracted from tomograms, using an improved version of our 3D ART VeSElecT workflow (see S1 Fig), and then manually assigned for vesicle types by two experts. In uncertain cases, a third expert was consulted. Since closer examination of tomograms revealed that a definite assignment of some vesicles to one of the two classes (CCV and DCV) was difficult or even impossible, we established a third, hypothetic group of non-determinable (ND) vesicles. Fig 1 shows a representative selection of clear core, non-determinable and dense core vesicles from 8 exemplary tomograms (4 dauer larva tomograms and 4 hermaphrodite tomograms).

Fig 1. Comparison of CCV, ND vesicles and DCV of dauer larvae and adult hermaphrodites. A selection of representative pictures is shown. Pictures of the same developmental state (young adult hermaphrodite (H) / dauer larva (D)) and with identical number (I – IV) are from the same tomogram (e.g. CCV DI, ND DI and DCV DI originate from exactly the same tomogram). Edge length of micrographs = 100 nm.

Together, we included 1072 segmented vesicles from the 7 young adult hermaphrodite tomograms, and 1108 segmented vesicles from the 8 dauer larva tomograms for analyses (Table 1). When we included all identified vesicle populations from all tomograms, the percentage of DCVs differed slightly for dauer larvae (13%) and adult hermaphrodite (10%) datasets. After filtering the datasets, comparing only the vesicle population from the same type of NMJ with the bouton fully reconstituted in

161 the tomograms (putative cholinergic NMJs with central cut through the AZ area and at

162 least 100 vesicles), the dauer larvae NMJs exhibited a higher DCV ratio of 16%

163 versus 9% in hermaphrodites.

165 Table 1: number and percentage of vesicle types in all tomograms[1]

Developmental state	tomograms (n)	Amount of vesicles	CCV	DCV	NA	total
Adult hermaphrodite	7	Sum	949	109	14	1072
		Ratio	89%	10%	1%	
	5	Sum [1]	837	85	11	933
		Ratio [1]	90%	9%	1%	
dauer larvae	8	sum	948	148	12	1108
		ratio	86%	13%	1%	
	5	Sum [1]	669	130	3	802
		Ratio [1]	83%	16%	0%	

[1] analysis with a selected set of tomograms where only putative cholinergic NMJs

with apparent dense projections and with > 100 vesicles were included.

3.2. Characterization based on image features

In order to perform a quantitative comparison of vesicles and for clarification about the

ND vesicle class, we extracted a number of image features based on qualitative

observations. Since DCVs appear larger, darker and further apart from the active

zone compared to CCVs, we used radius (r), mean gray value (gv) and distance to

active zone (distAZ) as primary features (Fig 2). In addition, the standard deviation of

the gray value distribution of a vesicle (GVSD) is a measure for the contrast between

membrane and core that is independent of the brightness of the vesicle.

Fig 2. Visualization of vesicle pool in a dauer larva tomogram with color

coded image features: inner radius, gray value, distance to active zone

(distAZ), gray value standard deviation (GVSD). Clear core like

characteristics are yellow-shifted, dense core like features are blue shifted.

The active zone is located at the bottom.

Statistical analysis and comparison of ND vesicles with CCV and DCV suggests that

synapses show no third type of vesicles and that NDs are in most cases CCVs (see

S3 Fig). Hence, for further analysis, we proceeded with the assumption of two existing

vesicle classes, CCVs and DCVs (Table 1). We tried to apply simple thresholding and

linear combinations of the four features in order to separate DCVs from CCVs, but

with limited success (see S3 Fig). Therefore, a machine learning approach was

developed and applied, as described below. To generate training data, ND vesicles

were first tried to be concordantly assigned to one group by two experts. In difficult

cases, a third expert was consulted. Still non-assignable (NA) vesicles, were excluded

from training (see table 1: n = 12, ~ 1.1 % in dauer larvae; n = 14, ~1.3 % in adult

hermaphrodites).

3.3. Machine learning can distinguish vesicle types

To automatically classify vesicles using the previously calculated image features, we

evaluated different machine-learning algorithms. We compared a support vector

machine (SVM), random forest (RF,) and k nearest neighbor (KNN) for their

performance on differentiating between CCVs and DCVs using four image-based

characteristics described above: radius (r), mean gray value (gv), standard deviation

of the gray value (GVSD) and distance to the active zone (distAZ) (see Fig 3). 2180

manually labelled vesicles from 15 different tomograms (7 young adult hermaphrodite

and 8 dauer larval tomograms) were used as training data. The features were

calculated using the modified 3D ART VeSELecT workflow, saved as a table, and

processed in a python script using the scikit-learn machine learning library [29]. Table

2 shows the results of the comparison of the different algorithms.

Table 2. Average performance of different machine learning classifiers

classifier	accuracy	precision$_{DCV}$	recall$_{DCV}$	F-score$_{DCV}$
SVM	0.98 ± 0.02	0.95 ± 0.06	0.91 ± 0.1	0.93 ± O.7
Random forest (k = 10)	0.96 ± 0.03	0.88 ± 0.13	0.86 ± 0.13	0.86 ± 0.1
Random forest (k = 1500)	0.97 ± 0.02	0.93 ± 0.09	0.84 ± 0.13	0.88 ± 0.09
KNN	0.97 ± 0.02	0.93 ± 0.1	0.89 ± 0.11	0.90 ± 0.07

The comparison between SVM, random forest and KNN revealed that the SVM

algorithm performs best in our classification problem. The SVM shows the lowest error

rates in all four categories (see Methods; values in the following always given for

dense core vesicles): accuracy, precision$_{DCV}$, recall$_{DCV}$ and F-score$_{DCV}$.

KNN shows second best results with about 1-3 percent worse results compared to

SVM. In both random forest variants, k = 1500 outperformed k = 10 in most cases

except for recall$_{DCV}$. Random forest (k = 1500) shows overall third best results with

comparable values in accuracy and precision$_{DCV}$ as KNN, but 5 percent worse results

for recall$_{DCV}$ and about 2 percent worse results for the F-score$_{DCV}$. Resulting values for

the SVM hyperplane delimiting DCVs from CCVs are shown in Table 3.

Table 3. SVM and standardization parameters transferred to the Fiji macro

parameter	r	gv	distAZ	GVSD
mean	10.4	129.1	259.1	5.9
SD	3.1	4.7	118.5	1.7
weight	-1.69	1.65	-0.76	1.21

The final trained SVM classifier used the following weighting of the four features:

$-4.36 = -1.69*r + 1.65*gv - 0.76*distAZ + 1.21*GVSD$ (1)

In summary, the SVM algorithm in its linear form without kernelization outperformed

all other classification methods including non-linear SVM and random forest

(k = 1500) classification. Additionally, using a linear SVM has the advantage that the

resulting classifier can easily be incorporated into a classification workflow by

extracting the coefficients after successful training. For this purpose, a new vesicle

classification macro in ImageJ/Fiji was developed to be consecutively applied after the

modified 3D ART VeSElecT registration macro. The workflow of the classification

macro is shown in Fig 3 and is described in detail in Materials and Methods.

Fig 3. Workflow of the classification macro. 1) preparation: StackSegmented (registered vesicles) and StackDuplicateScale (scaled tomogram) from 3D ART VeSElecT [22] are used as input; the user is requested to mark the active zone via point-selection (semi-automated). 2) readout of features: gray value (gv), radius (r), vesicles coordinates and gray value standard deviation (GVSD) in the middle slice of vesicles are determined; the distance to active zone (distAZ) is calculated. 3)

processing of features: offsetting, standardization and weighting of features followed

by final assignment of labels. 4) output: includes a logfile and visualization of results

as new image stack windows.

3.4. Application to *C. elegans* dauer and adult hermaphrodite NMJs

We applied the automated vesicle classification workflow to quantify differences in the

vesicle pool in electron tomograms of *C. elegans* dauer and adult hermaphrodite

NMJs. Visual results of 3D ART VeSElecT and the classification macro for one

exemplary tomogram of each analyzed developmental state are presented in Fig 4.

Fig 4. Comparison of the visual results from the adapted 3D ART VeSElecT and the consecutive classification macro. *Top:* I) and II) original tomograms of young adult hermaphrodite (I) and dauer larva (II). *Bottom:* a) and c) show *Composit* stacks of 3D ART VeSElecT (random colors), whereas a') and c') show the resulting *Composit* after classification of vesicles (CCV = magenta, DCV = green). b) and d) show the 3D model of the vesicle pool, b') and d') show the 3D model with color labels showing the assigned class. All scale bars = 100nm.

253   The 3D models show that DCVs most often localize on the rim of the CCV cluster.

254   CCVs themselves gather around the dense projections, electron dense structures

255   attached to the cell membrane that indicate the active zone where synaptic vesicle

256   fusion occurs. For further analysis of vesicle characteristics (radius, gray value, GVSD

257   and distAZ), results are shown as violin plots in Fig 5.

Fig 5. Comparison of DCVs (black) characteristics with CCVs (light gray) characteristics. Violin plots show a) inner vesicle radius in nm, b) gray value (gv), c) distance to active zone (distAZ) and d) gray value standard deviation in central slice of each vesicle (GVSD). Panels b)-d) show normalized values of n_h = 7 tomograms of N2 young adult hermaphrodites; n_d = 8 tomograms of N2 dauer larvae.

Fig 5a shows highly significant differences in CCV and DCV size in both developmental states. In the following, the radius always refers to our original measured results, which is the "inner radius" of all vesicles, as described in [22]. To make vesicle size comparable to other published results, the measured radius must be multiplied by two to receive the inner diameter and increased by 9 nm to include vesicle membrane thickness [22]. This "real diameter" is only an approximation and cannot be directly compared to manually measured data [22].

CCVs of dauer larvae show a median inner radius of 9.0 nm, whereas DCVs have a median inner radius of 14.7 nm, which results in a significant difference of 5.7 nm. However young adult hermaphrodites' median inner radius of DCVs is 16.7 nm, hence they have a ~ 5.4 nm bigger radius than CCVs with a median inner radius of 10.3 nm. Noticeably, both vesicle types are slightly but significantly bigger in hermaphrodites compared to dauer larvae. The radius of hermaphrodite CCVs is 1.3 nm larger compared to

dauer larvae, whereas the radius of hermaphrodite DCVs is 2 nm larger
compared to dauer larvae.

In all cases, DCVs have significantly darker normalized gray value intensities
in comparison to CCVs, as shown in Fig 5b. Hermaphrodites show a
normalized gv of 0.17 for CCVs and 0.45 for DCVs, whereas dauer larvae
have a normalized gv of 0.23 for CCVs and 0.56 for DCVs. Hence, vesicles of
dauer larvae tomograms possess significantly darker vesicles (both CCVs
and DCVs) in comparison to hermaphrodite tomograms.

In contrast, GVSD (gray value standard deviation in the central slice of each
vesicle) does not differ significantly between developmental states, but shows
highly significant differences of ~ 0.22 in GVSD values of CCV to DCV for
both developmental states (Fig 6d).

The normalized distance of CCVs to the active zone distAZ is smaller (0.42
for hermaphrodites and 0.4 for dauer larvae) than that of DCVs
(hermaphrodites: 0.64, dauer larvae: 0.72) in both larval stages (Fig 6c). The
normalized distances of DCVs to the AZ, of hermaphrodites in comparison to
dauer larvae, show a highly significant difference of 0.08 (normalized). For
better visualization of this observation, distances are visualized as histogram
plots in Fig 6.

Fig 6. Cumulative histogram (a, b) and kernel density estimation plots (c, d; bandwidth = 0.07*maximum value of total dataset) of distances to the active zone (distAZ). Given are normalized distances (left) and real values in nm (right). CCVs are shown as dashed line and DCVs as continuous line. Adult hermaphrodite vesicles are presented in shades of gray, whereas dauer larval vesicles are shown in violet.

Fig 6a and c illustrate that normalized distances of CCVs of both developmental states are almost identical, whereas distances of DCVs differ significantly in both plots. DCVs show a systematic discrepancy in their localization: DCVs in hermaphrodites cluster closer to the active zone than

DCVs of dauer larvae. Histograms in Fig 6b and d shows results of distAZ in

real distance measured in nm. Interestingly, this statistic shows that vesicle

pools in dauer larvae have a generally smaller maximal distAZ in adult

hermaphrodites. The largest distance of dauer larvae vesicles from the active

zone is about 550 nm whereas the largest distance of vesicles in young adult

hermaphrodites is about 700 nm.

3.5. Error rates and cross validation

To generate gold standard data, a consensus label was created (N = 15 tomograms, 8

dauer larva and 7 hermaphrodite tomograms): First, labels were assigned

independently by two biologists. Differing labels were re-examined. In case of no

agreement, a third examiner was employed. Still non assignable vesicles (~1 %) were

excluded from SVM training. The initial assignment by individuals was compared with

this consensus label to determine the manual error rate (Table 4).

Table 4 manual vs classification macro (SVM performance)

	$precision_{DCV}$	$recall_{DCV}$	$F\text{-}score_{DCV}$
Manual [1]	0.98 ± 0.05	0.96 ± 0.08	0.97 ± 0.05
SVM [2]	0.95 ± 0.07	0.90 ± 0.10	0.92 ±0.07

[1] Compares two independent manual assignments to the consensus label

(ground truth)

[2] Compares the SVM assignment to the ground truth

Manual annotation shows higher scores than the SVM. Manual assignment of

vesicle types shows 98 % $precision_{DCV}$, whereas the SVM reaches 95 % $precision_{DCV}$.

The results for recall$_{DCV}$ is 96 % for manual assignment and 90 % for the SVM. The F-

score is the harmonic mean of precision and recall (see Materials and Methods) and

shows a mean result for the error rate of 97% for manual and 92% for automated

classification. Results for precision and recall are additionally shown as scatterplot in

Fig 7.

Fig 7. SVM error rates as precision$_{DCV}$ and recall$_{DCV}$ plot. Each data point

represents one validation tomogram. Red squares: adult hermaphrodite, blue circles:

dauer larvae. Right: magnified detail plot. Red star indicates mean of hermaphrodite

data: precision$_{DCV}$ = 0.95, recall$_{DCV}$ = 0.85. Blue cross indicates mean of dauer data:

precision$_{DCV}$ = 0.96, recall$_{DCV}$ = 0.95. 3 dauer tomograms and 1 hermaphrodite

tomogram were classified without error.

The SVM classifier was cross-validated against each tomogram using all other

tomograms as training data. Fig 7 shows error rates of cross validation of automated

SVM classification. Hermaphrodite tomograms are represented as red squares and

dauer larvae tomograms are represented as blue circles. Mean classification results

are shown as red star for hermaphrodites ($precision_{DCV}$ = 0.95, $recall_{DCV}$ = 0.85) and

as blue cross for dauer larvae tomograms ($precision_{DCV}$ = 0.96, $recall_{DCV}$ = 0.95).

Dauer larvae tomograms have in average better results than hermaphrodites and

cluster in the right upper corner. 3 dauer tomograms and one hermaphrodite

tomogram were classified without any error and overlay in the right upper corner at

$precision_{DCV}$ = 1.0, $recall_{DCV}$ = 1.0.

5. Discussion

Automated methods for image analysis are of great importance to reduce manual labor in the long term and prevent biased analysis of data, hence making double blind analysis unnecessary. This is confirmed by our own experience, where one re-analysis of the same tomogram by the same expert revealed one differential interpretation of CCV and DCV and 6 other non-matching assignments (like CCV and ND or CCV and error) on n= 134 vesicles in total. Such deviations in manual analysis may have a great influence on the outcome. For this reason, we had two experts manually assign all vesicles independently. In case of inconsistency, both persons came to a mutual agreement or consulted a third expert in the decision process. Still unassignable vesicles were excluded from analysis. Since this evaluation process is very elaborate and time-consuming, a long-term automated solution is inevitable. As far as we know there is no other approach for the specific task of automated vesicle classification in electron tomograms. We present here a new methodological approach based on a modified version of the previously published Fiji macro 3D ART VeSElecT [22]. This tool was applied for automated recognition and read-out of important characteristics which are used to train a machine learning classifier. Subsequently, it was used to visualize results through color labelling of classified particles in 3D and can be used for automated vesicle classification in the future. We apply our approach to compare two different developmental states of *C. elegans*, young adult hermaphrodites and dauer larvae.

Manual analysis revealed a hypothetical third type of vesicles, called non-determinable vesicles (ND) which are most likely CCVs with no perfectly clear inner

core. This seems reasonable, since they resemble CCV characteristics in radius and distance to the active zone and only differ significantly in their gray value, which lies among gv intensities of DCVs and CCVs (S3 Fig). This observation can be explained by either biological or technical reasons. On the one hand, medium-gray vesicles could be DCVs with partially released cargo, e.g. after kiss-and-run [30–32]. Overall this does not explain why NDs have the same size as CCVs, since DCVs are known to be considerably larger [23]. On the other hand, vesicles may appear darker in electron micrographs because of lower quality of tomograms at some locations. Lacking precision of the tomogram overlay of the two single tilted tomograms to receive one double tilted tomogram could result in darker, more blurry vesicles with no perfectly clear core.

The comparison of DCV number in dauer larvae and adult hermaphrodites revealed differences in the ratio of DCVs (9 % in adult hermaphrodites and 16 % in dauer larvae), when comparing only cholinergic synapses with apparent dense projections and at least 100 vesicles in the NMJ. Furthermore we found that DCV ratio correlates with cell size change throughout the tomogram (S2 Fig). A large cell size change indicates inclusion of decentral regions of the synaptic bouton in the tomogram. We assume that inclusion of more remote areas in the tomogram is more likely in dauer larvae, since the overall size of the synaptic bouton is smaller (mean diameter ± SD of cross section: 452 ± 85 nm) compared to adult hermaphrodites (mean diameter ± SD of cross section: 542 ± 97 nm). Therefore standard sized 250 nm sections, typically prepared for electron tomography, naturally have a higher chance of catching decentral regions in dauer larvae NMJs. Regardless of the origin of differences in DCV ratio, we show that our approach can automatically quantify

such differences. For a more comprehensive understanding, serial section

tomography for visualization of complete synaptic boutons would be necessary.

Additionally, we found that hermaphrodites show significantly larger vesicles (CCVs

and DCVs) and diameters of synapses in comparison to dauer larvae. This

observation could either be a biological phenomenon of dauer larvae or a side effect

of the sample preparation. Dauer larvae have a distinct cuticle that shields the animal

from adverse environmental conditions, like heat and drought. This could lead to

adverse effects during fixation (freeze substitution) of the sample, even though we did

not observe any sign of deficient sample fixation in dauer larvae. Divergent synaptic

vesicle sizes have already been found in NMJs of other organisms, like rats [33] and

zebrafish [23]. For zebrafish a dependency of vesicle size on the maturation stage of

NMJs was assumed that might be related to the functional transition of these

synapses to control different swimming behavior [23]. Differences in synaptic vesicle

sizes during normal development have not been observed for *C. elegans*, to our

knowledge. However, divergent vesicle sizes could also be related to different moving

behavior of dauer larvae that are well known for their unique nictation behavior [9].

Since manual measurements are often inaccurate, smaller deviations in vesicle sizes

are only observable in extensive analysis, therefore automated approaches are

inevitable. Our previously published macro 3D ART VeSElecT proved to be

successful for distinguishing vesicle size differences [22]. This approach in

combination with our here presented extended version for automated classification

may be the key to reveal so far undistinguishable differences in the future.

Interestingly, although nervous cells in dauer larvae are smaller, segregation of

DCVs and CCVs is more distinct. Our interpretation is that DCVs in dauer larvae

locate further away from the active zone on the rim of the synaptic vesicle pool

compared to adult hermaphrodites. Differences in this characteristic of synaptic

architecture could be explained by a shift towards a higher ratio of DCVs in a non-

fusion competent resting state in dauer compared to adult hermaphrodites. Moreover,

the location of DCVs closer to the cell margin in dauer compared to adult

hermaphrodites could indicate the need for faster peptide release, and their increased

number could compensate for lower volume.

For automated classification of vesicles, different machine learning algorithms

were tested. The SVM showed best results in comparison to random forest or k-

nearest neighbor classification. Furthermore, the SVM works regardless of

dimensionality, can classify non-linearly separable classes by using the kernel trick,

enables an easy integration of the trained linear classifier into the macro and generally

can be used for assignment into more than two classes [34].

Cross-validation showed very good results for automated classification of

vesicles. Interestingly, dauer larval tomograms showed better results than

hermaphrodite tomograms, which could be related to the differing gray value

intensities of the tomograms. Hence high contrast vesicles were easier to

automatically recognize and showed more distinct results for GVSD and gv. Another

reason could be the more pronounced positional segregation of the two vesicle types

for dauer larvae.

We suggest that our pre-trained SVM can be used for DCV and CCV classification in

other, similar tomograms, and provide the Fiji classification macro including the

trained parameters. In addition, users can train a SVM on their own data using the

python script we provide, and integrate the parameters into the Fiji macro.

Our ImageJ classification macro and the python script, for retraining of the

classifier, a tutorial and test datasets are available at: https://www.biozentrum.uni-

wuerzburg.de/bioinfo/computing/3dart-veselect/.

6.   Conclusion

We developed a new automated approach to classify vesicles and to quantify their properties from electron tomograms. We combined machine learning with an extension of our previously developed vesicle segmentation workflow 3D ART VeSElecT to reliably distinguish clear core from dense core vesicles using image-based features. We apply this method to electron tomograms in *C. elegans* NMJs and the task to separate clear and dense core vesicles. We find an increased fraction of dense core vesicles (~ 16 % vs ~ 9 %) of tomograms matching certain characteristics (cholinergic synapses with > 100 vesicles of a central slice through the AZ area), as well as significantly reduced vesicle size and a higher distance of dense core vesicles to the active zone in dauer larvae compared to adult hermaphrodites. Our approach is not limited to *C. elegans* and can be applied to study differences in synaptic and other vesicles in electron tomograms in various settings and many other model organisms, e.g. in *Danio rerio* [22].

7. Material and Methods

6.1 Electron Microscopy

1. Animals

C. elegans wild-type (Bristol N2) were used and maintained using standard methods [4] . For the experiments, young adult hermaphrodite worms as well as dauer larvae were used.

2. Purification of dauer larvae

For segregation of dauer larvae from other states of the *C. elegans* life cycle, treatment with sodium dodecyl sulfate (SDS) buffer was used [35]. Firstly, worms were washed from the agar plates using M9 medium [36]. The worm solution was then transferred into 50 ml tubes and centrifuged for 5 min at 2000g at 4°C. Afterwards, the supernatant was removed and the pellet was dissolved in 50 ml 1 % SDS solution and incubated on a shaker for 15 min. SDS was then removed by centrifuging (2000g, 5 min, 4°C) and subsequent discarding of the supernatant while retaining 5 ml of solution. The tube was then refilled with distilled water and the procedure repeated 2-3 times. For separation of dauer larvae from dead material, the solution was pipetted into a pit on an unseeded agar plate. Once the solution had dried, living dauer larvae were able to crawl away from the pit. After 1-3 hours, emigrated worms were washed off with M9 medium.

3. High-pressure freezing, freeze-substitution and electron tomography

HPF, FS and electron tomography were performed as previously described [22]. Tilt

series were recorded from at least +65° to -65° but not exceeding +70° to -70°.

6.2 Improvements to 3D ART VeSElecT

We extended our tool 3D ART VeSElecT to calculate various image features of

detected vesicles to be used for classification. The workflow is shown in

Supplementary Figure S1, with changes compared to the previous version highlighted

by a black box. The overall workflow of 3D ART VeSElecT remained as previously

published [22]. One change in the preprocessing step is the determination of the

mean radius instead of volume, since the radius or diameter is typically used when

comparing vesicle size. Furthermore, automated calculation of the cell volume was

added. Since semi-automated selection of the cell outline by the user was already

included in the macro, this selection is now basis for calculating the analyzed volume

in 3D (which corresponds to cell volume if the selection is taken along the cell

membrane, as described in the 3D ART VeSElecT tutorial).

6.3 Classification Macro

The workflow of the new Fiji macro for classification of DCV and CCV in electron

tomograms of synapses comprises four main steps: 1) preparation, 2) reading out

features, 3) processing of features followed by labelling of vesicles, and 4) output. In

the preparation step, StackDuplicateScale and StackSegmented, which are results

from the 3D ART VeSElecT recognition macro, are read into the classification macro.

First of all, the macro opens a preparation window for changing the names of read-in

stacks (StackDuplicateScale and StackSegmented) and setting parameters for

standardization and adaption of SVM weights. In case of adaptation of the SVM

algorithm to another user's needs, values can easily be changed through the input

window. In a second step, four characterizing vesicle features are computed from the

images: radius, gv, GVSD and distAZ. GVSD is a value for standard deviation of gray

values in the central slice of each vesicle, since CCVs have a higher gv variance than

DCV. In the third step, features are processed, which includes applying an offset to gv

(if the darkest vesicle is lighter than a threshold), standardization of parameters,

weighting features based on SVM weights, and finally the assignment of labels

according to the identified class (CCV = magenta, DCV = green). In the last step, the

output is created, including a logfile and visual output shown in new windows. The

logfile includes all measured characteristics (e.g. features, classification labels (CCV =

1, DCV = 0), number of vesicles in each class). The visual output shows, besides the

input stacks (registered vesicles as "StackSegmented" and scaled, original tomogram

as "StackDuplicateScale"), an additional image stack, named "label" which

corresponds to the StackSegmented but displays color-coded vesicles according to

their classification. Furthermore a Composit stack, named "CompositLabel" (overlay of

original tomogram and the "label" stack) is created.

6.4 Comparison of different types of machine learning classifiers

We compared a support vector machine (SVM), random forest (RF, using k = 10, k =

1500; k indicates here the leaf parameter) and k nearest neighbor (KNN) for their

performance on differentiating between CCVs and DCVs based on four image-based

characteristics: radius (r), mean gray value (gv), standard deviation of the gray value

(GVSD) and distance to the active zone (distAZ).

Linear SVM (C = 1) showed the best average classification results (see Table 2).

Parameters (Table 3) were determined using all input tomograms as training data

and transferred to a Fiji-Macro (Figure 3) to enable classification downstream of

the 3D ART VeSElecT registration macro.

6.5 Support Vector Machine (SVM)

A linear support-vector machine (SVM) divides two classes by a hyperplane. As

such, the SVM is a binary classifier dealing with linear classification problems. A

one-versus-all approach and kernelization can extend classification to non-linear

problems and into more than two classes, but will not be discussed here since the

classification power of the SVM is sufficiently strong, see results. With the aim of

minimizing generalization error, the optimization objective of the SVM is to

maximize the margin between the hyperplane and the closest data points of the

two classes. Those eponymous support-vectors lie on the positive and negative

hyperplane, which are equidistant to the separating hyperplane [37]. As the linear

SVM is parametric, the equation of the separating hyperplane can be transferred

independently of the training-data (see Results, Table 3 and Equation 1), with x_i

being the features and ω_i the assigned feature-weights. All data points with

$\sum_{i=1}^{n} w_i x_i'' > 0$ are assigned to class 1 (CCV), whereas all data points with

$\sum_{i=1}^{n} w_i x_i'' < 0$ are assigned to class -1 (DCV).

6.6 Implementation of machine learning algorithms

A python 3 script (Supplementary Figure S2) reads in the CSV files containing

the training datasets and applies classifiers from the scikit-learn library [29]. The

manual labels are transformed into a feature array: Particles marked "D" (DCV)

are relabeled "-1". Particles marked "E" (error) are removed. All other markings

are relabeled "1", including "C" (CCV) and "N" (not determinable), since non-

determinable particles show more similarities to CCV (S1 Fig).

Classifiers are evaluated by cross-validation: Data from one CSV file are used as

validation data, all other files are concatenated and used as training data (there

are as many training-validation combinations as input CSV files). For each

combination, features are standardized using mean and standard deviation (sd)

of the training data and passed to the classifiers, namely linear SVM, random

forest and KNN. An evaluation function determines accuracy of all vesicles and

precision, recall and F-score of DCV only (see equations (2) - (4)). Evaluation

results are returned as pandas.DataFrame [38] and as CSV file.

Fig8. Format of input CSV file. First four columns are the four features, the

fifth column is the manual labels. The python script rewrites manual labels

into a binary label-array: "D" (DCV) is rewritten to -1, "E" (error) is removed,

and everything else is rewritten as "1" (CCV).

Equations:

$$precision_{DCV} = \frac{tp_{DCV}}{tp_{DCV}+fp_{DCV}} \qquad (2)$$

$$recall_{DCV} = \frac{tp_{DCV}}{tp_{DCV}+fn_{DCV}} \qquad (3)$$

$$F_{DCV} = 2 * \frac{DCV_{precision} * DCV_{recall}}{DCV_{precision}+DCV_{recall}} \qquad (4)$$

tp: true positive; fp: false positive; fn: false negative.

9.   References

1.   Salio C, Lossi L, Ferrini F, Merighi A. Neuropeptides as synaptic transmitters. Cell Tissue Res.
Springer-Verlag; 2006;326: 583–598.

2.   Li C. Neuropeptides. WormBook. 2008; 1–36.

3.   Zhai RG, Vardinon-Friedman H, Cases-Langhoff C, Becker B, Gundelfinger ED, Ziv NE, et al.
Assembling the presynaptic active zone: a characterization of an active one precursor vesicle.
Neuron. 2001;29: 131–43.

4.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. Genetics Society of America;
1974;77: 71–94.

5.   White JG, Southgate E, Thomson JN, Brenner S. The Structure of the Nervous System of the
Nematode Caenorhabditis elegans. Phil Trans R Soc Lond B. 1986;314 1165: 1–340.

6.   Varshney LR, Chen BL, Paniagua E, Hall DH, Chklovskii DB. Structural Properties of the
Caenorhabditis elegans Neuronal Network. PLoS Comput Biol. 2011;7: 1–21.

7.   Haueis P, Slaby J. Connectomes as constitutively epistemic objects: Critical perspectives on
modeling in current neuroanatomy. Progress in brain research. 2017. pp. 149–177.

8.   Golden JW, Riddle DL. The Caenorhabditis elegans dauer larva: Developmental effects of
pheromone, food, and temperature. Dev Biol. Academic Press; 1984;102: 368–378.

9.   Cassada RC, Russel RL. The dauerlarva, a post-embryonic developmental variant of the
nematode Caenorhabditis elegans. Dev Biol. Academic Press; 1975;46: 326–342.

10.   Erkut C, Kurzchalia T V. The C. elegans dauer larva as a paradigm to study metabolic
suppression and desiccation tolerance. Planta. Springer Berlin Heidelberg; 2015;242: 389–396.

11.   Pierce SB, Costa M, Wisotzkey R, Devadhar S, Homburger SA, Buchman AR, et al. Regulation
of DAF-2 receptor signaling by human insulin and ins-1, a member of the unusually large and
diverse C. elegans insulin gene family. Genes Dev. Cold Spring Harbor Laboratory Press;
2001;15: 672–86.

12.   Hung WL, Wang Y, Chitturi J, Zhen M. A Caenorhabditis elegans developmental decision
requires insulin signaling-mediated neuron-intestine communication. Development. Oxford
University Press for The Company of Biologists Limited; 2014;141: 1767–79.

13. Li W, Kennedy SG, Ruvkun G. daf-28 encodes a C. elegans insulin superfamily member that is
regulated by environmental cues and acts in the DAF-2 signaling pathway. Genes Dev. Cold
Spring Harbor Laboratory Press; 2003;17: 844–58.

14. Cornils A, Gloeck M, Chen Z, Zhang Y, Alcedo J. Specific insulin-like peptides encode sensory
information to regulate distinct developmental processes. Development. Oxford University Press
for The Company of Biologists Limited; 2011;138: 1183–93.

15. Ren P, Lim CS, Johnsen R, Albert PS, Pilgrim D, Riddle DL. Control of C. elegans larval
development by neuronal expression of a TGF-β homolog. Science. 1996;274: 1389–1391.

16. Schackwitz WS, Inoue T, Thomas JH. Chemosensory Neurons Function in Parallel to Mediate a
Pheromone Response in C. elegans. Neuron. Cell Press; 1996;17: 719–728.

17. Salio C, Averill S, Priestley JV, Merighi A. Costorage of BDNF and neuropeptides within
individual dense-core vesicles in central and peripheral neurons. Dev Neurobiol. Wiley
Subscription Services, Inc., A Wiley Company; 2007;67: 326–338.

18. Walch-Solimena C, Takei K, Marek KL, Midyett K, Südhof TC, De Camilli P, et al.
Synaptotagmin: a membrane constituent of neuropeptide-containing large dense-core vesicles.
J Neurosci. Society for Neuroscience; 1993;13: 3895–903.

19. Stigloher C, Zhan H, Zhen M, Richmond J, Bessereau J-L. The presynaptic dense projection of
the Caenorhabditis elegans cholinergic neuromuscular junction localizes synaptic vesicles at the
active zone through SYD-2/liprin and UNC-10/RIM-dependent interactions. J Neurosci. 2011;31:
4388–96.

20. Kittelmann M, Hegermann J, Goncharov A, Taru H, Ellisman MH, Richmond JE, et al. Liprin-
α/SYD-2 determines the size of dense projections in presynaptic active zones in C. elegans. J
Cell Biol. 2013;203: 849–863.

21. Lim MA, Chitturi J, Laskova V, Meng J, Findeis D, Wiekenberg A, et al. Neuroendocrine
modulation sustains the C. elegans forward motor state. Elife. 2016;5.

22. Kaltdorf KV, Schulze K, Helmprobst F, Kollmannsberger P, Dandekar T, Stigloher C. FIJI Macro
3D ART VeSElecT: 3D Automated Reconstruction Tool for Vesicle Structures of Electron
Tomograms. Dyer E, editor. PLOS Comput Biol. Public Library of Science; 2017;13: e1005317.

23. Helmprobst F, Frank M, Stigloher C. Presynaptic architecture of the larval zebrafish

639    neuromuscular junction. J Comp Neurol. 2015; 523;13: 1984-97

640 24. Phillips MI, M.Ian. Neuropeptides: Regulators of Physiological Processes. Regul Pept. Elsevier;

641    2000;90: 101.

642 25. Mastronarde DN. Dual-axis tomography: an approach with alignment methods that preserve

643    resolution. J Struct Biol. 1997;120: 343–52.

644 26. Zampighi GA, Fain N, Zampighi LM, Cantele F, Lanzavecchia S, Wright EM. Conical Electron

645    Tomography of a Chemical Synapse: Polyhedral Cages Dock Vesicles to the Active Zone. J

646    Neurosci. 2008;28.

647 27. Zampighi GA, Zampighi L, Fain N, Wright EM, Cantele F, Lanzavecchia S. Conical tomography

648    II: A method for the study of cellular organelles in thin sections. J Struct Biol. 2005;151: 263–

649    274.

650 28. Robertson JD. The Early Days of Electron Microscopy of Nerve Tissue and Membranes. Int Rev

651    Cytol. 1987;100: 129–201.

652 29. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:

653    Machine Learning in Python. J Mach Learn Res. 2012;12: 2825–2830.

654 30. Artalejo CR, Elhamdani A, Palfrey HC. Secretion: Dense-core vesicles can kiss-and-run too.

655    Curr Biol. Cell Press; 1998;8: R62–R65.

656 31. Rutter GA, Tsuboi T. Kiss and run exocytosis of dense core secretory vesicles. Neuroreport.

657    2004;15: 79–81.

658 32. Tsuboi T, McMahon HT, Rutter GA. Mechanisms of dense core vesicle recapture following "kiss

659    and run" ("Cavicapture") exocytosis in insulin-secreting cells. J Biol Chem. American Society for

660    Biochemistry and Molecular Biology; 2004;279: 47115–24.

661 33. Markus EJ, Petit TL, LeBoutillier JC. Synaptic structural changes during development and aging.

662    Brain Res. 1987;432: 239–48.

663 34. Shawe-Taylor J, Cristianini N. Kernel Methods for Pattern Analysis [Internet]. Elements. 2004.

664 35. Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant of the

665    nematode Caenorhabditis elegans. Dev Biol. 1975;46: 326–342.

666 36. Hope I. C. elegans a practical approach. Oxford Univ Press. 1999;

667 37. Raschka S. Python Machine Learning. Packt Publishing; 2015.

668   38.   McKinney W. Data Structures for Statistical Computing in Python. Proc 9th Python Sci Conf.

669   2010;1697900: 51–56.

10. Supplementary Information

S1 Figure: Improvements to the original vesicle registration workflow of 3D ART

VeSElecT

S1 Fig. Changes in the workflow of the 3D ART VeSElecT registration macro:

The figure shows the workflow of the previously published 3D ART VeSElecT

macro [22], with changes framed in black. In the yellow box only one small

change was made: instead of volume, the inner radius is now used for size

determination.

Supplementary S2 Figure: Schematic representation of the python-script used for

training the classifier

S2 Fig. Schematic representation of the python-script that trains and

cross-validates SVM, KNN and random forest. Functions are written bold,

descriptions of arguments italic. Horizontal, light gray arrow indicates

function calls, vertical black arrows passing of arguments.

 S3 Figure: Vesicle classification by thresholding a combination of features

d)

Distribution CCV + ND dauer [%]			
	< ◯	> ◯	
> □	5 ± 4	35 ± 22	39 ± 24
< □	7 ± 4	54 ± 21	61 ± 24
	11 ± 7	89 ± 7	

Distribution CCV + ND hermaphrodite [%]			
	< ◯	> ◯	
> □	26 ± 31	55 ± 25	81 ± 14
< □	4 ± 3	15 ± 16	19 ± 14
	29 ± 33	71 ± 33	

S3 Fig. Statistical analysis of CCV and DCV features. a + b) boxplots of inner radius and normalized gray value, distance to active zone and DCF of dauer larva and adult hermaphrodites. Significance levels were determined via Mann-Whitney-U: p-values > 0.05 = not significant (n.s.), < 0.05 = *, <

0.01 = **, < 0.001 = ***. c) vesicles standard deviation of gray value vs DCF.

Exemplary plots of one dauer (N2 stack01 bp) and one adult hermaphrodite

(F1 s1 vnc) tomogram. Marked in hermaphrodite tomogram: DCV with

highest GVSD (black square) and DCV with highest DCF (dense core factor,

black circle) and corresponding quadrant thresholds (black dotted lines). d)

Distribution of CCV and ND within the quadrants of all examined tomograms.

Thresholds were independently determined for each tomogram. Green

shaded quadrant (lower left) contains all DCV.

S1 Text: Vesicle classification by thresholding a combination of features

Features used for manual distinction between CCVs and DCVs were

obtained using the Fiji macro 3D ART VeSElecT. S3 Fig. shows boxplots

comparing gv, distAZ, radius and dense core factor (DCF) of CCVs, NDs and

DCVs. The DCF is a linear weighted combination of the normalized values for

distAZ, radius (r) and gv:

$$DCF = distAZ_{norm} * 0.25 + r_{norm} * 0.25 + gv_{norm} * 0.5$$

S3a Fig. shows highly significant differences of gv between all three types of

vesicles (CCVs, DCVs and NDs) for both developmental states. CCVs and

NDs of dauer larvae are significantly brighter than of hermaphrodites (p-value

< 0.001). Comparing dauer larvae and hermaphrodites, DCVs show no

significant difference in gv. CCVs and NDs show no (dauer larvae, p-value >

0.05) or only slight differences (hermaphrodite, p-value <0.05) in their distAZ.

DCVs and CCVs differ significantly (p-value < 0.01) in their distAZ in both

states. DCVs and NDs show obvious differences in dauer larvae and

significant differences in hermaphrodites. DCVs in dauer larvae have a

significantly bigger distAZ than DCVs in hermaphrodites. Whereas CCVs and

NDs in both developmental states are about the same size, DCVs show

clearly higher values for radius. Young adult hermaphrodite's vesicles are

generally bigger in size compared to dauer larvae vesicles. The DCF shows

that NDs and CCVs in dauer larvae differ only slightly. In young adult

hermaphrodites, all vesicle types show strong differences in DCF values.

Furthermore NDs differ highly significantly between developmental states,

whereas CCVs and DCVs show no significant disparity.

S3c Fig. shows results of one exemplary tomogram of dauer and young adult

hermaphrodites. Here, values of GVSD are plotted against the DCF. The

DCV with highest GVSD is marked with a black square, whereas DCV with

highest DCF is marked with a black circle. Using these values, corresponding

thresholds to separate the four quadrants are determined. All DCVs locate in

the lower left quadrant. Resulting percentages of CCVs and NDs are shown

in in S3d Fig. For dauer larvae, ~ 7 %, for adult hermaphrodites ~ 4 % CCVs

and NDs remain in in the lower left quadrant. This shows that GVSD and

DCF are important characteristics for DCVs separation, but setting two

thresholds is not sufficient for reliable partitioning. For that reason, we aimed

at using a machine learning classifier to obtain a better separation using gv,

radius, distAZ and GVSD.

S1 Table: CCV, NA and DCV ratio for each tomogram and measured

synaptic cell size in x and y direction

Hermaphrodit	%CCV	%NA	%DCV	x1	x2	y1	y2	x difference	y difference	SUM differenc
H_1_Tomo	84,78%	0,00%	15,22%	412	594	478	605	182	127	309
H_2_Tomo	89,34%	1,84%	8,82%	632	634	603	588	2	15	17
H_3_Tomo	93,15%	0,00%	6,85%	411	361	597	599	50	2	52
H_4_Tomo	88,06%	0,37%	11,57%	759	694	657	605	65	52	117
H_5_Tomo	87,22%	3,01%	9,77%	494	465	459	439	29	20	49
H_6_Tomo	92,98%	0,88%	6,14%	509	506	596	564	3	32	35
H_7_Tomo	72,34%	6,38%	21,28%	557	383	483	507	174	24	198
mean	86,84%	1,78%	11,38%							
sd	7,07%	2,30%	5,32%							
mean selected	90,15%	1,22%	8,63%							
sd selected	2,77%	1,21%	2,20%							
Dauer (8 tomograms)										
D_1_Tomo	90,67%	4,00%	5,33%	482	462	380	324	20	56	76
D_2_Tomo	79,23%	0,00%	20,77%	422	336	437	368	86	69	155
D_3_Tomo	88,59%	0,00%	11,41%	350	397	453	497	47	44	91
D_4_Tomo	90,57%	0,00%	9,43%	334	402	446	498	68	52	120
D_5_Tomo	79,19%	0,00%	20,81%	510	419	638	533	91	105	196
D_6_Tomo	80,10%	1,57%	18,32%	530	622	463	546	92	83	175
D_7_Tomo	87,88%	3,03%	9,09%	352	335	406	388	17	18	35
D_8_Tomo	93,94%	2,27%	3,79%	532	519	552	551	13	1	14
mean	86,27%	1,36%	12,37%					dorsal nerve cord		
sd	5,88%	1,61%	6,77%					< 100 vesicles		
mean selected	83,54%	0,31%	16,15%					decentral & < 100 vesicles		
sd selected	5,57%	0,70%	5,37%					GABAergic		

S1 Table shows an overview of the final results for each tomogram. All

selected tomograms (> 100 vesicles, cholinergic, central cut) are marked in

yellow. Other tomograms are colored after exclusion criterion in light gray (<

100 vesicles), dark gray (decentral cut plus < 100 vesicles) or blue

(GABAergic NMJ).

Since it is known that DCVs appear in higher concentration at the rim of the AZ [19],

we compared the cell size of each NMJ in x and y direction in the first and last picture

of each image stack. To receive the overall cell size change we added both

determined differences up. This measurements will give us a clue of how central the

cut through the NMJ really is. For a very central cut we expect a very low cell size

change rate, whereas for more remote cuts through the AZ (although they

demonstrably include dense projections) we expect a higher change of the cell size

since synapses tend to narrow at regions far-off the AZ. In S2 Fig we plotted the

determined cell size change against the DCV ratio for each tomogram.

S4 Fig. Scatterplot of cell size change [nm] compared to the DCV ratio.
Cell size change was measured for each NMJ in x and y direction in the first
and last picture of the tomogram. Tomograms of adult hermaphrodites are
plotted in dark red. Results of dauer larvae are plotted in teal green.
Tomograms of the selected set (see Table 1, only putative cholinergic
synapses with > 100 vesicles and apparent dense projections) are visualized
as circles, NMJs with divergent characteristics are shown as different
symbols (see legend).

S4 Fig shows that the number of DCVs included in a cell seems to be

proportional to the cell size change. As an example, 3 of the dauer larvae

tomograms show a DCV ratio of 18-21 % but also have a high change of cell

size. Hence, although apparent dense projections can be seen in the

tomograms, the sample also includes decentral regions, were higher DCV

abundance is expected.

770