Automated classification of synaptic vesicles in electron tomograms of *C. elegans* using machine learning

Kristin Verena Kaltdorf[1,2,3], Maria Theiss[2,3], Sebastian Matthias Markert[1], Mei

Zhen[4], Thomas Dandekar[2,&], Christian Stigloher[1,&], Philip Kollmannsberger[3,&]

[&] TD, CS and PK are Joint Senior Authors.

[1]Imaging Core Facility, Biocenter, University of Würzburg, 97074 Würzburg, Germany

[2]Department of Bioinformatics, Biocenter, University of Würzburg, 97074 Würzburg,
Germany

[3]Center for Computational and Theoretical Biology, Hubland North, building 32,
University of Würzburg, 97074 Würzburg, Germany

[4]Department of Molecular Genetics, University of Toronto, Toronto, M5G 1X5, Canada

Correspondence to Philip.Kollmannsberger@uni-wuerzburg.de.

1.   Abstract

Synaptic vesicles (SVs) are a key component of neuronal signaling and fulfil different roles depending on their composition. In electron micrograms of neurites, two types of vesicles can be distinguished by morphological criteria, the classical "clear core" vesicles (CCV) and the typically larger "dense core" vesicles (DCV), with differences in electron density due to their diverse cargos. Compared to CCVs, the precise function of DCVs is less defined. DCVs are known to store neuropeptides, which function as neuronal messengers and modulators [1]. In *C. elegans*, they play a role in locomotion, dauer formation, egg-laying, and mechano- and chemosensation [2]. Another type of DCVs, also referred to as granulated vesicles, are known to transport Bassoon, Piccolo and further constituents of the presynaptic density in the center of the active zone (AZ), and therefore are important for synaptogenesis [3].

To better understand the role of different types of SVs, we present here a new automated approach to classify vesicles. We combine machine learning with an extension of our previously developed vesicle segmentation workflow, the ImageJ macro 3D ART VeSElecT. With that we reliably distinguish CCVs and DCVs in electron tomograms of *C. elegans* NMJs using image-based features. Analysis of the underlying ground truth data shows an increased fraction of DCVs as well as a higher mean distance between DCVs and AZs in dauer larvae compared to young adult hermaphrodites. Our machine learning based tools are adaptable and can be applied to study properties of different synaptic vesicle pools in electron tomograms of diverse model organisms.

2.   Author summary

Vesicles are important components of the cell, and synaptic vesicles are central for neuronal signaling. Two types of synaptic vesicles can be distinguished by electron microscopy: the classical "clear core" vesicles (CCVs) and the typically larger "dense core" vesicles (DCVs). The distinct appearance of vesicles is caused by their different cargos. To rapidly distinguish between both vesicle types, we present here a new automated approach to classify vesicles in electron tomograms. We combine machine learning with an extension of our previously developed vesicle segmentation workflow, an ImageJ macro, to reliably distinguish CCVs and DCVs using specific image-based features. The approach was trained and validated using data-sets that were hand curated by microscopy experts. Our technique can be transferred to more extensive comparisons in both stages as well as to other neurobiology questions regarding synaptic vesicles.

3.   Introduction

Caenorhabditis elegans is a well-studied model organism. Its small nervous system, consisting of 302 neurons [4,5] allows for studies of the connectome in its entirety [5–7]. A focus of research is on *C. elegans* young adult hermaphrodites, which makes them a reference system for comparison with other developmental states in the life cycle. The most remarkable of all *C. elegans* developmental states is the dauer larva.

When the L1 stage larvae sense adverse environmental conditions such as heat, overpopulation and starvation, they activate an alternative state, diapause, instead of continuing the reproductive life cycle [8,9]. Dauer larvae are characterized by their ability to survive several months without any food uptake [10]. Although metabolic activity is reduced [10], dauer larvae are able to move quickly, if necessary, in particular to escape adverse environmental stimuli and in order to reach and colonize new habitats. Improvement of environmental conditions leads to dauers to exit diapause, resume development into pre-adult L4 larvae, and re-enter the reproductive life cycle [9]. The mechanisms behind this strikingly fast developmental transition, and the role of the dauer nervous system in sensing and responding to environmental cues, remain largely enigmatic.

Neuronal signaling plays a critical role in regulating the dauer stage. Neuronally released insulin-like peptides [11–14] and morphogens such as TGF-beta [15,16] regulate the activation of dauer diapause. These observations suggest potential changes of the nature of neuronal signaling in dauer larvae.

Neuropeptides are stored and released through dense core vesicles (DCVs) [1,17,18]. Recent studies, particularly the electron microscopic reconstruction of *C. elegans* prepared by high-pressure freezing, reveal that almost all synapses, including

the neuromuscular junctions (NMJs), contain two principal types of vesicles, a small

fraction of DCVs and the much more abundant clear core vesicles (CCVs) [19–21].

CCVs are characterized by their clear inner core in electron micrographs, which

stands in strong contrast to the stained vesicle membrane. CCVs typically contain

neurotransmitters such as acetylcholine and GABA in the *C. elegans* NMJs that are

released into the synaptic cleft after fusion of the CCV with the presynaptic cell

membrane at the release sites at active zones [2]. In contrast to CCVs, the role of

DCVs is much more enigmatic. DCVs are characterized by their eponymous dark,

electron dense core that typically contains neuropeptides, small protein molecules

with about 3 - 100 amino acids [1] or other protein cargos such as active zone

proteins [3]. DCVs have a bigger average size of about 60 - 80 nm in comparison to

CCVs, which have a much smaller diameter [19], that can vary in different organisms

or developmental states [22,23]. Whereas CCVs cluster around the active zone,

DCVs tend to locate further away from the release site, typically at the rim of the SV

pool as described for *C. elegans* NMJs [19].

DCV exocytosis is considered to mediate slow synaptic and non-synaptic

signaling in neurons [1]. It typically occurs independently of the active zone and only

at high rates of neuronal activity [1,2]. In *C. elegans*, neuropeptides participate in a

wide variety of behaviors such as egg-laying, locomotion, dauer formation, mechano-

and chemosensation depending on the expression pattern of the peptides [2]. Despite

the known critical involvement of neural peptides in dauer development, whether the

dauer states exhibit differences in DCV distribution in the nervous system remains

unknown.

We used high-resolution 3D transmission electron microscopy (electron

tomography) to systematically investigate differences of DCV and CCV in NMJs of

C. elegans young adult hermaphrodites and dauer larvae. Series of 2D images were

taken from different viewing angles by tilting the sample, and reconstructed *in silico* to

obtain one 3D image stack with near isotropic resolution of typically 4 – 5 nm,

determined by measuring the "unit membrane" thickness [19,25–28]. Quantification of

SV in 3D electron tomograms requires reconstruction of the vesicle pool of a sufficient

number of tomograms, which is very time-consuming and prone to mistakes when

performed manually.

To circumvent these shortcomings we previously developed *3D ART*

VeSElecT, a workflow for automated reconstruction of vesicles [22]: This ImageJ/Fiji

tool allows for automated reconstruction of synaptic vesicle pools in electron

tomograms, but is not able to differentiate between different types of vesicles. To

solve this issue, we here present a python-based machine learning approach that

trains vesicle classifiers, and a straightforward ImageJ/Fiji tool that applies a trained

classifier to previously detected vesicles.

We show that our technique can differentiate between DCV and CCV in two

developmental states, young adult hermaphrodites and dauer larvae. The application

of these image analysis and machine learning tools for electron tomograms is in

principle useful to study differences in synaptic vesicle pools of many other model

organisms.

4.   Results

4.1   Manual classification shows three distinct vesicle types

To compare vesicle properties in two different developmental states of *C. elegans*,

dauer larvae and young adult hermaphrodites, we used high-pressure freezing and

subsequent freeze substitution to immobilize and fix animals in a near-to-native state.

Using electron tomography on ~250 nm cross-sections of the worm, we obtained high

resolution 3D images that allow for investigation of the ultrastructure of the

nematode's NMJs. We used this information to compare synaptic vesicle pool

characteristics, aiming for an automated approach for vesicle classification.

Upon visual examination of electron microscopic datasets of the NMJs of

C. elegans dauer samples, we noted an increased fraction of DCV in dauer larvae

compared to young adult hermaphrodites. To systematically quantify this, vesicles

were automatically extracted from tomograms, using an improved version of our 3D

ART VeSElecT workflow (see S1 Fig), and then manually assigned for vesicle types

by two experts. In uncertain cases, a third expert was consulted. Since closer

examination of tomograms revealed that a definite assignment of some vesicles to

one of the two classes (CCV and DCV) was difficult, we initially established a third,

hypothetic group of non-determinable (ND) vesicles. Non-assignable (NA) vesicles

are a subgroup of ND that we could ultimately not classify as CCV or DCV. Fig 1

shows a representative selection of clear core, non-determinable and dense core

vesicles from 8 exemplary tomograms (4 dauer larva tomograms and 4 hermaphrodite

tomograms).

Fig 1. Comparison of CCV, ND vesicles and DCV of dauer larvae and young adult hermaphrodites. A selection of representative pictures is shown. Pictures of the same developmental state (young adult hermaphrodite (H) / dauer larva (D)) and with identical number (I – IV) are from the same tomogram (e.g. CCV DI, ND DI and DCV DI originate from exactly the same tomogram). Edge length of micrographs = 100 nm.

Together, we included 1072 segmented vesicles from the 7 young adult hermaphrodite tomograms, and 1108 segmented vesicles from the 8 dauer larva tomograms for analyses (Table 1). When we included all identified vesicle populations from all tomograms, the percentage of manually labeled DCVs differed slightly for dauer larvae (13%) and adult hermaphrodite (10%) datasets. After filtering the datasets, comparing only the vesicle population from the same type of NMJ with the bouton fully reconstituted in the tomograms (putative cholinergic NMJs with central cut

161 through the AZ area and at least 100 vesicles), the dauer larvae NMJs exhibited a

162 higher DCV ratio of 16% versus 9% in young adult hermaphrodites.

164 Table 1: number and percentage of vesicle types in all tomograms[1]

Developmental state	Tomograms (n)	Amount of vesicles	CCV	DCV	NA	Total
Adult hermaphrodite	7	Sum	949	109	14	1072
		Percentage	89%	10%	1%	
	5	Sum [1]	837	85	11	933
		Percentage [1]	90%	9%	1%	
Dauer larvae	8	Sum	948	148	12	1108
		Percentage	86%	13%	1%	
	5	Sum [1]	669	130	3	802
		Percentage [1]	83%	16%	0%	

[1] analysis with a selected set of tomograms where only putative cholinergic NMJs

with apparent dense projections and with > 100 vesicles were included.

4.2. Characterization based on image features

In order to perform a quantitative comparison of vesicles and for clarification about the

ND vesicle class, we extracted a number of image features based on qualitative

observations. Since DCVs appear larger, darker and further apart from the active

zone compared to CCVs, we used radius (r), mean gray value (gv) and distance to

active zone (distAZ) as primary features (Fig 2). In addition, the standard deviation of

the gray value distribution of a vesicle (GVSD) is a measure for the contrast between

membrane and core that is independent of the brightness of the vesicle.

Fig 2. Visualization of vesicle pool in a dauer larva tomogram with color coded image
176

features: inner radius, gray value, distance to active zone (distAZ), gray value standard

deviation (GVSD). Clear core like characteristics are yellow-shifted, dense core like features

are blue shifted. The active zone is located at the bottom.

Statistical analysis and comparison of ND vesicles with CCV and DCV suggests that

synapses show no third type of vesicles and that NDs are in most cases CCVs (see

S2 Fig). Hence, for further analysis, we proceeded with the assumption of two existing

vesicle classes, CCVs and DCVs (Table 1). We tried to apply simple thresholding and

manually weighted linear combinations of vesicle radius, distAZ and gv to separate

DCVs from CCVs, but with limited success (see S2 Fig and S1 Text). Therefore, a

machine learning approach was developed and applied, as described below. To

generate training data, ND vesicles were first tried to be concordantly assigned to one

group by two experts. Their label was only changed if they turned out to be DCV, as

we programmed classifiers to treat the label ND as CCV. In difficult cases, a third

expert was consulted. Still non-assignable vesicles (NA) were excluded from training

by labelling them as error (see table 1: n = 12, ~ 1.1 % in dauer larvae; n = 14, ~1.3 %

in young adult hermaphrodites).

4.3. Machine learning can distinguish vesicle types

To automatically classify vesicles using the previously calculated image features, we

evaluated different machine-learning algorithms. We compared a support vector

machine (SVM), random forest (RF), and k nearest neighbor (KNN) for their

performance on differentiating between CCVs and DCVs using four image-based

characteristics described above: radius (r), mean gray value (gv), standard deviation

of the gray value (GVSD) and distance to the active zone (distAZ) (see Fig 3). 1994

manually labelled vesicles from 15 different tomograms (7 young adult hermaphrodite

and 8 dauer larval tomograms) were used as training data. The features were

extracted as in the first two steps of our classification macro (see Fig 3 and Methods

section 6.3), saved as a table, and processed in a python script using the scikit-learn

machine learning library [29]. Table 2 shows the results of the comparison of the

different algorithms. Evaluation scores of the whole dataset were impractically high,

as CCV greatly outnumber DCV. Therefore precision$_{DCV}$, recall$_{DCV}$ and F-score$_{DCV}$

calculates scores for DCV specifically.

Table 2. Average performance of different machine learning classifiers

classifier	accuracy	precisionDCV	recallDCV	F-scoreDCV
SVM	0.98 ± 0.02	0.95 ± 0.07	0.90 ± 0.1	0.92 ± 0.07
Random forest (t = 10)	0.97 ± 0.03	0.88 ± 0.12	0.89 ± 0.09	0.88 ± 0.08
Random forest (t = 1500)	0.97 ± 0.02	0.93 ± 0.08	0.86 ± 0.1	0.89 ± 0.07

KNN (k = 10)	0.98 ± 0.02	0.92 ± 0.09	0.89 ± 0.12	0.90 ± 0.09

The comparison between SVM, random forest and KNN revealed that the SVM

algorithm performs best on our classification problem. The SVM shows the lowest

error rates in all four categories (see Methods): Accuracy, precision$_{DCV}$, recall$_{DCV}$ and

F-score$_{DCV}$. KNN shows second best results that were about 0-3 percent worse

compared to the SVM. For the random forest variants, t = 1500 trees performed

similarly or better than t = 10, except for recall$_{DCV}$. Random forest (t = 1500) shows

overall third best results with 3 percent worse results for recall$_{DCV}$ and about 1 percent

worse results for accuracy, precision$_{DCV}$ and F-score$_{DCV}$ compared to KNN. Resulting

values of the SVM hyperplane delimiting DCVs from CCVs are shown in Table 3.

Table 3. SVM and standardization parameters transferred to the Fiji macro

parameter	r	gv	distAZ	GVSD
mean	10.4	129.1	259.1	5.9
SD	3.1	4.7	118.5	1.7
weight	-1.69	1.65	-0.76	1.21

The final trained SVM classifier used the following weighting of the four features:

$\quad -4.36 = -1.69 * r + 1.65 * gv - 0.76 * distAZ + 1.21 * GVSD \qquad (1)$

The weight for distAZ is smaller than for the other three parameters, which

indicates that it is less important. Still, it contributes to the classifier, otherwise the

resulting weight after training would be very small or zero. In such cases, it could be

discarded from the training data.

In summary, the SVM algorithm in its linear form without kernelization

outperformed all other classification methods including non-linear SVM and random

forest (t = 1500) classification. Additionally, using a linear SVM has the advantage that

the resulting classifier can easily be incorporated into a classification workflow by

extracting the coefficients after successful training. For this purpose, a new vesicle

classification macro in ImageJ/Fiji was developed to be consecutively applied after the

modified 3D ART VeSElecT registration macro. The workflow of the classification

macro is shown in Fig 3 and is described in detail in Materials and Methods.

Fig 3. Workflow of the classification macro. 1) preparation: StackSegmented (registered

vesicles) and StackDuplicateScale (scaled tomogram) from 3D ART VeSElecT [22] are used

as input; the user is requested to mark the active zone via point-selection (semi-automated).

2) readout of features: gray value (gv), radius (r), vesicles coordinates and gray value

standard deviation (GVSD) in the middle slice of vesicles are determined; the distance to

active zone (distAZ) is calculated. 3) processing of features: offsetting, standardization and

weighting of features followed by final assignment of labels. 4) output: includes a logfile and

visualization of results as new image stack windows.

4.4. Application to *C. elegans* dauer and adult hermaphrodite NMJs

We used the ground truth classification to quantify differences in the vesicle pool of

C. elegans dauer and adult hermaphrodite NMJs in electron tomograms. Additionally,

visual results of 3D ART VeSElecT and the classification macro for one exemplary

tomogram of each analyzed developmental state are presented in Fig 4 to provide an

intuitive impression of DCV and CCV distribution.

Fig 4. Comparison of the visual results from the adapted 3D ART VeSElecT and the

consecutive classification macro. *Top:* I) and II) original tomograms of young adult

hermaphrodite (I) and dauer larva (II). *Bottom:* a) and c) show *Composit* stacks of 3D ART

VeSElecT (random colors), whereas a') and c') show the resulting *Composit* after

classification of vesicles (CCV = magenta, DCV = green). b) and d) show the 3D model of the

vesicle pool, b') and d') show the 3D model with color labels showing the assigned class. All

scale bars = 100 nm.

The 3D models show that DCVs most often localize on the rim of the CCV cluster.

CCVs themselves gather around the dense projections, electron dense structures

attached to the cell membrane that indicate the active zone where synaptic vesicle

fusion occurs. For further analysis of vesicle characteristics (radius, gray value, GVSD

and distAZ), results are shown as violin plots in Fig 5.

Fig 5. Comparison of DCVs (black) characteristics with CCVs (light gray) characteristics. Violin plots show a) inner vesicle radius in nm, b) gray value (gv), c) distance to active zone (distAZ) and d) gray value standard deviation in central slice of each vesicle (GVSD). Panels b)-d) show normalized values of $n_h = 7$ tomograms of N2 young adult hermaphrodites; $n_d = 8$ tomograms of N2 dauer larvae. All plots are based on expert curated manual ground truth classification.

Fig 5a shows highly significant differences in CCV and DCV size in both

developmental states. In the following, the radius always refers to our original

measured results, which is the "inner radius" of all vesicles, as described in [22]. To

make vesicle size comparable to other published results, the measured radius must

be multiplied by two to receive the inner diameter and increased by 9 nm to include

vesicle membrane thickness [22]. This estimated diameter is only an approximation

and cannot be directly compared to manually measured data due to the different way

how both are measured [22].

CCVs of dauer larvae show a median inner radius of 9.0 nm, whereas DCVs

have a median inner radius of 14.7 nm, which results in a significant difference of 5.7

nm. However young adult hermaphrodites' median inner radius of DCVs is 16.7 nm,

hence they have a ~ 5.4 nm bigger radius than CCVs with a median inner radius of

10.3 nm. Noticeably, both vesicle types are slightly but significantly bigger in young

adult hermaphrodites compared to dauer larvae. The radius of hermaphrodite CCVs is

1.3 nm larger compared to dauer larvae, whereas the radius of hermaphrodite DCVs

is 2 nm larger compared to dauer larvae.

In all cases, DCVs have significantly darker normalized gray value intensities in

comparison to CCVs, as shown in Fig 5b. Young adult hermaphrodites show a

normalized gv of 0.17 for CCVs and 0.45 for DCVs, whereas dauer larvae have a

normalized gv of 0.23 for CCVs and 0.56 for DCVs. Hence, vesicles of dauer larvae

tomograms possess significantly darker vesicles (both CCVs and DCVs) in

comparison to hermaphrodite tomograms. In contrast, GVSD (gray value standard

deviation in the central slice of each vesicle) does not differ significantly between

developmental states, but shows highly significant differences of ~ 0.22 in GVSD

values of CCV to DCV for both developmental states (Fig 6d).

The normalized distance of CCVs to the active zone distAZ is smaller (0.42 for

young adult hermaphrodites and 0.4 for dauer larvae) than that of DCVs (young adult

hermaphrodites: 0.64, dauer larvae: 0.72) in both larval stages (Fig 6c). The

normalized distances of DCVs to the AZ, of young adult hermaphrodites in

comparison to dauer larvae, show a highly significant difference of 0.08 (normalized).

For better visualization of this observation, distances are visualized as histogram plots

in Fig 6.

Fig 6. Cumulative histogram (a, b) and kernel density estimation plots (c, d; bandwidth =

0.07*maximum value of total dataset) of distances to the active zone (distAZ). Given are

normalized distances (left) and real values in nm (right). CCVs are shown as dashed line and

DCVs as continuous line. Adult hermaphrodite vesicles are presented in shades of gray,

whereas dauer larval vesicles are shown in violet. All plots are based on expert curated

manual ground truth classification.

Fig 6a and c illustrate that normalized distances of CCVs of both developmental

states are almost identical, whereas distances of DCVs differ significantly in both

plots. DCVs show a systematic discrepancy in their localization: DCVs in young adult

hermaphrodites cluster closer to the active zone than DCVs of dauer larvae.

Histograms in Fig 6b and d shows results of distAZ in real distance measured in nm.

Interestingly, this statistic shows that vesicle pools in dauer larvae have a generally

smaller maximal distAZ in young adult hermaphrodites. The largest distance of dauer

larvae vesicles from the active zone is about 550 nm whereas the largest distance of

vesicles in young adult hermaphrodites is about 700 nm. It should be noted that

distAZ as derived from tomograms should be considered as an estimate of the actual

distance of vesicles to the active zone, since a single tomographic slice may not

contain the full information on the actual position and shape of the active zone.

4.5. Error rates and cross validation

To generate the ground truth data, a consensus label was created (N = 15

tomograms, 8 dauer larva and 7 hermaphrodite tomograms): First, labels were

assigned independently by two biologists. Differing labels were re-examined. In case

of no agreement, a third examiner was employed. NA vesicles (~1 %) were excluded

from SVM training. The initial assignment by individuals was compared with this

consensus label to determine the manual error rate (Table 4).

Table 4 manual vs classification macro (SVM performance)

	presicion$_{DCV}$	recall$_{DCV}$	F-score$_{DCV}$
Manual [1]	0.98 ± 0.05	0.96 ± 0.08	0.97 ± 0.05
SVM [2]	0.95 ± 0.07	0.90 ± 0.10	0.92 ±0.07

$^{(1)}$ Compares two independent manual assignments to the consensus label

(ground truth)

$^{(2)}$ Compares the SVM assignment to the ground truth

Manual annotation shows higher scores than the SVM. Manual assignment of

vesicle types shows 98 % precision$_{DCV}$, whereas the SVM reaches 95 % precision$_{DCV}$.

The results for recall$_{DCV}$ is 96 % for manual assignment and 90 % for the SVM. The F-

score is the harmonic mean of precision and recall (see Materials and Methods) and

shows a mean result for the error rate of 97% for manual and 92% for automated

classification. Results for precision and recall are additionally shown as scatterplot in

Fig 7.

Fig 7. SVM error rates as precision$_{DCV}$ and recall$_{DCV}$ plot. Each data point represents one

validation tomogram. Red squares: adult hermaphrodite, blue circles: dauer larvae. Right:

magnified detail plot. Red star indicates mean of hermaphrodite data: precision$_{DCV}$ = 0.93,

recall$_{DCV}$ = 0.85. Blue cross indicates mean of dauer data: precision$_{DCV}$ = 0.97, recall$_{DCV}$ =

0.94. 3 dauer tomograms and 1 hermaphrodite tomogram were classified without error.

The SVM classifier was cross-validated against each tomogram using all other

tomograms as training data. Fig 7 shows error rates of cross validation of automated

SVM classification. Hermaphrodite tomograms are represented as red squares and

dauer larvae tomograms are represented as blue circles. Mean classification results

are shown as red star for young adult hermaphrodites (precision$_{DCV}$ = 0.93, recall$_{DCV}$ =

0.85) and as blue cross for dauer larvae tomograms (precision$_{DCV}$ = 0.97, recall$_{DCV}$ =

0.94). Dauer larvae tomograms have in average better results than hermaphrodites

and cluster in the right upper corner. 3 dauer tomograms and one young adult

hermaphrodite tomogram were classified without any error and overlay in the right

upper corner at precision$_{DCV}$ = 1.0, recall$_{DCV}$ = 1.0.

5.   Discussion

Automated methods for image analysis are of great importance to reduce manual

labor in the long term and prevent biased analysis of data, hence making double blind

analysis unnecessary. This is confirmed by our own experience, where one re-

analysis of the same tomogram by the same expert revealed one differential

interpretation of CCV and DCV and 6 other non-matching assignments on n = 134

vesicles in total. Such deviations in manual analysis may have a great influence on

the outcome. For this reason, we had two experts manually assign all vesicles

independently. In case of inconsistency, both persons came to a mutual agreement or

consulted a third expert in the decision process. Still unassignable vesicles were

excluded from analysis. Since this evaluation process is very elaborate and time-

consuming, a long-term automated solution is inevitable. As far as we know there is

no other approach for the specific task of automated vesicle classification in electron

tomograms. We present here a new methodological approach that is applied

subsequent to a modified version of our previously published Fiji macro 3D ART

VeSElecT [22]. First, a Fiji macro was written to read out important characteristics.

Those were, together with manual labelling, used to train machine learning classifiers

in python. Extracted machine learning parameters of an SVM completed the new Fiji

macro to classify vesicles and visualize classes through 3D color labelling. Our tool

can thus be used for automated vesicle classification. We used our ground truth label

to compare two different developmental states of *C. elegans*, young adult

hermaphrodites and dauer larvae.

Manual analysis revealed a hypothetical third type of vesicles, called non-

determinable vesicles (ND) which are most likely CCVs with no perfectly clear inner

core. This seems reasonable, since they resemble CCV characteristics in radius and

distance to the active zone and only differ significantly in their gray value, which lies

among gv intensities of DCVs and CCVs (S2 Fig). This observation can be explained

by either biological or technical reasons. On the one hand, medium-gray vesicles

could be DCVs with partially released cargo, e.g. after kiss-and-run [30–32]. Overall

this does not explain why NDs have the same size as CCVs, since DCVs are known

to be considerably larger [23]. On the other hand, vesicles may appear darker in

electron micrographs because of lower quality of tomograms at some locations.

Lacking precision of the tomogram overlay of the two single tilted tomograms to

receive one double tilted tomogram could result in darker, more blurry vesicles with no

perfectly clear core.

The comparison of DCV number in dauer larvae and young adult

hermaphrodites revealed differences in the ratio of DCVs (9 % in young adult

hermaphrodites and 16 % in dauer larvae), when comparing only cholinergic

synapses with apparent dense projections and at least 100 vesicles in the NMJ.

Furthermore we found that DCV ratio correlates with cell size change throughout the

tomogram (S4 Fig). A large cell size change indicates inclusion of decentral regions of

the synaptic bouton in the tomogram. We assume that inclusion of more remote areas

in the tomogram is more likely in dauer larvae, since the overall size of the synaptic

bouton is smaller (mean diameter ± SD of cross section: 452 ± 85 nm) compared to

young adult hermaphrodites (mean diameter ± SD of cross section: 542 ± 97 nm).

Therefore, standard sized 250 nm sections, typically prepared for electron

tomography, naturally have a higher chance of catching decentral regions in dauer

larvae NMJs. For a more comprehensive understanding, serial section tomography for

visualization of complete synaptic boutons would be necessary.

Additionally, we found that young adult hermaphrodites show significantly larger

vesicles (CCVs and DCVs) and diameters of synapses in comparison to dauer larvae.

This observation could either be a biological phenomenon of dauer larvae or a side

effect of the sample preparation. Dauer larvae have a distinct cuticle that shields the

animal from adverse environmental conditions, like heat and drought. This could lead

to adverse effects during fixation (freeze substitution) of the sample, even though we

did not observe any sign of deficient sample fixation in dauer larvae. Divergent

synaptic vesicle sizes have already been found in NMJs of other organisms, like rats

[33] and zebrafish [23]. For zebrafish a dependency of vesicle size on the maturation

stage of NMJs was assumed that might be related to the functional transition of these

synapses to control different swimming behavior [23]. Differences in synaptic vesicle

sizes during normal development have not been observed for *C. elegans*, to our

knowledge. However, divergent vesicle sizes could also be related to different moving

behavior of dauer larvae that are well known for their unique nictation behavior [9].

Since manual measurements are often inaccurate, smaller deviations in vesicle sizes

are only observable in extensive analysis, therefore automated approaches are

inevitable. Our previously published macro 3D ART VeSElecT proved to be

successful for distinguishing vesicle size differences [22]. This approach in

combination with our here presented extended version for automated classification

may be the key to reveal so far undistinguishable differences in the future.

Interestingly, although nervous cells in dauer larvae are typically smaller,

segregation of DCVs and CCVs is more distinct. Our interpretation is that DCVs in

dauer larvae locate further away from the active zone on the rim of the synaptic

vesicle pool compared to young adult hermaphrodites. Differences in this

characteristic of synaptic architecture could be explained by a shift towards a higher

ratio of DCVs in a non-fusion competent resting state in dauer compared to young

adult hermaphrodites. Moreover, the location of DCVs closer to the cell margin in

dauer compared to young adult hermaphrodites could indicate the need for faster

peptide release, and their increased number could compensate for lower volume.

For automated classification of vesicles, different machine learning algorithms

were tested. The SVM showed best results in comparison to random forest or k-

nearest neighbor classification. Furthermore, the SVM works regardless of

dimensionality, can classify non-linearly separable classes by using the kernel trick,

enables an easy integration of the trained linear classifier into the macro and generally

can be used for assignment into more than two classes [34].

Cross-validation showed very good results for automated classification of

vesicles. Interestingly, dauer larval tomograms showed better results than young adult

hermaphrodite tomograms, which could be related to the differing gray value

intensities of the tomograms. Hence high contrast vesicles were easier to

automatically recognize and showed more distinct results for GVSD and gv. Another

reason could be the more pronounced positional segregation of the two vesicle types

for dauer larvae.

We suggest that our pre-trained SVM can be used for DCV and CCV classification in

other, similar tomograms, and provide the Fiji classification macro including the

trained parameters. In addition, users can train a SVM on their own data using the

python script we provide, and integrate the parameters into the Fiji macro, which is

quickly achievable via an input dialog.

Our ImageJ classification macro and the python script, for retraining of the classifier, a

tutorial and test datasets are available at: https://www.biozentrum.uni-

wuerzburg.de/bioinfo/computing/3dart-veselect/.

6.   Conclusion

We developed a new automated approach to classify vesicles and to quantify their properties from electron tomograms. We combined machine learning with an extension of our previously developed vesicle segmentation workflow 3D ART VeSElecT to reliably distinguish clear core from dense core vesicles using image-based features. We apply this method to electron tomograms of *C. elegans* NMJs in young adult hermaphrodites and dauer larvae. Using our ground truth data, we find an increased fraction of dense core vesicles (~ 16 % vs ~ 9 %) in tomograms matching certain characteristics (cholinergic synapses with > 100 vesicles of a central slice through the AZ area), as well as significantly reduced vesicle size and a higher distance of dense core vesicles to the active zone in dauer larvae compared to young adult hermaphrodites. Our approach is not limited to *C. elegans* and can be easily adapted to study differences in synaptic and other vesicles in electron tomograms in various settings and many other model organisms, e.g. in *Danio rerio* [22].

7.   Material and Methods

7.1 Electron Microscopy

<u>Animals</u>

C. elegans wild-type (Bristol N2) were used and maintained using standard methods [4] . For the experiments, young adult hermaphrodite worms as well as dauer larvae were used.

<u>Purification of dauer larvae</u>

For segregation of dauer larvae from other states of the C. elegans life cycle, treatment with sodium dodecyl sulfate (SDS) buffer was used [35]. Firstly, worms were washed from the agar plates using M9 medium [36]. The worm solution was then transferred into 50 ml tubes and centrifuged for 5 min at 2000g at 4°C. Afterwards, the supernatant was removed and the pellet was dissolved in 50 ml 1 % SDS solution and incubated on a shaker for 15 min. SDS was then removed by centrifuging (2000g, 5 min, 4°C) and subsequent discarding of the supernatant while retaining 5 ml of solution. The tube was then refilled with distilled water and the procedure repeated 2-3 times. For separation of dauer larvae from dead material, the solution was pipetted into a pit on an unseeded agar plate. Once the solution had dried, living dauer larvae were able to crawl away from the pit. After 1-3 hours, emigrated worms were washed off with M9 medium.

<u>High-pressure freezing, freeze-substitution and electron tomography</u>

HPF, FS and electron tomography were performed as previously described [22]. Tilt series were recorded from at least +65° to -65° but not exceeding +70° to -70°.

7.2 Improvements to 3D ART VeSElecT

We extended our tool 3D ART VeSElecT to calculate various image features of

detected vesicles to be used for classification. The workflow is shown in

Supplementary Figure S1, with changes compared to the previous version highlighted

by a black box. The overall workflow of 3D ART VeSElecT remained as previously

published [22]. One change in the preprocessing step is the determination of the

mean radius instead of volume, since the radius or diameter is typically used when

comparing vesicle size. Furthermore, automated calculation of the cell volume was

added. Since semi-automated selection of the cell outline by the user was already

included in the macro, this selection is now basis for calculating the analyzed volume

in 3D (which corresponds to cell volume if the selection is taken along the cell

membrane, as described in the 3D ART VeSElecT tutorial).

7.3 Classification Macro

The workflow of the new Fiji macro for classification of DCV and CCV in electron

tomograms of synapses comprises four main steps: 1) preparation, 2) reading out

features, 3) processing of features followed by labelling of vesicles, and 4) output. In

the preparation step, StackDuplicateScale and StackSegmented, which result from

the 3D ART VeSElecT recognition macro, are read into the classification macro.

StackDuplicateScale is the scaled original tomogram, StackSegmented contains the

registered vesicles. First, the macro opens a preparation window for changing the

names of read-in stacks (StackDuplicateScale and StackSegmented) and setting

parameters for standardization and adaption of SVM weights. In case of adaptation of

the SVM algorithm to another user's needs, values can easily be changed through the

input window. In a second step, four characterizing vesicle features are computed

from the images: radius, gv, GVSD and distAZ. GVSD is a value for standard

deviation of gray values in the central slice of each vesicle, since CCVs have a higher

gv variance than DCV. In the third step, features are processed, which includes

applying an offset to gv (if the darkest vesicle is lighter than a threshold),

standardization of parameters, weighting features based on SVM weights, and finally

the assignment of labels according to the identified class (CCV = magenta, DCV =

green). In the last step, the output is created, including a logfile and visual output

shown in new windows. The logfile includes all measured characteristics (e.g.

features, classification labels (CCV = 1, DCV = 0), number of vesicles in each class).

The visual output shows, besides the input stacks ("StackSegmented" and

"StackDuplicateScale"), an additional image stack, named "label" which corresponds

to the StackSegmented but displays color-coded vesicles according to their

classification. Furthermore a Composit stack, named "CompositLabel" (overlay of

original tomogram and the "label" stack) is created.

7.4 Comparison of different types of machine learning classifiers

We compared a support vector machine (SVM), random forest (RF, using t = 10, t =

1500; t indicates the number of trees) and k nearest neighbor (KNN, k = 10

neighbors) for their performance on differentiating between CCVs and DCVs based

on four image-based characteristics: radius (r), mean gray value (gv), standard

deviation of the gray value (GVSD) and distance to the active zone (distAZ). Random

forest uses decision trees to classify samples: Comparable to a flowchart, samples

are divided based on their features. A random forest utilizes the majority classification

of t decision trees each grown on randomly selected training data and split using the

best randomly selected feature. Generalization error is lower than for individual

decision trees. Performance, but also runtime increases with t. KNN classifies

samples based on the majority of their k nearest training samples in feature space.

Linear SVM (C = 1) showed the best average classification results (see Table 2). C

determines the penalty for falsely classified training data points and therefore

indicates bias-variance trade-off, since bias decreases with C [37]. The SVM will be

described more accurately below. Parameters of the SVM (Table 3) were

determined using all input tomograms as training data and transferred to a Fiji-

Macro (Figure 3) to enable classification downstream of the 3D ART VeSElecT

registration macro.

7.5 Support Vector Machine (SVM)

A linear support-vector machine (SVM) divides two classes by a hyperplane. As

such, the SVM is a binary classifier dealing with linear classification problems. A

one‑versus-all approach and kernelization can extend classification to non-linear

problems and into more than two classes, but will not be discussed here since the

classification power of the SVM is sufficiently strong, see results. With the aim of

minimizing generalization error, the optimization objective of the SVM is to

maximize the margin between the hyperplane and the closest data points of the

two classes. Those eponymous support-vectors lie on the positive and negative

hyperplane, which are equidistant to the separating hyperplane [37]. As the linear

SVM is parametric, the equation of the separating hyperplane can be transferred

independently of the training-data (see Results, Table 3 and Equation 1), with x_i

being the features and ω_i the assigned feature-weights. All data points with

$\sum_{i=1}^{n} w_i x_i'' > 0$ are assigned to class 1 (CCV), whereas all data points with

$\sum_{i=1}^{n} w_i x_i'' < 0$ are assigned to class -1 (DCV).

7.6 Implementation of machine learning algorithms

A python 3 script (S3 Fig) reads in the CSV files containing the training datasets

and applies classifiers from the scikit-learn library [29]. The manual labels are

transformed into a feature array: Particles marked "D" (DCV) are relabeled "-1".

Particles marked "E" (error) are removed. All other markings are relabeled "1",

including "C" (CCV) and "N" (not determinable), since non-determinable

particles show more similarities to CCV (S2 Fig).

Classifiers are evaluated by cross-validation: Data from one CSV file are used as

validation data, all other files are concatenated and used as training data (there

are as many training-validation combinations as input CSV files). For each

combination, features are standardized using mean and standard deviation (sd)

of the training data and passed to the classifiers, namely linear SVM, random

forest and KNN. An evaluation function determines accuracy of all vesicles and

precision, recall and F-score of DCV only (see equations (2) - (4)). Evaluation

results are returned as pandas.DataFrame [38] and as CSV file.

r	gv	distAZ	GVSD	Manual_label		Label-array
12.685	134.98	461.932	4.266	D	→	-1
6.631	150.42	190.144	7.5	C		1
9.354	144.51	405.247	7.633	E		Deleted
7.911	134.485	473.051	4.539	N		1
...	...	...	...	...		...

Features

Fig8. Format of input CSV file. First four columns are the four features, the fifth column is

the manual labels. The python script rewrites manual labels into a binary label-array: "D"

(DCV) is rewritten to -1, "E" (error) is removed, and everything else is rewritten as "1" (CCV).

Equations:

$$precision_{DCV} = \frac{tp_{DCV}}{tp_{DCV} + fp_{DCV}} \qquad (2)$$

$$recall_{DCV} = \frac{tp_{DCV}}{tp_{DCV} + fn_{DCV}} \qquad (3)$$

$$F_{DCV} = 2 * \frac{DCV_{precision} * DCV_{recall}}{DCV_{precision} + DCV_{recall}} \qquad (4)$$

tp: true positive; fp: false positive; fn: false negative.

The runtime of the training script for the different classifiers was 5 s for t=10 trees and

140 s for t=1500 trees. The ImageJ classification macro takes 33s for the calculation,

plus the time required to manually click on the active zone. All computations were

performed on a 16-core AMD Opteron 1.7 GHz server with 128 GB RAM running

Ubuntu Linux 16.04. In comparison, manual classification of vesicles takes an

experienced user about 1 h per tomogram. The time required to set up the python

environment and prepare the training data depends on the level of experience of the

user, but overall there should still be at least a factor of 10 (one order of magnitude) of

speed-up with reasonably large data sets of a few tomograms, as in our case. In

addition, automated workflows as reported here increase reproducibility compared to

manual assignment.

8. References

1. Salio C, Lossi L, Ferrini F, Merighi A. Neuropeptides as synaptic transmitters. Cell Tissue Res. Springer-Verlag; 2006;326: 583–598.

2. Li C. Neuropeptides. WormBook. 2008; 1–36.

3. Zhai RG, Vardinon-Friedman H, Cases-Langhoff C, Becker B, Gundelfinger ED, Ziv NE, et al. Assembling the presynaptic active zone: a characterization of an active one precursor vesicle. Neuron. 2001;29: 131–43.

4. Brenner S. The genetics of Caenorhabditis elegans. Genetics. Genetics Society of America; 1974;77: 71–94.

5. White JG, Southgate E, Thomson JN, Brenner S. The Structure of the Nervous System of the Nematode Caenorhabditis elegans. Phil Trans R Soc Lond B. 1986;314 1165: 1–340.

6. Varshney LR, Chen BL, Paniagua E, Hall DH, Chklovskii DB. Structural Properties of the Caenorhabditis elegans Neuronal Network. PLoS Comput Biol. 2011;7: 1–21.

7. Haueis P, Slaby J. Connectomes as constitutively epistemic objects: Critical perspectives on modeling in current neuroanatomy. Progress in brain research. 2017. pp. 149–177.

8. Golden JW, Riddle DL. The Caenorhabditis elegans dauer larva: Developmental effects of pheromone, food, and temperature. Dev Biol. Academic Press; 1984;102: 368–378.

9. Cassada RC, Russel RL. The dauerlarva, a post-embryonic developmental variant of the nematode Caenorhabditis elegans. Dev Biol. Academic Press; 1975;46: 326–342.

10. Erkut C, Kurzchalia T V. The C. elegans dauer larva as a paradigm to study metabolic suppression and desiccation tolerance. Planta. Springer Berlin Heidelberg; 2015;242: 389–396.

11. Pierce SB, Costa M, Wisotzkey R, Devadhar S, Homburger SA, Buchman AR, et al. Regulation of DAF-2 receptor signaling by human insulin and ins-1, a member of the unusually large and diverse C. elegans insulin gene family. Genes Dev. Cold Spring Harbor Laboratory Press; 2001;15: 672–86.

12. Hung WL, Wang Y, Chitturi J, Zhen M. A Caenorhabditis elegans developmental decision requires insulin signaling-mediated neuron-intestine communication. Development. Oxford University Press for The Company of Biologists Limited; 2014;141: 1767–79.

13. Li W, Kennedy SG, Ruvkun G. daf-28 encodes a C. elegans insulin superfamily member that is regulated by environmental cues and acts in the DAF-2 signaling pathway. Genes Dev. Cold Spring Harbor Laboratory Press; 2003;17: 844–58.

14. Cornils A, Gloeck M, Chen Z, Zhang Y, Alcedo J. Specific insulin-like peptides encode sensory information to regulate distinct developmental processes. Development. Oxford University Press for The Company of Biologists Limited; 2011;138: 1183–93.

15. Ren P, Lim CS, Johnsen R, Albert PS, Pilgrim D, Riddle DL. Control of C. elegans larval development by neuronal expression of a TGF-β homolog. Science. 1996;274: 1389–1391.

16. Schackwitz WS, Inoue T, Thomas JH. Chemosensory Neurons Function in Parallel to Mediate a

Pheromone Response in C. elegans. Neuron. Cell Press; 1996;17: 719–728.

17.   Salio C, Averill S, Priestley JV, Merighi A. Costorage of BDNF and neuropeptides within
individual dense-core vesicles in central and peripheral neurons. Dev Neurobiol. Wiley
Subscription Services, Inc., A Wiley Company; 2007;67: 326–338.

18.   Walch-Solimena C, Takei K, Marek KL, Midyett K, Südhof TC, De Camilli P, et al.
Synaptotagmin: a membrane constituent of neuropeptide-containing large dense-core vesicles.
J Neurosci. Society for Neuroscience; 1993;13: 3895–903.

19.   Stigloher C, Zhan H, Zhen M, Richmond J, Bessereau J-L. The presynaptic dense projection of
the Caenorhabditis elegans cholinergic neuromuscular junction localizes synaptic vesicles at the
active zone through SYD-2/liprin and UNC-10/RIM-dependent interactions. J Neurosci. 2011;31:
4388–96.

20.   Kittelmann M, Hegermann J, Goncharov A, Taru H, Ellisman MH, Richmond JE, et al. Liprin-
α/SYD-2 determines the size of dense projections in presynaptic active zones in C. elegans. J
Cell Biol. 2013;203: 849–863.

21.   Lim MA, Chitturi J, Laskova V, Meng J, Findeis D, Wiekenberg A, et al. Neuroendocrine
modulation sustains the C. elegans forward motor state. Elife. 2016;5.

22.   Kaltdorf KV, Schulze K, Helmprobst F, Kollmannsberger P, Dandekar T, Stigloher C. FIJI Macro
3D ART VeSElecT: 3D Automated Reconstruction Tool for Vesicle Structures of Electron
Tomograms. Dyer E, editor. PLOS Comput Biol. Public Library of Science; 2017;13: e1005317.

23.   Helmprobst F, Frank M, Stigloher C. Presynaptic architecture of the larval zebrafish
neuromuscular junction. J Comp Neurol. 2015; 523;13: 1984-97

24.   Phillips MI, M.Ian. Neuropeptides: Regulators of Physiological Processes. Regul Pept. Elsevier;
2000;90: 101.

25.   Mastronarde DN. Dual-axis tomography: an approach with alignment methods that preserve
resolution. J Struct Biol. 1997;120: 343–52.

26.   Zampighi GA, Fain N, Zampighi LM, Cantele F, Lanzavecchia S, Wright EM. Conical Electron
Tomography of a Chemical Synapse: Polyhedral Cages Dock Vesicles to the Active Zone. J
Neurosci. 2008;28.

27.   Zampighi GA, Zampighi L, Fain N, Wright EM, Cantele F, Lanzavecchia S. Conical tomography
II: A method for the study of cellular organelles in thin sections. J Struct Biol. 2005;151: 263–
274.

28.   Robertson JD. The Early Days of Electron Microscopy of Nerve Tissue and Membranes. Int Rev
Cytol. 1987;100: 129–201.

29.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. J Mach Learn Res. 2012;12: 2825–2830.

30.   Artalejo CR, Elhamdani A, Palfrey HC. Secretion: Dense-core vesicles can kiss-and-run too.
Curr Biol. Cell Press; 1998;8: R62–R65.

31.   Rutter GA, Tsuboi T. Kiss and run exocytosis of dense core secretory vesicles. Neuroreport.
2004;15: 79–81.

32. Tsuboi T, McMahon HT, Rutter GA. Mechanisms of dense core vesicle recapture following "kiss
and run" ("Cavicapture") exocytosis in insulin-secreting cells. J Biol Chem. American Society for
Biochemistry and Molecular Biology; 2004;279: 47115–24.

33. Markus EJ, Petit TL, LeBoutillier JC. Synaptic structural changes during development and aging.
Brain Res. 1987;432: 239–48.

34. Shawe-Taylor J, Cristianini N. Kernel Methods for Pattern Analysis [Internet]. Elements. 2004.

35. Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental variant of the
nematode Caenorhabditis elegans. Dev Biol. 1975;46: 326–342.

36. Hope I. C. elegans a practical approach. Oxford Univ Press. 1999;

37. Raschka S. Python Machine Learning. Packt Publishing; 2015.

38. McKinney W. Data Structures for Statistical Computing in Python. Proc 9th Python Sci Conf.
2010;1697900: 51–56.