Adult influence on juvenile phenotypes by stage-specific pheromone production

Michael S. Werner[1,2], Marc H. Claaßen[1,2], Tess Renahan[1,2], Mohannad Dardiry[1], and Ralf. J. Sommer[1*]

[1]Department of Evolutionary Biology, Max Planck Institute for Developmental Biology, 72076 Tübingen, Germany

[2]These authors contributed equally
*Corresponding author: ralf.sommer@tuebingen.mpg.de

1 **Summary**
2
Many animal and plant species respond to high or low population densities by phenotypic

plasticity. To investigate if specific age classes and/or cross-generational signaling affect(s)

phenotypic plasticity, we developed a dye-based method to differentiate co-occurring nematode

age classes. We applied this method to *Pristionchus pacificus*, which develops a predatory

mouth form to exploit alternative resources and kill competitors in response to high population

densities. Remarkably, only adult, but not juvenile, crowding induces the predatory morph in

other juveniles. Profiling of secreted metabolites throughout development with HPLC-MS

combined with genetic mutants traced this result to the production of adult-specific pheromones.

Specifically, the *P. pacificus*-specific di-ascaroside#1 that induces the predatory morph exhibits

a binary induction in adults, even though mouth form is no longer plastic in adults. This cross-

generational signaling between adults and juveniles may serve as an indication of rapidly

increasing population size. Thus, phenotypic plasticity depends on critical age classes.

25 **Keywords:** Phenotypic plasticity, population density, *Pristionchus pacificus*, *Caenorhabditis*

26 *elegans,* nematode derived modular metabolites (NDMMs)

Introduction

Population density is an important ecological parameter, with higher densities corresponding to increased competition for resources (Hastings, 2013). In addition to density-dependent selection (MacArthur, 1962; Travis et al., 2013), which operates on evolutionary time scales, some organisms can respond dynamically to population density through phenotypic plasticity. For example, plants can sense crowding by detecting the ratio of red (chlorophyll absorbing) to far red (non-absorbing) light, and respond by producing higher shoots (Dudley and Schmitt, 2015). Locusts undergo solitary to swarm (i.e. gregarious) transition, and aphids can develop wings, both as results of increased physical contact (Pener and Simpson, 2009; Simpson et al., 2001; Slogget and Weisser, 2004). Intriguingly, population density can also have cross-generational effects. For example, adult crowding of the desert locust *Schistocerca gregaria* (Maeno and Tanaka, 2008; Simpson and Miller, 2007) and migratory locust *Locusta migratoria* (Chen et al., 2015; Hamouda et al.) also influences the egg size, number, and morphology of their progeny; and high population densities of red squirrels elicit hormonal regulation in mothers to influence faster-developing offspring (Ben Dantzer et al., 2013). In many species, population density and cross-generational signaling are detected through pheromones, however the precise nature, mechanisms of induction, age-specificity, and exact ecological role are not well understood.

Nematodes are a powerful model system to investigate the mechanisms of density-dependent plasticity because many small molecule pheromones that affect plastic phenotypes have been characterized (Butcher, 2017; Butcher et al., 2007; Reuss et al., 2012). For example, in the model organism *Caenorhabditis elegans,* high population densities induce entry into a stress-resistant dormant 'dauer' stage (Fielenbach and Antebi, 2008). The decision to enter dauer was revealed to be regulated by a family of small molecule nematode-derived modular metabolites (NDMMs) called ascarosides that act as pheromones (Butcher et al., 2007; 2008; Jeong et al., 2005). Ascarosides consist of an ascarylose sugar with a fatty acid side chain and

modular head and terminus groups (Figure 1A). The level and composition of ascarosides were

later shown to be dependent on sex (Chasnov et al., 2007; Izrayelit et al., 2012) and

development (Kaplan et al., 2011), although it is thought that dauer can be induced by all

developmental stages (Golden and Riddle, 1982). Subsequent studies revealed that specific

NDMMs also regulate other life history traits, such as mating (Chasnov et al., 2007; Izrayelit et

al., 2012), social behavior (Srinivasan et al., 2012) and developmental speed (Ludewig et al.,

2017). Although NDMMs are broadly conserved (Choe et al., 2012; Dong et al., 2018; Markov et

al., 2016), inter- and intraspecific competition have driven the evolution of distinct response

regimes for the same phenotypes (Bose et al., 2014; Choe et al., 2012; Diaz et al., 2014; Falcke

et al., 2018; Greene et al., 2016). In addition, more complex structures have been observed that

affect distinct plastic phenotypes (Bose et al., 2012).

In *Pristionchus pacificus,* a soil-associated nematode that is reliably found on scarab

beetles (Figure 1A)(Herrmann et al., 2006; 2007; Sommer and McGaughran, 2013), an

ascaroside dimer (dasc#1) that is not found in *C. elegans* regulates the development of a

predatory mouth form (Bento et al., 2010a; Bose et al., 2012; Sommer et al., 2017). Mouth-form

plasticity represents an example of a morphological novelty that results in predatory behavior to

exploit additional resources and kill competitors. Specifically, adult *P. pacificus* exhibit either a

narrow stenostomatous (St) mouth (Figure 1B), which is restricted to bacterial feeding, or a wide

eurystomatous (Eu) mouth with an extra denticle (Figure 1C), which allows for feeding on

bacteria and fungi (Sanghvi et al., 2016), and predation on other nematodes (Wilecki et al.,

2015). This type of phenotypic plasticity is distinct from direct *vs.* indirect (dauer) development

because it results in two alternative life history strategies in the adult (for review see Sommer &

Mayer, 2015). Recent studies in *P. pacificus* have begun to investigate the dynamics and

succession of nematodes on decomposing beetle carcasses to better understand the ecological

significance of mouth-form plasticity (Meyer et al., 2017). These studies revealed that on a

carcass (Figure 1D), *P. pacificus* exits the dauer diapause to feed on microbes, and then re-

enters dauer after food sources have been exhausted, displaying a 'boom-and-bust' ecology

(Meyer et al., 2017; Sommer and McGaughran, 2013). Presumably different stages of this

succession comprise different ratios of juveniles and adults, and recognizing the age-structure

of a population as a juvenile could provide predictive value for adulthood. However, it is

unknown whether the mouth-form decision is sensitive to crowding by different age classes.

More broadly, while age classes are known to be important for population growth and density-

dependent selection {Hastings:2013dn, Charlesworth:1994ww, Charlesworth:1970ks}, their role

in phenotypic plasticity has thus far been largely unexplored.

While nematodes have many experimental advantages, including easy laboratory culture

and advanced genetic, genomic, and the aforementioned chemical tools, their small size has

made investigations at the organismal level and in experimental ecology challenging. For

example, no *in vivo* methodologies are currently available to label distinct populations without

the need for transgenics, which is only available in select model organisms such as *C. elegans*,

P. pacificus, and some of their relatives. Here, we combine a novel dye-staining method with the

first developmental pheromone profiling in *P. pacificus* to study potential effects of age on

density-dependent plasticity. This vital-dye method allows tracking adults with juveniles, or

juveniles with juveniles, and can be applied to any nematode system that can be cultured under

laboratory conditions. In contrast to dauer, we found that mouth form is strongly affected by

cross-generational signalling. Specifically, only adult crowding induces the predatory morph,

which is controlled by adult-specific pheromones.

Results

A vital dye method for labeling nematode populations

To directly test if different age groups of nematodes influence plastic phenotypes, we required

two synchronized populations to co-habit the same space, yet still be able to identify worms

from different age groups. To do so, we developed a dye-staining methodology to robustly

differentiate between nematode populations. After trying several vital dyes, we identified that

Neutral Red (Thomas and Lana, 2008) and CellTracker Green BODIPY (Thermo) stain

nematode intestines brightly and specifically to their respective channels (Figures 2A-E and S1).

These dyes stain all nematodes tested including *C. elegans* (Figure S2) and dauer larvae

(Figure S3A,B). They also last more than three days (Figure S3C-G), allowing long-term

tracking of mixed nematode populations. Importantly, neither Neutral Red nor CellTracker

Green staining affects viability, developmental rate, or the formation of specific morphological

structures, such as *P. pacificus* mouth form (Figure S4). Thus, Neutral Red and CellTracker

Green allow specific labeling of worm populations to study age-dependent effects on

phenotypes.

Adult but not juvenile crowding induces the predatory mouth form in *P. pacificus*

To assess potential intra- or inter-generational influence on *P. pacificus* mouth form we stained

juveniles of the highly St strain RSC017 with Neutral Red, and added an increasing number of

CellTracker Green-stained RSC017 adults or juveniles (Figure 2F, 3A). Three days later we

phenotyped red animals that had developed into adults, but showed no green staining. To

ascertain potential differences between adding juveniles or adults, we performed a binomial

regression on Eu count data from multiple independent biological replicates (n>3), with age and

number of individuals added as fixed effects (Transparent Methods, Table S1). We observed a

significant increase in Eu worms in response to adults, but not juveniles (p=2.59 x10^{-2}; for

display summed percents are shown in Figure 3B,C). Almost half (48%) of the population

developed the Eu mouth form with just 500 adult animals, which is a greater than 50-fold

induction compared to side-by-side controls (Figure 3B,C). We were also curious if dauers,

which have a thickened cuticle and represent a distinct stage in the boom-and-bust life cycle of

nematodes, could still respond to adults. Indeed, the same trend that was observed with

juveniles was seen with dauers (p=2.96x10^{-3}), albeit to a more muted extent (Figure 3D,E).

Specifically, with a total of 200 dauers and 500 adults, 25.7% of dauers become Eu, whereas

only 1.8% of dauers become Eu on a plate containing 700 dauers (and no adults) (Figure 3D).

Collectively, these data indicate that adult crowding specifically induces the Eu mouth form.

Even though we did not detect a mouth-form switch in large populations of J2s or

dauers, and food was still visible on plates containing the most animals (500 adults and 200

juveniles), we could not completely rule out the possible effect of food availability on mouth

form. As a proxy for starvation, we conducted assays with greatly increased numbers of

juveniles from 1,000 to 10,000 that would rapidly deplete bacterial food. We noticed a stark cliff

in the fraction of juveniles that reach adulthood at 4,000-5,000 animals, arguing that food is a

limiting resource at this population density (Figure 3F). Importantly however, in these plates we

still did not see a shift in mouth form (Figure 3G) (p=0.99, binomial regression, Table S1). With

an overwhelming 10,000 worms on a plate, 5.8% were Eu, compared to 48% in the presence of

only 500 adults. While longer-term starvation may yet have an impact on mouth form, under our

experimental conditions it appears to be negligible.

Adult but not juvenile secretions induce the Eu mouth form

As mouth-form plasticity in *P. pacificus* is regulated by nematode-derived modular metabolites

(NDMMs)(Bose et al., 2012), we wondered if the difference between adults and juveniles

resulted from differences in secreted NDMMs. To test this hypothesis we added secretions from

adult or juvenile worms to RSC017 (highly St) juveniles. We found that adult secretions from

both the laboratory stain RS2333 (highly Eu) and RSC017 led to a significant increase in the Eu

morph relative to juvenile secretions (p=5.27x10^{-06}, 1.33x10^{-3}, respectively, Fisher's exact

test)(Figure 4). To confirm the effect was caused by ascaroside pheromones, we exposed

RSC017 juveniles to supernatant from a *daf-22.1;22.2* double mutant, which exhibits virtually no

ascaroside production in both *C. elegans* and *P. pacificus* (Golden and Riddle, 1985; Markov et

al., 2016). Again, juvenile secretion had no impact on Eu frequency, but in contrast to wild-type

supernatants, we observed no significant increase in Eu frequency with adult secretions

(p=0.8324, Fisher's exact test, Figure 4). Thus, adult-specific NDMMs induce development of

the Eu mouth form.

Developmental-staged NDMM profiles reveal adult-specific synthesis of dasc#1

Next, we investigated whether the difference between adult and juvenile pheromones is one of

dosage, or of identity. To answer this question and verify age-specific differences in

pheromones, we profiled *P. pacificus* NDMM levels in two strains and at three time points

throughout development. We used RS2333 and RSC017 and measured the exo-metabolomes

of juvenile stage 2 (J2s, 24 hrs), J3s (48 hrs) and J4/adults (72 hrs) from a constant culture with

excess OP50 bacterial food (Figures 5A,B, S5, Materials and methods). To assess potential

differences in pheromone levels we performed a linear regression with the area under the curve

for each NDMM (aoc) (Figure S5) as the response variable. Stage and strain were modeled as

fixed effects, and because we performed separate regression analyses for each pheromone, we

adjusted the resulting p values for multiple testing using false discovery rate (FDR)(see Table

S2 for p and *FDR* values between stage and strain). We observed that there was a significant

affect of developmental stage on the levels of ascr#9, ascr#12, npar#1, and dasc#1, and that

ubas#1 and #2 are strain and stage specific (FDR<0.05). Interestingly, dasc#1 is the most

potent known Eu-inducing compound when tested as a single synthesized compound, while

npar#1 is both Eu- and dauer-inducing (Figure 5C,D,F-I) (Bose et al., 2012). Closer inspection

revealed dasc#1, npar#1, and ascr#9 increase throughout development in both strains, and

dasc#1 peaks in adults in RS2333 (p<0.05, student's two-tailed *t*-test between 72 hrs and 24

hrs for each NDMM in both strains, and also 72 hrs and 48 hrs for dasc#1 in RS233, Table S3).

Intriguingly, the trajectory of dasc#1 appeared binary in both strains (Figure 5F,G). In fact our

statistical model for dasc#1 fits better if we assume cubic rather than linear growth

(ΔAIC=3.958). In contrast, ascr#9, which was also statistically up-regulated but does not affect

known plastic phenotypes (Bose et al., 2012), displays a more gradual increase in both strains

(Figure 5E,J,K), and the model fits better with linear growth ($AIC_{linear} - AIC_{cubic}$= -1.208).

Meanwhile, the trajectory of npar#1 appears strain-specific (Figure 5H,I). Hence the mode of

induction is NDMM-specific, and the kinetics of production may be related to their roles in

phenotypic plasticity.

In principle, the increase in abundance of certain pheromones could be a result of a

concomitant increase in body mass, however several observations indicate more targeted

regulation. First, no other compounds were significantly different in our linear model. Second, an

analysis of previously published RNA-seq data (Baskaran et al., 2015) reveals the increase in

NDMM abundance corresponds to an increase in transcription of the thiolase *Ppa-daf-22.1*

(Figure S6), the most downstream enzyme in the β-oxidation pathway of ascaroside synthesis.

Third, pasc#9 and pasc#12 actually exhibit a peak in abundance at the 48 hr/J3 time point,

rather than in 72 hrs/adults. Finally, we profiled the endo-metabolome of eggs, and found

appreciable amounts of ascr#1, #9, #12, and pasc#9, but little to no traces of other ascaroside

derivatives (Figure S5C), suggesting age-specific synthesis rather than release. Together, these

results suggest that the observed increase in ubas#1 and #2, ascr#9, npar#1, and dasc#1 over

time corresponds to age-specific production. The observation that dasc#1 is produced

specifically during the juvenile-to-adult transition is especially intriguing because adults are no

longer able to switch mouth forms, hinting at cross-generational signaling.

Discussion

Here, we introduce a novel dye-based method that allowed us to assess cross-generational

influence on mouth form. Our results demonstrate adult crowding induces the Eu predatory

morph, and that this effect is a result of age-specific pheromones. In doing so, we provide the

first multi-stage time series of pheromone production in *P. pacificus*, which shows that dasc#1

exhibits a surprising 'off-on' switch-like induction pattern. Collectively, our results argue that

adults represent a critical age group (Charlesworth, 1972) in nematode populations.

Our developmental profiling revealed an increase in two NDMMs that affect plastic

phenotypes. The observation that this trend mirrors the transcriptional regulation of enzymes

involved in NDMM synthesis argues that the stage-dependent increase is not simply a result of

an increase in body mass, but rather that these molecules are programmed for stage-specific

induction. The binary 'off-on' kinetics might reflect a population level feedback loop, such that

the production of density-sensing pheromones is based on a threshold level of previously

produced pheromones. It is also worth noting that while npar#1 is the major dauer-inducing

pheromone in *P. pacificus* (Bose et al., 2012), we did not observe dauers in any experimental

setup described herein. Thus, it seems that mouth-form phenotype is the first-level plastic

response to population density. Presumably higher concentrations are required for dauer

induction, reflecting a calculated response strategy depending on the level of crowding or

duration of starvation. Interestingly, the effect of adult supernatants was noticeably less (23%-

26% Eu) than of adult worms (up to 48% with only 500 adults). It is difficult to compare the

amount of pheromone concentrations between experiments, but presumably worms in the vital-

dye assay experienced a greater local concentration as they were in direct physical contact with

each other, compared to worms in the supernatant assay.

Among the many environmental influences on mouth form (Werner et al., 2017),

population density and starvation are perhaps the most ecologically relevant. However, teasing

apart these two factors has proven difficult (Bento et al., 2010b). Here, we demonstrate that

while a strong shift is observed with adult-specific pheromones, no such effect was seen under

limited resource conditions. Thus, age-specific crowding is sufficient to induce the Eu mouth

form. Nevertheless, this does not preclude that long-term starvation could also have an effect.

Determining the relative contributions of these factors to mouth form will be important to better

understand the sophisticated ecological response strategies of *P. pacificus*, nematodes, and

phenotypic plasticity in general.

Why do adults and not juveniles affect mouth form? Given that St animals can develop

faster (Serobyan et al., 2013), there may be a 'race' to sexual maturation in emergent

populations at low densities. But as the nematode population increases, there will likely be a

commensurate decrease in bacterial populations. When faced with competition from other

nematodes, *P. pacificus* has a particular advantage in developing the Eu morph; their expanded

dietary range includes their competition. Indeed, when nematode prey is the only available food

source, the Eu morph provides longer life spans and more progeny than the St morph

(Serobyan et al., 2014). When resources become depleted as population size increases, *C.*

elegans and other monomorphic nematodes may enter dauer and disperse (Frézal and Félix,

2015). But in St-biased dimorphic strains of *P. pacificus*, juveniles may switch to the Eu morph

in response to adults as a first-level indication of rapidly increasing population size (Figure 6).

Then, after prolonged starvation and crowding, worms will presumably enter dauer. By analogy

to economic models of population growth (Malthus et al., 1992; Trewavas, 2002) mouth-form

plasticity is a 'technological innovation' to temporarily escape a Malthusian resource trap. To

what extent this occurs in nature, or with *P. pacificus* strains that are highly Eu, remains to be

determined.

The evolution of dimorphic mouth forms is one among myriad nematode ecological

strategies. For example, entomopathogenic nematodes release their symbiont bacteria in insect

hosts to establish their preferred food source, and some release antibiotics to kill off competing

bacteria and fungi from other entomopathogenic species (Griffin, 2012). Some free-living

species, like those of the genus *Oscheius*, refrain from combat and stealthily feed and

reproduce amidst warring entomopathogenic species. Interspecific killing also occurs in

gonorchoristic species, in which both mated and virgin males are killed, implying fighting not just

for mates but for resources as well (O'Callaghan et al., 2014; Zenner et al., 2014). Reproductive

strategies also exist, and hermaphroditic species have an advantage over gonachristic species

when colonizing a new niche, such as an insect carcass (Campos-Herrera, 2015). Meanwhile

insect hosts and colonizing nematodes have their own distinct pheromone-based attraction and

toxicity (Cinkornpumin et al.; Renahan and Hong, 2017). Finally, the renaissance of *C. elegans*

sampling from around the world (Cook et al., 2017; Evans et al., 2016; Félix et al., 2013;

Petersen et al., 2014; Poullet and Braendle, 2015) is rapidly building a resource of wild isolates

that will almost certainly have different and fascinating ecologies. We hope our method for

labeling and then combining different nematode populations on the same plate will aid in studies

to identify these strategies. Perhaps the time is also ripe to complement these studies with more

sophisticated ecological modelling that can lead to testable hypotheses.

Although beyond the scope of this manuscript, the cross-generational communication we

observed could in principle reflect an intended signal from adults to juveniles, i.e. kin selection

(Bourke, 2014). However, we favor a more simplistic view that juveniles have evolved to

recognize adult-produced metabolites. Regardless of these interpretations, our results argue

that age classes are a critical factor in density-dependent plasticity, as has been theorized in

density-dependent selection (Charlesworth, 1994).

Limitations of the Study

Given the ubiquity of certain traits in reproductive adults and their contribution to population

growth, we suspect similar results will be found in other systems. However, it may depend on

the phenotype and system being studied. For example, the population dynamics of nematodes

(fast hermaphroditic reproduction) may be sufficiently different from other species such that our

findings are not extendable in every case. In addition, our method of staining different

populations, while fast and easy, is particular to nematodes.

Methods

*See Transparent Methods in Supplemental Information

Acknowledgements

We would like to thank all members of the Sommer lab, Dr. Talia Karasov, Dr. Hernan Burbano

and Moises Exposito-Alonso for guidance with statistical analysis, and Dr. Adrian Striet (Max

Planck Institute), and Dr. Cameron Weadick (University of Sussex) for thoughtful critique and

discussion.

Author Contributions

MSW and RJS conceived of the project. MC conducted pheromone profiling. MSW and TR

designed and conducted dye-labeling experiments. TR and MC performed supernatant

experiments. MD and MSW considered ecological implications. MSW and TR wrote the

manuscript with input and edits from all authors.

References

Baskaran, P., Rödelsperger, C., Prabh, N., Serobyan, V., Markov, G.V., Hirsekorn, A., and
Dieterich, C. (2015). Ancient gene duplications have shaped developmental stage-specific
expression in Pristionchus pacificus. BMC Evol. Biol. *15*, 185.

Ben Dantzer, Newman, A.E.M., Boonstra, R., Palme, R., Boutin, S., Humphries, M.M., and
McAdam, A.G. (2013). Density Triggers Maternal Hormones That Increase Adaptive Offspring
Growth in a Wild Mammal. Science *340*, 1235765–1217.

Bento, G., Ogawa, A., and Sommer, R.J. (2010a). Co-option of the hormone-signalling module
dafachronic acid–DAF-12 in nematode evolution. Nature *466*, 494–497.

Bento, G., Ogawa, A., and Sommer, R.J. (2010b). Co-option of the hormone-signalling module
dafachronic acid–DAF-12 in nematode evolution. Nature *466*, 494–497.

Bose, N., Meyer, J.M., Yim, J.J., Mayer, M.G., Markov, G.V., Ogawa, A., Schroeder, F.C., and
Sommer, R.J. (2014). Natural Variation in Dauer Pheromone Production and Sensing Supports
Intraspecific Competition in Nematodes. Current Biology *24*, 1536–1541.

Bose, N., Ogawa, A., Reuss, von, S.H., Yim, J.J., Ragsdale, E.J., Sommer, R.J., and
Schroeder, F.C. (2012). Complex Small-Molecule Architectures Regulate Phenotypic Plasticity
in a Nematode. Angewandte Chemie International Edition *51*, 12438–12443.

Bourke, A.F.G. (2014). Hamilton's rule and the causes of social evolution. Philosophical
Transactions of the Royal Society of London B: Biological Sciences *369*, 20130362–20130362.

Butcher, R.A. (2017). Small-molecule pheromones and hormones controlling nematode
development. Nature Chemical Biology *13*, 577–586.

Butcher, R.A., Fujita, M., Schroeder, F.C., and Clardy, J. (2007). Small-molecule pheromones
that control dauer development in Caenorhabditis elegans. Nature Chemical Biology *3*, 420–
422.

Butcher, R.A., Ragains, J.R., Kim, E., and Clardy, J. (2008). A potent dauer pheromone
component in Caenorhabditis elegans that acts synergistically with other components. Proc Natl
Acad Sci USA *105*, 14288–14292.

Campos-Herrera, R. (2015). Nematode Pathogenesis of Insects and Other Pests (Cham:
Springer).

Charlesworth, B. (1994). Evolution in age-structured populations.

Charlesworth, B. (1972). Selection in populations with overlapping generations. III. Conditions
for genetic equilibrium. Theoretical Population Biology.

Chasnov, J.R., So, W.K., Chan, C.M., and Chow, K.L. (2007). The species, sex, and stage
specificity of a Caenorhabditis sex pheromone. Proc Natl Acad Sci USA *104*, 6730–6735.

Chen, B., Li, S., Ren, Q., Tong, X., Zhang, X., and Le Kang (2015). Paternal epigenetic effects
of population density on locust phase-related characteristics associated with heat-shock protein
expression. Molecular Ecology *24*, 851–862.

Choe, A., Reuss, von, S.H., Kogan, D., Gasser, R.B., Platzer, E.G., Schroeder, F.C., and
Sternberg, P.W. (2012). Ascaroside Signaling Is Widely Conserved among Nematodes. Current
Biology *22*, 772–780.

Cinkornpumin, J.K., Wisidagama, D.R., Rapoport, V., elife, J.G., 2014 A host beetle pheromone
regulates development and behavior in the nematode Pristionchus pacificus.
Cdn.Elifesciences.org

.

Cook, D.E., Zdraljevic, S., Roberts, J.P., and Andersen, E.C. (2017). CeNDR, the
Caenorhabditis elegans natural diversity resource. Nucleic Acids Res. *45*, D650–D657.

Diaz, S.A., Brunet, V., Lloyd-Jones, G.C., Spinner, W., Wharam, B., and Viney, M. (2014).
Diverse and potentially manipulative signalling with ascarosides in the model nematode C.
elegans. BMC Evol. Biol. *14*, 46.

Dong, C., Reilly, D.K., Bergame, C., Dolke, F., Srinivasan, J., and Reuss, von, S.H. (2018).
Comparative Ascaroside Profiling of CaenorhabditisExometabolomes Reveals Species-Specific
(ω) and (ω – 2)-Hydroxylation Downstream of Peroxisomal β-Oxidation. J. Org. Chem. *83*,
7109–7120.

Dudley, S.A., and Schmitt, J. (2015). Testing the Adaptive Plasticity Hypothesis: Density-
Dependent Selection on Manipulated Stem Length in Impatiens capensis. The American
Naturalist *147*, 445–465.

Evans, K.S., Zhao, Y., Brady, S.C., Long, L., McGrath, P.T., and Andersen, E.C. (2016).
Correlations of Genotype with Climate Parameters Suggest Caenorhabditis elegans Niche
Adaptations. G3: Genes, Genomes, Genetics *7*, g3.116.035162–g3.116.035298.

Falcke, J.M., Bose, N., Artyukhin, A.B., Rödelsperger, C., Markov, G.V., Yim, J.J., Grimm, D.,
Claassen, M.H., Panda, O., Baccile, J.A., et al. (2018). Linking Genomic and Metabolomic
Natural Variation Uncovers Nematode Pheromone Biosynthesis. Cell Chem Biol *25*, 787–
796.e12.

Félix, M.-A., Jovelin, R., Ferrari, C., Han, S., Cho, Y.R., Andersen, E.C., Cutter, A.D., and
Braendle, C. (2013). Species richness, distribution and genetic diversity of Caenorhabditis
nematodes in a remote tropical rainforest. BMC Evol. Biol. *13*, 10.

Fielenbach, N., and Antebi, A. (2008). C. elegans dauer formation and the molecular basis of
plasticity. Genes & Development *22*, 2149–2165.

Frézal, L., and Félix, M.-A. (2015). The Natural History of Model Organisms: C. elegans outside
the Petri dish. Elife *4*, e05849.

Golden, J.W., and Riddle, D.L. (1982). A pheromone influences larval development in the
nematode Caenorhabditis elegans. Science *218*, 578–580.

Golden, J.W., and Riddle, D.L. (1985). A gene affecting production of the Caenorhabditis
elegans dauer-inducing pheromone. Mol. Gen. Genet. *198*, 534–536.

Greene, J.S., Brown, M., Dobosiewicz, M., Ishida, I.G., Macosko, E.Z., Zhang, X., Butcher, R.A.,
Cline, D.J., McGrath, P.T., and Bargmann, C.I. (2016). Balancing selection shapes density-
dependent foraging behaviour. Nature *539*, 254–258.

Griffin, C.T. (2012). Perspectives on the behavior of entomopathogenic nematodes from
dispersal to reproduction: traits contributing to nematode fitness and biocontrol efficacy. J.
Nematol. *44*, 177–184.

Hamouda, A.B., and, S.T.O.E., 2011 Density-dependent phenotypic plasticity in body coloration
and morphometry and its transgenerational changes in the migratory locust, Locusta migratoria.
Journal of Entomology and Nematology.

Hastings, A. (2013). Population biology: concepts and models.

Herrmann, M., Mayer, W.E., and Sommer, R.J. (2006). Nematodes of the genus Pristionchus
are closely associated with scarab beetles and the Colorado potato beetle in Western Europe.
Zoology *109*, 96–108.

Herrmann, M., Mayer, W.E., Hong, R.L., Kienle, S., Minasaki, R., and Sommer, R.J. (2007). The
Nematode Pristionchus pacificus (Nematoda: Diplogastridae) Is Associated with the Oriental
Beetle Exomala orientalis (Coleoptera: Scarabaeidae) in Japan (Zoological Society of Japan).

Izrayelit, Y., Srinivasan, J., Campbell, S.L., Jo, Y., Reuss, von, S.H., Genoff, M.C., Sternberg,
P.W., and Schroeder, F.C. (2012). Targeted metabolomics reveals a male pheromone and sex-
specific ascaroside biosynthesis in Caenorhabditis elegans. ACS Chem. Biol. *7*, 1321–1325.

Jeong, P.-Y., Jung, M., Yim, Y.-H., Kim, H., Park, M., Hong, E., Lee, W., Kim, Y.H., Kim, K., and
Paik, Y.-K. (2005). Chemical structure and biological activity of the <i>Caenorhabditis
elegans</i> dauer-inducing pheromone. Nature *433*, 541–545.

Kaplan, F., Srinivasan, J., Mahanti, P., Ajredini, R., Durak, O., Nimalendran, R., Sternberg,
P.W., Teal, P.E.A., Schroeder, F.C., Edison, A.S., et al. (2011). Ascaroside Expression in
Caenorhabditis elegans Is Strongly Dependent on Diet and Developmental Stage. PLoS ONE *6*,
e17804.

Ludewig, A.H., Gimond, C., Judkins, J.C., Thornton, S., Pulido, D.C., Micikas, R.J., Döring, F.,
Antebi, A., Braendle, C., and Schroeder, F.C. (2017). Larval crowding accelerates C. elegans
development and reduces lifespan. PLoS Genet *13*, e1006717.

MacArthur, R.H. (1962). SOME GENERALIZED THEOREMS OF NATURAL SELECTION. Proc
Natl Acad Sci USA *48*, 1893–1897.

Maeno, K., and Tanaka, S. (2008). Maternal effects on progeny size, number and body color in
the desert locust, Schistocerca gregaria: Density- and reproductive cycle-dependent variation.
Journal of Insect Physiology *54*, 1072–1080.

Malthus, T.R., Winch, D., and James, P. (1992). Malthus: "An Essay on the Principle of
Population" (Cambridge University Press).

Markov, G.V., Meyer, J.M., Panda, O., Artyukhin, A.B., Claaßen, M., Witte, H., Schroeder, F.C.,
and Sommer, R.J. (2016). Functional Conservation and Divergence of daf-22 Paralogs in
Pristionchus pacificus Dauer Development. Mol Biol Evol *33*, 2506–2514.

Meyer, J.M., Baskaran, P., Quast, C., Susoy, V., Rödelsperger, C., Glöckner, F.O., and
Sommer, R.J. (2017). Succession and dynamics of Pristionchus nematodes and their
microbiome during decomposition of Oryctes borbonicus on La Réunion Island. Environmental
Microbiology *19*, 1476–1489.

O'Callaghan, K.M., Zenner, A.N.R.L., Hartley, C.J., and Griffin, C.T. (2014). Interference
competition in entomopathogenic nematodes: male Steinernema kill members of their own and
other species. International Journal for Parasitology *44*, 1009–1017.

Pener, M.P., and Simpson, S.J. (2009). Locust Phase Polyphenism: An Update (Elsevier).

Petersen, C., Dirksen, P., Prahl, S., Strathmann, E.A., and Schulenburg, H. (2014). The
prevalence of Caenorhabditis elegans across 1.5 years in selected North German locations: the
importance of substrate type, abiotic parameters, and Caenorhabditis competitors. BMC Ecol.
14, 4.

Poullet, N., and Braendle, C. (2015). Sampling and Isolation of C. elegans from the Natural
Habitat. Methods Mol. Biol. *1327*, 221–229.

Renahan, T., and Hong, R.L. (2017). A species-specific nematocide that results in terminal
embryogenesis. J. Exp. Biol. *220*, jeb.159665–3247.

Reuss, von, S.H., Bose, N., Srinivasan, J., Yim, J.J., Judkins, J.C., Sternberg, P.W., and
Schroeder, F.C. (2012). Comparative metabolomics reveals biogenesis of ascarosides, a
modular library of small-molecule signals in C. elegans. J. Am. Chem. Soc. *134*, 1817–1824.

Sanghvi, G.V., Baskaran, P., Röseler, W., Sieriebriennikov, B., Rödelsperger, C., and Sommer,
R.J. (2016). Life History Responses and Gene Expression Profiles of the Nematode
Pristionchus pacificus Cultured on Cryptococcus Yeasts. PLoS ONE *11*, e0164881.

Serobyan, V., Ragsdale, E.J., and Sommer, R.J. (2014). Adaptive value of a predatory mouth-
form in a dimorphic nematode. Proceedings of the Royal Society of London B: Biological
Sciences *281*, 20141334–20141989.

Serobyan, V., Ragsdale, E.J., Müller, M.R., and Sommer, R.J. (2013). Feeding plasticity in the
nematode Pristionchus pacificusis influenced by sex and social context and is linked to
developmental speed. Evolution & Development *15*, 161–170.

Simpson, S.J., Despland, E., Hägele, B.F., and Dodgson, T. (2001). Gregarious behavior in
desert locusts is evoked by touching their back legs. Proc Natl Acad Sci USA *98*, 3895–3897.

Simpson, S.J., and Miller, G.A. (2007). Maternal effects on phase characteristics in the desert
locust, Schistocerca gregaria: A review of current understanding. Journal of Insect Physiology
53, 869–876.

Slogget, J.J., and Weisser, W.W. (2004). Aphids in a New Millennium (INRA

).

Sommer, R.J., and Mayer, M.G. (2015). Toward a synthesis of developmental biology with
evolutionary theory and ecology. Annu. Rev. Cell Dev. Biol. *31*, 453–471.

Sommer, R.J., and McGaughran, A. (2013). The nematode Pristionchus pacificus as a model
system for integrative studies in evolutionary biology. Molecular Ecology *22*, 2380–2393.

Sommer, R.J., Dardiry, M., Lenuzzi, M., Namdeo, S., Renahan, T., Sieriebriennikov, B., and
Werner, M.S. (2017). The genetics of phenotypic plasticity in nematode feeding structures.
Open Biology *7*, 160332–118.

Srinivasan, J., Reuss, von, S.H., Bose, N., Zaslaver, A., Mahanti, P., Ho, M.C., O'Doherty, O.G.,
Edison, A.S., Sternberg, P.W., and Schroeder, F.C. (2012). A Modular Library of Small Molecule
Signals Regulates Social Behaviors in Caenorhabditis elegans. PLOS Biol *10*, e1001237.

Thomas, M.C., and Lana, P.D.C. (2008). Evaluation of vital stains for free-living marine
nematodes. Brazilian Journal of Oceanography *56*, 249–251.

Travis, J., Leips, J., and Rodd, F.H. (2013). Evolution in Population Parameters: Density-
Dependent Selection or Density-Dependent Fitness? The American Naturalist *181*, S9–S20.

Trewavas, A. (2002). Malthus foiled again and again. Nature *418*, 668–670.

Werner, M.S., Sieriebriennikov, B., Loschko, T., Namdeo, S., Lenuzzi, M., Dardiry, M.,
Renahan, T., Sharma, D.R., and Sommer, R.J. (2017). Environmental influence on Pristionchus
pacificus mouth form through different culture methods. Sci Rep *7*, 7207.

Wilecki, M., Lightfoot, J.W., Susoy, V., and Sommer, R.J. (2015). Predatory feeding behaviour
in Pristionchus nematodes is dependent on phenotypic plasticity and induced by serotonin. J.
Exp. Biol. *218*, 1306–1313.

Zenner, A.N.R.L., O'Callaghan, K.M., and Griffin, C.T. (2014). Lethal Fighting in Nematodes Is
Dependent on Developmental Pathway: Male-Male Fighting in the Entomopathogenic
Nematode Steinernema longicaudum. PLoS ONE *9*, e89385.

Figure Legends

Figure 1. Life cycle and developmental plasticity of the model nematode *Pristionchus*

***pacificus*.**

(A) The life cycle of *P. pacificus* consists of four juvenile stages (J1-4) until sexual maturation

(adults). Like many nematodes *P. pacificus* can enter a long-living 'dormant' dauer state that is

resistant to harsh environmental conditions. The decision to continue through the direct life

cycle or enter dauer is regulated by small molecule excreted ascarosides (chemical structure

adapted from (Butcher, 2017)). (B) *P. pacificus* can also adopt one of two possible feeding

structures; either a microbivorous narrow-mouth (stenostomatous, St), or (C) an omnivorous

wide-mouth (eurystomatous, Eu) with an extra tooth that can be utilized to kill and eat other

nematodes or fungi. White lines indicate the presence of an extra tooth (right side) in the Eu

morph or its absence in the St morph, and the dorsal tooth (left side), which is flint-like in St and

hook-like in Eu. White scale bar indicates 5 µM. (D), *P. pacificus* is often found in a necromenic

association with beetles (ex. shown here *Oryctes borbonicus,* photo taken by Tess Renahan) in

the dauer state, and resumes the free living life cycle upon beetle death to feed on the ensuing

microbial bloom.

Figure 2. Vital-dye method in nematodes allows mixing different populations together. (A-

E) *P. pacificus* were stained with either 0.005% Neutral Red or 50 µM CellTracker Green Bodipy

(Thermo) and viewed using Cy3 and FITC filters. Images were merged with Differential

interference contrast (DIC), scale bar = 100 µM. An example of relative intensities of each

fluorescence channel are displayed in the histograms (right) with the chemical structure of

Neutral Red or CellTracker Green Bodipy. (F) Age-dependent functional pheromone assay:

Experimental juveniles were stained with Neutral Red, and challenged with CellTracker Green

Bodipy-stained juveniles or adults on standard condition NGM agar plates seeded with 300 µl

OP50 *E. coli*. Three days later, only red-positive and green-negative adults were phenotyped.

Figure 3. Vital-dye method confirms adult-specific density effect on mouth form.

(A) The wild isolate RSC017 grown in standard conditions (5 young adults passed to fresh

plates, progeny phenotyped 4 days later) are highly stenostomatous (<10%, *n*=102). (B,C)

Mouth form ratios of Neutral Red-stained J2s, and (D,E) dauers, with increasing number of

CellTracker Green-stained competitors, as described in Figure 2 (n=3-5 independent biological

replicates per experiment, with total n>100 per experiment). Overall significance between strain

and age was determined by a binomial linear regression (see Transparent Methods), and pair-

wise comparisons were assessed by Fisher's exact test on summed Eu counts (Significance

codes: '***' < 0.001, '**' <0.01, '*' <0.05). Mouth forms were phenotyped at 40-100x on a Zeiss

Axio Imager 2 light microscope. (F) Percent reaching adulthood and percent Eu of those that

reached adulthood (G) after increasing numbers of J2s are added to standard 6 cm NGM agar

plates with 300 μl OP50 *E.coli* bacteria (n=2 biological replicates, with total *n*>200 for percent

reaching adulthood, and total n>100 for mouth form. Significance was determined by a binomial

regression).

Figure 4. Adult-specific secretions induce predatory morph in juveniles.

Highly St strain RSC017 juveniles were exposed to J2 and adult supernatants of its own strain,

and to the J2 and adult supernatants of highly Eu strain RS2333. Mouth form was phenotyped

three days later. Worms exposed to J2 secretions remained highly St, while worms exposed to

adult secretions had a small but significant increase in Eu morphs (p<0.05, Fisher's exact test).

Supernatants from the double mutant *daf-22.1/2*, which has deficient ascaroside pheromone

production, did not elicit juvenile or adult increase in Eu. Worms exposed to the S-media control

also remained highly St. n=4 independent biological replicates for RS2333 and *daf-22.1/2*

secretions, and n=2 independent biological replicates for RSC017 adult and juvenile secretions,

with an average count of 55 animals per replicate. For display, total Eu and St counts are

presented as percentages (Significance codes: '***' < 0.001, '**' <0.01, '*' <0.05).

Figure 5. Time-resolved Nematode-Derived Modular Metabolites (NDMMs) in *Pristionchus*

pacificus.

(A) Time resolved secretion profile of nematode derived modular metabolites from the wild-type

laboratory strain RS2333 and (B) wild isolate RSC017. Data is presented as the mean of 8

(RS2333) and 9 (RSC017) biological replicates, and error bars represent standard error of the

mean (SEM). (C-E) Chemical structures of adult specific NDMMs dasc#1, npar#1, and ascr#9,

as described in the Small Molecule Identifier Database (http://www.smid-db.org/), produced in

ChemDraw. (F-K) Time-resolved abundance of dasc#1, npar#1, and ascr#9 NDMMs in RS2333

and RSC017. Each data point represents a biological replicate, and for comparison to (A-B)

lines represent mean abundance. *P* values calculated by a 2-tailed students *t*-test (Significance

codes: '***' < 0.001, '**' <0.01, '*' <0.05).

Figure 6. Conceptual model of the role of critical age classes in mouth-form

phenotypic plasticity. Conceptual life cycle models of monomorphic or dimorphic mouth

form nematodes. In an isolated niche such as a decaying insect carcass, at some point

microbial food supplies will run out, leading to a Malthusian catastrophe. Nematodes escape

this trap by entering the dauer state, hitchhiking to a new insect carrier, and re-starting the

cycle. Dimorphic nematodes may sense the impending catastrophe earlier by recognizing an

abundance of adults in the population, and switching to the Eu morph to exploit new resources

and kill competitors. By analogy to economic models, the mouth form switch is a technological

innovation to temporarily escape a Malthusian resource trap.

Figure_1

A

t = 0

adult

embryo

fatty acid side chain

ascarylose sugar   terminus
                        groups

head
groups

J1

72 hrs

24 hrs

J4   J2

J3
'direct'

48 hrs

dauer

B St

C Eu

D

Figure 1. Life cycle and developmental plasticity of the model nematode *Pristionchus pacificus*.

Figure 2. Vital-dye method in nematodes allows mixing different populations together.

Figure_3

Figure 3. Vital-dye method confirms adult-specific density effect on mouth form.

Figure 4. Adult-specific secretions induce predatory morph in juveniles.

Figure 5. Time-resolved Nematode-Derived Modular Metabolites (NDMMs) in *Pristionchus pacificus*.

Figure 6. Conceptual model of the role of critical-age classes in mouth-form phenotypic plasticity.

Werner, et al. Adult influence on juvenile phenotypes by stage-specific pheromone production

---

Supplemental Information

Transparent Methods

Nematode strains and husbandry

P. pacificus Wild-type RS2333 (California) and RSC017 (La Réunion) strains were kept on 6 cm nematode growth media (NGM) plates seeded with OP50 and kept at 20°C. RSC017 is highly St and does not predate on other nematodes, and thus was used for biological assays instead of the highly Eu, predatory RS2333. To induce dauer, mixed-stage plates with little to no OP50 were washed with M9 and the resulting worm pellets were used in a modified 'White Trap' method. Worm pellets were placed on killed *Tenebrio molitor* grubs and dispersing dauers were collected in surrounding MilliQ water. Age of dauers ranged from one week to one month.

Dye staining

A stock solution of Neutral Red was prepared by dissolving 0.5 mg in 10 ml 5% acetic acid and stored at -20° C. Working solutions were prepared by 100x dilution in M9, aliquoted, stored at -20°C, and thawed directly before use. Working solutions were kept for approximately 1 month. Stock solutions of 10 mM Green Bodipy were made in DMSO and stored -20. J2s were prepared from 20-40 x 6 cm plates 6 days after passaging 5 worms to each plate on 300 µl OP50. Worms were washed from plates with M9 into a conical tube, and then filtered through 2x20 µM filters (Millipore) placed between rubber gaskets. The flow-through contains mostly J2 and some J3, which were pelleted by centrifugation, 8 seconds on a table-top eppdendorf centrifuge 5424, reaching approximately 10,000 x g. The juvenile pellet was then either re-

suspended in 1 ml Neutral Red working solution, or in 1 ml M9 and split to two tubes, then re-centrifuged, and then re-suspended in either 1 ml working solution Neutral Red (0.005% in M9) or 1 ml 50 µM Green BODIPY (Thermo) in M9. Tubes were then rotated for 3 hours in the dark, then washed by centrifugation as before, and re-suspended in 1 ml M9. This was repeated 3-4x until the dye was no longer visible in the worm pellet. Then, the concentration of worms was determined by aliquoting 2 µl onto a glass coverslip in 5 technical replicates, and counted under a dissecting microscope. Finally the appropriate number of animals was added to 6 cm plates that had been previously seeded with 300 µl OP50, and incubated at 20°C. After 3 days, 100% of worms exhibited Neutral Red staining (*n*=50, Supplementary figure 4). Dauers and J2s recovered after Neutral Red staining developed at the same developmental speed (3-4 days) and with the same mouth-form ratio as control worms recovered side-by-side (100% St for both, Supplementary figure 5, *n*=30). Dauers and J2s stained with Cell tracker Green BODIPY (50 µM) (Thermo) were similar, although less efficiently stained compared to Neutral Red. On day 4, 90% retained intestinal fluorescence (Supplementary figure 4), although brightness decreased with the number of days. Mouth-form ratios of dauers or J2s in +/- 50 µM Cell tracker Green BODIPY also developed at equivalent rates and mouth-form ratios (Supplementary figure 5). Lower than 25 µM did not yield strongly fluorescent worms after three hours. Cell Tracker Blue CMAC (Thermo) was also used at 50 µM and imaged 3 days post-staining for *P. pacificus*, and one day post-staining for *C. elegans*. However, due to the higher fluorescent background in the blue light spectrum in both *P. pacificus* and *C. elegans*, we performed all experiments using only Neutral Red and Cell tracker Green BODIPY.

Microscopy

All images were taken on a Zeiss Axio Imager 2 with an Axiocam 506 mono, and processed using Zen2 pro software. Image brightness and contrast were enhanced in ImageJ with a

minimum displayed value of 10 and maximum of 100 for all images in Fig 2, and Supplementary figures 4 and 5, and a minimum of 21 and maximum of 117 for Supplementary figure 3. The following exposure times were used for all images: Cy3 (peak emission = 561, exposure = 80 ms), FITC (peak emission = 519, exposure = 150 ms), Dapi (peak emission = 465, exposure = 80 ms), DIC (exposure = 80-140 ms).

Mixed culture experiments and statistical analysis

We performed mixed culture experiment presented in figure 2 with 3 to 5 independent biological replicates, and a minimum total number of counts $n > 100$ (median counts per replicate for J2=29 and the median counts per replicate for dauers=27). J2 or dauers were stained with Neutral Red as described in the 'Dye Staining' method section, then added to green-stained J2, dauer, or adult populations on 6 cm plates with 300 µl OP50 and incubated at 20° C. To ensure consistent bacterial food supply, we added 1 ml more overnight OP50-LB to each plate on the following day, then air-dried under a chemical fume hood for 1 hour, then returned to 20° C. On days 3-4, we phenotyped 'red' adults that exhibited no 'green' staining. To assess whether the age of the 'green' surrounding population affects the mouth form of the dependent variable 'red' J2s we performed a binomial regression on Eu counts (i.e. "successes") weighted by the number of counts per replicate and the number added as a fixed effect, using a generalized linear model from the standard statistical package in R:

glm(formula=cbind(Eu,total)~'stage_added' * '#_added', data='J2/Da', family="binomial"))

See Supplementary figure 6a for a table containing the resulting p values. The AIC for our models (78.97 for J2s and 89.59 for dauers) was substantially lower than the null hypothesis (220.16 for J2s and 147.29 for dauers), arguing a reasonable fit. For pair-wise comparisons of the effect of age for a given number of added animals, we performed a post-hoc Fisher's exact test on a contingency table containing the summed counts of Eu and St observations. For

display, we converted Eu counts into percent of total in figure 2, with the *p* values between the same number of animals added indicated over the adult-added population (Significance codes: 0 '***' 0.001, '**' 0.01, '*' 0.05).

Measuring the effect of food depletion on mouth form

To verify that starvation was not a factor in our mixed culture experiments, we added increasing number of J2s to standard 6 cm plates with 300 µl OP50 to rapidly consume bacterial food, and measured both the amount of animals that reached adulthood, and the percent Eu in each population for two biological replicates. To assess the affects of added J2s to each dependent variable we performed a binomial regression with count data weighted by the total number of counts for each replicate:

glm(formula = cbind(reached_adult, total)~thousand_J2s, data=data_2, family="binomial"))

p values indicate a significant difference in percent reaching adult as a function of J2s added, but not in percent Eu (Table S1).

Supernatant collection and assays

Strains RS2333, RSC017, and RS2333-*daf-22.1;22.2* were raised in 10 mL liquid culture as in the time-resolved NDMM collections (see below). For each time point, 9 mL of the supernatant was lyophilized overnight, extracted again overnight with 90% ethanol (diluted in Millipore water) while being stirred, and centrifuged (4000g, 10 min, 4°C). The solvent was evaporated and the solid re-dissolved with 1 mL Millipore water. This clear extract was then directly used for the assays. One mL of the supernatant was cleaned for HPLC-MS analysis (refer to pheromone profiling: HPLC-MS sample preparation) for quality control. For the assays, RSC017 was synchronized by bleaching and added to plates seeded with 300 µl OP50. The supernatants were added to the RSCO17 J2s in two 500 µl increments (for a total of 1ml supernatant) and

dried for 30 minutes in a sterile hood after each addition. Plates were kept at 20°C and adult mouth forms were screened three days later.

HPLC-MS sample preparation for normal exo-metabolome and time resolved analysis

To collect staged phermone profiles, we seeded 35 x 6 cm plates with 5 worms each, and bleached 5-6 days later when gravid to collect eggs/J1s. These were then seeded in 6 x 10 mL flasks with OP50 as described in Werner et al., 2017 (Werner et al., 2017). Then at 24, 48, or 72 hr time intervals, supernatant was obtained by centriguation (>4,000 x g, 4°C for 10 minutes). 1 mL supernatant was adsorbed onto a SPE-C8 cartridge (Thermo Scinetific Hypersept C8 100 mg/1mL), conditioned with 1 mL MeOH followed by 2 mL Millipore water. The adsorbed material was then washed with 200 uL water and subsequently eluted with 200 uL MeOH. This extract was then measured directly via HPLC-qTof MS (Bruker ImpactII).

HPLC-MS measurement

20 uL extract was injected into a Thermo ultimate 3000 HPLC equipped with a Sigma-Alderich Ascentis Express C18 2.7um 10mm x 4.6mm column at 20 °C with a flow of 500 uL/min. All MS measurements have been performed in negative ion mode and molecules are detected as [M-H]⁻ Ions. The solvent gradient started with 5 % acetonitrile (ACN)/ 95 % water (both containing 0.1 % formic acid) for 2 min. After this equilibration step, the ACN proportion has been increased to 65 % over 8 min, then to 100 % ACN in 1.2 min followed by a hold step for 8.8 min. Afterwards, they system was flushed to 5 % ACN with 2 min equilibration for a total of 22 min. For calibration, a sodium formiat cluster building solution has been automatically injected in the first 2 minutes of each run. Data analysis was performed with TASQ version 1.0 from Bruker Daltonics. Extracted ion chromatograms for each well-known compound with a mass width of 0.1 m/z and time slices of 0.5 min around the expected retention time have been produced after

calibrating and baseline correction. Assignment errors have been corrected with the provided MRSQ value.

Statistical analysis of NDMMs

NDMM levels were compared simultaneously between strains and developmental stages by a linear model in R: lm('NDMM' ~ 'developmental stage' * 'strain', data='data.frame')). *P* values between stages and strains were adjusted for multiple testing by a false discovery rate correction. The level of fit between linear vs. exponential growth was determined by the Akaike information criterion (AIC). The lowest AIC for iterations of different exponents (n=1,2,3…) was used for comparison to the simple linear model. While significant in both cases, for consistency we present the original *p* values from the original linear model in Supplementary Figure 8.

Figure S1, related to Figure 2. Vital dye staining of *Pristionchus pacificus.*

Figure S2, related to Figure 2. Vital dye staining of *Caenorhabditis elegans*.

Figure S3, related to Figure 2. Vital dye staining of *Pristionchus pacificus* dauers, and duration of staining.

Figure S4, related to Figure 2. Vital dye staining does not affect P. pacificus mouth form or development.

affect of population age on mouth form of developing juveniles

binomial regression	*p* value J2s	*p* value dauers
age added	0.0259	0.002955
number added	4.28e-13	0.000404

affect of number of peers on development and mouth form (proxy for potential starvation effects on mouth form)

binomial regression	*p* value for development (relative to 1,000)	*p* value for Eu (relative to 1,000)
3,000 J2s added	0.3408	1.0
4,000 J2s added	0.0424	1.0
5,000 J2s added	6.06E-14	0.99
10,000 J2s added	4.09E-14	0.99

Table S1, related to Figure 3. Table of binomial regression p values for vital-dye method and excess crowding.

Figure S5, related to Figure 5. Pheromone profiling quality control

NDMM comparison	pvalue	fdr corrected
ascr1_stage	0.4733	0.774490909
ascr1_strain	0.0429	0.110314286
ascr1_stage:strain	0.031	0.085846154
ascr9_stage	3.79E-05	0.0002274
ascr9_strain	0.651	0.778064516
ascr9_stage:strain	0.272	0.50148
ascr12_stage	0.0029	0.01404
ascr12_strain	0.0897	0.201825
ascr12_stage:strain	0.0302	0.085846154
dasc1_stage	9.62E-08	8.66E-07
dasc1_strain	0.11363	0.240628235
dasc1_stage:strain	0.00351	0.01404
npar1_stage	0.0033	0.01404
npar1_strain	0.9426	0.984
npar1_stage:strain	0.6355	0.778064516
npar2_stage	0.0516	0.12384
npar_2strain	0.984	0.984
npar2_stage:strain	0.9716	0.984
pasc1_stage	0.449	0.769714286
pasc1_strain	0.753	0.847125
pasc1_stage:strain	0.564	0.778064516
pasc9_stage	0.616	0.778064516
pasc9_strain	0.267	0.50148
pasc9_stage:strain	0.523	0.778064516
pasc12_stage	0.6122	0.778064516
pasc12_strain	0.2786	0.50148
pasc12_stage:strain	0.67	0.778064516
tasc1_stage	0.522	0.778064516
tasc1_strain	0.862	0.940363636
tasc1_stage:strain	0.57	0.778064516
ubas1_stage	3.13E-12	1.13E-10
ubas1_strain	0.00538	0.019368
ubas1_stage:strain	6.69E-08	8.03E-07
ubas2_stage	1.34E-11	2.41E-10
ubas2_strain	0.00711	0.023269091
ubas2_stage:strain	6.18E-07	4.45E-06

Table S2, related to Figure 5. Table of linear regression *p* values with *FDR* correction for strain and stage comparison of NDMM levels.

RS2333	dasc#1	npar#1	ascr#9
72 hrs compared to 24 hrs	5.7511E-07	3.4672E-05	0.00010345
72 hrs compared to 48 hrs	0.00571003	0.17579174	0.19705545
RSC017	dasc#1	npar#1	ascr#9
72 hrs compared to 24 hrs	0.02548973	0.00365597	0.02028689
72 hrs compared to 48 hrs	0.21178072	0.36578067	0.10414329

Table S3, related to Figure 5. Pairwise comparison of dasc#1, npar#1, and ascr#9 throughout development.

Figure S6, related to Figure 5. Enzyme that synthesize NDMMs is transcriptionally regulated during development.

Supplemental Figure Legends

Figure S1, related to Figure 2. Vital dye staining of *Pristionchus pacificus*.

(A) Control *P. pacificus* imaged with Cy3, FITC, and DAPI filters, and a merge with Differential Interference Contrast (DIC). Histogram on the right represents quantification of intensity with each filter. (B) Same as (A) but stained with 0.005% Neutral Red, (C), 50 µM CellTracker Green Bodipy (Thermo Fischer), or (D) 50 µM CellTracker Blue CMAC Dye (Thermo Fischer). J2s were stained (see Materials and Methods), and ensuing adult animals were imaged 3 days later on a Zeiss Axio Imager 2 with an Axiocam 506 mono, and processed using Zen2 pro software. Image brightness and contrast were enhanced in ImageJ for display, with a minimum displayed value of 10 and maximum of 100 for all images. Note that while Neutral Red and CellTracker Green staining are bright and specific to their respective channels, CellTracker Blue is indistinguishable from background fluorescence.

Figure S2, related to Figure 2. Vital dye staining of *Caenorhabditis elegans.*

(A-D) Same as Supplementary Figure 1, but with *C. elegans*.

Figure S3, related to Figure 2. Vital dye staining of Pristionchus pacificus dauers, and duration of staining. (A) Control *P. pacificus* dauer imaged with DIC, Cy3, and FITC filters. (B) Dauers stained with either 0.005% Neutral Red or 50 µM CellTracker Green Bodipy and imaged immediately after staining with DIC, Cy3, and FITC filters and merged with DIC. Images were taken using Zeiss Axio Imager 2 with an Axiocam 506 mono, processed using Zen2pro software, and adjusted in ImageJ, with a display value minimum of 21 and maximum of 117.

(C-G) 50 µM Cell Tracker Green Bodipy and 0.005% Neutral Red-stained J2s were imaged every day for five days. Percent of individuals retaining the dyes are shown in panels next to each microscope image for each day. Both stains are seen in all organisms for three days; Neutral Red (NR) persists for at least five, while the number of Green Bodipy (GB) –stained worms drops on day four. All images are merged with DIC, n=31 GB, 63 NR day 1, 68 GB, 56 NR day 2, 50 GB, 50 NR day 3, 50 GB, 50 NR day 4, 50 GB, 50 NR day 5.

Figure S4, related to Figure 2. Vital dye staining does not affect *P. pacificus* mouth form or development.

(A) Neutral Red and CellTracker Green Bodipy-stained J2s reach adulthood at the same rate as unstained J2s (3 days). (B) All of the J2s stained retain the dye in adulthood in the intestine. (C) Neither dye affects mouth form; both unstained and stained worms remain 100% St (n=30). (D-F) Same as for (A-C) except with dauers instead of J2s, and only with Neutral Red.

Table S1, related to Figure 3. Table of binomial regression *p* values for vital-dye method and excess crowding.

Significance *p* values from binomial regression of vital-dye method for age and number added, and from binomial regression of number-reaching-adult and Eu counts for each number of individuals added relative to 1,000 individuals added.

Figure S5, related to Figure 5. Pheromone-profiling quality control.

(A) Extracted ion traces (width 0.1 m/z) of 11 of the 12 NDMMs used in this publication from a seven-day mixed-stage sample, double peak of 247.12 m/z indicate isomeric structures (Part#9/Ascr#9). (B) Example of an averaged spectrum over a calibration segment, sodium-

formiat cluster building solution has been used to ensure high mass accuracy in each run. (C) Comparison of an endometabolome sample from a seven day mixed-stage cultured compared to the endometabolome of eggs, produced by using bleached eggs from 80 x 60 mm plates.

Table S2, related to Figure 5. Table of linear regression *p* values with FDR corrections for strain and stage comparison of NDMM levels. FDR-corrected and uncorrected *p* values from linear regression of *P. pacificus* NDMMs (alternating grey background between NDMMs for clarity). Red values indicate FDR<0.05.

Table S3, related to Figure 5. Pairwise comparison of dasc#1, npar#1, and ascr#9 throughout development. Significance assessed with a two-tailed student's *t*-test. Red values indicate *p*<0.05.

Figure S6, related to Figure 5. Enzyme that synthesize NDMMs is transcriptionally regulated during development. Comparison of *daf-22.1* and *daf-22.2* expression (FPKM) by RNA-seq through different stages of development, data from Baskaran et al., 2015. A two-sided students *t*-test was performed between J4-adults and J2s (Significance codes: 0 '***' 0.001, '**' 0.01, '*' 0.05).