A vital dye method in nematodes reveals age is a critical parameter in density-dependent plasticity

Michael S. Werner[1,2], Marc H. Claaßen[1,2], Tess Renahan[1,2], Mohannad Dardiry[1], and Ralf. J. Sommer[1*]

[1]Department of Evolutionary Biology, Max Planck Institute for Developmental Biology, 72076 Tübingen, Germany

[2]These authors contributed equally

*Corresponding author: ralf.sommer@tuebingen.mpg.de

Abstract

Population size is a major ecological constraint on resources, which affects density-dependent

selection over evolutionary time scales. Yet many species can also respond dynamically

through phenotypic plasticity. In *Caenorhabditis elegans* and other nematodes, high population

density drives juveniles into a dormant dauer stage. In *Pristionchus pacificus* high population

density also induces the development of a predatory mouth form. However, it is unknown if such

phenotypic switches respond simply to density, or to the density of specific age classes, as in

density-dependent selection. Although nematodes have many experimental advantages, their

small size has made investigating these and other experimental ecology questions challenging.

In particular, no *in vivo* methodologies are currently available to label distinct populations

without the need for transgenics. To study potential cross-generational signaling we developed

a novel dye-based method to differentiate nematode populations on the same plate, allowing us

to track adults with juveniles, or juveniles with juveniles. Surprisingly, only adult crowding

induces the predatory morph, even though adult mouth-forms are no longer plastic. We then

profiled secreted metabolites through development and traced this result to adult-specific

pheromones. Thus, a novel dye-based method reveals age as a critical parameter controlling

density-dependent plasticity.

Keywords: Phenotypic plasticity, population density, *Pristionchus pacificus*, *Caenorhabditis*

elegans, nematode derived modular metabolites (NDMMs)

1. Introduction

Population density is an important ecological parameter, with higher densities corresponding to increased competition for resources [1]. In addition to density-dependent selection [2,3], which operates on evolutionary time scales, some organisms can respond dynamically to population density through phenotypic plasticity. For example, plants can sense crowding by detecting the ratio of red (chlorophyll absorbing) to far red (non-absorbing) light, and respond by producing higher shoots [4]. Locusts undergo solitary to swarm (i.e. gregarious) transition, and aphids can develop wings, both as results of increased physical contact [5-7]. Intriguingly, population density can also have cross-generational effects. For example, adult crowding of the desert locust *Schistocerca gregaria* [8,9] and migratory locust *Locusta migratoria* [10,11] also influences the egg size, number, and morphology of their progeny; and high population densities of red squirrels elicit hormonal regulation in mothers to influence faster developing offspring [12]. In many species population density and cross-generational signaling are detected through pheromones, however the precise nature, mechanisms of induction, age-specificity and exact ecological role are not well understood.

Nematodes are a powerful model system to investigate the mechanisms of density-dependent plasticity because many small molecule pheromones that affect plastic phenotypes have been characterized [13-15]. For example, in the model organism *Caenorhabditis elegans,* high population densities induce entry into a stress resistant dormant 'dauer' stage [16]. The decision to enter dauer was revealed to be regulated by a family of small molecule nematode-derived modular metabolites (NDMMs) called ascarosides that act as pheromones[14,17,18]. Ascarosides consist of an ascarylose sugar with a fatty acid side chain and modular head and terminus groups (figure 1*a*). The level and composition of ascarosides was later shown to be dependent on sex [19,20] and developmental stage [21], although it is thought that dauer can be induced by all developmental stages [22]. Subsequent studies revealed that specific NDMMs

also regulate other life history traits, such as mating [19,20], social behavior [23] and

developmental speed [24]. Although NDMMs are broadly conserved [25-27], inter- and

intraspecific competition has driven the evolution of distinct response regimes in different strains

and species for the same phenotypes [25,28-31]. Additionally, more complex structures have

been observed that affect distinct plastic phenotypes [32].

In *Pristionchus pacificus,* a soil-associated nematode that is reliably found on scarab

beetles (figure 1*a*)[33-35], an ascaroside dimer (dasc#1) that is not found in *C. elegans*

regulates the development of a predatory mouth form [32,36,37]. Mouth-form plasticity

represents an example of a morphological novelty that results in predatory behavior to exploit

additional resources and kill competitors. Specifically, adult *P. pacificus* exhibit either a narrow

stenostomatous (St) mouth, which is restricted to bacterial feeding (figure 1*b*), or a wide

eurystomatous (Eu) mouth with an extra denticle (figure 1*c*), which allows for feeding on

bacteria and fungi [38], and predation on other nematodes [39]. This type of phenotypic

plasticity is distinct from direct *vs.* indirect (dauer) development because it results in two

alternative life history strategies in the adult (for review see Sommer & Mayer, 2015 [40]).

Recent studies in *P. pacificus* have begun to investigate the dynamics and succession of

nematodes on decomposing beetle carcasses to better understand the ecological significance

of mouth-form plasticity [41]. These studies revealed that on a carcass (figure 1*d*), *P. pacificus*

exits the dauer diapause to feed on microbes, and then re-enters dauer after food sources have

been exhausted, displaying a 'boom-and-bust' ecology [33,41]. Presumably different stages of

this succession comprise different ratios of juveniles and adults, and recognizing the age-

structure of a population as a juvenile could provide predictive value for adulthood. In locusts

density-dependent plasticity is affected by the age of crowded individuals [5], however age has

not yet been demonstrated to have a role in density-dependent plasticity in nematodes.

Apropos, whether the mouth-form decision is sensitive to population age structure has never

been investigated, neither experimentally, nor chemically by performing NDMM profiles of

multiple developmental stages.

While nematodes have many experimental advantages, including easy laboratory culture

and advanced genetic, genomic and the aforementioned chemical tools, their small size has

made investigations at the organismal level and in experimental ecology challenging. For

example, no *in vivo* methodologies are currently available to label distinct populations without

the need for transgenics, which is only available in selected model organisms such as *C.*

elegans, *P. pacificus* and some of their relatives. Here, we combine a novel dye-staining

method with the first developmental pheromone profiling in *P. pacificus* to study potential affects

of age on density-dependent plasticity. This vital-dye method allows tracking adults with

juveniles, or juveniles with juveniles, and can be applied to any nematode system that can be

cultured under laboratory conditions. In contrast to dauer, we found that mouth form is strongly

affected by cross-generational signalling. Specifically, only adult crowding induces the predatory

morph, which is controlled by adult-specific pheromones.

2. Materials and methods

(a) Nematode strains and husbandry

P. pacificus Wild-type RS2333 (California) and RSC017 (La Réunion) strains were kept on 6 cm

nematode growth media (NGM) plates seeded with OP50 and kept at 20°C. RSC017 is highly

St and does not predate on other nematodes, and thus was used for biological assays instead

of the highly Eu, predatory RS2333.

(b1) Pheromone profiling

HPLC-MS sample preparation for normal exo-metabolome and time resolved analysis

To collect staged phermone profiles, we seeded 35 x 6 cm plates with 5 worms each, and

bleached 5-6 days later when gravid to collect eggs/J1s. These were then seeded in 6 x 10 mL

flasks with OP50 as described in Werner et al., 2017 [42]. Then at 24, 48, or 72 hr time

intervals, supernatant was obtained by centriguation (>4,000 x g, 4°C for 10 minutes). 1 mL

supernatant was adsorbed onto a SPE-C8 cartridge (Thermo Scinetific Hypersept C8 100

mg/1mL), conditioned with 1 mL MeOH followed by 2 mL Millipore water. The adsorbed material

was then washed with 200 uL water and subsequently eluted with 200 uL MeOH. This extract

was then measured directly via HPLC-qTof MS (Bruker ImpactII).

(b2) HPLC-MS measurement

20 uL extract was injected into a Thermo ultimate 3000 HPLC equipped with a Sigma-Alderich

Ascentis Express C18 2.7um 10mm x 4.6mm column at 20 °C with a flow of 500 uL/min. All MS

measurements have been performed in negative ion mode and molecules are detected as [M-

H]$^-$ Ions. The solvent gradient started with 5 % acetonitrile (ACN)/ 95 % water (both containing

0.1 % formic acid) for 2 min. After this equilibration step, the ACN proportion has been

increased to 65 % over 8 min, then to 100 % ACN in 1.2 min followed by a hold step for 8.8 min.

Afterwards, they system was flushed to 5 % ACN with 2 min equilibration for a total of 22 min.

For calibration, a sodium formiat cluster building solution has been automatically injected in the

first 2 minutes of each run. Data analysis was performed with TASQ version 1.0 from Bruker

Daltonics. Extracted ion chromatograms for each well-known compound with a mass width of

0.1 m/z and time slices of 0.5 min around the expected retention time have been produced after

calibrating and baseline correction. Assignment errors have been corrected with the provided

MRSQ value.

(c) Statistical analysis of NDMMs

NDMM levels were compared simultaneously between strains and developmental stages by a

linear model in R: lm('NDMM' ~ 'developmental stage' * 'strain', data='data.frame')). *P* values

between stages and strains were adjusted for multiple testing by a false discovery rate

correction. The level of fit between linear vs. exponential growth was determined by the Akaike

information criterion (AIC). The lowest AIC for iterations of different exponents (n=1,2,3…) was

used for comparison to the simple linear model. While significant in both cases, for consistency

we present the original *p* values from the original linear model in Supplementary Figure 8.

(d) Supernatant collection and assays

Strains RS2333, RSC017, and RS2333-*daf-22.1;22.2* were raised in 10 mL liquid culture as in

the time-resolved NDMM collections (b1). For each time point, 9 mL of the supernatant was

lyophilized overnight, extracted again overnight with 90% ethanol (diluted in Millipore water)

while being stirred, and centrifuged (4000g, 10 min, 4°C). The solvent was evaporated and the

solid re-dissolved with 1 mL Millipore water. This clear extract was then directly used for the

assays. One mL of the supernatant was cleaned for HPLC-MS analysis (refer to pheromone

profiling: HPLC-MS sample preparation) for quality control. For the assays, RSC017 was

synchronized by bleaching and added to plates seeded with 300 µl OP50. The supernatants

were added to the RSCO17 J2s in two 500 µl increments (for a total of 1ml supernatant) and

dried for 30 minutes in a sterile hood after each addition. Plates were kept at 20°C and adult

mouth forms were screened three days later.

(e) Dye Staining

A stock solution of Neutral Red was prepared by dissolving 0.5 mg in 10 ml 5% acetic acid and

stored at -20° C. Working solutions were prepared by 100x dilution in M9, aliquoted, stored at -

20°C, and thawed directly before use. Working solutions were kept for approximately 1 month.

Stock solutions of 10 mM Green Bodipy were made in DMSO and stored -20. J2s were

prepared from 20-40 x 6 cm plates 6 days after passaging 5 worms to each plate on 300 µl

OP50. Worms were washed from plates with M9 into a conical tube, and then filtered through

2x20 µM filters (Millipore) placed between rubber gaskets. The flow-through contains mostly J2

and some J3, which were pelleted by centrifugation, 8 seconds on a table-top eppdendorf

centrifuge 5424, reaching approximately 10,000 x g. The juvenile pellet was then either re-

suspended in 1 ml Neutral Red working solution, or in 1 ml M9 and split to two tubes, then re-

centrifuged, and then re-suspended in either 1 ml working solution Neutral Red (0.005% in M9)

or 1 ml 50 µM Green BODIPY (Thermo) in M9. Tubes were then rotated for 3 hours in the dark,

then washed by centrifugation as before, and re-suspended in 1 ml M9. This was repeated 3-4x

until the dye was no longer visible in the worm pellet. Then, the concentration of worms was

determined by aliquoting 2 µl onto a glass coverslip in 5 technical replicates, and counted under

a dissecting microscope. Finally the appropriate number of animals was added to 6 cm plates

that had been previously seeded with 300 µl OP50, and incubated at 20°C. After 3 days, 100%

of worms exhibited Neutral Red staining (*n*=50, Supplementary figure 4). Dauers and J2s

recovered after Neutral Red staining developed at the same developmental speed (3-4 days)

and with the same mouth-form ratio as control worms recovered side-by-side (100% St for both,

Supplementary figure 5, *n*=30). Dauers and J2s stained with Cell tracker Green BODIPY (50

µM) (Thermo) were similar, although less efficiently stained compared to Neutral Red. On day 4,

90% retained intestinal fluorescence (Supplementary figure 4), although brightness decreased

with the number of days. Mouth-form ratios of dauers or J2s in +/- 50 µM Cell tracker Green

BODIPY also developed at equivalent rates and mouth-form ratios (Supplementary figure 5).

Lower than 25 µM did not yield strongly fluorescent worms after three hours. Cell Tracker Blue

CMAC (Thermo) was also used at 50 µM and imaged 3 days post-staining for *P. pacificus*, and

one day post-staining for *C. elegans*. However, due to the higher fluorescent background in the

blue light spectrum in both *P. pacificus* and *C. elegans*, we performed all experiments using only

Neutral Red and Cell tracker Green BODIPY.

(f) Mixed culture experiments and statistical analysis

We performed mixed culture experiment presented in figure 2 with 3 to 5 independent biological

replicates, and a minimum total number of counts *n* > 100 (median counts per replicate for

J2=29 and the median counts per replicate for dauers=27). J2 or dauers were stained with

Neutral Red as described in the 'Dye Staining' method section, then added to green-stained J2,

dauer, or adult populations on 6 cm plates with 300 μl OP50 and incubated at 20° C. To ensure

consistent bacterial food supply, we added 1 ml more overnight OP50-LB to each plate on the

following day, then air-dried under a chemical fume hood for 1 hour, then returned to 20° C. On

days 3-4, we phenotyped 'red' adults that exhibited no 'green' staining. To assess whether the

age of the 'green' surrounding population affects the mouth form of the dependent variable 'red'

J2s we performed a binomial regression on Eu counts (i.e. "successes") weighted by the

number of counts per replicate and the number added as a fixed effect, using a generalized

linear model from the standard statistical package in R:

glm(formula=cbind(Eu,total)~'stage_added' * '#_added', data='J2/Da', family="binomial"))

See Supplementary figure 6a for a table containing the resulting *p* values. The AIC for our

models (78.97 for J2s and 89.59 for dauers) was substantially lower than the null hypothesis

(220.16 for J2s and 147.29 for dauers), arguing a reasonable fit. For pair-wise comparisons of

the effect of age for a given number of added animals, we performed a post-hoc Fisher's exact

test on a contingency table containing the summed counts of Eu and St observations. For

display, we converted Eu counts into percent of total in figure 2, with the *p* values between the

same number of animals added indicated over the adult-added population (Significance codes:

0 '***' 0.001, '**' 0.01, '*' 0.05).

(g) Measuring the effect of food depletion on mouth form

To verify that starvation was not a factor in our mixed culture experiments, we added increasing

number of J2s to standard 6 cm plates with 300 μl OP50 (see Figure 4) to rapidly consume

bacterial food, and measured both the amount of animals that reached adulthood, and the

percent Eu in each population for two biological replicates. To assess the affects of added J2s

to each dependent variable we performed a binomial regression with count data weighted by the

total number of counts for each replicate:

glm(formula = cbind(reached_adult, total)~thousand_J2s, data=data_2, family="binomial"))

p values indicate a significant difference in percent reaching adult as a function of J2s added,

but not in percent Eu (Supplementary Fig. 6b).

(h) Microscopy

All images were taken on a Zeiss Axio Imager 2 with an Axiocam 506 mono, and processed

using Zen2 pro software. Image brightness and contrast were enhanced in ImageJ with a

minimum displayed value of 10 and maximum of 100 for all images in Fig 2, and Supplementary

figures 4 and 5, and a minimum of 21 and maximum of 117 for Supplementary figure 3. The

following exposure times were used for all images: Cy3 (peak emission = 561, exposure = 80

ms), FITC (peak emission = 519, exposure = 150 ms), Dapi (peak emission = 465, exposure =

80 ms), DIC (exposure = 80-140 ms).

(i) Dauer induction

To induce dauer, mixed-stage plates with little to no OP50 were washed with M9 and the

resulting worm pellets were used in a modified 'White Trap' method. Worm pellets were placed

on killed *Tenebrio molitor* grubs and dispersing dauers were collected in surrounding MilliQ

water. Age of dauers ranged from one week to one month.

3. Results

(a) A vital dye method for labeling nematode populations

To directly test if different age groups of nematodes influence plastic phenotypes, we required

two synchronized populations to co-habit the same space, yet still be able to identify worms

from different age groups. To do so, we developed a dye-staining methodology to robustly

differentiate between nematode populations. After trying several vital dyes, we identified that

Neutral Red [43] and CellTracker Green BODIPY (Thermo) stain nematode intestines brightly

and specifically to their respective channels (figure 2*a-e*, Supplementary Fig. 1, Materials and

methods). These dyes stain all nematodes tested including *C. elegans* (Supplementary figure 2)

and dauer larvae (Supplementary figure 3). They last more than three days (Supplementary

figure 4), allowing long-term tracking of mixed nematode populations. Importantly, neither

Neutral Red nor CellTracker Green staining affect viability, developmental rate, or the formation

of specific morphological structures, such as *P. pacificus* mouth form (Supplementary figure 5).

Thus, Neutral Red and CellTracker Green allow specific labeling of worm populations to study

age-dependent effects on phenotypes.

(b) Adult but not juvenile crowding induces the predatory mouth form in *P. pacificus*

To assess potential intra-, or inter-generational influence on *P. pacificus* mouth form we stained

juveniles of the highly St strain RSC017 with Neutral Red, and added an increasing number of

CellTracker Green-stained RSC017 adults or juveniles (figure 2*f,g*). Three days later we

phenotyped red animals that had developed into adults, but showed no green staining. To

ascertain potential differences between adding juveniles or adults, we performed a binomial

regression on Eu count data from multiple independent biological replicates (n>3), with age and

number of individuals added as fixed effects (Materials and methods, Supplementary figure 6).

We observed a significant increase in Eu worms in response to adults, but not juveniles (*p*=2.59

x10^{-2}; for display summed percents are shown in figure 2*h,i*). Almost half (48%) of the

population developed the Eu mouth form with just 500 adult animals, which is a greater than 50-

fold induction compared to side-by-side controls (figure 2*h,i*). We were also curious if dauers,

which have a thickened cuticle and represent a distinct stage in the boom-and-bust life cycle of

nematodes, could still respond to adults. Indeed, the same trend that was observed with

juveniles was seen with dauers (*p*=2.96x10^{-3}), albeit to a more muted extent (figure 2*j,k*).

Specifically, with a total of 200 dauers and 500 adults, 25.7% of dauers become Eu, whereas

only 1.8% of dauers become Eu on a plate containing 700 dauers (and no adults) (figure 2*j*).

Collectively, these data indicate that adult crowding specifically induces the Eu mouth form.

Even though we did not detect a mouth-form switch in large populations of J2s or

dauers, and food was still visible on plates containing the most animals (500 adults and 200

juveniles), we could not completely rule out the possible effect of food availability on mouth

form. As a proxy for starvation, we conducted assays with greatly increased numbers of

juveniles from 1,000 to 10,000 that would rapidly deplete bacterial food. We noticed a stark cliff

in the fraction of juveniles that reach adulthood at 4,000-5,000 animals, arguing that food is a

limiting resource at this population density (figure 2*l*). Importantly however, in these plates we

still did not see a shift in mouth form (p=0.99, binomial regression, Supplementary figure 6).

With an overwhelming 10,000 worms on a plate, 5.8% were Eu, compared to 48% in the

presence of only 500 adults. While longer-term starvation may yet have an impact on mouth

form, under our experimental conditions it appears to be negligible.

(c) Adult but not juvenile secretions induce the Eu mouth form

As mouth-form plasticity in *P. pacificus* is regulated by nematode-derived modular metabolites

(NDMMs)[32], we wondered if the difference between adults and juveniles resulted from

differences in secreted NDMMs. To test this hypothesis we added secretions from adult or

juvenile worms to RSC017 (highly St) juveniles. We found that adult secretions from both the

laboratory stain RS2333 (highly Eu) and RSC017 led to a significant increase in the Eu morph

relative to juvenile secretions ($p=5.27\times10^{-06}$, 1.33×10^{-3}, respectively, Fisher's exact test)(figure

3). To confirm the effect was caused by ascaroside pheromones, we exposed RSC017 juveniles

to supernatant from a *daf-22.1;22.2* double mutant, which exhibits virtually no ascaroside

production in both *C. elegans* and *P. pacificus* [26,44]. Again, juvenile secretion had no impact

on Eu frequency, but in contrast to wild-type supernatants, we also observed no increase in Eu

frequency with adult secretions ($p=0.8324$, Fisher's exact test, figure 3). Thus, adult-specific

NDMMs induce development of the Eu mouth form.

(d) Developmental-staged NDMM profiles reveal adult-specific synthesis of dasc#1

Next, we investigated whether the difference between adult and juvenile pheromones is one of

dosage, or of identity. To answer this question and verify age-specific differences in

pheromones, we profiled *P. pacificus* NDMM levels in two strains and at three time points

through development. We used RS2333 and RSC017 and measured the exo-metabolomes of

juvenile stage 2 (J2s, 24 hrs), J3s (48 hrs) and J4/adults (72 hrs) from a constant culture with

excess OP50 bacterial food (figure 4*a,b*, Supplementary figure 7, Materials and methods). To

assess potential differences in pheromone levels we performed a linear regression with the area

under the curve for each NDMM (aoc) (Supplementary Figure 1a) as the response variable.

Stage and strain were modeled as fixed effects, and because we performed separate

regression analyses for each pheromone, we adjusted the resulting *p* values for multiple testing

using false discovery rate (FDR)(see Supplementary figure 8 for *p* and *FDR* values between

stage and strain). We observed that there was a significant affect of developmental stage on the

levels of ascr#9, ascr#12, npar#1, and dasc#1, and that ubas#1 and #2 are strain and stage

specific (FDR<0.05). Interestingly, dasc#1 is the most potent known Eu-inducing compound

when tested as a single synthesized compound, while npar#1 is both Eu- and dauer-inducing

(figure 4*c,d,f-i*)[32]. Closer inspection revealed dasc#1, npar#1, and ascr#9 increase throughout

development in both strains, and dasc#1 peaks in adults in RS2333 (p<0.05, student's two-

tailed *t*-test between 72 hrs and 24 hrs for each NDMM in both strains, and also 72 hrs and 48

hrs for dasc#1 in RS233). Intriguingly, the trajectory of dasc#1 appeared binary in both strains

(figure 4*f,g*). In fact our statistical model for dasc#1 fits better if we assume cubic rather than

linear growth (*ΔAIC*=3.958). In contrast, ascr#9, which was also statistically up-regulated but

does not affect known plastic phenotypes [32], displays a more gradual increase in both strains

(figure 4*e,j,k*), and the model fits better with linear growth ($AIC_{linear} - AIC_{cubic} = -1.208$).

Meanwhile, the trajectory of npar#1 appears strain specific (figure 4*h,i*). Hence the mode of

induction is NDMM-specific, and the kinetics of production may be related to their roles in

phenotypic plasticity.

In principle, the increase in abundance of certain pheromones could be a result of a

concomitant increase in body mass, however several observations indicate more targeted

regulation. First, no other compounds were significantly different in our linear model. Second, an

analysis of previously published RNA-seq data [45] reveals the increase in NDMM abundance

corresponds to an increase in transcription of the thiolase *Ppa-daf-22.1* (Supplementary figure

9) the most downstream enzyme in the β-oxidation pathway of ascaroside synthesis. Third,

pasc#9 and pasc#12 actually exhibit a peak in abundance at the 48 hr/J3 time point, rather than

in 72 hrs/adults. Finally, we profiled the endo-metabolome of eggs, and found appreciable

amounts of ascr#1, #9, #12, and pasc#9, but little to no traces of other ascaroside derivatives

(Supplementary figure 7), suggesting age-specific synthesis rather than release. Together,

these results suggest that the observed increase in ubas#1 and #2, ascr#9, npar#1, and dasc#1

over time corresponds to age-specific production. The observation that dasc#1 is produced

specifically during the juvenile-to-adult transition is especially intriguing because adults are no

longer able to switch mouth forms, hinting at cross-generational signaling.

4. Discussion

Here, we introduce a novel dye-based method that allowed us to determine cross-generational

influence on mouth form, and demonstrate this effect is a result of age-specific production of

plasticity affecting pheromones. In doing so, we provide the first multi-stage time series of

pheromone production in *P. pacificus*, which shows that dasc#1 exhibits a surprising 'off-on'

switch-like induction pattern. Collectively, our results argue that adults represent a critical age

group [46] in nematode populations.

Our developmental profiling revealed an increase in two NDMMs that affect plastic

phenotypes. The observation that this trend mirrors the transcriptional regulation of enzymes

involved in NDMM synthesis argues that the stage-dependent increase is not simply a result of

an increase in body mass, but rather that these molecules are programmed for stage-specific

induction. The binary 'off-on' kinetics might reflect a population level feed-back loop, such that

the production of density-sensing pheromones is based on a threshold level of previously

produced pheromones. It is also worth noting that while npar#1 is the major dauer-inducing

pheromone in *P. pacificus* [32], we did not observe dauers in any experimental setup described

herein. Thus, it seems that mouth-form phenotype is the first-level plastic response to

population density. Presumably higher concentrations are required for dauer induction,

reflecting a calculated response strategy depending on the level of crowding or duration of

starvation. Interestingly, the effect of adult supernatants was noticeably less (23%-26% Eu) than

with adult worms (up to 48% with only 500 adults). It is difficult to compare the amount of

pheromone concentrations between experiments, but presumably worms in the vital-dye assay

experienced a greater local concentration as they were in direct physical contact with each

other, compared to worms in the supernatant assay.

Among the many environmental influences on mouth form [42], population density and

starvation are perhaps the most ecologically relevant. However, teasing apart these two factors

has proven difficult [47]. Here, we demonstrate that while a strong shift is observed with adult-

specific pheromones, no such effect was seen under limited resource conditions. Thus, age-

specific crowding is sufficient to induce the Eu mouth form. Nevertheless, this does not preclude

that long term starvation could also have an effect. Determining the relative contributions of

these factors to mouth form will be important to better understand the sophisticated ecological

response strategies of *P. pacificus*, nematodes, and phenotypic plasticity in general.

Why do adults and not juveniles affect mouth form? Given that St animals can develop

faster [48], there may be a 'race' to sexual maturation in emergent populations at low densities.

But as the nematode population increases, there will likely be a commensurate decrease in

bacterial populations. When faced with competition from other nematodes, *P. pacificus* has a

particular advantage in developing the Eu morph; their expanded dietary range includes their

competition. Indeed, when nematode prey is the only available food source, the Eu morph

provides longer life spans and more progeny than the St morph [49]. When resources become

depleted as population size increases, *C. elegans* and other monomorphic nematodes may

enter dauer and disperse [50]. But in St-biased dimorphic strains of *P. pacificus*, juveniles may

switch to the Eu morph in response to adults as a first level indication of rapidly increasing

population size (Supplementary figure 10). Then, after prolonged starvation and crowding,

worms will presumably enter dauer. By analogy to economic models of population growth

[51,52] we view mouth-form plasticity and dauer formation as alternative 'technological

innovations' to escape a Malthusian resource trap. To what extent this occurs in nature, or with

P. pacificus strains that are highly Eu, remains to be determined.

The evolution of dimorphic mouth forms is one among myriad nematode ecological

strategies. For example, entomopathogenic nematodes release their symbiont bacteria in insect

hosts to establish their preferred food source, and some release antibiotics to kill off competing

bacteria and fungi from other entomopathogenic species [53]. Some free-living species, like

those of the genus *Oscheius*, refrain from combat and stealthily feed and reproduce amidst

warring entomopathogenic species. Interspecific killing also occurs in gonorchoristic species, in

which both mated and virgin males are killed, implying fighting not just for mates but for

resources as well [54,55]. Reproductive strategies also exist, and hermaphroditic species have

an advantage over gonachristic species when colonizing a new niche, such as an insect

carcass [56]. Meanwhile insect hosts and colonizing nematodes have their own distinct

pheromone-based attraction and toxicity [57,58]. Finally, the renaissance of *C. elegans*

sampling from around the world [59-63] is rapidly building a resource of wild isolates that will

almost certainly have different and fascinating ecologies. We hope our method for labeling and

then combining different nematode populations on the same plate will aid in studies to identify

these strategies. Perhaps the time is also ripe to complement these studies with more sophisted

ecological modelling that can lead to more testable hypotheses.

Although beyond the scope of this manuscript, the cross-generational communication we

observed could in principle reflect an intended signal from adults to juveniles, i.e. kin selection

[64]. However, we favor a more simplistic view that juveniles have evolved to recognize adult-

produced metabolites. Regardless of these interpretations, our results argue that age groups

are a critical factor in density-dependent plasticity, as has been theorized in density-dependent

selection [65].

Author Contributions

402   MSW and RJS conceived of the project. MC conducted pheromone profiling. MSW and TR

403   designed and conducted dye-labeling experiments. TR and MC performed supernatant

404   experiments. MD and MSW considered ecological implications. MSW and TR wrote the

405   manuscript with input and edits from all authors.

Acknowledgements

408   We would like to thank all members of the Sommer lab, Dr. Talia Karasov, Dr. Hernan Burbano

409   and Moises Exposito-Alonso for guidance with statistical analysis, and Dr. Adrian Striet (Max

410   Planck Institute), and Dr. Cameron Weadick (University of Sussex) for thoughtful critique and

411   discussion.

References

1.   Hastings, A. 2013 *Population biology: concepts and models*.

2.   MacArthur, R. H. 1962 Some generalized theorems of natural selection. *Proc Natl*
Acad Sci USA **48**, 1893–1897. (doi:10.1073/pnas.48.11.1893)

3.   Travis, J., Leips, J. & Rodd, F. H. 2013 Evolution in Population Parameters: Density-
Dependent Selection or Density-Dependent Fitness? *The American Naturalist* **181**,
S9–S20. (doi:10.1086/669970)

4.   Dudley, S. A. & Schmitt, J. 2015 Testing the Adaptive Plasticity Hypothesis: Density-
Dependent Selection on Manipulated Stem Length in Impatiens capensis. *The*
American Naturalist **147**, 445–465. (doi:10.1086/285860)

5.   Pener, M. P. & Simpson, S. J. 2009 Locust Phase Polyphenism: An Update.
Advances in insect physiology. (doi:10.1016/S0065-2806(08)36001-9)

6.   Simpson, S. J., Despland, E., Hägele, B. F. & Dodgson, T. 2001 Gregarious behavior
in desert locusts is evoked by touching their back legs. *Proc Natl Acad Sci USA* **98**,
3895–3897. (doi:10.1073/pnas.071527998)

7.   Slogget, J. J. & Weisser, W. W. 2004 *Aphids in a New Millennium. INRA*

.

8.   Simpson, S. J. & Miller, G. A. 2007 Maternal effects on phase characteristics in the
desert locust, Schistocerca gregaria: A review of current understanding. *Journal of*
Insect Physiology **53**, 869–876. (doi:10.1016/j.jinsphys.2007.05.011)

9.   Maeno, K. & Tanaka, S. 2008 Maternal effects on progeny size, number and body
color in the desert locust, Schistocerca gregaria: Density- and reproductive cycle-
dependent variation. *Journal of Insect Physiology* **54**, 1072–1080.
(doi:10.1016/j.jinsphys.2008.04.010)

10.   Chen, B., Li, S., Ren, Q., Tong, X., Zhang, X. & Le Kang 2015 Paternal epigenetic
effects of population density on locust phase-related characteristics associated with
heat-shock protein expression. *Molecular Ecology* **24**, 851–862.
(doi:10.1111/mec.13072)

11.   Hamouda, A. B., and, S. T. O. E. 2011. Density-dependent phenotypic plasticity in
body coloration and morphometry and its transgenerational changes in the migratory
locust, Locusta migratoria. *Journal of Entomology and Nematology.*

12.   Ben Dantzer, Newman, A. E. M., Boonstra, R., Palme, R., Boutin, S., Humphries, M.
M. & McAdam, A. G. 2013 Density Triggers Maternal Hormones That Increase
Adaptive Offspring Growth in a Wild Mammal. *Science* **340**, 1235765–1217.
(doi:10.1126/science.1235765)

13.   Butcher, R. A. 2017 Small-molecule pheromones and hormones controlling nematode
development. *Nature Chemical Biology* **13**, 577–586. (doi:10.1038/nchembio.2356)

14.   Butcher, R. A., Fujita, M., Schroeder, F. C. & Clardy, J. 2007 Small-molecule
pheromones that control dauer development in Caenorhabditis elegans. *Nature*
Chemical Biology **3**, 420–422.

15.   Reuss, von, S. H., Bose, N., Srinivasan, J., Yim, J. J., Judkins, J. C., Sternberg, P. W.
& Schroeder, F. C. 2012 Comparative metabolomics reveals biogenesis of
ascarosides, a modular library of small-molecule signals in C. elegans. *J. Am. Chem.*
Soc. **134**, 1817–1824. (doi:10.1021/ja210202y)

16.   Fielenbach, N. & Antebi, A. 2008 C. elegans dauer formation and the molecular basis
of plasticity. *Genes & Development* **22**, 2149–2165. (doi:10.1101/gad.1701508)

17.   Butcher, R. A., Ragains, J. R., Kim, E. & Clardy, J. 2008 A potent dauer pheromone

component in Caenorhabditis elegans that acts synergistically with other components.
Proc Natl Acad Sci USA **105**, 14288–14292. (doi:10.1073/pnas.0806676105)

18.   Jeong, P.-Y. et al. 2005 Chemical structure and biological activity of the
465   Caenorhabditis elegans dauer-inducing pheromone. *Nature* **433**, 541–545.
466   (doi:10.1038/nature03201)

467   19.   Chasnov, J. R., So, W. K., Chan, C. M. & Chow, K. L. 2007 The species, sex, and
468   stage specificity of a Caenorhabditis sex pheromone. *Proc Natl Acad Sci USA* **104**,
469   6730–6735. (doi:10.1073/pnas.0608050104)

470   20.   Izrayelit, Y., Srinivasan, J., Campbell, S. L., Jo, Y., Reuss, von, S. H., Genoff, M. C.,
471   Sternberg, P. W. & Schroeder, F. C. 2012 Targeted metabolomics reveals a male
472   pheromone and sex-specific ascaroside biosynthesis in Caenorhabditis elegans. *ACS*
473   *Chem. Biol.* **7**, 1321–1325. (doi:10.1021/cb300169c)

474   21.   Kaplan, F. et al. 2011 Ascaroside Expression in Caenorhabditis elegans Is Strongly
475   Dependent on Diet and Developmental Stage. *PLoS ONE* **6**, e17804.
476   (doi:10.1371/journal.pone.0017804)

477   22.   Golden, J. W. & Riddle, D. L. 1982 A pheromone influences larval development in the
478   nematode Caenorhabditis elegans. *Science* **218**, 578–580.
479   (doi:10.1126/science.6896933)

480   23.   Srinivasan, J. et al. 2012 A Modular Library of Small Molecule Signals Regulates
481   Social Behaviors in Caenorhabditis elegans. *PLOS Biol* **10**, e1001237.
482   (doi:10.1371/journal.pbio.1001237)

483   24.   Ludewig, A. H. et al. 2017 Larval crowding accelerates C. elegans development and
484   reduces lifespan. *PLoS Genet* **13**, e1006717. (doi:10.1371/journal.pgen.1006717)

485   25.   Choe, A., Reuss, von, S. H., Kogan, D., Gasser, R. B., Platzer, E. G., Schroeder, F. C.
486   & Sternberg, P. W. 2012 Ascaroside Signaling Is Widely Conserved among
487   Nematodes. *Current Biology* **22**, 772–780. (doi:10.1016/j.cub.2012.03.024)

488   26.   Markov, G. V., Meyer, J. M., Panda, O., Artyukhin, A. B., Claaßen, M., Witte, H.,
489   Schroeder, F. C. & Sommer, R. J. 2016 Functional Conservation and Divergence of
490   daf-22 Paralogs in Pristionchus pacificus Dauer Development. *Mol Biol Evol* **33**,
491   2506–2514. (doi:10.1093/molbev/msw090)

492   27.   Dong, C., Reilly, D. K., Bergame, C., Dolke, F., Srinivasan, J. & Reuss, von, S. H.
493   2018 Comparative Ascaroside Profiling of CaenorhabditisExometabolomes Reveals
494   Species-Specific (ω) and (ω – 2)-Hydroxylation Downstream of Peroxisomal β-
495   Oxidation. *J. Org. Chem.* **83**, 7109–7120. (doi:10.1021/acs.joc.8b00094)

496   28.   Diaz, S. A., Brunet, V., Lloyd-Jones, G. C., Spinner, W., Wharam, B. & Viney, M. 2014
497   Diverse and potentially manipulative signalling with ascarosides in the model
498   nematode C. elegans. *BMC Evol. Biol.* **14**, 46. (doi:10.1186/1471-2148-14-46)

29.   Bose, N., Meyer, J. M., Yim, J. J., Mayer, M. G., Markov, G. V., Ogawa, A.,

Schroeder, F. C. & Sommer, R. J. 2014 Natural Variation in Dauer Pheromone
Production and Sensing Supports Intraspecific Competition in Nematodes. *Current*
Biology **24**, 1536–1541. (doi:10.1016/j.cub.2014.05.045)

30. Greene, J. S. et al. 2016 Balancing selection shapes density-dependent foraging
behaviour. *Nature* **539**, 254–258. (doi:10.1038/nature19848)

31. Falcke, J. M. et al. 2018 Linking Genomic and Metabolomic Natural Variation
Uncovers Nematode Pheromone Biosynthesis. *Cell Chem Biol* **25**, 787–796.e12.
(doi:10.1016/j.chembiol.2018.04.004)

32. Bose, N., Ogawa, A., Reuss, von, S. H., Yim, J. J., Ragsdale, E. J., Sommer, R. J. &
Schroeder, F. C. 2012 Complex Small-Molecule Architectures Regulate Phenotypic
Plasticity in a Nematode. *Angewandte Chemie International Edition* **51**, 12438–12443.
(doi:10.1002/anie.201206797)

33. Sommer, R. J. & McGaughran, A. 2013 The nematode Pristionchus pacificus as a
model system for integrative studies in evolutionary biology. *Molecular Ecology* **22**,
2380–2393. (doi:10.1111/mec.12286)

34. Herrmann, M., Mayer, W. E. & Sommer, R. J. 2006 Nematodes of the genus
Pristionchus are closely associated with scarab beetles and the Colorado potato
beetle in Western Europe. *Zoology* **109**, 96–108. (doi:10.1016/j.zool.2006.03.001)

35. Herrmann, M., Mayer, W. E., Hong, R. L., Kienle, S., Minasaki, R. & Sommer, R. J.
2007 The Nematode Pristionchus pacificus (Nematoda: Diplogastridae) Is Associated
with the Oriental Beetle Exomala orientalis (Coleoptera: Scarabaeidae) in Japan.
Zoological Society of Japan. (doi:10.2108/zsj.24.883)

36. Bento, G., Ogawa, A. & Sommer, R. J. 2010 Co-option of the hormone-signalling
module dafachronic acid–DAF-12 in nematode evolution. *Nature* **466**, 494–497.
(doi:10.1038/nature09164)

37. Sommer, R. J., Dardiry, M., Lenuzzi, M., Namdeo, S., Renahan, T., Sieriebriennikov,
B. & Werner, M. S. 2017 The genetics of phenotypic plasticity in nematode feeding
structures. *Open Biology* **7**, 160332–118. (doi:10.1098/rsob.160332)

38. Sanghvi, G. V., Baskaran, P., Röseler, W., Sieriebriennikov, B., Rödelsperger, C. &
Sommer, R. J. 2016 Life History Responses and Gene Expression Profiles of the
Nematode Pristionchus pacificus Cultured on Cryptococcus Yeasts. *PLoS ONE* **11**,
e0164881. (doi:10.1371/journal.pone.0164881)

39. Wilecki, M., Lightfoot, J. W., Susoy, V. & Sommer, R. J. 2015 Predatory feeding
behaviour in Pristionchus nematodes is dependent on phenotypic plasticity and
induced by serotonin. *J. Exp. Biol.* **218**, 1306–1313. (doi:10.1242/jeb.118620)

40. Sommer, R. J. & Mayer, M. G. 2015 Toward a synthesis of developmental biology
with evolutionary theory and ecology. *Annu. Rev. Cell Dev. Biol.* **31**, 453–471.
(doi:10.1146/annurev-cellbio-102314-112451)

41. Meyer, J. M., Baskaran, P., Quast, C., Susoy, V., Rödelsperger, C., Glöckner, F. O. &
Sommer, R. J. 2017 Succession and dynamics of Pristionchus nematodes and their
microbiome during decomposition of Oryctes borbonicus on La Réunion Island.
Environmental Microbiology **19**, 1476–1489. (doi:10.1111/1462-2920.13697)

42. Werner, M. S., Sieriebriennikov, B., Loschko, T., Namdeo, S., Lenuzzi, M., Dardiry,
M., Renahan, T., Sharma, D. R. & Sommer, R. J. 2017 Environmental influence on
Pristionchus pacificus mouth form through different culture methods. *Sci Rep* **7**, 7207.
(doi:10.1038/s41598-017-07455-7)

43. Thomas, M. C. & Lana, P. D. C. 2008 Evaluation of vital stains for free-living marine
nematodes. *Brazilian Journal of Oceanography* **56**, 249–251. (doi:10.1590/S1679-
87592008000300009)

44. Golden, J. W. & Riddle, D. L. 1985 A gene affecting production of the Caenorhabditis
elegans dauer-inducing pheromone. *Mol. Gen. Genet.* **198**, 534–536.

45. Baskaran, P., Rödelsperger, C., Prabh, N., Serobyan, V., Markov, G. V., Hirsekorn, A.
& Dieterich, C. 2015 Ancient gene duplications have shaped developmental stage-
specific expression in Pristionchus pacificus. *BMC Evol. Biol.* **15**, 185.
(doi:10.1186/s12862-015-0466-2)

46. Charlesworth, B. 1972 Selection in populations with overlapping generations. III.
Conditions for genetic equilibrium. *Theoretical Population Biology*

47. Bento, G., Ogawa, A. & Sommer, R. J. 2010 Co-option of the hormone-signalling
module dafachronic acid–DAF-12 in nematode evolution. *Nature* **466**, 494–497.
(doi:10.1038/nature09164)

48. Serobyan, V., Ragsdale, E. J., Müller, M. R. & Sommer, R. J. 2013 Feeding plasticity
in the nematode Pristionchus pacificusis influenced by sex and social context and is
linked to developmental speed. *Evolution & Development* **15**, 161–170.
(doi:10.1111/ede.12030)

49. Serobyan, V., Ragsdale, E. J. & Sommer, R. J. 2014 Adaptive value of a predatory
mouth-form in a dimorphic nematode. *Proceedings of the Royal Society of London B:
Biological Sciences* **281**, 20141334–989. (doi:10.1098/rspb.2014.1334)

50. Frézal, L. & Félix, M.-A. 2015 The Natural History of Model Organisms: C. elegans
outside the Petri dish. *Elife* **4**, e05849. (doi:10.7554/eLife.05849)

51. Trewavas, A. 2002 Malthus foiled again and again. *Nature* **418**, 668–670.
(doi:10.1038/nature01013)

52. Malthus, T. R., Winch, D. & James, P. 1992 *Malthus: 'An Essay on the Principle of
Population'*. Cambridge University Press.

53. Griffin, C. T. 2012 Perspectives on the behavior of entomopathogenic nematodes
from dispersal to reproduction: traits contributing to nematode fitness and biocontrol
efficacy. *J. Nematol.* **44**, 177–184.

54. Zenner, A. N. R. L., O'Callaghan, K. M. & Griffin, C. T. 2014 Lethal Fighting in
Nematodes Is Dependent on Developmental Pathway: Male-Male Fighting in the
Entomopathogenic Nematode Steinernema longicaudum. *PLoS ONE* **9**, e89385.
(doi:10.1371/journal.pone.0089385)

55. O'Callaghan, K. M., Zenner, A. N. R. L., Hartley, C. J. & Griffin, C. T. 2014
Interference competition in entomopathogenic nematodes: male Steinernema kill
members of their own and other species. *International Journal for Parasitology* **44**,
1009–1017. (doi:10.1016/j.ijpara.2014.07.004)

56. Campos-Herrera, R. 2015 *Nematode Pathogenesis of Insects and Other Pests*.
Cham: Springer.

57. Renahan, T. & Hong, R. L. 2017 A species-specific nematocide that results in terminal
embryogenesis. *J. Exp. Biol.* **220**, jeb.159665–3247. (doi:10.1242/jeb.159665)

58. Cinkornpumin, J. K., Wisidagama, D. R., Rapoport, V., elife, J. G. 2014 A host beetle
pheromone regulates development and behavior in the nematode Pristionchus
pacificus. *Elife*.

59. Evans, K. S., Zhao, Y., Brady, S. C., Long, L., McGrath, P. T. & Andersen, E. C. 2016
Correlations of Genotype with Climate Parameters Suggest Caenorhabditis elegans
Niche Adaptations. *G3: Genes, Genomes, Genetics* **7**, g3.116.035162–298.
(doi:10.1534/g3.116.035162)

60. Félix, M.-A., Jovelin, R., Ferrari, C., Han, S., Cho, Y. R., Andersen, E. C., Cutter, A. D.
& Braendle, C. 2013 Species richness, distribution and genetic diversity of
Caenorhabditis nematodes in a remote tropical rainforest. *BMC Evol. Biol.* **13**, 10.
(doi:10.1186/1471-2148-13-10)

61. Cook, D. E., Zdraljevic, S., Roberts, J. P. & Andersen, E. C. 2017 CeNDR, the
Caenorhabditis elegans natural diversity resource. *Nucleic Acids Res.* **45**, D650–
D657. (doi:10.1093/nar/gkw893)

62. Petersen, C., Dirksen, P., Prahl, S., Strathmann, E. A. & Schulenburg, H. 2014 The
prevalence of Caenorhabditis elegans across 1.5 years in selected North German
locations: the importance of substrate type, abiotic parameters, and Caenorhabditis
competitors. *BMC Ecol.* **14**, 4. (doi:10.1186/1472-6785-14-4)

63. Poullet, N. & Braendle, C. 2015 Sampling and Isolation of C. elegans from the Natural
Habitat. *Methods Mol. Biol.* **1327**, 221–229. (doi:10.1007/978-1-4939-2842-2_16)

64. Bourke, A. F. G. 2014 Hamilton's rule and the causes of social evolution.
Philosophical Transactions of the Royal Society of London B: Biological Sciences **369**,
20130362–20130362. (doi:10.1098/rstb.2013.0362)

65. Charlesworth, B. 1994 *Evolution in age-structured populations*.

Figure Legends

Figure 1. Life cycle and developmental plasticity of the model nematode *Pristionchus*

***pacificus*.**

(a) The life cycle of *P. pacificus* consists of four juvenile stages (J1-4) until sexual maturation

(adults). Like many nematodes *P. pacificus* can enter a long-living 'dormant' dauer state that is

resistant to harsh environmental conditions. The decision to continue through the direct life

cycle or enter dauer is regulated by small molecule excreted ascarosides (chemical structure

adapted from [13]). (b) *P. pacificus* can also adopt one of two possible feeding structures; either

a microbivorous narrow-mouth (stenostomatous, St), or (c) an omnivorous wide-mouth

(eurystomatous, Eu) with an extra tooth that can be utilized to kill and eat other nematodes or

fungi. White lines indicate the presence of an extra tooth (right side) in the Eu morph or its

absence in the St morph, and the dorsal tooth (left side), which is flint-like in St and hook-like in

Eu. White scale bar indicates 5 µM. (d), *P. pacificus* is often found in a necromenic association

with beetles (ex. shown here *Oryctes borbonicus*) in the dauer state, and resumes the free living

life cycle upon beetle death to feed on the ensuing microbial bloom.

Figure 2. Vital-dye method confirms adult-specific density effect on mouth form.

(a-e) *P. pacificus* were stained with either 0.005% Neutral Red or 50 µM CellTracker Green

Bodipy (Thermo) and viewed using Cy3 and FITC filters. Images were merged with Differential

interference contrast (DIC), scale bar = 100 µM. An example of relative intensities of each

fluorescence channel are displayed in the histograms (right) with the chemical structure of

Neutral Red or CellTracker Green Bodipy. (f) Age-dependent functional pheromone assay:

Experimental juveniles were stained with Neutral Red, and challenged with CellTracker Green

Bodipy-stained juveniles or adults on standard condition NGM agar plates seeded with 300 μl

OP50 *E. coli*. Three days later, only red-positive and green-negative adults were phenotyped.

(g) The wild isolate RSC017 grown in standard conditions (5 young adults passed to fresh

plates, progeny phenotyped 4 days later) are highly stenostomatous (<10%, *n*=102). (h,i) Mouth

form ratios of Neutral Red-stained J2s, and (j,k) dauers, with increasing number of CellTracker

Green-stained competitors, as described in (f) (n=3-5 independent biological replicates per

experiment, with total n=104, 126, 134, and 109 for 200 NR-stained J2 with 0, 50, 200, and 500

CellTracker Green-stained J2s, and 115, 113, and 112 for 50, 200, and 500 CellTracker Green-

stained adults; n=104, 105, 105, and 102 for NR-stained dauers with 0, 50, 200, and 500

CellTracker Green-stained dauers, and 100, 103, and 105 for 50, 200, and 500 CellTracker

Green-stained adults. Overall significance between strain and age was determined by a

binomial linear regression (see Materials and methods), and pair-wise comparisons were

assessed by Fisher's exact test on summed Eu counts (Significance codes: '***' < 0.001, '**'

<0.01, '*' <0.05). Mouth forms were phenotyped at 40-100x on a Zeiss Axio Imager 2 light

microscope. (i) Percent reaching adulthood (top) and percent Eu of those that reached

adulthood (bottom) after increasing numbers of J2s are added to standard 6 cm NGM agar

plates with 300 μl OP50 *E.coli* bacteria (n=2 biological replicates, with total n for percent

reaching adulthood=216, 202, 206, 207, and 202 for 1, 3, 4, 5, and 10 thousand J2s, and total n

for mouth form=106, 106, 103, 110, and 104). Significance was determined by a binomial

regression (see Materials and methods).

Figure 3. Adult-specific secretions induce predatory morph in juveniles.

Highly St strain RSC017 juveniles were exposed to J2 and adult supernatants of its own strain,

and to the J2 and adult supernatants of highly Eu strain RS2333. Mouth form was phenotyped

three days later. Worms exposed to J2 secretions remained highly St, while worms exposed to

adult secretions had a small but significant increase in Eu morphs (p<0.05, Fisher's exact test).

Supernatants from the double mutant *daf-22.1/2*, which has deficient ascaroside pheromone

production, did not elicit juvenile or adult increase in Eu. Worms exposed to the S-media control

also remained highly St. n=4 independent biological replicates for RS2333 and *daf-22.1/2*

secretions, and n=2 independent biological replicates for RSC017 adult and juvenile secretions,

with an average count of 55 animals per replicate. For display, total Eu and St counts are

presented as percentages (Significance codes: '***' < 0.001, '**' <0.01, '*' <0.05).

Figure 4. Time-resolved Nematode-Derived Modular Metabolites (NDMMs) in *Pristionchus*

pacificus.

(a) Time resolved secretion profile of nematode derived modular metabolites from the wild-type

laboratory strain RS2333 and (b) wild isolate RSC017. Data is presented as the mean of 8

(RS2333) and 9 (RSC017) biological replicates, and error bars represent standard error of the

mean (SEM). (c-e) Chemical structures of adult specific NDMMs dasc#1, npar#1, and ascr#9,

as described in the Small Molecule Identifier Database (http://www.smid-db.org/), produced in

ChemDraw. (f-k) Time-resolved abundance of dasc#1, npar#1, and ascr#9 NDMMs in RS2333

and RSC017. Each data point represents a biological replicate, and for comparison to (a-b) lines

represent mean abundance. *P* values calculated by a 2-tailed students *t*-test (Significance

codes: '***' < 0.001, '**' <0.01, '*' <0.05).

A

t = 0

adult

embryo

72 hrs

fatty acid side chain

ascarylose sugar

terminus groups

head groups

J1

24 hrs

J2

J4

J3

'direct'

48 hrs

dauer

B St

C Eu

D

Figure 1. Life cycle and developmental plasticity of the model nematode *Pristionchus pacificus*.

Figure 2. Vital-dye method confirms adult-specific density effect on mouth form.

Figure 3. Adult-specific secretions induce predatory morph in juveniles.

Figure 4. Time-resolved Nematode-Derived Modular Metabolites (NDMMs) in *Pristionchus pacificus*.