Geometry-dependent instabilities in electrically excitable tissues

Harold M. McNamara[1,2], Stephanie Dodson[3], Yi-Lin Huang[4], Evan W. Miller[4], Björn Sandstede[3], Adam E. Cohen[1,5,6*]

1 Department of Physics, Harvard University
2 Harvard-MIT Division of Health Sciences and Technology
3 Division of Applied Mathematics, Brown University
4 Departments of Chemistry, Molecular and Cell Biology, and Hellen Wills
Neuroscience Institute, University of California, Berkeley
5 Department of Chemistry and Chemical Biology, Harvard University
6 Howard Hughes Medical Institute

* cohen@chemistry.harvard.edu

Abstract

Little is known about how individual cells sense the macroscopic geometry of their tissue environment. Here we explore whether long-range electrical signaling can convey information on tissue geometry to influence electrical dynamics of individual cells. First, we studied an engineered electrically excitable cell line where all voltage-gated channels were known. Cells grown in patterned islands of different shapes showed remarkably diverse firing patterns, including regular spiking, period-doubling alternans, and arrhythmic firing. A spatially resolved Hodgkin-Huxley numerical model quantitatively reproduced these effects, showing how the macroscopic geometry affected the single-cell electrophysiology via gap junction-mediated electrical coupling. Qualitatively similar geometry dependent dynamics were observed in human induced pluripotent stem cell (iPSC)-derived cardiomyocytes. The cardiac results urge caution in translating observations of arrhythmia in vitro to predictions in vivo where the tissue geometry is very different. These findings establish that tissue geometry is a fundamental determinant of dynamical stability in excitable cells.

Introduction

Cells in multicellular organisms sense their location within the tissue via diffusible, contact, and mechanical signals. Gap junction-mediated electrical signals can also, in principle, provide long-range positional cues (1), though mechanistic details have been difficult to determine. Electrical sensitivity to distant boundaries is best established in the heart, where, for instance, structural defects can act as nuclei for arrhythmias (2). However, interpreting such effects in the heart is difficult due to the multiple interacting factors that govern dynamics, including electrical coupling between myocytes (3-5), mechano-electrical feedbacks (6, 7), and the properties of the individual myocytes (8-10). Subtle shifts in any of these parameters can cause discontinuous changes in dynamics, e.g. from a stable beat to a possibly fatal arrhythmia. It is important to understand how long-range electrical signaling can convey positional cues generally, and how these signals govern cardiac stability specifically.

Early models of cardiac dynamics established a stability criterion based solely on the cell-autonomous relation of spike width to beat rate.(11) This thinking motivated studies of arrhythmias in isolated cells as a means to predict dynamics *in vivo* (12, 13). For instance, it has been widely claimed that if human induced pluripotent stem cell (iPSC)-derived myocytes

(hiPSC-CM) can be made to show mature patterns of ion channel expression (14-17), then *in vitro* cultures will be a useful substrate for studying arrhythmias (18-21). Cardiotoxicity screening in hiPSC-CM is one of the pillars of the ongoing FDA-sponsored Comprehensive *in vitro* Proarrhythmia Assay (CiPA) project (19). However, it is not known to what extent the difference in geometry between cultured cells and intact tissues affects the dynamics or stability. Gap junctions *in vivo* introduce electrical coupling through which each myocyte is influenced by its neighbors, and ultimately by the large-scale geometry of the tissue. One can then ask whether this coupling is a minor perturbation on the individual myocytes; or whether it fundamentally changes the dynamics. In condensed matter physics, the properties of a bulk solid can differ dramatically from the constituent atoms. In the presence of intercellular coupling, the stability of the bulk tissue might differ dramatically from that of individual cells.

Recent theoretical work showed that conduction could dramatically alter the stability conditions (22, 23). However, the wide diversity of cardiac models, combined with uncertainty in model parameters, presents a challenge for comparison to experiments (23, 24). Only a few experiments have explicitly probed the roles of intercellular coupling in cardiac dynamics (25, 26). In complex cells such as cardiomyocytes, one typically cannot vary one parameter without affecting many others. For instance, growing hiPSC-CM on different size islands affects their patterns of gene expression (27). Thus it can be difficult to determine the underlying cause of discrepancies between *in vitro* and *in vivo* results.

An alternative approach is to engineer synthetic excitable tissues where all elements are well understood, and then to test the necessity and sufficiency of specific components for different aspects of the dynamics. This synthetic approach has the further merit of being amenable to rigorous quantitative modeling. Optopatch Spiking Human Embryonic Kidney (OS-HEK) cells(28) are an engineered excitable cell type with just two voltage-dependent channels, the voltage-gated cardiac sodium channel, $Na_V1.5$, and the inward rectifier potassium channel, $K_{ir}2.1$. When electrically stimulated, these cells produced single action potentials. When grown into a confluent syncytium, endogenous gap junctions coupled cells to their neighbors, supporting bulk propagation of electrical waves. Expression of a channelrhodopsin permitted optogenetic stimulation, and application of a far-red voltage-sensitive dye reported electrical dynamics.

In prior work, we found that the OS-HEK cells showed many interesting attributes of excitable tissues, including wave conduction, curvature-dependent wavefront velocity, and re-entrant spiral waves (28). However, the OS-HEK cells did not show beat-rate dependent changes in the action potential (AP) width, nor did they show period-doubling alternans or arrhythmia at fast pacing rates. In these regards OS-HEK cells differed from cardiomyocytes (29, 30). Arrhythmias and chaotic dynamics have not previously been observed in a synthetic bioelectrical system, suggesting that a necessary ingredient for arrhythmia was missing.

We hypothesized that this discrepancy was due to the absence of slowly recovering channels in OS-HEK cells. Upon repolarization after a spike, OS-HEK cell ion channels recovered fully within 4 ms, so each beat was independent of its predecessors. In contrast, cardiomyocytes express slowly recovering conductances, principally the hERG channel,(31, 32) which introduce memory between beats. This memory is believed to be a necessary condition for arrhythmia.(33)

We used a pharmacological approach to introduce a slowly recovering conductance into the OS-HEK cells. The sodium channel blocker isradipine shows state-dependent block of $Na_V1.7$, with a recovery time of 200 ms at –100 mV.(34) We reasoned that in the presence of isradipine, the $Na_V1.5$ channels in OS-HEK cells would show a similar slow recovery after a beat, introducing a time-dependent recovery variable and thereby achieving more cardiac-like dynamics, including the possibility of arrhythmia.

Here we characterize in detail the frequency-dependent behavior of isradipine OS-HEK (iOS-HEK) cells. In the presence of isradipine, the cells showed transitions to alternans and chaos. We used patterned cell growth to define features with different geometries in a single dish. Remarkably, we found that the transitions to alternans and chaos depended sensitively on the culture geometry. At a single pacing frequency, we simultaneously observed regular rhythms,

alternans, chaos, or depolarization block in islands that were identical in all respects except for their ₆₉ geometry. A biophysically detailed Hodgkin Huxley-style model captured these geometric effects. ₇₀ The iOS-HEK cells further showed second-degree conduction block in regions of reduced electrical ₇₁ coupling, mimicking another emergent feature of cardiac dynamics. ₇₂

Finally, we show that a similar geometry-dependent transition to arrhythmia occurs in cultured ₇₃ human iPSC-derived cardiomyocytes. These findings suggest caution in extrapolating from ₇₄ measurements in cultures to predictions of *in vivo* dynamics. Together our findings show that ₇₅ macroscopic tissue geometry is a fundamental determinant of bioelectrical dynamics, and not just a ₇₆ perturbation on the cell-autonomous behavior. We discuss which parameters are sensitive or ₇₇ insensitive to tissue geometry, and propose scaling relations that can be used to design *in vitro* ₇₈ experiments that best match the *in vivo* dynamics. ₇₉

Results ₈₀

iOS-HEK cells show alternans and arrhythmias ₈₁

Figure 1. Isradipine renders Optopatch-Spiking HEK cells susceptible to arrhythmia. **A)** Molecular components of the synthetic excitable tissue. The voltage-gated sodium channel $Na_V1.5$ and inward-rectifying potassium channel $K_{ir}2.1$ are together sufficient to produce action potentials in response to a depolarizing stimulus. CheRiff is a blue-shifted channelrhodopsin which depolarizes the cells in response to blue light, and BeRST1 is a dye which reports voltage through changes in red fluorescence. Connexin channels (Cxn) introduce electrical coupling between neighboring cells. **B)** Left: microcontact printing defines patterns of cell growth. The vertical stripes connect in a serpentine pattern outside the field of view. Blue overlay shows regions of optogenetic stimulation. Zero-dimensional "islands" are tested alongside one-dimensional "tracks". Right: single frame from a movie of BeRST1 fluorescence showing traveling waves in the track region. See **Supplementary Movie 1**. Scale bars 500 µm. **C)** Fluorescence recordings of membrane voltage showing that isradipine (10 µM) induces alternans in OS-HEK cells paced at 4 Hz. **D)** At low pace frequency isradipine OS-HEK (iOS-HEK) cells beat periodically in synchrony with the pacing, but at high pace frequency the cells produce an irregular rhythm.

In a confluent monolayer of OS-HEK cells, we observed regular spiking when the cells were optogenetically paced at 4 Hz. Addition of isradipine (10 µM) converted the spiking to an alternans rhythm (Fig. 1b), consistent with a previous report.(34) We refer to the OS-HEK cells with 10 µM isradipine as iOS-HEK cells.

Using microcontact printing,(35) we patterned cell-adhesive fibronectin features onto cytophobic polyacrylamide surfaces. Features comprised square islands of linear size 100 µm, 200 µm and 500 µm, as well as serpentine tracks of width 100, 200, 500 µm and edge length of 1 mm and 5 mm. Overall track lengths were as long as 7 cm. Cell growth followed the printed patterns (Fig. 1c). We used a digital micromirror device (DMD) to target blue light stimulation to specific regions of the sample. In a change from previous studies on OS-HEK cells, we used the far red dye BeRST1 (36) to report membrane voltage (Fig. 1a). This dye had superior brightness to QuasAr2, and showed otherwise similar response. We recorded the voltage dynamics using a custom ultrawide-field 'Firefly' microscope.(37)

Localized stimulation of the serpentine tracks induced propagating electrical waves (**Supplementary Movie 1**). Optogenetically induced waves propagated with a typical conduction velocity of 3.3 cm/s, had an AP width at 50% repolarization of 110 ms, and thus had a depolarized action potential length, $\lambda = 3.6$ mm. The fluorescence in the paced region of the track showed complex beat rate-dependent dynamics. At 2 Hz pacing, the cells spiked regularly and in phase with the drive (Fig. 1d, left). At 8.5 Hz pacing, the cells showed a complex and irregular fluorescence pattern, indicative of arrhythmia (Fig. 1d, right).

Geometry dependent instabilities in iOS-HEK cells

We characterized in detail the dependence of the electrical dynamics on local geometry. Cells were grown either in small square 'islands' or on adjacent linear 'tracks', and paced simultaneously in both geometries at frequencies between 2 and 11 Hz. The islands were paced with spatially homogeneous illumination, so cells across each island spiked synchronously and gap junction-mediated conduction did not contribute to the dynamics. We refer to these as zero-dimensional (0D) dynamics.

Tracks were stimulated in small regions (200 µm wide, 100 µm long) to induce 1D propagating waves. We observed the response in both the directly stimulated region (near field) and in the conductively stimulated region (far field; Fig. 2a). Electrical waveforms stabilized to their far-field dynamics within a distance d $\approx 0.05 \times \lambda$ from the stimulus (d = 180 µm in our experiments), so we characterized the far-field dynamics at a distance 750 µm from the stimulus. Island and track features were intermixed within each dish, were seeded with the same stock of OS-HEK cells, and were measured simultaneously. Thus any differences in dynamics between regions could be ascribed to the island geometry.

At low pace frequencies (≤ 3 Hz), all three regions spiked with a stable rhythm in synchrony with the pacing (Fig. 2b). To our surprise, at higher pace frequencies we observed dramatically different dynamics in the three regions (**Supplementary Movie 2**). The 0D islands always produced regular electrical oscillations at the pace frequency. As the pace frequency increased, the amplitude of these oscillations diminished, up to 8 Hz, beyond which the responses were barely detectable (Fig. 2).

The 1D near-field response developed a 1:1 alternans pattern at frequencies > 3 Hz. Additional bifurcations arose at 6 Hz and 8 Hz. At 10 Hz the dynamics no longer showed any repeating pattern, suggesting a transition to chaos (Fig. 2). Remarkably, the 1D far-field showed no transitions to alternans or chaos (Fig. 2b). All spikes in the far-field had nearly equal amplitude and waveform. At pace frequencies between 3 and 6 Hz, only every other spike propagated to the far field, i.e. the local beat frequency was half the pace frequency. At higher pace frequencies, a smaller portion of spikes reached the far field. These spikes had irregular timing in the near-field but conducted at velocities which gradually evened out timing variations, such that the spiking appeared regular in the far-field, always at a frequency < 3 Hz. The 1D track acted as a filter which converted high-frequency arrhythmic spiking in the near-field into lower frequency rhythmic spiking in the far field. Together these experiments gave the unanticipated result that under regular pacing, iOS-HEK cells showed irregular dynamics only in the 1D near-field, not in the 0D or 1D far-field regions.

Figure 2. Stability of spiking in iOS-HEKs cells depends on sample geometry and location of pacing. A) Cell culture geometries, with locations of optogenetic stimulus shown in blue and regions of fluorescence voltage imaging shown with dotted lines. See **Supplementary Movie 2**. **B)** Simultaneously recorded electrical dynamics in 0D, 1D near field, and 1D far field. At pacing frequencies of 5 and 10 Hz the cells showed different dynamics in the three regions. **C)** Geometry-dependent spike amplitude as a function of beat frequency. The plots for 0D and 1D near field show spike amplitude as a function of pacing frequency. The plot for 1D far field shows spike amplitude as a function of local frequency, which can be a sub-harmonic of the pacing frequency.

A Hodgkin-Huxley model captures iOS-HEK dynamics

We developed a conductance-based Hodgkin Huxley-type model to simulate the dynamics of iOS-HEK cells. The properties of $Na_V1.5$, $K_{ir}2.1$, and the channelrhodopsin CheRiff are all well known, so we were able to constrain the model with a small number of free parameters (Fig. 3a). The sodium channel model comprised Hodgkin-Huxley activation and inactivation gates m and h with dynamics taken from the literature.(38, 39) To capture the use-dependent block by isradipine, we introduced an additional slowly activating and slowly recovering gate, j (Methods). The $K_{ir}2.1$ conductance was modeled as an instantaneous function of voltage, inferred from the shape of the action potential repolarization (Methods). The channelrhodopsin was modeled as a linear conductance with a 0 mV reversal potential, modulated in space and time by the blue light illumination. A diffusive term captured the nearest-neighbor gap junction coupling. The governing equation is:

$$C_m \frac{\partial V}{\partial t} = G_{cxn} \nabla^2 V - g_{NaV} m^3 h(1-j)(V - E_{Na}) - g_{Kir}(V)(V - E_K) - g_{ChR}(x,t)(V - E_{ChR}) \quad (1)$$

where C_m is the membrane capacitance of a cell, and $G_{cxn} = g_{cxn} \times l^2$, where g_{cxn} is the gap junction conductance between cells, and l is the linear dimension of a cell. The electrical diffusion coefficient is given by G_{cxn}/C_m. To simulate 0D dynamics, g_{cxn} was set to zero. Eq. 1 represents a continuum model, which treats the domain as homogeneous tissue. In simulations, the cell discreteness is recovered by setting the spatial discretization equal to cell length.

The unknown parameters (C_m, g_cxn, g_NaV, g_Kir, g_ChR) were determined by fitting to observed fluorescence waveforms, conduction velocities, and patch clamp measurements (Methods). The free parameters in the model enter in the dynamics of the isradipine variable, j, which was assumed to bind sodium channels in their open state with rate α, and unbind with rate μ:

$$\frac{dj}{dt} = \alpha m_\infty (1 - j) - \mu j \tag{2}$$

where m_∞ is the steady state value of the m gate. The kinetic parameters in Eq. 2 were chosen to fit the experimentally observed dynamical transitions.

Action potential waveforms were simulated in a 0D geometry and in a linear 1D track comprising 2000 cells, with pacing delivered to 40 cells on one end. The near-field response was monitored in the paced zone, and the far-field response was monitored at a distance of 3 mm from the stimulus.

Simulated electrical waveforms (Fig. 3b) captured the main geometry and frequency-dependent features of the data. Specifically, the simulations in 0D showed a smooth and monotonic decrease in spike amplitude with increasing pace frequency. Simulations in the 1D near-field showed a series of frequency-dependent bifurcations that became increasingly irregular at high pace frequency. In the 1D far-field, these bifurcations were suppressed: spikes that propagated to the far-field had full amplitude, regular spacing, and were within the narrow band of frequencies that supported far-field propagation. These simulations confirm that cells with identical conductances, paced at the same frequency, can show widely divergent behavior depending upon the geometry of the surrounding excitable tissue.

Mapping the transition between the near field and far field response

We next investigated the transition from near-field to far-field behavior. How does alternans, or even chaos, in the near-field lead to regular spiking in the far-field? We mapped the fluorescence dynamics of 1D tracks as a function of distance from the stimulated zone, across a range of stimulus frequencies (Fig. 4a,b). There was a clear bifurcation in the ability of spikes to propagate into the far-field. Near-field spikes with amplitude lower than a critical threshold decayed as a function of distance, while spikes above this threshold grew to become propagating far-field spikes. Spatially resolved simulations yielded similar results (Fig. 4d). The simulations showed that the conduction velocity of each far-field spike decreased as it approached the preceding spike such that irregularities in spike spacing gradually evened out.

Figure 3. Mapping the transition from near field to far field dynamics. **A)** 1D tracks were stimulated at one end and action potential waveforms were recorded as a function of distance from the pacing stimulus. **B)**Kymographs showing electrical dynamics as a function of distance from the pacing stimulus. Some spikes decay in amplitude, while others grow in amplitude. **C)**Quantification of spike amplitude as a function of distance from the stimulus (shown in cyan). Spikes that reach the far field are shown in blue and spikes that decay are shown in red. **D)**Numerical simulations under conditions matched to the experiments. **E)**Definition of the spike decay length d, corresponding to 50% loss of amplitude. **F)**Simulated decay length for near-field spikes as a function of pacing frequency. Below a critical frequency, f*, all spikes reach the far field.

The propagation of spikes in the far-field is governed by the nonlinear dispersion relation, which defines the relationship between wavenumber and frequency (Supplement and Supplementary Fig. 3). The numerically simulated dispersion relation indicated that the far-field dynamics only supported frequencies up to f_{max} = 3.85 Hz. Waves of greater frequency could not propagate into the far field.

At pace frequencies just above the transition to near-field alternans (experimentally observed between 3 and 4 Hz), one might expect period-doubling deviations from a regular spike train to arise slowly. We defined an alternans decay length, d, as the distance over which a near-field alternans beat decayed to 50% of its initial height (Fig. 4e). Simulations near the alternans transition indeed revealed a divergence in d near the critical frequency. At pace frequencies far above the critical frequency, the alternans decay length became a small fraction of the far-field action potential length, $d \approx 0.04 \times \lambda$ (Fig. 4f), consistent with experimental results (Fig. 4c).

Second-degree conduction block in iOS-HEK cells

Conduction block and consequent arrhythmias can arise when a region of the heart acts as a partial barrier to conduction.(40) This effect can arise from spatial variations in either ion channel expression (41), or gap junctional coupling.(42, 43) We thus explored the effect of local defects on spike propagation in iOS-HEK cell tracks. In a serpentine track with sharp turns, we observed that stably propagating far-field waves sometimes failed at the turns (Fig. 5a, **Supplementary Movie 3**). Furthermore, failures occurred in a regular temporal sequence, e.g. Fig. 5a shows a pattern that after the first few beats stabilized into a 3:2 block (3 upstream spikes triggered 2 downstream spikes). The waves developed a curved wavefront and slowed repolarization at the turns, a purely geometrical consequence of the increased electrotonic loading associated with bending a wavefront around a corner. Thus geometrical effects alone are sufficient to cause conduction block, even in a background of homogeneous ion channel levels and gap junction strengths. A related effect has been reported in cultured cardiomyocytes, where a junction of a thin strand of cells to a large island showed unidirectional conduction block due to the inability of the thin strand to drive the large island.(44)

Figure 4. Curvature-induced second-degree conduction block in iOS-HEK cell tracks. **A)** Left: image of an iOS-HEK cell track showing stimulus region highlighted in cyan and regions of fluorescence monitoring shown with dashed rectangles. Right: single frame from an optical voltage recording showing two action potentials propagating through the track. See **Supplementary Movie 3**. Scale bars 200 μm. **B)** Left: kymograph showing action potential propagation with a 4 Hz pacing frequency. The vertical axis represents contour coordinate along the track. All spikes conducted into the far-field, but conduction showed second-degree block at a corner where there was a slight defect in the pattern, corresponding to the red and orange rectangles. **C)** Numerical simulations of second-degree conduction block in a 1D track with a 10 cell zone of 3-fold reduced gap junctional coupling.

In our experiments, the conduction block was attributable to a 2D wavefront curvature effect. To capture this effect in computationally tractable 1D simulations, we simulated linear tracks in which a small region (10 cells) had a reduced gap junctional coupling ($G_{cxn}' = G_{cxn}/3$). The simulated waves showed a 1:1 conduction block (Fig. 5C), qualitatively similar to that observed experimentally. These experiments and simulations show that local perturbations in the electrotonic coupling, are sufficient to lead to second-degree conduction block.

Geometry dependent instabilities in human iPSC cardiomyocytes

Finally, we explored whether the geometry-dependent effects observed in iOS-HEK cells also occurred in human iPSC-derived cardiomyocytes (hiPSC-CM). (45) Due to the significant commercial interest in using these cells as an *in vitro* model for cardiotoxicity testing (18, 19, 46-48), the correspondence (or lack of) between *in vitro* and *in vivo* arrhythmias has some practical importance.

We used microcontact printing to define side-by-side patterns of 0D islands and 1D tracks and then plated hiPSC-CM onto these patterns (Fig. 6b). CheRiff was expressed using a lentiviral vector to allow patterned blue light stimulation (27), and BeRST1 was used to image changes in membrane

Figure 5. Geometry-dependent arrhythmias in cultured human iPSC-derived cardiomyocytes (hiPSC-CM). A) Image of hiPSC-CM grown on a patterned square island. Scale bar 100 µm. **B)** Patterned hiPSC-CM cell growth in a set of 0D islands and a long 1D track. The track geometry was designed to interpose long path lengths between returns to the field of view (2 cm between red and yellow regions of interest), to account for the high propagation speed of action potentials in hiPSC-CM compared to iOS-HEK cells. Scale bar 1 mm. **C)** Simultaneously recorded electrical dynamics in 0D, 1D near field, and 1D far field. At elevated pacing frequencies the cells showed different dynamics in the three geometrical regimes. See **Supplementary Movie 4**. **D)** Quantification of the action potential height as a function of frequency in the three geometrical regimes.

potential. We optogenetically paced 0D islands and 1D tracks simultaneously across a range of frequencies. In the 1D tracks, waves propagated with a conduction velocity of 7.2 cm/s and had an AP duration of 360 ms, corresponding to a depolarized action potential length of $\lambda = 2.6$ cm.

As with the iOS-HEK cells, we observed that the dynamics depended strongly on the geometry (Fig. 6c, **Supplementary Movie 4**). At high pace frequency (3.3 Hz), the 0D islands showed small regular oscillations, the 1D near-field showed an erratic pattern of large beats with small oscillations superposed, and the 1D far-field only showed the large beats at a sub-harmonic of the pace frequency. As in the iOS-HEK cultures, the alternans beats decayed over a distance much less than the action potential length (Figure 6e; decay length $d = 535$ µm, corresponding to $d = 0.2 \times \lambda$). Thus the qualitative geometry-dependent behavior of the hiPSC-CM largely mirrored the behavior of the iOS-HEK cells.

The hiPSC-CM cells differed from the iOS-HEK cells in several important regards. First, the hiPSC-CM were spontaneously active, imposing a minimum on the optogenetic pace frequency. Second, the 0D hiPSC-CM islands showed a transition to alternans (Fig. 6c,d) which disappeared at high drive frequencies (3 Hz). Third, the 1D bifurcation to alternans was continuous in the iOS-HEK cells but discontinuous in the hiPSC-CM. In iOS-HEK simulations, these last two differences can be captured simply by tuning the isradipine unbinding rate (Fig. S1b). Simulations of the Noble model also showed clear geometry-dependent differences in the onset of instabilities (Fig. S1c).(49) Finally, real cardiac tissue can support alternans in far-field traveling waves(50), while both the iOS-HEK cells and the hiPSC-CM seemed only to support this phenomenon in the near field. Thus there

remain important dynamical features of real cardiac tissue which appear not to be captured by ^{237}
either the iOS-HEk cells or the hiPSC-CM. ^{238}

Discussion ^{239}

Gap junction-mediated currents convey the effects of boundaries to all cells in the tissue. The highly ^{240}
simplified iOS-HEK 'toy' model revealed that the qualitative dynamics depend in a sensitive way on ^{241}
the overall geometry and the nature of the pacing. Intact myocardium differs in many ways from ^{242}
cultured hiPSC-CM. However, even if one could create hiPSC-CM with fully adult single-cell ^{243}
properties, their behavior in a cultured syncytium would differ dramatically from their behavior in ^{244}
the intact heart. Thus one cannot directly infer *in vivo* behavior from *in vitro* measurements. ^{245}

Are there parameters which are invariant to gap junction coupling or sample geometry? We ^{246}
explored this question by systematically varying the gap junction strength in iOS-HEK simulations ^{247}
of a 1-D track. As anticipated from dimensional analysis, the far-field conduction velocity scaled as ^{248}
$\text{CV} \propto \sqrt{g_{cxn}}$) (Fig. S2a). The action potential duration (APD) was largely insensitive to g_{cxn}, so ^{249}
the action potential length scaled as $\lambda \propto \sqrt{g_{cxn}}$ (Fig. S2b). The transition frequency for alternans, ^{250}
f*, was also largely insensitive to g_{cxn}, varying by $< 2\%$ over a 100-fold change in g_{cxn} (Fig. S2c). ^{251}
On the other hand, f* depended sensitively on the dynamics of the slow recovery variable (Fig. S3). ^{252}
Thus the maximum stable frequency is a parameter that should be largely independent of gap ^{253}
junction strength, and by extension, tissue geometry—provided that the cells are paced via gap ^{254}
junction-mediated conduction rather than by direct pacing. ^{255}

Can one extrapolate from *in vitro* measurements of small samples with relatively weak gap ^{256}
junction coupling to larger tissues with stronger gap junctional coupling? In iOS-HEK simulations, ^{257}
the alternans decay length, d, scaled almost linearly with λ (Fig. S2d). These simulations suggest ^{258}
that the action potential length, λ, sets a natural length scale for a bioelectric tissue. Moreover, ^{259}
using λ as a scaling parameter is advantageous since this parameter can be experimentally measured ^{260}
without knowledge of gap junction conductance or other molecular parameters. ^{261}

The conduction velocity in the atria and ventricles of the human heart *in vivo* is approximately ^{262}
50 cm/s (51, 52), and action potential durations are typically 350 ms, implying an action potential ^{263}
length of $\lambda = 18$ cm. An adult human heart is approximately L = 12 cm long, so L $\approx 0.7 \times \lambda$. ^{264}
Pacemaker-triggered action potentials *in vivo* are thus primarily in the near-field and close far-field ^{265}
regimes. Remarkably, the near-field is the only regime in which we observed arrhythmias in either ^{266}
the iOS-HEK cells or in the hiPSC-CMs, either in experiment or in simulation. ^{267}

In the cultured hiPSC-CM the action potential length was $\lambda = 2.6$ cm, so to best match the ^{268}
geometrical regime of the heart, the culture should have size L $\approx 0.7 \times \lambda$, or 1.7 cm. Others have ^{269}
reported conduction velocities *in vitro* ranging from 3.5 to 20 cm/s,(15, 37, 53) corresponding to $\lambda =$ ^{270}
1.2 to 7 cm (assuming a 350 ms AP width). While cultures of size L $\approx 0.7 \times \lambda$ are easily accessible ^{271}
on the smaller end of the range, 5 cm-wide hiPSC-CM cultures are impractical. This challenge can ^{272}
be addressed either by cell patterning to produce serpentine tracks, or by adding gap junction ^{273}
blockers such as 2-APB to diminish the strength of the gap junction coupling and thereby to slow ^{274}
the conduction velocity and decrease λ. ^{275}

To best mimic activation via the sinoatrial node, cultures of hiPSC-CM should be stimulated ^{276}
locally to launch propagating waves. Whole-field stimuli, e.g. as delivered by field stimulation ^{277}
electrodes or wide-area optogenetic stimulation, will induce synchronous depolarization of all cells, ^{278}
mimicking the 0D case and missing possibly important gap junction-mediated dynamical instabilities. ^{279}
One should ideally perform spatially-resolved voltage measurements to reveal the distinct dynamics ^{280}
in the near- and far-field regimes. ^{281}

Size scaling is only a coarse approximation of the differences between *in vitro* and *in vivo* cardiac ^{282}
dynamics. Dynamics *in vivo* are further affected by spatial heterogeneity. For example, the ^{283}
conduction velocity at the atrioventricular node can be as low as 2 cm/s (43), shrinking the ^{284}
wavelength below 1 cm. This effect may be recapitulated *in vitro* by patterning the density of cells ^{285}
or by introducing geometrical constrictions. Furthermore, transmural dispersion in conduction ^{286}

velocity or repolarization can be a trigger for arrhythmia *in vivo*. While this three-dimensional effect would be difficult to model *in vitro*, one could plausibly establish in-plane gradients in cell density or gap junction strength to study such gradient effects. Finally, anisotropic conduction is critically important *in vivo*. This effect might also be captured by culturing cells on grooved or oriented substrates.

While it is not possible to recapitulate the full spatial structure of connectivity of the adult myocardium *in vitro*, one can use *in vitro* measurements to benchmark dynamical regimes according to biophysical parameters (e.g. λ and L) which can be measured *in vivo*. We expect that these considerations will be important for ongoing efforts to develop *in vitro* models of cardiac dynamics.

In this work, the use of a simplified synthetic system was crucial to identifying geometry as a determinant of dynamical stability. By comparing experimental results to a biophysically based numerical model, we demonstrated that changing tissue geometry alone could produce dramatically different responses in systems of otherwise identical excitable cells. We anticipate that this synthetic strategy may be useful in clarifying interactions in other complex excitable tissues.(54)

Materials and Methods

iOS-HEK cell line generation, culture, and patterning

Clonal OS-HEK cell lines were generated as described previously (34) and maintained in DMEM-10 supplemented with antibiotics to maintain transgene expression. For functional imaging experiments, cells were first incubated for 30 minutes in Tyrode's solution (containing 125 mM NaCl, 2 mM KCl, 2 mM CaCl$_2$, 1 mM MgCl$_2$, 10 mM HEPES, 30 mM glucose; pH 7.3, osmolality 305-310 mOsm) supplemented with 10 μg/mL isradipine and 1 μM BeRST1 voltage-sensitive dye.

Patterned cell growth was achieved using previously described methods (28, 35). Cytophilic patterns of fibronectin were deposited on cytophobic polyacrylamide gels using microcontact printing. Following pattern definition, cells were gently seeded at high density (>500k cells/mL) and allowed to adhere and proliferate throughout the pattern over several days before imaging experiments.

Wide-field all-optical electrophysiology

All-optical electrophysiology of iOS-HEK and hiPSC-CM cells was performed using an adapted 'Firefly' ultrawidefield inverted microscope.(37) Spatially patterned blue excitation for optogenetic stimulation was achieved using digital micromirror device (DMD). Near-infrared voltage sensors were excited using widefield 635 nm illumination (DILAS 8 W diode laser, M1B-638.3-8C-SS4.3-T3) configured in a near-TIRF configuration. Custom LabView software allowed for synchronization of time-modulated signals controlling blue and red light excitation, DMD patterns, and camera acquisitions.

All data were processed and analyzed using custom software (Supplementary Methods). To investigate geometry-dependent dynamical regimes, spatial regions of interest (ROIs) were defined and averaged across pixels to give fluorescence time-traces. A single set of ROIs was defined for a given sample dish and was applied uniformly across movies at different drive frequencies to systematically investigate dynamical responses (Figs. 2B and 6C). Using these spatially resolved measurements, we extract information from multiple replicates of 0D, 1D near-field, and 1D far-field responses all in a single dish.

Numerical modeling of iOS-HEK cells

In the conductance-based model, the voltage dynamics are governed by the equation

$$C_m \frac{\partial V}{\partial t} = G_{cxn}\nabla^2 V - (I_{NaV} + I_{Kir} + I_{ChR}) \tag{3}$$

Dynamics of individual currents were modeled according to well-established numerical equations with conductance magnitudes fit to experimental data. An additional slow gating variable was added to capture activity-dependent sodium channel block by isradipine (Supplementary Methods). Numerical simulations were run on isolated cells and on 1D tracks under identical model parameters for comparison to experimental results.

Acknowledgements

We thank Christopher Werley and Miao-Ping Chien for the hiPSC-CM image in Fig. 6a. This work was supported by the Howard Hughes Medical Institute. HMM was supported by the Department of Defense (DoD) through the National Defense Science & Engineering Graduate Fellowship (NDSEG) program. SD was supported by the NSF Graduate Research Fellowship Program under Grant No 1644760. Y-LH and EM were supported by NIH grant R35GM119855.

Disclosure

AEC is a co-founder of Q-State Biosciences.

Contributions

HMM and AEC designed the study. HMM conducted and analyzed experiments. HMM and SD developed and simulated the iOS-HEK numerical models. SD and BS conducted dispersion analysis and Noble model simulations. YLH and EWM provided BeRST1 dye reagent. AEC and HMM wrote the manuscript, with input from SD and BS. AEC and BS oversaw the research.

References

1. Sundelacruz S, Levin M, Kaplan DL (2009) Role of membrane potential in the regulation of cell proliferation and differentiation. Stem Cell Rev 5:231-246.

2. Roes SD, et al. (2009) Infarct tissue heterogeneity assessed with contrast-enhanced MRI predicts spontaneous ventricular arrhythmia in patients with ischemic cardiomyopathy and implantable cardioverter-defibrillator. Circ Cardiovasc Imaging 2:183-190.

3. Lindemans FW, Denier Van der Gon JJ (1978) Current thresholds and liminal size in excitation of heart muscle. Cardiovasc Res 12:477-485.

4. Cabo C, et al. (1994) Wave-front curvature as a cause of slow conduction and block in isolated cardiac muscle. Circ Res 75:1014-1028.

5. Fast VG, Kléber AG (1997) Role of wavefront curvature in propagation of cardiac impulse. Cardiovasc Res 33:258-271.

6. Bers DM (2002) Cardiac excitation–contraction coupling. Nature 415:198-205.

7. Nitsan I, Drori S, Lewis YE, Cohen S, Tzlil S (2016) Mechanical communication in cardiac cell synchronized beating12:472.

8. Ng SY, Wong CK, Tsang SY (2010) Differential gene expressions in atrial and ventricular myocytes: insights into the road of applying embryonic stem cell-derived cardiomyocytes for future therapies. Am J Physiol Cell Physiol 299:C1234-49.

9. Amin AS, Tan HL, Wilde AA (2010) Cardiac ion channels in health and disease. Heart Rhythm 7:117-126.

10. Hume J, Uehara A (1985) Ionic basis of the different action potential configurations of single guinea-pig atrial and ventricular myocytes. J Physiol (Lond) 368:525-544.

11. Nolasco J, Dahlen RW (1968) A graphic method for the study of alternation in cardiac action potentials. J Appl Physiol 25:191-196.

12. Moretti A, et al. (2010) Patient-specific induced pluripotent stem-cell models for long-QT syndrome. N Engl J Med 363:1397-1409.

13. Yazawa M, et al. (2011) Using induced pluripotent stem cells to investigate cardiac phenotypes in Timothy syndrome. Nature 471:230-234.

14. Du DT, Hellen N, Kane C, Terracciano CM (2015) Action potential morphology of human induced pluripotent stem cell-derived cardiomyocytes does not predict cardiac chamber specificity and is dependent on cell density. Biophys J 108:1-4.

15. Denning C, et al. (2016) Cardiomyocytes from human pluripotent stem cells: from laboratory curiosity to industrial biomedical platform. Biochimica Et Biophysica Acta (BBA)-Molecular Cell Research 1863:1728-1748.

16. Yang X, Pabon L, Murry CE (2014) Engineering adolescence: maturation of human pluripotent stem cell-derived cardiomyocytes. Circ Res 114:511-523.

17. Protze SI, et al. (2017) Sinoatrial node cardiomyocytes derived from human pluripotent cells function as a biological pacemaker. Nat Biotechnol 35:56-68.

18. Hoekstra M, Mummery CL, Wilde AA, Bezzina CR, Verkerk AO (2012) Induced pluripotent stem cell derived cardiomyocytes as models for cardiac arrhythmias. Front Physiol 3:

19. Colatsky T, et al (2016) The Comprehensive in Vitro Proarrhythmia Assay (CiPA) initiative — Update on progress. Journal of Pharmacological and Toxicological Methods 81: 15-20.

20. Sharma A, Wu JC, Wu SM (2013) Induced pluripotent stem cell-derived cardiomyocytes for cardiovascular disease modeling and drug screening. Stem Cell Research & Therapy 4:150.

21. Birket MJ, et al. (2015) Expansion and patterning of cardiovascular progenitors derived from human pluripotent stem cells. Nat Biotechnol 33:970-979.

22. Cherry EM, Fenton FH (2004) Suppression of alternans and conduction blocks despite steep APD restitution: electrotonic, memory, and conduction velocity restitution effects. Am J Physiol Heart Circ Physiol 286:H2332-41.

23. Cytrynbaum E, Keener JP (2002) Stability conditions for the traveling pulse: Modifying the restitution hypothesis. Chaos: An Interdisciplinary Journal of Nonlinear Science 12:788-799.

24. Clayton R, et al. (2011) Models of cardiac tissue electrophysiology: progress, challenges and open questions. Prog Biophys Mol Biol 104:22-48.

25. Bub G, Shrier A, Glass L (2002) Spiral wave generation in heterogeneous excitable media. Phys Rev Lett 88:058101.

26. Bub G, Shrier A, Glass L (2005) Global organization of dynamics in oscillatory heterogeneous excitable media. Phys Rev Lett 94:028105.

27. Werley CA, et al. (2017) Geometry-dependent functional changes in iPSC-derived cardiomyocytes probed by functional imaging and RNA sequencing. PloS One 12:e0172671.

28. McNamara HM, Zhang H, Werley CA, Cohen AE (2016) Optically Controlled Oscillators in an Engineered Bioelectric Tissue. Physical Review X 6:031001.

29. Glass L, Guevara MR, Shrier A, Perez R (1983) Bifurcation and chaos in a periodically stimulated cardiac oscillator. Physica D 7:89-101.

30. Guevara M, Ward G, Shrier A, Glass L (1984) Electrical alternans and period doubling bifurcations. IEEE Comp.Cardiol 562:167-170.

31. Lin EC, et al. (2010) Properties of WT and mutant hERG K+ channels expressed in neonatal mouse cardiomyocytes. Am J Physiol Heart Circ Physiol 298:H1842-H1849.

32. Zhang H, et al. (2012) Modulation of hERG potassium channel gating normalizes action potential duration prolonged by dysfunctional KCNQ1 potassium channel. Proceedings of the National Academy of Sciences 109:11866-11871.

33. Sanguinetti MC, Tristani-Firouzi M (2006) hERG potassium channels and cardiac arrhythmia. Nature 440:463-469.

34. Zhang H, Reichert E, Cohen AE (2016) Optical electrophysiology for probing function and pharmacology of voltage-gated ion channels. ELife 5:e15202.

35. Qin D, Xia Y, Whitesides GM (2010) Soft lithography for micro- and nanoscale patterning. Nat Protocols 5:491-502.

36. Huang Y, Walker AS, Miller EW (2015) A Photostable Silicon Rhodamine Platform for Optical Voltage Sensing. J Am Chem Soc 137:10767-10776.

37. Werley CA, Nagle SF, Ferrante JM, Wasserman SC (2017) An ultra-widefield microscope for high-speed, all-optical electrophysiology

38. ten Tusscher KHWJ, Noble D, Noble PJ, Panfilov AV (2004) A model for human ventricular tissue. Am J Physiol Heart Circ Physiol 286:H1573-H1589.

39. ten Tusscher KHWJ, Panfilov AV (2006) Alternans and spiral breakup in a human ventricular tissue model. Am J Physiol Heart Circ Physiol 291:H1088-H1100.

40. Shrier A, et al. (1987) Prediction of complex atrioventricular conduction rhythms in humans with use of the atrioventricular nodal recovery curve. Circulation 76:1196-1205.

41. Kruse M, et al. (2009) Impaired endocytosis of the ion channel TRPM4 is associated with human progressive familial heart block type I. J Clin Invest 119:2737-2744.

42. Donoghue M, et al. Heart block, ventricular tachycardia, and sudden death in ACE2 transgenic mice with downregulated connexins. J Mol Cell Cardiol 35:1043-1053.

43. Temple IP, Inada S, Dobrzynski H, Boyett MR (2013) Connexins and the atrioventricular node. Heart Rhythm 10:297-304.

44. Fast VG, Kléber AG (1995) Cardiac tissue geometry as a determinant of unidirectional conduction block: assessment of microscopic excitation spread by optical mapping in patterned cell cultures and in a computer model. Cardiovasc Res 29:697-707.

45. Yamamoto W, et al. (2016) Electrophysiological Characteristics of Human iPSC-Derived Cardiomyocytes for the Assessment of Drug-Induced Proarrhythmic Potential. PLOS ONE 11:e0167348.

46. Karakikes I, Ameen M, Termglinchan V, Wu JC (2015) Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes: Insights into Molecular, Cellular, and Functional Phenotypes. Circ Res 117:80-88.

47. Burridge PW, Keller G, Gold JD, Wu JC (2012) Production of De Novo Cardiomyocytes: Human Pluripotent Stem Cell Differentiation and Direct Reprogramming. Cell Stem Cell 10:16-28.

48. Mordwinkin NM, Burridge PW, Wu JC (2013) A review of human pluripotent stem   450
cell-derived cardiomyocytes for high-throughput drug discovery, cardiotoxicity screening, and   451
publication standards. Journal of Cardiovascular Translational Research 6:22-30.   452

49. Noble D (1962) A modification of the Hodgkin—Huxley equations applicable to Purkinje   453
fibre action and pacemaker potentials. J Physiol (Lond) 160:317-352.   454

50. Mines GR (1914) On circulating excitations in heart muscles and their possible relation to   455
tachycardia and fibrillation. Transactions of the Royal Society of Canada 4:43-52.   456

51. Draper M, Mya-Tu M (1959) A comparison of the conduction velocity in cardiac tissues of   457
various mammals. Exp Physiol 44:91-109.   458

52. Hoffman BF, DE Carvalho AP, Mello WC, Cranefield PF (1959) Electrical activity of single   459
fibers of the atrioventricular node. Circ Res 7:11-18.   460

53. Zhu H, et al. (2017) Two dimensional electrophysiological characterization of human   461
pluripotent stem cell-derived cardiomyocyte system. Scientific Reports 7:43210.   462

54. Pietak A, Levin M (2017) Bioelectric gene and reaction networks: computational modelling   463
of genetic, biochemical and bioelectrical dynamics in pattern regulation. J R Soc Interface   464
14:10.1098/rsif.2017.0425.   465

55. Hochbaum DR, et al. (2014) All-optical electrophysiology in mammalian neurons using   466
engineered microbial rhodopsins. Nat Methods 11:825-833.   467

56. Marino S, Hogue IB, Ray CJ, Kirschner DE (2008) A methodology for performing global   468
uncertainty and sensitivity analysis in systems biology. J Theor Biol 254:178-196.   469

57. Clayton R, Panfilov A (2008) A guide to modelling cardiac electrical activity in anatomically   470
detailed ventricles. Prog Biophys Mol Biol 96:19-43.   471

Supplementary Information

Supplementary Figures

Figure S1. Geometry dependent instabilities in different numerical models of excitable cells. A) Hodgkin Huxley model of iOS-HEK cells (same as Fig. 3c, repeated here for comparison to other models). **B)** Same as A) with artificially accelerated israpidine unbinding kinetics ($\mu = 0.005$ ms^{-1}). 0D features show alternans at intermediate drive frequencies. The 1D near field shows a discontinuous alternans transition. **C)** Simulation of the cardiac Noble model (49), which also shows geometry-dependent changes in dynamics.

Figure S2. Effect of gap junction coupling on dynamics and instabilities in Hodgkin-Huxley simulations of iOS-HEK cells. **A)** Linear scaling of conduction velocity with $\sqrt{g_{cxn}}$. **B)** Linear scaling of depolarized pulse length, λ, with conduction velocity. **C)** Frequency of near-field alternans onset as a function of gap junction coupling strength. The onset frequency varies by $< 2\%$ over a 100-fold variation in g_{cxn}. Pseudocolor shows the spatial extent of alternans (measured as the distance over which alternans beats decay to half of their value in the stimulus region). **D)** Relation of the alternans decay length to propagating pulse wavelength at a pacing frequency of 5 Hz, far above the onset of alternans.

Figure S3. Dispersion relation for iOS-HEK cell model. Nonlinear dispersion relations show which frequencies are stable in far-field and give wave numbers (inverse of peak-to-peak wavelength) selected by each frequency. In all figures, bottom branches of nonlinear dispersion relations are stable and the top branches are unstable. **A)** Left: nonlinear dispersion relation under conditions corresponding to experimental data. Right: spatial structure of voltage waves in the far-field with frequencies 2 and 3.5 Hz. Position on nonlinear dispersion relation indicated by correspondingly colored dots. **B)** Effect of varying the isradipine binding rate, α, on the nonlinear dispersion relation. Increasing α leads to a smaller range of frequencies that propagate to the far field. **C)** Changes in the connexin strength, g_{cxn}, affect the wavelength but have little effect on the maximum frequency prior to alternans onset. Thus the maximum frequency prior to alternans onset is a parameter that is largely independent of gap junction strength, and by extension, tissue geometry.

Supplementary movies

Movie S1. Electrical waves propagating through patterned iOS-HEK cells. Optical recordings of optogenetically triggered action potentials propagating through confluent monolayers of patterned electrically excitable iOS-HEK cells. See Fig. 1 for details.

Movie S2. Geometry dependent action potential dynamics in iOS-HEK cells. The square islands on the right produced spikes in response to each stimulus. The 1D tracks on the left produced an alternans pattern in the near-field and conducting waves only on alternating stimuli.

Movie S3. Second degree conduction block near sharp turns in an iOS-HEK serpentine track. The first four action potentials conducted into the far-field, but the fifth failed to conduct at a defect in the middle of the track. Thereafter the waves showed a 3:2 conduction block at the defect in the track.

Movie S4. Geometry dependent action potential dynamics in hiPSC-CM. The square islands on the right produced spikes in response to each stimulus. The 1D tracks on the left produced an alternans pattern in the near-field and conducting waves only on alternating stimuli.

Supplementary methods

iOS-HEK cell line generation and cell culture

All HEK cells were maintained in Dulbecco's modified Eagle medium with 10% fetal bovine serum (DMEM-10), penicillin (100 U/mL), and streptomycin (100 µg/mL). OS-HEK cells were generated as described previously.(28) Briefly, a $Na_V1.5$ plasmid (Johns Hopkins University ChemCORE) containing a puromycin selection marker was transfected into HEK cells using TransIT-293 transfection reagent (Mirus Bio). 48 hours after transfection, 2 µg/mL puromycin was introduced into the culture medium for 14 days to select for stably expressing $Na_V1.5$ cells. Following this selection, an Optopatch construct containing coding sequences of CheRiff-eGFP and QuasAr2-mOrange (separated by a self-cleaving P2A peptide sequence) was introduced using a lentiviral construct.(55) GFP-expressing cells were enriched 10 days after infection via fluorescence-activated cell sorting (FACS). Finally, a $K_{ir}2.1$ lentiviral construct containing a blasticidin selection marker was introduced. After 2 days, cell culture media was supplemented with 5 µg/mL blasticidin (to select for $K_{ir}2.1$-expressing cells) and 2 µg/mL puromycin (to ensure continued $Na_V1.5$ expression). After 14 days, cells were dispersed as single cells into a 48 well plate, and clonal populations were selected on their ability to spike robustly in response to a blue light stimulus.

Clonal cell lines were maintained in DMEM-10 supplemented with 5 µg/mL blasticidin and 2 µg/mL puromycin in 10 cm tissue culture dishes up until 80% confluence, after which they were trypsinized and cryopreserved at 500,000 cells/vial in 90% DMEM-10, 10% dimethylsulfoxide (DMSO). Cryopreserved vials were then thawed and subcultured in antibiotic-supplemented DMEM-10 in 10 cm tissue culture dishes, and were passaged via trypsinization at 80% confluence at a 1:3 ratio. A given subculture typically maintained robust blue-light induced spiking for approximately seven passages, or 2 weeks, after which cells become less excitable (presumably due to decreased $K_{ir}2.1$ expression over time).

To prepare OS-HEK cells as iOS-HEK cells for functional experiments, cells were incubated in Tyrode's solution (containing 125 mM NaCl, 2 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$, 10 mM HEPES, 30 mM glucose; pH 7.3, osmolality 305-310 mOsm) further supplemented with 10 µg/mL isradipine and 1 µM BeRST1 voltage-sensitive dye for 30 minutes in a mammalian tissue-culture incubator prior to imaging.

Cell patterning

Patterned cell growth was achieved using previously described methods(28, 35). Cytophilic fibronectin patterns were defined on a functionalized cyotophobic polyacrylamide gel using microcontact printing with patterned PDMS stamps. Patterns were first designed in software (Inkscape) and printed onto a Mylar transparency mask (CAD Art Services). Pattern negatives were then transferred onto silicon wafers coated in SU-8 3025 via contact photolithography and subsequent development of unexposed photoresist. PDMS stamps were then cast from the silicon wafer template. Functionalized polyacrylamide dishes were prepared from MatTek 35 mm-glass bottom dishes. Glass coverslips were chemically activated by plasma cleaning and then incubated for 30 min in a nitrogen-purged glovebox with silane solution (v/v: 0.5% 3-methacryloxypropyltrimethoxysilane, 2% acetic acid, 97.5% anhydrous EtOH). A 40:1 acrylamide:bisacrylamide gel doped with 4.2 mg/mL acryl-NHS (to allow for covalent bonding of fibronectic matrix) was then gelled for 2-3 minutes in ambient air under siliconized coverslips (to ensure smooth gel surfaces). Functionalized acryl-NHS dishes were then sealed in nitrogen and drierite and transferred to -80 C for long-term storage.

To complete dish preparation, patterned PDMS stamps were first coated in fibronectin protein (Yo Proteins no. 663) dissolved in PBS (0.05 mg/mL final concentration) in a sterile tissue culture hood. Coating was allowed to settle for 30 minutes, after which excess fibronectin-PBS solution was carefully removed via aspiration. Stamps were further air-dried for 10 minutes to remove excess moisture that could blur pattern transfer. Functionalized dishes were then printed with

fibronectin-coated stamps for 1 hour in a tissue-culture incubator, after which stamps were gently ₅₄₀ removed and dishes re-sterilized for 10 minutes with UV illumination. Cells were deposited by gently ₅₄₁ pipetting 500 µL DMEM-10 droplet containing the desired cell density (typically between 500,000 ₅₄₂ and 1M cells/mL). Cells were allowed to adhere for 30 minutes in a tissue culture laminar hood ₅₄₃ before an additional 2 mL of DMEM-10 was added and cells were transferred to an incubator. ₅₄₄

Wide-field all-optical electrophysiology ₅₄₅

All-optical electrophysiology of iOS-HEK and hiPSC-CM cells was performed using an adapted ₅₄₆ 'Firefly' ultrawidefield inverted microscope.(37) Spatially patterned blue excitation for optogenetic ₅₄₇ stimulation was achieved using digital micromirror device (DMD) module with an onboard 460 nm ₅₄₈ LED (Wintech DLP Lightcrafter 4500). Pixels of linear dimension 7.637 µm were demagnified 2x for ₅₄₉ optical pattern resolution of approximately 3.8 µm. Action potentials in iOS-HEK cells were ₅₅₀ triggered with 10 ms pulses at 100 mW/cm2 ; action potentials in hiPSC cardiomyocytes were ₅₅₁ triggered with 50 ms stimuli. Patterns were defined using custom software (MATLAB). DMD pixels ₅₅₂ were mapped to sample pixels by calibrating a linear transformation to a test pattern on a ₅₅₃ fluorescent target. Near-infrared voltage sensors were excited using widefield 635 nm illumination ₅₅₄ (DILAS 8 W diode laser, M1B-638.3-8C-SS4.3-T3) configured in a near-TIRF configuration (to ₅₅₅ reduce background autofluorescence). Robust signals from BeRST1 were obtained at an illumination ₅₅₆ intensity of 2 W/cm2. Near-infrared fluorescence emission was filtered using emission filters ₅₅₇ (Chroma ET665lp and Semrock quadband 336/510/581/703) and reimaged onto a sCMOS camera ₅₅₈ (Hamamatsu Orca Flash 4.2). Movies were acquired at 100 Hz over a 5 mm x 5 mm field of view. ₅₅₉ Custom LabView software allowed for synchronization of time-modulated signals controlling blue ₅₆₀ and red light excitation, DMD patterns, and camera acquisitions over experimental runs. ₅₆₁

Image processing and experimental data analysis ₅₆₂

All data were processed and analyzed using custom software (MATLAB). For each pixel, a baseline ₅₆₃ fluorescence, F, was calculated from the first percentile of the values in the recorded time-trace. The ₅₆₄ movie was then converted into units of $\frac{\Delta F}{F}$. Pixels that did not contain cells were set to zero (using ₅₆₅ a criterion that the 99th percentile of a time trace must be over a user-specified threshold). A ₅₆₆ spatial median filter (5x5 pixel filter) was further applied to account for measurement noise. At the ₅₆₇ illumination intensities used, BeRST1 photobleaching was negligible. ₅₆₈

To investigate geometry-dependent dynamical regimes, spatial regions of interest (ROIs) were ₅₆₉ defined and averaged across pixels to give fluorescence time-traces. A single set of ROIs was defined ₅₇₀ for a given sample dish and was applied uniformly across movies at different drive frequencies to ₅₇₁ systematically investigate dynamical responses (Figs. 2B and 6C). Using these spatially resolved ₅₇₂ measurements, we extract information from multiple replicates of 0D, 1D near-field, and 1D far-field ₅₇₃ responses all in a single dish. ₅₇₄

We extracted action potential amplitudes for each spike detected in each feature across movies ₅₇₅ taken at different pacing frequencies. Spike upstrokes were detected via a threshold on the time ₅₇₆ derivative. Spike amplitudes (in units $\frac{\Delta F}{F}$) were defined as the difference between the maximum ₅₇₇ value after a given upstroke and the minimum value preceding the upstroke. Between 10 and 300 ₅₇₈ spikes per feature were collected in a given movie, depending on drive frequency and acquisition ₅₇₉ time. Beeswarm plots of spike amplitudes at each pacing frequency were used to visualize the ₅₈₀ dynamical patterns of activity (Figs. 2C, 6E). Only spikes taken from the second half of movies are ₅₈₁ visualized to avoid early transients which decay as patterns converges to a stable cycle. The drive ₅₈₂ frequency in 0D and 1D near field measurements was set by the frequency of the blue light-gated ₅₈₃ CheRiff activation; in 1D far-field measurements, the pacing frequency is defined as the local ₅₈₄ frequency (i.e., the reciprocal of the interval between successive spikes). ₅₈₅

The transition between the near-field and far-field region was mapped using kymographs (Figures ₅₈₆ 4B and 5B). Kymographs were generated by spatially averaging $\frac{\Delta F}{F}$ movies across the short ₅₈₇ dimension of a linear track. For serpentine tracks (Figure 5), tracks segments were first ₅₈₈

computationally aligned into a single long track, and then spatially averaged along the short 589
dimension. Alternans transitions were spatially mapped by performing spike detection (as described 590
above) across linear windows of 8 pixels (Fig. 4C). 591

Numerical modeling of iOS-HEK cells 592

In the conductance-based model, the voltage dynamics are governed by the equation 593

$$C_m \frac{\partial V}{\partial t} = G_{cxn} \nabla^2 V - (I_{NaV} + I_{Kir} + I_{ChR}) \tag{4}$$

where $V = V(x, t)$ is the voltage in mV. Recall $G_{cxn} = g_{cxn} \times l^2$, where g_{cxn} is the gap junction 594
conductance between cells, and l is the linear dimension of a cell. The ionic currents drive local 595
dynamics, while the diffusion term couples neighboring regions. Units of space are 10^{-5} m 596
(corresponding to linear size of one cell), and time is in ms. Conductances are measured in nS/pF 597
and ionic currents in pA/pF. The currents are: 598

$$I_{NaV} = g_{NaV} m_\infty^3(V) h (1 - j)(V - E_{Na}) \tag{5}$$

$$I_{Kir} = g_{Kir}(V)(V - E_K) \tag{6}$$

$$I_{ChR} = g_{ChR}(x, t)(V - E_{ChR}) \tag{7}$$

The gating variables for the sodium activation and inactivation gates, m and h respectively, and 599
for the effect of isradipine, j, are dimensionless variables which take values between 0 and 1. The 600
inactivation gate evolves following 601

$$\frac{\partial h}{\partial t} = \frac{1}{\tau_h(V)} (h_\infty(V) - h) \tag{8}$$

To capture the use-dependent sodium block by isradipine, we introduced a slowly activating and 602
recovering gate, j, with behavior governed by 603

$$\frac{\partial j}{\partial t} = \alpha m_\infty(V)(1 - j) - \mu j \tag{9}$$

That is, the drug binds sodium channels in its open state with rate α, and unbinds with rate μ. 604
These kinetic parameters were chosen to fit the experimentally observed dynamical transitions. 605
Since the time constant of the sodium activation gate, m, is orders of magnitude faster than the 606
other gates, the gate was replaced with the asymptotic limit, $m_\infty(V)$. This standard approximation 607
allowed for more efficient computations without impacting the simulation results. 608
The gating variables approach steady-state asymptotic values, which are functions of the voltage.

$$h_\infty(V) = \left(1 + \exp\left(\frac{V + 71.55}{7.43}\right)\right)^{-1} \tag{10}$$

$$j_\infty(V) = \frac{\alpha m_\infty(V)}{\alpha m_\infty(V) + \mu} \tag{11}$$

$$m_\infty(V) = \left(1 + \exp\left(\frac{-56.86 - V}{9.03}\right)\right)^{-2} \tag{12}$$

Time constants for each gate tell how quickly the behaviors approach the asymptotic values, and 609
are given by 610

$$\tau_h(V) = \frac{1}{2}\left(1 - \tanh\left(\frac{V+40}{10}\right)\right)\tau_{h1}(V) + \frac{1}{2}\left(1 + \tanh\left(\frac{V+40}{10}\right)\right)\tau_{h2}(V) \tag{13}$$

$$\tau_{h1}(V) = \left(0.057\exp\left(\frac{-V-80}{6.8}\right) + 2.7\exp(0.079V) + (3.1\times10^5)\exp(0.3485V)\right)^{-1} \tag{14}$$

$$\tau_{h2}(V) = \frac{0.13\left(1 + \exp\left(\frac{-V-10.66}{11.1}\right)\right)}{0.77} \tag{15}$$

$$\tau_j(V) = (\alpha m_\infty(V) + \mu)^{-1} \tag{16}$$

The equations governing the sodium channel activation, m, and inactivation, h, gates were set following literature values.(38, 39) Minor modifications were made to the h time constant, $\tau_h(V)$, to smooth the discontinuous function. The channelrhodopsin was modeled as a linear conductance, modulated in space, x and time, t, by the blue light illumination. The repolarization conductance g_{Kir} was modeled as an instantaneous function of voltage derived from a smoothing spline fit to the action potential waveform at low drive frequency (f = 2 Hz).

Numerical simulations were run on single-cells for the 0D case, and 2 cm 1D tracks with Neumann (no-flux) boundary conditions at each end. The 1D tracks were discretized with a spatial step size, $dx = 10$ µm, the length of one OS-HEK cell. With this discretization size, each grid point represented one cell, allowing the simulations to have similar spatial resolution as the experiments.

The Laplacian was modeled with a fourth-order centered finite difference scheme. Boundary conditions were implemented using the standard ghost-point method. A combination IMEX scheme was used for the time evolution, with implicit Crank-Nicholson applied to the diffusion terms and explicit Adams-Bashforth applied to the remaining nonlinear terms (ion currents and gating dynamics). A time step of 0.1 dx ms (10^{-6} ms in the case of 10 µm spatial step) was used, which is well within the accuracies of the numerical scheme and is significantly smaller than any time scales in the model. Single cell, 0D dynamics were modeled by setting g_{cxn} to zero, and time evolution was performed with the same Adams-Bashforth scheme. All computations were performed using MATLAB.

Initial model parameters were experimentally constrained and fit using a combination of patch clamp measurements and fluorescent voltage recordings. Reversal potentials of $E_{Na} = 75$ mV, $E_K = -107$ mV, and $E_{ChR} = 0$ mV were determined according to Nernst equation for the corresponding intracellular and extracellular ionic concentrations. All conductances were expressed as specific conductances per unit capacitance, allowing us to set $C_m = 1$. The sodium conductance g_{NaV} was set to 1.5 nS/pF, corresponding to a maximum transient current of 3 nA at -25 mV ($V - E_{Na} = -100$ mV) measured in a 20 pF patch clamp recording. The channelrhodopsin conductance was determined to be 15 pS/pF following similar methodology. The inward rectifying potassium current g_{Kir} was represented as an instantaneous function of voltage, determined by fitting to the experimentally observed fluorescence repolarization waveform at low drive frequency (2 Hz). Because fluorescence traces do not report absolute voltage, we assumed a resting potential of -90 mV and peak voltage of +30 mV, in accordance with patch clamp recordings. The diffusion coefficient G_{cxn} is given by $G_{cxn} = g_{cxn} \times l^2$. where $l = 1$ cell. The connexin conductance g_{cxn} was estimated according to measurements of the static electronic coupling length (28) as 40 nS/pF.

The parameters α and μ were chosen to match experimentally determined dynamical regimes. Setting $\alpha = 0.5$ ms^{-1} and $\mu = 0.015$ ms^{-1} gave close agreement with the complex landscape of dynamical transitions observed experimentally (Fig. 3c) as well as simulated voltage traces which closely resembled experimental observations (Fig. 3b). Sensitivity analysis was performed to validate these parameter choices (Supplementary Methods).

Model dynamics were characterized using analogous parameters as for experimental results. Simulations were conducted for both an isolated cell which was directly paced ('0D') as well as for linear tracks of 2000 cells (i.e. $L = 2$ cm) in which 40 cells were paced on one end. Simulations were

run across a range of stimulus intervals between 500 ms (2 Hz) and 100 ms (10 Hz), sampled at 10 ms intervals. For each numerical experiment, we detected spike upstrokes as upward deflections of the derivative of the simulated voltage trace.

Sensitivity analysis of model parameters

Parameter values for α, μ, g_{NaV}, and g_{cxn} were selected to match as closely as possible the experimental results in both the 0D and 1D settings. The model was simulated for 1,500 ms with a given set of parameters under 2 Hz stimuli, and the resulting voltage at times t_k, $V(x, t_k)$, was compared to analogous experimental results, $W(x, t_k)$, using the ordinary least squares (OLS) objective function

$$OLS_{Obj} = \sum_k |W(x, t_k) - V(x, t_k)|^2 \tag{17}$$

Simulated and experimental voltage values were scaled to be between 0 and 1 to allow for direct comparison. An optimal set of parameters minimizes the objective function. Minimization was performed with the MATLAB constrained optimization function fmincon. Realistic parameter constraints were chosen from the patch clamp data.

Sensitivity of model parameters was tested using Latin Hypercube Sampling (LHS)(56). The LHS results identified α, μ,and, g_{NaV} as the most sensitive parameters, with g_{cxn} as the least. That is, small changes in the connexin strength, g_{cxn}, had negligible impact on the simulated voltages. Regardless, changing any model parameters over a modest regime maintained the main geometry-dependent differences in stability, giving confidence in the appropriateness of the model.

Nonlinear dispersion analysis of far-field dynamics

In the far-field, the waves become spatially periodic structures of constant shape and velocity, both of which are uniquely determined by the frequency. The nonlinear dispersion relation of the far-field equation indicates which wave number (inverse of peak-to-peak wavelength) is selected by each frequency (Supplementary Fig. 3A). The nonlinear dispersion relation is computed by considering the far-field traveling waves as periodic stationary solutions in a moving frame, which allows for the traveling wave and wave number to be solved for as a function of frequency. The full dispersion relation curve is traced out with pseudo-arclength numerical continuation implemented in MATLAB, and using the spatial discretization scheme as in the numerical simulations.

Each frequency has two associated wave numbers, with the bottom branch stable and top branch unstable. Two examples of the spatial structure of the far-field waves are given in Supplementary Fig. 3A, and their positions along the dispersion relation marked. The curve does not extend past 3.85 Hz, indicating the far-field equation does not support waves of higher frequencies. Therefore, for drive frequencies above 3.85 Hz, each stimulus will not propagate into the far-field, leading to the appearance of near-field alternans, and aligning with the experimentally and numerically observed behavior.

Changing the drug binding rate, α, and connexin strength, g_{cxn}, have different impacts on the nonlinear dispersion relation (Supplementary Fig. 3B and 3C). Small changes in α lead to large shifts in the transition frequency, indicating that faster binding rates will induce near-field alternans at lower drive frequencies. However, large changes to the coupling strength change properties of the traveling waves, such as wave number, wave length, and speed, but do not appreciably change the range of allowed frequencies or transition frequency. Changes to model parameters μ (drug unbinding rate) and g_{NaV} (sodium conductance) are not shown, but have a similar effect as changing α.

The 2:1 conduction block observed from local changes in the connexin strength (Fig. 5) is also explained from the nonlinear dispersion relation. As the spike propagates through the small region of lower connexin strength, the wavenumber and frequency are adjusted to values that are no longer stable when the wave reenters the region of higher connexin strength (observe the differences in the

curves with g_{cxn} values of 40 and 14 in Supplementary Fig. 3C). The modification of wavenumber and frequency upon the reentry results in the observed 2:1 conduction block.

Noble model of cardiomyocyte dynamics

To demonstrate that the observed geometry effects are not unique to the iOS-HEK cell model and are of importance in cardiac dynamics, we performed 0D and 1D simulations in the Noble model. The Noble model is an ionic channel model for Purkinje fibers and the voltage dynamics are governed by sodium, potassium, and background chloride currents (I_{Na}, I_K, and I_{Cl}, respectively). Spatial coupling between cells was again incorporated through a Laplacian.

$$C_m \frac{\partial V}{\partial t} = \delta \nabla^2 V - (I_{Na} + I_{K1} + I_{K2} + I_{Cl} + I_{ext}) \tag{18}$$

$$I_{Na} = \left(\bar{g}_{Na} m^3 h + g_{Na} \right) (V - E_{Na}) \tag{19}$$

$$I_{K1} = \left[g_k \exp \left(\frac{-V - 90}{50} \right) + 0.015 \exp \left(\frac{V + 90}{60} \right) \right] (V - E_k) \tag{20}$$

$$I_{K2} = g_K n^4 (V - E_k) \tag{21}$$

$$I_{Cl} = g_{Cl}(V - E_{Cl}) \tag{22}$$

$$I_{ext} = g_{stim}(V - E_{ChR}) \tag{23}$$

The dynamics of the dimensionless ionic gates for sodium activation, m, sodium deactivation, h, and potassium, n, take the form

$$\frac{dz_i}{dt} = \alpha_i(V)(1 - z_i) - \beta_i(V)z_i \tag{24}$$

where z_i stands for m, h, n, and α_i and β_i similarly correspond to opening and closing rates of the m, h, and n channels. The rates are given by:

$$\alpha_m(V) = 0.1(V+48) \left(1 - \exp \left(\frac{-V - 48}{15} \right) \right)^{-1} ; \beta_m(V) = 0.12(V+8) \left(\exp \left(\frac{V + 8}{5} \right) - 1 \right)^{-1} \tag{25}$$

$$\alpha_h(V) = 0.17 \exp \left(\frac{-V - 90}{20} \right) ; \beta_h(V) = \exp \left(\frac{-V - 42}{10} \right)^{-1} \tag{26}$$

$$\alpha_n(V) = -0.0001(V + 50) \left(\exp \left(\frac{-V - 50}{10} \right) - 1 \right)^{-1} ; \beta_n(V) = 0.002 \left(\exp \left(\frac{-V - 90}{80} \right) \right)^{-1} \tag{27}$$

All ionic currents, dynamic gating variables, and parameter values were set from literature. Membrane capacitance, conductances, and Nernst potentials are given in Table 1.(49, 57) The form of the applied current, I_{ext}, was set to mimic the channelrhodopsin stimulus. The Noble model was chosen for its ability to reproduce realistic behaviors observed in cardiac cells and for the relatively low number of ionic currents it includes, allowing for efficient simulation.

Numerical simulations in 0D and 1D geometries used the same centered finite-difference Laplacian and IMEX time evolution schemes described above. Simulations were run on 1D tracks of up to 6 cm in length with spatial grid spacing set to 0.1 mm and time step of 0.01 ms. Due to the higher cell coupling and conduction values, the stimulus duration was increased to 50 ms (following

$C_m = 12\mu$ F/cm^2	$g_{\text{stim}} = 0.5$ mS/cm^2
$E_{\text{Na}} = 40$ mV	$\bar{g}_{Na} = 400$ mS/cm^2
$E_{\text{K}} = -100$ mV	$g_{\text{K}}=1.2$ mS/cm^2
$E_{\text{Cl}} = -60$ mV	$g_{\text{Cl}}=0.075$ mS/cm^2
$\delta = 0.001$ cm^2/ms	

Table 1. Values of membrane capacitance, conductances, and Nernst potentials used in simulations of the Noble model.

hiPSC-CM experiments). Simulations were run for at least two forcing cycles before recording results to remove any transient effects caused by the self-spiking behavior of the Noble model. Model output was analyzed in a similar fashion to the OS-HEK model results.

The Noble model simulations also exhibit geometry-dependent changes in stability (Supplementary Fig. 1C). The 1D near-field results show a dramatic transition first to alternans at 4.5 Hz, and then to chaotic-like dynamics at 8 Hz. Similar transitions are not observed in the 0D or far-field results, which instead display monotonically decreasing spike amplitude for higher frequencies. As before, the model selects a narrow range of wave frequencies which propagate into the far-field.