Title: Maturing Mycobacterial Peptidoglycan Requires Non-canonical Crosslinks

to Maintain Shape

Authors: Catherine Baranowski[1], Michael A. Welsh[2], Lok-To Sham[2,3], Haig A.

Eskandarian[4,5], Hoong C. Lim[2], Karen J. Kieser[1], Jeffrey C. Wagner[1], John D.

McKinney[4], Georg E. Fantner[5], Thomas R. Ioerger[6], Suzanne Walker[2], Thomas G.

Bernhardt[2], Eric J. Rubin[1,2]*, E. Hesper Rego[7]*

Affiliations:

[1] Department of Immunology and Infectious Disease, Harvard T. H. Chan School of Public

Health, Boston, MA 02115, USA; [2] Department of Microbiology and Immunobiology,

Harvard Medical School, Boston, MA 02115, USA; [3] Department of Microbiology and

Immunology, National University of Singapore, 117545 Singapore; [4] School of Life

Sciences, Swiss Federal Institute of Technology in Lausanne (EPFL), 1015 Lausanne,

Switzerland; [5] School of Engineering, Swiss Federal Institute of Technology in Lausanne

(EPFL), 1015 Lausanne, Switzerland; [6] Department of Computer Science and

Engineering, Texas A&M University, College Station, Texas, USA;[7] Department of

Microbial Pathogenesis, Yale University School of Medicine, New Haven, Connecticut,

06510, USA.

* to whom correspondence should be addressed: erubin@hsph.harvard.edu;

hesper.rego@yale.edu

Abstract:

In most well studied rod-shaped bacteria, peptidoglycan is primarily crosslinked by penicillin binding proteins (PBPs). However, in mycobacteria, L,D-transpeptidase (LDT)-mediated crosslinks are highly abundant. To elucidate the role of these unusual crosslinks, we characterized mycobacterial cells lacking all LDTs. We find that LDT-mediated crosslinks are required for rod shape maintenance specifically at sites of aging cell wall, a byproduct of polar elongation. Asymmetric polar growth leads to a non-uniform distribution of these two types of crosslinks in a single cell. Consequently, in the absence of LDT-mediated crosslinks, PBP-catalyzed crosslinks become more important. Because of this, *Mycobacterium tuberculosis* (Mtb) is more rapidly killed using a combination of drugs capable of PBP- and LDT- inhibition. Thus, knowledge about the single-cell distribution of drug targets can be exploited to more effectively treat this pathogen.

Introduction:

Peptidoglycan (PG) is an essential component of all bacterial cells (Vollmer, Blanot, & de Pedro, 2008a), and the target of many antibiotics. PG consists of linear glycan strands crosslinked by short peptides to form a continuous molecular cage surrounding the plasma membrane. This structure maintains cell shape and protects the plasma membrane from rupture. Our understanding of PG is largely derived from studies on laterally growing model rod-shaped bacteria like *Escherichia coli* and *Bacillus subtilis* (*Figure 1 - figure supplement 1A*). In these organisms, new PG is constructed along the

lateral side wall by the concerted effort of glycoslytransferases, which connect the glycan

of a new PG subunit to the existing mesh, and transpeptidases, which link peptide side

chains. An actin-like protein, MreB, positions this multi-protein complex along the short

axis of the cell so that glycan strands are inserted circumferentially, creating

discontinuous hoops of PG around the cell (Domínguez-Escobar et al., 2011; Garner et

al., 2011). This orientation of PG creates a mechanical anisotropy that is responsible for

rod shape (Hussain et al., 2018).

However, not all rod-shaped bacteria encode MreB. In fact, there are important

differences between model bacteria and Actinobacteria like mycobacteria, a genus of rod-

shaped bacteria that includes the major human pathogen *Mycobacterium tuberculosis*

(Mtb). In mycobacteria, new PG is inserted at the cell poles (at unequal amounts based

on pole age), rather than along the lateral walls (*Figure 1A*). Additionally, mycobacteria

are missing several factors, including MreB, that are important for cell elongation (Kieser

& Rubin, 2014). Furthermore, in *E. coli* and *B. subtilis* the vast majority (>90%) of the

peptide linkages are created by D,D-transpeptidases known as penicillin binding proteins

(PBPs) (Pisabarro, de Pedro, & Vázquez, 1985). PBPs, the targets of most β-lactams,

link the 4th amino acid of one peptide side chain to the 3rd amino acid of another, forming

4-3 crosslinks (*Figure 1- figure supplement 1B*). Peptide linkages can also be catalyzed

by L,D-transpeptidases (LDTs), which link peptide side chains by the 3rd amino acid

forming 3-3 linkages. In mycobacteria, these 3-3 crosslinks, are highly abundant,

accounting for ~60% of linkages (Kumar et al., 2012) (*Figure 1- figure supplement 1B*).

Because PG has been most well studied in bacteria where 3-3 crosslinks are rare, the

role of these linkages, and the enzymes that catalyze their formation, is poorly understood.

Tuberculosis remains an enormous global health problem, in part, because treating even drug susceptible disease is difficult. The standard regimen includes a cocktail of four drugs given over six months. Treatment of drug-resistant Mtb is substantially longer and includes combinations of up to seven drugs ("Global Tuberculosis Report 2017," 2017). While some of the most important anti-mycobacterials target cell wall synthesis, surprisingly, drugs that target PG are not part of the core treatment for either drug-susceptible or drug-resistant disease. However, carbapenems, β-lactam antibiotics that potently inhibit LDTs *in vitro*, are effective against drug resistant Mtb *in vitro* and drug sensitive Mtb in patients (Diacon et al., 2016; Hugonnet, Tremblay, Boshoff, Barry, & Blanchard, 2009).

But, why are LDTs important in mycobacteria? To explore this, we constructed strains that lack the ability to form 3-3 crosslinks. We find that 3-3 crosslinks are formed in maturing peptidoglycan and that they are necessary to stabilize the cell wall and prevent lysis. Cells that lose the ability to synthesize 3-3 crosslinks have increased dependence on 4-3 crosslinking. Thus, simultaneous inhibition of both processes results in rapid cell death.

Results:

Fluorescent D-amino acids are incorporated asymmetrically by LD-transpeptidases

PG uniquely contains D-amino acids, which can be labeled with fluorescent probes

(fluorescent D-amino acids, FDAAs) to visualize PG synthesis in live bacterial cells (Kuru

et al., 2012). When we incubated *M. smegmatis* (Msm), a non-pathogenic relative of Mtb,

with FDAAs for a short 2-minute pulse (< 2% of Msm's generation time) we observed

incorporation at both poles, the sites of new PG insertion in mycobacteria (*Figure 1A, B*)

(Aldridge et al., 2012). However, we also saw a gradient of fluorescence along the

sidewalls, extending from the old pole (the previously established growth pole) that fades

to a minimum at roughly mid-cell as it reaches the new pole (the pole formed at the last

cell division) (*Figure 1B*).

To identify the enzymes responsible for this unexpected pattern of lateral cell wall

FDAA incorporation, we performed a fluorescence-activated cell sorting (FACS)-based

transposon screen (*Figure 1C*). Briefly, we stained an Msm transposon library with FDAA

and repeatedly sorted the least fluorescent 12.5% of the population by FACS. After each

sort we regrew cells, extracted gDNA and used deep sequencing to map the location of

the transposons found in the low staining population.

From this screen, we identified three LDTs (*ldtA - MSMEG_3528, ldtB -*

MSMEG_4745, ldtE - MSMEG_0233) (*Figure 1D*) that were primarily responsible for

FDAA incorporation. Deleting these three LDTs significantly reduced FDAA incorporation

(*Figure 1- figure supplement 2A, B*). To further investigate the physiological role of LDTs,

we constructed a strain lacking all 6 LDTs (Δ*ldtAEBCGF*, hereafter ΔLDT). FDAA

incorporation and 3-3 crosslinking are both nearly abolished in ΔLDT cells and can be

partially restored by complementation with a single LDT (*ldtE*-mRFP; ΔLDT$_{comp}$) (*Figure*

1E, F, Figure 1- figure supplement 2C, 3). Thus, as might be the case in *Bdellovibrio*

(Kuru et al., 2017), FDAA incorporation in Msm is primarily LDT-dependent. LDTs have

previously been shown to exchange non-canonical D-amino acids onto PG tetrapeptides

in *Vibrio cholera* (Cava, de Pedro, Lam, Davis, & Waldor, 2011).

LDT catalyzed 3-3 crosslinks are required for rod shape maintenance at aging cell wall

As deletion of a subset of LDTs in Msm produces morphologic changes (Sanders,

Wright, & Pavelka, 2014), we visualized ΔLDT cells by time-lapse microscopy. We

observed that a subpopulation of cells loses rod shape progressively over time, resulting

in localized spherical blebs (*Figure 2A - top, Figure 2 - figure supplement 1A, Figure 2 -*

video 1). Complemented cells are able to maintain rod shape (*Figure 2 - figure*

supplement 1B). We reasoned that localized loss of rod shape may occur for two reasons:

1) spatially-specific loss of cell wall integrity and/or 2) cell wall deformation due to

uncontrolled, local PG synthesis. If the first hypothesis were true, high osmolarity should

protect cells against forming blebs. Indeed, switching cells from iso- to high- osmolarity

prevented bleb formation over time (*Figure 2A, bottom, Figure 2- video 2*). To test the

second hypothesis, we stained ΔLDT cells with an amine-reactive dye, and observed

outgrowth of new, unstained material (*Figure 2B*). Blebs retained stain, indicating a lack

of new cell wall synthesis in the region. Collectively, these results indicate that 3-3

crosslinks are required to counteract turgor pressure and maintain the rod shape of

mycobacteria. This led us to hypothesize that bleb formation is a result of a local defect

in cell wall rigidity.

To directly measure cell wall rigidity, we used atomic force microscopy (AFM) on

live ΔLDT and WT cells. We measured the rigidity of cells in relation to their height.

Generally, WT cells are stiffer than ΔLDT cells (*Figure 2D*). Blebs in ΔLDT cells can be

identified by a sharp increase in height (*Figure 2E, pink shaded*). Since circumferential

stress of the rod measured by AFM is proportional to the radius of the cell, and inversely

proportional to the thickness of the cell wall (an immeasurable quantity by AFM), we used

cell height, a proxy for radius, to normalize the stiffness measurement. We found that

stiffness drops in the area of blebs (*Figure 2E, pink shaded*).

Why does loss of rod shape occur locally and only in a subpopulation of cells?

Mycobacterial polar growth and division results in daughter cells with phenotypic

differences (Aldridge et al., 2012). For example, the oldest cell wall is specifically inherited

by the new pole daughter (*Figure 2- supplement 2A*, (Aldridge et al., 2012)). We

hypothesized that the loss of rod shape might occur in specific progeny generated by cell

division. Indeed, the daughter which inherited the new pole from the previous round of

division, and the oldest cell wall, consistently lost rod shape over time, while the old pole

daughter maintained rod shape (*Figure 2C, Figure 2- supplement 2B*). In addition, blebs

localized to the oldest cell wall (*Figure 2B*), as visualized by pulse-chase labeling of the

cell wall. Thus, 3-3 crosslinking is likely occurring in the oldest cell wall, which is non-

uniformly distributed along a single cell and in the population via asymmetric polar growth

and division. Taken together, these data suggest that LDTs act locally to reinforce aging

PG and to maintain rod shape in a subpopulation of Msm cells - specifically, new pole

daughters (*Figure 2F*).

Mycobacteria are hypersensitive to PBP inactivation in the absence of LDTs

Our observations lead to the following model (*Figure 6A*): PBP-catalyzed 4-3 crosslinks are formed at the poles where new PG is inserted and where pentapeptide substrates reside. These newly synthesized 4-3 crosslinks are then gradually cleaved (by D,D-endopeptidases) as PG ages and moves toward the middle of the cell, leaving tetrapeptide substrates for LDTs to create 3-3 crosslinks. This is consistent with the FDAA incorporation pattern, which reflects the abundance of tetrapeptide substrates available for LDT exchange. Specifically, there are more available tetrapeptides near the poles and fewer near mid-cell, the site of older PG (*Figure 1F*). In the absence of LDTs to catalyze 3-3 crosslinks, old cell wall loses integrity and turgor pressure causes bleb formation.

This model predicts that ΔLDT cells should be even more dependent on 4-3 crosslinking than wild type cells. To test this hypothesis, we used TnSeq (Long et al., 2015) to identify genes required for growth in cells lacking LDTs (*Figure 3A*). We found that mutants of two PBPs, *pbpA (MSMEG_0031c)* and *ponA2 (MSMEG_6201)*, were recovered at significantly lower frequencies in ΔLDT cells (*Figure 3B*). Likewise, using allele swapping (Kieser, Boutte, et al., 2015b) (*Figure 3C, Figure 3- figure supplement 1A*), a technique that tests the ability of various alleles to support viability, we found that the transpeptidase (TP) activity of PonA1, which is non-essential in WT cells (Kieser, Boutte, et al., 2015b), becomes essential in ΔLDT cells (*Figure 3D*). Thus, cells that lack 3-3 crosslinks are more dependent on 4-3 crosslinking enzymes.

Peptidoglycan synthesizing enzymes localize to differentially aged cell wall

Given our model, we hypothesized that enzymes catalyzing and processing

different types of crosslinks should be differentially localized along the length of the cell.

Specifically, we postulated that 4-3 crosslinking PBPs would localize at sites of new PG,

while 4-3 cleaving D,D-endopeptidases and 3-3 crosslinking LDTs would localize to sites

of older PG. To test whether 3-3 and 4-3 crosslinking enzymes localize differently, we

visualized fluorescent fusions of a PBP (PonA1), and an LDT (LdtE), (*Figure 4A*). We

found that PonA1-RFP largely localized to the old pole, where new PG is inserted (*Figure

4A, B, Figure 4 - video 1*). LdtE-mRFP localized farther from the poles, the sites of older

PG (*Figure 4A, B, Figure 4 - video 2*). Thus, enzymes responsible for 4-3 and 3-3

crosslinks exhibit distinctive subcellular localizations, consistent with the model that they

act on differentially aged PG.

We next sought to localize a D,D-endopeptidase. As no D,D-endopeptidase has

been clearly identified in mycobacteria, we used HHPRED (Zimmermann et al., 2017) to

find candidates. By homology to the *E. coli* protein AmpH, an enzyme with both D,D-

carboxy- and endopeptidase activity (González-Leiza, de Pedro, & Ayala, 2011), we

identified DacB2 (MSMEG_2433), a protein previously shown to have D,D-

carboxypeptidase activity in Msm (Bansal et al., 2015), as a candidate to also harbor D,D-

endopeptidase capability. We expressed and purified DacB2 and found that it, like AmpH,

had both D,D-carboxypeptidase and D,D-endopeptidase activity on peptidoglycan

substrates generated *in vitro* (*Figure 4C, Figure 4- figure supplement 1A-C*). Consistent

with our model, DacB2-mRFP localized closer to LDT-mRFP, farther from the poles, at

sites of older PG (*Figure 4A, B, Figure 4 – video 3*). Thus, bleb formation may result from

unchecked D,D-endopeptidase activity in ΔLDT cells.

Drugs targeting both PBPs and LDTs synergize to kill Mtb

The importance of 3-3 crosslinks in mycobacteria suggests a unique vulnerability. While Mtb can be killed by most non-carbapenem (N-C) β-lactams like penicillins, which largely target the PBPs, carbapenem β-lactams, which target both PBPs and LDTs (Mainardi et al., 2007; Papp-Wallace, Endimiani, Taracila, & Bonomo, 2011) are also effective against Mtb (Diacon et al., 2016; Hugonnet et al., 2009). As has been previously proposed (Gonzalo & Drobniewski, 2013; Gupta et al., 2010; Mainardi et al., 2007), our data suggest that more rapid killing of Mtb could be achieved with drug combinations that target both PBPs and LDTs (*Figure 5A*). In fact, we find that the combination of amoxicillin (a penicillin) and meropenem (a carbapenem), exhibits synergism in minimal inhibitory concentration (Σ Fractional Inhibitory Concentration < 0.5 ("Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution Methods," 2016), *Figure 5B, Figure 5- figure supplement 1A,B*). We also see that this combination leads to more rapid killing of Mtb *in vitro* (*Figure 5B, Figure 5- figure supplement 2A*) (Andreu, Fletcher, Krishnan, Wiles, & Robertson, 2012).

Discussion:

The success of antibiotics that target PG, like β-lactams, has led to decades of research of this critical bacterial polymer. Recently developed fluorescent probes (FDAAs) of PG synthesis have been used extensively to study PG-synthesis in live cells of numerous bacterial species. Intriguingly, these probes can be incorporated through diverse pathways in different bacteria and thus, their pattern can mark distinct processes (Kuru et al., 2012). Here, we find that, in mycobacteria, FDAA incorporation is primarily

LDT-dependent. FDAA incorporation in Msm shows an unusual gradient pattern (Botella

et al., 2017), suggesting an asymmetric distribution of tetrapeptide substrate for the LDT-

dependent exchange reaction. In addition to their ability to exchange D-amino acids onto

tetrapeptides, LDTs also catalyze non-canonical 3-3 crosslinks.

LDT-catalyzed crosslinks are rare in model rod-shaped bacteria like *E.coli* and *B.*

subtilis but, are abundant in polar growing bacteria like mycobacteria, *Agrobacterium*

tumefaciens and *Sinorhizobium meliloti* (Brown et al., 2012; Cameron, Zupan, &

Zambryski, 2015; Kumar et al., 2012; Lavollay et al., 2008; Pisabarro et al., 1985)*.* Here,

we find that Msm cells lacking 3-3 crosslinks cannot maintain rod shape at sites of aging

cell wall. PBP-catalyzed 4-3 crosslinks appear able to maintain rod shape near the poles,

the sites of newer cell wall (*Figure 6A*). Over time, older cell wall moves towards the

middle of the cell, loses structural stability, and begins to bleb. The gradual manner in

which rod shape is lost suggests that cell wall processing must occur to de-stabilize this

portion of the rod. Consistent with this idea, we find that an enzyme that cleaves 4-3

crosslinks, DacB2, also localizes to sites of old cell wall.

Why would Msm cells create 4-3 crosslinks to eventually cleave them? There are

many possibilities. For example, perhaps in the absence of lateral cell wall synthesis, the

creation of substrate for LDTs through the destruction of 4-3 crosslinks allows the cell to

engage the PG along the lateral cell body. This could be important for altering the

thickness of the PG layer or anchoring it to the membrane at sites of otherwise "inert" cell

wall. Additionally, it may be that as PG ages, it is being manicured for septal synthesis.

Supporting this idea, we find that the localization patterns of PonA1, LdtE and DacB2,

and FDAA all have local minima at mid-cell closer to the new pole, the asymmetric site of

division in mycobacteria (Aldridge et al., 2012; Dhar, Bousbaine, Wakamoto, McKinney,

& Santi, 2013; Eskandarian et al., 2017). This lack of localization suggests a lack of penta-

and tetra- peptide substrates, implying that that this region of the cell more abundantly

crosslinked, as crosslinking utilizes these peptide species. Mycobacteria are missing

known molecular septal placement mechanisms like the Noc and the Min system (Hett &

Rubin, 2008). Could 3-3 crosslinking be a signal for septal placement? Indeed, the major

septal PG hydrolase is RipA, a D,L-endopeptidase which cleaves the bond between the

second and third amino acid of PG side chains, a substrate available on 3-3 crosslinked

material (Böth, Schneider, & Schnell, 2011; Vollmer, Joris, Charlier, & Foster, 2008b).

Thus, it is intriguing to speculate that 3-3 crosslinks could be important for localizing cell

division machinery.

In well-studied rod-shaped bacteria like *E.coli* and *B. subtilis*, shape is maintained

by MreB-directed PG synthesis along the lateral cell body (Garner et al., 2011; Hussain

et al., 2018; Ursell et al., 2014). On the other hand, mycobacteria maintain shape in the

absence of an obvious MreB homolog, and in the absence of lateral cell wall elongation.

Furthermore, in contrast to lateral-elongating bacteria, in which new and old cell wall are

constantly intermingled during growth, polar growth segregates new and old cell wall

(*Figure 6B*). We find that mycobacteria appear to utilize 3-3 crosslinks at asymmetrically

distributed aging cell wall to provide stability along the lateral body, something that may

not be required in the presence of MreB-directed PG synthesis.

New drug combinations for TB are desperately needed. Indeed, there has been a

renewed interest in repurposing FDA-approved drugs for TB treatment (Diacon et al.,

2016). Our results suggest that a better understanding of the localization and distribution

of drug targets in single cells may lead to rational predictions of drug combinations. Indeed, we find that enzymes which do very similar chemistry – PG crosslinking – are distributed differentially in a single cell and across the population. In the absence of 3-3 crosslinks, 4-3 crosslinks become more important for cell viability. These data predict that a drug combination which inhibits both PBPs and LDTs will work synergistically to more quickly kill Mtb, a prediction we verified *in vitro*. Interestingly, meropenem combined with amoxicillin/clavulanate resulted in early clearance of Mtb from patient sputum (Diacon et al., 2016). In fact, the combination might be key to accelerated killing of Mtb (Gonzalo & Drobniewski, 2013).

Acknowledgments:

We thank Cara Boutte and Erkin Kuru for discussion and thorough manuscript review. We are grateful to the Rubin and Fortune laboratory for discussion and input. We also thank the Microbiology and Immunobiology department at Harvard Medical School for sharing equipment and reagents, as well as the BSL3 staff at the Harvard School of Public health for tremendous support. This work was supported by the National Institutes of Health U19 AI107774 to E.J.R and T.R.I., F32AI104287 to E.H.R (who is also supported by a Career Award at the Scientific Interface from BWF), R01 GM76710 to S.W., R01AI083365 and U19AI109764 to T.G.B.. M.A.W. is supported by an F32 GM123579. LT.S. was supported by an American Heart Association Postdoctoral fellowship (14POST18480014). H.C.L is funded by a Simons Foundation Fellow of the Life Sciences Research Foundation award. This work was also supported in part by the Swiss National Science Foundation (310030_156945) and the Innovative Medicines Initiative (115337)

to J.D.M., and from the Swiss National Science Foundation (205321_134786 and

205320_152675), the European Union FP7/2007-2013/ERC Grant Agreement No.

307338 (NaMic), and EU-FP7/Eurostars E!8213 to G.E.F. Support for H.A.E. comes from

a European Molecular Biology Organization Long Term Fellowships (191-2014 and 750-

2016). K.J.K. was supported by the National Science Foundation Graduate Research

Fellowship (DGE1144152, DGE0946799).

Figure 1: FDAAs are incorporated asymmetrically by LD-transpeptidases. (A)
Schematic of mycobacterial asymmetric polar growth. Green, old cell wall; grey, new
material; dotted line, septum; large arrows, old pole growth; small arrows, new pole
growth. **(B)** FDAA incorporation in log-phase WT Msm cell after 2-minute incubation.
Scale bar=5μm. Old pole marked with (*). **(C)** Schematic of Fluorescence Activated Cell
Sorting (FACS)-based FDAA transposon library enrichment. An Msm transposon (Tn)
library was stained with FDAAs, the dimmest and brightest cells were sorted, grown,
sorted again to enrich for transposon mutants that are unable or enhanced for FDAA
incorporation. **(D)** Results from 1C screen. For each gene, the contribution to low or
high staining population was calculated from transposon reads per gene. Plotted is the
ratio of the population contribution from the second sort of low FDAA staining (L2) over
the second sort of high FDAA staining (H2) cells compared to the Mann-Whitney *U* P-
value. **(E)** Representative image of FDAA incorporation in log-phase WT, ΔLDT and
ΔLDT$_{comp}$ cells. Scale bar= 5μm. **(F)** Profiles of FDAA incorporation in log-phase WT
(N=98), ΔLDT (N=40), and ΔLDT$_{comp}$ (N=77) cells. Thick lines represent mean
incorporation profile, thin lines are FDAA incorporation in single cells.

321
322

Figure 2: LDT catalyzed 3-3 crosslinks are required for rod shape maintenance at aging cell wall. (A) Msm ΔLDT time-lapse microscopy of cells switched from high- to iso- (top) osmolar media, or iso- to high osmolar media (bottom). (high=7H9+150mM sorbitol; iso=7H9). t= time in minutes post osmolarity switch. **(B)** ΔLDT cells were stained with Alexa 488 NHS-ester (green) to mark existing cell wall, washed, and visualized after outgrowth (unstained material). A, B scale bar=2µm. **(C)** Maximum cell width of ΔLDT cell lineages over time. Width of new pole daughters = blue circle; width of old pole daughters = orange circle. Division signs denote a division event. At each division, there are two arrows from the dividing cell leading to the resulting new and old pole daughter cell widths (blue and orange respectively). **(D)** Mean stiffness of WT (N=73) and ΔLDT (N=47) Msm cells as measured by atomic force microscopy. Mann-Whitney U P-Value **** < 0.0001. **(E)** Representative profile of cell height and height-normalized stiffness (modulus/height) in a single ΔLDT cell. Pink shaded portion highlights location of a bleb. **(F)** Model of rod shape loss in old cell wall of ΔLDT cells compared to WT. Green portions of the cell represents old cell wall; grey portion represents new cell wall. The larger arrows indicate more growth from the old pole, while smaller arrows show less relative growth from the new pole. Dotted lines represent septa. op= old pole, np=new pole.

340
341
342
343

344
345
Figure 3: Mycobacteria are hypersensitive to PBP inactivation in the absence of
LDTs. (A) Fold change in the number of reads for transposon insertion counts in ΔLDT
cells compared to WT Msm. P-value is derived from a rank sum test (DeJesus,
Ambadipudi, Baker, Sassetti, & Ioerger, 2015). **(B)** Transposon insertions per TA
dinucleotide in *pbpA* and *ponA2* in WT (grey) and ΔLDT (blue) cells. **(C)** Schematic of
L5 allele swapping experiment. Adapted from (Kieser, Boutte, et al., 2015b). **(D)** Results
of WT or transpeptidase null *ponA1* allele swapping experiment in ΔLDT cells.
353

Figure 4: **Peptidoglycan synthesizing enzymes localize to differentially aged cell wall. (A)** Representative fluorescence images of PonA1-RFP (magenta), LdtE-mRFP (cyan), and DacB2-mRFP (green). Scale bars=5µm. **(B)** Average PonA1-RFP (N=24),

LdtE-mRFP (N=23) or DacB2-mRFP (N=23) distribution in cells before division. **(C)** Schematic of the *in vitro* experiment to test D,D-carboxy- and D,D-endopeptidase activity of DacB2. Lipid II extracted from *B. subtilis* is first polymerized into linear (using SgtB) or crosslinked (using *B. subtilis* PBP1) peptidoglycan and then reacted with DacB2. The reaction products are analyzed by LC-MS. **(below)** Extracted ion chromatograms of the reaction products produced by incubation of DacB2 with peptidoglycan substrates.

Figure 5: Drugs targeting both PBPs and LDTs synergize to kill Mtb. (A) Schematic of synergistic targeting of PBPs and LDTs by pencillin and carbapenem β-lactams. **(B)** Minimum inhibitory concentration (MIC) of amoxicillin, meropenem, or the combination, in *M. tuberculosis*. FIC (fractional inhibitory concentration) = MIC of drug in combination/MIC of drug alone. Synergy is defined as ∑ FIC<=0.5 ("Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution Methods," 2016). **(C)** Killing dynamics of *M. tuberculosis* (expressing the *luxABCDE* operon from *Photorhabdus luminescens* (Andreu et al., 2010)) measured via luciferase production (RLU=relative light units). 5X MIC Amoxicillin (AM) (3.125 µg/mL); 5X MIC Meropenem (MR) (6.25 µg/mL); Amoxicillin+Meropenem 3.125 µg/mL AM; 6.25 µg/mL MR). Biological triplicate are plotted. All drugs were used in combination with 5µg/mL clavulanate.

Figure 6: Model for PG enzyme and substrate distribution as governed by polar growth and PG segregation by age. (A) A model for PG age, PG enzyme and crosslink segregation via polar growth in mycobacteria. **(B)** Schematic of PG segregation by age (top). 2-minute FDAA pulse (cyan), 45-minute outgrowth, followed by 2-minute FDAA chase (magenta) in WT Msm cells (bottom). Newest cell wall (magenta), older cell wall (cyan). Scale bar=5μm

  Supplementary Materials

Materials and Methods

Bacterial strains and culture conditions

Unless otherwise stated, *Mycobacterium smegmatis* (mc^{2}155) was grown shaking at

37°C in liquid 7H9 media consisting of Middlebrook 7H9 salts with 0.2% glycerol,

0.85g/L NaCl, ADC (5g/L albumin, 2g/L dextrose, 0.003g/L catalase), and 0.05% Tween

80 and plated on LB agar. *Mycobacterium tuberculosis* (H37Rv) was grown in liquid

7H9 with OADC (oleic acid, albumin, dextrose, catalase) with 0.2% glycerol and 0.05%

Tween 80*.* Antibiotic selection for *M. smegmatis* and *M. tuberculosis* were done at the

following concentrations in broth and on agar: 25μg/mL kanamycin, 50μg/mL

hygromycin, 20μg/mL zeocin and 20μg/mL nourseothricin and, twice those

concentrations for cloning in *Escherichia coli* (TOP10, XL1-Blue and DH5α).

Strain construction

M. smegmatis mc^{2}155 mutants lacking *ldtABECFG (*ΔLDT) was constructed using

recombineering to replace endogenous copies with zeocin or hygromycin resistance

cassettes flanked by lox sites as previously described (Boutte et al., 2016). Briefly, 500

base pairs of upstream and downstream sequence surrounding the gene of interest

were amplified via PCR (KOD Xtreme^{TM} Hot Start DNA polymerase (EMD Millipore,

Billerica, MA)). These flanking regions were amplified with overlaps to either a zeocin or

hygromycin resistance cassette flanked by loxP sites and these pieces were assembled

into a deletion construct via isothermal assembly (Gibson et al., 2009). Each deletion

cassette was transformed into Msm expressing inducible copies of RecET for

recombination (Murphy, Papavinasasundaram, & Sassetti, 2015). Once deletions were

verified by PCR using and sequencing, the antibiotic resistance cassettes were

removed by the expression of Cre recombinase. The order of deletion construction in

the ΔLDT strain was as follows (where arrows represent transformation of a Cre-

recombinase plasmid, followed by curing of the Cre-recombinase plasmid as it contains

the *sacB* gene for sucrose counter selection on LB supplemented with 10% sucrose,

and strain names are listed in parenthesis). This resulted in the removal of antibiotic

cassettes flanked by loxP sites:

1) mc^{2}155Δ*ldtA*:: zeo^R (KB103)→ mc^{2}155Δ*ldtA*::loxP (KB134)

2) mc^{2}155Δ*ldtA::loxP* + Δ*ldtE*:: zeo^R (KB156)

3) mc^{2}155Δ*ldtA::loxP* Δ*ldtE*:: zeo^R + Δ*ldtB*:: hyg^R (KB200) → mc^{2}155Δ*ldtA::loxP*

Δ*ldtE*::loxP Δ*ldtB*::loxP (KB207)

4) mc^{2}155Δ*ldtA::loxP* Δ*ldtE*::loxP Δ*ldtB*::loxP + Δ*ldtC*:: hyg^R (KB209)

5)mc^{2}155Δ*ldtA::loxP* Δ*ldtE*::loxP Δ*ldtB*::loxP Δ*ldtC*:: hyg^R Δ*ldtG*:: zeo^R (KB222)→

mc^{2}155Δ*ldtA::*loxP Δ*ldtE*::loxP Δ*ldtB*::loxP Δ*ldtC*:: loxP Δ*ldtG*:: loxP (KB241)

6) mc^{2}155Δ*ldtA::*loxP Δ*ldtE*::loxP Δ*ldtB*::loxP Δ*ldtC*:: loxP Δ*ldtG*:: loxP Δ*ldtF*:: hyg^R

(KB303 referred to as ΔLDT) .

M. tuberculosis H37Rv was transformed with a vector expressing the codon optimized

Photorhabdus luminescens luxABCDE operon (pMV306hsp+LuxG13 –Addgene

#26161)(Andreu et al., 2010). This strain is referred to as Mtb Lux.

Refer to Supplemental Table 1 for oligonucleotides, and Supplemental Table 2 for a

complete strain list.

ΔLDT complementation plasmid construction

To complement ΔLDT we placed a copy of *ldtE* (*MSMEG_0233*) under the constitutive

TetO promoter (a UV15 derivative within a pMC1s plasmid that is inducible with

anhydrous tetracycline in the presence of a tet-repressor TetR, which the ΔLDT$_{comp}$

strain lacks (Kieser, Boutte, et al., 2015b)) on vector that integrates at the L5 phage

integration site of the chromosome of the ΔLDT strain (the vector is marked with

kanamycin resistance). A glycine, glycine, serine linker was cloned between *ldtE* and

mRFP in this complementation construct.

PonA1 transpeptidase essentiality L5 allele swapping

To test essentiality of transpeptidation by PonA1 in the ΔLDT cells, L5 allele swapping

as described in (Kieser, Boutte, et al., 2015b) was performed. The plasmids used in

this experiment were previously published in (Kieser, Boutte, et al., 2015b). Briefly, a

wild-type copy of PonA1 (TetO driven expression, L5 integrating and kanamycin

marked) was transformed into ΔLDT. Then, the endogenous copy of *ponA1* was

replaced with zeocin using the above mentioned recombineering technique (amplifying

the construct from a previously published deletion mutation of *ponA1*(Kieser, Boutte, et

al., 2015b)). Swapping efficiency of either wildtype or transpeptidase mutant PonA1

marked with nourseothricin was tested with a transformation into ΔLDT//L5-TetO-ponA1

(WT)-kanamycin.

M. tuberculosis minimum inhibitory concentration (MIC) determination

Mtb Lux was grown to log phase and diluted to an OD_{600}=0.006 in each well of non-

treated 96 well plates (Genessee Scientific) containing 100µL of meropenem (Sigma

Aldrich) and/or amoxicillin (Sigma Aldrich) diluted in 7H9+OADC+5µg/mL clavulanate

(Sigma Aldrich). Cells were incubated in drug at 37°C shaking for 7 days, 0.002%

resazurin (Sigma Aldrich) was added to each well, and the plates were incubated for 24

hours before MICs were determined. Pink wells signify metabolic activity and blue

signify no metabolic activity. (Kieser, Baranowski, et al., 2015a) Checkerboard MIC

plates and fractional inhibitory concentrations were calculated as described in

("Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution

Methods," 2016).

M. tuberculosis drug killing assays

Mtb Lux was grown to log phase (kanamycin 25µg/mL) and diluted in 30mL inkwells

(Corning Lifesciences) to an OD_{600}=0.05 in 7H9+OADC+5µg/mL clavulanate with

varying concentrations of amoxicillin, meropenem, or both. 100µL of these cultures

were pipetted in duplicate into a white 96-well polystyrene plate (Greiner Bio-One) and

luminescence was read in a Synergy H1 microplate reader from BioTek Instrumenmts,

Inc. using the Gen5 Software (2.02.11 Installation version). The correlation between

relative light units (RLU) and colony forming units (CFU) is shown in Msm in Figure 5 –

figure supplement 2**.**

Fluorescent D-amino acid labeling

NADA (3-[7-nitrobenzofurazan]-carboxamide-D-alanine), HADA (3-[7-hydroxycoumarin]-

carboxamide-D-alanine) and TADA (3-[5-carboxytetramethylrhodamine]-carboxamide-

D-alanine) were synthesized by Tocris following the published procotol (Kuru, Tekkam,

Hall, Brun, & Van Nieuwenhze, 2014). To 1mL of exponentially growing cells 0.1mM of

FDAA final was added and incubated for 2 minutes before washing in 7H9 twice. For

still imaging, after the second wash, cells were fixed in 7H9 + 1% paraformaldehyde

before imaging. For pulse chase experiments, cells were stained, washed with 7H9 and

allowed to grow out for 40 minutes before being stained with a second dye and imaged.

Flow cytometry

An *M. smegmatis* transposon library was grown to mid-log phase, and stained with 2

µg/mL NADA for 2 minutes. Cells were centrifuged and half of the supernatant was

discarded. The pellet was resuspended in the remaining supernatant, passed through a

10µm filter and taken to be sorted (FACSAria; Excitation: 488nm; Emission filter:

530/30). Two bins were drawn at the lowest and highest staining end of the population,

representing 12.5% of the population. 600,000 cells from these bins were sorted into

7H9 medium. Half of this was directly plated onto LB agar supplemented with

kanamycin to select for cells harboring the transposon. The remaining 300,000 cells

were grown out in 7H9 to log phase, stained with FDAA and sorted again to enrich the

populations.

Transposon sequencing, mapping and FDAA flow cytometry enrichment analysis.

Genomic DNA (gDNA) was harvested from the sorted transposon library colonies and

transposon-gDNA junction libraries were constructed and sequenced using the Illumina

Hi-Seq platform (Long et al., 2015). Reads were mapped on the *M. smegmatis* genome,

tallied and reads at each TA site for the bins (low/high incorporating sort 1 and 2) were

imported into MATLAB and processed by a custom scripts as described in (Rego,

Audette, & Rubin, 2017).

Microscopy

Both still imaging and time lapse microscopy were performed on an inverted Nikon TI-E

microscope at 60x magnification. Time lapse was done using a CellASIC (B04A plate)

with constant liquid 7H9 flow in a 37°C chamber. For turgor experiment (*Figure 2A*),

cells were grown in either 7H9 or 7H9 500mM sorbitol overnight, and then switched to

either 7H9 with 150mM sorbitol (high osmolar) or to 7H9 alone (iso-osmolar).

Atomic force microscopy

AFM experimentation was conducted as previously(Eskandarian et al., 2017). In short,

polydimethylsiloxane (PDMS) – coated coverslips were prepared by spin-coating a

mixture of PDMS at a ratio of 15:1 (elastomer:curing agent) with hexane (Sigma

296090) at a ratio of 1:10 (PDMS:hexane) (Koschwanez, Carlson, & Meldrum, 2009;

Thangawng, Ruoff, Swartz, & Glucksberg, 2007). A 50 µl filtered (0.5 µm pore size

PVDF filter – Millipore) aliquot of bacteria grown to mid-exponential phase and

concentrated from 2-5 ml of culture was deposited onto the hydrophobic surface of a

PDMS-coated coverslip and incubated for ~20 min to increase surface interactions

between bacteria and the coverslip. 7H9 medium (~3 ml) was supplied to the sample

so as to immerse the bacterial sample and the AFM cantilever in fluid. The AFM

imaging mode, Peak Force QNM, was used to image bacteria with a Nanoscope 5

controller (Veeco Metrology) at a scan rate of 0.5 Hz and a maximum Z-range of 12 μm.

A ScanAsyst fluid cantilever (Bruker) was used. Height, peak force error, DMT

modulus, and log DMT modulus were recorded for all scanned images in the trace and

retrace directions. Images were processed using Gwyddion (Department of

Nanometrology, Czech Metrology Institute). ImageJ was used for extracting bacterial

cell profiles in a tabular form.

Correlated optical fluorescence and AFM

Correlated optical fluorescence and AFM images were acquired as described

(Eskandarian et al., 2017). Briefly, optical fluorescence images were acquired with an

electron-multiplying charge-coupled device (EMCCD) iXon Ultra 897 camera (Andor)

mounted on an IX81 inverted optical microscope (Olympus) equipped with an

UPLFLN100XO2PH x100 oil immersion objective (Olympus). Transmitted light

illumination was provided by a 12V/100W AHS-LAMP halogen lamp. An U-MGFPHQ

fluorescence filter cube for GFP with HQ-Ion-coated filters was used to detect GFP

fluorescence. The AFM was mounted on top of the inverted microscope, and images

were acquired with a Dimension Icon scan head (Bruker) using ScanAsyst fluid

cantilevers (Bruker) with a nominal spring constant of 0.7 N m^{-1} in Peak Force QNM

mode at a force setpoint ~1nN and typical scan rates of 0.5 Hz. Indentation on the cell

surface was estimated to be ~10 nm in the Z-axis. Optical fluorescence microscopy

was used to identify Wag31-GFP puncta expressed in a wild-type background(Dhar et

al., 2013) in order to distinguish them from cells of the ΔLDT mutant strains.

Calculating cell surface rigidity

A cell profile was extracted from AFM Height and DMT Modulus image channels as

sequentially connected linear segments following the midline of an individual cell. A

background correction was conducted to by dividing the DMT modulus values of the cell

surface by the mean value of the PDMS surface and rescaled to compare the cell

surface rigidity between individual cells from different experiments. The DMT modulus

reflects the elastic modulus (stress-strain relationship) for each cross-sectional

increment along the cell length.

Analysis of fluorescent protein distribution

Using a segmented line, profiles of cells from new to old pole were created at the frame

"pre-division" based on physical cell separation of the phase image. A custom FIJI

(Schindelin et al., 2012) script was run to extract fluorescence line profiles of each cell

and save them as .csv files. These .csv files were imported to Matlab where a custom

script was applied to normalize the fluorescence line profile to fractional cell length and

to interpolate the fluorescence values to allow for averaging.

Analysis of cell wall distribution

Cells were stained with Alexa488 NHS ester as described previously (Aldridge et al.,

2012) and followed via time-lapse microscopy in the CellASIC device. Briefly, 1mL of

log phase cells was pelleted at 8,000 rpm for 1 minute and washed with 1mL PBST.

The pellet was resuspended in 100uL of PBST and 10uL Alexa Fluor 488 carboxylic

acid succinimidyl ester was added for a final concentration of 0.05mg/mL. This was

incubated for 3 minutes at room temperature. Stained cells were pelleted for 1 minute at

13,000 rpm and washed with 500μL PBST. They were spun again and resuspended in

7H9 for outgrowth observation over time in the CellASIC device.

<u>Analysis of FDAAs</u>

Images were analyzed using a combination of Oufti (Paintdakhi et al., 2016) for cell

selection followed by custom coded Matlab scripts to plot FDAA fluorescence over

normalized cell length, calculate cell length and bin cells by existence of an FDAA

labeled septum.

<u>Generation of transposon libraries</u>

M. smegmatis cells were transduced at (OD$_{600}$ 1.1-1.7) with φMycoMarT7 phage

(temperature sensitive) that has a Kanamycin marked Mariner transposon as previously

described (Long et al., 2015). Briefly, mutagenized cells were plated at 37°C on LB

plates supplemented with Kanamycin to select for phage transduced cells. Roughly

100,000 colonies per library were scraped, and genomic DNA was extracted.

Sequencing libraries were generated specifically containing transposon disrupted DNA.

Libraries were sequenced on the Illumina platform. Data were analyzed using the

TRANSIT pipeline (DeJesus et al., 2015).

<u>Peptidoglycan isolation and analysis</u>

600mLs of wildtype and ΔLDT cells were grown to log phase and collected via

centrifugation at 5000 x g for 10 minutes at 4°C. The resulting pellet was resuspsended

in PBS and cells were lysed using a cell disruptor at 35,000psi twice. Lysed cells were

boiled in 10% SDS (sodium dodecyl sulfate) for 30 minutes and peptidoglycan was

collected via centrifugation at 17,000 x g. Pellets were washed with 0.01% DDM(*n*-

Dodecyl β-D-maltoside) to remove SDS and resuspended in 1XPBS + 0.01% DDM. PG

was digested with alpha amylase (Sigma A-6380) and alpha chymotrypsin (Amersco

0164) overnight. The samples were again boiled in 10% SDS and washed in 0.01%

DDM. The resulting pellet was resuspended in 400μL 25mM sodium phosphate pH6,

0.5mM MgCl2, 0.01% DDM. 20μL of lysozyme (10mg/mL) and 20μL 5U/μL

mutanolysoin (Sigma M9901) were added and incubated overnight at 37°C. Samples

were heated at 100°C and centrifuged at 100,000 x g. 128μL of ammonium hydroxide

was added and incubated for 5 hours at 37°C. This reaction was neutralized with 122μL

of glacial acetic acid. Samples were lyophilized, resuspended in 300μL 0.1% formic acid

and subjected to analysis by LC-MS/MS. Peptide fragments were separated with an

Agilent Technologies 1200 series HPLC on a Nucleosil C18 column (5μm 100A 4.6 x

250mm) at 0.5mL/min flow rate with the following method: Buffer A= 0.1% Formic Acid;

Buffer B=0.1% Formic Acid in acetonitrile; 0% B from 0-10 minutes, 0-20% B from 10-

100 minutes, 20% B from 100-120 minutes, 20-80% B from 120-130 minutes, 80% B

from 130-140 minutes, 80%-0% B from 140- 150minutes, 0% B from 150-170 minutes.

MS/MS was conducted in positive ion mode using electrospray ionization on an Agilent

Q-TOF (6520).

<u>Expression and purification of MSMEG_2433 (DacB2)</u>

MSMEG_2433 was expressed and purified using a modified method for purification of

low molecular weight PBPs that was previously published (Qiao et al., 2014). An N-

terminally truncated MSMEG_2433$_{(29-296)}$ was cloned into the pET28b vector for isopropyl

β-D-1-thiogalactopyranoside (IPTG) inducible expression in *E. coli* BL21 (DE3). 10mLs

of overnight culture grown in LB with Kanamycin (50μg/mL) were diluted 1:100 into 1 L of

LB with Kanamycin (50μg/mL) and grown at 37°C until an OD$_{600}$ of 0.5. The culture was

cooled to room temperature, induced with 0.5mM IPTG, and shaken at 16°C overnight.

Cells were pelleted via centrifugation at 4,000rpm for 20 min at 4°C. The pellet was

suspended in 20mL binding buffer (20mM Tris pH 8, 10mM MgCl$_2$, 160mM NaCl, 20mM

imidazole) with 1mM phenylmethylsulfonylfluoride (PMSF) and 500μg/mL DNase. Cells

were lysed via three passage through a cell disrupter at ≥ 10,000 psi. Lysate was pelleted

by ultracentrifugation (90,000 × g, 30 min, 4°C). To the supernatant, 1.0mL washed Ni-

NTA resin (Qiagen) was added and the mixture rocked at 4°C for 40 min. After loading

onto a gravity column, the resin was washed twice with 10mL wash buffer (20mM Tris pH

8, 500mM NaCl, 20mM imidazole, 0.1% Triton X-100). The protein was eluted in 10mL of

elution buffer (20 mM Tris pH8, 150mM NaCl, 300mM imidazole, 0.1% reduced Triton X-

100) and was concentrated to 1 mL with a 10kD MWCO Amicon Ultra Centrifuge Filter.

The final protein concentration was measured by reading absorbance at 280 nm and

using the estimated extinction coefficient (29459 M^{-1}cm^{-1}) calculate concentration. The

protein was diluted to 200µM in elution buffer with 10% glycerol, aliquoted, and stored at

-80°C.

Proper folding of purified MSMEG_2433$_{(29-296)}$ was tested via Bocillin-FL binding. Briefly,

20µM of purified protein was added to penicillin G (100, 1000 U/mL in 20mM K_2HPO_4,

140mM NaCl, pH7.5) in a 9µL reaction. The reaction was incubated at 37°C for 1hour.

10µM Bocillin-FL was added and incubated at 37°C for 30 minutes. SDS loading dye

was added the quench the reaction and samples were loaded onto a 4-20% gel.

MSMEG_2433$_{(29-296)}$ bound by Bocillin-FL was imaged using a Typhoon 9400 Variable

Mode Imager (GE Healthcare) (Alexa Excitation-488nm Emission-526nm).

<u>Lipid II extraction.</u>

B. subtilis Lipid II was extracted as previously published (Qiao et al., 2017).

<u>SgtB purification.</u>

S. aureus SgtB was purified as previously published (Rebets et al., 2014).

<u>Purification of *B. subtilis* PBP1.</u>

Purification of *B. subtilis* PBP1 was carried out as previously described (Lebar et al.,

2014).

<u>*In vitro* Lipid II polymerization and crosslinking.</u>

20 µM purified BS Lipid II was incubated in reaction buffer (50 mM HEPES pH 7.5, 10

mM $CaCl_2$) with either 5 µM PBP1 or 0.33 µM SgtB for 1 hour at room temperature. The

enzymes were heat denatured at 95°C for 5 minutes. Purified MSMEG_2433(29-296) was

added (20 uM, final) and the reaction was incubated at room temperature for 1 hour.

Mutanolysin (1 μL of a 4000 U/mL stock) was added and incubated for 1.5 hours at 37

°C (twice). The resulting muropeptides were reduced with 30 μL of NaBH$_4$ (10 mg/mL)

for 20 minutes at room temperature with tube flicking every 5 minutes to mix. The pH

was adjusted to ~4 using with 20% H$_3$PO$_4$ and the resulting product was lyophilized to

dryness. The residue was resuspended in 18μL of water and analyzed via LC-MS as

previously reported (Welsh et al., 2017).

Figure 1- figure supplement 1: Peptidoglycan synthesis and FDAA screen overview. (A) *Escherichia coli* and *Bacillus subtilis* lateral cell wall growth. Unlike mycobacteria, *E. coli* and *B. subtilis* insert new cell wall along the lateral cell body, mixing old and new peptidoglycan. Green portion represents old cell wall; grey portion represents new material. **(B)** Cartoon of penicillin binding proteins (PBPs), L,D-transpeptidases (LDTs), and both 4-3 and 3-3 crosslinking. PBPs utilize a pentapeptide substrate found on new peptidoglycan, ending in D-alanine-D-alanine. LDTs utilize a tetrapeptide substrate found on processed peptidoglycan. TP, transpeptidase; TG, transglycosylase

PEAK	Tripeptides observed	Description*
1	461.2	
	Tetrapeptides observed	
2	532.3	Tetra (2 NH2)
3	534.2	Tetra (OH)
4	533.2	Tetra (NH2)
	pentapeptides observed	
5	604.3	Penta (OH)
6	602.3	Penta(2 NH2)
	4-3 observed	
7	974.5	4--3 (2 NH2)
8	1045.4	4--4 (4 NH2)
9	1046.5	4--4 (3 NH2)
10	1049.4	4--4
11	975.4	4--3 (NH2)
12	974.3	4--3 (2 NH2)
13	1047.5	4--4 (2 NH2)
	3-3 observed	
14	974.4	4--3 (4 NH2)
15	904.4	3--3 (3 NH2)
16	975.5	4--3 (3 NH2)
	Tripeptide(anhydrous)	
17	865.4	
	4--4--4	
18	1559.7	
19	1560.0	

3-3 crosslink (tripeptide-tripeptide)
m/z: 904.44 (100.0%), 905.44 (38.9%),
906.44 (7.4%), 905.43 (4.1%), 906.44 (3.3%),
906.44 (1.6%), 907.44 (1.3%)

Figure 1- figure supplement 3: 3-3 crosslinks are not detectable in ΔLDT cells. (A)
Total ion chromatograms of WT, ΔLDT and ΔLDT_comp peptidoglycan. **(B)** Table of

muropeptide masses (Da) observed in (S3A). The molecular weight difference by one of the identified peptides is due to differential amidation. The descriptions include the peptide lengths in the crosslink (4= tetra-, 3= tri- peptide) and the following parenthesis specifies the number of amidation in the species according to mass. **(C)** Structure of a representative 3-3 crosslink with a m/z=904.4. **(D)** Extracted ion chromatograms from WT, ΔLDT and ΔLDT$_{comp}$ for a representative 3-3 crosslink with a m/z=904.4.

Figure 2- figure supplement 1: ΔLDT cell morphological characteristics. (A) Time-lapse microscopy montage of ΔLDT cells. The white stars mark new poles. The orange arrow points to the first new pole daughter cell of this series. The red arrow indicates the second resulting new pole daughter cell. In the last frame, white arrows point to all new pole daughter cells (besides the orange arrow and red arrow). **(B)** Time-lapse microscopy montage of ΔLDT$_{comp}$ cells expressing LdtE-mRFP.
All scale bars=5µm

A

B

Figure 2- figure supplement 2: Inheritance of old cell wall and occurrence of blebs
in new pole daughter cells. (A) WT Msm stained with Alexa Fluor™ 488 NHS ester,
washed and visualized over time. New material is unstained, old material is stained green.
Orange arrows indicate a new pole. Orange stars mark new pole daughter cells. All scale
bars=5µm **(B)** Maximum cell width of ΔLDT cell lineages over time. Width of new pole
daughters = blue circle; width of old pole daughters = orange circle. Division signs denote
a division event. At each division, there are two arrows from the dividing cell leading to
the resulting new and old pole daughter cell widths (blue and orange respectively).

A

Figure 3- figure supplement 1: L5 allele swapping to test essentiality of PonA1's ability to form 4-3 crosslinks (transpeptidation). (A) Schematic of L5 allele swapping experiment. Adapted from (Kieser, Boutte, et al., 2015b).

724
725

727
Figure 4- figure supplement 1: MSMEG_2433 (DacB2) functions as a D,D-carboxypeptidase and D,D-endopeptidase *in vitro*. (A) Coomassie-stained gel of purified His$_6$-DacB2. **(B)** Bocillin-FL and Penicillin G binding assay of purified DacB2.

Figure 4- figure supplement 1 (continued): MSMEG_2433 (DacB2) functions as a D,D-carboxypeptidase and D,D-endopeptidase *in vitro*. (C) Mass spectra of the reaction products of DacB2 digestion reactions.

736
737

A

	1	2	3	4	5	6	7	8	9	10	11	12
	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	
A	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	
B	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	2.50	
C	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	
D	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	0.63	
E	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	0.31	0.31	0.31	0.31	0.31	0.31	**0.31**	0.31	0.31	0.31	0.31	
F	10.00	5.00	2.50	1.25	0.63	0.31	**0.16**	0.08	0.04	0.02	0.01	
	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	
G	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	
	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	
H	10.00	5.00	2.50	1.25	0.63	0.31	0.16	0.08	0.04	0.02	0.01	

B

	1	2	3	4	5	6	7	8	9	10	11	12	
A	40.00	20.00	10.00	5.00	2.50	1.25	**0.63**	0.31	0.16	0.08	0.04	0.02	amoxicillin
B	40.00	20.00	10.00	5.00	2.50	1.25	**0.63**	0.31	0.16	0.08	0.04	0.02	
C	40.00	20.00	10.00	5.00	2.50	**1.25**	0.63	0.31	0.16	0.08	0.04	0.02	
													meropenem
D	40.00	20.00	10.00	5.00	2.50	**1.25**	0.63	0.31	0.16	0.08	0.04	0.02	

739
740 **Figure 5- figure supplement 1: Minimum inhibitory concentration (MIC) of**
741 **amoxicillin or meropenem alone or in combination against *Mycobacterium***

tuberculosis. **(A)** Resazurin MIC plate and dilution matrix of amoxicillin and meropenem in combination in a checkerboard MIC plate. The table below shows the concentration of each drug per well in μg/mL. The concentration of meropenem is in black text and the concentration of amoxicillin is in red text in each well. **(B)** Dilution matrix of amoxicillin or meropenem (alone). The concentrations of drugs are shown in the table. Pink indicates metabolically active cells, blue indicates not metabolically active. In both single drug and checkerboard MIC plates, 5μg/mL clavulanate was used.

CFU/mL · RLU · Negative Control RLU

Figure 5- figure supplement 2: Light production (RLU) correlated to colony forming units (CFU) in mycobacterial cells expressing *luxABCDE* in drug treatment. (A) *Mycobacterium smegmatis* colony forming units (CFU) and luminescence (RLU) during drug treatment.

Video captions

Figure 2-video1- Time-lapse of ΔLDT cells in iso-osmolar media.
This video corresponds to Figure 2A (top). This is a time-lapse microscopy video of ΔLDT Msm cells growing in 7H9 (iso-osmolar) media. Frames are 15 minutes apart, and the video is 5 frames/second.

Figure 2-video2- Time-lapse of ΔLDT cells in high-osmolar media.
This video corresponds to Figure 2A (bottom). This is a time-lapse microscopy video of ΔLDT Msm cells growing in 7H9 +150mM sorbitol (high-osmolar) media. Frames are 15 minutes apart, and the video is 5 frames/second.

Figure 4-video1- Time-lapse of ponA1-RFP.
This video corresponds to Figure 4A,B. This is a time-lapse microscopy video of Msm cells expressing ponA1-RFP growing in 7H9 media. Frames are 15 minutes apart, and the video is 5 frames/second.

Figure 4-video2- Time-lapse of ldtE-mRFP.
This video corresponds to Figure 4A,B. This is a time-lapse microscopy video of Msm cells expressing ldtE-mRFP growing in 7H9 media. Frames are 15 minutes apart, and the video is 5 frames/second.

Figure 4-video3- Time-lapse of dacB2-mRFP.
This video corresponds to Figure 4A,B. This is a time-lapse microscopy video of Msm cells expressing dacB2-mRFP growing in 7H9 media. Frames are 15 minutes apart, and the video is 5 frames/second.

Supplementary table 1: List of Primers

Primer description	Primer Sequence	5' or 3' features	Primer #
MSMEG_3528 (ldtA) upstream flanking region FOR	CGTTGTAAAACGACGGCCAG TGATGGCGGGCGTGATCTGG AATCTCT	pUC57 overlap	208
MSMEG_3528 (ldtA) upstream flanking region REV	ACCCAACTTAATCGCCTTGCA GCTCTTCCAGTGTAGGTTGT CGAAACG	Zeo cassette overlap	209
MSMEG_3528 (ldtA) downstream flanking region FOR	TAATCATGGTCATAGCTGTTT TCATCGTGCAGGCGTGACGT GCAG	Zeo cassette overlap	210
MSMEG_3528 (ldtA) downstream flanking region REV	CAGTCGACGGGCCCGGGAT CCGCGGTGGTGCCCTTGGTG ATGGTC	pUC57 overlap	211
MSMEG_0233 (ldtE) upstream flanking region FOR	CGTTGTAAAACGACGGCCAG TGGCTGTCCGCCCAGCCCCG GGCC	pUC57 overlap	220
MSMEG_0233 (ldtE) upstream flanking region REV	CCAACTTAATCGCCTTGCAG CGTGGTACCTCCAGAGCACA ACTG	Zeo cassette overlap	221
MSMEG_0233 (ldtE) downstream flanking region FOR	GTAATCATGGTCATAGCTGTT TCCGGACGTCATACGAAGAA CCCCC	Zeo cassette overlap	222
MSMEG_0233 (ldtE) downstream flanking region REV	GCAGTCGACGGGCCCGGGA TCTGGACCGACGCCGACCGC ACCG	pUC57 overlap	223
MSMEG_4745 (ldtB) upstream flanking region FOR	GGTCGAGATGCTCCTGGAAG AGGCCG		444

MSMEG_4745 (ldtB) upstream flanking region REV	GAGCTCCAATTCGCCCTACG TCGTTACCTGCCCCATCACG	hyg cassette overlap	445
MSMEG_4745 (ldtB) downstream flanking region FOR	GTACCTCGAGTCTAGAAGTA GCGCTATCGCACCGCGCGGT CCAG	hyg cassette overlap	446
MSMEG_4745 (ldtB) downstream flanking region REV	CGACCCGGCCCGTCACAAGG ACACCGAAC		447
MSMEG_0929 (ldtC) upstream flanking region FOR	CGTTGTAAAACGACGGCCAG TGAACTGGCGACGGCGCTGG GCGTGG	pUC57 overlap	216
MSMEG_0929 (ldtC) upstream flanking region REV	TGGAGCTCCAATTCGCCCTA GTGGATCTAGGGTACCGACA GCACGC	hyg cassette overlap	448
MSMEG_0929 (ldtC) downstream flanking region FOR	GGTACCTCGAGTCTAGAAGT AGTCCGGCGGCTAGGTCCG GCGGTTGAAG	hyg cassette overlap	449
MSMEG_0929 (ldtC) downstream flanking region REV	CAGTCGACGGGCCCGGGAT CCCAAGGGACTCGCGCCGGT CTCC	pUC57 overlap	219
MSMEG_0929 (ldtC) upstream flanking region FOR	GGCTCGTTCTTCACCAACC		507
MSMEG_0929 (ldtC) downstream flanking region REV	CTGCCCAAGCTCATCGAC		508
MSMEG_0674 (ldtG) upstream flanking region FOR	GTTGTAAAACGACGGCCAGT GCGGCGTCGACCTCCCGGC CGGGTC	pUC57 overlap	228

MSMEG_0674 (ldtG) upstream flanking region REV	GTGGAGCTCCAATTCGCCCT AGCGCATTGGCTTCCGATTT CCCTCG	hyg cassette overlap	454
MSMEG_0674 (ldtG) downstream flanking region FOR	CGGTACCTCGAGTCTAGAAG TACGCCGACGTGTATGCCCA CCCCCGCG	hyg cassette overlap	455
MSMEG_0674 (ldtG) downstream flanking region REV	GCAGTCGACGGGCCCGGGA TCGCCTGCGCCCGCGGGAG CGCCTGCC	pUC57 overlap	231
MSMEG_0674 (ldtG) upstream flanking region FOR	GCATCTGAGTTTCGGCAAG		513
MSMEG_0674 (ldtG) downstream flanking region REV	CAACTACCCCGCAGTTGAAT		514
MSMEG_1322 (ldtF) upstream flanking region FOR	GTTGTAAAACGACGGCCAGT GCGAGGTAAGGGTCTCGACG GTTTCT	pUC57 overlap	224
MSMEG_1322 (ldtF) upstream flanking region REV	GTGGAGCTCCAATTCGCCCT ATCCAATGTGCTTCGGCGAA AGCCAGTTTG	hyg cassette overlap	452
MSMEG_1322 (ldtF) downstream flanking region FOR	GTACCTCGAGTCTAGAAGTA GTTCCCCCGGCCCACATAT GTCTGGACG	hyg cassette overlap	453
MSMEG_1322 (ldtF) downstream flanking region REV	GCAGTCGACGGGCCCGGGA TCCACGACAACGCCAGCGCG AT	pUC57 overlap	227
MSMEG_1322 (ldtF) upstream flanking region FOR	GGTCGACGACGAACTGGT		511
MSMEG_1322 (ldtF) downstream	AACGGCACGTACATCAGGAC		512

flanking region REV			
MSMEG_0233 (ldtE) FOR with TetO overlap (vector)	CATGCTTAATTAAGAAGGAGA TATACAATGCCGAAATCGGC AAAACGCAG		323A
MSMEG_0233 (ldtE) REV (no stop codon) with ser-ser-gly linker	GATGACGTCCTCGGAGGAGG CCGAGCCGCCGAACATCTGC CAGTCGGATG		351
mRFP FOR with ser-ser-gly linker	CATCCGACTGGCAGATGTTC GGCGGCTCGGCCTCCTCCGA GGACGTCATC		352
mRFP REV with vector overlap	GTCCCCAATTAATTAGCTAAG TGATGGTGATGGTGATGACA GGCG		353
MSMEG_2433 (dacB2) FOR (first 27 amino acids truncated)	GGCCTGGTGCCGCGCGGCA GCCATCGCGCGGACGCCGA CATCCAG	with 5' overlaps to pet28b cut with NdeI	662
MSMEG_2433 (dacB2) REV	GCTGTCCACCAGTCATGCTA GCCATCAGAGCGCCCCGATG CTCG	with 3' overlaps to pet28b cut with NdeI	663

790
791

792 **Supplementary Table 2: List of Strains**
793

Strain	Description	Primers
KB134	mc2155Δ*ldtA*::loxP	208/209; 210/211
KB156	mc2155Δ*ldtA*::loxP + Δ*ldtE*:: zeoR	220/221; 222/223
KB200	mc2155Δ*ldtA*::loxP Δ*ldtE*:: zeoR + Δ*ldtB*:: hygR	444/445; 446/447
KB209	mc2155Δ*ldtA*::loxP Δ*ldtE*::loxP Δ*ldtB*::loxP + Δ*ldtC*:: hygR	216/448; 449/219 (create original hyg KO) but used 507/508; (amplify KO from strain within the flanks)
KB222	mc2155Δ*ldtA*::loxP Δ*ldtE*::loxP Δ*ldtB*::loxP Δ*ldtC*:: hygR Δ*ldtG*:: zeoR	228/454; 455/231 (create original hyg KO) but used 513/514; (amplify KO from strain within the flanks)
KB303 (ΔLDT)	mc2155Δ*ldtA*::loxP Δ*ldtE*::loxP Δ*ldtB*::loxP Δ*ldtC*:: loxP Δ*ldtG*:: loxP Δ*ldtF*:: hygR	224/452; 453/227 (create hyg KO) but used 511/512 (amplify KO from strain or gibson)
KB302	pTetO-ldtE(MSMEG_0233)-linker-mRFP in CT94 XH (XL1-Blue)	323A/351; 352/353
KB316 (ΔLDTcomp)	[mc2155ΔldtA::loxP ΔldtE::loxP ΔldtB::loxP ΔldtC:: loxP ΔldtG:: loxP ΔldtF:: hygR] + KB302	
KB428	*E.coli* BL21 + pet28b (dacB2)	662/663

794
795

Supplemental Table 3: FDAA FACs screen data used for Figure 1D

Supplemental Table 4: ΔLDT Tnseq data used for Figure 3A
Below are the column names with a brief description-
Orf   - ID of gene
Name - name of gene
Desc - annotation of gene
Sites - number of TA sites in gene
Mean Ctrl - mean insertion count averaged over TA sites and replicates for wildtype strain (mc^{2}155)
Mean Exp - mean insertion count averaged over TA sites and replicates for knockout strain (ΔLDT)
log2FC - log-fold-change, log2(meanExp/meanCtl)
Sum Ctrl - sum of insertion counts over TA sites and replicates for wildtype strain (mc^{2}155)
Sum Exp   - sum of insertion counts over TA sites and replicates for knockout strain (ΔLDT)
Delta Sum - difference of sums (sumExp-sumCtl)
p-value - probability of null hypothesis (i.e. no significant difference between strains) estimated from resampling distribution
Adj. p-value - p-values after applying Benjamini-Hochberg correction for multiple tests

References:

Aldridge, B. B., Fernandez-Suarez, M., Heller, D., Ambravaneswaran, V., Irimia, D., Toner, M., & Fortune, S. M. (2012). Asymmetry and aging of mycobacterial cells lead to variable growth and antibiotic susceptibility. *Science*, *335*(6064), 100–104. http://doi.org/10.1126/science.1216166

Andreu, N., Fletcher, T., Krishnan, N., Wiles, S., & Robertson, B. D. (2012). Rapid measurement of antituberculosis drug activity in vitro and in macrophages using bioluminescence. *The Journal of Antimicrobial Chemotherapy*, *67*(2), 404–414. http://doi.org/10.1093/jac/dkr472

Andreu, N., Zelmer, A., Fletcher, T., Elkington, P. T., Ward, T. H., Ripoll, J., et al. (2010). Optimisation of bioluminescent reporters for use with mycobacteria. *PLoS ONE*, *5*(5), e10777. http://doi.org/10.1371/journal.pone.0010777

Bansal, A., Kar, D., Murugan, R. A., Mallick, S., Dutta, M., Pandey, S. D., et al. (2015). A putative low-molecular-mass penicillin-binding protein (PBP) of Mycobacterium smegmatis exhibits prominent physiological characteristics of DD-carboxypeptidase and beta-lactamase. *Microbiology (Reading, England)*, *161*(Pt 5), 1081–1091. http://doi.org/10.1099/mic.0.000074

Botella, H., Yang, G., Ouerfelli, O., Ehrt, S., Nathan, C. F., & Vaubourgeix, J. (2017). Distinct Spatiotemporal Dynamics of Peptidoglycan Synthesis between Mycobacterium smegmatis and Mycobacterium tuberculosis. *mBio*, *8*(5), e01183–17. http://doi.org/10.1128/mBio.01183-17

Boutte, C. C., Baer, C. E., Papavinasasundaram, K., Liu, W., Chase, M. R., Meniche, X., et al. (2016). A cytoplasmic peptidoglycan amidase homologue controls mycobacterial cell wall synthesis. *eLife*, *5*, a021113. http://doi.org/10.7554/eLife.14590

Böth, D., Schneider, G., & Schnell, R. (2011). Peptidoglycan remodeling in Mycobacterium tuberculosis: comparison of structures and catalytic activities of RipA and RipB. *Journal of Molecular Biology*, *413*(1), 247–260. http://doi.org/10.1016/j.jmb.2011.08.014

Brown, P. J. B., de Pedro, M. A., Kysela, D. T., Van der Henst, C., Kim, J., De Bolle, X., et al. (2012). Polar growth in the Alphaproteobacterial order Rhizobiales. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(5), 1697–1701. http://doi.org/10.1073/pnas.1114476109

Cameron, T. A., Zupan, J. R., & Zambryski, P. C. (2015). The essential features and modes of bacterial polar growth. *Trends in Microbiology*, *23*(6), 347–353. http://doi.org/10.1016/j.tim.2015.01.003

Cava, F., de Pedro, M. A., Lam, H., Davis, B. M., & Waldor, M. K. (2011). Distinct pathways for modification of the bacterial cell wall by non-canonical D-amino acids. *The EMBO Journal*, *30*(16), 3442–3453. http://doi.org/10.1038/emboj.2011.246

DeJesus, M. A., Ambadipudi, C., Baker, R., Sassetti, C., & Ioerger, T. R. (2015). TRANSIT - A Software Tool for Himar1 TnSeq Analysis. *PLoS Computational Biology*, *11*(10), e1004401–17. http://doi.org/10.1371/journal.pcbi.1004401

Dhar, N., Bousbaine, D., Wakamoto, Y., McKinney, J. D., & Santi, I. (2013). Single-cell dynamics of the chromosome replication and cell division cycles in mycobacteria.

Nature Communications, *4*, 1–10. http://doi.org/10.1038/ncomms3470

Diacon, A. H., van der Merwe, L., Barnard, M., Groote-Bidlingmaier, von, F., Lange, C., García-Basteiro, A. L., et al. (2016). β-Lactams against Tuberculosis--New Trick for an Old Dog? *The New England Journal of Medicine*, *375*(4), 393–394. http://doi.org/10.1056/NEJMc1513236

Domínguez-Escobar, J., Chastanet, A., Crevenna, A. H., Fromion, V., Wedlich-Söldner, R., & Carballido-López, R. (2011). Processive movement of MreB-associated cell wall biosynthetic complexes in bacteria. *Science*, *333*(6039), 225–228. http://doi.org/10.1126/science.1203466

Eskandarian, H. A., Odermatt, P. D., Ven, J. X. Y., Hannebelle, M. T. M., Nievergelt, A. P., Dhar, N., et al. (2017). Division site selection linked to inherited cell surface wave troughs in mycobacteria. *Nature Microbiology*, *2*(9), 17094. http://doi.org/10.1038/nmicrobiol.2017.94

Garner, E. C., Bernard, R., Wang, W., Zhuang, X., Rudner, D. Z., & Mitchison, T. (2011). Coupled, Circumferential Motions of the Cell Wall Synthesis Machinery and MreB Filaments in B. subtilis. *Science*, *333*(6039), 222–225. http://doi.org/10.1126/science.1203285

Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., & Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods*, *6*(5), 343–345. http://doi.org/10.1038/nmeth.1318

Global Tuberculosis Report 2017. (2017). Global Tuberculosis Report 2017, 1–262.

Gonzalo, X., & Drobniewski, F. (2013). Is there a place for β-lactams in the treatment of multidrug-resistant/extensively drug-resistant tuberculosis? Synergy between meropenem and amoxicillin/clavulanate. *The Journal of Antimicrobial Chemotherapy*, *68*(2), 366–369. http://doi.org/10.1093/jac/dks395

González-Leiza, S. M., de Pedro, M. A., & Ayala, J. A. (2011). AmpH, a bifunctional DD-endopeptidase and DD-carboxypeptidase of Escherichia coli. *Journal of Bacteriology*, *193*(24), 6887–6894. http://doi.org/10.1128/JB.05764-11

Gupta, R., Lavollay, M., Mainardi, J.-L., Arthur, M., Bishai, W. R., & Lamichhane, G. (2010). The Mycobacterium tuberculosis protein LdtMt2 is a nonclassical transpeptidase required for virulence and resistance to amoxicillin. *Nature Medicine*, *16*(4), 466–469. http://doi.org/10.1038/nm.2120

Hett, E. C., & Rubin, E. J. (2008). Bacterial growth and cell division: a mycobacterial perspective. *Microbiology and Molecular Biology Reviews : MMBR*, *72*(1), 126–56– table of contents. http://doi.org/10.1128/MMBR.00028-07

Hugonnet, J.-E., Tremblay, L. W., Boshoff, H. I., Barry, C. E., & Blanchard, J. S. (2009). Meropenem-clavulanate is effective against extensively drug-resistant Mycobacterium tuberculosis. *Science*, *323*(5918), 1215–1218. http://doi.org/10.1126/science.1167498

Hussain, S., Wivagg, C. N., Szwedziak, P., Wong, F., Schaefer, K., Izore, T., et al. (2018). MreB filaments align along greatest principal membrane curvature to orient cell wall synthesis. *eLife*, *7*, e32471. http://doi.org/10.7554/eLife.32471

Kieser, K. J., & Rubin, E. J. (2014). How sisters grow apart: mycobacterial growth and division. *Nature Publishing Group*, *12*(8), 550–562. http://doi.org/10.1038/nrmicro3299

Kieser, K. J., Baranowski, C., Chao, M. C., Long, J. E., Sassetti, C. M., Waldor, M. K., et

al. (2015a). Peptidoglycan synthesis in Mycobacterium tuberculosis is organized into networks with varying drug susceptibility. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(42), 13087–13092. http://doi.org/10.1073/pnas.1514135112

Kieser, K. J., Boutte, C. C., Kester, J. C., Baer, C. E., Barczak, A. K., Meniche, X., et al. (2015b). Phosphorylation of the Peptidoglycan Synthase PonA1 Governs the Rate of Polar Elongation in Mycobacteria. *PLoS Pathogens*, *11*(6), e1005010. http://doi.org/10.1371/journal.ppat.1005010

Koschwanez, J. H., Carlson, R. H., & Meldrum, D. R. (2009). Thin PDMS films using long spin times or tert-butyl alcohol as a solvent. *PLoS ONE*, *4*(2), e4572. http://doi.org/10.1371/journal.pone.0004572

Kumar, P., Arora, K., Lloyd, J. R., Lee, I. Y., Nair, V., Fischer, E., et al. (2012). Meropenem inhibits D,D-carboxypeptidase activity in Mycobacterium tuberculosis. *Molecular Microbiology*, *86*(2), 367–381. http://doi.org/10.1111/j.1365-2958.2012.08199.x

Kuru, E., Hughes, H. V., Brown, P. J., Hall, E., Tekkam, S., Cava, F., et al. (2012). In Situ Probing of Newly Synthesized Peptidoglycan in Live Bacteria with Fluorescent D-Amino Acids. *Angewandte Chemie International Edition*, *51*(50), 12519–12523. http://doi.org/10.1002/anie.201206749

Kuru, E., Lambert, C., Rittichier, J., Till, R., Ducret, A., Derouaux, A., et al. (2017). Fluorescent D-amino-acids reveal bi-cellular cell wall modifications important for Bdellovibrio bacteriovorus predation. *Nature Microbiology*, *2*(12), 1648–1657. http://doi.org/10.1038/s41564-017-0029-y

Kuru, E., Tekkam, S., Hall, E., Brun, Y. V., & Van Nieuwenhze, M. S. (2014). Synthesis of fluorescent D-amino acids and their use for probing peptidoglycan synthesis and bacterial growth in situ. *Nature Protocols*, *10*(1), 33–52. http://doi.org/10.1038/nprot.2014.197

Lavollay, M., Arthur, M., Fourgeaud, M., Dubost, L., Marie, A., Veziris, N., et al. (2008). The Peptidoglycan of Stationary-Phase Mycobacterium tuberculosis Predominantly Contains Cross-Links Generated by L,D-Transpeptidation. *Journal of Bacteriology*, *190*(12), 4360–4366. http://doi.org/10.1128/JB.00239-08

Lebar, M. D., May, J. M., Meeske, A. J., Leiman, S. A., Lupoli, T. J., Tsukamoto, H., et al. (2014). Reconstitution of peptidoglycan cross-linking leads to improved fluorescent probes of cell wall synthesis. *Journal of the American Chemical Society*, *136*(31), 10874–10877. http://doi.org/10.1021/ja505668f

Long, J. E., DeJesus, M., Ward, D., Baker, R. E., Ioerger, T., & Sassetti, C. M. (2015). Identifying Essential Genes in <Emphasis Type="Italic">Mycobacterium tuberculosis</Emphasis> by Global Phenotypic Profiling. In *Gene Essentiality* (Vol. 1279, pp. 79–95). New York, NY: Humana Press, New York, NY. http://doi.org/10.1007/978-1-4939-2398-4_6

Mainardi, J.-L., Hugonnet, J.-E., Rusconi, F., Fourgeaud, M., Dubost, L., Moumi, A. N., et al. (2007). Unexpected inhibition of peptidoglycan LD-transpeptidase from Enterococcus faecium by the beta-lactam imipenem. *Journal of Biological Chemistry*, *282*(42), 30414–30422. http://doi.org/10.1074/jbc.M704286200

Murphy, K. C., Papavinasasundaram, K., & Sassetti, C. M. (2015). Mycobacterial recombineering. *Methods in Molecular Biology (Clifton, N.J.)*, *1285*, 177–199.

http://doi.org/10.1007/978-1-4939-2450-9_10

Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I., & Jacobs-Wagner, C. (2016). Oufti: an integrated software package for high-accuracy, high-throughput quantitative microscopy analysis. *Molecular Microbiology*, *99*(4), 767–777. http://doi.org/10.1111/mmi.13264

Papp-Wallace, K. M., Endimiani, A., Taracila, M. A., & Bonomo, R. A. (2011). Carbapenems: past, present, and future. *Antimicrobial Agents and Chemotherapy*, *55*(11), 4943–4960. http://doi.org/10.1128/AAC.00296-11

Pisabarro, A. G., de Pedro, M. A., & Vázquez, D. (1985). Structural modifications in the peptidoglycan of Escherichia coli associated with changes in the state of growth of the culture. *Journal of Bacteriology*, *161*(1), 238–242.

Qiao, Y., Lebar, M. D., Schirner, K., Schaefer, K., Tsukamoto, H., Kahne, D., & Walker, S. (2014). Detection of lipid-linked peptidoglycan precursors by exploiting an unexpected transpeptidase reaction. *Journal of the American Chemical Society*, *136*(42), 14678–14681. http://doi.org/10.1021/ja508147s

Qiao, Y., Srisuknimit, V., Rubino, F., Schaefer, K., Ruiz, N., Walker, S., & Kahne, D. (2017). Lipid II overproduction allows direct assay of transpeptidase inhibition by β-lactams. *Nature Chemical Biology*, *13*(7), 793–798. http://doi.org/10.1038/nchembio.2388

Rebets, Y., Lupoli, T., Qiao, Y., Schirner, K., Villet, R., Hooper, D., et al. (2014). Moenomycin resistance mutations in Staphylococcus aureus reduce peptidoglycan chain length and cause aberrant cell division. *ACS Chemical Biology*, *9*(2), 459–467. http://doi.org/10.1021/cb4006744

Rego, E. H., Audette, R. E., & Rubin, E. J. (2017). Deletion of a mycobacterial divisome factor collapses single-cell phenotypic heterogeneity. *Nature*, *546*(7656), 153–157. http://doi.org/10.1038/nature22361

Sanders, A. N., Wright, L. F., & Pavelka, M. S. (2014). Genetic characterization of mycobacterial L,D-transpeptidases. *Microbiology (Reading, England)*, *160*(Pt 8), 1795–1806. http://doi.org/10.1099/mic.0.078980-0

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. http://doi.org/10.1038/nmeth.2019

Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution Methods. (2016). Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution Methods. In *Clinical Microbiology Procedures Handbook, Fourth Edition* (pp. 5.16.1–5.16.23). American Society of Microbiology. http://doi.org/10.1128/9781555818814.ch5.16

Thangawng, A. L., Ruoff, R. S., Swartz, M. A., & Glucksberg, M. R. (2007). An ultra-thin PDMS membrane as a bio/micro-nano interface: fabrication and characterization. *Biomedical Microdevices*, *9*(4), 587–595. http://doi.org/10.1007/s10544-007-9070-6

Ursell, T. S., Nguyen, J., Monds, R. D., Colavin, A., Billings, G., Ouzounov, N., et al. (2014). Rod-like bacterial shape is maintained by feedback between cell curvature and cytoskeletal localization. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(11), E1025–34. http://doi.org/10.1073/pnas.1317174111

Vollmer, W., Blanot, D., & de Pedro, M. A. (2008a). Peptidoglycan structure and

architecture. *FEMS Microbiology Reviews*, *32*(2), 149–167.
http://doi.org/10.1111/j.1574-6976.2007.00094.x
Vollmer, W., Joris, B., Charlier, P., & Foster, S. (2008b). Bacterial peptidoglycan
(murein) hydrolases. *FEMS Microbiology Reviews*, *32*(2), 259–286.
http://doi.org/10.1111/j.1574-6976.2007.00099.x
Welsh, M. A., Taguchi, A., Schaefer, K., Van Tyne, D., Lebreton, F., Gilmore, M. S., et
al. (2017). Identification of a Functionally Unique Family of Penicillin-Binding
Proteins. *Journal of the American Chemical Society*, *139*(49), 17727–17730.
http://doi.org/10.1021/jacs.7b10170
Zimmermann, L., Stephens, A., Nam, S.-Z., Rau, D., Kübler, J., Lozajic, M., et al.
(2017). A Completely Reimplemented MPI Bioinformatics Toolkit with a New
HHpred Server at its Core. *Journal of Molecular Biology*.
http://doi.org/10.1016/j.jmb.2017.12.007
1020