An ultra-dense haploid genetic map for evaluating the highly

fragmented genome assembly of Norway spruce (*Picea abies*)

Carolina Bernhardsson[1,2,3,*], Amaryllis Vidalis[1,4], Xi Wang[1,3], Douglas G.

Scofield[1,5,6], Bastian Schiffthaler[7], John Basion[2], Nathaniel R. Street[7], M. Rosario

García-Gil[2], Pär K. Ingvarsson[1,3,*]

[1] Department of Ecology and Environmental Science, Umeå University, Umeå,

Sweden

[2] Umeå Plant Science Centre, Department of Forest Genetics and Plant

Physiology, Swedish University of Agricultural Science, Umeå, Sweden

[3] Department of Plant Biology, Uppsala BioCenter, Swedish University of

Agricultural Science, Uppsala, Sweden.

[4] Department of Population Genetics, Center of Life and Food Sciences

Weihenstephan, Technische Universität München, 85354 Freising,, Germany

[5] Uppsala Multidisciplinary Center for Advanced Computational Science,

Uppsala University, Uppsala, Sweden

[6] Department of Ecology and Genetics: Evolutionary Biology, Uppsala

University, Uppsala, Sweden

[7] Umeå Plant Science Centre, Department of Plant Physiology, Umeå

University, Umeå, Sweden

*Authors for correspondence: carolina.bernhardsson@umu.se,

par.ingvarsson@slu.se

Abstract

Norway spruce (*Picea abies* (L.) Karst.) is a conifer species of substanital

economic and ecological importance. In common with most conifers, the *P. abies*

genome is very large (~20 Gbp) and contains a high fraction of repetitive DNA. The

current *P. abies* genome assembly (v1.0) covers approximately 60% of the total

genome size but is highly fragmented, consisting of >10 million scaffolds. The

genome annotation contains 66,632 gene models that are at least partially validated

(www.congenie.org), however, the fragmented nature of the assembly means that

there is currently little information available on how these genes are physically

distributed over the 12 *P. abies* chromosomes. By creating an ultra-dense genetic

linkage map, we anchored and ordered scaffolds into linkage groups, which

complements the fine-scale information available in assembly contigs. Our ultra-

dense haploid consensus genetic map consists of 21,056 markers derived from 14,336

scaffolds that contain 17,079 gene models (25.6% of the validated gene models) that

we have anchored to the 12 linkage groups. We used data from three independent

component maps, as well as comparisons with previously published *Picea* maps to

evaluate the accuracy and marker ordering of the linkage groups. We demonstrate that

approximately 3.8% of the anchored scaffolds and 1.6% of the gene models covered

by the consensus map have likely assembly errors as they contain genetic markers that

map to different regions within or between linkage groups. We further evaluate the

utility of the genetic map for the conifer research community by using an independent

data set of unrelated individuals to assess genome-wide variation in genetic diversity

using the genomic regions anchored to linkage groups. The results show that our map

is sufficiently dense to enable detailed evolutionary analyses across the *P. abies*

genome.

Introduction

For over a century genetic linkage maps have been used to order genetic markers

and link phenotypic traits to genomic regions and chromosomes by calculating

recombination events in crosses (Sturtevant 1913a; Sturtevant 1913b). With the

advent of Next Generation Sequencing technologies (NGS), large numbers of markers

can now be scored at a relatively low cost and within a reasonable time, which has

enabled generation of high-density genetic maps consisting of thousands of markers

that, in combination with a sufficiently large mapping population, can achieve

unprecedented mapping resolution even in non-model systems and in species with

large genomes. Genetic maps represent a complementary approach to the local, fine-

scale genomic information that is available in scaffolds from a genome assembly,

with a genetic map providing information on genome organization over larger scales

(up to whole-chromosome level) (Fierst 2015). By grouping markers into linkage

groups and subsequently ordering them within each linkage group, it is possible to

anchor underlying scaffolds containing those markers to putative chromosomes with

high precision (Fierst 2015). If several genetic markers, derived from a single

genomic scaffold, are placed on the map, information on their relative placement in

the genetic map can be used to orient the scaffold and to evaluate scaffolding

decisions made in the genome assembly and hence to locate and resolve possible

assembly errors (Drost et al. 2009; Bartholomé et al. 2015). For instance, when two

markers originating from a single scaffold map to different linkage groups or to

different regions within a linkage group, the contigs comprising the scaffold are

candidates for having been wrongly joined during the assembly process. On the other

hand, if markers from the same scaffold map close to each other this increases the

likelihood that the scaffolding decisions were correct.

Norway Spruce (*Picea abies*) is one of the most important conifer species in

Europe, both from an ecological and economic perspective. The natural distribution

range of *P. abies* extends from the west coast of Norway to the Ural mountains and

across the Alps, Carpathians and the Balkans in central Europe. *P. abies* composes,

together with *Pinus sylvestris*, the majority of the continuous boreal forests of the

Northern hemisphere where it is considered a keystone species (Farjon 1990). *P.*

abies has a genome size of ~20 Gbp that is characterized by a very high fraction of

repetitive sequences. Like most conifers, *P. abies* has a karyotype consisting of $2n=24$

and with chromosomes that are all uniformly sized (Sax and Sax 1933). Due to the

large and complex genome of conifers, this important group of plants was, until

recently, lacking species with available reference genomes. In 2013 the first draft

assembly of the *P. abies* genome was published (Nystedt et al. 2013). Despite

extensive whole-genome shotgun sequencing derived from both haploid and diploid

tissues, the *P. abies* genome assembly is still highly fragmented due to the complex

nature and size of the genome. The current *P. abies* genome assembly (v1.0) consists

of 10.3 million scaffolds >500 bp and contains 70,736 annotated gene models of

which 66,632 are at least partially validated by supporting evidence (ESTs or UniProt

proteins) (Nystedt et al. 2013; De La Torre et al. 2014). Although the current genome

assembly only covers about two thirds of the total genome size (12 Gbp out of the 20

Gbp *P. abies* genome), it is expected to contain the majority of expressed genes.

In this paper, we used sequence capture to identify segregating SNP markers

in megagametophytes from three open-pollinated mother trees. These markers were

used to create an ultra-dense haploid genetic map consisting of 21,056 probe-markers

derived from 14,336 gene-bearing scaffolds in the *P. abies* genome assembly. Our

aim with creating the genetic map was to 1) anchor, and where possible, order

scaffolds to assign as many gene models as possible to linkage groups, and 2) to

evaluate the accuracy of the *P. abies* genome assembly v1.0 on the basis of anchored

scaffolds. To evaluate the accuracy of the map itself, we compared scaffold order to

previously published genetic maps for *P. abies* and the closely related *Picea glauca*.

Finally, we evaluated utility of the genetic map for population genomic studies by

performing genome-wide analyses of genetic diversity for the genomic regions

anchored in the map using a sample of c. 500 unrelated *P. abies* trees.

Material and Methods

DNA extraction and sequence capture

In the autumn of 2013, seeds were collected for linkage map construction from five of

30 putative ramets of Z4006, the genotype used to generate the reference genome for

Picea abies (Nystedt et al. 2013). Megagametophytes were dissected from 2,000

seeds by removing the diploid seed coat surrounding the haploid megagametophyte

tissue. DNA extraction from megagametophytes was performed using a Qiagen Plant

Mini Kit. Each extracted sample was measured for DNA quality using a Qubit® ds

DNA Broad Range (BR) Assay Kit, and all samples with a total amount of DNA

>354 ng were kept. The remaining 1,997 samples were sent to RAPiD Genomics©

(Gainesville, Florida, USA) in September 2014 for sequence capture using 31,277

capture probes that had been specifically designed to target 19,268 partially-validated

gene models from the *P. abies* genome assembly. Where possible, probes were

designed to flank regions of known contig joins in the v1.0 genome assembly (for

further detail of the probe design, see Vidalis et al. 2018).

The capture data was sequenced by RAPiD Genomics© on an Illumina HiSeq

2000 using 1x75 bp sequencing and was delivered in October 2015. The raw reads

were mapped against the complete *P. abies* reference genome v.1.0 using BWA-

MEM v.0.7.12 and default settings (Li and Durbin 2009). Following read mapping,

the genome was subset to only contain the probe-bearing scaffolds (a total of 18,461

scaffolds) using Samtools v.1.2 (Li and Durbin 2009; Li et al. 2009). Duplicates were

marked and local realignment around insertion/deletions (indels) was performed using

Picard   (http://broadinstitute.github.io/picard/)   and   GATK

(https://software.broadinstitute.org/gatk/) (McKenna et al. 2010; DePristo et al. 2011).

Genotyping was performed using GATK Haplotypecaller (version 3.4-46, (DePristo

et al. 2011; Van der Auwera et al. 2013) with a diploid ploidy setting and gVCF

output format. We used a diploid ploidy setting to increase the likelihood of detecting

possible sample contamination from diploid tissue for the haploid megagametophyte

samples. CombineGVCFs was then run on batches of ~200 gVCFs to hierarchically

merge them into a single gVCF and a final SNP call was performed using

GenotypeGVCFs jointly on the 10 combined gVCF files, using default read mapping

filters, a standard minimum confidence threshold for emitting (stand-emit-conf) of 10,

and a standard minimum confidence threshold for calling (stand_call_conf) of 20. See

Vidalis   et   al.   (2018)   and   the   script   "per_sample_gvcf.sh"   (available   at

https://github.com/parkingvarsson/HaploidSpruceMap) for a full description of the

pipeline used for calling variants.

SNP filtration and megagametophyte relationships

After SNP filtering, we performed a principle component analysis (PCA) to evaluate

the relationship among samples (see Supplementary file for details on the PCA

analysis and subsequent filtering steps). Based on the PCA and a hierarchical

clustering approach, we divided samples into three clusters representing putative

maternal families (Supplementary, Figure S1-3) that were then analyzed

independently. In the end we obtained 9,073 probe-markers from 7,101 scaffolds for

Cluster 1 (314 samples), 11,648 probe-markers from 8,738 scaffolds for Cluster 2

(270 samples) and 19,006 probe-markers from 13,301 scaffolds for Cluster 3 (842

samples) with a total of 21,056 probe-markers from 14,336 scaffolds across all three

clusters (Table 1). In total, these scaffolds cover 0.34 Gbp of the *P. abies* genome and

contain 17,079 partially validated gene models.

Table 1: Overview of the three component maps and the total number of probe-markers

available in the consensus map. Cluster: Name of each putative maternal family that

was identified in the principal component analysis. Samples: Number of

megagametophytes in each cluster. Markers: Number of probe-markers in each

component map with number of unique segregating bins within brackets (one marker

for each bin was used to anchor the bin markers to the genetic map). Scaffolds: Number

of scaffolds represented in each component map.

Cluster	Samples	Markers	Scaffolds
Cluster 1	314	9,073 (3,924)	7,101
Cluster 2	270	11,647 (5,311)	8,738
Cluster 3	842	19,006 (11,479)	13,301
Total	1,426	21,056	14,336

Component and consensus maps

We created genetic linkage maps using the R-package BatchMap (Schiffthaler et al.

2017), a parallel implementation of the R-package Onemap (Margarido, Souza, and

Garcia 2007). All probe-markers were recoded using the D1.11 cross-type (Wu et al.

2002), tested for segregation distortion (p < 0.05 after Bonferroni correction)

(Supplementary, Figure S4) and grouped into marker bins. The probe-marker with

lowest amount of missing data in each bin was then used to represent the bin when

constructing the genetic map. Bin markers were grouped into LGs using LOD = 8 and

a maximum recombination fraction = 0.35. LGs were then ordered using the

RECORD algorithm (Van Os et al. 2005) with 16 times counting, parallelized over 16

cores, reordered in a 10 marker sliding window with 1 marker incremental steps using

the command 'ripple' and finally mapped using the Kosambi mapping function and

the 'map batches' approach (Schiffthaler et al. 2017) over four parallel cores. Finally,

heat maps with pairwise recombination fraction (lower triangular) and phase LOD

score (upper triangular) for the ordered markers were created to evaluate the ordering

accuracy of independent linkage groups (Supplementary, Figure S5 and S6A-L). We

observed 183 probe-marker bins showing signs of segregation distortion. These bins

were, however, randomly distributed over the linkage groups and did not appear to

affect marker ordering and map distance and were therefore retained in subsequent

analyses.

To evaluate correspondence between LGs in maps derived from the three PCA

clusters, the number of unique scaffolds shared between cluster LGs were counted

(Supplementary, Figure S5). We then created a consensus map for each linkage group

from the three independent component maps using the R-package LPmerge

(Endelman and Plomion 2014) with component maps ranked according to marker

numbers (Cluster 3, Cluster 2, Cluster 1), a maximum interval setting ranging from

one to 10 and map weights proportional to the size of the mapping population (Cluster

3= 0.5, Cluster 2 and Cluster 1 = 0.25). From all possible consensus maps generated

by LP merge, for each linkage group we selected the map with the lowest mean root

mean square error (RMSE) to serve as the consensus map (Endelman and Plomion

2014). Order correlations between individual component maps and the consensus

maps (Table 2 and Supplementary, Figure S7A-L) as well as between the three

component maps (Supplementary, Figure S8A-L) were estimated using Kendall's τ.

For visual representation of the consensus map we created a Circos plot using the R-

package omicCircos (Hu et al. 2014), available from Bioconductor

(https://bioconductor.org/biocLite.R).

To evaluate the inflation of map distances due to possible genotyping errors,

we performed 100 rounds of random subsampling of 100 probe-marker bins per LG

and component map. The following marker ordering and genetic distance calculation

were performed with 10 rounds of RECORD and the Kosambi mapping function.

Accuracy of the reference Picea abies genome assembly

To evaluate the accuracy of scaffolds from the v1.0 *P. abies* reference genome

containing at least two probe-markers (here after called multi-marker scaffolds) we

determine whether probe-markers from the same genomic scaffold mapped to the

same region of an LG, on different regions within a single LG or on different LGs. In

the consensus map, we considered markers to be positioned in the same region on an

LG if all probe-markers from a scaffold mapped within a 5 cM interval of each other.

If any marker from the scaffold was positioned further apart, the scaffold was tagged

as containing a putative assembly error. The same considerations were made for

scaffolds with probe-markers positioned on different LGs.

Comparative analyses of Picea linkage maps

To evaluate the consistency of our genetic map with earlier maps from *P. abies* we

compared our haploid consensus map to the *P. abies* linkage map from Lind et al.

(2014). The Lind et al. map was created using genetic markers generated using an

Illumina 3072 SNP Golden Gate Assay. We performed using `tblastn` sequence

homology searches against the *P. abies* v1.0 genome assembly for the SNP array

sequences of the makers mapped in the Lind et al. map and extracted reciprocal best

hits with >95% identity, which were then assigned to the corresponding scaffold in

the *P. abies* genome. We performed similar analyses to compare the synteny between

our consensus map and the *P. glauca* composite map from Pavy et al. (2017). Again,

we used `tblastn` sequence homology search comparisons of array sequences from

the *P. glauca* SNP array (Pavy et al. 2013) with scaffolds from the *P. abies* v1.0

genome assembly to assign corresponding map positions between *P. abies* and *P.*

glauca. In order to evaluate correspondence between LGs from the different genetic

maps, we assessed the number of shared scaffolds between our consensus map, the

Lind et al. and Pavy et al. maps. Consistency of scaffold ordering was then evaluated

using visual comparisons (Figure 4 and 5) and by calculating correlations of marker

orders using Kendall's τ.

Population genetic analysis of the consensus genetic map

In order to independently evaluate the utility of the consensus map for downstream

research, we used a subset of the data from Baison et al. (2018) to estimate patterns of

nucleotide diversity across the Norway spruce genome. The data from Baison et al.

originally contained 517 individuals sequenced with 40,018 probes designed for

diploid spruce samples (Vidalis et al 2018). We extracted data for all probes that we

had anchored in our genetic map from the VCF file containing the data from Baison et

al.. We further hard-filtered the resulting VCF file by only considering bi-allelic SNPs

within the extended probe regions (120 bp probes ±100 bp) with a QD >5, MQ >50

and a overall DP between 3000 and 16000. Samples containing >25% missing data

were removed from further analysis. We used the data to calculate nucleotide

diversity (π), the number of segregating sites and Tajima's D (Tajima 1989). We used

the R package vcfR (Knaus and Grünwald 2017) to read the VCF-file into R and then

used in-house developed scripts to perform all calculations (available at

https://github.com/parkingvarsson/HaploidSpruceMap). We assigned probes to LGs

and map positions by assigning them the coordinates of the physically closest (in bp)

probe. We also calculated pairwise linkage disequilibrium (LD) between markers

within probes using vcftools (Danecek et al. 2011) and imported the results into R

where they were used to calculate Z_{nS} scores (Kelly 1997) per probe using an in-house

developed script (available at https://github.com/parkingvarsson/HaploidSpruceMap).

Finally, we ran sliding window analyses along the linkage groups for the different

summary statistics using 10 cM windows that were moved in 1 cM incremental steps.

Results

We generated a *P. abies* consensus linkage map from three haploid component maps containing a total of 21,056 unique probe-markers from 14,336 scaffolds in the *P. abies* genome assembly v1.0. The consensus map anchored 0.34 Gbp of the *P. abies* 1.0 assembly, corresponding to 1.7% of the complete *P. abies* genome or 2.8% of the genome assembly. However, these scaffolds anchor 25.6% of all validated gene models with these anchored scaffolds containing 31.7%, 20.6% and 25.8% of the High-, Medium- and Low confidence gene models from Nystedt et al (2013), respectively. The consensus map had a total length of 3,556 centiMorgan (cM), distributed over 12 linkage groups (LGs), corresponding to the haploid chromosome number (Sax and Sax 1933), and with an average distance of 0.17 cM between probe-markers (Table 2, Figure 1A).

Correlations of probe-marker order between the three component maps and the consensus map ranged from 0.96 to 0.998, while the correlations between marker orders between individual component maps ranged from 0.943 to 0.993 (Supplementary, Figure S7 and S8). 183 probe-marker bins showed evidence of segregation distortion in Cluster 3, but these were randomly distributed over all linkage groups and we did not observe regions showing clusters of markers with segregation distortion or with conflicting marker orders between clusters (Supplementary, Figure S8). LG XI, which displayed the largest discrepancy in marker order between component maps, has a region at the distal end of the LG, covering 252 probe-markers, where the resolution was too low to identify the correct marker order and where the entire region was positioned at 36.115 cM (Supplementary, Figure S7K and S8K), explaining the lower correlations in marker order between individual maps for this LG.

We used a random subsampling approach to evaluate potential inflation of map

distances due to possible genotyping errors. From these analyses, total map size for

Cluster 1 ranged between 2,166.8 and 2,450.0 cM with an average size of 2,294.2 cM

and a standard deviation (SD) of 3.6- 5.8 cM per LG. Cluster 2 ranged between

2,304.2 and 2,663.6 cM with an average of 2,478.3 cM and a SD of 4.4 – 9.1 cM per

LG, while Cluster 3 ranged between 1,855.4 and 2,093.2 cM with an average of

1,971.0 cM and a SD of 2.7 – 7.3 cM per LG. The estimated inflation was therefore

predicted to be 0.15 – 0.31 cM per probe-marker bin across the three component maps

(Table 3). This inflation per probe-marker bin roughly corresponded to the map

resolution of the clusters (Cluster 1- 0.32 cM: Cluster 2 - 0.37 cM: Cluster 3 – 0.12

cM) and yielded an error estimate of ~1 genotype error per marker-bin or 11-17

genotype errors per sample.

Table 2: Marker density and size of each component genetic map created from the three clusters as well as for the consensus map. LG: Linkage

group. Cluster 1-3: Component maps for cluster 1-3 with number of probe-markers (marker-bins) assigned, map size (in cM) and maximum gap in map

(in cM) for each of the LGs. Consensus: Number of markers and map size of the LGs in the consensus map.

LG	Cluster 1			Cluster 2			Cluster 3			Consensus	
	Markers	Length (cM)	Max gap (cM)	Markers	Length (cM)	Max gap (cM)	Markers	Length (cM)	Max gap (cM)	Markers	Length (cM)
I	975 (421)	385.5	8.0	1,159 (553)	439.9	21.1	1,967 (1,185)	414.1	8.8	2,172	414.1
II	701 (305)	249.2	9.6	863 (366)	289.0	9.4	1,456 (864)	289.8	10.9	1,608	250.3
III	859 (394)	324.0	4.6	1,069 (479)	381.1	7.1	1,738 (1,075)	346.4	5.2	1,940	342.5
IV	771 (323)	298.7	14.5	970 (452)	350.9	8.6	1,531 (916)	303.0	27.0	1,704	303.0
V	761 (311)	273.2	8.9	1,116 (499)	395.6	9.5	1,649 (1,032)	342.6	15.1	1,865	275.0
VI	648 (292)	241.0	8.4	915 (399)	270.7	4.6	1,456 (894)	269.5	8.4	1,622	240.2
VII	682 (331)	314.0	8.4	923 (443)	380.8	13.4	1,625 (1,013)	321.9	7.9	1,769	321.0
VIII	775 (339)	307.0	5.6	943 (454)	367.26	9.8	1,465 (904)	315.6	6.6	1,609	305.9
IX	792 (332)	283.3	5.4	786 (364)	295.6	5.9	1,589 (911)	285.1	7.4	1,738	285.0
X	648 (289)	231.6	7.0	960 (454)	342.7	6.9	1,564 (917)	272.7	7.1	1,709	273.1
XI	677 (253)	200.6	3.7	1,025 (411)	269.2	4.0	1,440 (818)	233.6	3.0	1,608	233.4
XII	784 (334)	281.6	9.3	919 (437)	360.7	11.1	1,526 (950)	312.3	14.3	1,712	312.3
Total	9,073 (3,924)	3,389.4	14.5	11,648 (5,311)	4,143.4	21.1	19,006 (11,479)	3,706.7	27.0	21,056	3,555.8

304   **Table 3:** Estimated genetic length of each Linkage Group (LG) in the three component maps. LG: linkage group in the consensus map; Observed
305   genetic length (cM): The genetic length of the LG calculated from all probe-marker bins (same as in table 2); Mean estimated genetic length (cM): the
306   average length of the LG when using 100 random probe-marker bins in 100 map calculations; SD (cM): Standard deviation of the estimated length;
307   Inflation/Marker bin: The difference between observed genetic length and the estimated length divided by the number of probe-marker bins in the
308   linkage group.

LG	Cluster 1				Cluster 2				Cluster 3			
	Observed genetic length (cM)	Mean estimated genetic length (cM)	SD (cM)	Inflation / Marker bin (cM)	Observed genetic length (cM)	Mean estimated genetic length (cM)	SD (cM)	Inflation / Marker bin (cM)	Observed genetic length (cM)	Mean estimated genetic length (cM)	SD (cM)	Inflation / Marker bin (cM)
I	385.5	245.5	5.2	0.33	439.9	252.3	6.1	0.34	414.2	204.8	7.3	0.18
II	249.2	168.8	4.7	0.26	289.0	192.9	4.4	0.26	289.8	166.4	2.8	0.14
III	324.0	195.8	5.8	0.33	381.1	218.6	5.8	0.34	346.4	168.5	3.9	0.17
IV	298.7	204.7	5.0	0.29	350.9	215.6	5.7	0.30	303.0	167.0	3.5	0.15
V	273.2	195.7	4.6	0.25	395.6	218.4	9.0	0.36	342.6	180.0	5.1	0.16
VI	241.0	161.8	4.7	0.27	270.7	170.0	4.7	0.25	269.5	142.2	2.9	0.14
VII	314.0	223.6	5.3	0.27	380.8	248.7	6.3	0.30	321.9	175.9	3.7	0.14
VIII	307.0	203.6	4.9	0.31	367.26	226.7	5.7	0.31	315.6	179.2	4.3	0.15
IX	283.3	194.0	4.9	0.27	295.6	185.3	6.8	0.30	285.1	157.2	3.0	0.14
X	231.6	164.4	3.6	0.23	342.7	193.6	4.5	0.33	272.7	141.5	2.7	0.14
XI	200.6	141.8	4.7	0.23	269.2	147.0	4.8	0.30	233.6	119.6	3.0	0.14
XII	281.6	194.4	4.9	0.26	360.7	209.6	9.1	0.35	312.3	168.7	3.1	0.15
Total	3,389.4	2,294.2	-	0.28	4,143.4	2,478.3	-	0.31	3,706.7	1,971.0	-	0.15

Figure 1: Circos plot of the consensus map. A) Marker distribution over the 12 linkage groups (LG I-LG XII). Each black vertical line represents a marker (21,056 in total) in the map and is displayed according to the marker positions in cM. Track B-C visualizes multi marker scaffolds, where each line is a pairwise position comparison of probe-markers from the same scaffold. B) Position comparisons of probe-markers from the same scaffold that are located on the same LG. Light grey lines indicate probe-markers that are located < 5cM from each other, dark grey lines indicate probe-markers located 5-10 cM apart and red lines indicate probe-markers >10 cM apart. C) Position comparisons of probe-markers from the same scaffold that are mapping to different LGs. Orange lines indicated probe-markers from the same scaffold split over 2 LGs, while dark blue lines indicated probe-markers split over 3 LGs.

Evaluation of the Picea abies genome assembly v1.0

4,859 scaffolds (33.9%) contained more than one unique probe-marker combined

over all three component maps. 185 of these multi-marker scaffolds contained

markers that were located in more than one LG (*inter-split scaffolds*) or over different

parts of the same LG (*intra-split scaffolds*). 26 scaffolds (0.18% of mapped scaffolds

and 0.54% of multi-marker scaffolds) contained markers that were positioned on the

same LG but at distances exceeding 5 cM in the consensus map. When exploring the

individual component maps, it was apparent that for two of these scaffolds

(MA_281725 on LG X and MA_10431182 on LG I) the probe-markers in the

consensus map all came from different component maps. The consensus map thus

contain a gap that we can not verify using any of the individual component maps

(Figure 1 and Supplementary, Figure S9). Three other scaffolds (MA_9458 on LG IX,

MA_10431315 on LG II and MA_10432328 on LG III) all have multiple probe-

markers present in at least one component map and were these component maps do

not support the split we observe in the consensus map (Supplementary, Figure S9). It

thus appears that these splits are artifacts arising from the construction of the

consensus map.

There were 164 scaffolds (1.14% of mapped scaffolds and 3.38% of multi-

marker scaffolds) containing markers that were mapped to two or three different LGs

(Figure 2 and Supplementary, Figure S10). All LGs contained inter-split scaffolds,

while 10 LGs (LGII and LGXI are the exceptions) contained intra-split scaffolds

supported by the component maps (Figure 1B-C and Supplementary, Figure S9).

Figure 2: Fraction of scaffolds that are being represented by 1-11 unique markers in the consensus map. Insert: Fraction of scaffolds that have multiple probe-markers (2-11) that are distributed over 1-3 linkage groups (inter-split scaffolds). Red dot indicate the fraction of scaffolds with multiple probe-markers which are positioned > 5cM apart on the same linkage group (intra-split scaffolds).

The scaffolds covered by the consensus map ranged in length from 0.22 to 208.1 Kbp with a median of 17.1 Kbp, while multi-marker scaffolds ranged from 0.39 to 161.5 Kbp (median of 21 Kbp). The 185 scaffolds that are split within or across LGs ranged in size from 2.5 to 121.6 Kbp, with a median length of 36.9 Kbp. Split scaffolds were significantly longer than multi-marker scaffolds in general (t = -7.7, df

= 193.4, p-value = 7.0e-13; Figure 3), suggesting that longer scaffolds are more likely

to contain assembly errors compared to shorter scaffolds. Split scaffolds mostly

contained high- and medium confidence gene models (Table 4). A visual inspection

of the split scaffolds revealed that for 75 and 10 of the inter-split and intra-split

scaffolds, respectively, the predicted position of the split(s) occurred between

different gene models on the same scaffold. Of greater concern, for 88 of the inter-

split scaffolds and 11 of the intra-split scaffolds the predicted position of the split was

located within a single gene model (Supplementary, Figure S9 and S10). In addition,

21 inter-split scaffolds showed an even more complicated picture, where an interior

region of the gene model (most often containing an intron > 5kb) mapped to another

chromosome whereas the 5' and 3' regions of the gene model mapped to the same

chromosome location (Supplementary, Figure S10). However, 84% (184 out of a total

of 219 splits) appear to occur between contig joins (where a sequence of N's appear in

the assembly) of the scaffold. Of the 17,079 gene models that were anchored to the

consensus genetic map, 330 were positioned on inter- or intra-split scaffolds (5.4% of

gene models that were positioned on multi-marker scaffolds) and 100 showed a split

within gene models (1.6% of gene models from multi-marker scaffolds) (Table 4).

Figure 3: Box plot of scaffold lengths for all multi-marker scaffolds (dark gray

box) and for scaffolds showing a split within or across LGs (light gray box). The split

scaffolds are significantly longer than the multi-marker scaffolds in general (t = -7.70,

df = 193.39, p-value = 7.00e-13).

Table 4: Overview of annotated gene models anchored to the genetic map. Gene

models: Annotated protein coding gene models with High-, Medium- and Low

confidence level (Nystedt et al. 2013). Mapped scaffolds: Number of gene models

positioned on scaffolds that are anchored to the genetic map (Percentage of total

number of gene models for each confidence level). Multi-marker scaffolds: Number of

gene models positioned on scaffolds with multiple markers in the genetic map

(Percentage of gene models on mapped scaffolds). Inter-split scaffolds: Number of

gene models positioned on the 164 scaffolds that are split between LGs in the genetic

map (Percentage of gene models on mapped scaffolds / Percentage of gene models on

multi-marker scaffolds). Intra-split scaffolds: Number of gene models positioned on the
22 scaffolds that are split between different regions of the same LG (Percentage of
gene models on mapped scaffolds / Percentage of gene models on multi-marker
scaffolds). Split within gene models: Number of gene models that have an internal split
(Percentage of gene models on mapped scaffolds / Percentage of gene models on multi-
marker scaffolds).

Gene models	Mapped scaffolds	Multi-marker scaffolds	Inter-split scaffolds	Intra-split scaffolds	Split within gene models
High confidence	8,379 (31.7%)	3,122 (37.3%)	145 (1.7% / 4.6%)	15 (0.18% / 0.48%)	58 (0.69% / 1.9%)
Medium confidence	6,624 (20.6%)	2,215 (33.4%)	114 (1.7% / 5.1%)	16 (0.23% / 0.68%)	29 (0.44% / 1.3%)
Low confidence	2,076 (25.8%)	762 (36.7%)	35 (1.7% / 4.6%)	5 (0.29% / 0.79%)	13 (0.63% / 1.7%)
Total	17,079 (25.6%)	6,099 (35.7%)	294 (1.7% / 4.8%)	36 (0.21% / 0.59%)	100 (0.59% / 1.6%)

Comparative analyses to other Picea linkage maps

In order to assess the accuracy and repeatability of the *P. abies* genetic maps we
compared our consensus map to the *P. abies* map presented in Lind et al. (2014). 353
comparisons between 298 markers from Lind et al. and 288 scaffolds contained in our
consensus map were identified at a > 95 % identity threshold. Of these markers,
96.7% grouped to the same LG in the two maps while the remaining 3.3% (11 out of
353) were distributed across several LGs (Figure 4). Correlations of marker order
between the two *P. abies* maps ranged from 0.53 to 0.99 across the 12 LGs. The

comparison between the haploid consensus map for LG I and LG 7 from Lind et.al,

which had the lowest correlation of marker order, showed inconsistencies of marker

order where a contiguous subset of markers were arranged in the opposite order from

the rest of the markers for that LG. The remaining LGs showed high synteny, with

consistent marker ordering between the two genetic maps.

Figure 4: Marker order comparison between Linkage Groups (LGs) from the
haploid consensus map presented here and the *Picea abies* map from Lind et al. (2014).
Consensus LG I - LG XII are located on the x-axis from left to right. Lind et al. LG 1 -
LG 12 are located on the y-axis from top to bottom. Each dot represents a marker
comparison from the same scaffold, where black coloration represents the LG where
the majority of marker comparisons are mapped. Grey coloration represents markers

mapping to a different LG compared to the majority of markers. Turquoise coloration

represents markers located on split scaffolds, which are indicative of assembly errors.

Synteny between *P. abies* and *P. glauca* species was assessed by comparing LG

location and marker order between our *P. abies* consensus map and the composite

map of *P. glauca* from Pavy et al. (2017). 14,112 comparisons of 4,053 gene models

in the composite map in *P. glauca* (Pavy et al. 2017) and 4,310 scaffolds in the *P.*

abies consensus map were identified at a > 95% identity threshold. 92.7% (13,084 out

of 14,112 comparisons) of these were located on homologous LGs while the

remaining 7.3% (1,028 comparisons from 388 *P.abies* scaffolds) were distributed

across the 12 LGs (Figure 5). 8.2% of all comparisons from multi-probe scaffolds

were between non-homologous LGs while 44.3% of all comparisons from split

scaffolds were between non-homologous LGs.　 31.9% of all non-homologous LG

comparisons involved split scaffolds. The correlations of marker order between the

two maps were comparable to the correlations we observed between individual

component maps in *P. abies* (0.96-0.99), showing that synteny is highly conserved

between *P. abies* and *P glauca*.

Figure 5: Marker order comparison of Linkage Groups (LGs) between the *Picea abies* haploid consensus map presented here and the *Picea glauca* map from Pavy et al. (2017). Consensus LG I - LG XII are located on the x-axis from left to right. Pavy et al. LG 1 - LG 12 are located on the y-axis from top to bottom. Each dot represents a marker comparison from the same scaffold, where black coloration represents markers mapping to the same LG in the two species, grey coloration represents markers mapping to different LGs. Turquoise coloration represents markers located on split scaffolds, indicating an assembly error.

Population genetic analyses based on the consensus map

22,413 probes, covering 12,908 scaffolds, were used in the population genetic analyses based on the consensus genetic map. On a per-probe basis, we observed substantial variation in all neutrality statistics, with the number of segregating sites ranging from 0 - 77 (mean 15.9), nucleotide diversity (π) from 0 - 0.4 (0.005), Zns from 0 - 1 (mean 0.04) and Tajima's D from -2.4 – 3.5 (mean -0.85). To study large-scale trends and possible chromosomal differences we performed sliding window analyses across the LGs for the different summaries (Figure 6). One interesting large-scale feature we observed was that SNP densities were often highest at the distal or central regions of LGs, indicating the possible location of centromeres and telomeres, for which recombination rates are expected to be reduced (Gaut et al. 2007) and where we hence would expect higher densities of probes per cM (Figure 6a). The large-scale analyses also revealed several instances where entire chromosomal arms might be experiencing different evolutionary patterns (Figure 6b-c). Finally, we identified regions that appear to be evolving under the influence of natural selection. For instance, several regions showed higher than average levels of nucleotide diversity and positive Tajima's D (e.g. on LG IV, V and XII), suggesting that they might harbor genes under balancing selection. Similarly, regions with low nucleotide diversity, an excess of rare alleles and strong linkage disequilibrium (i.e. negative Tajima's D and high Z_{ns} scores, e.g. on LG III) could indicate regions harboring possible selective sweeps (Figure 6c-d).

Figure 6. Sliding window analysis of neutrality statistics. Analyses were performed using 10 cM windows with 1 cM incremental steps along the consensus map linkage groups and visualized using coloring alternates between adjacent LGs. A) Number of segregating sites. Dashed horizontal line indicates the overall average of 1017. B) Pairwise nucleotide diversity (π). Dashed horizontal line indicates the overall average of 0.005. C) Tajima's D. Dashed horizontal line indicates the overall average of -0.852. D) Linkage disequilibrium Zn scores. Dashed horizontal line indicates the overall average of 0.040.

Discussion

This is, to our knowledge, the densest genetic linkage map ever created for a conifer species and possible for any tree species. We successfully used this genetic map to

anchor 1.7% of the 20 Gbp *P. abies* genome, corresponding to 2.8% of the v1.0

genome assembly (Nystedt et al. 2013), to 12 LGs, constituting the haploid

chromosome number (Sax and Sax 1933). The *P. abies* genome has a very large

proportion of gene-poor heterochromatin, so while the fraction of the genome that we

successfully anchored to the assembly is relatively small, those anchored scaffolds

cover 24% of all gene-containing assembly scaffolds and 25% of all partially

validated gene models from Nystedt et al. (2013).

The individual LGs from the three component maps (36 LGs from three

independent maps) consisted of 648-1,967 probe-markers and 305-1,185 probe-

marker bins and, as such, it was not feasible to analyze the maps using an exhaustive

ordering algorithm (Mollinari et al. 2009). Instead, we used RECORD (Van Os et al.

2005) with 16 times counting, parallelized over 16 cores and with reordering of

markers within 10 marker windows, for each LG to determine the most likely marker

order. An heuristic approach, such as RECORD, will undoubtedly introduce some

errors in marker ordering (Mollinari et al. 2009), but analyses from simulated data

suggested that the average distance between estimated and true marker position is

small (< 5 markers) for data sets of similar size to ours (Schiffthaler et al. 2017).

However, reliable marker ordering requires robust data and the more genotyping

errors and missing data that are present, the harder it will be to determine the true

order. This in turn will impact the final size of the map, where both errors in marker

order and genotyping results in inflation in the size of the map (Cartwright et al.

2007).

By collecting our 2,000 megagametophytes from what we initially thought were

five different ramets of Z4006, we accidentally sampled material from at least three

unrelated families. This error stemmed from a mix-up of genotypes due to wrong

assignment of ramet ID to the different ramets in the seed orchard. Unfortunately, we

were not able to assess which megagametophytes were collected from the different

putative ramets since the seed bags were pooled prior to DNA extraction and the

sampling errors were not detected until after all sequencing was completed. We used

a PCA and hierarchical clustering approach to assign samples into three independent

clusters, representing three putative maternal families. We also used PCAs of the

putative individual families to verify that these clusters were consistent with offspring

derived from a single mother tree (Supplementary, Figure S3). However, we

nevertheless cannot completely rule out that a small fraction of samples have been

incorrectly assigned to the three families and this would lead to inflated map sizes by

introducing an excess of recombination events. Another potential confounding issue is

tissue contamination. *P. abies* megagametophytes are very small and are surrounded

by a diploid seed coat that needs to be removed prior to DNA extraction. If traces of

the diploid seed coat remain in the material used for DNA extractions, the haploid

samples will be contaminated with diploid material. To identify and eliminate this

possibility, we called sequence variants using a diploid model and any heterozygous

SNP calls were subsequently treated as missing data. Samples with a high proportion

of heterozygous (>10 %) or missing calls (>20%) were excluded from further

analyses to reduce the possibilities of genotyping error due to tissue contamination

influencing downstream analyses. We estimated map lengths from 100 rounds of

subsampling of 100 random probe-marker bins per component LG and used this to

demonstrate that individual maps showed size inflations of 0.15-0.31 cM per probe-

marker bin. This inflation is on the same order as the map resolutions for the different

clusters and, therefore, indicated an average of ~1 genotyping error per probe-marker

bin or 11-17 genotyping errors per sample.

Both sample- and tissue contaminations can influence the accuracy of the genetic

map, both with regards to marker order and map size. The smaller family sizes

resulting from dividing our original 2,000 samples into three independent families

yielded lower resolution of the three component maps. Fortuitously enough, however,

this also enabled us to incorporate more markers into the consensus map since

different markers were segregating in the different mother trees from which the three

families were derived. Furthermore, it also allowed us to evaluate marker ordering

across three independently derived maps. Although our consensus map was 70-90%

(60-120% for the individual component maps) larger than previously estimated *Picea*

maps (3,556 cM vs. 1,889-2,083 cM), it also contained 2-31 times more markers than

earlier maps (Pavy et al. 2012; Lind et al. 2014; Pavy et al. 2017). When comparing

marker order between our three independent component maps (Cluster 1-3), we found

overall high correlations of marker order (0.94-0.99, Supplementary, Figure S8),

which is similar to what has previously been observed between estimated and true

positions in maps derived from simulated data without genotyping errors but with

20% missing data (Mollinari et al 2009; Schiffthaler et al. 2017). Also, earlier *Picea*

maps were all based on diploid F_1 crosses with even the densest composite map

containing only 2,300-2,800 markers per framework map (Table 1 - Pavy et al. 2017),

compared to our haploid component maps that contained between 3,924 and 11,479

probe-marker bins each (Table 2).

The comparisons between our haploid consensus map and earlier maps in *Picea*

showed an overall high correlation of marker order, which is in line with previous

studies suggesting highly conserved synteny within *Picea* and in conifers in general

(de Miguel et al. 2015; Pavy et al. 2017). LG I from our haploid consensus map and

LG 7 from Lind et al. (2014) showed an inverted order for approximately half of the

markers compared (Figure 4). Whether this inversion is due to ordering errors in one

of the maps or represents true biological differences between the parents used for the

respective maps is, however, not currently known and further investigations are

needed to resolve this issue.

A small percentage of the marker comparisons in both the intra- and inter-

specific maps did not co-align to homologous LGs. Some of these errors likely arose

form the repetitive nature of the *P. abies* genome (and conifer genomes in general),

where regions with high sequence similarity can often be found interspersed

throughout the genome. If the true homologous region between different maps is

missing or has been collapsed in the genome assembly due to high sequence similarity,

pairwise sequence comparisons may end up assigning homology to regions that are

located on different chromosomes. However, it might also be hat these errors

represent scaffold assembly errors for scaffolds containing only a single probe-marker

or where one region of the scaffold is not captured by the probes, therefore negating

evaluation. Approximately 72% of all non-homologues LG comparisons between *P.*

abies and *P. glauca* were from multi-markers scaffolds (of which 45% were from

probe-markers on split scaffolds in the consensus map (turquoise points in Figure 5).

The remaining 28% were comparisons with scaffolds that were only represented by a

single probe in the consensus map.

Four percent of the scaffolds containing multiple makers showed a pattern where

different markers mapped to different regions, either within or between LGs in the

consensus map. This indicates possible errors in scaffolding during the assembly of

the v1.0 *P. abies* genome (Nystedt et al. 2013). If this estimate represents the overall

picture for the entire assembly, as many as 400,000 of the ~10 million total scaffolds,

and 2,400 of the ~60,000 gene-containing scaffolds, may suffer from assembly errors.

Most worryingly, 2% of the multi-marker scaffolds (100/4,859) contained splits that occurred within a single gene model. It is likely that many of these problematic scaffolds stem from incorrect scaffolding of exons from paralogous genes with a high sequence similarity. Since the *P. abies* genome contains a high proportion of repetitive content, that also includes a large number of pseudo-genes, this is perhaps not surprising. Additional work is needed to disentangle these issues and to resolve any assembly errors. False scaffold joins in a genome assembly are not a unique feature for *P. abies*, rather it appears to be a frequent problem in the assembly process. For instance, dense genetic maps in both *Eucalyptus* and *Crassostrea* have identified and resolved false scaffold joins, thereby improving the genome assemblies in these species (Bartholomé et al. 2015; Hedgecock et al. 2015). Our goal for the *P. abies* genetic map was not only to identify incorrect scaffolding decisions in the v1.0 genome assembly, but to also help improve future iterations of the genome. Long-read sequencing technologies (e.g. Pacific Bioscience or Oxford Nanopore) could be used to resolve these problematic scaffolds and help disentangle the reasons for their ambiguous localization in the genetic map. A future reference genome for *P. a*bies, based on long read technologies will also be able to utilize this genetic map in a much more efficient way since the resulting assembled scaffolds will be substantially longer and would hence enable anchoring a greater fraction of the genome to LGs, ultimately to the point that chromosome-scale assemblies may be achieved.

Our population genetic analyses based on the scaffolds anchored to the consensus map demonstrates the utility of having a dense, accurate genetic map and suggest that the map will facilitate further analyses of genome-wide patterns of variation and selection in *P. abies* in addition to facilitating comparative analyses among spruce species. Assigning even a small fraction of the genome to LGs enabled us to analyze

patterns of genetic diversity in approximately a quarter of all predicted genes. This

allowed for analyses of broad-scale patterns of variation across the genome and, as

the genome assembly is further improved and an even greater proportion of the

assembly if physically anchored to the genetic map, will allow for even more fine-

scaled analyses of how different evolutionary forces have interacted in shaping

patterns of genetic diversity across the *P. abies* genome.

Acknowledgements

This study was supported by Knut and Alice Wallenberg's foundation through

funding to the Norway spruce genome project. AV was partially supported by a grant

from the Stiftelsen Gunnar och Birgitta Nordins fond through the Kungl. Skogs- och

Lantbruksakademien (KSLA). NRS was supported by the Trees and Crops for the

Future (TC4F) project. All computations were performed on resources provided by

SciLifeLab and SNIC at the Uppsala Multidisciplinary Center for Advanced

Computational Science (UPPMAX) under project b2010042.

Author contribution

PKI and MRGG conceived the study. AV collected cones and extracted DNA. CB,

AV, DS and JB set up bioinformatics pipeline for analyzing sequence capture data.

AV and CB performed PCA and identified samples belonging to the three clusters.

CB, DS and BS created the genetic maps. CB and PKI performed intra- and

interspecific map comparisons. CB, XW and PKI performed population genetic

analysis. CB performed all remaining analyses and wrote the first draft of the

manuscript. NRS contributed to manuscript writing and development of the map

construction approach. All authors commented on the manuscript at various stages

during the writing.

Data availability

BatchMap input files for the three clusters, component maps and consensus map files

are available from zenodo.org at https://doi.org/10.5281/zenodo.1209841. All scripts

needed to recreate the analyses described in the paper are publically available at

https://github.com/parkingvarsson/HaploidSpruceMap. Raw sequence data for all

samples included in this study are available though the European Nucleotide Archive

under accession number PRJEB25757.

References

Baison, J., Vidalis, A., Zhou, L., Chen, Z-Q., Li, Z, Sillanpää, M.J., Bernahrdsson, C.,

Scofield, D.G., Forsberg, N., Olsson, L., Karlsson, B., Wu, H., Ingvarsson, P.K.,

Lundqvist, S-O., Niittylä, T., Garcia Gil, M.R. 2018. Association mapping

identified novel candidate loci affecting wood formation in Norway spruce.

bioRxiv https://doi.org/10.1101/292847

Bartholomé, Jérôme, Eric Mandrou, André Mabiala, Jerry Jenkins, Ibouniyamine

Nabihoudine, Christophe Klopp, Jeremy Schmutz, Christophe Plomion, and

Jean-Marc Gion. 2015. High-Resolution Genetic Maps of Eucalyptus Improve

Eucalyptus Grandis Genome Assembly. *New Phytologist* 206: 1283–96.

doi:10.1111/nph.13150.

Cartwright, Dustin A, Michela Troggio, Riccardo Velasco, and Alexander Gutin.

2007. Genetic Mapping in the Presence of Genotyping Errors. *Genetics* 176:

2521–27. doi:10.1534/genetics.106.063982.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.

Handsaker, et al. 2011. The Variant Call Format and VCFtools. *Bioinformatics*

27: 2156–58. doi:10.1093/bioinformatics/btr330.

De La Torre, Amanda R., Inanc Birol, Jean Bousquet, Pär K. Ingvarsson, Stefan

Jansson, Steven J.M. Jones, Christopher I. Keeling, et al. 2014. Insights into

Conifer Giga-Genomes. *Plant Physiology* 166: 1724 – 1732.

http://www.plantphysiol.org/content/166/4/1724.short.

de Miguel, Marina, Jérôme Bartholomé, François Ehrenmann, Florent Murat,

Yoshinari Moriguchi, Kentaro Uchiyama, Saneyoshi Ueno, et al. 2015. Evidence

of Intense Chromosomal Shuffling during Conifer Evolution. *Genome Biology*

and Evolution 7: 2799–2809. doi:10.1093/gbe/evv185.

DePristo, Mark A, Eric Banks, Ryan Poplin, Kiran V Garimella, Jared R Maguire,

Christopher Hartl, Anthony A Philippakis, et al. 2011. A Framework for

Variation Discovery and Genotyping Using next-Generation DNA Sequencing

Data. *Nat Genet* 43: 491–98. doi:10.1038/ng.806.

Drost, Derek R., Evandro Novaes, Carolina Boaventura-Novaes, Catherine I.

Benedict, Ryan S. Brown, Tongming Yin, Gerald A. Tuskan, and Matias Kirst.

2009. A Microarray-Based Genotyping and Genetic Mapping Approach for

Highly Heterozygous Outcrossing Species Enables Localization of a Large

Fraction of the Unassembled *Populus Trichocarpa* Genome Sequence. *The Plant*

Journal 58: 1054–67. doi:10.1111/j.1365-313X.2009.03828.x.

671 Endelman, Jeffrey B., and Christophe Plomion. 2014. LPmerge: An R Package for

672   Merging Genetic Maps by Linear Programming. *Bioinformatics* 30: 1623–24.

673   doi:10.1093/bioinformatics/btu091.

674 Farjon, A. 1990. Pinaceae. Drawings and Descriptions of the Genera Abies, Cedrus,

675   Pseudolarix, Keteleeria, Nothotsuga, Tsuga, Cathaya, Pseudotsuga, Larix and

676   Picea. *Pinaceae. Drawings and Descriptions of the Genera Abies, Cedrus,*

677   *Pseudolarix, Keteleeria, Nothotsuga, Tsuga, Cathaya, Pseudotsuga, Larix and*

678   *Picea.* Koeltz Scientific Books.

679   https://www.cabdirect.org/cabdirect/abstract/19920656698.

680 Fierst, Janna L. 2015. Using Linkage Maps to Correct and Scaffold de Novo Genome

681   Assemblies: Methods, Challenges, and Computational Tools. *Frontiers in*

682   *Genetics* 6: 220. doi:10.3389/fgene.2015.00220.

683 Gaut, Brandon S., Stephen I. Wright, Carène Rizzon, Jan Dvorak, and Lorinda K.

684   Anderson. 2007. Recombination: An Underappreciated Factor in the Evolution

685   of Plant Genomes. Nature Reviews Genetics 8: 77–84.

686 Hedgecock, Dennis, Grace Shin, Andrew Y Gracey, David Van Den Berg, and Manoj

687   P Samanta. 2015. Second-Generation Linkage Maps for the Pacific Oyster

688   Crassostrea Gigas Reveal Errors in Assembly of Genome Scaffolds. G3: Genes,

689   Genomes, Genetics: 5: 2007–19. doi:10.1534/g3.115.019570.

690 Hu, Ying, Chunhua Yan, Chih-Hao Hsu, Qing-Rong Chen, Kelvin Niu, George

691   Komatsoulis, and Daoud Meerzaman. 2014. OmicCircos: A Simple-to-Use R

692   Package for the Circular Visualization of Multidimensional Omics Data. *Cancer*

693   *Informatics* 13: 13. doi:10.4137/CIN.S13495.

694 Kelly, J. K. 1997. "A Test of Neutrality Based on Interlocus Associations." *Genetics*

146: 1197–1206.

Knaus, Brian J., and Niklaus J. Grünwald. 2017. vcfR : A Package to Manipulate and
    Visualize Variant Call Format Data in R. *Molecular Ecology Resources* 17: 44–
    53. doi:10.1111/1755-0998.12549.

Li, H., and R. Durbin. 2009. Fast and Accurate Short Read Alignment with Burrows-
    Wheeler Transform. *Bioinformatics* 25: 1754–60.
    doi:10.1093/bioinformatics/btp324.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.
    Abecasis, R. Durbin, and 1000 Genome Project Data Processing Subgroup. 2009.
    The Sequence Alignment/Map Format and SAMtools. *Bioinformatics* 25: 2078–
    79. doi:10.1093/bioinformatics/btp352.

Lind, Mårten, Thomas Källman, Jun Chen, Xiao-Fei Ma, Jean Bousquet, Michele
    Morgante, Giusi Zaina, et al. 2014. A Picea Abies Linkage Map Based on SNP
    Markers Identifies QTLs for Four Aspects of Resistance to Heterobasidion
    Parviporum Infection. *PloS One* 9: e101049. doi:10.1371/journal.pone.0101049.

Margarido, G R A, A P Souza, and A A F Garcia. 2007. OneMap: Software for
    Genetic Mapping in Outcrossing Species. *Hereditas* 144: 78–79.
    doi:10.1111/j.2007.0018-0661.02000.x.

McKenna, Aaron, Matthew Hanna, Eric Banks, Andrey Sivachenko, Kristian
    Cibulskis, Andrew Kernytsky, Kiran Garimella, et al. 2010. The Genome
    Analysis Toolkit: A MapReduce Framework for Analyzing next-Generation
    DNA Sequencing Data. *Genome Research* 20: 1297–1303.
    doi:10.1101/gr.107524.110.

Mollinari, M, G R A Margarido, R Vencovsky, and A A F Garcia. 2009. Evaluation

of Algorithms Used to Order Markers on Genetic Maps. *Heredity* 103: 494–502.

doi:10.1038/hdy.2009.96.

Nystedt, Björn, Nathaniel R. Street, Anna Wetterbom, Andrea Zuccolo, Yao-Cheng

Lin, Douglas G. Scofield, Francesco Vezzi, et al. 2013. The Norway Spruce

Genome Sequence and Conifer Genome Evolution. *Nature* 497: 579–84.

doi:10.1038/nature12211.

Pavy, Nathalie, Astrid Deschênes, Sylvie Blais, Patricia Lavigne, Jean Beaulieu,

Nathalie Isabel, John Mackay, and Jean Bousquet. 2013. The Landscape of

Nucleotide Polymorphism among 13,500 Genes of the Conifer Picea Glauca,

Relationships with Functions, and Comparison with Medicago Truncatula.

Genome Biology and Evolution 5: 1910–25. doi:10.1093/gbe/evt143.

Pavy, Nathalie, Manuel Lamothe, Betty Pelgas, France Gagnon, Inanç Birol, Joerg

Bohlmann, John Mackay, Nathalie Isabel, and Jean Bousquet. 2017. A High-

Resolution Reference Genetic Map Positioning 8.8 K Genes for the Conifer

White Spruce: Structural Genomics Implications and Correspondence with

Physical Distance. *The Plant Journal* 90: 189–203. doi:10.1111/tpj.13478.

Pavy, Nathalie, Betty Pelgas, Jérôme Laroche, Philippe Rigault, Nathalie Isabel, and

Jean Bousquet. 2012. A Spruce Gene Map Infers Ancient Plant Genome

Reshuffling and Subsequent Slow Evolution in the Gymnosperm Lineage

Leading to Extant Conifers. *BMC Biology* 10: 84. doi:10.1186/1741-7007-10-84.

R Core Team. 2013. R: A Language and Environment for Statistical Computing. *R*

Foundation for Statistical Computing, Vienna, Austria. http://www.r-project.org.

Sax, Karl, and Hally Jolivettte Sax. 1933. Chromosome Number and Morphology in

the Conifers. *Journal of the Arnold Arboretum* 14: 356-375.

Schiffthaler B, Bernhardsson C, Ingvarsson PK, Street NR (2017) BatchMap: A

parallel implementation of the OneMap R package for fast computation of

F_1 linkage maps in outcrossing species. PLoS ONE 12(12): e0189256.

https://doi.org/10.1371/journal.pone.0189256

Sturtevant, A. H. 1913a. The Linear Arrangement of Six Sex-Linked Factors in

Drosophila, as Shown by Their Mode of Association. *Journal of Experimental*

Zoology 14: 43–59. doi:10.1002/jez.1400140104.

Sturtevant, A. H. 1913b. A Third Group of Linked Genes in Drosophila Ampelophila.

Science 37: 990–92. doi:10.1126/science.37.965.990.

Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis by

DNA polymorphism. Genetics 123: 585–595.

Van der Auwera, Geraldine A., Mauricio O. Carneiro, Christopher Hartl, Ryan Poplin,

Guillermo del Angel, Ami Levy-Moonshine, Tadeusz Jordan, et al. 2013. From

FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit

Best Practices Pipeline. In *Current Protocols in Bioinformatics*, 11.10.1-

11.10.33. Hoboken, NJ, USA: John Wiley & Sons, Inc.

doi:10.1002/0471250953.bi1110s43.

Van Os, Hans, Piet Stam, Richard G F Visser, and Herman J Van Eck. 2005.

RECORD: A Novel Method for Ordering Loci on a Genetic Linkage Map.

Theoretical and Applied Genetics. 112: 30–40. doi:10.1007/s00122-005-0097-x.

Vidalis, A. Scofield, D.G., Neves, L-G., Bernhardsson, C., García-Gil, M.R.,

Ingvarsson, P.K. 2018. Design and evaluation of a large sequence-capture

probe set and associated SNPs for diploid and haploid samples of Norway

spruce (*Picea abies*) *BioRxiv* doi: https://doi.org/10.1101/291716

Wu, Rongling, Chang-Xing Ma, Ian Painter, and Zhao-Bang Zeng. 2002.

Simultaneous Maximum Likelihood Estimation of Linkage and Linkage Phases

in Outcrossing Species. *Theoretical Population Biology* 61: 349–63.

doi:10.1006/tpbi.2002.1577.