1

Human pathogenic bacterium *Shigella* infects *Arabidopsis* plants using type-III effectors

that suppress conserved MAP kinase signaling

Sung Hee Jo[1,2], Jiyoung Lee[1,3], Dong Wook Kim[4,5], Dae Hee Lee[2,6], Choong Min Ryu[2,7], and Jeong Mee Park[1,2]

[1]*Plant Systems Engineering Research Center, Korea Research Institute of Bioscience & Biotechnology (KRIBB), Daejeon, South Korea;* [2]*Department of Biosystems and Bioengineering, KRIBB School of Biotechnology, Korea University of Science and Technology (UST), Daejeon, South Korea;* [3]*Biological Resource Center, KRIBB, Jeongeup, South Korea;* [4]*Department of Pharmacy, College of Pharmacy, Hanyang University, Ansan, South Korea;* [5]*Institute of Pharmacological Research, Hanyang University, Ansan, South Korea;* [6]*Synthetic Biology and Bioengineering Research Center, KRIBB, Daejeon, South Korea;* [7]*Infectious Disease Research Center, KRIBB, Daejeon, South Korea.*

Author for correspondence: Jeong Mee Park

Tel: +82-42-860-4346

E-mail: jmpark@kribb.re.kr

Abstract

Although human enteropathogenic bacteria can survive both on the surface of and inside

plants, little is known about the molecular interactions between bacteria and plants. Here, we

examined the interaction between the human pathogenic bacterium *Shigella* and the plant

Arabidopsis. Four *Shigella* strains actively proliferated in, and showed virulence against,

Arabidopsis. Using mutants lacking type-III secretion systems (T3SS), we showed that T3SS

that regulate the pathogenesis of shigellosis in humans also play a central role in pathogenesis

in *Arabidopsis* by inhibiting mitogen-activated protein kinase (MAPK)-mediated plant

immune responses. Using *Shigella* mutants deficient for a single T3S effector, we showed

that the virulence activity of T3S effectors (OspF or OspG) was needed for pathogenicity of

Shigella in *Arabidopsis*. Production of OspF in plant cells suppressed a constitutively active

form of tobacco mitogen-activated protein kinase kinase 2 (MEK2^{DD})-triggered cell death by

inhibiting phosphorylation of MAPK, indicating that suppression of MAPK signaling

activation by OspF is conserved in both human and plants. These findings demonstrate for

the first time that determinants of *Shigella*-mediated virulence and pathogenic symptoms are

expressed in both plant and mammalian models.

Keywords: *Arabidopsis thaliana*, Effectors, Enteropathogenic bacteria, MAPK signaling,

Shigella, Type-III secretion system

Introduction

Shigella species are Gram-negative bacteria from the *Enterobacteriaceae* family. They are non-motile, rod-shaped, facultative intracellular and invasive pathogens, very closely related to *Escherichia coli*. Based on the carbohydrate composition of the O-antigen, i.e., the polysaccharide component of the lipopolysaccharide molecule that is the major bacterial surface antigen, *Shigella* is classified into four serogroups. These have been given species designations, namely, *S. dysenteriae* 1 (serogroup A), *S. flexneri* (serogroup B), *S. boydii* (serogroup C), and *S. sonnei* (serogroup D) (Lindberg *et al.*, 1991; Schroeder & Hilbi, 2008). *Shigella* spp. are important epidemic pathogens and a serious public health concern in developed and developing countries. Although about 164,300 deaths in all age groups, and 54,900 deaths in children younger than 5 years, were reported globally in 2015, the actual number of infections might be higher because mild symptoms are not reported (Mortality & Causes of Death, 2016). This means that the microorganism may employ a variety of survival strategies not only for infection of animals but also for survival in a restricted host such as a plant. In recent years, the morbidity or mortality of the infection caused by the emergence of *Shigella* that produces β-lactamase, an enzyme that underpins the resistance to the third-generation antibiotics cephalosporins, has been increasing (Zhang *et al.*, 2011).

Shigella is a human-adapted pathogen that infects the host via multiple transmission routes. Human hosts are primarily infected by the ingestion of water or food contaminated with bacteria from feces (the fecal-oral route), including ingestion of undercooked contaminated meats and improperly washed contaminated vegetables (Franz *et al.*, 2008; Ferens & Hovde, 2011). Fresh fruits and vegetables, such as lettuce, tomatoes and green peppers, are responsible for the widespread transmission of food-borne infections by *Salmonella* or *Shigella* (Guchi & Ashenafi, 2010; Semenov *et al.*, 2010; Gu *et al.*, 2013b). These observations suggest that human pathogenic bacteria use plants as alternative hosts as a stable environmental niche. Several human pathogenic bacteria, including *Salmonella enterica* serovar Typhimurium, *Escherichia coli* O157:H7, and *Pseudomonas aeruginosa*, are known to use plants as alternative hosts (Plotnikova *et al.*, 2000; Semenov *et al.*, 2010). These bacteria can attach to the plant surface, and actively invade and proliferate in plant tissues (Cevallos-Cevallos *et al.*, 2012; Martínez-Vaz *et al.*, 2014). In particular, several enteropathogenic bacteria, including *E. coli* O157:H7 and *Salmonella*, spread within plants through vascular tissues after infection via contaminated water (Solomon *et al.*, 2002; Cevallos-Cevallos *et al.*, 2012; Gu *et al.*, 2013a). Although *Shigella* contamination has also

been reported in plants, it is not yet known whether the bacterium actively invades and/or

proliferates inside the plant.

Unlike animals, plants have no adaptive immune system; instead, each cell possesses

an innate immune system. The innate immune system in plants and animals recognizes and

suppresses pathogens and has common features that are preserved throughout evolution. Plant

pattern recognition receptors recognize conserved microbial or pathogen- associated

molecular patterns; the pattern-triggered immunity (PTI) is activated via the mitogen-

activated protein kinase (MAPK) cascades (Jones & Dangl, 2006). To suppress PTI, bacteria

inject effector proteins into plant cells using type-III secretion systems (T3SS). To counteract

this PTI evasion response, the plant nucleotide binding-leucine rich repeat proteins recognize

the pathogen effectors; effector-triggered immunity is then activated to accompany the

hypersensitive response (Jones & Dangl, 2006). For human or animal intestinal bacteria to

infect plants, the PTI must first be disabled. *S. enterica* serovar Typhimurium, similar to its

activity in the mammalian host, uses T3SS to suppress plant immune responses (Schikora *et*

al., 2011; Schikora *et al.*, 2012). In particular, SpvC, one of the T3S effector proteins of *S.*

enterica, targets the MAPK signaling system in both plant and animal hosts to suppress the

host PTI (Neumann *et al.*, 2014).

Here, we examined the ability of four *Shigella* strains [*S. sonnei* (*S. s*) (Holt *et al.*,

2012), *S. boydii* (*S. b*), *S. flexneri* 2a (*S. f* 2a) (Wei *et al.*, 2003), and *S. flexneri* 5a (*S. f* 5a)

(Onodera *et al.*, 2012)] to proliferate in *Arabidopsis* plants. We found that the bacteria

actively invaded and proliferated in plant root and leaf tissues. Proliferation of mutants

lacking the T3S effectors, i.e., noninvasive human strains, was reduced both *in vitro* and *in*

planta. Reverse genetics and molecular biology experiments demonstrated that the *Shigella*

T3S effectors OspF and OspG were delivered to plant cells and functioned as pathogenic

proteins. These observations indicate that *Arabidopsis* may be useful as a model host for

studying the pathogenesis of *Shigella*.

Materials and Methods

Plant materials and growth

Arabidopsis thaliana accession Columbia (Col-0) was used for *Shigella* infection.

Briefly, *Arabidopsis* seeds were surface-sterilized for 2 min in 70% (v/v) ethanol, incubated

in 50% household bleach for 10 min, washed extensively with sterile deionized water, and

planted on 1/2 Murashige and Skoog (MS) medium (Duchefa Biochemie, Haarlem,

Netherlands) supplemented with 1% sucrose and solidified with 0.6% (w/v) agar (Murashige

& Skoog, 1962). *Nicotiana benthamiana* plants were germinated and grown at $22 \pm 3°C$

under a 16 h light/8 h dark cycle in plastic pots containing steam-sterilized mixed soil (2:1:1,

v/v/v, soil/vermiculite/perlite) (Moon *et al.*, 2016). To measure the plant immune response in

terms of MAPK activity, 1 μM flg22 peptide (#FLG22-P-1; Alpha Diagnostics, Inc., San

Antonio, TX, USA) was used as a positive control in the experiments (Bethke *et al.*, 2009).

Bacterial strains, growth conditions, and plasmids

The bacterial strains and plasmids used in the study are described in Supporting

Table S1. *Shigella and Pseudomonas* strains harboring plasmid pDSK-GFPuv were generated

by electroporation as previously described (Wang *et al.*, 2007; Hong *et al.*, 2016).

Shigella spp. were grown at 37°C in Luria-Bertani (LB) medium or tryptic soy agar

containing 0.003% (w/v) Congo red dye (Sigma-Aldrich, St. Louis, MO, USA) (Runyen-

Janecky & Payne, 2002). *Pseudomonas syringae* strains were grown at 28°C with shaking at

200 rpm in King's B liquid medium (Sigma-Aldrich) containing appropriate antibiotics (King

et al., 1954). *Escherichia coli* DH5α was grown in LB medium at 37°C with shaking

(Kennedy, 1971) and used as a non-pathogenic control. *Agrobacterium tumefaciens* GV2260

was grown at 28°C in LB broth with shaking at 200 rpm (Shamloul *et al.*, 2014).

The coding region of *ospF* or *ospG* was PCR-amplified using *attB*-containing PCR primers

(Supporting Table S2), and the PCR fragments were cloned into the pDONR™207 vector

using the pENTR Directional TOPO Cloning kit (Invitrogen, Carlsbad, CA, USA). The

products were then transferred to the pBAV178 (for AvrRpt2 fusion) or pBAV179 (for HA

fusion) vectors by LR recombination (Gateway® LR Clonase™ II, Invitrogen). pBAV178,

pBAV179, and pME6012 (empty vector control) were kindly provided by Dr. Jean T.

Greenberg (University of Chicago) (Vinatzer *et al.*, 2005). The products were also transferred

to pK7FWG2 (obtained from Ghent University, Belgium) by LR recombination to produce

the green fluorescent protein (GFP)-fused *Shigella* effectors OspF:GFP and OspG:GFP

(Karimi *et al.*, 2002).

Bacterial inoculation assay *in planta*

Leaves (4 weeks old) from *Arabidopsis* plants grown in soil were infected with each

strain of bacteria by syringe infiltration (5×10^7 or 5×10^5 cfu/ml) or spray-inoculation ($5 \times$

10^7 cfu/ml) (Katagiri *et al.*, 2002). After infection, plants were covered with a clear lid to

maintain humidity and transferred to a growth room (22 ± 3°C, 16 h light/8 h dark). The

bacterial populations were measured via serial dilution assays (Katagiri *et al.*, 2002). Colony

forming units (cfu) were normalized to cfu/mg using the total weight of inoculated leaves.

Bacterial populations were evaluated on three plants per treatment (as biological replicates)

and in three independent experiments conducted successively.

Arabidopsis seedlings (2 weeks old) grown in 1/2 MS medium were used for flood

inoculation (Ishiga *et al.*, 2011). Briefly, 10 *Arabidopsis* seedlings in one container were

incubated for 3 min with 35 ml of each bacterial strain suspension (5×10^5 cfu/ml) containing

0.02% Silwet L-77 (Lehle Seeds, Round Rock, TX, USA) or buffer. After the bacterial

suspensions were removed by decantation, plates containing inoculated plants were incubated

in a growth room (23 ± 2°C, 16 h light/8 h dark). Bacterial cell counts from inoculated plants

were monitored as described previously (Ishiga *et al.*, 2011). Three inoculated seedlings in

one container were harvested by cutting the hypocotyls, and total fresh weight was measured.

The cfu were normalized to cfu/mg using sample weight. The cfu of seedlings in three

separate containers (as biological replicates) were measured. In addition, the bacterial

population was evaluated in more than three independent experiments conducted successively

under the same conditions.

To assess root pathogenicity, 10-d-old *Arabidopsis* seedlings grown vertically in 1/2

MS medium were infected by dropping 2.0 μl of bacterial suspension (5×10^7 cfu/ml) onto

the root tips. Disease symptoms were observed under white light (WL) at 7 or 14 dpi, and

bacterial proliferation was monitored at 7 dpi by observation of GFP-expressing bacteria

under ultra violet (UV) light. Three biological replicates were generated in separate plates,

and three independent experiments were conducted under the same conditions.

H$_2$O$_2$ and cell death staining

Bacterial suspensions (5×10^7 cfu/ml) were injected into one half of each

Arabidopsis leaf using a syringe infiltration method. Accumulation of H$_2$O$_2$ in infected leaves

was monitored using a 3,3′-diaminobenzidine (DAB) staining method (Thordal-Christensen

et al., 1997). At 12 h post-infiltration, the leaves were cut and incubated with 1 mg/ml DAB

(Sigma-Aldrich) solution for 12 h as previously described (Thordal-Christensen *et al.*, 1997).

Cell death was observed by trypan blue staining as described previously (Koch & Slusarenko,

1990).

Microscopy

For scanning electron microscopy (SEM), flood-inoculated *Arabidopsis* leaves were fixed in 4% (w/v) paraformaldehyde and dehydrated in an ethanol series (30%, 50%, 70%, 96%, and 100%). The fixed leaves were then dried, coated with gold-palladium, and visualized using a scanning electron microscope (LEO 1455VP, Oberkochen, Germany) (Plotnikova *et al.*, 2000). For transmission electron microscopy (TEM), flood-inoculated *Arabidopsis* leaves were cut off, fixed overnight in 2.5% (w/v) glutaraldehyde, post-fixed in 2% (w/v) osmium tetroxide, dehydrated in ethanol, and embedded in the resin. After staining in 2% (w/v) uranyl acetate and lead citrate, samples were observed under an electron microscope (Bio-TEM; Tecnai G2 Spirit Twin; FEI, USA) (Chae & An, 2016).

 For fluorescence confocal microscopy, expression of GFP-labeled bacteria or GFP-tagged *Shigella* effector proteins in plants was observed under a Nikon laser scanning confocal microscope C2 (Nikon, Tokyo, Japan) using filter sets for GFP (λ_{ex}, 488 nm; λ_{em}, 505–530 nm) or RFP (λ_{ex}, 561 nm; λ_{em}, 570–620 nm). For each microscopic method, three leaves were used per treatment and at least three microscopic fields were observed for each leaf, including the control.

Expression of *Shigella* virulence genes in *Arabidopsis* plants

Total RNA was extracted from *Shigella*-infected leaves (from three plants) using RNAiso plus (#9108; TaKaRa, Otsu, Japan), according to the manufacturer's protocol. RT-PCR was performed using M-MLV reverse transcriptase (Invitrogen), according to the manufacturer's instructions. Quantitative RT-PCR was carried out in a CFX Connect™ Real Time System (BioRad, Hercules, CA, USA) using iQ™ SYBR® Green Supermix (BioRad) and primers specific for target genes (*ipaB*, *ipaC*, *icsA*, *icsB*, *virB*, and *virF*) (Bando *et al.*, 2010).

Immunoblotting

Total protein was extracted from *Shigella*- or *Agrobacterium*-infected leaves (from three plants) using denaturing extraction buffer [150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% Triton-X, 1× protease inhibitor cocktail (Roche, Basel, Switzerland), 0.4 M DTT, 1 M NaF, and 1 M Na_3VO_3]. The extracted proteins were separated on 12% SDS-PAGE gels and transferred to a PVDF membrane (Pierce, Rockford, IL, USA). Antibodies

specific for phospho-p44/p42 MAPK ERK1/2 (#4377; Cell Signaling Technology, Danvers, MA, USA), ERK1 (#sc-94; Santa Cruz, Dallas, TX, USA), hemagglutinin (HA) (#S2930; Clontech Laboratories, Mountain View, CA, USA), or GFP (#sc-9996; Santa Cruz) were used for immunoblot analyses. Target proteins were detected using ECL plus reagent (GE Healthcare, Wauwatosa, WI, USA) and visualized using an Alliance 9.7 chemiluminescence imaging system (UVITEC, Cambridge, UK).

Agrobacterium-mediated transient gene expression

A. tumefaciens strain GV2260 harboring the *GFP*, *OspF:GFP*, or *OspG:GFP* genes driven by the 35S promoter was prepared as described previously (Lee *et al.*, 2013). The inoculum ($OD_{600} = 0.4$) was infiltrated into 4-week-old *N. benthamiana* leaves using a 1 ml needleless syringe. To observe MEK2^{DD}-triggered cell death suppression by *Shigella* effectors, *Agrobacterium* ($OD_{600} = 0.4$) expressing HA:MEK2^{DD} was mixed with *Agrobacterium* containing *GFP*, *OspF:GFP*, or *OspG:GFP* at a 1:1 ratio and infiltrated into the leaves of *N. benthamiana*. Each experiment was repeated using at least three leaves from the plant, and each experiment included at least three different plants.

Statistical analysis

All data are expressed as the mean ± SD. The statistical significance of bacterial cell growth in infected plants was examined using the Student's t-test (Microsoft Office Excel) and an ANOVA (SPSS v.18, IBM, Armonk, NY, USA) (Moon *et al.*, 2016). Asterisks and letters indicate significant differences between samples ($P < 0.05$).

Results

Disease severity and symptomatic development in *A. thaliana* according to the infection method and *Shigella* infection load

To test whether the human pathogen *Shigella* is also a phytopathogen, we investigated the interaction of four *Shigella* strains representing three serogroups (*S. b*, *S. s*, *S. f* 2a, and *S. f* 5a) with the model plant *A. thaliana*. Inoculation of *Arabidopsis* plant leaves with high-density suspensions of *Pseudomonas syringae* pv. *tomato* DC3000 cells enables observation of the plant immune response (e.g., production of reactive oxygen species) (Chinchilla *et al.*, 2007; Smith & Heese, 2014) and necrosis (Katagiri *et al.*, 2002). Hence, we first inoculated *Arabidopsis* leaves with high-density bacterial cell suspensions ($\cong 5 \times 10^7$

cfu/ml) using syringe-pressure infiltration. This method is an artificial method of injecting

pathogens directly into the intercellular space of the plant cells; it is widely used to reduce

variation in plant cell immune responses among infiltrated plants (Liu *et al.*, 2015). *Pst hrcC*,

a mutant lacking the T3SS of *Pst*, and the non-pathogenic bacterium *E. coli* DH5α were used

as negative controls. *Arabidopsis* leaves infected with *Pst* showed an apparent accumulation

of H_2O_2 [at 1 d post-infection (dpi)] and necrosis (at 3 dpi), whereas no such responses were

observed in leaves treated with the buffer or either of the two non-pathogenic bacteria (Fig.

1a). In the majority of leaves that had been syringe-infiltrated with *S. s* and *S. f* 2a, cell death,

i.e., wilting and obvious accumulation of H_2O_2, was observed (Fig. 1a). On the other hand,

infection with *S. b* and *S. f* 5a resulted in a slight production of H_2O_2 compared with the

buffer treatment or the non-pathogenic bacterium treatments; unlike the other *Shigella* strains

investigated, no apparent cell death was observed (Fig. 1a). These observations indicated that

the bacterium can elicit an immune response in plants, and that the extent of the response is

strain-dependent.

Shigella is commonly known as a water-borne pathogen, with infection spread

through contaminated water (Pandey *et al.*, 2014). Hence, we chose to use a flood-inoculation

approach (Ishiga *et al.*, 2011), which is thought to closely mimic natural infection. Similarly

to *Pst*, when 2-week-old *A. thaliana* seedlings were inoculated with *S. s* or *S. f* 2a, clear

disease phenotypes were observed, such as yellowing and necrosis of the leaves (Fig. 1b). On

the other hand, obvious symptoms of disease were not apparent with *S. b* and *S. f* 5a, such as

after syringe-pressure inoculation (Fig. 1b). In addition to observing the symptoms, the

bacterial growth *in planta* was also evaluated, to detect initial plant adherence and

proliferation. Early attachment of all *Shigella* strains was more than 10 times lower than that

of *Pst* and *Pst hrcC* (Fig. 1c). However, 3 d after the inoculation, the extent of cell

amplification differed depending on the *Shigella* strain; nevertheless, the cell numbers of all

strains increased over 1 000 times. Notably, *S. s* cell number increased over 10^5 times. This

level of proliferation was comparable to that of plant pathogens.

Previously published studies demonstrated that several human pathogenic bacteria,

including *E. coli* O157:H7 and *Enterococcus faecalis*, can infect the leaves and roots of *A.*

thaliana (Jha *et al.*, 2005; Deering *et al.*, 2012). To determine whether the selected *Shigella*

strains could be transmitted through *A. thaliana* roots, we investigated if these strains existed

within the *Arabidopsis* roots, and if the plants could be infected via the roots. The tissues of

the flood-infected *Arabidopsis* plants were split into the aerial and root sections, and the

propagation of *Shigella* was assessed in each tissue type; a large number of bacteria were

detected in the former and also latter tissue types (Fig. S1). In addition, dropping the bacterial

solutions onto the root tips elicited similar disease symptoms to those caused by the *Shigella*

strains flooding, supporting these observations (Fig. S2).

Apparently different disease phenotypes and bacterial proliferation rates were

observed for different *Shigella* strains. This suggested that a variability of the infection

mechanisms among the strains, e.g., the adherence and invasion efficiency, might contribute

to the differences in pathogenesis. Differences in the nutritional requirements of bacterial

strains may constitute another reason for the differences in the growth rates within the

Arabidopsis host.

Plant cell entrance and internalization of *Shigella*

Since we observed that *Shigella* proliferate in *Arabidopsis*, we next directly

examined the plant cell entry and intercellular presence of the bacterium using SEM and

TEM. As anticipated, *Pst*, which infects plants via open stomata (Panchal *et al.*, 2016),

colonized the guard cells (Fig. 2). All *Shigella* strains invaded the plant through the stomata

(Fig. 2). Most plant pathogenic bacteria *in planta* infect and colonize the intercellular spaces

(Abramovitch *et al.*, 2006; Gao *et al.*, 2016). To investigate whether *Shigella* is capable of

intercellular colonization and induces plant responses, e.g., alteration of host cell structures,

we analyzed *A. thaliana* leaves infected with *Shigella* by TEM (Fig. 3). Indeed, the TEM

images revealed *Shigella* colonization of the intercellular spaces and bacterial attachment to

the host cell walls. The presence of the microbes in the intercellular space resulted in the

alteration of the host organelle structure, such as the separation of the plasma membrane from

the cell wall, the liberation of cell organelles, and disruption of chloroplasts (Fig. 3). This

effect was most pronounced in plant cells infected with *S. b* and *S. s*; further, *S. s* and *S. b*

were more commonly found in the intercellular spaces than *S. f* 2a and *S. f* 5a.

In a previous study, *Salmonella* was labeled with GFP and its presence in plant

cytoplasm was reported (Schikora *et al.*, 2008). We also labeled four *Shigella* strains with

GFP and then observed them in live plant tissues to see whether plant penetration by *Shigella*

occurs via the guard cells. *S. b*, *S. f* 2a and *S. f* 5a strains were successfully labeled with GFP

(the *S. s* strain was not) and then inoculated into the plant using the flood-inoculation method.

The leaf surface was then observed under a confocal microscope (Fig. S3). Similar to the

previous SEM results, GFP-labeled *Shigella* were observed in the guard cells, and strain *S. b*

was more abundant than strains *S. f* 2a and *S. f* 5a.

Taken together, these results indicated that, although the proliferative capacities of

the *Shigella* strains differ, the cells can infect the plants through the stomata, proliferate in the

intercellular spaces, and cause structural damage to the host during an infection.

T3S *Shigella* effectors are necessary for both epiphytic and endophytic growth in

Arabidopsis

GFP was used to label the *Shigella* strains to facilitate the observation of bacterial

migration in living plants. In addition, to study the role of *Shigella* T3SS in plant infection,

we used noninvasive variants of *S. f* 2a and *S. f* 5a, strains: Δvp and BS176, respectively

(Sansonetti *et al.*, 1982; Wenneras *et al.*, 2000; Shim *et al.*, 2007). First, we verified that the

GFP labeling did not visibly affect *Shigella* growth and pathogenicity (Figs 4a, S1, S4).

Strains *S. f* 2a, *S. f* 5a and their GFP-labeled derivatives were used to flood-inoculate

Arabidopsis seedlings, and their growth *in planta* was compared. Bacterial proliferation after

infection with Δvp or BS176 strains was 10 times lower than that after infection by parental

Shigella strains; similar results were confirmed when we observed growth of GFP-labeled

Shigella strains and mutants (Fig. 4a). To more readily observe bacterial growth, taking

advantage of the fluorescent tagging, the bacteria were dropped onto transparent root tissues

and their proliferation was compared (Fig. 4b). Under UV light, we observed that the GFP-

producing *S. f* 2a and *S. f* 5a strains colonized and multiplied along the whole root system 7 d

after the inoculation (Fig. 4b). Notably, *S. f* 2a:GFP cells formed biofilm-like communities on

the root surface to a greater extent than *S. f* 5a:GFP (Fig. 4b). Unlike *S. f* 2a:GFP and *S. f*

5a:GFP, strong fluorescence was not observed in the root tissues of plants treated with

Δvp:GFP or BS176:GFP, suggesting that T3S effectors of *Shigella* indeed play an important

role in the bacteria epiphytic growth in plants (Fig. 4b). We also confirmed that none of the

Shigella strains grew in the plant culture medium (Fig. S5).

To determine if these microorganisms were also present inside the root cells *per se*,

the roots of the infected plants were sampled and their surface was disinfected, and they were

observed under a fluorescence microscope (Fig. 4c). *S. f* 2a:GFP and *S. f* 5a:GFP were both

found in the proximal vascular tissue; the former also appeared to form communities in the

root epidermal tissue. On the other hand, communities inside the root cells formed by the

mutant strains were not readily apparent.

Altogether, these results suggested that the T3S effectors of *Shigella* that operate

during an infection in human are also required for the bacterial pathogenicity in plants.

Similar observations were also made for the plant pathogen *Pst* and its T3SS deletion mutant

Pst hrcC⁻ (Fig. S6a, b). However, *Pst*:GFP cells proliferated along the root surface but no

growth was observed in the vascular tissue inside the root. The *Pst hrcC⁻*:GFP cells were

rarely observed on either the root surface or inside the root.

Expression of *Shigella* T3S effector genes and activation of *Arabidopsis* MAPK signaling

during *Shigella* infection in *Arabidopsis* plants

To further investigate T3S effectors involved in plant pathogenesis, we measured

expression of key virulence effectors related to human pathogenesis, including *ipaB*, *ipaC*,

icsA, *icsB*, *virB*, and *virF*, in *S. f* 2a-, *S. f* 5a-, Δvp-, and BS176-infected *Arabidopsi*s plants.

These effectors play a role in mammalian cell lysis (*ipaB, ipaC*) and intracellular spread (*icsA*,

icsB), and regulation of virulence factor (*virF* and *virB*) expression (Ogawa *et al.*, 2008).

Total RNAs were isolated from *A. thaliana* leaves at 6 and 12 h after *Shigella* inoculation and

changes in expression of virulence genes were confirmed by quantitative RT-PCR (Fig. 5a).

Expression of all *Shigella* virulence genes examined in WT *S. f* 2a- or *S. f* 5a-infected

Arabidopsis leaves increased. In particular, induction was higher and faster in *S. f* 2a-infected

plants than in *S. f* 5a-infected plants, which is in agreement with the earlier results showing

that *S. f* 2a was more pathogenic to plants than *S. f* 5a. Expression of virulence genes was not

detected in plants infected with non-pathogenic mutants Δvp or BS176, similar to buffer-

treated control plants. These results indicate that common virulence factors regulate

pathogenesis in both plants and human intestinal cells.

To investigate the plant innate immune responses to *Shigella* infection, MAPK

phosphorylation was evaluated. The activation of MAPK by phosphorylation is a conserved

response of the earliest pathogen-triggered immune signaling in both plants and animals

(Zipfel, 2009). The flg22 peptide is a representative microbe-associated molecular pattern in

plants (Bethke *et al.*, 2009). In plants treated with flg22, pronounced MAPK phosphorylation

was apparent within 5 min of treatment and this response lasted up to 30 min (Fig. 5b). On

the other hand, MAPK phosphorylation in plants treated with *S. f* 2a or *S. f* 5a was reduced;

from 15 min on, it was strongly suppressed and almost completely disappearing after ca. 30

min (Fig. 5b). Meanwhile, in plants treated with the virulence plasmid-deficient mutants, Δvp

or BS176, MAPK activation was recovered, in contrast to *S. f* 2a- or *S. f* 5a-treated plants,

although the degree and duration of the phosphorylation were lower than those elicited by the

flg22 treatment (Fig. 5b). These observations indicated that *Shigella* suppresses the innate

immunity of *Arabidopsis* via its T3S virulence effectors.

Two *S. f* 5a effectors, OspF and OspG, function as virulence factors in *Arabidopsis*

To see if the *Shigella* effector protein really has pathogenic functions in plants as in

human cells, we next investigated its effector proteins OspF and OspG. These two proteins

interfere with the human innate immune response during infection (Kim *et al.*, 2005; Arbibe

et al., 2007; Ashida & Sasakawa, 2014; Pruneda *et al.*, 2014). To test whether OspF and

OspG are required for plant infection by *Shigella*, we tested single *S. f.* 5a Δ*ospF* and *S. f 5a*

Δ*ospG* deletion mutants. We observed a clear reduction in bacterial proliferation in plants

infected with *S. f.* 5a Δ*ospF* or *S. f* 5a Δ*ospG* when compared with wild-type *S. f* 5a (Fig. 6a).

Pathogenicity was fully restored by complementation of the mutations, indicating that these

effector proteins indeed play an important role in *Shigella* pathogenesis in plants (Fig. 6a).

We then monitored activation of MAPKs to determine whether plant immune suppression

was altered by single mutation of *OspF* and *OspG*. As shown in Figure 6B, *S. f.* 5a Δ*ospF*

induced stronger phosphorylation of MAPKs than wild-type *S. f* 5a; this induction was offset

by complementation with OspF. On the other hand, *S. f 5a*-deficient OspG showed little

change in the inhibition of MAPK phosphorylation compared to wild-type *S. f* 5a, but rather a

slight increase in MAPK activation due to overexpression of OspG (Fig. 6b). These results

suggest that OspG - induced plant immunosuppression may be present in the lower part of the

MAPK signaling or may be involved in the plant immune response other than inhibition of

MAPK activation. Taken together, these results indicate that *Shigella* effectors OspF or OspG

play an important role in increasing bacterial invasion and proliferation in plants as well as in

animal hosts.

To understand how the *Shigella* effectors OspF and OspG affect the pathogenesis of

plant pathogens, each gene was introduced into *Pst*. First, we used an AvrRpt2-derived T3SS

reporter system to determine whether the *Shigella* effector was delivered into plant cells via

Pst T3SS (Mudgett, 2005). In this system, AvrRpt2 (a T3S effector protein from

Pseudomonas syringae pv. tomato) interacts with the RPS2 protein of *Arabidopsis* plants to

induce plant cell death (Mudgett, 2005). We introduced the *ospF* and *ospG* genes into the

pBAV178 vector encoding the AvrRpt[101–255] fragment (Fig. S7a). When *Arabidopsis* leaves

were syringe-infiltrated with *Pst* producing various AvrRpt[101–255] constructs, *Pst* producing

$OspF_{S.f5a}$:AvrRpt2$^{101-255}$ or $OspG_{S.f5a}$:AvrRpt2$^{101-255}$ induced a strong cell death response at 1

dpi, while control *Pst* cells containing the empty vector did not, indicating that the *Shigella*

effectors OspF and OspG are successfully delivered into *Arabidopsis* cells via *Pst* T3SS (Fig.

7a). Full-length AvrRpt2 was used as a positive control for the cell death response.

Furthermore, we tested whether the pathogenicity of *Pst* increased in the presence of

Shigella OspF or OspG. Plants infected with *Pst* cells producing OspF:HA or OspG:HA

showed more severe disease symptoms than plants infected with the empty vector control;

furthermore, bacterial cell numbers were 10 times higher than those in the control (Fig. 7b, c).

Production of OspF:HA and OspG:HA in *Pst* was also confirmed by immunoblotting with an

anti-HA antibody (Fig. S7b). Overall, these observations suggest that the *Shigella* effectors

OspF and OspG function as virulence factors in plants.

OspF completely suppresses the MEK2^{DD}-dependent hypersensitive response

We next attempted to produce the OspF and OspG proteins in the plant cells, to

verify whether their known function in the human host is different from that in the plant cell.

To do this, subcellular localization of the two proteins fused with GFP was evaluated in the

plant cells. The DNA fragments encoding the OspF:GFP and OspG:GFP fusions were

inserted into a binary plant expression vector and transiently expressed in *N. benthamiana*

leaves using the *Agrobacterium* system (Fig. 8a). The fluorescence signal for both proteins

was strong in the plant cell nucleus; OspF:GFP fluorescence was also observed along the

cytoplasmic membrane, and punctate OspG:GFP fluorescence was observed in the cytosol

(Fig. 8b). In the host human cell, OspF localization is nuclear (Zurawski *et al.*, 2006; Arbibe

et al., 2007) and that of OspG is nuclear and cytoplasmic (Kim *et al.*, 2005; Zhou *et al.*, 2013;

de Jong & Alto, 2014), i.e., their subcellular localization is similar to that of plant cells.

Previously, *in vitro* studies demonstrated that OspF exerts a phosphothreonine lyase

activity and irreversibly removes phosphate groups from MAPK (Arbibe *et al.*, 2007; Li *et al.*,

2007). We used a phenotypic screening system involving a MEK2 (a tobacco MAP kinase

kinase2) mutant, MEK2^{DD}, to investigate whether the virulence associated with *Shigella*

OspF in plants was linked to the same mechanism of action as in animals. MEK2^{DD}, a

constitutively active mutant of MEK2, induces cell death when overproduced in plant leaves

(Yang *et al.*, 2001; Kim & Zhang, 2004). In this experiment, an HA:MEK2^{DD} clone fused to

an HA epitope tag at the N-terminus was used to monitor expression of MEK2^{DD}. As

expected, co-production of HA:MEK2^{DD} and GFP (control) resulted in a pronounced cell

death (Fig. 8c). Surprisingly, co-production of OspF:GFP, but not OspG:GFP, and MEK2^{DD}

completely inhibited the MEK2^{DD}-induced cell death (Fig. 8c). The production of the two

effectors fused with GFP was then evaluated in the MEK2^{DD}-producing plant leaves; the

production of OspF:GFP was apparent, but that of OspG:GFP was not (Fig. 8d), even though

both proteins were stably produced in the absence of MEK2^{DD} (Fig. 8a). The degradation of

the OspG protein may have been associated with the activated MAPK. Indeed, as assessed by

immunoblotting with specific anti-phosphorylated MAPK antibodies, MAPK

phosphorylation was very weak in the OspF:GFP-producing plant samples in comparison

with GFP- or OspG:GFP-producing plant samples (Fig. 8d). The production of MEK2^{DD}

protein was confirmed in all samples using anti-HA antibodies (Fig. 8d). These observations

strongly suggested that the *Shigella* effector OspF suppressed plant MAPK activation through

its dephosphorylation, as in the human host.

DISCUSSION

In the current study, we investigated the interaction of the human pathogenic

bacterium, *Shigella*, with a novel host, the *Arabidopsis* plant. We demonstrated that four

Shigella strains, *S. b*, *S. s*, *S. f* 2a, and *S. f* 5a, promote proliferation and confer virulence on

Arabidopsis. It has been clearly demonstrated that, in human, *Shigella* initially enters the

epithelial layer via the M cells through transcytosis, leading to the invasion of the basolateral

surfaces of the intestinal epithelial cells. A subsequent gut inflammation leads to the

activation of the innate immune response (Phalipon & Sansonetti, 2007). We demonstrated in

the current study that, in addition to the human host, *Shigella* infects *Arabidopsis*, through the

vascular tissues of the root and leaf stomata pores (Figs 2, 4). In particular, *S. s* and *S. b*

formed relatively wide clusters in the surrounding areas, including the guard cells (Figs 2, S3).

Interestingly, we found that the four studied strains of *Shigella* associate with the plant cells

and induce different plant responses. The bacterial loads of *S. s in planta* were relatively

higher than those of the other serogroups. By contrast, *S. f* 5a infection was associated with

reduced bacterial proliferation and disease symptoms compared with the other strains (Figs 1,

S1, S2). *S. f* 2a and *S. f* 5a, which belong to the same serogroup (Lindberg *et al.*, 1991),

elicited distinctly different plant responses with respect to disease symptoms. The specific

plant responses, which were different for bacterial strains within the same serogroup, suggest

that virulent effectors may play a relatively more important role in plant pathogenesis than

PTI. These observations also indicate that specific plant immune systems may be useful in

the search for novel virulence factors expressed by different *Shigella* serogroups.

471   Many Gram-negative bacterial pathogens utilize common infection strategies to

colonize and invade plant and animal cells, and pathogenicity appears to depend on highly

conserved T3SSs, which deliver the effector proteins to host cells (Buttner & Bonas, 2003).

By using avirulent mutant strains, we were able to show that effectors that regulate the

pathogenesis of shigellosis in humans also play a central role in regulating pathogenesis in

Arabidopsis. We showed that secretion of T3S effectors is required to trigger both pathogenic

symptoms and expression of the invasive phenotypes in plants (Fig. 4). Furthermore, the

effector proteins impacted MAPK-dependent innate immune responses in *Arabidopsis* (Figs 5,

6). Although reduced colonization of T3SS-deficient pathogenic *E. coli* in plants was

previously reported, in contrast with the present study, they suggested that *E. coli* uses the

T3SS apparatus for attachment to leaves, rather than for delivery of effector proteins (Shaw *et*

al., 2008). The relevance of T3SS for infection of plants with *Salmonella* remains unclear due

to the different effects of T3SS function on *Salmonella*-plant interactions (Iniguez *et al.*, 2005;

Schikora *et al.*, 2011; Shirron & Yaron, 2011). Until now, the transfer of effector proteins

into plant cells by T3SS of animal bacterial pathogens has not been reported. Our results

indirectly suggest that *Shigella* effectors can be transferred into plant cells; however, direct

evidence, such as visualization of the delivery of effectors using split fluorescent proteins

(Mach, 2017), is required.

489   Expression of the T3SS of *Shigella* is regulated at the transcriptional level and is

activated at a permissive temperature ($\geq 32°C$) (Erickson, 2012). We were able to observe

expression of the T3SS genes of *Shigella* under temperatures at which plants grow ($22 \pm 3°C$)

(Fig. 5a). A recent study showed that elevation of the temperature increases T3SS-mediated

virulence of the phytopathogen *Pst* in plants, which is in contrast with the negative effect of

high temperature on expression of the T3SS of *Pst in vitro* (Huot *et al.*, 2017). Regardless of

the temperature of host cells, it will be interesting to determine whether *Shigella* regulates

T3SS gene expression *in vivo* and to identify factors that influence T3SS gene expression

other than plant temperature.

498   The current study provides new insights into host invasion mechanisms utilized by

499   *Shigella* to infect an alternative host, the plant *Arabidopsis*. Studying trans-kingdom

500   pathogenesis involving human-adapted pathogens, such as *Shigella*, may uncover novel

501   pathogenic mechanisms uniquely activated in response to specific hosts. When we isolated

502   the two *Shigella* effectors OspF and OspG, and produced them in plant cells, their

localization coincided with that in the animal cells (Fig. 8b), and it was apparent that the

production of both proteins increased the virulence of plant pathogens (Fig. 7b, c). In addition,

we confirmed that OspF inhibits the innate immune response of plants via the same

enzymatic activity as in animals (Fig. 8c, d). OspG inhibits the host immune response by

inhibiting the activity of NF-κB by blocking the degradation of IκB in animals (Ashida *et al.*,

2015). Plants possess an IκB-like protein called NIM1 (Ryals *et al.*, 1997), but no other

published studies have investigated whether a NF-κB–induced immune response exists in

plants. In the current study, we demonstrated the ability of OspG to enhance the

phytopathogenicity of non-*Shigella* bacteria (Fig. 7b, c) and also observed the degradation of

OspG:GFP upon the constitutive activation of MAPK signaling (Fig. 8c, d). The existence of

a plant immune signaling pathway similar to that of animal NF-κB, which would also be the

target of OspG, may hence be assumed. Characterization of the previously unrecognized

stress-activated mediators of the innate immunity in plants upon infection with food-borne

pathogens would help better define the defensive functions of plants. Finally, the

characterization of plants as an alternative host for food-borne pathogens will be critical in

developing effective means to prevent their transmission and disease.

Acknowledgments

We thank Dr. Myung Hee Kim for initial support with the preparation of pathogenic *Shigella*

strains, Dr. Jean T. Greenberg for providing the pBAV178, pBAV179, and pME6012

plasmids, and Dr. Cha Young Kim for providing the HA:MEK2^{DD} clone.

Authors' contributions

JMP, SHJ, and JL conceived and designed the study and wrote the manuscript. SHJ, JL, and

DHL carried out the experiments. JMP, SHJ, JL, DWK, and CMR analyzed and discussed the

data.

References

Abramovitch RB, Anderson JC, Martin GB. 2006. Bacterial elicitation and evasion of

plant innate immunity. *Nat Rev Mol Cell Biol* **7**(8): 601-611.

**Arbibe L, Kim DW, Batsche E, Pedron T, Mateescu B, Muchardt C, Parsot C,

Sansonetti PJ. 2007.** An injected bacterial effector targets chromatin access for transcription

factor NF-kappaB to alter transcription of host genes involved in immune responses. *Nat Immunol* **8**(1): 47-56.

Ashida H, Mimuro H, Sasakawa C. 2015. Shigella Manipulates Host Immune Responses by Delivering Effector Proteins with Specific Roles. *Frontiers in Immunology* **6**(219).

Ashida H, Sasakawa C. 2014. Shigella hacks host immune responses by reprogramming the host epigenome. *EMBO J* **33**(22): 2598-2600.

Bando SY, Moreno AC, Albuquerque JA, Amhaz JM, Moreira-Filho CA, Martinez MB. 2010. Expression of bacterial virulence factors and cytokines during in vitro macrophage infection by enteroinvasive Escherichia coli and Shigella flexneri: a comparative study. *Mem Inst Oswaldo Cruz* **105**(6): 786-791.

Bethke G, Unthan T, Uhrig JF, Poschl Y, Gust AA, Scheel D, Lee J. 2009. Flg22 regulates the release of an ethylene response factor substrate from MAP kinase 6 in Arabidopsis thaliana via ethylene signaling. *Proc Natl Acad Sci U S A* **106**(19): 8067-8072.

Buttner D, Bonas U. 2003. Common infection strategies of plant and animal pathogenic bacteria. *Curr Opin Plant Biol* **6**(4): 312-319.

Cevallos-Cevallos JM, Gu G, Danyluk MD, van Bruggen AH. 2012. Adhesion and splash dispersal of Salmonella enterica Typhimurium on tomato leaflets: effects of rdar morphotype and trichome density. *Int J Food Microbiol* **160**(1): 58-64.

Chae Y, An YJ. 2016. Toxicity and transfer of polyvinylpyrrolidone-coated silver nanowires in an aquatic food chain consisting of algae, water fleas, and zebrafish. *Aquat Toxicol* **173**: 94-104.

Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nurnberger T, Jones JD, Felix G, Boller T. 2007. A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* **448**(7152): 497-500.

de Jong MF, Alto NM. 2014. Thinking outside the Osp(G)--kinase activation by E2-ubiquitin. *EMBO J* **33**(5): 403-404.

Deering AJ, Mauer LJ, Pruitt RE. 2012. Internalization of E. coli O157:H7 and Salmonella spp. in plants: A review. *Food Research International* **45**(2): 567-575.

Erickson MC. 2012. Internalization of Fresh Produce by Foodborne Pathogens. *Annual Review of Food Science and Technology* **3**(1): 283-310.

Ferens WA, Hovde CJ. 2011. Escherichia coli O157:H7: animal reservoir and sources of human infection. *Foodborne Pathog Dis* **8**(4): 465-487.

Franz E, Semenov AV, van Bruggen AH. 2008. Modelling the contamination of lettuce with

Escherichia coli O157:H7 from manure-amended soil and the effect of intervention strategies. *J Appl Microbiol* **105**(5): 1569-1584.

Gao X, Huang Q, Zhao Z, Han Q, Ke X, Qin H, Huang L. 2016. Studies on the Infection, Colonization, and Movement of Pseudomonas syringae pv. actinidiae in Kiwifruit Tissues Using a GFPuv-Labeled Strain. *PLoS One* **11**(3): e0151169.

Gu G, Cevallos-Cevallos JM, Vallad GE, van Bruggen AH. 2013a. Organically managed soils reduce internal colonization of tomato plants by Salmonella enterica serovar Typhimurium. *Phytopathology* **103**(4): 381-388.

Gu G, Cevallos-Cevallos JM, van Bruggen AH. 2013b. Ingress of Salmonella enterica Typhimurium into tomato leaves through hydathodes. *PLoS One* **8**(1): e53470.

Guchi B, Ashenafi M. 2010. Microbial load, prevalence and antibiograms of salmonella and Shigella in lettuce and green peppers. *Ethiop J Health Sci* **20**(1): 41-48.

Holt KE, Baker S, Weill F-X, Holmes EC, Kitchen A, Yu J, Sangal V, Brown DJ, Coia JE, Kim DW, et al. 2012. Shigella sonnei genome sequencing and phylogenetic analysis indicate recent global dissemination from Europe. *Nat Genet* **44**(9): 1056-1059.

Hong CE, Kwon SY, Park JM. 2016. Biocontrol activity of Paenibacillus polymyxa AC-1 against Pseudomonas syringae and its interaction with Arabidopsis thaliana. *Microbiological Research* **185**: 13-21.

Huot B, Castroverde CDM, Velásquez AC, Hubbard E, Pulman JA, Yao J, Childs KL, Tsuda K, Montgomery BL, He SY. 2017. Dual impact of elevated temperature on plant defence and bacterial virulence in Arabidopsis. *Nature Communications* **8**(1): 1808.

Iniguez AL, Dong Y, Carter HD, Ahmer BMM, Stone JM, Triplett EW. 2005. Regulation of Enteric Endophytic Bacterial Colonization by Plant Defenses. *Molecular Plant-Microbe Interactions* **18**(2): 169-178.

Ishiga Y, Ishiga T, Uppalapati SR, Mysore KS. 2011. Arabidopsis seedling flood-inoculation technique: a rapid and reliable assay for studying plant-bacterial interactions. *Plant Methods* **7**: 32.

Jha AK, Bais HP, Vivanco JM. 2005. Enterococcus faecalis mammalian virulence-related factors exhibit potent pathogenicity in the Arabidopsis thaliana plant model. *Infect Immun* **73**(1): 464-475.

Jones JD, Dangl JL. 2006. The plant immune system. *Nature* **444**(7117): 323-329.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY™ vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci* **7**(5): 193-195.

Katagiri F, Thilmony R, He SY. 2002. The Arabidopsis thaliana-pseudomonas syringae

interaction. *Arabidopsis Book* **1**: e0039.

Kennedy CK. 1971. Induction of colicin production by high temperature or inhibition of

protein synthesis. *J Bacteriol* **108**(1): 10-19.

Kim CY, Zhang S. 2004. Activation of a mitogen-activated protein kinase cascade induces

WRKY family of transcription factors and defense genes in tobacco. *The Plant Journal* **38**(1):

142-151.

Kim DW, Lenzen G, Page AL, Legrain P, Sansonetti PJ, Parsot C. 2005. The Shigella

flexneri effector OspG interferes with innate immune responses by targeting ubiquitin-

conjugating enzymes. *Proc Natl Acad Sci U S A* **102**(39): 14046-14051.

King EO, Ward MK, Raney DE. 1954. Two simple media for the demonstration of

pyocyanin and fluorescin. *J Lab Clin Med* **44**(2): 301-307.

Koch E, Slusarenko A. 1990. Arabidopsis is susceptible to infection by a downy mildew

fungus. *Plant Cell* **2**(5): 437-445.

Lee JH, Kim YC, Choi D, Park JM. 2013. Identification of novel pepper genes involved in

Bax- or INF1-mediated cell death responses by high-throughput virus-induced gene silencing.

Int. J. Mol. Sci. **14**(11): 22782-22795.

Li H, Xu H, Zhou Y, Zhang J, Long C, Li S, Chen S, Zhou JM, Shao F. 2007. The

phosphothreonine lyase activity of a bacterial type III effector family. *Science* **315**(5814):

1000-1003.

Lindberg AA, Karnell A, Weintraub A. 1991. The lipopolysaccharide of Shigella bacteria

as a virulence factor. *Rev Infect Dis* **13 Suppl 4**: S279-284.

Liu X, Sun Y, Korner CJ, Du X, Vollmer ME, Pajerowska-Mukhtar KM. 2015. Bacterial

Leaf Infiltration Assay for Fine Characterization of Plant Defense Responses using the

Arabidopsis thaliana-Pseudomonas syringae Pathosystem. *J Vis Exp*(104).

Mach J. 2017. Tracking the Bacterial Type III Secretion System: Visualization of Effector

Delivery Using Split Fluorescent Proteins. *The Plant Cell* **29**(7): 1547-1548.

Martínez-Vaz BM, Fink RC, Diez-Gonzalez F, Sadowsky MJ. 2014. Enteric Pathogen-

Plant Interactions: Molecular Connections Leading to Colonization and Growth and

Implications for Food Safety. *Microbes and Environments* **29**(2): 123-135.

Moon JY, Lee JH, Oh C-S, Kang H-G, Park JM. 2016. Endoplasmic reticulum stress

responses function in the HRT-mediated hypersensitive response in Nicotiana benthamiana.

Molecular Plant Pathology **17**(9): 1382-1397.

Mortality GBD, Causes of Death C. 2016. Global, regional, and national life expectancy,

all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a

systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **388**(10053): 1459-

1544.

Mudgett MB. 2005. New insights to the function of phytopathogenic bacterial type III

effectors in plants. *Annu Rev Plant Biol* **56**: 509-531.

Murashige T, Skoog F. 1962. A Revised Medium for Rapid Growth and Bio Assays with

Tobacco Tissue Cultures. *Physiologia Plantarum* **15**(3): 473-497.

Neumann C, Fraiture M, Hernandez-Reyes C, Akum FN, Virlogeux-Payant I, Chen Y,

Pateyron S, Colcombet J, Kogel KH, Hirt H, et al. 2014. The Salmonella effector protein

SpvC, a phosphothreonine lyase is functional in plant cells. *Front Microbiol* **5**: 548.

Ogawa M, Handa Y, Ashida H, Suzuki M, Sasakawa C. 2008. The versatility of Shigella

effectors. *Nat Rev Micro* **6**(1): 11-16.

Onodera NT, Ryu J, Durbic T, Nislow C, Archibald JM, Rohde JR. 2012. Genome

Sequence of Shigella flexneri Serotype 5a Strain M90T Sm. *Journal of Bacteriology* **194**(11):

3022.

Panchal S, Roy D, Chitrakar R, Price L, Breitbach ZS, Armstrong DW, Melotto M. 2016.

Coronatine Facilitates Pseudomonas syringae Infection of Arabidopsis Leaves at Night. *Front*

Plant Sci **7**: 880.

Pandey PK, Kass PH, Soupir ML, Biswas S, Singh VP. 2014. Contamination of water

resources by pathogenic bacteria. *AMB Express* **4**: 51.

Phalipon A, Sansonetti PJ. 2007. Shigella's ways of manipulating the host intestinal innate

and adaptive immune system: a tool box for survival? *Immunol Cell Biol* **85**(2): 119-129.

Plotnikova JM, Rahme LG, Ausubel FM. 2000. Pathogenesis of the human opportunistic

pathogen Pseudomonas aeruginosa PA14 in Arabidopsis. *Plant Physiol* **124**(4): 1766-1774.

Pruneda JN, Smith FD, Daurie A, Swaney DL, Villen J, Scott JD, Stadnyk AW, Le

Trong I, Stenkamp RE, Klevit RE, et al. 2014. E2~Ub conjugates regulate the kinase

activity of Shigella effector OspG during pathogenesis. *EMBO J* **33**(5): 437-449.

Runyen-Janecky LJ, Payne SM. 2002. Identification of chromosomal Shigella flexneri

genes induced by the eukaryotic intracellular environment. *Infect Immun* **70**(8): 4379-4388.

Ryals J, Weymann K, Lawton K, Friedrich L, Ellis D, Steiner HY, Johnson J, Delaney

TP, Jesse T, Vos P, et al. 1997. The Arabidopsis NIM1 protein shows homology to the

mammalian transcription factor inhibitor I kappa B. *Plant Cell* **9**(3): 425-439.

Sansonetti PJ, Kopecko DJ, Formal SB. 1982. Involvement of a plasmid in the invasive

ability of Shigella flexneri. *Infect Immun* **35**(3): 852-860.

Schikora A, Carreri A, Charpentier E, Hirt H. 2008. The dark side of the salad:

Salmonella typhimurium overcomes the innate immune response of Arabidopsis thaliana and

shows an endopathogenic lifestyle. *PLoS One* **3**(5): e2279.

Schikora A, Garcia AV, Hirt H. 2012. Plants as alternative hosts for Salmonella. *Trends*

Plant Sci **17**(5): 245-249.

Schikora A, Virlogeux-Payant I, Bueso E, Garcia AV, Nilau T, Charrier A, Pelletier S,

Menanteau P, Baccarini M, Velge P, et al. 2011. Conservation of Salmonella infection

mechanisms in plants and animals. *PLoS One* **6**(9): e24112.

Schroeder GN, Hilbi H. 2008. Molecular pathogenesis of Shigella spp.: controlling host cell

signaling, invasion, and death by type III secretion. *Clin Microbiol Rev* **21**(1): 134-156.

Semenov AM, Kuprianov AA, van Bruggen AH. 2010. Transfer of enteric pathogens to

successive habitats as part of microbial cycles. *Microb Ecol* **60**(1): 239-249.

Shamloul M, Trusa J, Mett V, Yusibov V. 2014. Optimization and utilization of

Agrobacterium-mediated transient protein production in Nicotiana. *J Vis Exp*(86).

Shaw RK, Berger CN, Feys B, Knutton S, Pallen MJ, Frankel G. 2008.

Enterohemorrhagic Escherichia coli Exploits EspA Filaments for Attachment to Salad Leaves.

Applied and Environmental Microbiology **74**(9): 2908-2914.

Shim DH, Suzuki T, Chang SY, Park SM, Sansonetti PJ, Sasakawa C, Kweon MN. 2007.

New animal model of shigellosis in the Guinea pig: its usefulness for protective efficacy

studies. *J Immunol* **178**(4): 2476-2482.

Shirron N, Yaron S. 2011. Active suppression of early immune response in tobacco by the

human pathogen Salmonella Typhimurium. *PLoS One* **6**(4): e18855.

Smith JM, Heese A. 2014. Rapid bioassay to measure early reactive oxygen species

production in Arabidopsis leave tissue in response to living Pseudomonas syringae. *Plant*

Methods **10**(1): 6.

Solomon EB, Yaron S, Matthews KR. 2002. Transmission of Escherichia coli O157:H7

from contaminated manure and irrigation water to lettuce plant tissue and its subsequent

internalization. *Appl Environ Microbiol* **68**(1): 397-400.

Thordal-Christensen H, Zhang Z, Wei Y, Collinge DB. 1997. Subcellular localization of

H2O2 in plants. H2O2 accumulation in papillae and hypersensitive response during the

barley—powdery mildew interaction. *The Plant Journal* **11**(6): 1187-1194.

Vinatzer BA, Jelenska J, Greenberg JT. 2005. Bioinformatics correctly identifies many

type III secretion substrates in the plant pathogen Pseudomonas syringae and the biocontrol

isolate P. fluorescens SBW25. *Mol Plant Microbe Interact* **18**(8): 877-888.

Wang K, Kang L, Anand A, Lazarovits G, Mysore KS. 2007. Monitoring in planta

bacterial infection at both cellular and whole-plant levels using the green fluorescent protein

variant GFPuv. *New Phytologist* **174**(1): 212-223.

Wei J, Goldberg MB, Burland V, Venkatesan MM, Deng W, Fournier G, Mayhew GF,

Plunkett G, Rose DJ, Darling A, et al. 2003. Complete Genome Sequence and Comparative

Genomics of Shigella flexneri Serotype 2a Strain 2457T. *Infection and Immunity* **71**(5): 2775-

2786.

Wenneras C, Ave P, Huerre M, Arondel J, Ulevitch RJ, Mathison JC, Sansonetti P. 2000.

Blockade of CD14 increases Shigella-mediated invasion and tissue destruction. *J Immunol*

164(6): 3214-3221.

Yang KY, Liu Y, Zhang S. 2001. Activation of a mitogen-activated protein kinase pathway

is involved in disease resistance in tobacco. *Proc Natl Acad Sci U S A* **98**(2): 741-746.

Zhang R, Zhou HW, Cai JC, Zhang J, Chen GX, Nasu M, Xie XY. 2011. Serotypes and

extended-spectrum beta-lactamase types of clinical isolates of Shigella spp. from the

Zhejiang province of China. *Diagn Microbiol Infect Dis* **69**(1): 98-104.

Zhou Y, Dong N, Hu L, Shao F. 2013. The Shigella type three secretion system effector

OspG directly and specifically binds to host ubiquitin for activation. *PLoS One* **8**(2): e57558.

Zipfel C. 2009. Early molecular events in PAMP-triggered immunity. *Curr Opin Plant Biol*

12(4): 414-420.

Zurawski DV, Mitsuhata C, Mumy KL, McCormick BA, Maurelli AT. 2006. OspF and

OspC1 are Shigella flexneri type III secretion system effectors that are required for

postinvasion aspects of virulence. *Infect Immun* **74**(10): 5964-5976.

Figure legends

Fig. 1 *Shigella* cells proliferate and induce disease symptoms in *Arabidopsis* plants. (a) *Arabidopsis* leaves were syringe-infiltrated with buffer or bacterial suspensions (5×10^7 cfu/ml). *S. s*, and *S. f* 2a induced wilting of infected leaves. Accumulation of reactive oxygen species was measured by 3,3′-diaminobenzidine staining 1 d after infiltration. The fractions indicate the number of times that the symptoms were observed relative to the number of tests. (b, c) *Arabidopsis* seedlings in 1/2 MS medium were flood-inoculated with distilled H_2O containing 0.025% Silwet L-77 (buffer) or bacterial suspensions (5×10^5 cfu/ml). (b) The

disease phenotypes of *Arabidopsis* infected with *Shigella* spp. by flood-inoculation.

Photographs of representative disease symptoms were taken at 3 dpi. (c) Bacterial cell

numbers were evaluated on Days 0 and 3 after flood-inoculation. The bars represent the mean

± SD of three replicates and asterisks indicate significant differences between samples (*$P <$

0.05, *t*-test). All experiments were repeated three times independently and representative

results are shown.

Fig. 2 *Shigella* cells adhere to the *Arabidopsis* epidermis and enter the plant via open stomata.

Arabidopsis seedlings were flood-inoculated with a buffer or bacterial suspensions (5×10^6

cfu/ml). After 24 h, the leaves were fixed and analyzed under a SEM. The *Pst* cells were

observed over the stomata. *Shigella* strains were associated with the stomata. Bar, 10 μm.

Representative image showing bacteria around open stomata (indicated by white arrows). The

results are representative of two independent experiments.

Fig. 3 *Shigella* cells associate with and colonize plant cell structures. *Arabidopsis* seedlings

were flood-inoculated with a buffer or bacterial suspensions (5×10^6 cfu/ml). After 24 h, the

leaves were evaluated under a TEM. The *Shigella* strains colonized the intercellular spaces.

Bar, 5 μm (left panels). Bar, 1 μm (right panels). Each column of micrographs represents a

different magnification. TEM images revealed the *Shigella* (white arrows) attached to the cell

wall in the intercellular spaces and altered mesophyll cells. Black arrows indicate separation

of the plasma membrane from the cell wall and disruption of chloroplasts. The results are

representative of two independent experiments.

Fig. 4 The virulence of a plasmid-deficient mutant of *Shigella* is impaired in the *Arabidopsis*

model. (a) *Arabidopsis* seedlings in 1/2 MS medium were flood-inoculated with *S. f* 2a, Δvp,

S. f 5a, or BS176, and their GFP-labeled variants (5×10^5 cfu/ml). The bacterial cell numbers

in planta were determined at 3 dpi. Bars represent the mean ± SD of three replicates and the

different letters indicate significant differences between samples ($P < 0.05$, one-way

ANOVA). (b, c) The root tips of *Arabidopsis* seedlings were drop-inoculated with

suspensions of GFP-labeled bacteria (5×10^7 cfu/ml). (b) Biofilm formation and colonization

of *Arabidopsis* seedling roots by GFP-labeled *Shigella* was observed under white (WL) and

ultra violet (UV) light at 7 dpi. (c) Images of GFP-labeled *Shigella* wild types in the

epidermis or vascular tissue of the *Arabidopsis* roots acquired using a fluorescence

microscope at 7 dpi. Bar, 100 μm. All experiments were repeated at least three times, and

representative results are shown.

Fig. 5 Expression of *Shigella* effectors and MAPK activation in *Arabidopsis* plants in
response to virulence plasmid-deficient *Shigella* mutants. (a, b) *Arabidopsis* leaves were
syringe-infiltrated with buffer, 1 μM flg22, or bacterial suspension (5×10^6 cfu/ml), and
samples were collected at the indicated times. (a) Transcription of *Shigella* effectors (*ipaB*,
ipaC, *icsA*, *icsB*, *virB*, and *virF*) in infected *A. thaliana* leaves was analyzed by qRT-PCR.
qRT-PCR results were normalized to expression of 16s rRNA. Expression of effectors by
plasmid-deficient mutant strains was compared with that in the WT. Data are expressed as the
mean ± SD of three replicates. (b) Immunoblotting was performed using either anti-phospho-
p44/42 (ERK1/2, upper panels) or anti-ERK1 (lower panels) antibodies. All experiments
were repeated at least three times, each with similar results.

Fig. 6 The *Shigella* effector proteins OspF and OspG underpin the bacterial virulence in
plants. (a) The virulence of the OspF- or OspG-deficient *Shigella* mutants is impaired in the
Arabidopsis model. *Arabidopsis* seedlings in 1/2 MS medium were flood-inoculated with *S. f*
5a, Δ*ospF*, Δ*ospF*+p*ospF*, Δ*ospG*, or Δ*ospG*+p*ospG* (5×10^5 cfu/ml). Photographs of
representative disease symptoms were taken, and bacterial cell numbers *in planta* were
determined, at 3 dpi. The bars represent the mean ± SD of three replicates. The different
letters indicate significant differences between samples ($P < 0.05$, one-way ANOVA). (b)
MAPK activation in *Arabidopsis* plants in response to infection by ospF or ospG deletion
mutants. *Arabidopsis* leaves were infiltrated with each bacterial strain as described previously
and then collected at the indicated times. Anti-phospho-p44/42 (ERK1/2, upper panels) or
anti-ERK1 (lower panels) antibodies were used for immunoblotting. All experiments were
repeated at least three times, with similar results.

Fig. 7 Increased virulence of phytopathogen *Pst* in the presence of *Shigella* effectors OspF or
OspG. (a) *Arabidopsis* leaves were syringe-infiltrated with *Pst* cells (5×10^7 cfu/ml)
producing AvrRpt2, carrying the empty vector (pME6012), and producing OspF:AvrRpt2^{101-}
^{255} or OspG:AvrRpt2$^{101-255}$. The hypersensitive response in the infiltrated leaves was
monitored using the trypan blue staining and photographed at 1 dpi. (b, c) The *Arabidopsis*
plants were sprayed with *Pst* cells carrying an empty vector (pME6012), or producing
OspF:HA or OspG:HA (5×10^7 cfu/ml). (b) The disease symptoms were monitored 4 d after
spraying. (c) The proliferation of *Pst* cells producing the *Shigella* effector proteins in

Arabidopsis, 0 and 4 d after spray-inoculation. The bars represent the mean ± SD of six

replicates, and different letters indicate significant differences between samples ($P < 0.05$,

two-way ANOVA). All experiments were repeated at least three times.

Fig. 8 Co-expression of OspF:GFP suppresses tobacco MEK2^{DD}-triggered cell death. (a) GFP,

OspF:GFP, or OspG:GFP was produced in *N. benthamiana* leaves upon infiltration with

Agrobacterium carrying the appropriate expression constructs (OD$_{600}$ 0.4). The production of

the *Shigella* effector proteins in *N. benthamiana* 2 d after the infiltration was evaluated by

immunoblotting using anti-GFP antibodies. The RuBisCo protein was stained by Ponceau S

as a loading control. (b) Subcellular localization of GFP, OspF:GFP, OspG:GFP, and

NLS:RFP (a nuclear localization marker) in *N. benthamiana* leaves. Bar, 50 μm. (c, d) GFP,

OspF:GFP or OspG:GFP was co-produced in *N. benthamiana* leaves harboring HA:MEK2^{DD}.

(c) Production of OspF:GFP suppressed MEK2^{DD}-dependent cell death by 5 d after the co-

infiltration. (d) Production of specific proteins in samples from panel **C** was analyzed by

immunoblotting using anti-GFP, anti-HA, anti-phospho-p44/42 (ERK1/2), and anti-ERK1

antibodies. Ponceau S was used to stain the RuBisCo protein (loading control). The asterisk

indicates the size of the GFP protein not associated with effector proteins.

(a)

(b)

3 dpi

(c)

Buffer

Pst

S. b

S. s

S. f 2a

S. f 5a

(a)

(b)

	WL	UV	UV (GFP filter)
S. f 2a:GFP			
Δvp:GFP			
S. f 5a:GFP			
BS176:GFP			

7 dpi

(c)

S. f 2a:GFP

Δvp:GFP

S. f 5a:GFP

BS176:GFP

7 dpi

(a)

Relative expression (Fold)

(b)

(a)

(b)

(a)

Pst

AvrRpt2

pME6012
(E.V)

OspF$_{S.f5a}$:
AvrRpt2$^{101-255}$

OspG$_{S.f5a}$:
AvrRpt2$^{101-255}$

(b)

Pst

pME6012
(E.V)

OspF$_{S.f5a}$:HA

OspG$_{S.f5a}$:HA

(c)

Bacterial cell numbers
(log cfu/leaf disc)

■ Day 0
■ Day 4

pME6012
(E.V)

OspF$_{S.f5a}$:HA

OspG$_{S.f5a}$:HA

Pst

(a)

kDa

GFP OspF:GFP OspG:GFP

50 —

37 —

anti-GFP

25 — *

Ponceau S

(b)

Merged   GFP   NLS:RFP   BP

GFP

OspF:GFP

OspG:GFP

2 dpi

(c)

HA:MEK2^{DD}

GFP   OspF:GFP   GFP   OspG:GFP

WL

UV

5 dpi

(d)

HA:MEK2^{DD}

kDa   GFP   OspF:GFP   GFP   OspG:GFP

50 —

37 —

anti-GFP

25 — *

anti-HA

Ponceau S

anti-phospho
-p44/42 (ERK1/2)

anti-ERK1

Ponceau S