1
13 **Title:**
14
15 # Small RNA-mediated genomic silencing promotes
16 telomere stability in the absence of telomerase in *C.*
17 *elegans*
23 **Authors:**
24

25 Charlie Longtine[1,2,#], Stephen Frenk[1,2,#], Evan H. Lister-Shimauchi[1,2], and Shawn
26 Ahmed[1,2,3]*
27
28 **Affiliations:**
29
30 [1]Department of Genetics, [2]Department of Biology, [3]Lineberger Comprehensive Cancer
31 Center, University of North Carolina, Chapel Hill, NC, USA
32
33
34 # Equal contributions
35 *Correspondence: shawn@med.unc.edu

36
37
38 Running title: small RNAs promote telomere stability

39   Keywords: telomere, telomerase, siRNA, sRNA, RNA interference

Abstract

We define an endogenous small RNA pathway in *Caenorhabditis elegans* that protects

telomeres via Dicer, the perinuclear Argonaute protein WAGO-1, the nuclear Argonaute

protein HRDE-1, and the HP1 protein HPL-2 that stabilizes the heterochromatin mark

H3K9me2. Loss of telomerase induces biogenesis of subtelomeric small RNAs,

whereas loss of both telomerase and small RNA-mediated genome silencing induces

expression of the telomeric lncRNA TERRA, DNA damage, and accelerated sterility.

These phenotypes were rescued by exogenous telomeric small RNAs and by loss of the

DNA damage response protein EXO-1. Although ATRX promotes heterochromatin

deposition and is lost in human tumors that maintain their telomeres via ALT, we

detected only a modest role for the *C. elegans* ATRX ortholog XNP-1 in telomere

protection in the absence of telomerase. Moreover, neither XNP-1 nor small RNA

factors that promote telomere stability in the absence of telomerase repress ALT in *C.

elegans*. We propose that small RNA-mediated heterochromatin formation and

telomerase function redundantly to repress a form of telomeric DNA damage that

induces TERRA expression.   We conclude that small RNA-mediated genome silencing

promotes telomere stability in the absence of telomerase.

Article Summary

We found that a number of small RNA biogenesis factors and heterochromatin proteins

promoted telomere stability in the absence of telomerase. The telomeric non-coding

RNA TERRA and DNA damage signaling were induced upon loss of small RNAs and

telomerase, indicating that small RNAs directly or indirectly repress TERRA expression

and telomeric DNA damage, both of which were suppressed when telomerase acts to

heal telomeric damage. These findings reveal a functional interaction between small

RNAs and telomere biology that is relevant to the fields of aging, genome stability and

small RNAs.

Introduction

Telomeres are repetitive nucleoprotein structures that protect the ends of linear

eukaryotic chromosomes from degradation and recognition as double-stranded DNA

breaks (Maciejowski and de Lange, 2017, Shay, 2016). Deficiency for telomerase

reverse transcriptase in most human somatic cells results in progressive telomere

erosion over cell divisions, which eventually triggers replicative senescence (Holohan et

al., 2014, Shay, 2016). This telomere clock acts as a barrier to tumorigenesis, but may

also contribute to organismal aging through impaired stem cell function, regeneration,

and organ maintenance (Donate and Blasco, 2011, Shay, 2016). Regulating the rate of

telomere erosion in telomerase-negative cells is therefore crucial for maintaining the

balance between tumor suppression and age-related tissue decline (Li et al., 2017,

Maciejowski and de Lange, 2017). Tumor cells escape telomere-triggered senescence

by stabilizing telomere length, either through reactivation of telomerase or through

activation of the telomerase-independent recombination-mediated telomere

maintenance pathway termed Alternative Lengthening of Telomeres (ALT) (Pickett and

Reddel, 2015).

Telomeres have features characteristic of heterochromatin in that they associate with

the heterochromatin protein HP1 and contain transcriptionally repressive histone marks

such as methylation of H3K9 and H4K20 (Benetti et al., 2007, McMurchy et al., 2017).

Nonetheless, telomeres are transcribed into the G-rich long noncoding RNA (lncRNA)

termed TERRA, a feature widely conserved among eukaryotes (Azzalin et al., 2007).

TERRA is transcribed from the subtelomere into the telomere by RNA polymerase II

and is upregulated at shortened and damaged telomeres, where it may recruit

telomerase and chromatin modifiers to promote telomere elongation (Azzalin and

Lingner, 2015, Cusanelli et al., 2013). Telomere elongation represses expression of

TERRA through increased heterochromatin marks at telomeres such as histone H3K9

methylation and HP1 binding (Arnoult et al., 2012). In contrast, shortened telomeres of

mice deficient for telomerase or of mouse cells that overexpress the telomere-binding

protein TRF2 display loss of heterochromatic marks at telomeres and subtelomeres

(Benetti et al., 2007, Benetti et al., 2008). Telomere length has therefore been inversely

correlated with TERRA expression and the heterochromatic state of telomeres, and

TERRA expression in mouse cells has recently been linked to H3K9me3 and other

heterochromatic histone marks present at mammalian telomeres (Montero et al., 2018).

Transcriptional silencing and heterochromatin deposition at repetitive segments of the

genome such as the pericentromere require long non-coding RNA (lncRNA) transcripts

that promote biogenesis of and interact with small RNAs (sRNAs) that direct

heterochromatin factors to target these genomic loci (Holoch and Moazed, 2015,

Reinhart and Bartel, 2002). Endogenous sources of double-stranded RNA such as

sense-antisense transcript hybrids can be processed into small RNAs that play roles in

creating and maintaining sRNA-mediated heterochromatin. In this context, RNA

expression is paradoxically required for genomic silencing (Holoch and Moazed, 2015).

Bidirectional transcription of telomeres has been linked to production of telomeric

sRNAs that are associated with the presence of subtelomeric DNA methylation in

Arabidopsis (Vrbsky et al., 2010). Similar telomeric sRNAs have also been observed in

Tetrahymena, mouse stem cells, and human cell lines (Cao et al., 2009, Couvillion et

al., 2009, Rossiello et al., 2017), and can promote the DNA damage response at

uncapped telomeres in mammals (Cao et al., 2009, Couvillion et al., 2009, Rossiello et

al., 2017). However, their functional relevance at canonical telomeres and their

interaction with telomerase is not well understood.

Germline transcription in *C. elegans* is regulated in part by a network of small RNA

pathways that promote silencing or expression of target loci in a sequence-specific

manner (Billi et al., 2014). Primary sRNA classes trigger production of secondary

sRNAs that can promote transcriptional silencing or mRNA destruction. One class of

primary sRNAs are 21 nucleotide Piwi-Interacting RNAs (piRNAs) that are created from

thousands of tiny genes that repress expression of potentially harmful foreign genetic

elements including transposons and transgenes (Batista et al., 2008, Das et al., 2008,

Kasper et al., 2014). piRNAs interact with PRG-1, a member of the Piwi clade of

Argonaute proteins, which can recognize piRNA targets based on sequence

complementarity with up to 3 mismatches (Bagijn et al., 2012). RNA target recognition

triggers the recruitment of RNA-dependent RNA polymerases (RdRPs) that act in

Mutator bodies, which are associated with perinuclear P granules, to create *de novo* 22

nucleotide secondary sRNAs with a 5' guanine (22G RNAs) (Pak and Fire, 2007).

These 22G RNAs are loaded onto worm-specific Argonaute proteins (WAGOs) such as

the nuclear WAGO HRDE-1, which promotes heterochromatin deposition at target loci

(Ashe et al., 2012, Buckley et al., 2012). Germline transcription is protected by a

separate set of 22G RNAs that interact with the anti-silencing Argonaute protein CSR-1

(Avgousti et al., 2012, Wedeles et al., 2013). Aside from the piRNA pathway, 22G RNAs

can be created in response to primary sRNAs derived from other sources. For example,

DCR-1, the *C. elegans* homologue of the highly conserved RNAse class III enzyme

Dicer, can cleave endogenous double stranded RNA to create primary 26G RNAs,

which bind distinct Argonaute proteins to trigger RDRP-dependent silencing through the

22G RNA pathway described above (Bernstein et al., 2001, Ketting et al., 2001, Knight

and Bass, 2001).

Here we describe an endogenous sRNA and heterochromatin pathway that functions

redundantly with telomerase to repress TERRA and promote telomere stability.

Materials and Methods

Strains: Unless otherwise noted, all strains were maintained at 20°C on nematode

growth medium plates seeded with *Escherichia coli* OP50. Strains used include Bristol

N2 ancestral, *wago-1(tm1414), trt-1(ok410), unc-29(e193), dpy-5(e61), dcr-1(mg375),*

hrde-1(tm1200), rde-1(ne219), rde-4(ne299), rrf-3(mg373), rrf-2(ok210), eri-1(mg366),

mut-7(pk204), mut-7(pk720), mut-14(pk738), rbr-2(tm1231), spr-5(by101), xnp-

1(ok1823), xnp-1(tm678), hpl-1(tm1624), hpl-2(ok916), hpl-2(tm1479), and *exo-*

1(tm1842). A *trt-1 unc-29* double mutant, where *unc-29* is tightly linked to *trt-1*, was

used to create strains used in this study. Double, triple, and quadruple mutants were

created using standard marker mutations and verified based on phenotype or genotype.

mut-7 and *mut-14* mutants are known to accumulate transposon-induced mutations, so

these strains were outcrossed several times before use.

<u>DAPI staining</u>: One-day-old adults were washed twice in M9, soaked in 150 µL of

400ng/mL DAPI in ethanol solution for 30 min or until evaporated, rehydrated in 2 mL of

M9 solution overnight at 4°C, and mounted in 8 µL VECTASHIELD mounting medium

(Vector Laboratories). Chromosome counts were performed under 60x magnification

and a 359nm excitation wavelength using a Nikon Eclipse E800 microscope.

<u>qPCR</u>: Non-starved mixed-stage worms were washed with M9 and RNA was extracted

using 1 mL TRIZOL according to the manufacturer's instructions. For RT-PCR analysis,

total RNA was treated twice with DNase I and was reverse transcribed with primers for

a reference gene *act-1* and a telomere-specific primer $(GCCTAA)_3$ using SuperScript II

reverse transcriptase (Invitrogen). For SYBR green reactions, subtelomere specific

primers were designed within 200bp of the telomere-subtelomere junction.

<u>RNA FISH</u>: RNA FISH was performed as previously described (Sakaguchi et al., 2014)

with several modifications. In brief, non-starved, mixed-stage animals were washed

twice in M9 and three times in 1x DEPC-treated PBS. Worms were fixed in 1mL fixation

buffer (1x DEPC-treated PBS with 3.7% formaldehyde) for 45 min rotating at room

temperature then washed twice in 1mL 1x DEPC-treated PBS. Fixed animals were

permeabilized overnight at 4°C in 70% ethanol in DEPC-treated H_2O, then washed in 1

mL wash buffer [10% (vol/vol) formamide in 2x RNases-free SSC]. Aliquots of each

sample were removed prior to hybridization and treated with RNaseA for 1h at 37°C to

ensure probe specificity to RNA. Hybridization was performed at 30°C overnight in 100

µL hybridization buffer (10% dextran sulfate, 2x RNase-free SSC, 10% deionized

formamide, 0.02% RNase-free BSA, 50 µg *E. coli* tRNA) with 1.25 µM probe. Cy5-

labelled $(GCCTAA)_3$ probe was used to detect TERRA transcripts. Animals were

washed twice with wash buffer, counterstained with 25 ng/mL DAPI for 30 min at room

temperature, and mounted on glass slides in VECTASHIELD mounting medium (Vector

Laboratories) before epifluorescence microscopy.

<u>IF:</u> IF was carried out as previously described (Phillips et al., 2009) by fixing dissected

gonads of 1 day old adults in 2% PFA for 10 minutes followed by a 1 minute postfixation

in methanol. Rabbit anti-pS/TQ (Cell Signaling Technology) was used at a 1:500 dilution

and mouse anti-RNA:DNA hybrid S9.6 at 1:500 dilution. Stained gonads were

counterstained with DAPI and mounted with VECTASHIELD (Vector laboratories)

before epifluorescence microscopy.

RNAi: RNAi clone expressing dsRNA targeting *C. elegans* telomeres was constructed

using NEB HiFi DNA assembly mix according to the manufacturer's instructions. 200

bp of telomeric repeats were amplified from cTel55x by PCR with 20 bp overlap to SpeI

digested L4440. Correct constructs were transformed into HT115 and RNAi was

performed as previously described (McMurchy et al., 2017).

Southern Blot: *C. elegans* genomic DNA was digested overnight with Hinf1 (New

England Biolabs), separated on a 0.8% agarose gel at 1.5 V/cm, and transferred to a

neutral nylon membrane (Hybond-N, GE Healthcare Life Sciences). A digoxigenin-

labelled telomere probe was hybridized and detected as described (Ahmed and

Hodgkin, 2000). Migration distance was measured using ImageJ.

Mrt and ALT assays: When telomerase-deficient *trt-1(ok410) C. elegans* mutants are

passaged by using the standard Mortal Germline (Mrt) assay developed in our

laboratory, which involves passaging six L1 larval animals to fresh plates weekly, 100%

become sterile within 20–30 generations. When *trt-1(ok410)* strains are passaged

weekly by transferring a 1cm^2 chunk of agar populated with ≈200–400 animals, 10-20%

of the strains survive for hundreds of generations. Mantel-Cox Log Rank Analysis was

used to determine differences of transgenerational lifespan between strains. A z-test

was performed to compare proportion of ALT survivors in chunked strains.

RNA-seq: Animals were grown at 20°C on 100 mm NGM plates seeded with OP50

bacteria. Two biological replicates were prepared for each genotype. RNA was

extracted using Trizol (Ambion) followed by isopropanol precipitation. RNA samples

were treated with 20U RNA 5' polyphosphatase (Lucigen, RP8092H) for 30 minutes at

37°C to remove 5' triphosphate groups. Library preparation and sequencing was

performed at the UNC School of Medicine High-Throughput Sequencing Facility

(HTSF). RNA-seq libraries were prepared using the NEXTflex Small RNA Library Prep

Kit for Illumina (Bio Scientific). Libraries were sequenced on an Illumina HiSeq 2500 to

produce approximately twenty million 50 bp single-end reads per library.

Analysis of high throughput sequencing data: Adapter trimming was performed as

required using the bbduk.sh script from the bbmap suite3 and custom scripts (Bushnell,

2016). Reads were then filtered for 22G species or all species between 18 and 30

nucleotides in length using a custom python script. Reads were then converted to fasta

format and mapped to the *C. elegans* genome (WS251) using bowtie (Langmead et al.,

2009) with the following options: -M 1 -v 0 -best -strata. Uniquely mapping sRNAs were

detected by replacing the -M 1 parameter with -m 1. To detect telomere-mapping

sRNAs, reads were mapped to seven repeats of perfect telomere sequence (TTAGGC),

allowing for zero mismatches. Unmapped reads were re-mapped to the telomere

sequence, this time allowing for one mismatch. This step was repeated to detect reads

that map to telomeres with two or three mismatches. Genes with differential sRNA

abundance in *trt-1* were identified using DESeq2 (Love et al., 2014). To detect changes

in the abundance of subtelomeric sRNAs in *trt-1* and *trt-1; dcr-1*, 22G reads mapping to

the 2kb of sequence adjacent to the telomere were counted for each chromosome arm.

Counts were normalized by dividing by the total number of mapped reads for each

library. A pseudocount of 1 was added to each normalized value to avoid division by

zero errors. Replicates were averaged and the log2-fold change between mutant and

wildtype was calculated for each subtelomere. Analysis of sequencing data was

performed using the R statistical computing environment (Team, 2017). Genome

regions were visualized using the Gviz R package (Hahne and Ivanek, 2016).

Immunoprecipitation datasets were used in Figure 3G, H but were excluded from the

analysis shown in Figure 3D-F. To detect sRNAs targeting transposons, 22G reads

were mapped to the *C. elegans* transposon consensus sequences downloaded from

Repbase with bowtie, allowing for up to two mismatches. ChIP-seq reads were mapped

to the genome as fastq files as described above without the size-filtering step. Read

coverage was calculated at each base using bedtools (Quinlan and Hall, 2010).

Coverage values were normalized by dividing by the number of total mapped reads for

each library and enrichment was calculated by dividing the normalized coverage for IP

libraries by the normalized coverage of their respective input libraries.

Results

Loss of endogenous sRNAs or XNP-1/ATRX does not promote ALT

High levels of TERRA and loss of the histone H3.3 chaperone subunits ATRX and

DAXX commonly occur in human ALT cancers (Flynn et al., 2015, Lovejoy et al., 2012).

The ATRX/DAXX complex mediates histone H3.3 incorporation at telomeres and

interacts with methylated H3K9 and HP1 to promote heterochromatin formation (Wong

et al., 2010), which has been proposed to repress ALT by protecting telomeres from

replication stress and TERRA expression (Udugama et al., 2015). We have previously

shown that 10-20% of *C. elegans* strains that are deficient for telomerase can be

maintained indefinitely through the activation of ALT (Cheng et al., 2012). Because

human ALT cancers are characterized by ATRX-mediated heterochromatin dysfunction,

we reasoned that small RNA factors that promote heterochromatin silencing might

regulate the establishment of ALT. We tested this hypothesis by passaging *trt-1*

telomerase mutant strains that were deficient for biogenesis of either primary or

secondary sRNAs by transferring hundreds of worms per strain weekly, which reliably

allows survival of 10-20% of strains via ALT (Cheng et al., 2012). Loss of either primary

sRNAs (*trt-1; dcr-1)* or downstream secondary sRNAs (*trt-1; mut-7* and *trt-1; mut-14)* did

not significantly alter the proportion of strains that survive via ALT (Figure 1A, Table

S1). Similarly, loss of the histone H3.3 chaperone XNP-1/ATRX itself did not affect the

establishment or maintenance of ALT (Table S1), indicating that loss of neither sRNA-

nor XNP-1/ATRX-mediated heterochromatin formation is sufficient to promote ALT in *C.*

elegans.

Small RNA and heterochromatin factors repress rapid sterility when telomerase is

deficient

C. elegans trt-1 telomerase mutant strains do not remain fertile via ALT when 6 worms

are passaged weekly. Instead, when 20-40 strains per genotype are passaged,

telomere erosion and chromosome fusion results in uniform sterility after growth for 20-

30 generations (Ahmed and Hodgkin, 2000, Cheng et al., 2012, Meier et al., 2006). The

sterility of late-generation *trt-1* mutants is analogous to crisis observed in checkpoint-

deficient telomerase-negative mammalian cells. Intriguingly, *trt-1* mutant lines lacking

primary or secondary sRNA biogenesis factors became sterile significantly faster than

trt-1 single mutants when they were passaged by transferring 6 L1 larvae per week

(Figure 1B, Table S2). sRNA-deficient *trt-1* mutant strains rapidly accumulated end-to-

end chromosome fusions by generation F12, a characteristic of telomerase mutants that

have grown for more than 20 generations (Figure 1G). *C. elegans* chromosome fusions

cause non-disjunction events during meiosis that elicit dominant embryonic lethality or

X0 male phenotypes in progeny of heterozygous F1 animals. We asked if early-

generation *trt-1; dcr-1* mutant animals possessed chromosome fusions by performing

genetic crosses that would be capable of revealing a single heterozygous end-to-end

chromosome fusion (Ahmed and Hodgkin, 2000). We crossed four independently

derived *trt-1 dcr-1* F4 hermaphrodites with N2 wild-type males but failed to detect fused

chromosomes that were present in generation F24 *trt-1* mutant controls (p = 3.841e-05

for males and p = 3.61e-05 for dead embryos, Mann Whitney U test) (Table S3).

However, early-generation *trt-1; dcr-1* mutant strains displayed high levels of DNA

damage that were not observed for *trt-1* single mutant controls (Figure 1H). Together,

these results suggest that loss of small RNAs and telomerase elicits telomere damage

but not telomere fusions in early generations. The rapid sterility of these strains in later

generations occurs in response to end-to-end chromosome fusions that arise from

critically shortened telomeres.

We created a number of sRNA-deficient *trt-1* double mutants to define additional small

RNA factors that promote telomere stability in the absence of telomerase. The telomeric

sRNA pathway included RDE-4, which binds endogenous dsRNA and forms a complex

with the Dicer nuclease DCR-1, and the Argonaute RDE-1, which acts with the RNA-

dependent RNA polymerase RRF-3 to promote production of dsRNA-derived primary

sRNAs (Gent et al., 2010, Steiner et al., 2009, Tabara et al., 2002) (Figure 1B). The

MUTATOR proteins MUT-7 and MUT-14, which promote amplification of 22G

secondary sRNAs that can promote transcriptional silencing in the nucleus (Billi et al.,

2014, Ketting et al., 1999, Tijsterman et al., 2002), were also required to prevent rapid

sterility in *trt-1* mutants (Figure 1B). Two other small RNA factors, RRF-2 and ERI-1,

did not affect *trt-1* mutant sterility (Figure 1B), indicating that the endogenous 26G

sRNA pathway that requires ERI-1 (Gent et al., 2010, Han et al., 2009) is dispensable

for telomere stability in the absence of telomerase.

We crossed *dcr-1* and *mut-7* single mutants with wild-type males, singled F2 progeny

that were homozygous mutant for *dcr-1* or *mut-7*, and isolated genomic DNA from F4,

F6 and F8 generation descendants. Southern blotting revealed discrete telomere bands

that display modest telomere erosion for freshly homozygous *dcr-1* and *mut-7* mutant

telomeres (Figure S1A, B), although none of the small RNA single mutants we tested

became progressively sterile or accumulated chromosome fusions (Figure 1G). We

asked if the accelerated sterility of small RNA-deficient *trt-1* mutant strains might be

explained by short telomeres inherited from gametes derived from small RNA-deficient

grandparents using genetics, by crossing a *trt-1; dcr-1* double mutant hermaphrodite

with *dcr-1* mutant males, singling *trt-1 +/-; dcr-1 -/-* F1 hermaphrodites and isolating *trt-*

1; dcr-1 double mutant F2. This cross was repeated 15 times to create a *trt-1; dcr-1*

double mutant strain whose telomeres should be exclusively derived from the *dcr-1*

mutant background. The *trt-1; dcr-1* 15x strain became sterile slightly slower than

freshly created *trt-1; dcr-1* double mutants, although this difference was not statistically

significant (p = 0.07) (Figure 1F). These results indicate that shortened telomeres

transmitted from the *dcr-1* mutant cannot explain the rapid sterility of *trt-1; dcr-1* double

mutants. We analyzed the dynamics of telomere erosion in sRNA-deficient *trt-1* mutant

strains using terminal restriction fragment analysis but did not observe a significant

difference in the rate of telomere erosion when passaging multiple worms per plate, in

comparison to *trt-1* single mutant controls (p > 0.7 in both cases) (Figure S1A, B).

However, when we passaged individual worms for each genotype, we observed

significant levels of telomere instability in sRNA-deficient *trt-1* mutant strains in

comparison to *trt-1* single mutant controls. Southern blotting revealed that *trt-1; mut-7*

and *trt-1; dcr-1* mutants had a significantly larger variance in the rate of telomere

erosion than *trt-1* mutants, which resulted from atypical shortening and elongation

events (Figure S2A, B).

We hypothesized that small RNAs might repress rapid sterility by directing chromatin

factors to telomeres to promote deposition of heterochromatic marks. *C. elegans*

telomeres have previously been reported to be enriched for H3K9me2 and bound by the

HP1 homolog HPL-2, which promotes H3K9me2 deposition and heterochromatin

formation, but not by the other *C. elegans* HP1 homolog HPL-1, which interacts with

H3K9me3 (Canzio et al., 2014, McMurchy et al., 2017). We confirmed that H3K9me2

but not H3K9me3 was enriched at *C. elegans* telomeres using high throughput

sequencing data sets generated from chromatin immunoprecipitation (ChIP)

experiments performed by independent groups (McMurchy et al., 2017, Zeller et al.,

2016). Moreover, we observed small H3K9me2 domains that extended into the

subtelomere at all *C. elegans* chromosome ends (Figure 3A-C). Similar to primary and

secondary sRNA-deficient *trt-1* strains, loss of HPL-2 but not HPL-1 caused accelerated

senescence in *trt-1* mutants (Figure 1C). H3K4 methylation, which is associated with

regions of active transcription, must be removed by demethylases prior to deposition of

repressive H3K9 methylation (Greer et al., 2010, Li et al., 2008). We found that loss of

either of the two *C. elegans* H3K4 demethylases RBR-2/KDM5 or SPR-5/LSD1 caused

rapid sterility of *trt-1* mutants (Figure 1C). Thus, H3K4 demethylation and H3K9me2

deposition prevent telomere instability in the absence of telomerase.

The histone remodeling protein XNP-1/ATRX has previously been implicated in

telomeric heterochromatin formation and interacts with unmethylated H3K4 and

methylated H3K9 deposited on the histone variant H3.3 (Udugama et al., 2015). *trt-1*

xnp-1 double mutants became sterile significantly faster than *trt-1* single mutants but not

as fast as small RNA deficient *trt-1; mut-7* double mutants (Figure 1D). As loss of both

xnp-1 and *mut-7* did not exacerbate the accelerated sterility phenotype of sRNA-

deficient *trt-1* mutant strains (Figure 1D), these genes function in the same telomere

stability pathway. This suggests that XNP-1 mediates some, but not all, of the small

RNA response to DNA damage at telomeres, possibly because sRNA-dependent

methylation of histones other than H3.3, such as canonical histone H3, can promote

telomere stability.

Increased expression of TERRA in small RNA and chromatin deficient *trt-1* strains

We hypothesized that DCR-1 and RDE-4 may interact with telomeric dsRNA produced

from bidirectional transcription of telomeres or interstitial telomere sequences to create

primary sRNAs that are then amplified into a secondary sRNA population that triggers

small RNA-dependent silencing of TERRA (Rossiello et al., 2017, Vrbsky et al., 2010).

To test whether small RNA and chromatin factors regulate telomeric transcription we

measured the abundance of TERRA by qRT-PCR at three chromosome ends using

primers within their H3K9me2-coated subtelomere domains (Figure 3A-C). Compared

to *trt-1* single mutant controls, TERRA was upregulated in early-generations in three

strains - *trt-1; rde-4* and *trt-1; mut-14* and *trt-1; spr-5* double mutants - but not in *trt-1* or

sRNA-deficient single mutant controls (Figure 2A). These results are distinct from

previous observations that show that TERRA transcription can be specifically induced at

damaged or shortened telomeres of mammalian cells that possess telomerase (Porro et

al., 2014, Rossiello et al., 2017).

We next used RNA Fluorescence *In Situ* Hybridization (FISH) to confirm the effects on

TERRA expression and to examine its cell and tissue localization. In wild-type control

and *trt-1* mutant hermaphrodite and male worms, TERRA was expressed at low levels

in the soma as well as in the pachytene region of the meiotic germline (Figures 2B-D,

S3-4), which corresponds to the location of several lncRNA-mediated germline silencing

processes (Taylor et al., 2015), including Meiotic Silencing of Sex Chromosomes in

mammals and *C. elegans* (Heard and Turner, 2011) and a transcriptional burst of

piRNAs in mammals that primes genomic silencing in germ cells prior to gametogenesis

(Gou et al., 2014*). Compared to both wild-type control and *trt-1* or *dcr-1* single mutant

strains, *trt-1; dcr-1* double mutant strains had increased expression of TERRA in the

germline (Figures 2B-D, S3) and soma (Figures S3-4) by the F6 generation. We

observed an increase in the number of TERRA foci per nucleus in sRNA-deficient *trt-1*

mutants, which are likely to occur at telomeres (Figure 2D). We also observed TERRA

expression by RNA FISH in the cytoplasm and nucleoplasm of sRNA-deficient *trt-1* cells

(Figure 2B-C, S3), which might be consistent with possible non-telomeric functions of

TERRA (Chu et al., 2017, Roake and Artandi, 2017) or with a role for TERRA in

protecting telomeres *in trans* (Montero et al., 2018). This suggests that endogenous

sRNAs may repress telomeric transcription in the absence of telomerase, although it is

possible that subtelomeric transcription increases in sRNA-deficient telomerase mutants

in response to telomeric DNA damage (Figure 1H).

Subtelomeric sRNAs are upregulated in telomerase mutants

Telomeric sRNAs have been observed in protists, plants, and mammals (Cao et al.,

2009, Couvillion et al., 2009, Vrbsky et al., 2010). We recently interrogated 92 publicly

available small RNA sequencing libraries created from RNA isolated from various *C.*

elegans strains and found rare reads that mapped perfectly to the *C. elegans* telomere

repeat $(TTAGGC)_n$, occurring at a frequency of 1 per 10,000,000 reads (Frenk et al.,

2019). piRNAs with up to 3 mismatches can target regions of the *C. elegans* genome for

silencing (Bagijn et al., 2012), and we found that telomeric sRNAs with up to 3

mismatches were substantially more abundant than those with perfect telomere repeat

sequence (1 per 100,000) (Frenk et al., 2019). We compared levels of telomeric sRNAs

with mismatches from wildtype with those of mutants for small RNA genes that promote

telomere stability in the absence of telomerase, such as *dcr-1*, *rde-4*, *mut-7*, and *mut-*

14, and found that levels of telomeric sRNAs were sometimes depleted in these

backgrounds, but not always (File S1).

We asked if loss of telomerase might alter levels of telomeric sRNAs in a manner that

might depend on Dicer by isolating RNA from wild-type control, *trt-1* single mutant, and

trt-1; dcr-1 double mutant strains and preparing small RNA libraries. We sequenced

these libraries deeply in an effort to detect rare perfect telomeric sRNAs. We obtained

~20,000,000 reads each for 6 data sets and found that our *trt-1; dcr-1* data sets were

strongly depleted of 22G RNAs that were previously reported to depend on *dcr-1*

(Figure S5A) (Gent et al., 2010). To assess TRT-1-dependent changes in sRNAs

throughout the genome, we counted the number of 22G reads mapping uniquely

antisense to protein-coding genes and found upregulation of sRNAs at 720 loci and

downregulation at 456 loci in *trt-1* animals compared with wild-type controls (Figure

S5B). Of the upregulated loci, 465 were not upregulated in *trt-1; dcr-1* animals

compared with wildtype, indicating DCR-1 dependence. We did not find significant GO

terms in loci with either up or downregulated sRNA production. Given that *trt-1* is

expressed at high levels in *C. elegans* germ cells (Meier et al., 2006), we asked whether

the loci with differential sRNA production were enriched for germline-enriched genes.

We used previously published data on germline expression (Ortiz et al., 2014) and

found significant enrichment for germline-enriched genes in both upregulated and

downregulated loci (p < 1e-19 and p < 1e-33 respectively, Pearson's Chi Square test

with Bonferroni correction) (Figure S5C). This may reflect an epigenetic response to

deficiency for telomerase in germ cells, even though early-generation *trt-1* mutants from

which the RNA was prepared had normal levels of fertility. sRNAs mapping to

transposons were unaffected by loss of TRT-1, with the exception of the CER-10 LTR

retrotransposon, for which sRNAs were three-fold downregulated (Figure S5D).

We found that levels of telomeric sRNAs with 1 to 3 mismatches did not change

substantially in sRNAs from *trt-1* single mutants or *trt-1; dcr-1* double mutants, but

detected no sRNAs composed of perfect telomere repeats in any genetic background

(Figure 4A). We next examined 22G RNAs at subtelomeric regions (defined as 2kb of

sequence adjacent to each telomere) and found that these were present in wild-type

samples at the left end of chromosome *III*, and that their abundance was increased two-

fold in *trt-1* single mutants and their presence almost entirely eliminated in *trt-1; dcr-1*

double mutants (Figure 4C, E). Subtelomeric sRNAs were also observed within 1kb of

the telomere at the left ends of chromosomes *I* and *IV* in *trt-1* mutants and not in

wildtype or *trt-1; dcr-1* double mutants (Figure 4B, D). The right ends of Chromosomes *I*

and *III* and the left end of Chromosome *II* all showed at least a two-fold increase in

subtelomeric sRNAs in *trt-1* and/or *trt-1; dcr-1* double mutants (Figure 4E). We

observed a significant bias for subtelomeric sRNAs whose sequences correspond to the

C-rich telomere strand (antisense to TERRA) for the left ends of Chromosomes *I* and *III*

(p < 1e-13 and p < 1e-43 respectively, binomial test with Bonferroni correction) and for

the right arm of Chromosome *I* (p < 1e-172, binomial test with Bonferroni correction)

while the nucleotide sequences of subtelomeric sRNAs at the left arm of Chromosome

IV and the right arm of Chromosome *III* showed a bias for subtelomeric TERRA,

suggesting that they were created from RNA containing the C-rich telomeric sequence

(p < 1e-11, p = 0.006 respectively, binomial test with Bonferroni correction) (Figure 4F).

These observations suggest that telomere dysfunction triggers the upregulation of small

RNA biogenesis at multiple subtelomeres through both DCR-1-dependent and -

independent mechanisms. sRNA biogenesis might be induced at a subset of

subtelomeres when telomerase is dysfunctional because 1) shortened telomeres

inherited from *trt-1* mutant gametes that created *trt-1* single mutants might trigger sRNA

biogenesis, or 2) DNA damage that occurs stochastically at specific subtelomeres elicits

sRNA biogenesis. Subtelomeric siRNAs that correspond to the G- or C-rich telomeric

strand at specific subtelomeres could be related to the presence of transcripts that can

be targeted for sRNA biogenesis at these telomeres.

We next interrogated data sets of small RNAs that immunoprecipitate with Argonaute

proteins that are known to promote silencing in the germline, HRDE-1 and WAGO-1, as

well as with the germline anti-silencing Argonaute CSR-1 (Gu et al., 2009, Claycomb et

al., 2009, Ashe et al., 2012). Enrichment of small RNAs for each protein was calculated

by dividing the number of reads in the immunoprecipitate library by the number of reads

in the corresponding input library. Telomeric sRNAs were enriched in WAGO-1

immunoprecipate relative to input, but this was not the case for HRDE-1 or CSR-1

(Figure 3D). Enrichment of subtelomeric small RNAs in WAGO-1 was detected at all

chromosome ends tested apart from the right arm of Chromosome *I* (some

chromosome ends were not tested due to an absence of reads in the input library).

Enrichment was also detected for HRDE-1 at all chromosome ends excluding the right

arm of Chromosome *I* and the left arm of Chromosome *III* (Figure 3E). Interestingly,

CSR-1 showed the opposite pattern of enrichment to WAGO-1 and HRDE-1, with

enrichment at none of the chromosome ends apart from the right end of Chromosome *I*.

The ribosomal DNA, the most highly expressed locus in the *C. elegans* genome, is

immediately adjacent to this telomere (Wicky et al., 1996), and it is therefore possible

that sRNA activity takes a predominantly anti-silencing function at this chromosome

end. We asked if either Argonaute that associates with telomeric or sub-telomeric

sRNAs, WAGO-1 or to a lesser degree HRDE-1, promotes telomere maintenance in the

absence of telomerase and found that *wago-1 trt-1* and *trt-1; hrde-1* double mutants

both displayed rapid sterility phenotypes in comparison to *trt-1* single mutants (p < 1e-

11 and p < 1e-12, respectively, log rank test) (Figure 1E).

Telomeric sRNAs and loss of EXO-1 repress sterility in the absence of telomerase

We considered simultaneously targeting the 11 unique subtelomeres of the *C. elegans*

genome with exogenous subtelomeric sRNAs in an effort to show that this class of

sRNAs promotes telomere stability in the absence of telomerase, but felt that this might

be difficult. Therefore, we decided to test the hypothesis that telomere repeat sRNAs

can promote telomere stability in the absence of telomerase. We asked if introduction

of exogenous telomeric sRNAs could rescue the rapid sterility of sRNA-deficient *trt-1*

mutants using a viable *dcr-1* allele, *mg375*, which is deficient for endogenous sRNA

production but is hypersensitive to exogenous RNAi when fed *E. coli* expressing dsRNA

(Welker et al., 2010). *trt-1; dcr-1(mg375)* double mutant strains that were passaged on

bacteria expressing telomeric dsRNA remained fertile significantly longer than those

expressing dsRNA from an empty vector control (Figure 4A). In contrast, exogenous

telomeric dsRNA did not affect the onset of sterility for *trt-1* single mutant controls.

Moreover, *trt-1; dcr-1(mg375)* strains fed telomeric dsRNA had markedly reduced levels

of DNA damage at F6 in comparison to *trt-1; dcr-1(mg375)* strains grown on control

RNAi or on regular OP50 bacteria that do not express dsRNA (Figure 1F, 5C).

In budding yeast, artificially-induced TERRA overexpression promotes telomere

shortening *in cis* by forming RNA:DNA hybrids that stall telomeric replication forks in a

manner that stimulates telomere resection by the exonuclease Exo1 (Maicher et al.,

2012, Pfeiffer and Lingner, 2012). We therefore hypothesized that loss of EXO-1 might

rescue the accelerated sterility phenotype in sRNA-deficient telomerase mutants. *trt-1;*

exo-1; mut-14 triple mutant lines survived significantly longer than *trt-1; mut-14* double

mutants (Figure 5B) and displayed reduced levels of DNA damage and moderate levels

of TERRA at generation F6 (Figures 5B, S5). Moreover, *trt-1; exo-1* double mutant

controls remained fertile even longer than *trt-1* single mutants (Figure 5B), indicating

that EXO-1-mediated telomere resection promotes telomere-shortening-induced

senescence in the absence of telomerase, both at normal telomeres and at telomeres

lacking sRNA-mediated genomic silencing. While *trt-1; exo-1* double mutant controls

remained fertile longer than *trt-1* single mutants, this was not true for *trt-1* single

mutants grown on exogenous telomeric dsRNA (Figure 5A, B). Thus, *exo-1* has at least

two roles at telomeres, one role that promotes sterility when telomerase is absent and a

second that promotes sterility when telomerase and sRNA silencing is absent.

Discussion

Small RNAs promote genomic silencing of many types of transposons (Holoch and

Moazed, 2015), including those found at *Drosophila melanogaster* telomeres, which are

composed of non-LTR retrotransposons that are distinct from the simple repetitive

sequences that are maintained by telomerase and cap chromosome ends in most

organisms (Pardue and DeBaryshe, 2003). Components of the RNAi machinery

function at *Drosophila* telomeres to repress expression of HeT-A and TART

retrotransposons and their accumulation at chromosome termini (Pane et al., 2007,

Savitsky et al., 2006), even though they do not affect telomere capping (Perrini et al.,

2004). In contrast, loss of the *Drosophila* homologs of the heterochromatin-mediated

genome silencing proteins HP1 and ATRX causes immediate telomere deprotection and

fusion (Chavez et al., 2017, Fanti et al., 1998, Fulcher et al., 2014). H3K9me2 but not

H3K9me3 is present at wild-type *C. elegans* telomeres (Figure 3A-C) (McMurchy et al.,

2017, Zeller et al., 2016), and we found that the *C. elegans* HP1 homolog HPL-2 that

binds to and specifically stabilizes H3K9me2 (McMurchy et al., 2017) promotes

telomere stability in the absence of telomerase (Figure1C). We also identified factors

that create rare small RNAs that may guide the deposition of histone silencing marks in

response to damaged telomeres (Figure 6A). Therefore, small RNAs and

heterochromatin proteins represent conserved features of both non-canonical and

canonical telomeres, even though their functions are distinct.

Loss of telomerase and sRNA factors in *C. elegans* caused high levels of DNA damage

response signaling in germ cells, but did not result in telomere fusions for a number of

generations, each generation representing 8-10 cell divisions. Therefore, the

accelerated sterility phenotype of sRNA-deficient *trt-1* mutants is not a consequence of

a defect in acute telomere uncapping, but rather an acceleration of the onset of crisis.

Biochemical analysis of telomere erosion in sRNA-deficient *trt-1* mutants revealed

increased variance in telomere size, which could be due to an error-prone

recombinational response to a form of DNA damage such as a stalled replication fork

(Figure 6C-D). Although both elongation and shortening of telomeres were observed

(Figure S1,S2), we propose that rapid telomere shortening events are likely responsible

for the premature end-to-end chromosome fusion and sterility phenotypes of sRNA-

deficient *trt-1* mutants. In telomerase single mutants that are proficient for sRNA

biogenesis and heterochromatin deposition, telomeric DNA damage may be healed in a

manner that more accurately maintains telomere length and is coupled to expression

and subsequent sRNA-mediated silencing of TERRA (Figure 6A-B).

Heterochromatin is known to suppress replication-based DNA damage at repetitive

segments of the genome (Zeller et al., 2016, Janssen et al., 2018). However, repeated

crossing of a *trt-1* mutation into a *dcr-1* mutant background failed to exacerbate the

accelerated sterility of *trt-1; dcr-1* double mutants (Figure 1F), indicating that *dcr-*

1(mg375) mutants have telomeres of normal length. Therefore, telomerase-mediated

telomere repeat addition is likely to be an effective method of healing telomeric DNA

damage in the absence of sRNA-mediated heterochromatin deposition.

Loss of telomeric sRNA biogenesis in telomerase mutants resulted in TERRA

expression, which could create R-loops that induce homologous recombination in a

manner that creates telomere shortening events that are toxic (Figure 6C). As

exogenous telomeric sRNAs were sufficient to suppress telomere instability, DNA

damage signaling and TERRA expression in the absence of telomerase and

endogenous sRNAs (Figure 5), silencing of TERRA could be an integral part of a

telomeric DNA damage response (Figure 6B,D) (Toubiana and Selig, 2018). These

results are consistent with a role for TERRA expression in response to a distinct form of

telomeric damage that occurs in the presence of telomerase – acutely deprotected, non-

replicating telomeres – and is accompanied by production of telomeric sRNAs that

promote DNA damage response factor recruitment (Rossiello et al., 2017). In addition,

work in *S. cerevisiae* showed that increased TERRA expression and persistent

telomeric R-loops in telomerase mutants can repress senescence by promoting

recombination, indicating that increased levels of TERRA are coupled to induction of a

telomeric DNA damage response (Graf et al., 2017). Finally, elevated levels of TERRA

have been shown to contribute to senescence in *S. cerevisiae* (Wanat et al., 2018).

We observed reduced levels of TERRA and DNA damage signaling in sRNA-deficient

telomerase mutants that lack *exo-1*. Consistently, loss of EXO1 in telomerase-negative

mice has been shown to extend lifespan and improve organ maintenance by repressing

telomeric ssDNA formation and DNA damage response (Schaetzlein et al., 2007),

suggesting a conserved role for EXO1 in the metazoan response to dysfunctional

telomeres.   Although expression of TERRA at mammalian telomeres induces EXO1-

dependent telomere shortening in a manner that independent of RNase H (Pfeiffer and

Lingner, 2012), we did not test a role for RNase H in the telomere instability phenotype

of sRNA-deficient *C. elegans* telomerase mutants.

We previously observed that *C. elegans* telomeric sRNAs with mismatches can be

mapped to Interstitial Telomere Sequences, which are degenerate telomere tracts that

are abundant on chromosomes arms and are distinct from telomeres (Frenk et al.,

2019). Interstitial Telomere Sequences were often heterochromatic and were markedly

depleted from exons and 5' or 3' UTRs (Frenk et al., 2019). Telomeric sRNAs with

mismatches were not altered by the *dcr-1(mg375)* mutation that causes telomere

instability in the absence of telomerase (Figure 4A). Although telomeric sRNAs with

mismatches contained a mixture of all four 5' nucleotides, very rare sRNAs composed

of perfect telomere repeats were strongly depleted for 5' G nucleotides (Frenk et al.,

2019). The unique structural property suggests a mechanism of biogenesis that is

distinct from known classes of *C. elegans* sRNAs (Billi et al., 2014). We found that

perfect telomeric sRNAs from the environment were sufficient to promote telomere

stability in *trt-1; dcr-1* mutants (Figure 5A). This suggests that very rare perfect

telomeric small RNAs, which might appear fleeting or artefactual based on analysis of

RNA sequencing data sets, promote telomere stability in the absence of telomerase.

Similar classes of rare small RNAs, which may be difficult (Frenk et al., 2019) or

impossible (Figure 4A) to measure, could be relevant to other segments of the genome.

As both Dicer and the dsRNA-binding protein RDE-4 promoted telomere stability in the

absence of telomerase (Figure 1B), we propose that biogenesis of telomeric small

RNAs could involve bidirectional transcripts that emanate from ITS tracts at internal

segments of the genome to create dsRNAs that contain both perfect and imperfect

telomere repeats (Frenk et al., 2019). ITS dsRNAs could interact with RDE-4 and Dicer

to create primary and secondary sRNA populations that could interact with TERRA

(Figure 6A). We detected subtelomeric sRNAs that were dependent on Dicer and that

interacted with the Argonaute proteins WAGO-1 and HRDE-1, both of which promoted

telomere stability in the absence of telomerase. These Argonaute proteins are known to

act in germ cells to target overlapping groups of transcripts, including genes,

transposons, and pseudogenes (Gu et al., 2009). Although WAGO-1 localizes to

perinuclear P-granules (germ granules), where it might promote either transcriptional or

post-transcriptional gene silencing (Gu et al., 2009), HRDE-1 localizes to the nucleus

where it promotes transcriptional gene silencing. Moreover, HRDE-1 and WAGO-1

have been shown to function redundantly to promote a form of permanent

transgenerational transcriptional silencing (Ashe et al., 2012, Buckley et al., 2012).

Although proteomics has been performed for both HRDE-1 and WAGO-1, they do not

physically interact with one another or possess common interacting proteins (Akay et

al., 2017, Shirayama et al., 2014).

We were unable to measure perfect telomeric sRNAs by deeply sequencing small

RNAs from several genetic backgrounds (Figure 4A), but we observed upregulation of

sRNAs at some subtelomeres in telomerase mutants, and in some cases this increase

was DCR-1-dependent (Figure 4). We also found that HRDE-1 and WAGO-1 interact

with 22G sRNAs that map to distinct and overlapping subtelomeres. Given that DNA

damage responses induce histone modifications that can be propagated for long

distances from sites of DNA damage (Polo and Jackson, 2011), we propose that the

biological response to telomeric DNA damage initially may spread such that abundant

sRNAs are created from RNA transcripts at some subtelomeric loci. Therefore, the

variable sRNA biogenesis patterns observed at distinct subtelomeres of telomerase

mutants may reflect a wider epigenetic consequence of the biological response to

telomeric DNA damage.

In mammals, TERRA facilitates transcriptional regulation of the epigenome (Chu et al.,

2017). Moreover, both TERRA and telomerase have been implicated in regulation of

genes that promote stem cell proliferation (Marion et al., 2019, Sarin et al., 2005).

Therefore, the interplay of telomerase and TERRA in the context of deficiency for small

RNA-mediated genome silencing could reflect more general roles for these factors in

epigenetic regulation of cell fate.

sRNA-mediated telomeric silencing was promoted by the *C. elegans* secondary sRNA

biogenesis factor MUT-7 (Figure 1A, B), which is homologous to the exonuclease

domain of human WRN1 Werner's syndrome protein whose loss results in early-onset

aging of multiple tissues (Ketting et al., 2001). WRN1 deficiency is accompanied by

accelerated telomere erosion and senescence phenotypes that can be rescued by

telomerase expression (Shimamoto et al., 2015). While WRN1 and its homologs have

been intensively studied for their roles in DNA recombination, WRN1 safeguards the

stability of heterochromatin (Zhang et al., 2015), which our study suggests may play a

critical role in maintaining telomere stability in most human somatic cells. Consistently,

deficiency for DNA methylation at repetitive segments of the genome, including

telomeres, occurs in ICF syndrome cells that undergo rapid telomere shortening and

senescence accompanied by increased TERRA expression, telomeric R-loops and DNA

damage (Sagie et al., 2017). As global dysfunction of the epigenome is a hallmark of

human aging (Zhang et al., 2015), our findings may be generally relevant to

understanding how sRNA-driven chromatin states established in germ cells or the early

embryo might contribute to telomere instability during aging of adult somatic cells.

Data availability statement

RNA-seq data reported in this study are available under the accession number GEO:

GSE111800.

Acknowledgements

We thank R. Dowen for technical advice with qRT-PCR, K. Billmyre for advice on

immunofluorescence, and members of the S.A. laboratory for critical review of the

manuscript. Some strains were provided by the *Caenorhabditis* Genetics Center, which

is funded by National Institutes of Health Office of Research Infrastructure Programs

(P40 OD010440).

Funding

S.F., C.L. and S.A. were supported by National Institute of Health grant R01 GM135470

and R01 ES035777.

Conflict of Interest

The authors declare no competing interests.

References

AHMED, S. & HODGKIN, J. 2000. MRT-2 checkpoint protein is required for germline immortality and telomere replication in *C. elegans*. *Nature,* 403**,** 159-64.

AKAY, A., DI DOMENICO, T., SUEN, K. M., NABIH, A., PARADA, G. E., LARANCE, M., MEDHI, R., BERKYUREK, A. C., ZHANG, X., WEDELES, C. J., RUDOLPH, K. L. M., ENGELHARDT, J., HEMBERG, M., MA, P., LAMOND, A. I., CLAYCOMB, J. M. & MISKA, E. A. 2017. The Helicase Aquarius/EMB-4 Is Required to Overcome Intronic Barriers to Allow Nuclear RNAi Pathways to Heritably Silence Transcription. *Dev Cell,* 42**,** 241-255 e6.

ARNOULT, N., VAN BENEDEN, A. & DECOTTIGNIES, A. 2012. Telomere length regulates TERRA levels through increased trimethylation of telomeric H3K9 and HP1alpha. *Nat Struct Mol Biol,* 19**,** 948-56.

ASHE, A., SAPETSCHNIG, A., WEICK, E. M., MITCHELL, J., BAGIJN, M. P., CORDING, A. C., DOEBLEY, A. L., GOLDSTEIN, L. D., LEHRBACH, N. J., LE PEN, J., PINTACUDA, G., SAKAGUCHI, A., SARKIES, P., AHMED, S. & MISKA, E. A. 2012. piRNAs Can Trigger a Multigenerational Epigenetic Memory in the Germline of C. elegans. *Cell,* 150**,** 88-99.

AVGOUSTI, D. C., PALANI, S., SHERMAN, Y. & GRISHOK, A. 2012. CSR-1 RNAi pathway positively regulates histone expression in C. elegans. *EMBO J,* 31**,** 3821-32.

AZZALIN, C. M. & LINGNER, J. 2015. Telomere functions grounding on TERRA firma. *Trends Cell Biol,* 25**,** 29-36.

AZZALIN, C. M., REICHENBACH, P., KHORIAULI, L., GIULOTTO, E. & LINGNER, J. 2007. Telomeric repeat containing RNA and RNA surveillance factors at mammalian chromosome ends. *Science,* 318**,** 798-801.

BAGIJN, M. P., GOLDSTEIN, L. D., SAPETSCHNIG, A., WEICK, E. M., BOUASKER, S., LEHRBACH, N. J., SIMARD, M. J. & MISKA, E. A. 2012. Function, targets, and evolution of Caenorhabditis elegans piRNAs. *Science,* 337**,** 574-8.

BATISTA, P. J., RUBY, J. G., CLAYCOMB, J. M., CHIANG, R., FAHLGREN, N., KASSCHAU, K. D., CHAVES, D. A., GU, W., VASALE, J. J., DUAN, S., CONTE, D., JR., LUO, S., SCHROTH, G. P., CARRINGTON, J. C., BARTEL, D. P. & MELLO, C. C. 2008. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in C. elegans. *Mol Cell,* 31**,** 67-78.

BENETTI, R., GARCIA-CAO, M. & BLASCO, M. A. 2007. Telomere length regulates the epigenetic status of mammalian telomeres and subtelomeres. *Nat Genet,* 39**,** 243-50.

BENETTI, R., SCHOEFTNER, S., MUNOZ, P. & BLASCO, M. A. 2008. Role of TRF2 in the assembly of telomeric chromatin. *Cell Cycle,* 7**,** 3461-8.

BERNSTEIN, E., CAUDY, A. A., HAMMOND, S. M. & HANNON, G. J. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature,* 409**,** 363-6.

BILLI, A. C., FISCHER, S. E. & KIM, J. K. 2014. Endogenous RNAi pathways in C. elegans. *WormBook***,** 1-49.

BOERCKEL, J., WALKER, D. & AHMED, S. 2007. The Caenorhabditis elegans Rad17 homolog HPR-17 is required for telomere replication. *Genetics,* 176**,** 703-9.

BUCKLEY, B. A., BURKHART, K. B., GU, S. G., SPRACKLIN, G., KERSHNER, A., FRITZ, H., KIMBLE, J., FIRE, A. & KENNEDY, S. 2012. A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature,* 489**,** 447-51.

BUSHNELL, B. 2016. BBMap short-read aligner, and other bioinformatics tools. http://sourceforge.net/projects/bbmap/.

CANZIO, D., LARSON, A. & NARLIKAR, G. J. 2014. Mechanisms of functional promiscuity by HP1 proteins. *Trends Cell Biol,* 24**,** 377-86.

CAO, F., LI, X., HIEW, S., BRADY, H., LIU, Y. & DOU, Y. 2009. Dicer independent small RNAs associate with telomeric heterochromatin. *RNA,* 15**,** 1274-81.

CHAVEZ, J., MURILLO-MALDONADO, J. M., BAHENA, V., CRUZ, A. K., CASTANEDA-SORTIBRAN, A., RODRIGUEZ-ARNAIZ, R., ZURITA, M. & VALADEZ-GRAHAM, V. 2017. dAdd1 and dXNP prevent genome instability by maintaining HP1a localization at Drosophila telomeres. *Chromosoma,* 126**,** 697-712.

CHENG, C., SHTESSEL, L., BRADY, M. M. & AHMED, S. 2012. Caenorhabditis elegans POT-2 telomere protein represses a mode of alternative lengthening of telomeres with normal telomere lengths. *Proc Natl Acad Sci U S A,* 109**,** 7805-10.

CHU, H. P., CIFUENTES-ROJAS, C., KESNER, B., AEBY, E., LEE, H. G., WEI, C., OH, H. J., BOUKHALI, M., HAAS, W. & LEE, J. T. 2017. TERRA RNA Antagonizes ATRX and Protects Telomeres. *Cell,* 170**,** 86-101 e16.

CLAYCOMB, J.M., BATISTA, P.J., PANG, K.M., GU, W., VASALE, J.J., VAN WOLFSWINKEL, J.C., CHAVES, D.A., SHIRAYAMA, M., MITANI, S., KETTING, R.F., CONTE D., MELLO C.C. (2009). The Argonaute CSR-1 and its 22G-RNA cofactors are required for holocentric chromosome segregation. *Cell* 139, 123-134.

COUVILLION, M. T., LEE, S. R., HOGSTAD, B., MALONE, C. D., TONKIN, L. A., SACHIDANANDAM, R., HANNON, G. J. & COLLINS, K. 2009. Sequence, biogenesis, and function of diverse small RNA classes bound to the Piwi family proteins of Tetrahymena thermophila. *Genes Dev,* 23**,** 2016-32.

CUSANELLI, E., ROMERO, C. A. & CHARTRAND, P. 2013. Telomeric noncoding RNA TERRA is induced by telomere shortening to nucleate telomerase molecules at short telomeres. *Mol Cell,* 51**,** 780-91.

DAS, P. P., BAGIJN, M. P., GOLDSTEIN, L. D., WOOLFORD, J. R., LEHRBACH, N. J., SAPETSCHNIG, A., BUHECHA, H. R., GILCHRIST, M. J., HOWE, K. L., STARK, R., MATTHEWS, N., BEREZIKOV, E., KETTING, R. F., TAVARE, S. & MISKA, E. A. 2008. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the Caenorhabditis elegans germline. *Mol Cell,* 31**,** 79-90.

DONATE, L. E. & BLASCO, M. A. 2011. Telomeres in cancer and ageing. *Philos Trans R Soc Lond B Biol Sci,* 366**,** 76-84.

FANTI, L., GIOVINAZZO, G., BERLOCO, M. & PIMPINELLI, S. 1998. The heterochromatin protein 1 prevents telomere fusions in Drosophila. *Mol Cell,* 2**,** 527-38.

FLYNN, R. L., COX, K. E., JEITANY, M., WAKIMOTO, H., BRYLL, A. R., GANEM, N. J., BERSANI, F., PINEDA, J. R., SUVA, M. L., BENES, C. H., HABER, D. A.,

BOUSSIN, F. D. & ZOU, L. 2015. Alternative lengthening of telomeres renders
cancer cells hypersensitive to ATR inhibitors. *Science,* 347**,** 273-7.
FRENK, S., LISTER-SHIMAUCHI, E. H. & AHMED, S. 2019. Telomeric small RNAs in
the genus Caenorhabditis. *RNA*.
FULCHER, N., DERBOVEN, E., VALUCHOVA, S. & RIHA, K. 2014. If the cap fits, wear
it: an overview of telomeric structures over evolution. *Cell Mol Life Sci,* 71**,** 847-
65.
GENT, J. I., LAMM, A. T., PAVELEC, D. M., MANIAR, J. M., PARAMESWARAN, P.,
TAO, L., KENNEDY, S. & FIRE, A. Z. 2010. Distinct phases of siRNA synthesis
in an endogenous RNAi pathway in C. elegans soma. *Mol Cell,* 37**,** 679-89.
GOU, L. T., DAI, P., YANG, J. H., XUE, Y., HU, Y. P., ZHOU, Y., KANG, J. Y., WANG,
X., LI, H., HUA, M. M., ZHAO, S., HU, S. D., WU, L. G., SHI, H. J., LI, Y., FU, X.
D., QU, L. H., WANG, E. D. & LIU, M. F. 2014. Pachytene piRNAs instruct
massive mRNA elimination during late spermiogenesis. *Cell Res,* 24**,** 680-700.
GRAF, M., BONETTI, D., LOCKHART, A., SERHAL, K., KELLNER, V., MAICHER, A.,
JOLIVET, P., TEIXEIRA, M. T. & LUKE, B. 2017. Telomere Length Determines
TERRA and R-Loop Regulation through the Cell Cycle. *Cell,* 170**,** 72-85 e14.
GREER, E. L., MAURES, T. J., HAUSWIRTH, A. G., GREEN, E. M., LEEMAN, D. S.,
MARO, G. S., HAN, S., BANKO, M. R., GOZANI, O. & BRUNET, A. 2010.
Members of the H3K4 trimethylation complex regulate lifespan in a germline-
dependent manner in C. elegans. *Nature,* 466**,** 383-7.
GU, W., SHIRAYAMA, M., CONTE, D., JR., VASALE, J., BATISTA, P. J., CLAYCOMB,
J. M., MORESCO, J. J., YOUNGMAN, E. M., KEYS, J., STOLTZ, M. J., CHEN,
C. C., CHAVES, D. A., DUAN, S., KASSCHAU, K. D., FAHLGREN, N., YATES,
J. R., 3RD, MITANI, S., CARRINGTON, J. C. & MELLO, C. C. 2009. Distinct
argonaute-mediated 22G-RNA pathways direct genome surveillance in the C.
elegans germline. *Mol Cell,* 36**,** 231-44.
HAHNE, F. & IVANEK, R. 2016. Visualizing Genomic Data Using Gviz and
Bioconductor. *Methods Mol Biol,* 1418**,** 335-51.
HAN, T., MANOHARAN, A. P., HARKINS, T. T., BOUFFARD, P., FITZPATRICK, C.,
CHU, D. S., THIERRY-MIEG, D., THIERRY-MIEG, J. & KIM, J. K. 2009. 26G
endo-siRNAs regulate spermatogenic and zygotic gene expression in
Caenorhabditis elegans. *Proc Natl Acad Sci U S A,* 106**,** 18674-9.
HEARD, E. & TURNER, J. 2011. Function of the sex chromosomes in mammalian
fertility. *Cold Spring Harb Perspect Biol,* 3**,** a002675.
HOLOCH, D. & MOAZED, D. 2015. RNA-mediated epigenetic regulation of gene
expression. *Nat Rev Genet,* 16**,** 71-84.
HOLOHAN, B., WRIGHT, W. E. & SHAY, J. W. 2014. Cell biology of disease:
Telomeropathies: an emerging spectrum disorder. *J Cell Biol,* 205**,** 289-99.
JANSSEN, A., COLMENARES, S. U. & KARPEN, G. H. 2018. Heterochromatin:
Guardian of the Genome. *Annu Rev Cell Dev Biol,* 34**,** 265-288.
KASPER, D. M., GARDNER, K. E. & REINKE, V. 2014. Homeland security in the C.
elegans germ line: insights into the biogenesis and function of piRNAs.
Epigenetics, 9**,** 62-74.
KETTING, R. F., FISCHER, S. E., BERNSTEIN, E., SIJEN, T., HANNON, G. J. &
PLASTERK, R. H. 2001. Dicer functions in RNA interference and in synthesis of

small RNA involved in developmental timing in C. elegans. *Genes Dev,* 15**,** 2654-
9.
KETTING, R. F., HAVERKAMP, T. H., VAN LUENEN, H. G. & PLASTERK, R. H. 1999.
Mut-7 of C. elegans, required for transposon silencing and RNA interference, is a
homolog of Werner syndrome helicase and RNaseD. *Cell,* 99**,** 133-41.
KNIGHT, S. W. & BASS, B. L. 2001. A role for the RNase III enzyme DCR-1 in RNA
interference and germ line development in Caenorhabditis elegans. *Science,*
293, 2269-71.
LANGMEAD, B., TRAPNELL, C., POP, M. & SALZBERG, S. L. 2009. Ultrafast and
memory-efficient alignment of short DNA sequences to the human genome.
Genome Biol, 10**,** R25.
LI, F., HUARTE, M., ZARATIEGUI, M., VAUGHN, M. W., SHI, Y., MARTIENSSEN, R. &
CANDE, W. Z. 2008. Lid2 is required for coordinating H3K4 and H3K9
methylation of heterochromatin and euchromatin. *Cell,* 135**,** 272-83.
LI, J. S., MIRALLES FUSTE, J., SIMAVORIAN, T., BARTOCCI, C., TSAI, J.,
KARLSEDER, J. & LAZZERINI DENCHI, E. 2017. TZAP: A telomere-associated
protein involved in telomere length control. *Science,* 355**,** 638-641.
LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
LOVEJOY, C. A., LI, W., REISENWEBER, S., THONGTHIP, S., BRUNO, J., DE
LANGE, T., DE, S., PETRINI, J. H., SUNG, P. A., JASIN, M., ROSENBLUH, J.,
ZWANG, Y., WEIR, B. A., HATTON, C., IVANOVA, E., MACCONAILL, L.,
HANNA, M., HAHN, W. C., LUE, N. F., REDDEL, R. R., JIAO, Y., KINZLER, K.,
VOGELSTEIN, B., PAPADOPOULOS, N., MEEKER, A. K. & CONSORTIUM, A.
L. T. S. C. 2012. Loss of ATRX, genome instability, and an altered DNA damage
response are hallmarks of the alternative lengthening of telomeres pathway.
PLoS Genet, 8**,** e1002772.
MACIEJOWSKI, J. & DE LANGE, T. 2017. Telomeres in cancer: tumour suppression
and genome instability. *Nat Rev Mol Cell Biol,* 18**,** 175-186.
MAICHER, A., KASTNER, L., DEES, M. & LUKE, B. 2012. Deregulated telomere
transcription causes replication-dependent telomere shortening and promotes
cellular senescence. *Nucleic Acids Res,* 40**,** 6649-59.
MARION, R. M., MONTERO, J. J., LOPEZ DE SILANES, I., GRANA-CASTRO, O.,
MARTINEZ, P., SCHOEFTNER, S., PALACIOS-FABREGA, J. A. & BLASCO, M.
A. 2019. TERRA regulate the transcriptional landscape of pluripotent cells
through TRF1-dependent recruitment of PRC2. *Elife,* 8.
MCMURCHY, A. N., STEMPOR, P., GAARENSTROOM, T., WYSOLMERSKI, B.,
DONG, Y., AUSSIANIKAVA, D., APPERT, A., HUANG, N., KOLASINSKA-
ZWIERZ, P., SAPETSCHNIG, A., MISKA, E. A. & AHRINGER, J. 2017. A team
of heterochromatin factors collaborates with small RNA pathways to combat
repetitive elements and germline stress. *Elife,* 6.
MEIER, B., BARBER, L. J., LIU, Y., SHTESSEL, L., BOULTON, S. J., GARTNER, A. &
AHMED, S. 2009. The MRT-1 nuclease is required for DNA crosslink repair and
telomerase activity in vivo in Caenorhabditis elegans. *Embo J,* 28**,** 3549-63.

MEIER, B., CLEJAN, I., LIU, Y., LOWDEN, M., GARTNER, A., HODGKIN, J. &
AHMED, S. 2006. trt-1 is the Caenorhabditis elegans catalytic subunit of
telomerase. *PLoS Genet,* 2**,** e18.
MONTERO, J. J., LOPEZ-SILANES, I., MEGIAS, D, M, F. F., CASTELLS-GARCIA, A.
& BLASCO, M. A. 2018. TERRA recruitment of polycomb to telomeres is
essential for histone trymethylation marks at telomeric heterochromatin. *Nat*
Commun, 9**,** 1548.
MORAVEC, M., WISCHNEWSKI, H., BAH, A., HU, Y., LIU, N., LAFRANCHI, L., KING,
M. C. & AZZALIN, C. M. 2016. TERRA promotes telomerase-mediated telomere
elongation in Schizosaccharomyces pombe. *EMBO Rep,* 17**,** 999-1012.
ORTIZ, M. A., NOBLE, D., SOROKIN, E. P. & KIMBLE, J. 2014. A new dataset of
spermatogenic vs. oogenic transcriptomes in the nematode Caenorhabditis
elegans. *G3 (Bethesda),* 4**,** 1765-72.
PAK, J. & FIRE, A. 2007. Distinct populations of primary and secondary effectors during
RNAi in C. elegans. *Science,* 315**,** 241-4.
PANE, A., WEHR, K. & SCHUPBACH, T. 2007. zucchini and squash encode two
putative nucleases required for rasiRNA production in the Drosophila germline.
Dev Cell, 12**,** 851-62.
PARDUE, M. L. & DEBARYSHE, P. G. 2003. Retrotransposons provide an
evolutionarily robust non-telomerase mechanism to maintain telomeres. *Annu*
Rev Genet, 37**,** 485-511.
PERRINI, B., PIACENTINI, L., FANTI, L., ALTIERI, F., CHICHIARELLI, S., BERLOCO,
M., TURANO, C., FERRARO, A. & PIMPINELLI, S. 2004. HP1 controls telomere
capping, telomere elongation, and telomere silencing by two different
mechanisms in Drosophila. *Mol Cell,* 15**,** 467-76.
PFEIFFER, V. & LINGNER, J. 2012. TERRA promotes telomere shortening through
exonuclease 1-mediated resection of chromosome ends. *PLoS Genet,* 8**,**
e1002747.
PHILLIPS, C. M., MCDONALD, K. L. & DERNBURG, A. F. 2009. Cytological analysis of
meiosis in Caenorhabditis elegans. *Methods Mol Biol,* 558**,** 171-95.
PICKETT, H. A. & REDDEL, R. R. 2015. Molecular mechanisms of activity and
derepression of alternative lengthening of telomeres. *Nat Struct Mol Biol,* 22**,**
875-80.
POLO, S. E. & JACKSON, S. P. 2011. Dynamics of DNA damage response proteins at
DNA breaks: a focus on protein modifications. *Genes Dev,* 25**,** 409-33.
PORRO, A., FEUERHAHN, S., DELAFONTAINE, J., RIETHMAN, H., ROUGEMONT, J.
& LINGNER, J. 2014. Functional characterization of the TERRA transcriptome at
damaged telomeres. *Nat Commun,* 5**,** 5379.
QUINLAN, A. R. & HALL, I. M. 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics,* 26**,** 841-2.
REINHART, B. J. & BARTEL, D. P. 2002. Small RNAs correspond to centromere
heterochromatic repeats. *Science,* 297**,** 1831.
ROAKE, C. M. & ARTANDI, S. E. 2017. Approaching TERRA Firma: Genomic
Functions of Telomeric Noncoding RNA. *Cell,* 170**,** 8-9.
ROSSIELLO, F., AGUADO, J., SEPE, S., IANNELLI, F., NGUYEN, Q., PITCHIAYA, S.,
CARNINCI, P. & D'ADDA DI FAGAGNA, F. 2017. DNA damage response

inhibition at dysfunctional telomeres by modulation of telomeric DNA damage
response RNAs. *Nat Commun,* 8**,** 13980.
SAGIE, S., TOUBIANA, S., HARTONO, S. R., KATZIR, H., TZUR-GILAT, A.,
HAVAZELET, S., FRANCASTEL, C., VELASCO, G., CHEDIN, F. & SELIG, S.
2017. Telomeres in ICF syndrome cells are vulnerable to DNA damage due to
elevated DNA:RNA hybrids. *Nat Commun,* 8**,** 14015.
SAKAGUCHI, A., SARKIES, P., SIMON, M., DOEBLEY, A. L., GOLDSTEIN, L. D.,
HEDGES, A., IKEGAMI, K., ALVARES, S. M., YANG, L., LAROCQUE, J. R.,
HALL, J., MISKA, E. A. & AHMED, S. 2014. Caenorhabditis elegans RSD-2 and
RSD-6 promote germ cell immortality by maintaining small interfering RNA
populations. *Proc Natl Acad Sci U S A,* 111**,** E4323-31.
SARIN, K. Y., CHEUNG, P., GILISON, D., LEE, E., TENNEN, R. I., WANG, E.,
ARTANDI, M. K., ORO, A. E. & ARTANDI, S. E. 2005. Conditional telomerase
induction causes proliferation of hair follicle stem cells. *Nature,* 436**,** 1048-52.
SAVITSKY, M., KWON, D., GEORGIEV, P., KALMYKOVA, A. & GVOZDEV, V. 2006.
Telomere elongation is under the control of the RNAi-based mechanism in the
Drosophila germline. *Genes Dev,* 20**,** 345-54.
SCHAETZLEIN, S., KODANDARAMIREDDY, N. R., JU, Z., LECHEL, A.,
STEPCZYNSKA, A., LILLI, D. R., CLARK, A. B., RUDOLPH, C., KUHNEL, F.,
WEI, K., SCHLEGELBERGER, B., SCHIRMACHER, P., KUNKEL, T. A.,
GREENBERG, R. A., EDELMANN, W. & RUDOLPH, K. L. 2007. Exonuclease-1
deletion impairs DNA damage signaling and prolongs lifespan of telomere-
dysfunctional mice. *Cell,* 130**,** 863-77.
SHAY, J. W. 2016. Role of Telomeres and Telomerase in Aging and Cancer. *Cancer*
Discov, 6**,** 584-93.
SHIMAMOTO, A., YOKOTE, K. & TAHARA, H. 2015. Werner Syndrome-specific
induced pluripotent stem cells: recovery of telomere function by reprogramming.
Front Genet, 6**,** 10.
SHIRAYAMA, M., STANNEY, W., 3RD, GU, W., SETH, M. & MELLO, C. C. 2014. The
Vasa Homolog RDE-12 engages target mRNA and multiple argonaute proteins to
promote RNAi in C. elegans. *Curr Biol,* 24**,** 845-51.
STEINER, F. A., OKIHARA, K. L., HOOGSTRATE, S. W., SIJEN, T. & KETTING, R. F.
2009. RDE-1 slicer activity is required only for passenger-strand cleavage during
RNAi in Caenorhabditis elegans. *Nat Struct Mol Biol,* 16**,** 207-11.
TABARA, H., YIGIT, E., SIOMI, H. & MELLO, C. C. 2002. The dsRNA binding protein
RDE-4 interacts with RDE-1, DCR-1, and a DExH-box helicase to direct RNAi in
C. elegans. *Cell,* 109**,** 861-71.
TAYLOR, D. H., CHU, E. T., SPEKTOR, R. & SOLOWAY, P. D. 2015. Long non-coding
RNA regulation of reproduction and development. *Mol Reprod Dev,* 82**,** 932-56.
TEAM, R. C. 2017. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
TIJSTERMAN, M., KETTING, R. F., OKIHARA, K. L., SIJEN, T. & PLASTERK, R. H.
2002. RNA helicase MUT-14-dependent gene silencing triggered in C. elegans
by short antisense RNAs. *Science,* 295**,** 694-7.
TOUBIANA, S. & SELIG, S. 2018. DNA:RNA hybrids at telomeres - when it is better to
be out of the (R) loop. *FEBS J,* 285**,** 2552-2566.

UDUGAMA, M., FT, M. C., CHAN, F. L., TANG, M. C., PICKETT, H. A., JD, R. M.,
MAYNE, L., COLLAS, P., MANN, J. R. & WONG, L. H. 2015. Histone variant
H3.3 provides the heterochromatic H3 lysine 9 tri-methylation mark at telomeres.
Nucleic Acids Res, 43**,** 10227-37.
VRBSKY, J., AKIMCHEVA, S., WATSON, J. M., TURNER, T. L., DAXINGER, L.,
VYSKOT, B., AUFSATZ, W. & RIHA, K. 2010. siRNA-mediated methylation of
Arabidopsis telomeres. *PLoS Genet,* 6**,** e1000986.
WANAT, J. J., LOGSDON, G. A., DRISKILL, J. H., DENG, Z., LIEBERMAN, P. M. &
JOHNSON, F. B. 2018. TERRA and the histone methyltransferase Dot1
cooperate to regulate senescence in budding yeast. *PLoS One,* 13**,** e0195698.
WEDELES, C. J., WU, M. Z. & CLAYCOMB, J. M. 2013. Protection of germline gene
expression by the C. elegans Argonaute CSR-1. *Dev Cell,* 27**,** 664-71.
WELKER, N. C., PAVELEC, D. M., NIX, D. A., DUCHAINE, T. F., KENNEDY, S. &
BASS, B. L. 2010. Dicer's helicase domain is required for accumulation of some,
but not all, C. elegans endogenous siRNAs. *RNA,* 16**,** 893-903.
WICKY, C., VILLENEUVE, A. M., LAUPER, N., CODOUREY, L., TOBLER, H. &
MULLER, F. 1996. Telomeric repeats (TTAGGC)n are sufficient for chromosome
capping function in Caenorhabditis elegans. *Proc Natl Acad Sci U S A,* 93**,** 8983-
8.
WONG, L. H., MCGHIE, J. D., SIM, M., ANDERSON, M. A., AHN, S., HANNAN, R. D.,
GEORGE, A. J., MORGAN, K. A., MANN, J. R. & CHOO, K. H. 2010. ATRX
interacts with H3.3 in maintaining telomere structural integrity in pluripotent
embryonic stem cells. *Genome Res,* 20**,** 351-60.
ZELLER, P., PADEKEN, J., VAN SCHENDEL, R., KALCK, V., TIJSTERMAN, M. &
GASSER, S. M. 2016. Histone H3K9 methylation is dispensable for
Caenorhabditis elegans development but suppresses RNA:DNA hybrid-
associated repeat instability. *Nat Genet,* 48**,** 1385-1395.
ZHANG, W., LI, J., SUZUKI, K., QU, J., WANG, P., ZHOU, J., LIU, X., REN, R., XU, X.,
OCAMPO, A., YUAN, T., YANG, J., LI, Y., SHI, L., GUAN, D., PAN, H., DUAN,
S., DING, Z., LI, M., YI, F., BAI, R., WANG, Y., CHEN, C., YANG, F., LI, X.,
WANG, Z., AIZAWA, E., GOEBL, A., SOLIGALLA, R. D., REDDY, P., ESTEBAN,
C. R., TANG, F., LIU, G. H. & BELMONTE, J. C. 2015. Aging stem cells. A
Werner syndrome stem cell model unveils heterochromatin alterations as a driver
of human aging. *Science,* 348**,** 1160-3.

Figure 1: Endogenous small RNA and chromatin factors repress telomere stability in the absence of telomerase. (A) Survival curve of telomerase-deficient strains passaged at ~100 worms weekly for 40 independent lines. One allele of *xnp-1(ok1823)* extended survival but did not increase survival via ALT (Table S1). The same wild type control data is shown in panels B-F. (B) Survival curve of primary or secondary sRNA deficient telomerase mutants when 6 larval L1s are passaged weekly. (C) Survival curve of double mutants for *trt-1* and chromatin factors passaged by 6 L1s weekly. (D) Histone H3.3 remodeling protein XNP-1 represses senescence in *trt-1* mutants epistatically to the small RNA factor MUT-7. (E) Loss of HRDE-1 or WAGO-1 Argonaute proteins mimics the accelerated sterility observed in a *trt-1; dcr-1* mutant background. (F) A *trt-*

1; dcr-1 strain was crossed with *dcr-1* males fifteen times to yield a *trt-1; dcr-1* 15x strain
that displayed the same accelerated sterility observed for the initial *trt-1; dcr-1* double
mutants. (G) Small RNA and heterochromatin mutant *trt-1* lines have high levels of
chromosome fusions, comparable to *trt-1* single mutants at F24. Fusions were assayed
by diamidino-2-phenylindole (DAPI) staining of oocytes in diakinesis. Each bivalent in
wildtype represents a single paired chromosome. Representative images of DAPI-
stained oocyte nuclei are shown in panels below the graph. (H) High levels of phospho-
(Ser/Thr) staining in the meiotic gonad of *trt-1; dcr-1* at F6 compared to N2 or *trt-1*.
Scale bar, 25 m. (B)-(D), (F) ***p < 0.001; ****p < 0.00001; ns, not significant, (Log
rank test, n=40 independent lines for each genotype). (E) *trt-1; wago-1* = 39 lines per
genotype. *trt-1; hrde-1* and *trt-1; dcr-1* = 33 lines per genotype.
1045

Figure 2: (A) Expression of TERRA from subtelomeres *IIIL*, *VR*, and *XL* is increased in sRNA and heterochromatin deficient telomerase mutants. cDNA was reverse transcribed with a telomere specific primer and qPCR was performed using subtelomere-specific primer pairs and normalized to *act-1*. Expression data are plotted relative to wildtype and represent the mean ± SEM of three biological replicates. (B) RNA FISH signals of TERRA in meiotic germlines of hermaphrodites of the indicated genotypes. TERRA foci are present in wildtype and *trt-1* in the pachytene region and increased in *trt-1; dcr-1.* Scale bar, 10um. (C) Representative RNA FISH images of TERRA in males show foci in the pachytene region of the germline which are increased in *trt-1; dcr-1* compared to *trt-1.* Scale bar, 20 um. (D) Quantification of germline TERRA foci (mean ± SEM, n=10 germlines).

Figure 3: Analysis of telomeric chromatin and small RNAs in published datasets. (A)-(C) ChIP-seq enrichment profiles at three chromosome ends constructed using published data (McMurchy et al., 2017, Zeller et al., 2016). Small orange arrows between gene models and data tracks show the position of primers used in TERRA qRT-PCR experiments. (D) Enrichment (IP/input) of reads mapping to perfect telomere sequence with up to three mismatches in published immunoprecipitation datasets for three different Argonaute proteins. (E) Enrichment for subtelomeric sRNAs for datasets used in (D).

1071

1072

Figure 4: Upregulation of subtelomeric sRNAs in telomerase mutants. (A) Number of
telomere-mapping reads in wild-type, *trt-1,* and *trt-1;dcr-1* libraries. Numbers 1-3
indicate number of mismatches. (B)-(D) 22G sRNAs at three subtelomeres. (E) Log2-
fold change in sRNA abundance at subtelomeres relative to wildtype in *trt-1* and *trt-1;*
dcr-1 animals. (F) Proportion of subtelomeric sRNAs mapping to the strand
corresponding to the G-rich (TTAGGC) and C-rich (TAAGCC) telomere strand in *trt-1*
animals. **** $p < 0.0001$; ** $p < 0.01$; ns, not significant (Binomial test with Bonferroni
correction). The p values refer to the probability of observing the given proportion of G-rich
or C-rich telomeric reads at each chromosome end under the null hypothesis that G- and
C-rich reads are present in equal proportions.

1083

1084

1085
Figure 5: Fast senescence in sRNA-deficient *trt-1* is rescued by exogenous telomeric
dsRNA or loss of Exonuclease-1. (A) Survival of *trt-1* or *trt-1; dcr-1(mg375)* on either
empty vector control (EV) or bacteria expressing telomeric double-stranded RNA (telo
dsRNA). Log rank test relative to EV, ***P < 0.001. n=40 lines per genotype. (B) Loss
of *exo-1* extends survival of *trt-1; mut-14* lines. n=40 lines per genotype. (C) Telomeric
dsRNA and loss of *exo-1* suppress accumulation of phospho-(Ser/Thr) in sRNA-
deficient *trt-1* by F6. Scale bar=20um.
1093
1094
1095

Figure 6: Model for TERRA, small RNAs and telomerase. A) In *trt-1* telomerase
mutants, telomeric erosion induces expression of the lncRNA TERRA, which engages
production of sub-telomeric and telomeric sRNAs. Small RNAs promote deposition of
H3K9me2 and H3K4 demethylation in order to repress TERRA expression DNA
damage, and *trt-1* mutant sterility. B) TERRA expression and silencing may occur in the
context of a DNA damage response. C) In *trt-1; dcr-1* double mutants, TERRA
expression is not repressed, which may cause RNA:DNA hybrids and a form of
recombination that is dependent on the nuclease EXO-1 and leads to rapid sterility. D)
Lack of sRNAs may induce an aberrant recombinational response to DNA damage that
results in telomere elongation or shortening.

Supplemental Information
1112
1113 5 Supplemental Figures
1114
1115 3 Supplemental Tables
1116
1117

1118

Figure S1. Related to Figure 1. Terminal Restriction Fragment (TFR) analysis of
single-mutant *trt-1*, *mut-7*, and *dcr-1* and double-mutant *trt-1; mut-7* and *trt-1; dcr-1*
strains using a telomeric probe. (A) Southern blot of samples generated by passaging
small groups of worms for multiple generations. Two independent *trt-1* lines were used.
Long telomeres observed in strains derived from *mut-7* or *dcr-1* mutant background
could be due to rare stochastic telomere elongation events in these backgrounds. (B)
Quantification of telomere shortening rate for each genotype shown in panel A. No
significant difference was found between any genotypes by Mann Whitney U test.
1127
1128

1129

Figure S2. Related to Figure 1. TRF analysis of RNAi-deficient telomerase mutant
strains using telomeric probe. A) A single parental F2 plate was used to generate
multiple F4 plates derived from single worms. F4 *trt-1* single mutant lanes show
consistent bands corresponding to slight shortening of parental bands. In contrast, F4
lanes from *trt-1; mut-7* and *trt-1; dcr-1* show highly variable bands that in some cases
cannot be traced to an obvious parental band. B) Quantification of telomere shortening
rate based on pairing bands appearing at F2 with the closest bands appearing at F4.
Highly smeared lanes were excluded. Each point represents one pair of bands.
Comparing *trt-1* and *trt-1; mut-7* or *trt-1; dcr-1*, mean shortening rate was not
significantly different (p= 0.84 and 0.77, Welch's t-test) but variance was significantly
higher for both double-mutants (p=7.2e-15 and 1.5e-12, F-test). Variance was not
significantly different between the double-mutants (p=0.32).

1142
1143

1144

1145   **Figure S3. Related to Figure 2.** TERRA expression (yellow) in *trt-1* and *trt-1; dcr-*
1146   *1(mg375)* 1-day-old adults detected with a Cy5-labelled telomeric probe.
1147
1148
1149

1150

Figure S4. Related to Figure 2. RNA FISH of TERRA expression in *trt-1* and *trt-1; dcr-1* males. Note TERRA foci in the Pachytene region in both *trt-1* and *trt-1; dcr-1*, as well as lack of expression in the mitotic region and spermatocytes

1151
1152
1153

1154

1155

1156

1157

1158
1159

1160

1161

Figure S5. Related to Figure 3. Analysis of 22G small RNA species in telomerase and Dicer mutants. (A) Number of reads mapping to DCR-1 targets identified by Gent *et al.* (Gent et al., 2010). (B) log2 normalized mean read counts at every *C. elegans* gene for *trt-1* and wildtype. Genes identified as significantly differentially expressed by DESeq2 are shown in red. (C) Proportion of all genes or genes downregulated (trt1_dn) or upregulated (trt1_up) belonging to expression categories defined by Ortiz *et al.* (see reference 54 in main text). (D) log2 normalized mean read counts at transposon consensus sequences for *trt-1* and wildtype.

1170

1171
1172

	Lines tested	Number ALT survivors	Z score	p-value (two tailed)
trt-1	40	6		
trt-1; mut-7(pk204)	40	4	0.68	0.50
trt-1; mut-14(pk738)	40	4	0.68	0.50
trt-1 xnp-1(ok1823)	40	6	0	1
trt-1 xnp-1(tm678)	40	4	0.68	0.50
trt-1; dcr-1 (mg375)	40	5	0.32	0.75

Supplemental Table S1. Related to Figure 1. Statistical analysis of ALT survival in
double mutant strains compared to single mutant *trt-1* control. The Z score represents
how many standard deviations from the number of ALT survivors in double mutants is
from the *trt-1* single mutant control mean survival rate. The p-value represents the
probability that the number of survivors via ALT is not significantly different from the
control *trt-1* single mutant strain.

1199
1200

1201
1202

Strain	Log Rank test (Z=)	p-value
trt-1; mut-7(pk204)	9.79	<0.00001
trt-1; mut-14(pk738)	9.96	<0.00001
trt-1; dcr-1(mg375)	9.37	<0.00001
trt-1; rde-1	8.23	<0.00001
trt-1; rde-4	9.21	<0.00001
trt-1; rrf-3	8.4	<0.00001
trt-1; rrf-2	1.7	0.089
trt-1; exo-1	7.38	<0.00001
trt-1 xnp-1(ok1823)	4.3	<0.00001
trt-1 xnp-1(tm678)	4.23	<0.0001
trt-1 xnp-1(tm678); mut-7	9.21	<0.00001
trt-1; sid-1	6.52	<0.00001
trt-1; rbr-2(tm1231)	8.64	<0.00001
trt-1; spr-5 (by101)	8.89	<0.00001

Supplemental Table S2. Related to Figure 1. Log rank analysis of survival curves of double or triple mutants compare to single mutant *trt-1* controls. n=40 independent lines for each genotype. Strains were passaged by transferring 6 larval L1s weekly and scored as sterile when plates contained no new L1 larvae. The p-value represents the probability that there is no difference in the probability of sterility at any time point between the double or triple mutant strain and the *trt-1* single mutant strain.

1247

1248

trt-1 F1	males	dead embryos	trt-1; dcr-1 F1	males	dead embryos
A1	13	6	A1	0	0
A2	4	26	A2	0	0
A3	8	24	A3	1	0
D1	10	11	A4	0	0
D2	11	4	B1	0	0
D3	4	9	B2	0	0
D4	13	15	B3	0	0
			B4	0	0
			C1	0	0
			C2	0	6
			C3	1	0
			C4	0	0
			D1	0	0
			D2	0	0
			D3	0	0
			D4	0	0

1249

1250

1251

Supplemental Table S3. Related to Figure 1. Counts of total F2 males and dead
embryos scored for F1 progeny from crosses of wild-type males with either late-
generation F16 *trt-1* mutant controls (n=2 independently grown strains, three F1 scored
for strain A and four F1 for strain D) or early generation F4 *trt-1; dcr-1* double mutant
(n=4 strains, 4 F1 each scored for strains A, B, C, and D) hermaphrodites. If a
chromosome fusion is present in any of the F1 animals, then it will cause weak non-
disjunction of the fused chromosomes, which leads to F2 dead embryos if an autosome
is fused or to F2 males if the X chromosome is fused.

1260

1261

1262