Association mapping identified novel candidate loci affecting wood formation in Norway spruce

Baison J[1], Amarylis Vidalis[3], Linghua Zhou[1], Zhi-Qiang Chen[1], Zitong Li[9], Mikko J. Sillanpää[10], Carolina Bernhardsson[1,2], Douglas Scoffield[5], Nils Forsberg[1], Lars Olsson[8], Bo Karlsson[11], Harry Wu[1], Pär K. Ingvarsson[4,6], Sven-Olof Lundqvist[8], Totte Niittylä[1]*, M Rosario García Gil[1]*

*Double last authorship.

Corresponding author: m.rosario.garcia@slu.se

1. Department of Forest Genetics and Plant Physiology, Umeå Plant Science Centre, Swedish University of Agricultural Science, Umeå, Sweden

2. Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

3. Section of Population Epigenetics and Epigenomics, Center of Life and Food Sciences Weihenstephan, Technische Universität München, München, Germany

4. Department of Plant Physiology, Umeå Plant Science Centre, Umeå University, Umeå, Sweden

5. Uppsala Multidisciplinary Center for Advanced Computational Science, Uppsala University, Uppsala, Sweden

6. Department of Ecology and Genetics: Evolutionary Biology, Uppsala University, Uppsala, Sweden

7. Department of Plant Biology, Uppsala BioCenter, Swedish University of Agricultural Science, Uppsala, Sweden.

8. INNVENTIA AB, Drottning Kristinas väg 61, Stockholm, Sweden

9. Ecological Genetics Research Unit, Department of Biosciences, University of Helsinki, P.O. Box 65, FI-00014 Helsinki, Finland

10. Department of Mathematical Sciences, Biocenter Oulu, University of Oulu, Finland

11. Skogforsk, Ekebo 2250 SE-268 90 Svalöv, Sweden

Summary

➤ Norway spruce (*Picea abies*) is an important boreal forest tree species of significant

ecological and economic importance. Hence there is a strong imperative to dissect the

genetics controlling important wood quality traits in Norway spruce.

➤ We performed a functional genome-wide association mapping of 17 wood quality

traits in Norway spruce using 178101 single-nucleotide polymorphisms (SNPs)

generated from exome genotyping of 517 mother trees. The wood quality traits were

defined using functional modelling of wood properties across annual growth rings.

➤ Association mapping was performed using a multilocus LASSO penalized regression

method and we detected a total of 51 significant SNPs from 39 candidate genes that

are involved in wood formation.

➤ Our study represents the first functional multi-locus genome-wide association

mapping (AM) in Norway spruce. The results advance our understanding of the

genetics influencing wood traits, identify novel candidate genes for further functional

studies and support current Norway spruce breeding efforts.

Key Words: Association mapping, Sequence capture, Functional mapping, Single

nucleotide polymorphisms, Candidate genes

Introduction

Norway spruce (*Picea abies* (L.) Karst.) is a dominant boreal softwood species of significant

economic and ecological importance (Hannrup *et al.*, 2004). Long-term Norway spruce

breeding programmes for improvement of growth and survival were initiated in the 1940s and

recently, wood quality has become one of the priority traits (Bertaud & Holmbom, 2004;

Hannrup *et al.*, 2004). Norway spruce breeding in Sweden complete one cycle in about 20

years and such long generation times make improvements in growth and wood quality very

slow. Among wood quality traits, wood density is considered a key indicator of stability,

strength and stiffness of sawn timber (Hauksson *et al.*, 2001). Several studies of wood quality

observed that fast growth conflicts with high quality wood, as shown by the negative genetic

correlation between wood volume growth and density in Norway spruce (Chen *et al.*, 2014;

Dutilleul *et al.*, 1998; Olesen, 1977). In order to combine fast growth and desirable wood

properties through breeding, and to shorten the breeding cycle, it is therefore imperative to

design effective early selection methods and breeding strategies. In an effort to design optimal

breeding and selection strategies for reducing or breaking negative genetic correlations

between traits it is essential to identify alleles that are responsible for generating favourable or

unfavourable genetic correlations (Hallingbäck *et al.*, 2014).

When DNA markers were first introduced in 1980s, tree breeders were provided a

possibility to correlate phenotypes with polymorphic DNA markers and to conduct selection

using genotypes instead of phenotypes (Lande & Thompson, 1990). Groover et al. (1994) first

identified quantitative trait loci (QTL) for wood density variation in loblolly pine using

linkage analyses based on segregating family pedigrees. However, maker-aided selection

(MAS) based on results from QTL analyses was never implemented in practical tree breeding

due to the so-called Beavis effect (e.g. inflated estimates of allelic effects and underestimation

of QTL number for economically important traits) (Beavis, 1998), inconsistent associations

among different families and the low transferability of markers (Strauss *et al.*, 1992).

Association Mapping (AM) is a more powerful QTL detection method that was introduced to

tree genetics using a candidate gene approach (Thumma *et al.*, 2010). AM overcomes the

limited resolution of family-based QTL mapping by relying on historical recombination in the

mapping population (Huang & Han, 2014; Neale & Savolainen, 2004; Thavamanikumar *et*

al., 2013). The effectiveness of AM relies on genome-wide levels of LD, which decays

rapidly within coding regions in conifer species, however, it may be extensive in certain non-

coding regions (Moritsuka *et al.*, 2012). Fast-decaying LD, coupled with complex polygenic

nature for both growth and wood quality traits (Hall *et al.,* 2016) implies that a large number

of genomic regions need to be investigated to identify significant QTL (Beaulieu *et al.*, 2011).

The availability of a draft genome sequence for Norway spruce (Nystedt *et al.*, 2013)

has opened new possibilities for the development of genetic markers to conduct both AM at

the genome-wide level (genome-wide association, GWAS) and genomic selection (GS).

Several reduced representation-based approaches such as sequence capture and transcriptome

sequencing (Hirsch *et al.*, 2014) have been developed as complexity-reduction methods suited

for studying large genomes, such as the 20Gb Norway spruce genome. These approaches

reduce the sequence space by decreasing the repetitive sequence content of the genome. In

this study we employed a solution-based sequence capture method.

Several AM studies have been performed in trees and have identified genetic loci

linked to, for instance, wood properties in *Populus trichocarpa (Porth et al., 2013)*, adaptive

traits in *Pinus contorta* (Parchman *et al.*, 2012) and to wood quality traits in *Eucalyptus*

(Porth *et al.*, 2013; Resende *et al.*, 2012). Such studies aimed at dissecting the genetic basis of

wood properties can benefit from the application of mathematical functions that account for

year-to-year variation across annual growth rings, cambial age and distance from pith (Li *et*

al., 2014). Mathematical modelling allows the incorporation of phenotypic growth trends that

increase the precision and resolution of QTL detection through the integration of the

phenotype information over multiple time points and reduction of residual variance (Ma *et al.*,

2002). Such functional mapping analysis can be conducted using a multistage approach

(Heuven & Janss, 2010). First, the phenotype trends of each individual are modelled using

curve-fitting methods and the parameters describing the curve are then considered as latent

traits. The latent traits are then used in an independent association analyses to search for

genomic regions affecting the trait and to estimate genetic marker effects (Li *et al.*, 2014).

In this study, we applied a functional genome-wide association mapping (AM)

approach to identify genomic regions contributing to wood quality traits in Norway spruce

[*Picea abies* (L.) Karst.]. Estimated breeding values (EBVs) were calculated for growth and

wood quality traits at the resolution of annual growth rings and were then used to extract

latent traits from fitting quadratic splines. We applied quadratic splines since traditional

analyses that utilise a single point data across annual growth rings may confound the analyses

by averaging across a full sample. Such averaging may obscure mechanisms acting at specific

time points during wood formation and will make identification of underlying genes more

difficult. In this study, we have refined our data in order to cover features both in earlywood

and latewood, as well as different phases of juvenility (transitionwood). This study has also

performed the first analysis of number of cells per ring calculated from SilviScan data.

Penalized LASSO regression (Tibshirani, 1996) and the stabilizing selection probability

method of (Meinshausen & Bühlmann, 2010) were then used to detect significant associations

between latent traits derived from EBVs and 178101 SNP markers covering the Norway

spruce genome.

Materials and Methods

Plant material and phenotype data

Plant material and phenotype data used in this study have previously been described in Chen

et al. (2014). In brief, two progeny trails were established in 1990 in Southern Sweden

(S21F9021146 aka F1146 (trial1) and S21F9021147 aka F1147 (trial2)). These trials were

composed of 1373 and 1375 open pollinated families, respectively, and form the basis of our

analyses. We selected 517 families in 112 sampling stands to use in the investigation of wood

properties. At each site, increment wood cores of 12 mm were collected from six trees of the

selected families at breast height (1.3 m) (6 progeny × 2 sites = 12 progenies in total). A total

of 5618 trees, 2973 and 2645 trees from the F1146 and F1147 trails respectively, were

analysed. The pith to bark profiling of the wood physical attributes was analysed using the

SilviScan technology (Evans & Ilic, 2001) at Innventia, Stockholm, Sweden, where the initial

data evaluations were performed using customized methods. These methods focus on the

identification and dating of all annual rings and their compartments of earlywood,

transitionwood and latewood. For the current study, Innventia also calculated three additional

traits, number of cells per ring (NC), wood percentage (WP) and a phenomenon characterised

by trees with a positive correlation between density and growth that we termed Mass Index

(MI) which were all derived from the raw SilviScan data. The traits included in the current

study are listed in Table 1.

149 Table 1 List of the phenotypes, their abbreviations and measurement unit.

Phenotype	Abbreviation	Unit
Ring wood density	WD	kg m^{-3}
Early wood density	EWD	kg m^{-3}
Transition wood density	TWD	kg m^{-3}
Latewood density	LWD	kg m^{-3}
Ring width	RW	μm
Early wood ring width	ERW	μm
Transition wood ring width	TRW	μm
Latewood ring width	LRW	μm
Ring number of cells	NC	
Early wood number of cells	ENC	
Transition wood number of cells	TNC	
Latewood number of cells	LNC	
Early wood percentage	EP	%
Transition wood percentage	TP	%
Latewood percentage	LP	%
Early/Latewood percentage	EP/LP	%
Modulus of elasticity	MOE	GPa
Mass Index (Density/Growth)	MI	

Statistical analysis

EBVs were calculated for each cambial age (annual ring) separately and used for statistical

modelling to derive latent traits. The variance and covariance components were estimated

using ASREML 4.0 (Gilmour *et al.*, 2014) as described in Chen et al., (2014). In brief, the

EBVs at each cambial age were estimated using a univariate, bivariate or multivariate mixed

linear models. The fit of different models were evaluated using the Akaike Information

Criteria (AIC) and the optimal model was selected based on a compromise of model fit and

complexity. Breeding values were then centred in order to obtain within genotype trends. A

univariate linear mixed model for joint-site analysis was used for the analysis as described in

Methods S1.

A number of trees were observed that broke the negative correlation that is usually

observed between density and growth, as these trees exhibited both high density and fast

growth. In order to identify putative genes involved in this favourable combination of trait,

we developed a new dynamic trait that link high density and fast growth, termed, Mass Index

(MI), that we subsequently used in the association mapping. The MI was defined as follows:

*Mass index = (Individual average density/population average density) * (individual*

average disc area / population average disc area).

The index was then treated as a new dynamic trait in the AM analyses where

individuals with an index > 1 indicate a relative wood density to the relative growth rate that

was larger than the average correlation in the population. The index was calculated for all

progeny and were used to calculate BVs for the 517 mother trees.

EBVs were plotted against cambial age (annual ring number) to produce time

trajectories for each trait (Fig. S1) and used to estimate latent curve parameters. At the first

stage, all the trajectories versus cambial age were fitted with a quadratic spline in order to

define latent traits. In this study, the values of four parameters obtained from the spline fitting

were used to describe the dynamics of the EBVs across age: the intercept, slope and two knot

parameters (K1 and K2). The intercept and slopes were used to evaluate the mean and rate of

change for the trait along the annual rings, respectively. K1 and K2 represent inflection points

in the cambial age trajectories. These two points (K1 and K2) were studied due to their

biological significance, as they indicate cambial ages where the development of the EBVs

enters a new phase, thus warranting a closer analysis of the genes imparting this shift in the

EBVs dynamics. The four latent traits show lower correlations compared to the direct

measurements on the original scales and also have constant variances, thereby reducing the

need to account for residual dependencies in the model (Li *et al.*, 2014). The study utilized a

quadratic spline with multiple knots (Methods S2). The latent traits were then analysed using

the LASSO model in order to identify SNPs showing significant associations to the traits

(Methods S3) and to estimate the phenotypic variances explained (PVE) per QTL (Methods

S4).

Sequence capture, genotyping and SNP annotation

Total genomic DNA was extracted from 517 unrelated individuals using the Qiagen Plant

DNA extraction protocol with DNA quantification performed using the Qubit® ds DNA

Broad Range (BR) Assay Kit (Oregon, USA). Probe design and evaluation is described in

Vidalis et al (2017). Sequence capture was performed using the 40 018 diploid probes

previously designed and evaluated for *P. abies* (Vidalis *et al.*, 2018) and samples were

sequenced to an average depth of 15x using an Illumina HiSeq 2500 (San Diego, USA). Raw

reads were mapped against the *P.abies* reference genome v1.0 using BWA-mem (Langmead

& Salzberg, 2012; Li & Durbin, 2009). SAMTools (Li *et al.*, 2009) and Picard (McKenna *et*

al., 2010) were used for sorting and removal of PCR duplicates. Variant calling was

performed using GATK HaplotypeCaller (McKenna *et al.*, 2010) in gVCF output format.

Samples were then merged into batches of ~200 before all 517 samples were jointly called.

Variant Quality Score Recalibration (VQSR) method was performed in order to avoid

the use of hard filtering for exome/sequence capture data. For the VQSR analysis two datasets

were created, a training subset and input file. The training dataset was derived from a Norway

spruce genetic mapping population with loci showing expected segregation patterns

(Bernhardsson *et al.*, 2018) and assigned a prior value of 15.0. The input file was derived

from the raw sequence data using GATK best practices with the following parameters:

extended probe coordinates by +100 excluding INDELS, excluding LowQual sites, and

keeping only bi-allelic sites. The following annotation parameters QualByDepth (QD),

MappingQuality (MQ) and BaseQRankSum, with tranches 100, 99.9, 99.0 and 90.0 were then

applied for the determination of the good versus bad variant annotation profiles. After

obtaining the variant annotation profiles, the recalibration was then applied to filter the raw

variants. Using VCFTools (Danecek *et al.*, 2011), SNP trimming and cleaning involved the

removal of any SNP with a minor allele frequency (MAF) and "missingness" of < 0.05 and

>20%, respectively.

The resultant SNPs were annotated using default parameters for snpEff 4 (Cingolani *et*

al., 2012). Ensembl general feature format (GTF, gene sets) information was utilized to build

the *P. abies* snpEff database.

Genetic Structure

A principal component analysis (PCA) was performed on the sampled trees using SNPs

derived from the sequence capture data. SNPs with missing values following VQSR were

imputed using the nearest neighbour principle in TASSEL (Bradbury *et al.*, 2007). This

approach was essential considering that PCA demands no missing data points. The covariate

matrix derived from the PCA was then displayed by plotting principal component 1 scores

against principal component 2 scores in Figure 2. The PCA plot was used to make inference

about the population structure. The first two components of the PCA covariate matrix

explaining most of the variation were then applied to the AM to account for population

structure and correcting for any stratification within the study.

Linkage disequilibrium was calculated using VCFtools software using the squared

correlation coefficient between genotypes (r2) within scaffolds using the "geno-r2". The

trend-line of LD decay with physical distance was fitted using nonlinear regression (Hill &

Weir, 1988) and the regression line was displayed using R (Team, 2015).

Trait Association Mapping

To determine the association between latent traits and SNPs we implemented a LASSO

penalized regression method as described by Li *et al*. (2014). The LASSO model was then

applied to all traits for the detection of QTLs (Methods S3).

We then calculated the stability selection probability (SSP) of each SNP being

selected to the model, which is applied as a way to control the false discovery rate and

determine significant SNPs (Gao *et al.*, 2014; Li & Sillanpää, 2015). For the quantification of

QTL uncertainty and significance, we utilized hypothesis testing performed at a single level.

For a marker to be declared significant, a SSP inclusion ratio (Frequency) was used with an

inclusion frequency of at least 0.52 for all traits. This frequency inferred that the expected

number of falsely selected markers was less than one (1), according to the formula of

Buhlmann *et al,* (2014). Population structure was accounted for in all analyses by including

the first two principal components based on the genotype data as covariates into the model.

An adaptive LASSO approach (Zou et al. 2006) was used to determine the percentage of

phenotypic variance (PVE) (H^2_{QT}) of all the QTLs (Methods S4).

Candidate gene mining

To assess homology of contigs with significant associations, a BLAST search was performed

against ConGenIE and public sequence databases. After the identification of significant SNPs,

the complete *P. abies* contigs that harboured the QTLs were then BLASTed against the

ConGenIE database and if no significant hit were detected the whole contig was then

extracted. The complete contigs in fasta format were then used to perform a nucleotide

BLAST (Blastn) search using the option for only highly similar sequences (megablast) in the

National Center for Biotechnology Information (NCBI) nucleotide collection database

(https://blast.ncbi.nlm.nih.gov/Blast.cgi?).

Results

Norway spruce SNP identification and mapping population structure

All of the 517 Norway spruce mother trees in the study were considered for variant detection

and an average of 1.5 million paired end reads were sequenced per individual for the 40019

exome capture probes. This resulted in the identification of 178101 high confidence SNPs. In

order to account for effects derived from population stratification we performed a PCA and

identified two separate main population groups as well as a number of individuals scattered in

between these two main groups. Nevertheless, the differences due to population structure

were small with the first two principal components cumulatively explaining only 2.18% of the

genetic variation observed (Figure 2).

Significant SNPs affecting wood traits

Employing a SSP inclusion frequency of at least 0.52 on the intercept, slope and two knots

(K1 and K2) as latent traits, we detected 51 significant QTL across 17 individual traits with

the phenotypic variances explained QTL (H^2_{QTL}) ranging from 0.01 to 4.93% (Table 2).

Several appreciable QTLs were identified with WD and RW having the highest number of

associations, at a total of 13 and 14 QTLs, respectively. This was followed by EP/LP-ratio,

which had six QTLs. WD, RW and EP/LP were the only three observed traits that have QTLs

detected in all four latent traits. For these three phenotypes, the majority of the QTLs were

detected when the average ring phenotype was used to derive the latent traits (Table 2). NC

associated with one QTL that was detected for the entire ring, whilst six QTLs were identified

when EW, TW and LW were analysed separately, with H^2_{QTL} ranging from 0.01-4.93%

(Table 2).

293   Table 2 SNP features, Latent traits, QTL, Alleles and Frequencies

Phenotype	Latent Trait	Marker	Allele	SNP Feature	Frequency	PVE
WD	Intercept	167610	A/G	Downstream gene variant	0.71	4.64%
	Slope	30469	A/G	Upstream gene variant	0.72	4.50%
	K1	30469	A/G	Upstream gene variant	0.551	4.15%
	K2	157442	G/A	Upstream gene variant	0.567	2.43%
EWD	Intercept	167610	A/G	Downstream gene variant	0.545	3.38%
	Slope	23798	C/T	Upstream gene variant	0.53	0.69%

		70955	T/A	Upstream gene variant	0.644	0.40%
TWD	Slope	131698	G/A	Stop gained	0.672	1.58%
		160208	T/C	Intron variant	0.595	3.41%
	K1	89044	T/C	Upstream gene variant	0.534	3.34%
LWD	Slope	43797	T/C	Missense variant	0.524	1.81%
		165481	C/T	Intron variant	0.588	1.21%
		171223	G/A	Intron variant	0.712	4.03%
RW	Intercept	11535	A/C	Synonymous variant	0.545	1.95%
		112391	C/T,A	Stop gained	0.532	1.45%
		112394	C/A	Splice region variant	0.692	2.56%
	Slope	165481	C/T	Intron variant	0.521	2.66%
	K1	23808	T/G	Synonymous variant	0.554	1.78%
		165481	C/T	Intron variant	0.533	0.18%
	K2	23808	T/G	Synonymous variant	0.55	1.20%
		165481	C/T	Intron variant	0.615	1.79%
TRW	Slope	111057	T/A	Missense variant	0.685	
	K1	33110	T/A	Upstream gene variant	0.657	3.23%
		89295	G/A	Upstream gene variant	0.688	4.51%
	K2	111057	T/A	Missense variant	0.672	
LRW	Intercept	143628	C/T	Downstream gene variant	0.668	0.5%
	K2	164772	C/A	Downstream gene variant	0.571	0.06%
MOE	Slope	165481	C/T	Intron variant	0.602	1.00%
NC	K1	145839	G/C	Upstream variant	0.645	3.82%
ENC	Slope	98508	A/C	Upstream gene variant	0.667	0.03%
		167610	A/G	Downstream gene variant	0.685	0.01%

TNC	Intercept	95870	T/C	Upstream gene variant	0.667	3.78%
		126785	A/C	Upstream gene variant	0.68	4.93%
LNC	Intercept	143628	C/T	Downstream gene variant	0.66	3.14%
	Slope	143628	C/T	Downstream gene variant	0.672	4.77%
EP	Intercept	16868	G/T	Intron variant	0.609	3.32%
		91242	G/A	Synonymous variant	0.594	3.41%
TP	Intercept	101203	A/C	Upstream gene variant	0.594	1.16%
		132014	A/C	Upstream gene variant	0.601	3.22%
LP	K1	162397	C/T	Upstream gene variant	0.892	
EP/LP	Intercept	51657	G/A	Downstream gene variant	0.63	0.81%
		60787	C/T	Intron variant	0.655	1.80%
		123639	A/C	Upstream gene variant	0.628	0.75%
	K1	59480/36496	A/G	Synonymous	0.6	2.37%
	K2	117333	T/C	Missense variant	0.523	
	K3	72414	A/T	Non-Coding	0.617	4.05
Growth/De nsity	Intercept	166235	G/A	Intergenic variant	0.533	0.65%
	Slope	61096	G/A	Synonymous variant	0.66	0.01%
		67181	G/A	Missense variant	0.612	0.05%
		1401	C/A	Exon/stop gained	0.588	1.19%
		138744	G/T	Missense variant	0.58	1.80%
		162397	C/T	Upstream gene variant	0.627	1.44%
	K1	21924	A/G	Upstream gene variant	0.71	1.82%

Several QTLs shared *within* each trait and *across* traits were observed in the analysis.

WD, RW, TRW and LNC had one (30469), two (165481 and 23808), one (111057) and one

(143628) QTL shared by two latent traits, respectively. One of the common QTL (30469) for

WD had a frequency of 0.72 with an H^2_{QTL} of 4.50% for the slope trait, which indicates that it

is highly significant for the phenotypes. Common QTLs *within* RW were observed for slope,

K1 and K2 latent traits, with moderate frequencies ranging from 0.521 to 0.615 and

influenced their respective traits to modest degree (H^2_{QTL} in ranges of 0.18-2.66%).

For QTLs common *across* the different latent traits, QTL 165481 was shared between

LWD, RW and MOE; this is not surprising because of the close correlation between MOE

and wood density, with its generally negative correlation to RW. Intron variant

MA_10434805g0010_165481 explained between 0.18-2.66% of the H^2_{QTL} observed in the

respective traits. The SNP associated with this QTL also had high frequencies of 0.602 and

0.615 in MOE and RW explaining H^2_{QTL} of 1.00 and 2.66%, respectively. It was also

observed that the SNP MA_10434805g0010_165481 is a common QTL *within* the wood

traits related to Width (Table 2). SNP MA_10435406g0010_167610 was shared between

WD, EWD and ENC. This SNP was characterized by having high frequencies in WD (0.71)

and ENC (0.685), however it had a moderate frequency of 0.545 for EWD. This QTL was

detected by the intercept latent trait for WD and EWD, and the slope latent trait in ENC

(Table 2), with H^2_{QTL} ranging from 0.01-4.64%. The QTL had a high influence on the density

related traits as it explained 4.64% (WD) and 3.38% (EWD).

Trees showing a positive correlation between growth and density had seven QTL

specific for this observed phenomenon (MI) and had modest influence on the trait (H^2_{QTL} in

the ranges 0.05-1.82%). Five of the QTL were detected using the slope as the latent trait with

high frequencies ranging between 0.58 to 0.66 for SNP 61096 (Table 2).

Genetic associations with phenotypes

Sequence capture and the SNP-trait associations allowed the mining of candidate genes

involved in spruce wood formation coupled with the identification of orthologous annotations

and descriptions from *Populus* and *Arabidopsis*.

Width related traits, RW, TRW, and LRW were associated with nine gene models. For RW

five genes, endoglucanase 11-like, Alpha-dioxygenase 1 (DIOX1), Proliferating cell nuclear

antigen (PCNA), B3-DNA-binding and E3 ubiquitin-protein ligase were identified. The SNP

MA_879270g0010_112391, a splice region variant, explained 2.56% of the H^2_{QTL} and is

associated with DIOX1. Marker MA_20322g0010_23808 for RW is associated with the

protein domain for a plant specific B3-DNA binding protein, explaining 1.78% variation, with

similar orthologous genes in *Arabidposis* and *Populus* (Table S1). The three putative genes

associated with TRW are a Serine/threonine-protein kinase, a Homeodomain protein (HB2)

and a Senescence-associated protein, and all have high H^2_{QTL} ranging from 2.14 to 4.50%.

Contig MA_10434624 is homologos to a Pectinesterase and was associated with the

downstream variant MA_10434624g0010_164772 for LRW. This may suggest a link between

LRW and pectin modification. QTL associated with gene MA_214776g0010 for the TRW

may be linked with serine/threonine-protein kinase gene (Os01g0689900), this occurrence of

kinase-like related genes was also observed across TRW, NC, EP, EP/LP and EWD (Table

S1).

NC as well as the three compartments (ENC, TNC and LNC) are associated with a

total of three putative genes and three protein domains. Of the three putative genes, two are

associated with serine/kinase activity and one is involved in cysteine and methionine

synthesis (Table 1 S1). All the SNPs associated with these traits were either downstream or

upstream of coding regions and may thus act as modifiers of gene expression. The SNP

MA_402880g0010_98508 (an upstream gene variant) significantly associated with ENC

located on gene MA_402880g0010 is homologous to a *Populus* sphingolipid biosynthesis

protein. SNP MA_9447489g0010_126785 associated with TNC was located in the gene

MA_9447489g0010 which is homologous to a peptidase domain from *Arabidopsis* and

showed the highest H^2_{QTL} in the dataset (4.93%). This domain is similar to an orthologous

zinc carboxypeptidase enzyme of *Oryza sativa* (Zn-dependent exopeptidases superfamily

protein) (Table S1).

Wood percentage traits, EP, LP, TP and the ratio of EP/LP had significant associations

with ten SNPs. Four of the six significant SNP variants for EP/LP are modifiers with the other

two SNPs, being a synonymous (MA_96191g0010_59480) and missense

(MA_1045136g0010_117333) variant. The synonymous SNP MA_96191g0010_59480 was

associated with the gene model MA_96191g0010, which is homologous to a *P. sitchensis*

Glycosyltransferase (GT), similar to UDP-glucosyltransferase 73B2 (AT4G34135) from

Arabidopsis. Five protein domains were also detected, that were linked to phytochrome

kinase substrate 1, TIR/NBS/LRR and zein binding domains (Table S1). The significant SNP

MA_15729g0010_16868, an intron variant, that is associated with EP, is located in the gene

MA_15729g0010, which is homologous to a DNA-3-methyladenine glycosylase II enzyme.

The SNPs identified for TP and LP are all downstream gene variants (Table 2).

Wood density based properties, WD, EWD, TWD and LWD had a total of 12

significant associations. For the associations with WD we identified the SNP

MA_10435406g0010_167610 that is a 3′-gene variant which explained the highest H^2_{QTL}

observed (4.64%) and is located in a gene that is homologous to a Phosphoadenosine

phosphosulfate reductase gene cysH_2. This locus was also detected for EWD and ENC

explaining H^2_{QTL} of 3.38% and 0.01%, respectively. A missense SNP,

MA_33109g0010_30469, was associated with WD and located within the gene

MA_33109g0010 homologous to an *Arabidopsis* senescence associated gene 24 (Table S1).

The three significant SNPs identified for EWD were all modifiers, upstream and downstream

gene variants. Of the three significant SNP associations for TWD, two, SNP

MA_10235390_131698 (stop gained) and SNP MA_212523g0010_89044 (upstream gene

variant), were identified within genes. The intron variant MA_10433411g0010_160208

associated with TWD and is found in the gene MA_10433411g0010 that is homologous to an

Arabidopsis Transducin/WD40 repeat-like superfamily protein. Two of the three significant

SNPs identified for LWD were intron variants (MA_10434805g0010_165481 and

MA_10436058g0010_171223) with the third being a missense variant

(MA_62987g0010_43797). The SNP MA_10434805g0010_165481 was found in the gene

MA_10434805g0010, which is homologous to an *Arabidopsis* Proliferating Cell Nuclear

Antigen Protein (PCNA). This SNP is also associated with RW and explained 1.21% and

2.66% $H^2_{QTL,}$ respectively.

The Mass Index trait, that is linked to a positive effect of wood volume growth and

increased density (growth-density) yielded a total of seven associated SNPs, with two

upstream gene variants, two missense variants one intergenic variant, one stop gained variant

and one synonymous nucleotide replacement (Table 2). The slope latent trait had five genes

with modest influence on the phenotype ranging from 0.01-1.80%. The genes were

homologous to *Arabidopsis* GRAS transcription factor, Aluminium induced protein, Protein

virilizer, ARM repeat superfamily protein and an uncharacterized protein. The SNP

MA_1378g0010_1401 encodes for a premature stop codon (stop gained, a high impact

variant) on gene MA_1378g0010, which is homologous to an *Arabidopsis* protein virilizer

involved in mRNA splicing regulation. The gene homologous to the GRAS transcription

factor was associated with the SNP MA_99004g0010_61096 (a synonymous variant). The

SNP MA_19222g0010_21921, an upstream gene variant, was located in the gene

MA_19222g0010 which is homologous to a *Picea sitchensis* ADP (NB-ARC domain) and

explained the highest H^2_{QTL} of 1.82% (Table S1).

Wood density traits were associated with a total of 12 genes, the largest number of

genes identified from the contigs. Percentage of wood was linked to ten putative genes, cell

width had nine putative genes and number of cells was associated with six genes. Two genes

were shared *across* multiple traits, PCNA was common *across* RW and LWD, and

phophoadenosine phosphosulfate reductase was shared *across* WD, EWD and ENC. Genes

with the Serine/threonine-protein phosphatase and TIR-NBS-LRR domains were also

identified *across* width, wood density and cell percentage traits.

Discussion

We applied a functional mapping approach in a genome-wide association mapping context and identified 51 significant QTLs that were associated with wood formation in Norway spruce. Previous work utilizing a functional mapping analysis in forest trees have used a limited number of molecular markers (Li *et al.*, 2014; Ma *et al.*, 2002). Li *et al.,* applied this analysis in a bi-parental Scots pine (*P. sylvestris* L.) cross using 319 markers. Hence, our work represents a major advance in that we have been able to apply this approach at a genome-wide scale (178101 SNPs) on unrelated mother trees, with a dynamic functional trait dataset comprising 15-time points/annual growth rings (*i.e.,* cambial age).

The number of detected QTLs is relatively small compared with several recent studies in *Populus* (Evans *et al.*, 2014; McKown *et al.*, 2014; Porth *et al.*, 2013). The sample size, number of SNPs used and the stringency with which we accepted significant SNPs contributed to the modest number of QTL. Previous functional mapping studies, (Li *et al.*, 2014) involving SNPs in conifers have used two levels of evaluating QTLs, whereby they have suggestive and significant QTL. In our study, we only reported significant QTL. As indicated in Hall *et al* (2016), there should be hundreds to thousands of QTL of moderate to very small effect related to growth and wood quality traits in trees. Hence, a large population and accurate phenotyping are required for a reliable identification of most QTLs (Korte & Farlow, 2013). However, the sample size of our study allowed the detection of the largest/most significant QTL. The study identified significant associations explaining relatively small proportions of the phenotypic variance being observed, ranging from 0.01-4.93%. This is in line with other studies of QTL for wood traits (González-Martínez *et al.*, 2007; Porth *et al.*, 2013).

Genetic associations with potential to improve wood properties

With all the SNPs, having been derived from known genomic positions, it was

possible to identify genes linked to the associated QTLs and infer their potential function in

wood formation.

The gene MA_10694g0010 is homologous to an enzyme involved in cell wall

biosynthesis, endoglucanase 11-like, and was associated with RW (Table S1). This enzyme is

a vital component of xylogenesis and is involved in the active digestion of the primary cell

wall (Goulao *et al.*, 2011). The endoglucanase 11-like, was associated with a synonymous

SNP MA_10694g0010_11535 for (RW) suggesting an involvement in cell expansion and cell

wall loosening during wood formation. Endoglucanases have been proposed as enzymes

involved in controlling cell wall loosening (Cosgrove, 2005). Endoglucanase 11-like gene is

part of the endo-1 family in which the eno-1-4-β-glucanase *Korrigan* gene belongs.

Characterisation of the *Korrigan* gene in *P. glauca* has identified it to be functionally

conserved and essential for cellulose synthesis (Maloney *et al.*, 2012). Hence,

MA_10694g0010 is a candidate for the remodelling of cell walls that affects the mechanical

and growth properties of wood cells, and consequently annual ring width.

The synonymous SNP MA_20322g0010_23808 is associated with RW located on

gene MA_20322g0010 which is homologous with a plant specific B3-DNA binding protein

domain, that is shared among various plant-specific transcription factors. This includes

transcription factors involved in auxin and abscisic acid responsive transcription (Yamasaki *et*

al., 2004). Auxin is one of the central phytohormones in the control of plant growth and

development (Abel & Theologis, 1996), and also known to be involved in cell wall loosening

and elongation (Cosgrove, 2016).   This suggests a possible functional role for

MA_20322g0010 in influencing RW.

An intron variant located in the MA_10434805g0020 gene, which is homologous to

PCNA was detected *across* LWD, RW and MOE. PCNA proteins function as integral

enzymes in the regulatory pathways of cell cycle regulation and DNA metabolism (Maga &

Hübscher, 2003). PCNA has been associated with chromatin remodeling, DNA repair, sister-

chromatid cohesion and cell cycle control, which are all vital processes in plant growth

(Strzalka & Ziemienowicz, 2010), but it has not been previously associated with wood

formation traits.

In our study we detected a significant downstream SNP

(MA_10434624g0010_164772) associated with LRW on gene MA_10434624g0020,

homologous to pectinmethylesterases (PMEs), which are cell wall associated enzymes

responsible for demethylation of polygalacturonans (Phan *et al.*, 2007). This enzyme has been

shown to be linked with many developmental processes in plants, such as, cellular adhesion

and stem elongation (Micheli, 2001). An association study in White spruce identified a

significant nonsynonymous SNP coding for cysteine associated with earlywood and total

wood cell wall thickness associated with pectinmethylesterase (Beaulieu *et al.*, 2011). Our

study identified a PME SNP association in the latewood stage, supporting the importance of

PMEs in wood cell development.

A SNP (MA_10435406g0010_167910) downstream on gene MA_10435406g0010

was detected *across* the traits ENC, WD and EWD. The gene is homologous to

Phosphoadenosine phosphosulfate reductase (PAPS), which plays a central role in the

reduction of sulphur in plants. An analysis of PAPS enzymes in *Arabidopsis* (Klein &

Papenbrock, 2004) and *Populus* (Kopriva *et al.*, 2004) revealed that enzymes involved in

sulphate-conjugation, play an important role in plant growth and development (Klein &

Papenbrock, 2004). Reduced sulphur is utilized by the sulphate assimilation pathway for the

synthesis of essential amino acids cysteine and methionine (Kopriva & Koprivova, 2004).

Methione acts as a methyl donor in both lignin, hemicellulose and pectin biosynthesis

providing a possible mechanism of how PAPS could influence wood density and number of

cells.

When analyzing QTLs detected for traits linked to the percentage of cells (EP, LP and

EP/LP) we identified three putative candidate genes, DNA-3-methyladenine glycosylase II

enzyme, phytochrome kinase substrate 1 and glycosyltransferase. DNA-3-methyladenine

glycosylase II enzyme is responsible for carrying out base excision repairs (BER) in the

genome in order to maintain genomic integrity. This enzyme has the ability to initiate a broad

substrate recognition and provides a wide resistance to DNA damaging agents (Wyatt *et al.*,

1999). This DNA repair capacity can be expected to be essential for the process of cell

propagation and growth.

A synonymous SNP (MA_96191_59480) within the gene MA_96191g0010, which is

homologous to Glucosyltransferase in *P. sitchensis* was associated with EP/LP. Glycosyl

transferases operate by facilitating the catalytic sequential transfer of sugars from activated

donors to acceptor molecules that form region and stereospecific glycosidic linkages (Lairson

et al., 2008). The Arabidopsis ortholog (UDP-glucosyltransferase 73B2) encodes for a

putative flavonol 7-O-glucosyltransferase involved in stress responses. In our study, this

significant association was associated with EP/LP, however a nonsynonymous variant in a

gene coding for a Glycosyl transferase in *Populus* was associated with fibre development and

elongation (Porth *et al.*, 2013). Therefore, gene MA_96191g0010 is a novel candidate for

further investigation of how flavonol metabolism may influence the proportion of early and

late wood in Norway spruce.

Two genes concerning wood formation, PAPS and PCNA, were also detected across

related traits density, growth number of cells and MOE. Significant SNP

(MA_10435406_167610) in the PAPS reductase gene is common *across* ENC, WD and

EWD, with SNP MA_10434805_165481 located in an intron for a gene encoding for PCNA

protein being detected *across* WD, RW and MOE (Table S1). The presence of these common

QTL suggests that these traits might be under the control of the same genes or genetic

pathways. Chen et al (2014) reported a significant positive genetic correlation between wood

density and MOE, which increased with tree age. However, wood volume growth and density

have a negative correlation (Chen *et al.*, 2014), with our study being the first to detect QTLs

for trees exhibiting a positive correlation for this phenomenon (MI). The common QTL

observed *across* WD, EWD and ENC indicates that the number of cells during the juvenile

wood development stages has a significant impact on the overall density. The seasonal

changes in EWD to LWD has been speculated to be due to a change in auxin levels leading to

the initiation of wall-thickening phase, which has a direct impact on the wood quality traits

such as MOE. This phase coincides with the cessation of height growth and where available

resources are used for cell-wall thickening (Sewell *et al.*, 2000), which may explain the

common QTL between LWD, RW and MOE, as part of the same feedback loop mechanism.

We identified two associations to homologous genes related to nucleic acid repair

functions, DICER-LIKE3 (DCL3) and DNA mismatch repair protein (MSH5), which are

concerned with RNA processing as well as DNA repair, respectively. These genes are

involved in ensuring the fidelity of DNA replication and to preserve genomic integrity (Hsieh

& Yamane, 2008). These genes are possibly associated with cambial cell division and endo-

reduplication during wood formation and can conceivably have effects on wood density.

An association for TWD with a SNP located upstream of gene MA_212523g0010, is

homologous to Kinesin-related protein 13 (gene-L484_021891). Kinesin-related proteins are

known to be involved in secondary wall deposition, cell wall strength and oriented deposition

of cellulose microfibrils, which can impact wood density (Zhong *et al.*, 2002).

Several   receptor-like   Kinases   (TIR/NBS/LRR   and   Serine/threonine-protein

phosphatase) homologs were identified *across* traits (TRW, NC, EP, EP/LP and EWD) (Table

S1). These protein domains control a large range of processes including hormone perception

and plant development. Approximately 2.5% of the annotated genes in *Arabidopsis* genome

are RLK homologs (Shiu & Bleecker, 2001), where they among other functions play an

important role in the differentiation and separation of xylem and phloem cells (Fisher &

Turner, 2007). Similar to our study a synonymous SNP in a RLK gene was associated with

early wood proportion (EP) in White spruce (Beaulieu *et al.*, 2011), hence RLKs seem to be

involved in modifying a number of different wood properties from density to cell identity and

number.

Norway spruce trees that possess the ability of fast growth and high wood density are

very rare, but such trees and associated SNPs were discovered in our study. Trees combining

these traits are of high interest to both breeders and foresters. Of the seven genes significantly

linked to this phenomenon of particular interest was a synonymous SNP on MA_99004g0100

gene homologous to a transcription factor from the GRAS family (Table S1). GRAS is an

important class of plant-specific proteins derived from three members: GIBBERELLIC-ACID

INSENSITIVE (GAI), REPRESSOR of GAI (RGA) and SCARECROW (SCR) (GRAS)

(Hirsch & Oldroyd, 2009). GRAS genes are known to be involved in the regulation of plant

development through the regulation of gibberellic acid (GA) and light signalling (Cenci &

Rouard, 2017; Hirsch & Oldroyd, 2009). Furthermore GA signalling has also been shown to

stimulate wood formation in *Populus* (Mauriat & Moritz, 2009). Thus, the GRAS

transcription factor identified here and the other six genes positively associated with MI

provide interesting genetic markers and tools to understand this phenomenon.

Conclusion

This work has used "functional" association mapping to dissect the genetic basis of wood

properties in Norway spruce. In total, we identified 51 Significant QTLs for wood properties

and mining of candidate genes located in the vicinity of significant QTLs identified a host of

genes that could be directly or indirectly responsible for the observed traits. These genes are

candidates for further functional verification in Norway spruce. This work provides a

framework to dissect the genetic nature of these traits and will add to our understanding of

some of the fundamentals of tree development.

Acknowledgements

We acknowledge the support of the Bio4Energy research organization for wood property

analyses. All genetic data was obtained through funding from the Knut and Alice Wallenberg

foundation. SSF project support of continuing this work. JB is supported though a postdoc

position funded by the Kempe foundation.

References

Abel, S., & Theologis, A. (1996). Early Genes and Auxin Action. *Plant Physiology, 111*(1), 9.

Beaulieu, J., Doerksen, T., Boyle, B., Clément, S., Deslauriers, M., Beauseigle, S., Blais, S., Poulin, P.-L., Lenz, P., Caron, S., Rigault, P., Bicho, P., Bousquet, J., & MacKay, J. (2011). Association Genetics of Wood Physical Traits in the Conifer White Spruce and Relationships With Gene Expression. *Genetics, 188*(1), 197-214. doi: 10.1534/genetics.110.125781

Beavis, W. D. (1998). QTL analyses: power, precision, and accuracy. *Molecular dissection of complex traits, 1998*, 145-162.

Bernhardsson, C., Vidalis, A., Wang, X., Scofield, D. G., Shiffthaler, B., Baison, J., Street, N. R., Garcia Gil, M. R., & Ingvarsson, P. K. (2018). An ultra-dense haploid genetic map for evaluating the highly fragmented genome assembly of Norway spruce (Picea abies). *bioRxiv*. doi: 10.1101/292151

Bertaud, F., & Holmbom, B. (2004). Chemical composition of earlywood and latewood in Norway spruce heartwood, sapwood and transition zone wood. *Wood Science and Technology, 38*(4), 245-256. doi: 10.1007/s00226-004-0241-9

Bradbury, P. J., Zhang, Z., Kroon, D. E., Casstevens, T. M., Ramdoss, Y., & Buckler, E. S. (2007). TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics, 23*(19), 2633-2635.

Cenci, A., & Rouard, M. (2017). Evolutionary Analyses of GRAS Transcription Factors in Angiosperms. *Frontiers in Plant Science, 8*, 273. doi: 10.3389/fpls.2017.00273

Chen, Z.-Q., Gil, M. R. G., Karlsson, B., Lundqvist, S.-O., Olsson, L., & Wu, H. X. (2014). Inheritance of growth and solid wood quality traits in a large Norway spruce population tested at two locations in southern Sweden. *Tree Genetics & Genomes, 10*(5), 1291-1303.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X., & Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly, 6*(2), 80-92.

Cosgrove, D. J. (2005). Growth of the plant cell wall. *Nat Rev Mol Cell Biol, 6*(11), 850-861. doi: http://www.nature.com/nrm/journal/v6/n11/suppinfo/nrm1746_S1.html

Cosgrove, D. J. (2016). Catalysts of plant cell wall loosening. *F1000Research, 5*, F1000 Faculty Rev-1119. doi: 10.12688/f1000research.7180.1

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., & Sherry, S. T. (2011). The variant call format and VCFtools. *Bioinformatics, 27*(15), 2156-2158.

Dutilleul, P., Herman, M., & Avella-Shaw, T. (1998). Growth rate effects on correlations among ring width, wood density, and mean tracheid length in Norway spruce (Picea abies). *Canadian Journal of Forest Research, 28*(1), 56-68.

Evans, L. M., Slavov, G. T., Rodgers-Melnick, E., Martin, J., Ranjan, P., Muchero, W., Brunner, A. M., Schackwitz, W., Gunter, L., Chen, J.-G., Tuskan, G. A., & DiFazio, S. P. (2014). Population genomics of Populus trichocarpa identifies signatures of selection and adaptive trait associations. *Nat Genet, 46*(10), 1089-1096. doi: 10.1038/ng.3075 http://www.nature.com/ng/journal/v46/n10/abs/ng.3075.html#supplementary-information

Evans, R., & Ilic, J. (2001). Rapid prediction of wood stiffness from microfibril angle and density. *Forest products journal, 51*(3), 53.

Fisher, K., & Turner, S. (2007). PXY, a receptor-like kinase essential for maintaining polarity
during plant vascular-tissue development. *Current Biology, 17*(12), 1061-1066.
Gao, H., Wu, Y., Li, J., Li, H., Li, J., & Yang, R. (2014). Forward LASSO analysis for high-order
interactions in genome-wide association study. *Briefings in Bioinformatics, 15*(4),
552-561.
Gilmour, A., Gogel, B., Cullis, B., Welham, S., Thompson, R., Butler, D., Cherry, M., Collins, D.,
Dutkowski, G., & Harding, S. (2014). ASReml user guide. Release 4.1 structural
specification. *VSN International Ltd, Hemel Hempstead, HP1 1ES, UK www. vsni. co.
uk*.
González-Martínez, S. C., Wheeler, N. C., Ersoz, E., Nelson, C. D., & Neale, D. B. (2007).
Association genetics in Pinus taeda Ll Wood property traits. *Genetics, 175*(1), 399-
409.
Goulao, L. F., Vieira-Silva, S., & Jackson, P. A. (2011). Association of hemicellulose- and
pectin-modifying gene expression with Eucalyptus globulus secondary growth. *Plant
Physiology   and   Biochemistry,   49*(8),   873-881.   doi:
https://doi.org/10.1016/j.plaphy.2011.02.020
Hallingbäck, H. R., Sánchez, L., & Wu, H. X. (2014). Single versus subdivided population
strategies in breeding against an adverse genetic correlation. *Tree Genetics &
Genomes, 10*(3), 605-617.
Hannrup, B., Cahalan, C., Chantre, G., Grabner, M., Karlsson, B., Bayon, I. L., Jones, G. L.,
Müller, U., Pereira, H., & Rodrigues, J. C. (2004). Genetic parameters of growth and
wood quality traits in Picea abies. *Scandinavian Journal of Forest Research, 19*(1), 14-
29.
Hauksson, J. B., Bergqvist, G., Bergsten, U., Sjöström, M., & Edlund, U. (2001). Prediction of
basic wood properties for Norway spruce. Interpretation of Near Infrared
Spectroscopy data using partial least squares regression. *Wood Science and
Technology, 35*(6), 475-485. doi: 10.1007/s00226-001-0123-3
Heuven, H. C., & Janss, L. L. (2010). *Bayesian multi-QTL mapping for growth curve
parameters*. Paper presented at the BMC proceedings.
Hill, W., & Weir, B. (1988). Variances and covariances of squared linkage disequilibria in
finite populations. *Theoretical population biology, 33*(1), 54-78.
Hirsch, C. D., Evans, J., Buell, C. R., & Hirsch, C. N. (2014). Reduced representation
approaches to interrogate genome diversity in large repetitive plant genomes.
Briefings in Functional Genomics, elt051.
Hirsch, S., & Oldroyd, G. E. D. (2009). GRAS-domain transcription factors that regulate plant
development. *Plant Signaling & Behavior, 4*(8), 698-700.
Hsieh, P., & Yamane, K. (2008). DNA mismatch repair: molecular mechanism, cancer, and
ageing. *Mechanisms of ageing and development, 129*(7), 391-407.
Huang, X., & Han, B. (2014). Natural variations and genome-wide association studies in crop
plants. *Annual review of plant biology, 65*, 531-551.
Klein, M., & Papenbrock, J. (2004). The multi-protein family of Arabidopsis
sulphotransferases and their relatives in other plant species. *Journal of experimental
botany, 55*(404), 1809-1820. doi: 10.1093/jxb/erh183
Kopriva, S., Hartmann, T., Massaro, G., Hönicke, P., & Rennenberg, H. (2004). Regulation of
sulfate assimilation by nitrogen and sulfur nutrition in poplar trees. *Trees, 18*(3), 320-
326. doi: 10.1007/s00468-003-0309-4

Kopriva, S., & Koprivova, A. (2004). Plant adenosine 5′-phosphosulphate reductase: the past,
the present, and the future. *Journal of experimental botany, 55*(404), 1775-1783. doi:
10.1093/jxb/erh185
Korte, A., & Farlow, A. (2013). The advantages and limitations of trait analysis with GWAS: a
review. *Plant methods, 9*(1), 1.
Lairson, L., Henrissat, B., Davies, G., & Withers, S. (2008). Glycosyltransferases: structures,
functions, and mechanisms. *Annu. Rev. Biochem., 77*, 521-555.
Lande, R., & Thompson, R. (1990). Efficiency of marker-assisted selection in the
improvement of quantitative traits. *Genetics, 124*(3), 743-756.
Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature
Methods, 9*(4), 357-359.
Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics, 25*(14), 1754-1760.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., &
Durbin, R. (2009). The sequence alignment/map format and SAMtools.
Bioinformatics, 25(16), 2078-2079.
Li, Z., Hallingbäck, H. R., Abrahamsson, S., Fries, A., Gull, B. A., Sillanpää, M. J., & García-Gil,
M. R. (2014). Functional multi-locus QTL mapping of temporal trends in Scots pine
wood traits. *G3: Genes| Genomes| Genetics, 4*(12), 2365-2379.
Li, Z., & Sillanpää, M. J. (2015). Dynamic Quantitative Trait Locus Analysis of Plant Phenomic
Data. *Trends in Plant Science, 20*(12), 822-833. doi:
http://dx.doi.org/10.1016/j.tplants.2015.08.012
Ma, C.-X., Casella, G., & Wu, R. (2002). Functional mapping of quantitative trait loci
underlying the character process: a theoretical framework. *Genetics, 161*(4), 1751-
1762.
Maga, G., & Hübscher, U. (2003). Proliferating cell nuclear antigen (PCNA): a dancer with
many partners. *Journal of Cell Science, 116*(15), 3051.
Maloney, V. J., Samuels, A. L., & Mansfield, S. D. (2012). The endo-1, 4-β-glucanase Korrigan
exhibits functional conservation between gymnosperms and angiosperms and is
required for proper cell wall formation in gymnosperms. *New Phytologist, 193*(4),
1076-1087.
Mauriat, M., & Moritz, T. (2009). Analyses of GA20ox-and GID1-over-expressing aspen
suggest that gibberellins play two distinct roles in wood formation. *The Plant Journal,
58*(6), 989-1003.
McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,
Altshuler, D., Gabriel, S., & Daly, M. (2010). The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome
Research, 20*(9), 1297-1303.
McKown, A. D., Klápště, J., Guy, R. D., Geraldes, A., Porth, I., Hannemann, J., Friedmann, M.,
Muchero, W., Tuskan, G. A., & Ehlting, J. (2014). Genome-wide association implicates
numerous genes underlying ecological trait variation in natural populations of
Populus trichocarpa. *New Phytologist, 203*(2), 535-553.
Meinshausen, N., & Bühlmann, P. (2010). Stability selection. *Journal of the Royal Statistical
Society: Series B (Statistical Methodology), 72*(4), 417-473.

Micheli, F. (2001). Pectin methylesterases: cell wall enzymes with important roles in plant physiology. *Trends in Plant Science, 6*(9), 414-419. doi: http://doi.org/10.1016/S1360-1385(01)02045-3

Moritsuka, E., Hisataka, Y., Tamura, M., Uchiyama, K., Watanabe, A., Tsumura, Y., & Tachida, H. (2012). Extended linkage disequilibrium in noncoding regions in a conifer, Cryptomeria japonica. *Genetics, 190*(3), 1145-1148.

Neale, D. B., & Savolainen, O. (2004). Association genetics of complex traits in conifers. *Trends in Plant Science, 9*(7), 325-330.

Nystedt, B., Street, N. R., Wetterbom, A., Zuccolo, A., Lin, Y.-C., Scofield, D. G., Vezzi, F., Delhomme, N., Giacomello, S., & Alexeyenko, A. (2013). The Norway spruce genome sequence and conifer genome evolution. *Nature, 497*(7451), 579-584.

Olesen, P. (1977). The variation of the basic density level and tracheid width within the juvenile and mature wood of Norway spruce. *For. Tree Improv, 12*, 1-22.

Parchman, T. L., Gompert, Z., Mudge, J., Schilkey, F. D., Benkman, C. W., & Buerkle, C. (2012). Genome-wide association genetics of an adaptive trait in lodgepole pine. *Molecular Ecology, 21*(12), 2991-3005.

Phan, T. D., Bo, W., West, G., Lycett, G. W., & Tucker, G. A. (2007). Silencing of the Major Salt-Dependent Isoform of Pectinesterase in Tomato Alters Fruit Softening. *Plant Physiology, 144*(4), 1960-1967. doi: 10.1104/pp.107.096347

Porth, I., Klapště, J., Skyba, O., Hannemann, J., McKown, A. D., Guy, R. D., DiFazio, S. P., Muchero, W., Ranjan, P., & Tuskan, G. A. (2013). Genome-wide association mapping for wood characteristics in Populus identifies an array of candidate single nucleotide polymorphisms. *New Phytologist, 200*(3), 710-726.

Resende, M. D., Resende, M. F., Sansaloni, C. P., Petroli, C. D., Missiaggia, A. A., Aguiar, A. M., Abad, J. M., Takahashi, E. K., Rosado, A. M., & Faria, D. A. (2012). Genomic selection for growth and wood quality in Eucalyptus: capturing the missing heritability and accelerating breeding for complex traits in forest trees. *New Phytologist, 194*(1), 116-128.

Sewell, M., Bassoni, D., Megraw, R., Wheeler, N., & Neale, D. (2000). Identification of QTLs influencing wood property traits in loblolly pine (Pinus taeda L.). I. Physical wood properties. *TAG Theoretical and Applied Genetics, 101*(8), 1273-1281.

Shiu, S.-H., & Bleecker, A. B. (2001). Plant receptor-like kinase gene family: diversity, function, and signaling. *Sci stke, 113*(113), re22.

Strauss, S., Lande, R., & Namkoong, G. (1992). Limitations of molecular-marker-aided selection in forest tree breeding. *Canadian Journal of Forest Research, 22*(7), 1050-1061.

Strzalka, W., & Ziemienowicz, A. (2010). Proliferating cell nuclear antigen (PCNA): a key factor in DNA replication and cell cycle regulation. *Annals of botany, 107*(7), 1127-1140.

Team, R. (2015). RStudio: integrated development for R. *RStudio, Inc., Boston, MA URL http://www. rstudio. com*.

Thavamanikumar, S., Southerton, S. G., Bossinger, G., & Thumma, B. R. (2013). Dissection of complex traits in forest trees—opportunities for marker-assisted selection. *Tree Genetics & Genomes, 9*(3), 627-639.

Thumma, B. R., Southerton, S. G., Bell, J. C., Owen, J. V., Henery, M. L., & Moran, G. F. (2010).
Quantitative trait locus (QTL) analysis of wood quality traits in Eucalyptus nitens. *Tree*
Genetics & Genomes, 6(2), 305-317.
Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal*
Statistical Society. Series B (Methodological), 267-288.
Vidalis, A., Scofield, D. G., Neves, L. G., Bernhardsson, C., García-Gil, M. R., & Ingvarsson, P.
(2018). Design and evaluation of a large sequence-capture probe set and associated
SNPs for diploid and haploid samples of Norway spruce (Picea abies). *bioRxiv*. doi:
10.1101/291716
Wyatt, M. D., Allan, J. M., Lau, A. Y., Ellenberger, T. E., & Samson, L. D. (1999). 3-
methyladenine DNA glycosylases: structure, function, and biological importance.
Bioessays, 21(8), 668-676.
Yamasaki, K., Kigawa, T., Inoue, M., Tateno, M., Yamasaki, T., Yabuki, T., Aoki, M., Seki, E.,
Matsuda, T., Tomo, Y., Hayami, N., Terada, T., Shirouzu, M., Osanai, T., Tanaka, A.,
Seki, M., Shinozaki, K., & Yokoyama, S. (2004). Solution Structure of the B3 DNA
Binding Domain of the Arabidopsis Cold-Responsive Transcription Factor RAV1. *The*
Plant Cell, 16(12), 3448.
Zhong, R., Burk, D. H., Morrison, W. H., & Ye, Z.-H. (2002). A kinesin-like protein is essential
for oriented deposition of cellulose microfibrils and cell wall strength. *The Plant Cell,*
14(12), 3101-3117.
765

Fig. 1 Outline of the association mapping approach.

Fig 2. PCA plot of all the 517 individuals.