Natural selection on gene-specific codon usage bias is common across
eukaryotes

Zhen Peng[1], Hani Zaher[1], Yehuda Ben-Shahar[1,*]

[1]Department of Biology, Washington University in St. Louis, MO 63130, USA.

[*]Corresponding Author: Dr. Yehuda Ben-Shahar, Department of Biology, Washington

University in St. Louis, MO 63130, USA. benshahary@wustl.edu

Running title: Gene-specific codon usage bias

Keywords: *Drosophila melanogaster*, fruit fly, codon usage bias, synonymous mutation,
neutral mutation

ABSTRACT

Genomic data from diverse taxa indicate that synonymous codon usage frequencies are

often not random. Although the molecular evolutionary forces that shape codon usage

remain poorly understood, synonymous mutations are generally assumed to be neutral

because they do not affect protein sequences. However, empirical evidence indicates

that some synonymous mutations can have phenotypic consequences. Here we show

that in contrast to the current dogma, natural selection on gene-specific codon usage

bias is common across Eukaryota. Furthermore, by using bioinformatic and experimental

approaches, we demonstrate that specific combinations of rare codons contribute to

spatial and sex-related regulation of protein expression in *Drosophila melanogaster*.

These data indicate that natural selection shapes gene-specific codon usage biases,

and therefore, it represents an overlooked genomic feature that plays an important role

in the spatial and temporal regulation of gene functions. Hence, the broadly accepted

dogma that synonymous mutations are generally functionally neutral should be

reevaluated.

INTRODUCTION

Most amino acids are encoded by at least two interchangeable synonymous DNA codons (Nirenberg and Leder, 1964; Crick, 1966; Söll et al., 1966). Although synonymous codons do not impact the amino-acid sequences of proteins, analyses of genomic DNA indicate that protein-coding genes often exhibit codon usage bias, which is defined as usage frequencies of specific synonymous codons that deviate from expected random usage frequencies (Ikemura, 1981; Sharp et al., 1988; Plotkin and Kudla, 2011; Doherty and McInerney, 2013). While the specific patterns of codon usage bias often vary across species, this phenomenon seems to be a universal feature of most protein-coding DNA sequences (Sharp et al., 1988; Kanaya et al., 1999; Wang and Hickey, 2007; Quax et al., 2015; Pouyet et al., 2016). Nevertheless, the actual molecular forces that shape gene-specific codon usage bias at the evolutionary timescale remain poorly understood.

Previous theoretical models derived from population genetics suggested that observed codon usage frequencies are the result of an interplay between three forces: 1) Mutational bias, which refers to the asymmetric occurrence of mutations; 2) Genetic drift, whereby specific synonymous codons are randomly fixed; 3) Natural selection for favored specific synonymous codon usage patterns associated with specific gene functions. This theoretical framework was mathematically formalized as the "selection-mutation-drift model" (Bulmer, 1991; Yang and Nielsen, 2008). Although the selection-mutation-drift model does not assign different mathematical weights to these three forces, it is generally assumed that mutational bias is likely the dominant force in shaping codon usage bias across diverse species (Stenico, Lloyd, and Sharp, 1994; Chen et al., 2004; Fu, 2010; Sharp, Emery, and Zeng, 2010; Rao et al., 2011; Doherty

and McInerney, 2013; Yang et al., 2015). In contrast, whether genetic drift and natural

selection play equally important roles in driving gene-specific codon usage patterns

remains controversial (Akashi, 1995; Rao et al., 2011; Doherty and McInerney, 2013;

Kessler and Dean, 2014). Specifically, since synonymous mutations do not affect protein

sequences, and therefore are usually assumed to have no impact on their structure or

biochemical function, the effect of natural selection on gene-specific codon usage bias is

often dismissed as weak (King and Jukes, 1969; Kimura, 1968; Kimura and Ohta, 1974;

Echave, Spielman, and Wilke, 2016).

Nonetheless, a series of biochemical and molecular studies of specific protein-coding

genes seem to present mechanistic challenges to the broadly accepted dogma that

synonymous mutations are generally functionally neutral. First, gene-specific

synonymous codon usage patterns have been shown to be important for the function of

some protein-coding genes by affecting features such as translational efficacy (Ikemura,

1981; Akashi, 1994; Ko et al., 2005), mRNA secondary structures (Hasegawa,

Yasunaga, and Miyata, 1979), mRNA splicing patterns (Pagani, Raponi, and Baralle,

2005; Parmley and Hurst, 2007), miRNA binding accessibility (Gu et al., 2012), co-

translational protein folding (Zhang, Hubalewska, and Ignatova, 2009; Zhou et al., 2013;

Fu et al., 2016), and post-translational modifications (Kimchi-Sarfaty et al., 2007; Zhou et

al., 2013; Quax et al., 2015; Fu et al., 2016). Second, the temporal regulation of protein

synthesis in several viruses in the *Herpesviridae* group has been shown to depend on

viral-specific codon usage patterns that are different from those of the host cell (Shin et

al., 2015). Third, high protein levels of some housekeeping genes depend on the biased

usage of common synonymous codons (Stenico, Lloyd, and Sharp, 1994; Ma et al.,

2014). Some of these anecdotal examples could be explained by the "translational

selection theory", which stipulates that "optimal" protein translation rate and accuracy

partly depend on the availability and affinity of cognate tRNAs, and therefore,

necessitates the evolution of codon usage patterns that match the relative abundance of

specific tRNAs in the relevant cellular pools (Ikemura, 1981; Shields and Sharp, 1987;

Slimko and Lester, 2003; dos Reis, Savva, and Wernisch, 2004). Together, these

studies suggest that gene-specific codon usage biases might be playing a general, yet

underappreciated, role in the spatial and temporal regulation of gene activity, and

therefore should be affected by natural selection.

Here we combine theoretical and experimental approaches to test the hypothesis that

natural selection has played a significant and important role in shaping gene-specific

codon usage patterns. To test this hypothesis, we developed a statistical approach for

identifying signatures of natural selection on gene-specific codon usage patterns. Using

this approach, we demonstrate that a considerable proportion of protein-coding genes

from diverse eukaryotic species display molecular signatures of natural selection on

biased gene-specific codon usage patterns. Additionally, we use the well-annotated

genome of *Drosophila melanogaster* to identify genome-wide signatures of selection on

gene-specific codon usage patterns, and demonstrate that some combinations of unique

rare synonymous codons are associated with specific biological functions, including

spatial and male-biased protein expression patterns. Together, our results indicate that

in contrast to the broadly accepted dogma, natural selection plays a significant role in

determining gene-specific codon usage biases in diverse eukaryotic species, and that

gene-specific codon usage biases are important for the spatial and temporal regulation

of some protein-coding genes. Together, these data indicate that the broadly accepted

assumption that that synonymous mutations are in general functionally and evolutionarily

neutral should be reevaluated.

RESULTS

Signatures of natural selection on gene-specific codon usage are common across

eukaryotes

Testing the hypothesis that gene-specific codon usage bias plays a general role in

regulating gene function requires a broadly applicable method for statistically identifying

signatures of selection across protein-coding genes throughout genomes.

Mathematically, the previously published "selection-mutation-drift model" could be

theoretically used for identifying such signatures (Bulmer, 1991). However, applying this

specific model to diverse species in order to identify signatures of natural selection on

codon usage bias requires *a priori* empirical information about population-level genetic

variations, temporal and spatial gene expression patterns, and related estimates of

fitness (Yang and Nielsen, 2008; Rodrigue, Philippe and Lartillot, 2010; Tamuri, dos Reis

and Goldstein, 2012; Gilchrist et al., 2015; Kubatko et al. 2016). Since such data are not

available for most species, we developed an alternative statistical approach that is solely

based on the analyses of DNA sequences of protein-coding genes from publicly

available genomic data. In contrast to the "selection-mutation-drift" model, which

evaluates the relative contribution of different molecular forces to the observed codon

usage bias by estimating the selection coefficients for each possible nucleotide

substitution, our approach is based on statistically rejecting the null hypothesis that

'Synonymous mutations are neutral'. While our approach does not provide specific

quantitative estimates for the level of selection on each individual gene, it statisticaly

identifies specific genes whose biased codon usage patterns have been likely impacted

by natural selection.

Our statistical model is based on several key assumptions: 1) Reference genomic data

represent a "wild type" genome; 2) Observed gene-specific codon usage patterns are at

equilibrium; 3) For each codon, no more than one nucleotide substitution per generation

is possible; 4) Nonsynonymous mutations are mostly deleterious, and therefore, the

probabilities of observed nonsynonymous substitutions are negligible relative to those of

synonymous ones; 5) The contributions of individual alleles of a single gene to fitness

are additive; 6) Gene-specific mutational biases are independent, which also means that

the background nucleotide composition of nearby chromosomal regions is negligible

(Oliver et al, 2001); 7) For each individual open reading frame, the mutation rate for each

of the 12 possible nucleotide substitutions (A-to-T, T-to-A, *etc.*) is constant (Rodriguez et

al., 1990), which means that the effects of adjacent nucleotides on the mutation rates of

each focal nucleotide can be ignored (Bérard and Guéguen, 2012); 8) The lowest

mutation rate is at least 1/100 of the highest one (Drake et al., 1998).

Next, we used the previously published "selection-mutation-drift model" (Bulmer, 1991;

Yang and Nielsen, 2008) to describe the relationships between the various evolutionary

molecular processes that may have shaped observed gene-specific codon usage biases

(Equation (1)).

140
$$\sum_i \frac{2s_{pqi \to pqr}}{1-e^{-2Ns_{pqi \to pqr}}} \times x_{pqi} \times \mu_{i \to r} + \sum_j \frac{2s_{jqr \to pqr}}{1-e^{-2Ns_{jqr \to pqr}}} \times x_{jqr} \times \mu_{j \to p}$$
$$= \sum_i \frac{2s_{pqr \to pqi}}{1-e^{-2Ns_{pqr \to pqi}}} \times x_{pqr} \times \mu_{r \to i} + \sum_j \frac{2s_{pqr \to jqr}}{1-e^{-2Ns_{pqr \to jqr}}} \times x_{pqr} \times \mu_{p \to j} \quad (1)$$

Variables p, q, and r denote the specific nucleotide identities (A, C, G, or T) of the first,

second, and third positions respectively in each codon. Codons synonymous to codon

pqr that vary at either the first or third positions are denoted by codons jqr or pqi

respectively. Therefore, $s_{pqi-pqr}$, for example, denotes the selection coefficient of the pqi-

to-pqr mutation. N denotes effective chromosomal population size (Yang and Nielsen,

2008). x_{pqr} denotes the count of codon pqr in a focal gene across the entire population,

and $\mu_{i \to r}$ denotes an estimate for the i-to-r mutation rate. Consequently, the probability

that an i-to-r mutation is fixed in a population is $\frac{2s_{pqi \to pqr}}{1-e^{-2Ns_{pqi \to pqr}}}$ (Yang and Nielsen, 2008).

The amino acid serine represents a unique case because it is encoded by six

synonymous codons that belong to two independent codon groups, which are not

interchangeable via single synonymous mutations. Therefore, in our model we treat the

ser codon groups AGC, AGT and TCA, TCC, TCG, TCT as if they are coding two

independent amino acids (Gilchrist et al., 2015).

Since the null hypothesis of our model specifies that all synonymous mutations are

neutral, the selection coefficient of each synonymous mutation should have a zero value,

which simplifies Equation (1) as follows:

157
$$\sum_i \lim_{s_{pqi \to pqr} \to 0} \frac{2s_{pqi \to pqr}}{1-e^{-2Ns_{pqi \to pqr}}} \times x_{pqi} \times \mu_{i \to r} + \sum_j \lim_{s_{jqr \to pqr} \to 0} \frac{2s_{jqr \to pqr}}{1-e^{-2Ns_{jqr \to pqr}}} \times x_{jqr} \times \mu_{j \to p}$$
$$= \sum_i \lim_{s_{pqr \to pqi} \to 0} \frac{2s_{pqr \to pqi}}{1-e^{-2Ns_{pqr \to pqi}}} \times x_{pqr} \times \mu_{r \to i} + \sum_j \lim_{s_{pqr \to jqr} \to 0} \frac{2s_{pqr \to jqr}}{1-e^{-2Ns_{pqr \to jqr}}} \times x_{pqr} \times \mu_{p \to j}$$

158 $\Rightarrow$

159    $$\sum_i \frac{1}{N} \times x_{pqi} \times \mu_{i \to r} + \sum_j \frac{1}{N} \times x_{jqr} \times \mu_{j \to p} = \sum_i \frac{1}{N} \times x_{pqr} \times \mu_{r \to i} + \sum_j \frac{1}{N} \times x_{pqr} \times \mu_{p \to j} \quad (2)$$

160   Furthermore, since each term in Equation (2) includes the factor $\frac{1}{N}$, Equation (2) could be

161   further simplified as:

162
$$\sum_i x_{pqi} \times \mu_{i \rightarrow r} + \sum_j x_{jqr} \times \mu_{j \rightarrow p} = \sum_i x_{pqr} \times \mu_{r \rightarrow i} + \sum_j x_{pqr} \times \mu_{p \rightarrow j} \quad (3)$$

Subsequently, if the null hypothesis that all synonymous mutations are neutral is true

then the impacts of genetic drift, represented by the effective chromosomal population

size N, and natural selection, represented by selection coefficient s, are canceled as

shown above, and therefore, the observed codon usage pattern could be explained

solely by mutational bias. In contrast, if the null hypothesis is rejected then we must

accept the alternative hypothesis, which indicates natural selection has had a significant

impact on the observed gene-specific codon usage bias.

To apply our derived statistical model to actual genomic data, we initially generated a

dataset of expected values for gene-specific codon usage patterns across all protein-

coding genes in each analyzed genome, assuming they are driven solely by mutational

bias. To achieve this, Equation (3) was first used to estimate gene-specific μ values by

using available reference genomic sequences. We then calculated expected codon

counts for each gene based on the μ values estimated by the model. We then tested

whether the model-generated (expected) and the actual (observed) gene-specific codon

usage patterns are similar by using a χ^2 test (df = 40) (Davis and Olsen, 2009; Xiang et

al., 2015; Bera et al., 2017). Subsequently, all genes with observed codon usage

patterns that were significantly different from the expected patterns generated by the

model were classified as genes that carry signatures of natural selection on synonymous

codon usage. As this approach only requires genetic code and the nucleotide sequences

of open reading frames as input, in principle, it can be applied to any native protein-

coding genes from species with sequenced genomes.

Next, we used our approach to analyze the well-annotated genomes of five

representative eukaryotic species, *Drosophila melanogaster*, *Arabidopsis thaliana*,

Saccharomyces cerevisiae, *Caenorhabditis elegans*, and *Homo sapiens*. Initial analyses

of the overall relative synonymous codon usage (RSCU) (Sharp et al., 1988) revealed

that these species utilize markedly different genome-level patterns of synonymous

codon usage biases (Fig. 1A). Next, we searched for protein-coding genes that exhibit

signatures of selection on gene-specific codon usage bias by utilizing our new statistical

approach. We found that after correcting for false discovery rate (FDR = 0.05)

(Benjamini and Hochberg, 1995), the percentage of protein-coding genes that carry

signatures of selection on gene-specific synonymous codon usage bias varied between

14% and 64% across these five representative genomes (Fig. 1B). Additional analyses

of available eukaryotic genomes revealed that in 35 out of 40 species analyzed, at least

10% of the protein-coding genes carried signatures of natural selection on synonymous

codon usage patterns, independent of whether they are unicellular or multicellular (Fig.

S1). These results indicate that the overall impact of natural selection on gene-specific

patterns of codon usage is broader than would be expected if synonymous substitutions

were mostly functionally neutral. Our results also suggest that the relative impact of

selection on gene-specific codon usage bias is not constant across the eukaryotic

phylogeny, and therefore, might be the result of diverse, clade-specific selective forces.

As currently framed, our quantitative approach is based on a set of conservative

assumptions, which should make the model applicable to as many genomes as possible.

However, we acknowledge that some of these assumptions might not represent the

correct biological context for all species or ecological conditions. We also anticipate that

the assumption that adjacent nucleotides have no effect on the mutation rates of focal

nucleotides could lead to increased false-positive rates (Bérard and Guéguen, 2012).

However, we anticipate that this possible source for false-positives is compensated for

by the assumption that background nucleotide composition of nearby chromosomal

regions is constant (Oliver et al, 2001), which is further alleviated by the conservative

use of relaxed constraints for estimating μ values (Drake et al., 1998). Nevertheless,

false positive rates produced by our model could be further reduced by having more

precise estimates of the μ parameters, which could be achieved by using population-

level genetic variations data, and more specific information about species-specific

mutation rates, and how they might be impacted by the identities of adjacent nucleotides.

Selection on gene-specific codon usage bias in *D. melanogaster* is associated

with specific biological functions

Our statistical approach for identifying genes that carry signatures of natural selection on

synonymous codon usage cannot directly address which specific synonymous codons

might be under selection in each selected gene. Therefore, we next used a clustering

approach to determine whether the genes we have identified in our initial screen share

similar codon usage patterns, or alternatively, represent a heterogeneous population

comprised of multiple gene clusters, each defined by a unique pattern of biased codon

usage.

Due to its well-annotated genome and wealth of available genetic and phenotypic data,

we chose to use the genetically tractable species *D. melanogaster* for functional

analyses of genes that are under selection for biased codon usage patterns. Hierarchical

gene-clustering analysis revealed that genes with identified signatures of selection on

biased codon usage could be assigned to one of two major clusters (Clusters A and B;

Fig. 2). Further analysis of Cluster A revealed that synonymous codon usage in these

genes is enriched for the rare codons Lys-AAA, Glu-GAA, Gln-CAA, Phe-TTT, Tyr-TAT,

and His-CAT, relative to their usage in Cluster B genes. Similar clustering analyses of

the genomes of *A. thaliana*, *S. cerevisiae*, *C. elegans*, and *H. sapiens* revealed gene-by-

codon clustering in all five species, likely representing species-specific patterns of

selection (Figs. S2-5).

Next, we tested the hypothesis that the identified gene clusters in *D. melanogaster*

represent specific biological functions. Analyses of previously published tissue-specific

gene expression data (Chintapalli, Wang, and Dow, 2007) revealed that genes that are

preferentially expressed in the male accessory glands are significantly overrepresented

in Cluster A and underrepresented in Cluster B (Table 1). Similarly, analysis of

previously published data on sexually dimorphic genes (Graveley et al., 2010; Gelbart

and Emmert, 2013) revealed that genes with male-enriched expression are significantly

overrepresented in Cluster A and underrepresented in Cluster B (Table 2). These data

suggest that a specific combination of rare codons contributes to the function of some

genes associated with the male reproductive system.

In addition, analyses of Gene Ontology terms (Gene Ontology Consortium, 2015)

revealed that genes that encode extracellular matrix proteins are overrepresented, and

genes that encode cytoplasmic proteins are underrepresented, in Cluster A (Table S1).

We also found that genes annotated as encoding odorant-binding proteins, a class of

secreted proteins, as well as extracellular space proteins, are underrepresented in

Cluster B. Together, these analyses suggest that in *D.* melanogaster, natural selection

on the increased usage of specific combinations of rare synonymous codons might be

important for male-specific functions of some genes.

Biased gene-specific codon usage could modulate spatial protein expression

patterns

One possible explanation for the association between certain biased gene-specific

codon usage patterns and tissue-specific expression is that the selected specific rare

synonymous codons match tissue-specific tRNA pools, and therefore, enhances the

translational efficiency of some protein in specific tissues or cell types (Zalucki, Gittins

and Jennings, 2008; Pechmann, Chartron and Frydman, 2014). Similar associations

between gene-specific codon usage bias and organ-enriched gene expression patterns

have been previously reported in humans (Plotkin, Robins, and Levine, 2004; Dittmar,

Goodenbour and Pan, 2006). Therefore, we next tested the hypothesis that the specific

pattern of codon usage bias exhibited by Cluster A genes contributes to increased

translation efficiency in the male reproductive system.

Our clustering analysis revealed that male-enriched genes in Cluster A preferentially use

the rare codons Lys-AAA, Gln-CAA, Glu-GAA, Phe-TTT, Tyr-TAT, and His-CAT (Fig. 2).

However, only the first three codons are recognized by exactly matched tRNA

anticodons, while the other three rare codons share the same tRNAs with their more

commonly used synonymous codons, with a mismatch (wobble) at the third codon

position (Crick, 1966; Chan and Lowe, 2009), which suggests these codons are less

likely to contribute to tissue-specific translation efficiency. Therefore, we next analyzed

the specific usage of rare codons AAA, CAA, and GAA in all genes that show enriched

expression in the male reproductive system, independent of whether these genes have

passed the initial statistical threshold for the detection of selection on overall codon

usage patterns. We found that these specific rare synonymous codons are indeed

overrepresented in male-reproductive-system-specific genes (Table 3). Thus, not only

are genes preferring AAA, CAA, and GAA more likely to be male-reproductive-system-

specific, but also genes specifically transcribed in male reproductive system use AAA,

CAA, and GAA more often than their expected genome average usage.

Selection on the biased usage of codons AAA, CAA, and GAA in associate with male-

specific gene activity should depend, at least in part, on the increased expression of their

cognate tRNAs in the relevant tissues. In accordance with this, northern blot analyses of

tissue-specific tRNA expression revealed that $tRNA^{Lys}_{TTT}$ is enriched in the male

reproductive organs, further supporting the hypothesis that spatial regulation of some

tRNA genes that correspond to rare codons could contribute to tissue-specific increase

in protein translation efficiency (Fig. 3A-C). The expression patterns of the tRNAs that

match Glu-GAA and Gln-CAA were not investigated because they are too similar to their

common tRNA counterparts to allow detection by hybridization probes (de Muro, 2008).

Nevertheless, these data indicate that some *Drosophila* genes have evolved a specific

codon usage bias that specifically drives their levels and/or activity in the male

reproductive system.

Next, we empirically tested whether specific combinations of rare codons are sufficient to

drive tissue-specific translational efficiency by generating transgenic flies that express

different alleles of the EGFP reporter under the control of the UAS-GAL4 binary

expression system (Brand and Perrimon, 1993). One allele was comprised of common

codons for all amino acids (EGFP*Common*), and the second allele was comprised of rare

codons AAA, GAA, and CAA for amino acids Lys, Glu, and Gln respectively

(EGFP^{RareKEQ}), which represent about 18% of the total residues in EGFP. Each EGFP

allele was also co-expressed with an RFP reporter encoded by common codons, which

served as a transgene expression control. Both transgenes were driven by the

ubiquitous *Act*5*C*-GAL4 driver (Kopp, Blackman, and Duncan, 1999). To evaluate the

effect of codon usage bias on the spatial pattern of both EGFP alleles *in vivo*, we

compared the two genotypes by measuring the ratios of RFP-normalized EGFP signals

in the accessory gland secondary cells (AGSC) and in the hindgut proliferation zone

(HPZ) (Takashima et al., 2013) in the pylorus region. We found that the signal from the

EGFP^{RareKEQ} allele in the AGSCs was significantly higher than that of the *EGFP*^{Common}

allele. In contrast, both alleles produced similar signals in the HPZ (Fig. 3D-I). Because

the overall transcript levels of both alleles are similar (Fig. 3J), these results suggest that

some specific combinations of synonymous rare codons could impact the spatial

regulation of protein translation in the *Drosophila* male reproductive system.

DISCUSSION

In contrast to the broadly accepted dogma in evolutionary biology and population

genetics, our theoretical and empirical studies indicate that synonymous mutations

should not be considered neutral in general. By using multiple complementary

approaches, we demonstrate here that the impact of natural selection on gene-specific

codon usage bias is common across the eukaryotic phylogeny, and is likely to play a

much more prominent role in defining gene-specific codon usage bias than was

previously assumed. Furthermore, our data indicate that in multicellular organisms such

as the fruit fly, natural selection is likely to optimize codon usage patterns to match

specific spatial and temporal constraints on gene function, and therefore, what might

represent an optimal codon usage for individual genes does not necessarily mean

common codons should be expected at every position. Our findings also indicate that

previous reports about the phenotypic consequences of synonymous mutations are not

anecdotal rare cases. Instead, these cases likely represent a fundamental biological

principle associated with the spatial and temporal gene activity landscapes in eukaryotes.

In addition, our analyses of gene-specific codon usage biases in *D. melanogaster*

suggest that gene-specific codon usage bias is heterogeneous across protein coding

genes in a single species, and therefore, likely to be the outcome of diverse selective

pressures associated with different biological functions. Furthermore, our data suggest

that different classes of rare synonymous codons exhibit differential sensitivities to

natural selection.

Consequently, our discovery that gene-specific codon usage patterns are often shaped

by natural selection indicates that alternative synonymous codons serve important, yet

often overlooked, biological functions. Although several previous studies have used

statistical approaches to suggest that synonymous mutations in specific genes and

species might not be neutral (Berger, 1977; Eyre-Walker, 1991; Stenico, Lloyd, and

Sharp, 1994; Lawrie et al., 2013), our current study provides the first genome-scale

comprehensive view of the overall impact of natural selection on gene-specific

synonymous codon usage patterns across diverse eukaryotic taxa. Thus, our unbiased

and broadly applicable statistical approach has uncovered a fundamental genomic

organizational principle that is sensitive to natural selection, and likely represents an

important component of the genotype-phenotype axis in diverse eukaryotic species and

biological contexts. Therefore, it is likely that investigations, which rely on the

assumption that synonymous mutations are neutral, often overestimate or underestimate

the prevalence of selection on specific protein-coding genes, which could impact the

interpretations of the role of these genes in genotype-phenotype relationships at the

level of populations and the evolutionary timescale in particular (Chen et al., 2010), and

the genome evolution in general (Campos et al., 2012; Matsumoto et al., 2016).

Specifically, the prevailing assumption that most synonymous mutations are neutral, and

therefore represent a robust and quantitative measure of selection-independent

evolutionary rates of nucleotide substitutions, needs to be revised. This is especially

important in the context of the commonly used tests for identifying molecular signatures

of selection in specific protein-coding genes, and the classification of modes of natural

selection (*e.g.*, "positive selection" vs. "negative selection") (McDonald and Kreitman,

1991; Nielsen, 2001; Echave, Spielman, and Wilke, 2016).

Our data also indicate that, similarly to *Drosophila*, many human protein-coding genes

carry signatures of selection on their biased codon usage patterns (Fig. 1). This finding

could have broad implications for studies of genetic variants underlying quantitative

phenotypes in human populations. Consequently, assuming that synonymous mutations

are functionally neutral could lead to a biased view that associations between

nonsynonymous SNPs and specific phenotypic values are the only ones important for

understanding overall trait variations (Hampe et al., 2007; WTCCC and TASC, 2007; Xu

and Taylor, 2009; Fong et al., 2010; Huang et al., 2012). This is supported by a recent

data survey of human GWAS data from over 2000 independent studies, which

suggested that synonymous and nonsynonymous SNPs have similar likelihoods and

effect sizes in terms of association with disease phenotypes (Chen et al., 2010).

Although our analysis of human gene-specific codon usage bias was based on data from

a single reference genome, the simplest interpretation of our results is that both

synonymous and nonsynonymous mutations are likely influenced by natural selection in

human populations, and therefore, many synonymous SNPs could still have a significant

quantitative and/or qualitative impacts on gene functions and associated phenotypes in

health and disease.

We also found that gene-specific codon usage bias is not uniform across the genome.

Instead, our data suggest that different groups of genes in different species often exhibit

enrichment for specific combinations of rare or common codons, which suggests that

different combinations of synonymous codons may have been selected for independent

biological reasons. In this regard, we show that at least one group of genes with

enriched expression in the reproductive system of male *Drosophila* preferentially use the

rare codons Lys-AAA, Gln-CAA, and Glu-GAA. By using allelic variants of a reporter

gene, we show that the selective usage of these specific three rare codons is sufficient

for generating spatially biased protein expression patterns. These findings are consistent

with some of the assumptions about the possible roles of codon usage bias, first put

forward by the "translational selection theory" (Ikemura, 1981; dos Reis, Savva, and

Wernisch, 2004; Plotkin, Robins, and Levine, 2004; Dittmar, Goodenbour, and Pan,

2006). Consequently, our study provides a broader context to this fundamental

evolutionary theory by emphasizing the possible role of gene-specific codon usage bias

in the spatial regulation of proteins in multicellular eukaryotes. However, regulation of

spatial protein expression cannot explain all cases of selection on gene-specific codon

usage bias because genomes of some unicellular eukaryotes such as the

choanoflagellate *Monosiga brevicollis* and the green algae *Chlamydomonas reinhardtii*

also include a significant proportion of protein-coding genes exhibit signatures of

selection on their codon usage patterns (Fig. S1).

Our studies also have important practical implications for experimental systems that

depend on the use of heterologous transgenic gene expression. We found that for some

genes, the selective usage of synonymous rare codons could increase their translational

efficacy in specific tissues and cell types, and therefore, designing experiments with the

assumption that transgene codon optimization necessitates the usage of only common

synonymous codons could lead to unexpected phenotypic outcomes, and could affect

data interpretation, especially for genes with restricted spatial and/ or temporal

expression patterns.

Our studies also highlight the need for a better understanding of the mechanisms that

regulate the spatial expression patterns of tRNAs. Some previous studies have argued

that tRNA gene copy number is likely the primary mechanism that regulates the relative

levels of individual tRNAs in cellular pools (dos Reis, Savva and Wernisch, 2004). Yet,

variations in gene copy number alone cannot explain the tissue-specific expression

patterns of some rare tRNAs observed by us (Fig. 3A-C) and others (Dittmar,

Goodenbour and Pan, 2006; Gingold et al., 2014). Therefore, although tRNAs are

thought to be exclusively transcribed by the constitutively-active RNA Pol III complex,

there must be additional molecular mechanisms that enable the temporal and spatial

regulation of some tRNAs at the transcriptional and/ or posttranscriptional levels

(Yoshihisa et al., 2007; Wichtowska, Turowski, and Boguta, 2013; Turowski and

Tollervey, 2016).

Although our empirical work here has mostly focused on a specific set of rare codons

that bind unique tRNA anti-codons, it should be noted that genes in Cluster A (Fig. 2)

also preferentially use the rare codons Phe-TTT, Tyr-TAT, and His-CAT. These rare

synonymous codons share their tRNA anticodons with their more common synonymous

codons via a wobble mechanism (Chan and Lowe, 2009). Theoretical models and

empirical data suggest that this specific class of rare codons is likely to slowdown

translation due to the lower affinity of common tRNAs to the rare synonymous codons

(Ikemura, 1981; dos Reis, Savva, and Wernisch, 2004; Savir and Tlusty, 2013; Brule

and Grayhack, 2017). Therefore, we speculate that in contrast to rare codons that bind

unique perfectly-matching tRNAs, rare codons that share tRNAs with their more

common synonymous codons may have evolved to broadly restrict translational

efficiency of some low-abundance proteins, or proteins that require slow folding kinetics

associated with specific post-translational modifications, or the binding of other protein

and small-molecule partners, chaperones, or co-factors (Komar, Lesnik and Reiss, 1999;

Weygand-Durasevic and Ibba, 2010; Angov, 2011; Zhou et al., 2013; Fluman et al.,

2014; Pechmann, Chartron and Frydman, 2014; Chaney and Clark, 2015; Quax et al.,

2015; Yu et al., 2015; Fu et al., 2016; Zhao, Yu, and Liu, 2017).

In summary, the results presented here suggest that in contrast to a widely accepted

dogma, synonymous mutations should not be considered functionally and evolutionarily

neutral by default. Instead, as was previously shown for a few specific protein-coding

genes (Hasegawa, Yasunaga, and Miyata, 1979; Ikemura, 1981; Akashi, 1994; Stenico,

Lloyd, and Sharp, 1994; Ko et al., 2005; Pagani, Raponi, and Baralle, 2005; Kimchi-

Sarfaty et al., 2007; Parmley and Hurst, 2007; Zhang, Hubalewska, and Ignatova, 2009;

Gu et al., 2012; Zhou et al., 2013; Ma et al., 2014; Shin et al., 2015; Fu et al., 2016), our

data indicate that selection on synonymous codon usage is likely to serve a more

general function in diverse biological processes across Eukaryotes. Nevertheless,

although our data indicate that 'synonymous' and 'neutral' mutations are not necessarily

the same, we do not suggest that the overall concept of evolutionary neutrality, first

described by Darwin in the *On the Origin of Species* as "*Variations neither useful nor

injurious would not be affected by natural selection*", should be rejected. Instead, we

argue that the commonly accepted specific assumption that all synonymous mutations

are generally neutral should be reevaluated.

ACKNOWLEDGEMENTS

Z.P. dedicates this manuscript to Dr. Yang Zhong of Fudan University, a prominent

botanist and evolutionary biologist who has had a great influence on his choice to pursue

a research career in mathematical and evolutionary biology. The authors also wish to

thank Alan Templeton, Allan Larson, and members of the Ben-Shahar and Zaher

laboratories for helpful discussions. They also thank Paula Kiefel, Dianne Duncan, and

Ian Duncan for technical assistance with generating transgenic animals and microscopic

imaging. Z.P. is supported by the training grant 2T32HG000045 from the NIH. This

research has been supported in part by NIH grant NS089834, and NSF grants 1545778

and 1707221 to YB, and NIH grant 5R01GM112641 to HZ.

AUTHOR CONTRIBUTIONS

ZP and YB conceived the study; ZP performed the studies under the guidance of YB and
HZ; HZ assisted with the design and execution of molecular biology studies; ZP, HZ, and
YB wrote the manuscript and approved its final version.

REFERENCES

Akashi, H. (1994). Synonymous codon usage in Drosophila melanogaster: natural

selection and translational accuracy. *Genetics, 136*(3), 927-935.

Akashi, H. (1995). Inferring weak selection from patterns of polymorphism and

divergence at" silent" sites in Drosophila DNA. *Genetics, 139*(2), 1067-1076.

Angov, E. (2011). Codon usage: nature's roadmap to expression and folding of

proteins. *Biotechnology journal, 6*(6), 650-659.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical

and powerful approach to multiple testing. *Journal of the royal statistical society.Series B*

(Methodological), 289-300.

Bera, B. C., Virmani, N., Kumar, N., Anand, T., Pavulraj, S., Rash, A., Elton, D., Rash, N.,

Bhatia, S., Sood, R., Singh, R. K., & Singh, R. K. (2017).Genetic and codon usage bias

analyses of polymerase genes of equine influenza virus and its relation to evolution.*BMC*

genomics, 18(1), 652.

Bérard, J., &Guéguen, L. (2012). Accurate estimation of substitution rates with neighbor-

dependent models in a phylogenetic context. *Systematic biology, 61*(3), 510-521.

Berger, E. M. (1977). Are Synonymous Mutations Adaptively Neutral?.*The American*

Naturalist, 111(979), 606-607.

Bischof, J., Maeda, R. K., Hediger, M., Karch, F., & Basler, K. (2007). An optimized

transgenesis system for Drosophila using germ-line-specific φC31 integrases.

Proceedings of the National Academy of Sciences, 104(9), 3312-3317.

Brand, A. H., &Perrimon, N. (1993). Targeted gene expression as a means of altering

cell fates and generating dominant phenotypes. *Development, 118*(2), 401-415.

Brown, A., Shao, S., Murray, J., Hegde, R. S., &Ramakrishnan, V. (2015).Structural

basis for stop codon recognition in eukaryotes. *Nature*, *524*(7566), 493-496.

Brule, C. E., &Grayhack, E. J. (2017). Synonymous codons: choose wisely for

expression. *Trends in Genetics, 33*(4), 283-297.

Bulmer, M. (1991). The selection-mutation-drift theory of synonymous codon

usage. *Genetics*, *129*(3), 897-907.

Campos, J. L., Zeng, K., Parker, D. J., Charlesworth, B., &Haddrill, P. R. (2012). Codon

usage bias and effective population sizes on the X chromosome versus the autosomes

in Drosophila melanogaster.*Molecular biology and evolution, 30*(4), 811-823.

Chan, P. P., & Lowe, T. M. (2009). GtRNAdb: a database of transfer RNA genes

detected in genomic sequence. *Nucleic acids research, 37*(suppl 1), D93-D97.

Chaney, J. L., & Clark, P. L. (2015). Roles for synonymous codon usage in protein

biogenesis. *Annual review of biophysics*, *44*, 143-166.

Chen, R., Davydov, E. V., Sirota, M., & Butte, A. J. (2010). Non-synonymous and

synonymous coding SNPs show similar likelihood and effect size of human disease

association. *PloS one, 5*(10), e13574.

Chen, S. L., Lee, W., Hottes, A. K., Shapiro, L., & McAdams, H. H. (2004). Codon usage

between genomes is constrained by genome-wide mutational processes. *Proceedings of*

the National Academy of Sciences of the United States of America, 101(10), 3480-3485.

Chintapalli, V. R., Wang, J., & Dow, J. A. (2007). Using FlyAtlas to identify better

Drosophila melanogaster models of human disease. *Nature genetics, 39*(6), 715-720.

Crick, F. H. (1966). Codon—anticodon pairing: the wobble hypothesis. *Journal of*

molecular biology, 19(2), 548-555.

Davis, J. J., & Olsen, G. J. (2009). Modal codon usage: assessing the typical codon

usage of a genome. *Molecular biology and evolution, 27*(4), 800-810.

de Muro, M. A. (2008). Probe design, production, and applications. In Molecular

Biomethods Handbook, J. M. Walker & R. Rapleyeds. (Totowa, NJ, U.S.A.: Humana

Press. 2nd edition), pp. 41-53.

Dittmar, K. A., Goodenbour, J. M., & Pan, T. (2006). Tissue-specific differences in

human transfer RNA expression. *PLoS Genet, 2*(12), e221.

Doherty, A., &McInerney, J. O. (2013). Translational selection frequently overcomes

genetic drift in shaping synonymous codon usage patterns in vertebrates. *Molecular*

biology and evolution, 30(10), 2263-2267.

dos Reis, M., Savva, R., &Wernisch, L. (2004). Solving the riddle of codon usage

preferences: a test for translational selection. *Nucleic acids research, 32*(17), 5036-5044.

Drake, J. W., Charlesworth, B., Charlesworth, D., & Crow, J. F. (1998). Rates of

spontaneous mutation. *Genetics, 148*(4), 1667-1686.

Echave, J., Spielman, S. J., &Wilke, C. O. (2016).Causes of evolutionary rate variation

among protein sites.*Nature Reviews. Genetics, 17*(2), 109.

Eyre-Walker, A. C. (1991). An analysis of codon usage in mammals: selection or

mutation bias?.*Journal of molecular evolution, 33*(5), 442-449.

Fluman, N., Navon, S., Bibi, E., &Pilpel, Y. (2014).mRNA-programmed translation

pauses in the targeting of E. coli membrane proteins. *Elife*, *3*, e03440.

Fong, C., Ko, D. C., Wasnick, M., Radey, M., Miller, S. I., &Brittnacher, M. (2010).

GWAS analyzer: integrating genotype, phenotype and public annotation data for

genome-wide association study analysis. *Bioinformatics, 26*(4), 560-564.

Fu, M. (2010). Codon usage bias in herpesvirus.*Archives of Virology, 155*(3), 391-396.

Fu, J., Murphy, K. A., Zhou, M., Li, Y. H., Lam, V. H., Tabuloc, C. A., Chiu, J. C., & Liu, Y.

(2016). Codon usage affects the structure and function of the Drosophila circadian clock

protein PERIOD. *Genes & Development*, *30*(15), 1761-1775.

Gelbart, W. M., &Emmert, D. B. (2013). FlyBase High Throughput Expression Pattern

Data

Gene Ontology Consortium. (2015). Gene ontology consortium: going forward. *Nucleic

acids research*, *43*(D1), D1049-D1056.

Gilchrist, M. A., Chen, W. C., Shah, P., Landerer, C. L., &Zaretzki, R. (2015). Estimating

Gene Expression and Codon-Specific Translational Efficiencies, Mutation Biases, and

Selection Coefficients from Genomic Data Alone. *Genome biology and evolution*, *7*(6),

1559-1579.

Gingold, H., Tehler, D., Christoffersen, N. R., Nielsen, M. M., Asmar, F., Kooistra, S. M.,

Christophersen, N. S., Christensen, L. L., Borre, M., Sørensen, K. D., Andersen, L. D.,

Andersen, C. L., Hulleman, E., Wurdinger, T., Ralfkiær, E., Helin, K., Grønbæk, K.,

Ørntoft, T., Waszak, S. M., Dahan, O., Pedersen, J. S., Lund, A. H.,&Pilpel, Y. (2014). A

dual program for translation regulation in cellular proliferation and differentiation.*Cell,*

158(6), 1281-1292.

Graveley, B.R., Brooks, A.N., Carlson, J.W., Cherbas, L., Choi, J., Davis, C.A., Dobin, A.,

Duff, M., Eads, B., Hansen, K.D., Landolin, J., Langton, L., Malone, J., Miller, D.,

Roberts, J., Sandler, J., Sturgill, D., Tang, H., van Baren, M.J., Wan, K.H., Xiao, S.,

Yang, L., Zhang, D., Zhang, Y., Zou, Y., Andrews, J., Brenner, S.E., Brent, M., Cherbas,

P., Dudoit, S., Gingeras, T.R., Hoskins, R., Kaufman, T., Oliver, B, &Celniker, S.E.

(2010.03.15). The *D. melanogaster*transcriptome: modENCODE RNA-Seq data

Gu, W., Wang, X., Zhai, C., Xie, X., & Zhou, T. (2012).Selection on synonymous sites for

increased accessibility around miRNA binding sites in plants.*Molecular biology and*

evolution, 29(10), 3037-3044.

Hampe, J., Franke, A., Rosenstiel, P., Till, A., Teuber, M., Huse, K., Albrecht, M., Mayr,

G., De La Vega, F. M., Briggs, J., Günther, S., Prescott, N. J., Onnie, C. M., Häsler, R.,

Sipos, B., Fölsch, U. R., Lengauer, T., Platzer, M., Mathew, C. G., Krawczak, M., &

Schreiber, S. (2007). A genome-wide association scan of nonsynonymous SNPs

identifies a susceptibility variant for Crohn disease in ATG16L1. *Nature genetics, 39*(2),

207.

Hasegawa, M., Yasunaga, T., & Miyata, T. (1979).Secondary structure of MS2 phage

RNA and bias in code word usage.*Nucleic acids research, 7*(7), 2073-2079.

Hill, A., Zheng, X., Li, X., McKinney, R., Dickman, D., & Ben-Shahar, Y. (2017). The

Drosophila postsynaptic DEG/ENaC channel ppk29 contributes to excitatory

neurotransmission. *Journal of Neuroscience, 37*(12), 3171-3180.

Huang, X., Kurata, N., Wang, Z. X., Wang, A., Zhao, Q., Zhao, Y., Liu, K., Lu, H., Li, W.,

Guo, Y., Lu, Y., Zhou, C., Fan, D., Weng, Q., Zhu, C., Huang, T., Zhang, L., Wang, Y.,

Feng, L., Furuumi, H., Kubo, T., Miyabayashi, T., Yuan, X., Xu, Q., Dong, G., Zhan, Q.,

Li, C., Fujiyama, A., Toyoda, A., Lu, T., Feng, Q., Qian, Q., Li, J., & Han, B. (2012). A

map of rice genome variation reveals the origin of cultivated rice. *Nature, 490*(7421), 497.

Ikemura, T. (1981). Correlation between the abundance of *Escherichia coli* transfer

RNAs and the occurrence of the respective codons in its protein genes: a proposal for a

synonymous codon choice that is optimal for the *E. coli* translational system. *Journal of*

molecular biology, 151(3), 389-409.

Kanaya, S., Yamada, Y., Kudo, Y., &Ikemura, T. (1999). Studies of codon usage and

tRNA genes of 18 unicellular organisms and quantification of *Bacillus subtilis*tRNAs:

gene expression level and species-specific diversity of codon usage based on

multivariate analysis. *Gene, 238*(1), 143-155.

Kessler, M. D., & Dean, M. D. (2014). Effective population size does not predict codon

usage bias in mammals. *Ecology and evolution, 4*(20), 3887-3900.

Kimchi-Sarfaty, C., Oh, J. M., Kim, I. W., Sauna, Z. E., Calcagno, A. M., Ambudkar, S. V.,

&Gottesman, M. M. (2007). A" silent" polymorphism in the MDR1 gene changes

substrate specificity. *Science, 315*(5811), 525-528.

Kimura, M. (1968). Evolutionary rate at the molecular level.*Nature, 217*(5129), 624-626.

Kimura, M., &Ohta, T. (1974).On some principles governing molecular

evolution.*Proceedings of the National Academy of Sciences, 71*(7), 2848-2852.

King, J. L., & Jukes, T. H. (1969). Non-darwinian evolution. *Science, 164*(3881), 788-798.

Ko, H. J., Ko, S. Y., Kim, Y. J., Lee, E. G., Cho, S. N., & Kang, C. Y. (2005).

Optimization of codon usage enhances the immunogenicity of a DNA vaccine encoding

mycobacterial antigen Ag85B. *Infection and immunity, 73*(9), 5666-5674.

Komar, A. A., Lesnik, T., & Reiss, C. (1999). Synonymous codon substitutions affect

ribosome traffic and protein folding during in vitro translation. *FEBS letters*, *462*(3), 387-

391.

Kopp, A., Blackman, R. K., & Duncan, I. (1999). Wingless, decapentaplegic and EGF

receptor signaling pathways interact to specify dorso-ventral pattern in the adult

abdomen of Drosophila. *Development, 126*(16), 3495-3507.

Kraft, D. (1988).A software package for sequential quadratic programming. Technical

report DFVLR-FB. InstitutfürDynamik der Flugsysteme; Oberpfaffenhofen, pp. 88–28.

Kubatko, L., Shah, P., Herbei, R., & Gilchrist, M. A. (2016). A codon model of nucleotide

substitution with selection on synonymous codon usage. *Molecular Phylogenetics and*

Evolution, *94*, 290-297.

Lawrie, D. S., Messer, P. W., Hershberg, R., &Petrov, D. A. (2013). Strong purifying

selection at synonymous sites in *D. melanogaster.PLoS genetics, 9*(5), e1003527.

Lu, B., LaMora, A., Sun, Y., Welsh, M. J., & Ben-Shahar, Y. (2012). ppk23-Dependent

chemosensory functions contribute to courtship behavior in Drosophila melanogaster.

PLoS Genet, *8*(3), e1002587.

Ma, L., Cui, P., Zhu, J., Zhang, Z., & Zhang, Z. (2014). Translational selection in human:

more pronounced in housekeeping genes. *Biology direct*, 9(1), 17.

Matsumoto, T., John, A., Baeza-Centurion, P., Li, B., & Akashi, H. (2016). Codon usage

selection can bias estimation of the fraction of adaptive amino acid fixations. *Molecular*

biology and evolution, 33(6), 1580-1589.

McDonald, J. H., &Kreitman, M. (1991). Adaptive protein evolution at the Adh locus in

Drosophila.*Nature, 351*(6328), 652-654.

Meurer A., Smith C. P., Paprocki M., Čertík O., Kirpichev S. B., Rocklin M., Kumar A.,

Ivanov S., Moore J. K., Singh S., Rathnayake T., Vig S., Granger B. E., Muller R. P.,

Bonazzi F., Gupta H., Vats S., Johansson F., Pedregosa F., Curry M. J., Terrel A. R.,

Roučka Š., Saboo A., Fernando I., Kulal S., Cimrman R., Scopatz A. (2017). SymPy:

symbolic computing in Python. *PeerJ Computer Science, 3*, e103.

Nielsen, R. (2001). Statistical tests of selective neutrality in the age of

genomics. *Heredity, 86*(6), 641-647.

Nirenberg, M., &Leder, P. (1964). RNA codewords and protein synthesis.*Science,*

145(3639), 1399-1407.

Oliver, J. L., Bernaola-Galván, P., Carpena, P., &Román-Roldán, R. (2001). Isochore

chromosome maps of eukaryotic genomes.*Gene, 276*(1), 47-56.

Pagani, F., Raponi, M., &Baralle, F. E. (2005). Synonymous mutations in CFTR exon 12

affect splicing and are not neutral in evolution. *Proceedings of the National Academy of*

Sciences of the United States of America, *102*(18), 6368-6372.

Parmley, J. L., & Hurst, L. D. (2007). Exonic splicing regulatory elements skew

synonymous codon usage near intron-exon boundaries in mammals. *Molecular biology*

and evolution, 24(8), 1600-1603.

Pechmann, S., Chartron, J. W., &Frydman, J. (2014). Local slowdown of translation by

nonoptimal codons promotes nascent-chain recognition by SRP in vivo. *Nature structural*

& molecular biology, 21(12), 1100-1105.

Plotkin, J. B., Robins, H., & Levine, A. J. (2004). Tissue-specific codon usage and the

expression of human genes. *Proceedings of the National Academy of Sciences of the*

United States of America, *101*(34), 12588-12591.

Plotkin, J. B., &Kudla, G. (2011). Synonymous but not the same: the causes and

consequences of codon bias. *Nature reviews. Genetics, 12*(1), 32.

Pouyet, F., Bailly-Bechet, M., Mouchiroud, D., &Guéguen, L. (2016). SENCA: A

Multilayered Codon Model to Study the Origins and Dynamics of Codon Usage. *Genome*

Biology and Evolution, *8*(8), 2427-2441.

Quax, T. E., Claassens, N. J., Söll, D., & van der Oost, J. (2015). Codon bias as a

means to fine-tune gene expression. *Molecular cell*, *59*(2), 149-161.

Rao, Y., Wu, G., Wang, Z., Chai, X., Nie, Q., & Zhang, X. (2011). Mutation bias is the

driving force of codon usage in the *Gallus gallus* genome. *DNA research, 18*(6), 499-512.

Rodrigue, N., Philippe, H., &Lartillot, N. (2010). Mutation-selection models of coding

sequence evolution with site-heterogeneous amino acid fitness profiles. *Proceedings of*

the National Academy of Sciences, *107*(10), 4629-4634.

Rodriguez, F. J. L. O. J., Oliver, J. L., Marin, A., & Medina, J. R. (1990). The general

stochastic model of nucleotide substitution.*Journal of theoretical biology, 142*(4), 485-

501.

Savir, Y., &Tlusty, T. (2013). The ribosome as an optimal decoder: A lesson in molecular

recognition. *Cell, 153*(2), 471-479.

Sharp, P. M., Cowe, E., Higgins, D. G., Shields, D. C., Wolfe, K. H., & Wright, F. (1988).

Codon usage patterns in *Escherichia coli*, *Bacillus subtilis*, *Saccharomyces cerevisiae*,

Schizosaccharomycespombe, *Drosophila melanogaster* and *Homo sapiens*; a review of

the considerable within-species diversity. *Nucleic acids research*, *16*(17), 8207-8211.

Sharp, P. M., Emery, L. R., &Zeng, K. (2010).Forces that influence the evolution of

codon bias.*Philosophical Transactions of the Royal Society of London B: Biological*

Sciences, 365(1544), 1203-1212.

Shields, D. C., & Sharp, P. M. (1987). Synonymous codon usage in *Bacillus subtilis*

reflects both translational selection and mutational biases. *Nucleic Acids Research,*

15(19), 8023-8040.

Shin, Y. C., Bischof, G. F., Lauer, W. A., &Desrosiers, R. C. (2015). Importance of codon

usage for the temporal regulation of viral gene expression.*Proceedings of the National*

Academy of Sciences, 112(45), 14030-14035.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,

T.,Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,

Hartenstein, V., Eliceiri, K., &Tomancak, P. (2012). Fiji: an open-source platform for

biological-image analysis. *Nature methods, 9*(7), 676.

Slimko, E. M., & Lester, H. A. (2003). Codon optimization of

CaenorhabditiselegansGluCl ion channel genes for mammalian cells dramatically

improves expression levels. *Journal of neuroscience methods*, *124*(1), 75-81.

Söll, D., Jones, D. S., Ohtsuka, E., Faulkner, R. D., Lohrmann, R., Hayatsu, H., Khorana,

H.G., Cherayil, J.D., Hampel, A., & Bock, R. M. (1966). Specificity of sRNA for

recognition of codons as studied by the ribosomal binding technique. *Journal of*

molecular biology, *19*(2), 556-573.

Stenico, M., Lloyd, A. T., & Sharp, P. M. (1994). Codon usage in *Caenorhabditiselegans*:

delineation of translational selection and mutational biases. *Nucleic Acids*

Research, *22*(13), 2437-2446.

Takashima, S., Paul, M., Aghajanian, P., Younossi-Hartenstein, A., & Hartenstein, V.

(2013). Migration of Drosophila intestinal stem cells across organ boundaries.

Development, 140(9), 1903-1911.

Tamuri, A. U., dos Reis, M., & Goldstein, R. A. (2012). Estimating the distribution of

selection coefficients from phylogenetic data using sitewise mutation-selection

models. *Genetics*, *190*(3), 1101-1115.

Turowski, T. W., &Tollervey, D. (2016). Transcription by RNA polymerase III: insights

into mechanism and regulation. *Biochemical Society Transactions, 44*(5), 1367-1375.

Wang, H. C., & Hickey, D. A. (2007). Rapid divergence of codon usage patterns within

the rice genome. *BMC evolutionary biology, 7*(1), S6.

Warnes, G. R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W. H. A., & Lumley, T.

(2015). gplots: Various R Programming Tools for Plotting Data. R package version

3.0.1.*https://cran.r-project.org/web/packages/gplots/gplots.pdf.*

Wellcome Trust Case Control Consortium (WTCCC) & The Australo-Anglo-American

Spondylitis Consortium (TASC). (2007). Association scan of 14,500 nsSNPs in four

common diseases identifies variants involved in autoimmunity. *Nature genetics, 39*(11),

1329.

Weygand-Durasevic, I., &Ibba, M. (2010).New roles for codon

usage. *Science, 329*(5998), 1473-1474.

Wichtowska, D., Turowski, T. W., &Boguta, M. (2013). An interplay between transcription,

processing, and degradation determines tRNA levels in yeast. *Wiley Interdisciplinary*

Reviews: RNA, 4(6), 709-722.

Xiang, H., Zhang, R., Butler III, R. R., Liu, T., Zhang, L., Pombert, J. F., & Zhou, Z.

(2015). Comparative Analysis of Codon Usage Bias Patterns in Microsporidian

Genomes. *PloS one, 10*(6), e0129223.

Xu, Z., & Taylor, J. A. (2009). SNPinfo: integrating GWAS and candidate gene

information into functional SNP selection for genetic association studies. *Nucleic acids*

research, 37(suppl_2), W600-W605.

Yang, Z., & Nielsen, R. (2008).Mutation-selection models of codon substitution and their

use to estimate selective strengths on codon usage. *Molecular biology and*

evolution, *25*(3), 568-579.

Yang, X., Ma, X., Luo, X., Ling, H., Zhang, X., &Cai, X. (2015). Codon usage bias and

determining forces in Taeniasolium genome. *The Korean journal of parasitology, 53*(6),

689.

Yoshihisa, T., Ohshima, C., Yunoki-Esaki, K., & Endo, T. (2007).Cytoplasmic splicing of

tRNA in Saccharomyces cerevisiae.*Genes to Cells, 12*(3), 285-297.

Yu, C. H., Dang, Y., Zhou, Z., Wu, C., Zhao, F., Sachs, M. S., & Liu, Y. (2015). Codon

usage influences the local rate of translation elongation to regulate co-translational

protein folding. *Molecular cell, 59*(5), 744-754.

Zalucki, Y. M., Gittins, K. L., & Jennings, M. P. (2008). Secretory signal sequence non-

optimal codons are required for expression and export of β-lactamase. *Biochemical and*

biophysical research communications, 366(1), 135-141.

Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M.R., Barrell, D., Bhai, J., Billis, K.,

Cummins, C., Gall, A., Girón, C.G., Gil, L., Gordon, L., Haggerty, L., Haskell, E., Hourlier,

T., Izuogu, O.G., Janacek, S.H., Juettemann, T., To, J.K., Laird, M.R., Lavidas, I., Liu, Z.,

Loveland, J.E., Maurel, T., McLaren, W., Moore, B., Mudge, J., Murphy, D.N., Newman,

V., Nuhn, M., Ogeh, D., Ong, C.K., Parker, A., Patricio, M., S., H., Schuilenburg, H.,

Sheppard, D., Sparrow, H., Taylor, K., Thormann, A., Vullo, A., Walts, B., Zadissa, A.,

Frankish, A., Hunt, S.E., Kostadima, M., Langridge, N., Martin, F.J., Muffato, M., Perry,

E., Ruffier, M., Staines, D.M., Trevanion, S.J., Aken, B.L., Cunningham, F., Yates, A.,

&Flicek, P. (2017). Ensembl 2018. *Nucleic acids research, 46*(D1), D754-D761.

Zhang, G., Hubalewska, M., &Ignatova, Z. (2009). Transient ribosomal attenuation

coordinates protein synthesis and co-translational folding. *Nature structural & molecular*

biology, 16(3), 274-280.

Zhao, F., Yu, C. H., & Liu, Y. (2017). Codon usage regulates protein structure and

function by affecting translation elongation speed in Drosophila cells. *Nucleic Acids*

Research.45(14), 8484-8492.

Zhou, M., Guo, J., Cha, J., Chae, M., Chen, S., Barral, J. M., Sachs, M. S., & Liu, Y.

(2013). Non-optimal codon usage affects expression, structure and function of clock

protein FRQ. *Nature, 495*(7439), 111-115.

 FIGURE LEGENDS

Figure 1. Signatures of natural selection on gene-specific synonymous codon

usage bias in representative eukaryotic species. (A) Analyses of the relative

synonymous codon usage (RSCU) as a measure of the overall codon usage bias across

five representative Eukaryotic species. (B) Total number of protein-coding genes

analyzed for each species were: *D. melanogaster*, 12099; *A. thaliana*, 23694;

S.cerevisiae, 5366; *C. elegans*, 17210; *H. sapiens*, 20261. The estimated percentages

of genes carrying selection signatures are corrected by a 0.05 false discovery rate. See

also Figure S1.

Figure 2. Clustered codon usage patterns across genes carrying signatures of

selection on gene-specific codon usage bias in *D. melanogaster*. Hierarchical

clustering of all genes (horizontal axis) identified as carrying significant signatures of

selection on gene-specific codon usage bias. The relative usage frequencies of each

codon in each gene are color-coded (vertical axis). Euclidean distances were used to

measure dissimilarities between codons. Spearman's correlation coefficients were used

to measure similarities between genes. Complete linkage was used as the clustering

criterion. Clusters A (red bar) and B (black bar) represent the two major gene groups

identified. See also Figure S2-5, Table S1.

Figure 3. Gene-specific codon usage bias affects spatial protein expression in *D.*

***melanogaster*.** (A) Representative northern blots and (B-C) summary data of the

relative abundance of tRNA$^{Lys}_{CTT}$ (common) and tRNA$^{Lys}_{TTT}$ (rare) codons across male

tissues. *, $p<0.05$ (n = 4, ANOVA followed by SNK *post hoc* tests). Error bars denote

standard deviation. †: Reproductive system is excluded. (D-E) Representative images

showing EGFP and RFP expression in HPZ (yellow lines marks tissue boundaries). (F-G)

Representative images showing EGFP and RFP expression in the AGSC (yellow lines

marks secondary cells). (H-I) Summary data of normalized EGFP signals in HPZ and

AGSC. *, $p<0.01$ (n=5, unpaired Student's *t*-test). Error bars denote standard deviation.

(J) Real-time qRT-PCR mRNA expression data of $EGFP^{Common}$ and $EGFP^{RareKEQ}$

(unpaired Student's *t*-test, n = 4, NS). Error bars denote SEM.

TABLES

Tissue	Cluster A		Cluster B	
	Fold enrichment	p-value	Fold enrichment	p-value
Adult Central Nervous System			1.13	0.040
Midgut			1.19	0.003
Testis			0.56	<0.001
Male Accessory Gland	4.45	<0.001	0.36	<0.001
Carcass			0.34	<0.001
Salivary Gland			0.49	0.024
Eye			0.80	0.050

785   **Table 1. Tissue-specific expression patterns of genes in Clusters A and B.** Genes

786   included in Clusters A and B are as in Fig. 2. Only tissues with a significant over- or

787   underrepresentation in at least one of the two clusters are shown. Fold enrichment is

788   relative to the entire genome.

Gene cluster	Sex	Fold enrichment	p-value
Cluster A	Male	1.66	<0.001
	Female	1.67	0.091
Cluster B	Male	0.61	<0.001
	Female	1.07	0.060

790 **Table 2. Sex-biased expression patterns of genes in Clusters A and B.** Genes

791 included are as in Table 1. Fold enrichment is relative to the entire genome.

Codons	Count of codons in male-reproductive-organ-specific genes	Count of codons in all analyzed genes	Fold enrichment in male-reproductive-organ-specific genes	p-value
Lys-AAA	12936	117591	1.18	<0.001
Lys-AAG	22139	258907	0.92	
Gln-CAA	10181	108912	1.26	<0.001
Gln-CAG	15727	241374	0.88	
Glu-GAA	16093	151650	1.22	<0.001
Glu-GAG	21843	285332	0.88	

Table 3. Usage frequencies of rare codons Lys-AAA, Gln-CAA, and Glu-GAA in

male-specific tissues relative to whole genome averages.

Counts of relevant codons were calculated for the entire genome and male-reproductive-

organ-specific genes (including genes with enriched expression in the testis and the

accessory-glands). Hypergeometric tests were used to test whether a specific

synonymous codon is over- or under-represented in male-reproductive-organ-specific

genes. Fold enrichment is calculated by dividing the relative frequency of a focal codon

in male-reproductive-organ-specific genes by the relative frequency of the focal codon

usage across all protein-coding genes.

METHODS

Genomic and transcriptomic data. Protein-coding DNA sequences were from

Ensembl (Zerbino et al., 2017) (https://www.ensembl.org/). Reference coding sequences

included in the analysis of gene-specific codon usage bias were chosen according to the

following criteria: 1) The sequence length is a multiple of three. 2) The sequence uses

standard genetic code. 3) For each gene, only the longest mRNA isoform was used for

analysis. If there were multiple isoforms of the same length, then the first record shown

in the fasta file was used. 4) The encoded protein includes all 19 amino acids that have

degenerate codons; for the amino acid serine, the two-fold and four-fold degenerate

codon groups were treated as if they encoded two different amino acids. Transcriptomic

data were from the FlyAtlas microarray database (Chintapalli, Wang, and Dow, 2007)

and the modENCODE RNA-seq database (Graveley et al., 2010; Gelbart and Emmert,

2013).

Estimating μ and expected codon counts. The relationships between μ values and

codon counts are described by Equation (3) (Results section). Based on the standard

genetic code and all possible combinations for synonymous nucleotide substitutions, we

classified all degenerate codons into six categories. For each category, we used

Equation (3) to generate a homogeneous linear equation system. Subsequently, we

treated one of the x variables as a known parameter and analytically solved all other x

variables. Finally, these solutions were used to calculate the expected codon counts of a

protein-coding gene.

Category one – Codons for Arginine:

827
$$
\begin{cases}
x_{AGA} \times \mu_{A \to C} + x_{AGA} \times \mu_{A \to G} = x_{CGA} \times \mu_{C \to A} + x_{AGG} \times \mu_{G \to A} \\
x_{AGG} \times \mu_{A \to C} + x_{AGG} \times \mu_{G \to A} = x_{CGG} \times \mu_{C \to A} + x_{AGA} \times \mu_{A \to G} \\
x_{CGA} \times \mu_{C \to A} + x_{CGA} \times (\mu_{A \to C} + \mu_{A \to G} + \mu_{A \to T}) \\
\qquad = x_{AGA} \times \mu_{A \to C} + x_{CGC} \times \mu_{C \to A} + x_{CGG} \times \mu_{G \to A} + x_{CGT} \times \mu_{T \to A} \\
x_{CGG} \times \mu_{C \to A} + x_{CGG} \times (\mu_{G \to A} + \mu_{G \to C} + \mu_{G \to T}) \\
\qquad = x_{AGG} \times \mu_{A \to C} + x_{CGA} \times \mu_{A \to G} + x_{CGC} \times \mu_{C \to G} + x_{CGT} \times \mu_{T \to G} \\
x_{CGC} \times (\mu_{C \to A} + \mu_{C \to G} + \mu_{C \to T}) = x_{CGA} \times \mu_{A \to C} + x_{CGG} \times \mu_{G \to C} + x_{CGT} \times \mu_{T \to C} \\
x_{CGT} \times (\mu_{T \to A} + \mu_{T \to C} + \mu_{T \to G}) = x_{CGA} \times \mu_{A \to T} + x_{CGC} \times \mu_{C \to T} + x_{CGG} \times \mu_{G \to T}
\end{cases}. \quad (4)
$$

828   Category two – Codons for Leucine:

829
$$
\begin{cases}
x_{TTA} \times \mu_{T \to C} + x_{TTA} \times \mu_{A \to G} = x_{CTA} \times \mu_{C \to T} + x_{TTG} \times \mu_{G \to A} \\
x_{TTG} \times \mu_{T \to C} + x_{TTG} \times \mu_{G \to A} = x_{CTG} \times \mu_{C \to T} + x_{TTA} \times \mu_{A \to G} \\
x_{CTA} \times \mu_{C \to T} + x_{CTA} \times (\mu_{A \to C} + \mu_{A \to G} + \mu_{A \to T}) \\
\qquad = x_{TTA} \times \mu_{T \to C} + x_{CTC} \times \mu_{C \to A} + x_{CTG} \times \mu_{G \to A} + x_{CTT} \times \mu_{T \to A} \\
x_{CTG} \times \mu_{C \to T} + x_{CTG} \times (\mu_{G \to A} + \mu_{G \to C} + \mu_{G \to T}) \\
\qquad = x_{TTG} \times \mu_{T \to C} + x_{CTA} \times \mu_{A \to G} + x_{CTC} \times \mu_{C \to G} + x_{CTT} \times \mu_{T \to G} \\
x_{CTC} \times (\mu_{C \to A} + \mu_{C \to G} + \mu_{C \to T}) = x_{CTA} \times \mu_{A \to C} + x_{CTG} \times \mu_{G \to C} + x_{CTT} \times \mu_{T \to C} \\
x_{CTT} \times (\mu_{T \to A} + \mu_{T \to C} + \mu_{T \to G}) = x_{CTA} \times \mu_{A \to T} + x_{CTC} \times \mu_{C \to T} + x_{CTG} \times \mu_{G \to T}
\end{cases}. \quad (5)
$$

830   Category three – Four-fold degenerate codons:

831
$$
\begin{cases}
x_{pqA} \times (\mu_{A \to C} + \mu_{A \to G} + \mu_{A \to T}) = x_{pqC} \times \mu_{C \to A} + x_{pqG} \times \mu_{G \to A} + x_{pqT} \times \mu_{T \to A} \\
x_{pqC} \times (\mu_{C \to A} + \mu_{C \to G} + \mu_{C \to T}) = x_{pqA} \times \mu_{A \to C} + x_{pqG} \times \mu_{G \to C} + x_{pqT} \times \mu_{T \to C} \\
x_{pqG} \times (\mu_{G \to A} + \mu_{G \to C} + \mu_{G \to T}) = x_{pqA} \times \mu_{A \to G} + x_{pqC} \times \mu_{C \to G} + x_{pqT} \times \mu_{T \to G} \\
x_{pqT} \times (\mu_{T \to A} + \mu_{T \to C} + \mu_{T \to G}) = x_{pqA} \times \mu_{A \to T} + x_{pqC} \times \mu_{C \to T} + x_{pqG} \times \mu_{G \to T}
\end{cases}. \quad (6)
$$

832   Category four – Codons for Isoleucine:

833
$$
\begin{cases}
x_{ATA} \times (\mu_{A \to C} + \mu_{A \to T}) = x_{ATC} \times \mu_{C \to A} + x_{ATT} \times \mu_{T \to A} \\
x_{ATC} \times (\mu_{C \to A} + \mu_{C \to T}) = x_{ATA} \times \mu_{A \to C} + x_{ATT} \times \mu_{T \to C} \\
x_{ATT} \times (\mu_{T \to A} + \mu_{T \to C}) = x_{ATA} \times \mu_{A \to T} + x_{ATC} \times \mu_{C \to T}
\end{cases}. \quad (7)
$$

834   Category five – C/T-ended two-fold degenerate codons:

835
$$
x_{pqC} \times \mu_{C \to T} = x_{pqT} \times \mu_{T \to C}. \quad (8)
$$

836   Category six– A/G-ended two-fold degenerate codons:

837
$$
x_{pqA} \times \mu_{A \to G} = x_{pqG} \times \mu_{G \to A}. \quad (9)
$$

838   Information about the analytical solutions of the above equations is shown in the

839   Supplemental Information (Analytical Solutions of Equation Systems Used to Estimate μ

840   Values). Using these solutions, with a given set of μ values and counts of amino acid

841   residues, we can calculate the expected counts of synonymous codons. For example,

842   the expected counts of codons for Lys can be calculated by

843
$$\begin{cases} Ex_{AAA} = y_{Lys} \times \dfrac{x_{AAA}}{x_{AAA}+x_{AAG}} = y_{Lys} \times \dfrac{x_{AAA}}{x_{AAA}+x_{AAA}\times\frac{\mu_{A\to G}}{\mu_{G\to A}}} = y_{Lys} \times \dfrac{1}{1+\frac{\mu_{A\to G}}{\mu_{G\to A}}} \\ Ex_{AAG} = y_{Lys} \times \dfrac{x_{AAG}}{x_{AAA}+x_{AAG}} = y_{Lys} \times \dfrac{x_{AAA}\times\frac{\mu_{A\to G}}{\mu_{G\to A}}}{x_{AAA}+x_{AAA}\times\frac{\mu_{A\to G}}{\mu_{G\to A}}} = y_{Lys} \times \dfrac{\frac{\mu_{A\to G}}{\mu_{G\to A}}}{1+\frac{\mu_{A\to G}}{\mu_{G\to A}}} \end{cases} , (10)$$

where Ex_{AAA} is the expected count of AAA codon, and y_{Lys} is the count of Lys residues.

Then we can use the expected counts and the observed real counts of all degenerate

codons to calculate a χ^2 value. Since for a given protein-coding sequence, the χ^2 value

is a function of μ values, we can define this function as $\chi^2(\Theta)$, where Θ is a vector

describing all μ values,

$$\Theta = (\mu_{A\to C}, \mu_{C\to A}, \mu_{A\to G}, \mu_{G\to A}, \mu_{A\to T}, \mu_{T\to A}, \mu_{C\to G}, \mu_{G\to C}, \mu_{C\to T}, \mu_{T\to C}, \mu_{G\to T}, \mu_{T\to G}). (11)$$

For a gene, to estimate μ values, we try to minimize the value of $\chi^2(\Theta)$ by changing the

elements in Θ, using the sequential least squares programming (SLSQP) algorithm

(Kraft, 1988).

Since equations generated from Equation (3) form a system of homogeneous linear

equations, it is meaningless to infer exact μ values from the minimization of $\chi^2(\Theta)$; rather,

only the relative magnitudes of different μ values are important for calculating the

expected codon counts later. As we used the assumption that the lowest μ value is at

least 1/100 of the highest, we set the range of μ values between 0.001 and 0.1 during

the minimization of $\chi^2(\Theta)$. For each gene, the minimum $\chi^2(\Theta)$ is used to calculate p-value.

Since we assumed that reference genomic data represent a "wild type" genome, the

procedure mentioned above was applied to each individual reference gene.

Codon usage heatmaps

For a protein-coding gene g, the relative usage frequency f_{gd} of a codon d is defined as

$$f_{gd} = \frac{n_{gd}}{y_{ga}}, \text{(12)}$$

where n_{gd} is the count of d in g, and y_{ga} is the count of amino acid a encoded by d and its

synonymous codons in g. It should be noted that codons for Ser are not treated as two

codon groups in codon usage heatmaps. As the mechanism of recognizing stop codons

is fairly different from recognizing other codons (Brown et al., 2015), and methionine and

tryptophan respectively have only one codon, the analysis is restricted to the other 59

codons. Therefore, a 59-dimension vector B_g is used to describe the codon usage

pattern of g,

$$B_g = \left(f_{g1}, f_{g2}, f_{g3}, \cdots, f_{gd}, \cdots, f_{g59}\right)^T. \text{(13)}$$

For a genome containing M protein-coding genes, an M-dimension vector H_d is used to

describe how often a codon d is used across these genes,

$$H_d = \left(f_{1d}, f_{2d}, f_{3d}, \cdots, f_{gd}, \cdots, f_{Md}\right). \text{(14)}$$

Both B_g's and H_d's are hierarchically clustered. The values of all f_{gd}'s are then color-

coded to generate a codon usage heatmap.

The "gplots" (Warnes et al., 2015) package in R was used to generate heatmaps. The

method of hierarchical clustering was complete linkage with Euclidean distance

measuring the dissimilarities between codons and Spearman's correlation coefficient

measuring the similarities between genes.

Identifying genes with tissue-specific expression patterns

Mean adult gene expression data from the FlyAtlas database were used to identify

genes with tissue-specific expression patterns. A gene was classified as tissue-specific if

its average mRNA level in a specific tissue was at least ten-fold to the tissue with the

second highest mRNA level. To reduce redundancy, "head" expression values were

excluded from the analysis. Specifically, since "brain" and "thoracic-abdominal ganglion"

both belong to the central nervous system (CNS) and many CNS-specific genes are

expressed in both places, "adult central nervous system"-specific mRNA level is

represented by the higher mRNA level between "brain" and "thoracic-abdominal

ganglion". When "adult central nervous system" was involved in the analysis, the "brain"

and "thoracic-abdominal ganglion" were excluded, and *vice versa*.

Identifying genes with sex-biased expression patterns

Adult fly expression data from the modENCODE RNA-seq database were used to

identify genes with either male- or female-biased expression patterns. Only genes that

showed at least 10-fold higher expression in one sex relative to the other were defined

as sex-biased.

Hypergeometric tests for gene set enrichment

The hypergeometric tests were performed using the online tool at

http://www.rothsteinlab.com/tools/apps/hyper_geometric_calculator.

Gene ontology analysis

Gene ontology (GO) analysis (Gene Ontology Consortium, 2015) of *Drosophila* genes

was performed by using online tools (http://geneontology.org/). Category enrichments

were determined by comparing term frequencies between each gene cluster and the

whole genome, followed by a Bonferroni correction.

Analyses of tRNA gene expression analyses

Northern blots were used to measure relative tRNA abundance in different body parts.

Total RNA was extracted from four pools of 10 dissected male tissues (head,

reproductive system, and remaining thorax and abdominal parts) and 10 whole male

flies with the TRIzol reagent (Invitrogen Catalog # 15596-026). Probe sequences were:

$tRNA^{Lys}_{CTT}$, 5'-AACGTGGGGCTCGAACCCACGACCCTGA-3'; $tRNA^{Lys}_{TTT}$, 5'-

GAACAGGGACTTGAACCCTGGACCCTTG-3'. Probes were labeled by ^{32}P using T4

PNK (NEB Catalog # M0201S). Signals were measured and normalized to total tRNA

signals using the BIO-RAD Quantity One 1-D analysis software. ANOVA followed by

SNK *post hoc* test was used to compare tRNA levels between samples.

Animals

Fruit flies (*D. melanogaster*) were raised on corn syrup-soy food (Archon Scientific) at

25°C and 60% relative humidity with a 12-hour light/dark cycle. Synthetic cDNAs

encoding EGFP and mCherry fluorescent proteins were from IDT Inc. Full sequences of

mCherry^{Common}, *EGFP^{Common}*, and *EGFP^{RareKEQ}* alleles are in the Supplemental

Information (Sequences of Fluorescent Reporter Genes). To generate animals that

express each transgenic allele under UAS control, synthetic cDNAs were cloned into the

EcoRI/NotI cloning sites of the pUASTattB plasmid (Bischof et al., 2007). Since the

EGFP^{RareKEQ} allele contains one EcoRI site in the middle, digestion time was shortened

to less than 20 minutes to allow incomplete digestion. The UAS-*RFP^{Common}* transgene

was inserted into a chromosome II landing site (*y^1 M{vas-int.Dm}ZH-2A w*; M{3xP3-*

RFP.attP}ZH-51D; Bloomington #24483), and the UAS-*EGFP^{Common}* and UAS-

EGFP^{RareKEQ} transgenes were inserted into the same chromosome III landing site

(*y^{1}M{vas-int.Dm}ZH-2A w*; M{3xP3-RFP.attP}ZH-86Fb*; Bloomington #24749). Double

homozygotic lines UAS-*RFP^{Common}*; UAS-*EGFP^{Common}* and UAS-*RFP^{Common}*; UAS-

EGFP^{RareKEQ} were generated and then crossed to *Act*5C-GAL4*/SM*6 (Duncan lab,

Washington University).

Real-time qRT-PCR

The mRNA expression levels of reporter genes in whole four-day-old male flies were

quantified by using real-time qRT-PCR, following previously published methods (Lu et al.,

2012; Hill et al., 2017). For *RpL32*, the forward primer was 5'-

CACCAAGCACTTCATCCG-3', and the reverse primer was 5'-

TCGATCCGTAACCGATGT-3'. For *EGFP^Common^* and *EGFP^RareKEQ^*, the forward primers

were respectively 5'-AACTTCAAGATCCGCCACAAC-3' and 5'-

AACTTCAAAATCCGCCACAAC-3', while these alleles shared the same reverse primer

5'-GTGCTCAGGTAGTGGTTATCG-3'.

Quantitative reporter gene imaging

Male reproductive and gut tissues from four-day-old adult male flies that express either

the *EGFP^Common^* or *EGFP^RareKEQ^* were dissected in chilled PBS and mounted for imaging

on a Nikon A1Si laser scanning confocal microscope with a 20X oil objective (n=5

samples per genotype). All images were taken within 10 minutes of dissection. Single

plane fluorescent images of the AGSC were used to estimate EGFP expression levels of

each allele. Similar images of the HPZ were used as generic tissue controls. The NIS-

Element Ar software was used to capture EGFP and RFP signals and generate channel-

merged images. Normalized EGFP signals from each image were quantified using the

Fiji image processing software (Schindelin et al., 2012). The tissue-specific effect of the

EGFP codon usage was analyzed by comparing the normalized EGFP signals in either

the AGSC or HPZ between genotypes with an unpaired Student's *t*-test.

A

B

A

tRNA$^{Lys}_{CTT}$

tRNA$^{Lys}_{TTT}$

Total tRNA

Head   Reproductive system   Thorax and abdomen   Whole body

B

tRNA$^{Lys}_{CTT}$

Relative Abundance of tRNA

*

C

tRNA$^{Lys}_{TTT}$

*   *   *

Head   Reproductive system   Thorax and abdomen   Whole body

Act5C > RFP^{Common}, EGFP^{Common}

D   50 μm

F   50 μm

Act5C > RFP^{Common}, EGFP^{RareKEQ}

E   50 μm

G   50 μm

H

Normalized EGFP Signal of HPZ

I

Normalized EGFP Signal of AGSC

*

Act5C > RFP^{Common}, EGFP^{Common}

Act5C > RFP^{Common}, EGFP^{RareKEQ}

J

Relative Abundance of EGFP mRNA

Act5C > RFP^{Common}, EGFP^{Common}

Act5C > RFP^{Common}, EGFP^{RareKEQ}