1 **Many rice genes are differentially spliced between roots and shoots but cytokinin has**
2 **minimal effect on splicing**
3

Running title: Differential splicing in rice

Nowlan H. Freese[1*], April R. Estrada[1*], Ivory C. Blakley[1], Jinjie Duan[2], Ann E. Loraine[1**]

[1]Department of Bioinformatics and Genomics, University of North Carolina at Charlotte,

Charlotte, NC, USA

[2]Bioinformatics Research Center, Aarhus University, C.F. Moellers Alle 8, Aarhus C, Denmark.

[*]Equal contribution.

[**]Corresponding author.

Nowlan Freese: nowlan.freese@gmail.com

April R. Estrada: april.estrada@duke-energy.com

Ivory C. Blakley: ieclabau@uncc.edu

Jinjie Duan: jjduan.birc@gmail.com

Ann E. Loraine: ann.loraine@uncc.edu

Corresponding author phone: 704-687-0144

ABSTRACT

Alternatively spliced genes produce multiple spliced isoforms, called transcript variants. In

differential alternative splicing, transcript variant abundance differs across sample types.

Differential alternative splicing is common in animal systems and influences cellular

development in many processes, but its extent and significance is not as well known in plants. To

investigate alternative splicing in plants, we examined RNA-Seq data from rice seedlings. The

data included three biological replicates per sample type, approximately 30 million sequence

alignments per replicate, and four sample types: roots and shoots treated with exogenous

cytokinin delivered hydroponically or a mock treatment. Cytokinin treatment triggered

expression changes in thousands of genes but had negligible effect on splicing patterns.

However, many genes were alternatively spliced between mock-treated roots and shoots,

indicating that our methods were sufficiently sensitive to detect differential splicing in a data set.

Quantitative fragment analysis of reverse transcriptase-PCR products made from newly prepared

rice samples confirmed nine of ten differential splicing events between rice roots and shoots.

Differential alternative splicing typically changed the relative abundance of splice variants that

co-occurred in a data set. Analysis of a similar (but less deeply sequenced) RNA-Seq data set

from *Arabidopsis* showed the same pattern. In both the *Arabidopsis* and rice RNA-Seq data sets,

most genes annotated as alternatively spliced had small minor variant frequencies. Of splicing

choices with abundant support for minor forms, most alternative splicing events were located

within the protein-coding sequence and maintained the annotated reading frame. A tool for

visualizing protein annotations in the context of genomic sequence (ProtAnnot) together with a

genome browser (Integrated Genome Browser) were used to visualize and assess effects of

differential splicing on gene function. In general, differentially spliced regions coincided with

conserved regions in the encoded proteins, indicating that differential alternative splicing is

likely to affect protein function between root and shoot tissue in rice.

Key words: rice, alternative splicing, cytokinin, root, shoot, RNA-Seq

Abbreviations: SR - Serine/Arginine, BA - benzyladenine, L - long, S - short, FDR - false
discovery rate, PSI - percent spliced in, AS - acceptor site, DS - donor site, ES - exon skipping,
RI - retained intron, ABA - abscisic acid, IGB - Integrated Genome Browser, CGE - capillary gel
electrophoresis, RPKM - Reads Per Kilobase per Million mapped reads

INTRODUCTION

Differential splicing of pre-mRNA transcripts, called alternative splicing, enables one gene to produce multiple transcript variants encoding different functions. Alternative splicing is an almost universal phenomenon in higher eukaryotes, occurring to varying degrees in every animal and plant genome examined to date (Kalsotra and Cooper, 2011; Reddy *et al.*, 2013). In animals, differential expression of splice variants has been recruited as a regulatory mechanism in multiple processes, such as sex determination in invertebrates and neuronal differentiation in mammals (Kalsotra and Cooper, 2011; Salz, 2011; Barbosa-Morais *et al.*, 2012).

In plants, less is known about the functional significance and patterns of alternative splicing. However, several trends are apparent. Genes involved in circadian regulation are highly alternatively spliced, often producing multiple splice variants that fluctuate in concert with day/night cycling along with overall transcript abundance (Filichkin *et al.*, 2015b). The serine and arginine-rich (SR) family of RNA-binding, splicing regulatory proteins is greatly expanded compared to mammals and includes many plant-specific forms (Kalyna and Barta, 2004; Barbosa-Morais *et al.*, 2006; Plass *et al.*, 2008; Filichkin *et al.*, 2015a). SR transcripts themselves are also highly alternatively spliced in plants, with the relative abundance of these alternative transcripts varying according to environmental stresses and hormones (Palusa *et al.*, 2007; Gulledge *et al.*, 2012; Filichkin *et al.*, 2015a; Keller *et al.*, 2016; Mei *et al.*, 2017).

A growing body of evidence indicates that cell and tissue specific regulation of alternative splicing occurs in plants, but its significance and extent is not well established (Vitulo *et al.*, 2014; Li *et al.*, 2016a; Sun *et al.*, 2018). We previously found through analysis of RNA-Seq data from *Arabidopsis* pollen that the relative abundance of splice variants was similar between leaves and pollen, despite the differences between the two tissues (Loraine *et al.*, 2013). However, this latter analysis was limited by having just one biological replicate for pollen and only two biological replicates for leaves. A more comprehensive analysis of multiple *Arabidopsis* data sets found a high incidence of isoform switching, in which the identity of the most prevalent variant differs between sample types (Vaneechoutte *et al.*, 2017). However, this splicing diversity may have arisen in part from the heterogeneity of the data sets used, which were produced using rapidly changing (and improving) sequencing technologies at different times by different groups.

In this study, we used a well-replicated RNA-Seq data set from rice to re-examine prevalence

of alternative splicing between tissues and hormone (cytokinin) treatment. This data set was

previously generated to investigate cytokinin regulation of gene expression in roots and shoots

from 10-day old rice seedlings (Raines *et al.*, 2016). A parallel study produced an analogous data

set from *Arabidopsis* for comparison, but was less deeply sequenced (Zubo *et al.*, 2017). Both

the rice and *Arabidopsis* RNA-Seq data sets included three biological replicates per sample type

and four sample types – roots and shoots treated with exogenous cytokinin or a mock, vehicle-

only treatment. In both data sets, the treatment triggered differential expression of thousands of

genes, with roots affected to a greater degree than shoots.

For most alternatively spliced genes, regardless of whether or not they were differentially

spliced, the relative abundance of splicing forms was highly skewed, with most alternatively

spliced genes producing one major isoform. Nonetheless, there was a large minority of

alternatively spliced genes where minor isoforms were more abundant and therefore seemed

likely to affect gene function. We found that the relative abundance of splice variants for most

alternatively spliced genes was remarkably stable, with very few differentially spliced genes

between cytokinin treated and control samples. By contrast, many more genes were differentially

spliced between roots and shoot, and most differential splicing occurred within the protein-

coding sequence. Moreover, nearly every differential splicing event detected merely change the

relative abundance of splice variants that co-occurred in the same sample. These results suggest

differential alternative splicing likely contributes to gene function diversification between roots

and shoots by moderating the relative abundance of splice co-expressed splice variants, but

alternative splicing plays little role in cytokinin signaling.

MATERIALS & METHODS

RNA-Seq library preparation and sequencing

Rice and *Arabidopsis* samples were prepared and sequenced as described in (Raines *et al.*,

2016; Zubo *et al.*, 2017). Rice seedlings (Nipponbare) were grown hydroponically for ten days in

a growth chamber set to 14 hours light (28°C) and 8 hours of dark (23°C) with light intensity 700

mmol m^{-2} s^{-1}. Around six to ten seedlings were grown in the same pot, in four pots. On the tenth

day, culture media was replaced with new media containing 5 μM of the cytokinin

benzyladenine (BA) or 0.05 mN NaOH as a control. After 120 minutes, roots and shoots were

harvested separately. Roots and shoots from treatment or control pots were pooled to form three

replicates per treatment. RNA was extracted and used to synthesize twelve libraries from BA-

treated and mock-treated roots and shoots. Libraries were sequenced on an Illumina HiSeq

instrument for 100 cycles, yielding 100 base, single end reads. Sequence data are available from

the Sequence Read Archive under accession SRP04905. Aligned, processed data are available

from the Oct. 2011 rice genome assembly IgbQuickload directorie at

http://igbquickload.org/quickload/O_sativa_japonica_Oct_2011/.

Arabidopsis plants were grown vertically on 1× Murashige and Skoog agar with 1% sucrose

for ten days under continuous illumination in a temperature-controlled growth chamber (Zubo *et*

al., 2017). BA was applied to ten-day-old plants by immersing seedlings in MS solution

containing 5 µM BA dissolved using dimethyl sulfoxide (DMSO), or DMSO without BA as a

control, and gently shaken for two hours. Plants were harvested, and shoots and roots were

collected separately. Libraries were sequenced on an Illumina HiSeq 2000 instrument for 100x2

cycles, yielding 100 base, paired end reads, to a depth of around 10 million sequenced fragments

per library. Sequence data are available from the Sequence Read Archive under SRP049054 (rice

data) and SRP059384 (*Arabidopsis* data). Aligned, processed data are available from the

TAIR10 (June 2009) *Arabidopsis* genome assembly IgbQuickload directories at http://lorainelab-

quickload.scidas.org/rnaseq/.

Data processing

Rice sequences were aligned onto the *O. sativa japonica* genome assembly Os-Nipponbare-

Reference-IRGSP-1.0 (Kawahara *et al.*, 2013) and *Arabidopsis* sequences were aligned onto the

TAIR10 June 2009 release of the *Arabidopsis* genome (Lamesch *et al.*, 2012) using TopHat

(Kim *et al.*, 2013) and BowTie2 (Langmead and Salzberg, 2012) with maximum intron size set

to 5,000 bases. A command-line, Java program called "FindJunctions" was used to identify

exon-exon junctions from gapped read alignments in the RNA-Seq data. FindJunctions produces

BED format files containing junction features, and the score field of the BED file lists the

number of read alignments that supported the junction. Only reads that aligned to a unique

location in the genome were considered. Source code and compiled versions of FindJunctions are

available from https://bitbucket.org/lorainelab/findjunctions.

Identification of alternative splicing events and differential splicing

To date, there have been two major releases of *O. sativa japonica* gene models: the MSU7

gene set (Kawahara *et al.*, 2013) and the RAP-Db gene set (Sakai *et al.*, 2013). The two gene

model sets contain mostly the same data, but the MSU7 gene models appear to be the most

heavily used and annotated with Gene Ontology terms. For simplicity, and to take advantage of

available functional annotations, we used the MSU7 annotations here. For analysis of

Arabidopsis data, we used TAIR10 (Lamesch *et al.*, 2012) and Araport11 (Cheng *et al.*, 2017)

gene models.

Annotated alternative splicing events and the number of reads supporting each alternative

were identified using the exon-intron overlap method introduced in (English *et al.*, 2010) and

further developed here for use with RNA-Seq data. Exons and introns from pairs of gene models

from the same locus were compared to identify alternatively spliced regions. Regions where an

intron in one model overlapped an exon in another model on the same strand were identified and

used to define mutually exclusive splicing choices (Fig. 1). Gene models that included an

alternatively spliced region were designated the "L" form (for "Long") with respect to the

splicing choice. Likewise, models that lacked an alternatively spliced region were designated "S"

(for "Short"). Alternatively spliced regions were labeled according to the type of alternative

splicing, as follows. Regions flanked by alternative donor sites were designated "DS" for

alternative donor site. Regions flanked by alternative acceptor sites were labeled "AS" for

alternative acceptor site. AS and DS events that coincided with exon skipping were labeled

"AS/ES" and "DS/ES". Alternatively spliced regions arising from introns that the spliceosome

sometimes failed to excise were designated "RI" for retained intron.

Fig. 1. Alternative splicing annotation. The overlap between an intron in one gene model and an exon in another gene model defines an alternatively spliced region. Arrows indicate splice sites, named AS for acceptor site and DS for donor site. Use of sites named AS-L or DS-L causes inclusion of the differentially spliced region, generating the longer (L) isoform. Similarly, DS-S and AS-S refer to sites that exclude the differentially spliced region and generate the shorter (S) isoform. (**A**) Alternative donor sites, in which the U2 snRNP complex forms at alternative locations on the 5' end of introns. (**B**) Alternative acceptor sites, in which the U1 snRNP complex forms at alternative sites near the 3' end of alternatively spliced introns. (**C**) Alternatively spliced intron, in which a donor/acceptor site pairing can either be used or not. used, forming a retained intron (RI). (**D**) Alternatively spliced, skipped exon. In exon skipping, alternative splicing involves four sites, indicated by DS-S/L, AS-L, DS-L, and SD-S/L. Exon inclusion requires assembly of two spliceosome complexes linking DS-S/L with AS-L and DS-L with AS-S/L, while exon skipping requires linking DS-S/L and AS-S/L only.

180
181

For each alternatively spliced region representing two mutually exclusive splicing choices,

RNA-Seq read alignments that unambiguously supported one or the other splicing choice were

counted. For AS and DS events, only gapped reads that aligned across intron junctions were

counted. For RI events, gapped reads that aligned across the retained intron were counted as

support for the intron-removed (S) form, and un-gapped reads that overlapped at least 20 bases

within the intron were counted as support for the intron-retained (L) form.

For each alternatively spliced region in each biological replicate, the number of reads

supporting L or S, but not both, were used to calculate percent-spliced-in (PSI) as N/M*100,

where N was the number of reads supporting the L form and M was the number of reads that

supported S or L but not both. This is the same as the splicing index described in (Katz *et al.*,

2010). To identify differentially spliced regions, a two-sided t-test was used to compare PSI

between sample types. Because PSI variance was large for events with small M (very few

informative reads), only alternatively spliced regions where M was 10 or more in at least three

replicate libraries were tested. A false discovery rate (FDR) was calculated for each test using the

method of Benjamini and Hochberg (Benjamini and Hochberg, 1995), as implemented in the R

programming language "p.adjust" method. Alternative splicing events with FDR less than or

equal to 0.1 were considered differentially alternatively spliced.

Software used to identify and quantify alternative events is available from

https://bitbucket.org/lorainelab/altspliceanalysis. Data analysis code used to analyze RNA-Seq

data is available from https://bitbucket.org/lorainelab/ricealtsplice. Data analysis code is

implemented as R Markdown files designed to be run in the RStudio development environment.

Readers interested in experimenting with different analysis parameters can clone the repository,

modify the code, and re-run analyses as desired. RNA-Seq alignments, coverage graphs, and

junctions data are available for visualization in Integrated Genome Browser (Freese *et al.*, 2016).

RT-PCR and capillary gel electrophoresis analysis of alternative splicing

Differential alternative splicing detected by analysis of RNA-Seq was re-tested using the

reverse transcriptase, PCR-based fragment analysis method described in (Stamm *et al.*, 2012).

Differentially spliced regions identified computationally were PCR-amplified using fluorescently

labeled primers and quantified using capillary gel electrophoresis. One benefit of the method is

that the results are expressed as relative abundances of splice variants within a sample, thus

eliminating the need to normalize using reference genes as in traditional qRT-PCR experiments

aimed at measuring overall gene expression.

For splicing validation, new rice seedlings equivalent to the mock-treated (control) samples

from the RNA-Seq experiment were grown and harvested. Seedlings were grown hydroponically

in pots containing either liquid media only or calcined clay granules watered with liquid media

as recommended in (Eddy *et al.*, 2012). After twelve days, plants were removed from the pots

and roots and shoots were collected separately. Roots and shoots from the same pot were

combined to form paired biological replicates. Samples were flash frozen in nitrogen and stored

at -80°C prior to RNA extraction.

RNA was extracted using the RNeasy Plant Mini Kit from Qiagen following the

manufacturer's instructions. First strand cDNA was synthesized using oligo dT primers and 1 μg

of total RNA per 20 μL reaction. PCR amplification of cDNA was performed using primers

flanking differentially spliced regions, including one primer labeled with 6-carboxyfluorescein

(6-FAM) for amplicon detection during fragment analysis. Cycle parameters included

denaturation at 94°C for 2 minutes, followed by 24 cycles of 94°C for 15 sec, 50°C for 30 sec

and 70°C for 1 min, with a final elongation step of 72°C for 10 minutes. This was essentially the

same regime described in (Stamm *et al.*, 2012) but with fewer cycles to ensure reactions were

stopped before exiting the logarithmic phase. PCR products were combined with size standards

and separated on a 3730 Genetic Analyzer (Life Technologies). Amplicons were quantified using

manufacturer-provided software by calculating the area under each amplicon peak. The

percentage of the variant containing the alternatively spliced region (%L, see above) was

calculated by dividing the long form area by the total area for both long and short forms.

Spreadsheets with data exported from the instrument, along with PSI calculations, are available

in the project git repository (https://bitbucket.org/lorainelab/ricealtsplice) in a subfolder named

"Experimental Testing."

RESULTS

Most genes annotated as alternatively spliced favored one dominant isoform

Using the exon-intron overlap method described previously (English *et al.*, 2010) and Fig. 1,

alternative splicing events within each gene were identified and annotated. Following annotation

of alternative splicing events, RNA-Seq read alignments from the rice and *Arabidopsis* libraries

described in (Raines *et al.*, 2016; Zubo *et al.*, 2017) were used to assess alternative splicing in

four sample types: roots and shoots from seedlings treated with the cytokinin compound

benzyladenine (BA) or with a mock, control treatment. For each alternative splicing event, the

number of sequence alignments unambiguously supporting each alternative was counted. These

counts were used to calculate percent-spliced-in (PSI), the percentage of read alignments

supporting the longer (L) isoform.

In the combined data from all libraries from the rice data set, 77% of AS events had at least

one read supporting each of the two splicing choices, and 19.8% had support for just one splicing

choice. Only 2.8% of AS events has no reads supporting either form; these corresponded to

genes with low or no expression in any of the sample types tested. Most genes annotated as

alternatively spliced had small minor variant frequencies, i.e., the less frequently observed forms

were supported by fewer than 20% of informative RNA-Seq sequences (Fig. 2). Nevertheless,

there was a large minority of alternative splicing events (around one third) where the minor, less-

frequently observed form was more abundant and was supported by at least two out of ten

informative alignments. These alternatively spliced regions correspond to the middle, trough-like

region of Fig. 2.

Fig. 2. Distribution of percent-spliced-in (PSI) for annotated splicing events in rice where each choice was supported by at least one RNA-Seq alignment. PSI was calculated as 100*L/(S+L), where L and S were the number of reads that supported the splicing choice that included (L) or excluded (S) the differentially spliced region. Read alignment counts from all twelve libraries were combined to obtain a global view of alternative splicing occurrence in rice seedlings. The U-shaped character of the distribution persisted whether lower or higher thresholds of informative reads were used.

Rice genes with abundant support for both alternative splicing choices perform many diverse functions

We used Gene Ontology term enrichment to determine if the subset of genes in rice for which alternatively spliced forms were unusually abundant exhibited enrichment with specific functions or processes, e.g., circadian cycling, in which alternative splicing might play a prominent regulatory role. We asked if some Gene Ontology terms were significantly enriched with genes containing alternative splicing events in which the minor form frequency was between 20 and 50%, corresponding to the central trough region of Fig. 2. Interestingly, we found that these genes exhibited a diversity of gene functions, with no significant enrichment of functional categories. Thus, alternative splicing in which minor forms are highly prevalent appears to affect genes with many functions in rice.

Many rice genes are differentially spliced between roots and shoots but cytokinin hormone application has minimal effect on splicing

In animals, differential splicing between cell or tissue types contributes to cellular differentiation, especially in the nervous system (Naftelberg *et al.*, 2015). Less is known about the role of alternative splicing in regulating cellular differentiation and other processes in plants. Rice shoots and roots are profoundly different tissues, but our previous analysis of this same data set found that many of the same genes were expressed in both (Raines *et al.*, 2016). This raises the question of how these two different tissues are able to carry out their specialized roles, and suggest the hypothesis that differential splicing could enable differential functions in genes expressed in both tissues, as proposed in (Reddy *et al.*, 2013).

Analysis of the effects of cytokinin treatment on this same data set from rice identified many thousands of genes that were differentially expressed in response to cytokinin (Raines *et al.*, 2016). However, little is known about the role of alternative splicing during cytokinin response, except for one study in *Arabidopsis* that reported a shift in splicing of SR protein genes following cytokinin hormone treatment (Palusa *et al.*, 2007). Therefore, we examined differential splicing in the rice RNA-Seq data set comparing root and shoot tissue with or without cytokinin.

First, we asked: When an alternatively spliced gene was expressed in two different sample types, was the relative abundance of splice variants the same or different? To address this, we examined correlation of PSI between roots and shoots or between BA-treated versus mock-

treated samples (Fig. 3). We found that PSI was similar between treated and untreated samples,

as revealed by the tighter clustering of scatter plot points (Fig. 3A-B). This indicated that genes

that were alternatively spliced in BA-treated samples were also alternatively spliced in the

controls, and that the relative abundance of splice variants was similar. Thus, the cytokinin

hormone treatment had minimal effect on splicing. By contrast, there were many genes where the

relative abundance of splice variants was different between roots and shoots (Fig. 3C).

Consistent with Fig. 3, statistical testing of PSI differences between sample types identified 90

genes where PSI was significantly different between roots and shoots (FDR ≤ 0.1) but only four

and two genes where PSI was different between cytokinin-treated samples and controls in roots

and shoots, respectively (Supplementary Table S1). Thus, we observed limited but non-trivial

levels of differential alternative splicing between roots and shoots but minimal differential

alternative splicing between control and BA-treated samples.

Fig. 3. Scatter plots comparing percent- spliced-in (PSI) between sample types in rice for annotated splicing events.
PSI was calculated from RNA-Seq reads obtained from sequencing rice seedling shoots and roots grown
hydroponically and subjected to a two-hour treatment with BA, a cytokinin analog, or a mock-treatment (control).
PSI is the average of three biological replicates. Only events with at least 15 informative read alignments in all six
samples being compared were included. (A) BA-treated rice roots (y axis) compared to mock roots (x axis). (B) BA-
treated rice shoots (y axis) compared to mock shoots (x axis). (C) Mock shoots (y axis) compared to mock roots (x
axis).

Comparison of *Arabidopsis* differential splicing shows similar patterns to rice

To determine whether the observed patterns of differential splicing are similar in other plants,

we analyzed splicing in *Arabidopsis* roots and shoots that had also been treated with cytokinin

(Zubo *et al.*, 2017). Due to the *Arabidopsis* libraries not being sequenced to the same depth as

the rice libraries, many more splicing events had little or no support. Using the same FDR

threshold as with the rice data set (FDR ≤ 0.1), we identified few differentially spliced regions

between shoots and roots (3) and none in the control to treatment comparisons (Supplementary

Table S2). However, PSI was distributed similarly to rice in that most alternatively spliced genes

expressed one major isoform (Supplementary Fig. S1A). In addition, scatter plots showing

average PSI in treated versus untreated samples showed a much tighter clustering of points as

compared to scatter plots comparing roots and shoots (Supplementary Fig. S1B-D). Statistical

testing of PSI differences confirmed the cytokinin hormone treatment had minimal effect on

splicing in *Arabidopsis*. Thus, the general pattern of more differential splicing between tissue

types as compared to treatment with exogenous cytokinin appears conserved between rice and

Arabidopsis.

Alternative splicing remodeled protein-coding sequence more often than disrupting it in

rice

Alternative splicing can occur anywhere in a gene, including UTR and protein-coding

regions. Most differential splicing between roots and shoots (67%) occurred within protein-

coding regions (Table 1 and Supplementary Table S1), suggesting that differential splicing is

likely to affect gene function at the level of the protein product. In every instance of differential

alternative splicing, major and minor isoforms were both detected, with differential splicing

observed as a change in the relative abundance of the two forms.

Because three bases encode one amino acid, the lengths of spliced coding regions in a

transcript are multiples of three. Thus, when alternatively spliced regions occur in coding regions

and are not multiples of three, then inclusion of these regions in transcripts is likely to introduce

a frame shift, resulting in a premature stop codon and a truncated protein product. As shown in

Table 1, there was an enrichment of alternatively spliced regions in rice that were evenly

divisible by three in coding regions versus non-coding in all subsets of the data. These subsets

included all annotated alternatively spliced regions, regions where the minor form was unusually

prevalent (the trough region of Fig. 2), and differentially spliced regions. Thus, alternative

splicing within the coding regions of genes was biased against introducing frame shifts and

promoted protein remodeling rather than truncation.

Table 1. *Counts of alternative splicing choices in rice that produce difference regions evenly divisible by three or with remainder of 1 or 2.*

Alternative Splicing		Divisible	Remainder	Remainder	*P-value
Location	Event	by 3	of 1	of 2	
Coding region	Annotated as alternatively spliced	3,248	2,466	2,411	3e-36
UTR	Annotated as alternatively spliced	1,152	1,127	1,113	1
Coding region	Minor form is expressed	173	149	153	8e-6
UTR	Minor form is expressed	34	20	13	0.03
Coding region	Differentially spliced	33	24	18	0.03
UTR	Differentially spliced	6	5	11	0.79

*P-value obtained from binomial test of the null hypothesis that the true probability of a differentially spliced region having a length divisible by three is 1 in 3 and an alternative hypothesis that the probability is greater than 1 in 3.

To further understand the effects of splicing on protein-coding sequences, we visualized
differentially spliced regions together with RNA-Seq alignments, coverage graphs, and inferred
junctions using genome browsers. Two genome browsers were used to visualize the data -
Integrated Genome Browser (Freese *et al.*, 2016) and ProtAnnot (Mall *et al.*, 2016). Integrated
Genome Browser (IGB) was used to examine RNA-Seq read alignments and compare
alignments to the annotated gene structures. ProtAnnot, an IGB App, was used to search the
InterPro database of conserved protein motifs to find out how (or if) splicing inferred from RNA-
Seq data was likely to affect gene function through remodeling of protein motifs as detected by
the InterProScan Web service (Finn *et al.*, 2017).
Of the 105 differentially spliced regions, 71 overlapped protein-coding sequence regions,
suggesting that in these cases, alternative splicing affected protein function. All but one (70/71)
of the differentially spliced regions embedded in coding regions overlapped a predicted
functional motif (e.g., a predicted transmembrane helix) or a region found by protein
classification systems (e.g., Pfam (Finn *et al.*, 2016) or PANTHER (Thomas *et al.*, 2003)) to be
conserved among members of the same protein family (Supplementary Table S1 and Fig. 4).

Fig. 4. ProtAnnot and IGB images showing difference in splicing between rice shoot and root. ProtAnnot (upper panel) shows coding region exons color-coded by frame, with regions matching InterPro profiles indicated by green, linked rectangles. Asterisk highlights difference in the PANTHER InterPro profile PTHR11516 between isoforms 1 and 5 of the LOC_Os12g08260 gene. Integrated Genome Browser (lower panels) shows a zoomed-in view of RNA-Seq coverage graphs from rice root (blue) and shoot (green). Y-axis is the number of RNA-Seq aligned sequences with MSU7 gene models in black below.

RT-PCR with capillary gel electrophoresis confirmed differential splicing between rice roots and shoots for nine of ten genes tested

We used a method based on capillary gel electrophoresis of fluorescently tagged PCR products to assay alternative splicing of ten genes detected as differentially spliced between rice roots and shoots (Stamm *et al.*, 2012). New rice seedlings were grown under a close-to-identical replication of the RNA-Seq experiment. Primers were designed to amplify differentially spliced regions, including one primer that was conjugated to a fluorescent tag. Following PCR amplification of cDNA prepared from the new rice samples, products were resolved on a capillary-based sequencer and PSI calculated (Table 2). In nine out of ten genes, differential alternative splicing was confirmed. In the one case where differential alternative splicing was not confirmed, there were very few RNA-Seq read alignments covering the differentially spliced region, suggesting this was likely a false positive result. The FDR cutoff used to detect differential splicing in the RNA-Seq data was 0.1, corresponding to one in ten false discoveries, in line with results from the microcapillary-based analysis.

Table 2. *Differential splicing detected using RNA-Seq and re-tested using capillary gel electrophoresis (CGE) in rice.*

Gene	AS type	Avg. RPKM Expression		RNA-Seq PSI (%L)		CGE PSI (%L)		*P-value
		Root	Shoot	Root	Shoot	Root	Shoot	
LOC_Os01g25484, ferrodoxin nitrite reductase	RI	300	131	74.2	31.7	29.9	13	5e-05
LOC_Os01g35580, unknown	AS	55.7	34.9	44.0	66.3	49.9	66.9	2.47e-04
LOC_Os01g45274, carbonic anhydrase	ES	171	1,380	96.8	24.3	97.4	15.4	3e-09
LOC_Os01g51290, protein kinase	RI	49.6	49.9	88.4	95.1	13.3	17.2	0.03466
LOC_Os03g05390, citrate transporter	RI	219	174	86.0	96.3	86	95.5	1.2e-04
LOC_Os12g08260, dehydrogenase E1	RI	8.03	30.3	55.7	2.9	4.1	0.87	3e-05
LOC_Os01g61670, ureidoglycolate hydrolase	DS	59	37.8	78.0	31.0	59	37.8	1.6e-09
LOC_Os05g48040, MATE efflux family protein	DS	7.05	6.72	88.1	100.0	89.4	88.27	0.632
LOC_Os02g05830, ribulose bisphosphate carboxylase	RI	4.79	12.3	88.2	10.1	23.9	3.06	8e-04
LOC_Os06g05110, superoxide dismutase	RI	12.5	39.7	38.5	13.6	22.7	6.5	1.3e-07

*P-value obtained from comparing roots and shoots PSI from CGE.

DISCUSSION

This study profiled alternative splicing using a high coverage RNA-Seq data from 10-day

old, hydroponically-grown rice seedlings treated with a cytokinin hormone. A less-deeply

sequenced data set from similarly treated *Arabidopsis* seedlings provided comparison with

another plant species. We found that cytokinin treatment induced very few splicing changes

between treated and untreated controls. However, there were many differences in splicing

between untreated roots and shoots, and most of these changed the protein coding region of

genes.

Palusa et al. found that BA-treatment of *Arabidopsis* seedlings triggered splicing changes in

multiple SR genes (Palusa *et al.*, 2007), encoding RNA-binding proteins whose counterparts in

metazoans regulated alternative splice site choice. Their study used PCR amplification of cDNA

followed by agarose gel electrophoresis to detect changes in splicing and focused on SR protein

genes only. Thus, we expected to observe changes in SR gene splicing due to the cytokinin

treatment, leading to changes in splicing for many downstream genes. However, no such

differential splicing was apparent in either RNA-Seq data set tested. It is possible that the

differences in methodology used to measure splicing changes between the two studies (RNA-Seq

versus visualization of PCR amplification of cDNA) could account for the differences in

observations. However, close examination of SR splicing genes in our dataset revealed no

significant differences.

One possible explanation for why the cytokinin treatment had minimal effect on splicing was

that the treatment itself was ineffective. However, differential expression analysis showed that

many genes were up- or down- regulated by the treatment in the two data sets tested – rice and

Arabidopsis (Raines *et al.*, 2016). Fewer genes were detected as differentially expressed in the

Arabidopsis data set, most likely reflecting higher variability between biological replicates

combined with lower sequencing depth as compared to the rice data set. Nevertheless, known

cytokinin response genes were differentially regulated in both data sets, showing the cytokinin

treatment penetrated plant cells and induced stereotypical cytokinin signaling without also

triggering changes in splicing.

The relative lack of differential splicing between cytokinin-treated and mock-treated samples

suggests that cytokinin signaling does not employ alternative splicing as a regulatory mechanism

to the same degree as with other plant hormones, notably abscisic acid (ABA). ABA plays a role

420 in perception and response to stresses, especially desiccation stress (Maia *et al.*, 2014). ABA also

421 plays a role in regulating splicing of SR45, an SR-like protein, and SR45 plays a role in

422 regulating downstream splicing of multiple genes (Cruz *et al.*, 2014). Thus, there is a clear

423 connection between stress signaling and the plant stress hormone ABA.

424  By contrast, cytokinin signaling involves transfer of phosphate groups between successive

425 elements of a phosphorelay signaling pathway culminating in phosophorylation-dependent

426 activation of Myb-type transcription factor proteins called type B ARRs. Cytokinin treatment has

427 no or little effect on transcription of type B ARRs, the key regulators of cytokinin signaling

428 (Argyros *et al.*, 2008; Kieber and Schaller, 2018). In addition, type B transcriptional regulators

429 are not highly alternatively spliced. By contrast, a closely related family of similar genes

430 encoding so-called "pseudo-response regulators" have similar sequence to type B ARRs and are

431 highly alternatively spliced (James *et al.*, 2012). These genes are involved in regulating the

432 circadian clock and have nothing to do with cytokinin signaling.

433  Using the same methods and data set, we identified a relatively large number of genes in rice

434 (90) that were differentially spliced genes between shoots and roots, and we validated nine of ten

435 using fragment analysis of independently produced rice samples. This observation of differential

436 splicing between roots and shoots is important for two key reasons. First, it shows that our data

437 analysis methods can identify differential alternative splicing in a data set. In other words, the

438 roots versus shoots comparison provided a positive control for differential splicing detection.

439 Second, the detection of differential splicing between roots and shoots illuminates the function of

440 alternative splicing in plant cells. For most of the differentially spliced regions, both forms were

441 present, and the difference in relative abundance between forms was often slight, rarely more

442 than five or ten percent (Supplementary Table S1). Our data supports the growing body of

443 evidence that alternative splicing is cell, tissue, and stage specific in plants (Vitulo *et al.*, 2014;

444 Gupta *et al.*, 2015; Li *et al.*, 2016a; Sun *et al.*, 2018), including in roots (Li *et al.*, 2016b). It is

445 likely that alternative splicing plays a role in fine-tuning gene function to meet the needs of

446 different plant tissues or cell types where a gene is expressed.

447  We also examined the prevalence of alternative splicing, independent from differential

448 splicing. That is, we used RNA-Seq read alignments to assess how often annotated alternative

449 splice sites were used in our RNA-Set data sets. For most genes annotated as alternatively

450 spliced, the minor form frequency was typically low, accounting for less than 20% of sequence

read alignments across the differentially spliced region. Genes where minor form frequency

exceeded 20% exhibited a diversity of functions. Thus, many diverse processes in rice involved

alternatively spliced genes in which splice variants were expressed at levels likely to affect gene

function in different ways.

A major limitation of this study was that we limited our analysis to annotated splice forms

and did not attempt to form new transcript models based on the RNA-Seq data. This was done

mainly because the libraries used were not strand-specific and attempts to assemble transcripts

using transcript assembly tools led to incorrect fusion of neighboring genes and other artifacts

(not shown). Future studies will therefore benefit greatly from using better library preparation

protocols to simplify and streamline data analysis. Nonetheless, this analysis provides new

insight into the role of alternative splicing in plant tissues and hormone response.

In conclusion, by analyzing the number of reads that supported different splice variants, we

identified examples of differential splicing with confirmation by RT-PCR with capillary gel

electrophoresis. There were 90 genes differentially spliced between rice root and shoot tissues,

but only four between cytokinin-treated and non-treated samples. For most differential splicing

events, the protein-coding regions were affected, strongly suggesting that differential splicing is

playing a role in modulating gene function between roots and shoots.

Supplementary Table S1. Spreadsheet containing the rice mock root vs mock shoot, mock root

vs treated root, and mock shoot vs treated shoot splicing data.

Supplementary Table S2. Spreadsheet containing the *Arabidopsis* mock root vs mock shoot

splicing data.

ACKNOWLEDGEMENTS

This material is based upon work supported by the National Science Foundation through the

Postdoctoral Research Fellowship in Biology under Grant No.1523814 awarded to NHF and the

Plant Genome Research Program award number IOS-1238051. NIH award 2R01GM103463-

06A1 to AEL supports development of the IGB software.

Fig. S1. Percent-spliced-in (PSI) in *Arabidopsis* **RNA-Seq data set between sample types for**
annotated splicing events. (A) Counts of PSI events in all samples combined. PSI was calculated from
RNA-Seq reads obtained from sequencing *Arabidopsis* seedling shoots and roots grown with or without
BA, a cytokinin analog. PSI is the average of three biological replicates. Only events with at least 10
informative read alignments in all six samples being compared were included. (B) BA-treated
Arabidopsis roots (y axis) compared to mock roots (x axis). (C) BA-treated *Arabidopsis* shoots (y axis)
compared to mock shoots (x axis). (D) Mock shoots (y axis) compared to mock roots (x axis).
488

REFERENCES CITED

Argyros RD, Mathews DE, Chiang Y-H, Palmer CM, Thibault DM, Etheridge N, Argyros DA, Mason MG, Kieber JJ, Schaller GE. 2008. Type B Response Regulators of *Arabidopsis* Play Key Roles in Cytokinin Signaling and Plant Development. The Plant Cell **20**, 2102-2116.

Barbosa-Morais NL, Carmo-Fonseca M, Aparício S. 2006. Systematic genome-wide annotation of spliceosomal proteins reveals differential gene family expansion. Genome Research **16**, 66-77.

Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, Slobodeniuc V, Kutter C, Watt S, Colak R, Kim T, Misquitta-Ali CM, Wilson MD, Kim PM, Odom DT, Frey BJ, Blencowe BJ. 2012. The evolutionary landscape of alternative splicing in vertebrate species. Science **338**, 1587-1593.

Benjamini Y, Hochberg Y. 1995. *Controlling The False Discovery Rate - A Practical And Powerful Approach To Multiple Testing.*

Cheng CY, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD. 2017. Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. Plant J **89**, 789-804.

Cruz TM, Carvalho RF, Richardson DN, Duque P. 2014. Abscisic acid (ABA) regulation of *Arabidopsis* SR protein gene expression. Int J Mol Sci **15**, 17541-17564.

Eddy R, Hahn DT, Moulton T. 2012. Optimizing Greenhouse Rice Production: What Is the Best Root Medium? Purdue Methods for Rice Growth.

English AC, Patel KS, Loraine AE. 2010. Prevalence of alternative splicing choices in *Arabidopsis thaliana*. BMC Plant Biol **10**, 102.

Filichkin S, Priest HD, Megraw M, Mockler TC. 2015a. Alternative splicing in plants: directing traffic at the crossroads of adaptation and environmental stress. Curr Opin Plant Biol **24**, 125-135.

Filichkin Sergei A, Cumbie Jason S, Dharmawardhana P, Jaiswal P, Chang Jeff H, Palusa Saiprasad G, Reddy ASN, Megraw M, Mockler Todd C. 2015b. Environmental Stresses Modulate Abundance and Timing of Alternatively Spliced Circadian Transcripts in *Arabidopsis*. Molecular Plant **8**, 207-227.

Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, Chang H-Y, Dosztányi Z, El-Gebali S, Fraser M, Gough J, Haft D, Holliday GL, Huang H, Huang X, Letunic I, Lopez R, Lu S, Marchler-Bauer A, Mi H, Mistry J, Natale DA, Necci M, Nuka G, Orengo CA, Park Y, Pesseat S, Piovesan D, Potter SC, Rawlings ND, Redaschi N, Richardson L, Rivoire C, Sangrador-Vegas A, Sigrist C, Sillitoe I, Smithers B, Squizzato S, Sutton G, Thanki N, Thomas PD, Tosatto Silvio C E, Wu CH, Xenarios I, Yeh L-S, Young S-Y, Mitchell AL. 2017. InterPro in 2017—beyond protein family and domain annotations. Nucleic Acids Research **45**, D190-D199.

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families database: towards a more sustainable future. Nucleic Acids Research **44**, D279-D285.

Freese NH, Norris DC, Loraine AE. 2016. Integrated genome browser: visual analytics platform for genomics. Bioinformatics **32**, 2089-2095.

Gulledge AA, Roberts AD, Vora H, Patel K, Loraine AE. 2012. Mining *Arabidopsis thaliana* RNA-seq data with Integrated Genome Browser reveals stress-induced alternative splicing of the putative splicing regulator SR45a. American journal of botany **99**, 219-231.

Gupta V, Estrada AD, Blakley I, Reid R, Patel K, Meyer MD, Andersen SU, Brown AF, Lila MA, Loraine AE. 2015. RNA-Seq analysis and annotation of a draft blueberry genome assembly identifies candidate genes involved in fruit ripening, biosynthesis of bioactive compounds, and stage-specific alternative splicing. Gigascience **4**, 5.

James AB, Syed NH, Bordage S, Marshall J, Nimmo GA, Jenkins GI, Herzyk P, Brown JW, Nimmo HG. 2012. Alternative splicing mediates responses of the *Arabidopsis* circadian clock to temperature changes. Plant Cell **24**, 961-981.

Kalsotra A, Cooper TA. 2011. Functional consequences of developmentally regulated alternative splicing. Nature Reviews. Genetics **12**, 715-729.

Kalyna M, Barta A. 2004. A plethora of plant serine/arginine-rich proteins: redundancy or evolution of novel gene functions? Biochemical Society transactions **32**, 561-564.

Katz Y, Wang ET, Airoldi EM, Burge CB. 2010. Analysis and design of RNA sequencing experiments for identifying isoform regulation. Nat Methods **7**, 1009-1015.

Kawahara Y, de la Bastide M, Hamilton JP, Kanamori H, McCombie WR, Ouyang S, Schwartz DC, Tanaka T, Wu J, Zhou S, Childs KL, Davidson RM, Lin H, Quesada-Ocampo L, Vaillancourt B, Sakai H, Lee SS, Kim J, Numa H, Itoh T, Buell CR, Matsumoto T. 2013. Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data. Rice **6**, 1-10.

Keller M, Hu Y, Mesihovic A, Fragkostefanakis S, Schleiff E, Simm S. 2016. Alternative splicing in tomato pollen in response to heat stress. DNA Research, dsw051.

Kieber JJ, Schaller GE. 2018. Cytokinin signaling in plant development. Development **145**.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol **14**, R36.

Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, Muller R, Dreher K, Alexander DL, Garcia-Hernandez M, Karthikeyan AS, Lee CH, Nelson WD, Ploetz L, Singh S, Wensel A, Huala E. 2012. The Arabidopsis Information Resource (TAIR): improved gene annotation and new tools. Nucleic Acids Research **40**, D1202-D1210.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods **9**, 357.

Li S, Yamada M, Han X, Ohler U, Benfey PN. 2016a. High resolution expression map of the *Arabidopsis* root reveals alternative splicing and lincRNA regulation. Developmental cell **39**, 508-522.

Li S, Yamada M, Han X, Ohler U, Benfey PN. 2016b. High-Resolution Expression Map of the Arabidopsis Root Reveals Alternative Splicing and lincRNA Regulation. Dev Cell **39**, 508-522.

Loraine AE, McCormick S, Estrada A, Patel K, Qin P. 2013. RNA-seq of *Arabidopsis* pollen uncovers novel transcription and alternative splicing. Plant Physiol **162**, 1092-1109.

Maia J, Dekkers BJ, Dolle MJ, Ligterink W, Hilhorst HW. 2014. Abscisic acid (ABA) sensitivity regulates desiccation tolerance in germinated *Arabidopsis* seeds. New Phytol **203**, 81-93.

Mall T, Eckstein J, Norris D, Vora H, Freese NH, Loraine AE. 2016. ProtAnnot: an App for Integrated Genome Browser to display how alternative splicing and transcription affect proteins. Bioinformatics **32**, 2499-2501.

Mei W, Boatwright L, Feng G, Schnable JC, Barbazuk WB. 2017. Evolutionarily Conserved Alternative Splicing Across Monocots. Genetics **207**, 465-480.

Naftelberg S, Schor IE, Ast G, Kornblihtt AR. 2015. Regulation of alternative splicing through coupling with transcription and chromatin structure. Annual review of biochemistry **84**, 165-198.

Palusa SG, Ali GS, Reddy ASN. 2007. Alternative splicing of pre-mRNAs of *Arabidopsis* serine/arginine-rich proteins: regulation by hormones and stresses. The Plant Journal **49**, 1091-1107.

Plass M, Agirre E, Reyes D, Camara F, Eyras E. 2008. Co-evolution of the branch site and SR proteins in eukaryotes. Trends in Genetics **24**, 590-594.

Raines T, Blakley IC, Tsai Y-C, Worthen JM, Franco-Zorrilla JM, Solano R, Schaller GE, Loraine AE, Kieber JJ. 2016. Characterization of the cytokinin-responsive transcriptome in rice. BMC Plant Biology **16**, 260.

Reddy ASN, Marquez Y, Kalyna M, Barta A. 2013. Complexity of the Alternative Splicing Landscape in Plants. The Plant Cell **25**, 3657-3683.

Sakai H, Lee SS, Tanaka T, Numa H, Kim J, Kawahara Y, Wakimoto H, Yang CC, Iwamoto M, Abe T, Yamada Y, Muto A, Inokuchi H, Ikemura T, Matsumoto T, Sasaki T, Itoh T. 2013. Rice Annotation Project Database (RAP-DB): an integrative and interactive database for rice genomics. Plant Cell Physiol **54**, e6.

Salz HK. 2011. Sex Determination in Insects: a binary decision based on alternative splicing. Current opinion in genetics & development **21**, 395-400.

Stamm S, Smith C, Lührmann R. 2012. *Alternative pre-mRNA Splicing: Theory and Protocols*: Wiley-Blackwell.

Sun Y, Hou H, Song H, Lin K, Zhang Z, Hu J, Pang E. 2018. The comparison of alternative splicing among the multiple tissues in cucumber. BMC Plant Biology **18**, 5.

Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A, Narechania A. 2003. PANTHER: A Library of Protein Families and Subfamilies Indexed by Function. Genome Research **13**, 2129-2141.

Vaneechoutte D, Estrada AR, Lin YC, Loraine AE, Vandepoele K. 2017. Genome-wide characterization of differential transcript usage in Arabidopsis thaliana. Plant J **92**, 1218-1231.

Vitulo N, Forcato C, Carpinelli EC, Telatin A, Campagna D, D'Angelo M, Zimbello R, Corso M, Vannozzi A, Bonghi C, Lucchin M, Valle G. 2014. A deep survey of alternative splicing in grape

reveals changes in the splicing machinery related to tissue, stress condition and genotype. BMC Plant Biology **14**, 99-99.

Zubo YO, Blakley IC, Yamburenko MV, Worthen JM, Street IH, Franco-Zorrilla JM, Zhang W, Hill K, Raines T, Solano R, Kieber JJ, Loraine AE, Schaller GE. 2017. Cytokinin induces genome-wide binding of the type-B response regulator ARR10 to regulate growth and development in *Arabidopsis*. Proceedings of the National Academy of Sciences of the United States of America **114**, E5995-E6004.