1 **Many rice genes are differentially spliced between roots and shoots but cytokinin hormone**
2 **application has minimal effect on splicing**
3

4 Nowlan Freese[1*], April R. Estrada[1*], Ivory C. Blakley[1], Jinjie Duan[2], Ann Loraine[1**]
5 [1]Department of Bioinformatics and Genomics, University of North Carolina at Charlotte, Charlotte, NC,
6 USA
7 [2]Bioinformatics Research Center, Aarhus University, C.F. Moellers Alle 8, Aarhus C, Denmark.
8
9 [*]Equal contribution.
10 [**]Corresponding author.
11

ABSTRACT

Alternatively spliced genes produce multiple spliced isoforms, called transcript variants. In differential
alternative splicing, the relative abundance of transcript variants differs across sample types. Differential
alternative splicing is common in animal systems and influences cellular development in many processes,
but its extent and significance is not as well known in plants. To investigate alternative splicing in plants,
we examined rice seedling RNA-Seq data that included approximately 40 million sequence alignments
per library, three libraries per sample type, and four sample types: roots and shoots, each treated with
exogenous cytokinin delivered hydroponically, plus mock-treated controls. Cytokinin treatment triggered
expression changes in thousands of genes, but had minimal effect on alternative splicing. However, there
were many splicing differences between roots and shoots. Analysis of a similar (but less deeply
sequenced) mRNA-Seq data set from Arabidopsis showed the same pattern. Quantitative fragment
analysis of reverse transcriptase-PCR products made from newly prepared rice samples confirmed nine of
ten differential splicing events between rice roots and shoots. In both the Arabidopsis and rice mRNA-
Seq data sets, most genes annotated as alternatively spliced had small minor variant frequencies, i.e., the
less frequently observed forms were supported by fewer than 20% of informative mRNA-Seq sequences.
Of splicing choices where there was more abundant support for minor forms, most alternative splicing
events were located within the protein-coding sequence and also maintained the annotated reading frame.
A tool for visualizing protein annotations in the context of genomic sequence (ProtAnnot) together with a
genome browser (Integrated Genome Browser) were used to visualize and assess effects of differential
splicing on gene function. In general, differentially spliced regions coincided with conserved regions in
the encoded proteins, indicating that differential alternative splicing is likely to affect protein function
between root and shoot tissue in rice.
34

INTRODUCTION
Differential splicing of pre-mRNA transcripts, called alternative splicing, enables one gene to
produce multiple transcript variants encoding different functions. Alternative splicing is an almost
universal phenomenon in higher eukaryotes, occurring to varying degrees in every animal and plant
genome examined to date [1,2]. In animals, differential expression of splice variants has been recruited as
a regulatory mechanism in multiple processes, such as sex determination in invertebrates and neuronal
differentiation in mammals [2-4].
In plants, less is known about the functional significance and patterns of alternative splicing.
However, several trends are apparent. Genes involved in circadian regulation are highly alternatively
spliced, often producing multiple splice variants that fluctuate in concert with day/night cycling along
with overall transcript abundance [5]. The SR family of RNA-binding, splicing regulatory proteins is
greatly expanded compared to mammals and includes many plant-specific forms [6-9]. SR transcripts
themselves are also highly alternatively spliced in plants, with the relative abundance of these alternative
transcripts varying according to environmental stresses and hormones [6,10-13]. Our work on cytokinin
hormone response in rice and *Arabidopsis* found significant differences in gene expression following
cytokinin exposure [14], and previous work in *Arabidopsis* by Palusa et al. found that some SR transcripts

exhibit differential splicing when exposed to cytokinin [11], suggesting that alternative splicing may play
a role in cytokinin response.
There is a growing body of evidence indicating cell and tissue specific regulation of alternative
splicing in plants [15-17]. We previously found through analysis of RNA-Seq data from *Arabidopsis*
pollen that the relative abundance of splice variants was similar between leaves and pollen, despite the
differences between the two tissues [18]. However, this latter analysis was limited by having just one
biological replicate for pollen and only two biological replicates for leaves. A more comprehensive
analysis of multiple *Arabidopsis* data sets found a high incidence of isoform switching, in which the
identity of the most prevalent variant differs between sample types [19]. However, this splicing diversity
may have arisen in part from the heterogeneity of the data sets used, which were produced using rapidly
changing (and improving) sequencing technologies at different times by different groups.
In this study, we used a well-replicated RNA-Seq data set from rice to re-examine prevalence of
alternative splicing between tissues and hormone (cytokinin) treatment. This data set was previously
generated to investigate cytokinin regulation of gene expression in roots and shoots from 10-day old rice
seedlings [14]. The same study also produced an analogous data set from Arabidopsis for comparison, but
was less deeply sequenced. Both the rice and Arabidopsis mRNA-Seq data sets included three biological
replicates per sample type and four sample types – roots and shoots treated with exogenous cytokinin or a
mock, vehicle-only treatment. In both data sets, the treatment triggered differential expression of
thousands of genes, with roots affected to a greater degree than shoots.
We found that the relative abundance of splice variants for most alternatively spliced genes was
stable, with very few differentially spliced genes between cytokinin treated and control samples. There
was nonetheless a small but significant number of genes that were differentially spliced between roots and
shoots, with the majority of the differential splicing occurring within the protein-coding sequence. These
results provide new evidence that differential alternative splicing likely contributes to gene function
diversification between roots and shoots while playing little role in cytokinin signaling.
76
77 **MATERIALS & METHODS**
78
RNA-Seq library preparation and sequencing
Rice and Arabidopsis samples were prepared and sequenced as described in [14]. Rice seedlings
(Nipponbare) were grown hydroponically for ten days in a growth chamber set to 14 hours light (28°C)
and 8 hours of dark (23°C) with light intensity 700 mmol m^{-2} s^{-1}. Around six to ten seedlings were grown
in the same pot, in four pots. On the tenth day, culture media was replaced with new media containing 5
mM of the cytokinin benzyladenine (BA) or 0.05 mN NaOH as a control. After 120 minutes, roots and
shoots were harvested separately. Roots and shoots from treatment or control pots were pooled to form
three replicates per treatment. RNA was extracted and used to synthesize twelve libraries from BA-treated
and mock-treated roots and shoots. Libraries were sequenced on an Illumina HiSeq instrument for 100
cycles, yielding 100 base, single end reads. Sequence data are available from the Sequence Read Archive
under accession SRP04905. Aligned, processed data are available from the Oct. 2011 rice genome
assembly IgbQuickload directories at http://www.igbquickload.org.
Arabidopsis plants were grown vertically on 1X MS agar with 1% sucrose for ten days under
continuous illumination in a temperature-controlled growth chamber (Raines et al., 2016). BA was
applied to ten-day-old plants by immersing seedlings in MS solution containing 5 mM BA or vehicle
control and gently shaken for two hours. Plants were then harvested and shoots and roots collected
separately. Libraries were sequenced on an Illumina HiSeq 2000 instrument for 100x2 cycles, yielding
100 base, paired end reads, to a depth of around 10 million sequenced fragments per library. Sequence
data are available from the Sequence Read Archive under accession SRP059384. Aligned, processed data
are available from the TAIR10 (June 2009) Arabidopsis genome assembly IgbQuickload directories at
http://lorainelab-quickload.scidas.org/rnaseq/.
100
Data processing

Rice sequences were aligned onto the *O. sativa japonica* genome assembly Os-Nipponbare-
Reference-IRGSP-1.0 [20] and Arabidopsis sequences were aligned onto the TAIR10 June 2009 release
of the Arabidopsis genome (the latest) using TopHat and BowTie2 with maximum intron size was set to
5,000 bases. A command-line, Java program called "FindJunctions" was used to identify exon-exon
junctions from gapped read alignments in the RNA-Seq data. FindJunctions produces BED format files
containing junction features, and the score field of the BED file lists the number of read alignments that
supported the junction. Only reads that aligned to a unique location in the genome were considered.
Source code and compiled versions of FindJunctions are available from
https://bitbucket.org/lorainelab/findjunctions.
111
Identification of alternative splicing events and differential splicing
To date, there have been two major releases of *O. sativa japonica* gene models: the MSU7 gene set
[20] and the RAP-Db gene set [21]. The two gene model sets contain mostly the same data, but the MSU7
gene models appear to be the most heavily used and annotated with Gene Ontology terms. For simplicity,
and to take advantage of available functional annotations, we used the MSU7 annotations here. For
analysis of Arabidopsis data, we used TAIR10 and Araport11 gene models.
Annotated alternative splicing events and the number of reads supporting each alternative were
identified using the exon-intron overlap method described in [22]. Exons and introns from pairs of gene
models from the same locus were compared to identify alternatively spliced regions. Regions where an
intron in one model overlapped an exon in another model on the same strand were identified and used to
define mutually exclusive splicing choices (Figure 1). Gene models that included an alternatively spliced
region were designated the "L" form (for "Long") with respect to the splicing choice. Likewise, models
that lacked an alternatively spliced region were designated "S" (for "Short"). Alternatively spliced regions
were labeled according to the type of alternative splicing, as follows. Regions flanked by alternative
donor sites were designated "DS" for alternative donor site. Regions flanked by alternative acceptor sites
were labeled "AS" for alternative acceptor site. AS and DS events that coincided with exon skipping were
labeled "AS/ES" and "DS/ES". Alternatively spliced regions arising from introns that the spliceosome
sometimes failed to excise were designated "RI" for retained intron.
For each alternatively spliced region representing two mutually exclusive splicing choices, RNA-Seq
read alignments that unambiguously supported one or the other splicing choice were counted. For AS and
DS events, only gapped reads that aligned across intron junctions were counted as support. For RI events,
gapped reads that aligned across the retained intron were counted as support for the intron-removed (S)
form, and un-gapped reads that overlapped at least 20 bases within the intron were counted as support for
the intron-retained (L) form.
For each alternatively spliced region in each biological replicate, the number of reads supporting L or
S, but not both, were used to calculate percent-spliced-in (PSI) as $N/M*100$, where N was the number of
reads supporting the L form and M was the number of reads that supported S or L but not both. This is the
same as the splicing index described in [23]. To identify differentially spliced regions, a two-sided t-test
was used to compare PSI between sample types. Because PSI variance was large for events with small M
(very few informative reads), only alternatively spliced regions where M was 10 or more in at least three
replicate libraries were tested. A false discovery rate (FDR) was calculated for each test using the method
of Benjamini and Hochberg, as implemented in the R programming language "p.adjust" method.
Alternative splicing events with FDR less than or equal to 0.1 were considered differentially alternatively
spliced.
Software used to identify and quantify alternative events is available from
https://bitbucket.org/lorainelab/altspliceanalysis. Data analysis code used to analyze RNA-Seq data is
available from https://bitbucket.org/lorainelab/ricealtsplice. Data analysis code is implemented as R
Markdown files designed to be run in the RStudio development environment. Readers interested in
experimenting with different analysis parameters can clone the repository, modify the code, and re-run
analyses as desired. RNA-Seq alignments, coverage graphs, and junctions data are available for
visualization in Integrated Genome Browser.

153
RT-PCR and capillary gel electrophoresis analysis of alternative splicing
Differential alternative splicing detected by analysis of RNA-Seq was re-tested using the reverse
transcriptase, PCR-based fragment analysis method described in [24]. Differentially spliced regions
identified computationally were PCR-amplified using fluorescently labeled primers and quantified using
capillary gel electrophoresis. One benefit of the method is that the results are expressed as relative
abundances of splice variants within a sample, thus eliminating the need to normalize using reference
genes as in traditional qRT-PCR experiments aimed at measuring overall gene expression.
For splicing validation, new rice seedlings equivalent to the mock-treated (control) samples from the
RNA-Seq experiment were grown and harvested. Seedlings were grown hydroponically in pots
containing either liquid media only or calcined clay granules watered with liquid media as recommended
in [25]. After twelve days, plants were removed from the pots and roots and shoots were collected
separately. Roots and shoots from the same pot were combined to form paired biological replicates.
Samples were frozen on liquid nitrogen and stored at -80 degrees C prior to RNA extraction.
RNA was extracted using the RNeasy Plant Mini Kit from Qiagen following the manufacturer's
instructions. First strand cDNA was synthesized using oligo dT primers and 1 μg of total RNA per 20 μL
reaction. PCR amplification of cDNA was performed using primers flanking differentially spliced
regions, including one primer labeled with 6-carboxyfluorescein (6-FAM) for amplicon detection during
fragment analysis. Cycle parameters included denaturation at 94°C for 2 minutes, followed by 24 cycles
of 94°C for 15 sec, 50°C for 30 sec and 70°C for 1 min, with a final elongation step of 72°C for 10
minutes. This was essentially the same regime described in [24] but with fewer cycles to ensure reactions
were stopped before exiting the logarithmic phase. PCR products were combined with size standards and
separated on a 3730 Genetic Analyzer (Life Technologies). Amplicons were quantified using
manufacturer-provided software by calculating the area under each amplicon peak. The percentage of the
variant containing the alternatively spliced region (%L, see above) was calculated by dividing the long
form area by the total area for both long and short forms. Spreadsheets with data exported from the
instrument, along with PSI calculations, are available in the project git repository in a subfolder named
"Experimental Testing."
181
182 **RESULTS**
183
Most genes annotated as alternatively spliced favored one dominant isoform
Using the exon-intron overlap method described previously [22], alternative splicing events within
each gene were identified and annotated as shown in Figure 1. Following annotation of alternative
splicing events, RNA-Seq read alignments from the rice and Arabidopsis libraries described in [14] were
used to assess alternative splicing in four sample types: roots and shoots from seedlings treated with the
cytokinin compound benzyladenine (BA) or with a mock, control treatment. For each alternative splicing
event, the number of sequence alignments unambiguously supporting each alternative was counted. These
counts were used to calculate percent-spliced-in (PSI), the percentage of read alignments supporting the
longer (L) isoform.
In the combined data from all libraries from the rice data set, 77% of AS events had at least one read
supporting each of the two splicing choices, and 19.8% had support for just one splicing choice. Only
2.8% of AS events has no reads supporting either form; these corresponded to genes with low or no
expression in any of the sample types tested.
In the rice mRNA-Seq data set, most genes annotated as alternatively spliced had small minor variant
frequencies, i.e., the less frequently observed forms were supported by fewer than 20% of informative
mRNA-Seq sequences (Figure 2). In 70% of annotated alternative splicing events, only 20% or fewer
alignments supported the lower frequency choice.
Among events with support for mutually exclusive splicing choices, the distribution of read support
skewed toward supporting a single choice (Figure 2). In 54% of annotated alternative splicing events,
only 10% or fewer of informative read alignments supported the minor choice. Conversely, 30% of

alternative splicing choices showed simultaneous expression of multiple choices in non-trivial amounts,
e.g., at least 2 in 10 alignments supported the minor form.
206
Rice genes with abundant support for both alternative splicing choices exhibit a variety of functions
We used Gene Ontology term enrichment to determine if the subset of genes in rice for which
alternatively spliced forms were unusually abundant (the central "trough" region of Figure 2) were
enriched with certain functions or processes, e.g., circadian cycling, in which alternative splicing might
play a prominent regulatory. Specifically, we asked if some Gene Ontology terms were significantly
enriched with genes containing alternative splicing events in which the minor form frequency was
between 20 and 50%. Interestingly, we found that these genes exhibited a diversity of gene functions,
with no significant enrichment of functional categories. Thus, alternative splicing in which minor forms
are highly prevalent appears to affect genes with many functions in rice.
216
Many rice genes are differentially spliced between roots and shoots but cytokinin hormone
application has minimal effect on splicing
In animals, differential splicing between cell or tissue types contributes to cellular differentiation,
especially in the nervous system [26]. Less is known about the role of alternative splicing in regulating
cellular differentiation and other processes in plants. Rice shoots and roots are profoundly different
tissues, but our previous analysis of this same data set found that many of the same genes were expressed
in both [14]. This raises the question of how these two different tissues are able to carry out their
specialized roles, and suggest the hypothesis that differential splicing could enable differential functions
in genes expressed in both tissues, as proposed in [1].
Analysis of the effects of cytokinin treatment on this same data set from rice identified many
thousands of genes of that were differentially expressed in response to cytokinin [14]. However, little is
known about the role alternative splicing during cytokinin response, except for one study in Arabidopsis
that reported a shift in splicing of SR protein genes following cytokinin hormone treatment [11].
Therefore, we examined differential splicing in the rice RNA-Seq data set comparing root and shoot
tissue with or without cytokinin.
First, we asked: When an alternatively spliced gene was expressed in two different sample types, was
the relative abundance of splice variants the same or different? To address this, we examined correlation
of PSI between roots and shoots or between BA-treated versus mock-treated samples (Figure 3). We
found that PSI was similar between treated and untreated samples, as revealed by the tighter clustering of
scatter plot points in Figures 3A and 3B. This indicated that genes that were alternatively spliced in BA-
treated samples were also alternatively spliced in the controls, and that the relative abundance of splice
variants was similar. Thus, the cytokinin hormone treatment had minimal effect on splicing. By contrast,
there were many genes where the relative abundance of splice variants was different between roots and
shoots (Figure 3C). Consistent with Figure 3, statistical testing of PSI differences between sample types
identified 90 genes where PSI was significantly different between roots and shoots (FDR ≤ 0.1) but only
four and two genes where PSI was different between cytokinin-treated samples and controls in roots and
shoots, respectively (See Supplemental Data I). Thus, we observed limited but non-trivial levels of
differential alternative splicing between roots and shoots but minimal differential alternative splicing
between control and BA-treated samples.
246
Comparison of *Arabidopsis* differential splicing shows similar patterns to rice
To determine whether the observed patterns of differential splicing are similar in other plants, we
analyzed splicing in *Arabidopsis* roots and shoots that had also been treated with cytokinin [14]. Due to
the *Arabidopsis* libraries not being sequenced to the same depth as the rice libraries, many more splicing
events had little or no support. However, PSI was distributed similarly to rice in that most alternatively
spliced genes expressed on major isoform (Supplemental Figure 1A). In addition, scatter plots showing
average PSI in treated versus untreated samples showed a much tigher cluster of points as compared to
scatter ployd comparing roots and shoots (Supplemental Figure 1B). Thus, the cytokinin hormone

treatment had minimal effect on splicing in Arabidopsis. Statistical testing of PSI differences confirmed
this trend. Using the same FDR threshold as with the rice data set (FDR ≤ 0.1), we identified few
differentially spliced regions between shoots and roots (3) and none in the control to treatment
comparisons. Thus, the general pattern of more differential splicing between tissue types as compared to
treatment with exogenous cytokinin appeared conserved between rice and *Arabidopsis*.
260
Alternative splicing remodeled protein-coding sequence more often than disrupting it in rice
Alternative splicing can occur within the UTR or protein-coding regions of genes. Interestingly, 67%
of the differential splicing between roots and shoots occurred within protein-coding regions (Table I and
Supplemental Data I), suggesting that differential splicing is likely to affect gene function at the level of
the protein product. In nearly every instance, major and minor isoforms were both detected, with
differential splicing observed as a change in the relative abundance of the two forms.
In general, when alternatively spliced regions overlap the coding region of genes and the lengths of
these regions are not multiples of three, then inclusion of these differential regions in transcripts is likely
to introduce a frame shift, resulting in a premature stop codon and a truncated protein product. As shown
in Table I, there was an enrichment of alternatively spliced regions in rice that were evenly divisible by
three in coding regions versus non-coding in all subsets of the data. These subsets included all annotated
alternatively spliced regions, regions where the minor form was unusually prevalent (the trough region of
Figure 2), and differentially spliced regions. Thus, alternative splicing within the coding regions of genes
was biased against introducing frame shifts and promoted protein remodeling rather than truncation.
To further understand the effects of splicing on protein-coding sequences, we visualized differentially
spliced regions together with RNA-Seq alignments, coverage graphs, and inferred junctions using
genome browsers. Two genome browsers were used to visualize the data - Integrated Genome Browser
[27] and ProtAnnot [28]. Integrated Genome Browser (IGB) was used to examine RNA-Seq read
alignments and compare alignments to the annotated gene structures. ProtAnnot, an IGB App, was used
to search the InterPro database of conserved protein motifs to find out how (or if) splicing inferred from
RNA-Seq data was likely to affect gene function through remodeling of protein motifs as detected by the
InterProScan Web service [29].
Of the 105 differentially spliced regions, 71 overlapped protein-coding sequence regions, suggesting
that in these cases, alternative splicing affected protein function. All but one (70/71) of the differentially
spliced regions embedded in coding regions overlapped a predicted functional motif (e.g., a predicted
transmembrane helix) or a region found by protein classification systems (e.g., Pfam [30] or PANTHER
[31]) to be conserved among members of the same protein family (Supplemental Data I and Figure 4).
288
RT-PCR with capillary gel electrophoresis confirmed differential splicing between rice roots and
shoots for nine of ten genes tested
We used a method based on capillary gel electrophoresis of fluorescently tagged PCR products to
assay alternative splicing of ten genes detected as differentially spliced between rice roots and shoots
[24]. New rice seedlings were grown under a close-to-identical replication of the RNA-Seq experiment.
Primers were designed to amplify differentially spliced regions, including one primer that was conjugated
to a fluorescent tag. Following PCR amplification of cDNA prepared from the new rice samples, products
were resolved on a capillary-based sequencer and PSI calculated (Table II). In nine out of ten genes,
differential alternative splicing was confirmed. In the one case where differential alternative splicing was
not confirmed, there were very few RNA-Seq read alignments covering the differentially spliced region,
suggesting this was likely a false positive result. The FDR cutoff used to detect differential splicing in the
RNA-Seq data was 0.1, corresponding to 1 in 10 false discoveries, in line with results from the
microcapillary-based analysis.
302
DISCUSSION
This study profiled alternative splicing using a high coverage RNA-Seq data from 10-day old,
hydroponically-grown rice seedlings treated with a cytokinin hormone. A less-deeply sequenced data set

from similarly treated Arabidopsis seedlings provided a comparison. We found that cytokinin treatment
induced very few splicing changes between treated and untreated controls. However, there were many
differences in splicing between untreated roots and shoots, and most of these changed the protein coding
region of genes.
Palusa et al. found that BA-treatment of Arabidopsis seedlings triggered splicing changes in multiple
SR genes [11], encoding RNA-binding proteins whose counterparts in metazoans regulated alternative
splice site choice. Their study used PCR amplification of cDNA followed by agarose gel electrophoresis
to detect changes in splicing and focused on SR protein genes only. Thus, we expected to observe
changes in SR gene splicing due to the cytokinin treatment, leading to changes in splicing for many
downstream genes. However, no such differential splicing was apparent in either RNA-Seq data set
tested. It is possible that the differences in methodology used to measure splicing changes between the
two studies (RNA-Seq versus visualization of PCR amplification of cDNA) could account for the
differences in observations. However, close examination of SR splicing genes in our dataset revealed no
significant differences.
One possible explanation of why the cytokinin treatment had minimal effect on splicing was that the
treatment itself was ineffective. However, differential expression analysis showed that many genes were
up- or down- regulated by the treatment in the two data sets tested – rice and *Arabidopsis* [14]. Fewer
genes were detected as differentially expressed in the *Arabidopsis* data set, most likely reflecting higher
variability between biological replicates combined with lower sequencing depth as compared to the rice
data set. Nevertheless, known cytokinin sensitive genes were regulated in both data sets, showing the
cytokinin treatment penetrated plant cells and induced stereotypical cytokinin signaling without also
triggering changes in splicing.
The relative lack of differential splicing between cytokinin-treated and mock-treated samples suggests
that cytokinin signaling does not employ alternative splicing as a regulatory mechanism to the same
degree as with other plant hormones, notably abscisic acid (ABA). ABA plays a role in perception and
response to stresses, especially desiccation stress [32]. ABA also plays a role in regulating splicing of
SR45, an SR-like protein, and SR45 plays a role in regulating downstream splicing of multiple genes
[33]. Thus, there is a clear connection between stress signaling and the plant stress hormone ABA.
By contrast, cytokinin signaling involves transfer of phosphate groups between successive elements
of a phosphorelay signaling pathway culminating in phosophorylation-dependent activation of Myb-type
transcription factor proteins called type B ARRs. Cytokinin treatment has no or little effect on
transcription of type B ARRs, the key regulators of cytokinin signaling [34,35]. In addition, type B
transcriptional regulators are not highly alternatively spliced. By contrast, a closely related family of
similar genes encoding so-called "pseudo-response regulators" have similar sequence to type B ARRs and
are highly alternatively spliced [36]. These genes are involved in regulating the circadian clock and have
nothing to do with cytokinin signaling.
Using the same methods and data set, we identified a relatively large number of genes in rice (90) that
were differentially spliced genes between shoots and roots, and we validated nine of ten using fragment
analysis and an independent data set. This observation of differential splicing between roots and shoots is
important for two key reasons. First, it shows that our data analysis methods can identify differential
alternative splicing in a data set. In other words, the roots versus shoots comparison provided a positive
control for differential splicing detection. Second, the detection of differential splicing between roots and
shoots illuminates the function of alternative splicing in plant cells. For most of the differentially spliced
regions, both forms were present, and the difference in relative abundance between forms was often
slight, rarely more than 5 or 10% (Supplemental Data I). Our data supports the growing body of evidence
that alternative splicing is cell, tissue, and stage specific in plants [15-17,37], including in roots [38]. It is
likely that alternative splicing plays a role in fine-tuning gene function to meet the needs of different plant
tissues or cell types where a gene is expressed.
We also examined the prevalence of alternative splicing, independent from differential splicing. That
is, we used RNA-Seq read alignments to assess how often annotated alternative splice sites were used in
our RNA-Set data sets. For most genes annotated as alternatively spliced, the minor form frequency was

typically low, accounting for less than 20% of sequence read alignments across the differentially spliced
region. Genes where minor form frequency exceeded 20% exhibited a diversity of functions. Thus, many
diverse processes in rice involved alternatively spliced genes in which splice variants were expressed at
levels likely to affect gene function.
A major limitation of this study was that we limited our analysis to annotated splice forms and did not
attempt to form new transcript models based on the RNA-Seq data. This was done mainly because the
libraries used were not strand-specific and attempts to assemble transcripts using transcript assembly tools
led to incorrect fusion of neighboring genes and other artifacts (not shown). Future studies will therefore
benefit greatly from using better library preparation protocols to simplify and streamline data analysis.
Nonetheless, this analysis provides new insight into the role of alternative splicing in plant tissues and
hormone response.
368
CONCLUSION
By analyzing the number of reads that supported different splice variants, we identified examples of
differential splicing with confirmation by RT-PCR with capillary gel electrophoresis. There were 90
genes differentially spliced between rice root and shoot tissues, but only four between cytokinin-treated
and non-treated samples. For most differential splicing events, the protein-coding regions were affected,
strongly suggesting that differential splicing is playing a role in modulating gene function between roots
and shoots.
376
FUNDING AND SUPPORT
This material is based upon work supported by the National Science Foundation through the Postdoctoral
Research Fellowship in Biology under Grant No.1523814 awarded to Nowlan Freese and the Plant
Genome Research Program award number IOS-1238051. NIH award 2R01GM103463-06A1 to AL
supports development of the IGB software.

Table I. Alternative splicing choices producing difference regions evenly divisible by three or with
remainder of 1 or 2. P-value obtained from a binomial test of the null hypothesis that the true probability
of a differentially spliced region having a length divisible by three is 1 in 3 and an alternative hypothesis
that the probability is greater than 1 in 3.
386

Alternative Splicing		Divisible by 3	Remainder of 1	Remainder of 2	P-value
Location	Event				
Coding region	Annotated as alternatively spliced	3,248	2,466	2,411	3e-36
UTR	Annotated as alternatively spliced	1,152	1,127	1,113	1
Coding region	Minor form is expressed	173	149	153	8e-6
UTR	Minor form is expressed	34	20	13	0.03
Coding region	Differentially spliced	33	24	18	0.03
UTR	Differentially spliced	6	5	11	0.79

388 **Table II.** Differential splicing detected using RNA-Seq and re-tested using capillary gel electrophoresis
389 (CGE). P is the p-value obtained from comparing roots and shoots PSI from CGE.
390

Gene ID	Description	AS type	Avg. RPKM Expression		RNA-Seq PSI (%L)		CGE PSI (%L)		P-value
			Root	Shoot	Root	Shoot	Root	Shoot	
LOC_Os01g25484	ferrodoxin nitrite reductase	RI	300	131	74.2	31.7	29.9	13	5e-05
LOC_Os01g35580	unknown	AS	55.7	34.9	44.0	66.3	49.9	66.9	2.47e-04
LOC_Os01g45274	carbonic anhydrase	ES	171	1,380	96.8	24.3	97.4	15.4	3e-09
LOC_Os01g51290	protein kinase	RI	49.6	49.9	88.4	95.1	13.3	17.2	0.03466
LOC_Os03g05390	citrate transporter	RI	219	174	86.0	96.3	86	95.5	1.2e-04
LOC_Os12g08260	dehydrogenase E1	RI	8.03	30.3	55.7	2.9	4.1	0.87	3e-05
LOC_Os01g61670	ureidoglycolate hydrolase	DS	59	37.8	78.0	31.0	59	37.8	1.6e-09
LOC_Os05g48040	MATE efflux family protein	DS	7.05	6.72	88.1	100.0	89.4	88.27	0.632
LOC_Os02g05830	ribulose bisphosphate carboxylase	RI	4.79	12.3	88.2	10.1	23.9	3.06	8e-04
LOC_Os06g05110	superoxide dismutase	RI	12.5	39.7	38.5	13.6	22.7	6.5	1.3e-07

391
392

Supplemental Data I. Excel spreadsheet file (SuppTable_PSI.xlsx) is provided as a supplement
containing the rice mock root vs mock shoot, mock root vs treated root, and mock shoot vs treated shoot
splicing data. It is available from the "Manuscript" folder in the project source code repository
(https://bitbucket.org/lorainelab/ricealtsplice). The spreadsheet contains output of R script
SplicingAnalysis.Rmd run with "coverage_threshold" parameter set to 10, where coverage_threshold is
the number of reads overlapping a differentially spliced region which supports one or the other splicing
choice when comparing mutually exclusive splicing choices. The R script resides in the project source
code repository in the "AltSplice" subfolder. Data files produced by the script reside in subfolder
"AltSplice/results". The spreadsheet SuppTable_PSI.xlsx contains worksheets copied from script output
files rootsVshoots_MSU7_10.xlsx, roots_MSU7_10.xlsx, and shoots_MSU7_10.xlsx. The worksheet
comparing roots and shoots alternative splicing (from rootsVshoots_MSU7_10.xlsx) was edited by hand
to include information obtained from manual examination of differentially spliced genes and regions
using Integrated Genome Browser and ProtAnnot. A key describing each column and its meaning is also
included. The first column of each data worksheet contains hyperlinked alternatively spliced region
identifiers. To use the hyperlinks, users should first download and launch Integrated Genome Browser
from http://bioviz.org. Clicking the links sends a message to Integrated Genome Browser instructing it to
zoom and scroll to the alternatively spliced region. Users can then open and view RNA-Seq data sets from
the study by selecting the RNA-Seq folder in the **Available Data Sets** section of the IGB **Data Access**
tab.
412

413
Figure 1. Alternative splicing annotation. The overlap between an intron in one gene model and an
exon in another gene model defines an alternatively spliced region. Arrows indicate splice sites, named
AS for acceptor site and DS for donor site. Use of sites named AS-L or DS-L causes inclusion of the
differentially spliced region, generating the longer (L) isoform. Similarly, DS-S and AS-S refer to sites
that exclude the differentially spliced region and generate the shorter (S) isoform. (**A**) Alternative donor
sites, in which the U2 snRNP complex forms at alternative locations on the 5' end of introns. (**B**)
Alternative acceptor sites, in which the U1 snRNP complex forms at alternative sites near the 3' end of
alternatively spliced introns. (**C**) Alternatively spliced intron, in which a donor/acceptor site pairing can
either be used or not used, forming a retained intron (RI). (**D**) Alternatively spliced, skipped exon. In
exon skipping, alternative splicing involves four sites, indicated by DS-S/L, AS-L, DS-L, and SD-S/L.
Exon inclusion requires assembly of two spliceosome complexes linking DS-S/L with AS-L and DS-L
with AS-S/L, while exon skipping requires linking DS-S/L and AS-S/L only.
426

427
Figure 2. Distribution of percent-spliced-in (PSI) for annotated splicing events in rice where each
choice was supported by at least one RNA-Seq alignment. PSI was calculated as 100*L/(S+L), where
L and S were the number of reads that supported the splicing choice that included (L) or excluded (S) the
differentially spliced region. Read alignment counts from all twelve libraries were combined to obtain a
global view of alternative splicing occurrence in rice seedlings. The U-shaped character of the distribution
persisted whether lower or higher thresholds of informative reads were used.
434

435
Figure 3. Scatter plots comparing percent- spliced-in (PSI) between sample types in rice for
annotated splicing events. PSI was calculated from RNA-Seq reads obtained from sequencing rice
seedling shoots and roots grown hydroponically and subjected to a two-hour treatment with BA, a
cytokinin analog, or a mock-treatment (control). PSI is the average of three biological replicates. Only
events with at least 15 informative read alignments in all six samples being compared were included. (A)
BA-treated rice roots (y axis) compared to mock roots (x axis). (B) BA-treated rice shoots (y axis)
compared to mock shoots (x axis). (C) Mock shoots (y axis) compared to mock roots (x axis)
443

444
445
Figure 4. ProtAnnot and IGB images showing difference in splicing between rice shoot and root.
ProtAnnot (upper panel) shows coding region exons color-coded by frame, with regions matching
InterPro profiles indicated by green, linked rectangles. Asterisk highlights difference in the PANTHER
InterPro profile PTHR11516 between isoforms 1 and 5 of the LOC_Os12g08260 gene. Integrated
Genome Browser (lower panels) shows a zoomed-in view of RNA-Seq coverage graphs from rice root
(blue) and shoot (green). Y-axis is the number of RNA-Seq aligned sequences with MSU7 gene models in
black below.
453

Supplemental Figure 1. Percent-spliced-in (PSI) in Arabidopsis RNA-Seq data set between sample

types for annotated splicing events in *Arabidopsis*. PSI was calculated from RNA-Seq reads obtained

from sequencing *Arabidopsis* seedling shoots and roots grown hydroponically and subjected to a two-

hour treatment with BA, a cytokinin analog, or a mock-treatment (control). PSI is the average of three

biological replicates. Only events with at least 10 informative read alignments in all six samples being

compared were included. (B) BA-treated *Arabidopsis* roots (y axis) compared to mock roots (x axis). (C)

BA-treated *Arabidopsis* shoots (y axis) compared to mock shoots (x axis). (C) Mock shoots (y axis)

compared to mock roots (x axis).

REFERENCES CITED

1. Reddy ASN, Marquez Y, Kalyna M, Barta A (2013) Complexity of the Alternative Splicing Landscape in Plants. The Plant Cell 25: 3657-3683.
2. Kalsotra A, Cooper TA (2011) Functional consequences of developmentally regulated alternative splicing. Nature Reviews Genetics 12: 715-729.
3. Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, et al. (2012) The evolutionary landscape of alternative splicing in vertebrate species. Science 338: 1587-1593.
4. Salz HK (2011) Sex Determination in Insects: a binary decision based on alternative splicing. Current opinion in genetics & development 21: 395-400.
5. Filichkin Sergei A, Cumbie Jason S, Dharmawardhana P, Jaiswal P, Chang Jeff H, et al. (2015) Environmental Stresses Modulate Abundance and Timing of Alternatively Spliced Circadian Transcripts in Arabidopsis. Molecular Plant 8: 207-227.
6. Filichkin S, Priest HD, Megraw M, Mockler TC (2015) Alternative splicing in plants: directing traffic at the crossroads of adaptation and environmental stress. Current Opinion in Plant Biology 24: 125-135.
7. Plass M, Agirre E, Reyes D, Camara F, Eyras E (2008) Co-evolution of the branch site and SR proteins in eukaryotes. Trends in Genetics 24: 590-594.
8. Kalyna M, Barta A (2004) A plethora of plant serine/arginine-rich proteins: redundancy or evolution of novel gene functions? Biochemical Society transactions 32: 561-564.
9. Barbosa-Morais NL, Carmo-Fonseca M, Aparício S (2006) Systematic genome-wide annotation of spliceosomal proteins reveals differential gene family expansion. Genome Research 16: 66-77.
10. Keller M, Hu Y, Mesihovic A, Fragkostefanakis S, Schleiff E, et al. (2016) Alternative splicing in tomato pollen in response to heat stress. DNA Research: dsw051.
11. Palusa SG, Ali GS, Reddy ASN (2007) Alternative splicing of pre-mRNAs of Arabidopsis serine/arginine-rich proteins: regulation by hormones and stresses. The Plant Journal 49: 1091-1107.
12. Mei W, Boatwright L, Feng G, Schnable JC, Barbazuk WB (2017) Evolutionarily Conserved Alternative Splicing Across Monocots. Genetics 207: 465-480.
13. Gulledge AA, Roberts AD, Vora H, Patel K, Loraine AE (2012) Mining Arabidopsis thaliana RNA-seq data with Integrated Genome Browser reveals stress-induced alternative splicing of the putative splicing regulator SR45a. American journal of botany 99: 219-231.
14. Raines T, Blakley IC, Tsai Y-C, Worthen JM, Franco-Zorrilla JM, et al. (2016) Characterization of the cytokinin-responsive transcriptome in rice. BMC Plant Biology 16: 260.
15. Sun Y, Hou H, Song H, Lin K, Zhang Z, et al. (2018) The comparison of alternative splicing among the multiple tissues in cucumber. BMC Plant Biology 18: 5.
16. Vitulo N, Forcato C, Carpinelli EC, Telatin A, Campagna D, et al. (2014) A deep survey of alternative splicing in grape reveals changes in the splicing machinery related to tissue, stress condition and genotype. BMC Plant Biology 14: 99-99.
17. Li S, Yamada M, Han X, Ohler U, Benfey PN (2016) High resolution expression map of the Arabidopsis root reveals alternative splicing and lincRNA regulation. Developmental cell 39: 508-522.

18. Loraine AE, McCormick S, Estrada A, Patel K, Qin P (2013) RNA-seq of Arabidopsis pollen
uncovers novel transcription and alternative splicing. Plant Physiol 162: 1092-1109.
19. Vaneechoutte D, Estrada AR, Lin YC, Loraine AE, Vandepoele K (2017) Genome-wide
characterization of differential transcript usage in Arabidopsis thaliana. Plant J 92: 1218-
1231.
20. Kawahara Y, de la Bastide M, Hamilton JP, Kanamori H, McCombie WR, et al. (2013)
Improvement of the Oryza sativa Nipponbare reference genome using next generation
sequence and optical map data. Rice 6: 1-10.
21. Sakai H, Lee SS, Tanaka T, Numa H, Kim J, et al. (2013) Rice Annotation Project Database
(RAP-DB): an integrative and interactive database for rice genomics. Plant Cell Physiol
54: e6.
22. English AC, Patel KS, Loraine AE (2010) Prevalence of alternative splicing choices in
Arabidopsis thaliana. BMC Plant Biol 10: 102.
23. Katz Y, Wang ET, Airoldi EM, Burge CB (2010) Analysis and design of RNA sequencing
experiments for identifying isoform regulation. Nat Methods 7: 1009-1015.
24. Stamm S, Smith C, Lührmann R (2012) Alternative pre-mRNA Splicing: Theory and
Protocols: Wiley-Blackwell.
25. Eddy R, Hahn DT, Moulton T (2012) Optimizing Greenhouse Rice Production: What Is the
Best Root Medium? Purdue Methods for Rice Growth.
26. Naftelberg S, Schor IE, Ast G, Kornblihtt AR (2015) Regulation of alternative splicing
through coupling with transcription and chromatin structure. Annual review of
biochemistry 84: 165-198.
27. Freese NH, Norris DC, Loraine AE (2016) Integrated genome browser: visual analytics
platform for genomics. Bioinformatics 32: 2089-2095.
28. Mall T, Eckstein J, Norris D, Vora H, Freese NH, et al. (2016) ProtAnnot: an App for
Integrated Genome Browser to display how alternative splicing and transcription affect
proteins. Bioinformatics 32: 2499-2501.
29. Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, et al. (2017) InterPro in 2017—
beyond protein family and domain annotations. Nucleic Acids Research 45: D190-D199.
30. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, et al. (2016) The Pfam protein
families database: towards a more sustainable future. Nucleic Acids Research 44: D279-
D285.
31. Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, et al. (2003) PANTHER: A
Library of Protein Families and Subfamilies Indexed by Function. Genome Research 13:
2129-2141.
32. Maia J, Dekkers BJ, Dolle MJ, Ligterink W, Hilhorst HW (2014) Abscisic acid (ABA)
sensitivity regulates desiccation tolerance in germinated Arabidopsis seeds. New Phytol
203: 81-93.
33. Cruz TM, Carvalho RF, Richardson DN, Duque P (2014) Abscisic acid (ABA) regulation of
Arabidopsis SR protein gene expression. Int J Mol Sci 15: 17541-17564.
34. Argyros RD, Mathews DE, Chiang Y-H, Palmer CM, Thibault DM, et al. (2008) Type B
Response Regulators of Arabidopsis Play Key Roles in Cytokinin Signaling
and Plant Development. The Plant Cell 20: 2102-2116.
35. Kieber JJ, Schaller GE (2018) Cytokinin signaling in plant development. Development 145.

36. James AB, Syed NH, Bordage S, Marshall J, Nimmo GA, et al. (2012) Alternative splicing
mediates responses of the Arabidopsis circadian clock to temperature changes. Plant Cell
24: 961-981.
37. Gupta V, Estrada AD, Blakley I, Reid R, Patel K, et al. (2015) RNA-Seq analysis and
annotation of a draft blueberry genome assembly identifies candidate genes involved in
fruit ripening, biosynthesis of bioactive compounds, and stage-specific alternative
splicing. Gigascience 4: 5.
38. Li S, Yamada M, Han X, Ohler U, Benfey PN (2016) High-Resolution Expression Map of
the Arabidopsis Root Reveals Alternative Splicing and lincRNA Regulation. Dev Cell
39: 508-522.
566