1 **Cytokinin treatment of rice induces minimal effect on differential splicing compared to splicing**
2 **between tissues**
3

4 Nowlan Freese[1*], April R. Estrada[1*], Ivory C. Blakley[1], Jinjie Duan[2], Ann Loraine[1**]
5 [1]Department of Bioinformatics and Genomics, University of North Carolina at Charlotte, Charlotte, NC,
6 USA
7 [2]Bioinformatics Research Center, Aarhus University, C.F. Moellers Alle 8, Aarhus C, Denmark.
8
9 [*]Equal contribution.
10 [**]Corresponding author.

11
ABSTRACT

Alternatively spliced genes produce multiple spliced isoforms, called transcript variants. In differential
alternative splicing, the relative abundance of transcript variants differs across sample types. Differential
alternative splicing is common in animal systems, but its extent and significance is not as well known in
plants. We examined differential alternative splicing in rice seedlings using RNA-Seq data that included
approximately 40 million sequence alignments per library, three libraries per sample type, and four
sample types: roots and shoots plus and minus treatment with exogenous cytokinin, a versatile plant
hormone that controls a myriad of developmental and stress-related processes in plants. Comparing read
alignments to gene model annotations found that for 77% of alternative splicing events proposed in the
gene models, each splicing choice was supported by at least one RNA-Seq read alignment. Most genes
annotated as alternatively spliced favored one dominant isoform. Of splicing choices where there was
abundant support for minor forms, most alternative splicing events affected protein-coding sequence.
Statistical testing of read count proportions identified 90 genes as differentially spliced between rice roots
and shoots. By contrast, only four genes were detected as differentially spliced in response to cytokinin.
Ten differential splicing events were selected for validation via capillary gel electrophoresis analysis of
reverse transcriptase-PCR products, using newly prepared samples from an independent experiment. In
nine of ten cases, differential splicing between tissue types was confirmed. A tool for visualizing protein
annotations in the context of genomic sequence (ProtAnnot) together with a genome browser (Integrated
Genome Browser) were used to visualize and assess effects of differential splicing on gene function. In
general, differentially spliced regions coincided with conserved regions in the encoded proteins,
indicating that differential alternative splicing is likely to affect protein function between root and shoot
tissue in rice.

34
INTRODUCTION
Differential splicing of pre-mRNA transcripts, called alternative splicing, enables one gene to
produce multiple transcript variants encoding different functions. Alternative splicing is an almost
universal phenomenon in higher eukaryotes, occurring to varying degrees in every animal and plant
genome examined to date (Kalsotra and Cooper, 2011; Reddy et al., 2013). In animals, differential
expression of splice variants has been recruited as a regulatory mechanism in multiple processes, such as
sex determination in invertebrates and neuronal differentiation in mammals (Kalsotra and Cooper, 2011;
Salz, 2011; Barbosa-Morais et al., 2012).
In plants, less is known about the functional significance and patterns of alternative splicing.
However, several trends are apparent. Genes involved in circadian regulation are highly alternatively
spliced, often producing multiple splice variants that fluctuate in concert with day/night cycling along
with overall transcript abundance (Filichkin et al., 2015). The SR family of RNA-binding, splicing
regulatory proteins is greatly expanded compared to mammals and includes many plant-specific forms
(Kalyna and Barta, 2004; Barbosa-Morais et al., 2006; Plass et al., 2008; Filichkin et al., 2015). SR
transcripts themselves are also highly alternatively spliced in plants, with the relative abundance of these
alternative transcripts varying according to environmental stresses and hormones (Palusa et al., 2007;
Gulledge et al., 2012; Filichkin et al., 2015; Keller et al., 2016; Mei et al., 2017).

There is a growing body of evidence indicating cell and tissue specific regulation of alternative
splicing in plants (Vitulo et al., 2014; Li et al., 2016; Sun et al., 2018). We previously found through
analysis of expressed sequence tags (ESTs) that alternative splicing in the model plant *Arabidopsis*
thaliana was highly skewed toward expression of a single dominant isoform (English et al., 2010).
Analysis of RNA-Seq data from *Arabidopsis* pollen found the relative abundance of splice variants to be
similar between leaves and pollen, despite the differences between the two tissues (Loraine et al., 2013).
However, this latter analysis was limited by having just one biological replicate for pollen and only two
biological replicates for leaves. A more comprehensive analysis of multiple *Arabidopsis* data sets found a
high incidence of isoform switching, in which the identity of the most prevalent variant differs between
sample types (Vaneechoutte et al., 2017). However, this splicing diversity may have arisen in part from
the heterogeneity of the data sets used, which were produced using different technologies at different
times.
In this study, we used a well-replicated RNA-Seq data set from rice to re-examine prevalence of
alternative splicing between tissues and sample types. This data set was previously generated to
investigate cytokinin regulation of gene expression in roots and shoots from 10-day old rice seedlings
(Raines et al., 2016). The data set included three biological replicates per sample type and four sample
types – roots and shoots treated with exogenous cytokinin or a mock, vehicle-only treatment. The
previous study found that cytokinin treatment triggered differential expression of around 5,000 and 2,300
genes in roots and shoots, respectively. Previous work in *Arabidopsis* found that SR transcripts exhibit
differential splicing when exposed to cytokinin, however, less is known about the effect of cytokinin on
alternative splicing in rice (Palusa et al., 2007).
We found that the relative abundance of splice variants for most alternatively spliced genes was
stable, with very few differentially spliced genes between cytokinin treated and control samples. There
was nonetheless a small but significant number of genes that were differentially spliced between roots and
shoots, with the majority of the differential splicing occurring within the protein-coding sequence. These
results provide new evidence that differential alternative splicing likely contributes to gene function
diversification between roots and shoots while playing little role in cytokinin response in rice.
79
80   **MATERIALS & METHODS**
81
RNA-Seq library preparation and sequencing
Samples were prepared and sequenced as described in (Raines et al., 2016). Rice seedlings
(Nipponbare) were grown hydroponically for ten days in a growth chamber set to 14 hours light (28°C)
and 8 hours of dark (23°C) with light intensity 700 mmol m^{-2} s^{-1}. Around six to ten seedlings were grown
in the same pot, in four pots. On the tenth day, culture media was replaced with new media containing 5
mM of the cytokinin benzyladenine (BA) or 0.05 mN NaOH as a control. After 120 minutes, roots and
shoots were harvested separately. Roots and shoots from treatment or control pots were pooled to form
three replicates per treatment. RNA was extracted and used to synthesize twelve libraries from BA-treated
and mock-treated roots and shoots. Libraries were sequenced on an Illumina HiSeq instrument for 100
cycles, yielding 100 base, single end reads. Sequence data are available from the Sequence Read Archive
under accession SRP04905. Aligned, processed data are available from the Oct. 2011 rice genome
assembly IgbQuickload directories at http://www.igbquickload.org.
94
Data processing
Read sequences were aligned onto the *O. sativa japonica* genome assembly Os-Nipponbare-
Reference-IRGSP-1.0 (released October 2011) using TopHat version and BowTie2 with default
parameter settings except that the maximum intron size was set to 5,000 bases (Kawahara et al., 2013).
A command-line, Java program called "FindJunctions" was used to identify exon-exon junctions from
gapped read alignments in the RNA-Seq data. FindJunctions produces BED format files containing
junction features, and the score field of the BED file lists the number of read alignments that supported

the junction. Only reads that aligned to a unique location in the genome were considered. Source code and
compiled versions of FindJunctions are available from https://bitbucket.org/lorainelab/findjunctions.
104
Identification of alternative splicing events and differential splicing
To date, there have been two major releases of *O. sativa japonica* gene models: the MSU7 gene set
(Kawahara et al., 2013) and the RAP-Db gene set (Sakai et al., 2013). The two gene model sets contain
mostly the same data, but the MSU7 gene models appear to be the most heavily used and are well-
supported by informatics resources such as Gramene and Gene Ontology Consortium. For simplicity, and
to take advantage of available functional annotations, we used the MSU7 annotations here.
Annotated alternative splicing events and the number of reads supporting each alternative were
identified using the exon-intron overlap method described in (English et al., 2010). Exons and introns
from pairs of gene models from the same locus were compared to identify alternatively spliced regions.
Regions where an intron in one model overlapped an exon in another model on the same strand were
identified and used to define mutually exclusive splicing choices (Figure 1). Gene models that included an
alternatively spliced region were named "L" (for "Long"). Likewise, models that lacked an alternatively
spliced region were designated "S" (for "Short"). Alternatively spliced regions were classified according
to the type of alternative splicing, as follows. Regions flanked by alternative donor sites were designated
"DS" for alternative donor site. Regions flanked by alternative acceptor sites were labeled "AS" for
alternative acceptor site. AS and DS events that coincided with exon skipping were labeled "AS/ES" and
"DS/ES". Alternatively spliced regions arising from introns that the spliceosome sometimes failed to
excise were designated "RI" for retained intron.
For each alternatively spliced region, read alignments that unambiguously supported one or the other
splicing choice were counted. For AS and DS events, only gapped reads that aligned across intron
junctions were counted as support. For RI events, gapped reads that aligned across the retained intron
were counted as support for the intron-removed (S) form, and un-gapped reads that overlapped at least 20
bases within the intron were counted as support for the intron-retained (L) form.
For each alternatively spliced region in each biological replicate, the number of reads supporting L or
S, but not both, were used to calculate the percent-spliced-in (PSI) as $N/M*100$, where N was the number
of reads supporting the L form and M was the number of reads that supported S or L but not both. This is
the same as the splicing index described in (Katz et al., 2010). A two-sided t-test with equal variance
assumption was used to compare PSI between sample types. Because PSI variance was large for events
with small M (very few informative reads), only alternatively spliced regions where M was 10 or more in
at least three replicate libraries were tested. A false discovery rate (FDR) was calculated for each test
using the method of Benjamini and Hochberg, as implemented in the R programming language "p.adjust"
method. Alternative splicing events with FDR less than or equal to 0.1 were considered differentially
alternatively spliced.
Software used to identify and quantify alternative events is available from
https://bitbucket.org/lorainelab/altspliceanalysis. Data analysis code used to analyze RNA-Seq data is
available from https://bitbucket.org/lorainelab/ricealtsplice. Data analysis code is implemented as R
Markdown files designed to be run in the RStudio development environment. Readers interested in
experimenting with different analysis parameters can clone the repository, modify the code, and re-run
analyses as desired.
144
RT-PCR and capillary gel electrophoresis analysis of alternative splicing
Differential alternative splicing detected by analysis of RNA-Seq was re-tested using RT-PCR-based
fragment analysis method described in (Stamm et al., 2012). Differentially spliced regions were PCR-
amplified and quantified using capillary gel electrophoresis. One benefit of the method is that the results
are expressed as relative abundances of splice variants within a sample, thus eliminating the need to
normalize using reference genes as in traditional qRT-PCR experiments aimed at measuring overall gene
expression.

For splicing validation, new rice seedlings equivalent to the non-BA treated (control) samples used to
generate RNA-Seq data were grown and harvested. Seedlings were grown hydroponically in pots
containing either liquid media only or calcined clay granules (recommended in (Eddy et al., 2012))
watered with liquid media. After twelve days of cultivation, plants were removed from the pots and roots
and shoots were collected separately. Roots and shoots from the same pot were combined to form paired
biological replicates. Samples were frozen on liquid nitrogen and stored at -80 degrees C prior to RNA
extraction.
RNA was extracted using the RNeasy Plant Mini Kit from Qiagen following the manufacturer's
instructions. First strand cDNA was synthesized using oligo dT primers and 1 µg of total RNA per 20 µL
reaction. PCR amplification of cDNA was performed using primers flanking differentially spliced
regions, including one primer labeled with 6-carboxyfluorescein (6-FAM) to enable amplicon detection
via microcapillary electrophoresis. Cycle parameters included denaturation at 94°C for 2 minutes,
followed by 24 cycles of 94°C for 15 sec, 50°C for 30 sec and 70°C for 1 min, with a final elongation step
of 72°C for 10 minutes. This was essentially the same regime described in (Stamm et al., 2012) but with
fewer cycles to ensure reactions were stopped before exiting the logarithmic phase. PCR products were
combined with size standards and separated on a 3730 Genetic Analyzer (Life Technologies). Amplicons
were quantified using manufacturer-provided software by calculating the area under each amplicon peak.
The percentage of the variant containing the alternatively spliced region (%L, also called PSI for
"percent-spliced-in") was calculated by dividing the long form area by the total area for both long and
short forms. Spreadsheets with data exported from the instrument, along with PSI calculations, are
available in the project git repository in a subfolder named "Experimental Testing."
173
174   **RESULTS**
175
Most genes annotated as alternatively spliced favored one dominant isoform
Using the exon-intron overlap method described previously (English et al., 2010), alternative splicing
events within each gene were identified and annotated as shown in Figure 1. Following annotation of
alternative splicing events, RNA-Seq read alignments from the libraries described in (Raines et al., 2016)
were used to assess alternative splicing in the four sample types: roots and shoots from seedlings treated
with the cytokinin compound benzyladenine (BA) or with a mock, control treatment. For each alternative
splicing event, the number of sequence alignments unambiguously supporting each alternative was
counted. These counts were used to calculate percent-spliced-in (PSI), the percentage of read alignments
supporting the longer (L) isoform.
In the combined data from all libraries, 77% of AS events had at least one read supporting each of the
two splicing choices, and 19.8% had support for just one splicing choice. Only 2.8% of AS events has no
reads supporting either form; these corresponded to genes with low or no expression in any of the sample
types tested.
Among the events with support for both choices, the distribution of read support skewed toward
supporting a single choice (Figure 2). In 54% of annotated alternative splicing events, only 10% or fewer
of informative read alignments supported the minor choice. In 70% of annotated alternative splicing
events, only 20% or fewer alignments supported the lower frequency choice. Conversely, 30% of
alternative splicing choices showed simultaneous expression of multiple choices in non-trivial amounts,
e.g., at least 2 in 10 alignments supported the minor form.
195
Genes with abundant support for both alternative splicing choices exhibited a wide range of
functions
We used standard methods for Gene Ontology term enrichment to determine if specific categories of
genes were enriched with genes in which alternative isoform expression was high. Specifically, we asked
if there some Gene Ontology terms with significantly enriched with genes containing alternative splicing
events in which the minor form frequency was at least 20%, i.e., genes occupying the center trough in
Figure 2. Interestingly, we found that these genes exhibited a diversity of gene functions, but no

significant enrichment of functional categories were identified (Young et al., 2010). Thus, alternative
splicing in which minor forms are highly prevalent affects genes with many functions in rice.
205
Many rice genes are differentially spliced between roots and shoots but cytokinin hormone
application has minimal effect on splicing
In animals, differential splicing between cell or tissue types contributes to cellular differentiation,
especially in the nervous system (Naftelberg et al., 2015). Less is known about the role of alternative
splicing in regulating cellular differentiation and other processes in plants. Rice shoots and roots are
profoundly different tissues, but our previous analysis of this same data set found that many of the same
genes were expressed in both (Raines et al., 2016). This raises the question of how these two different
tissues are able to carry out their specialized roles, and suggest the hypothesis that differential splicing
could enable differential functions in genes expressed in both tissues (Reddy et al., 2013). Our previous
study of cytokinin on rice roots and shoots identified significant differences in gene expression in
response to cytokinin exposure (Raines et al., 2016). However, little is known about the role alternative
splicing during cytokinin response, except for one study in Arabidopsis that reported a shift in splicing of
SR protein genes following cytokinin hormone treatment (Palusa et al., 2007). Therefore, we examined
differential splicing in the rice RNA-Seq data set comparing root and shoot tissue with or without
cytokinin.
First, we asked: When an alternatively spliced gene was expressed in two different sample types, was
the relative abundance of splice variants the same or different? To address this, we examined correlation
of PSI between roots and shoots or between BA-treated versus mock-treated samples (Figure 3). We
found that PSI was similar between treated and untreated samples, as revealed by the tighter clustering of
scatter plot points in Figures 3A and 3B. This indicated that genes that were alternatively spliced in BA-
treated samples were also alternatively spliced in the controls, and that the relative abundance of splice
variants was similar. Thus, the cytokinin hormone treatment had minimal effect on splicing. By contrast,
there were many genes where the relative abundance of splice variants was different between roots and
shoots (Figure 3C). Consistent with Figure 3, statistical testing of PSI differences between sample types
identified 90 genes where PSI was significantly different between roots and shoots (FDR ≤ 0.1) but only
four and two genes where PSI was different between cytokinin-treated samples and controls in roots and
shoots, respectively (See Supplemental Table). Thus, we observed limited but non-trivial levels of
differential alternative splicing between roots and shoots but minimal differential alternative splicing
between control and BA-treated samples.
235
Alternative splicing remodeled protein-coding sequence more often than disrupting it
Alternative splicing can occur within the UTR or protein-coding regions of genes. Interestingly, 67%
of the differential splicing between roots and shoots occurred within protein-coding regions (Table I and
Supplemental Table I), suggesting that differential splicing is likely to affect gene function at the level of
the protein product. In nearly every instance, major and minor isoforms were both detected, with
differential splicing observed as a change in the relative abundance of the two forms.
In general, when alternatively spliced regions overlap the coding region of genes and the lengths of
these regions are not multiples of three, then inclusion of these differential regions in transcripts is likely
to introduce a frame shift, resulting in a premature stop codon and a truncated protein product. As shown
in Table I, there was an enrichment of alternatively spliced regions in rice that were evenly divisible by
three in coding regions versus non-coding in all subsets of the data. These subsets included all annotated
alternatively spliced regions, regions where the minor form was unusually prevalent (the trough region of
Figure 2), and differentially spliced regions. Thus, alternative splicing within the coding regions of genes
was biased against introducing frame shifts and promoted protein remodeling rather than truncation.
To further understand the effects of splicing on protein-coding sequences, we visualized differentially
spliced regions together with RNA-Seq alignments, coverage graphs, and inferred junctions using
genome browsers. Two genome browsers were used to visualize the data - Integrated Genome Browser
(Freese et al., 2016) and ProtAnnot (Mall et al., 2016). Integrated Genome Browser (IGB) was used to

examine RNA-Seq read alignments and compare alignments to the annotated gene structures. ProtAnnot,
an IGB App, was used to search the InterPro database of conserved protein motifs to find out how (or if)
splicing inferred from RNA-Seq data was likely to affect gene function through remodeling of protein
motifs as detected by the InterProScan Web service (Finn et al., 2017).
Of the 105 differentially spliced regions, 71 overlapped protein-coding sequence regions, suggesting
that in these cases, alternative splicing affected protein function. All but one (70/71) of the differentially
spliced regions embedded in coding regions overlapped a predicted functional motif (e.g., a predicted
transmembrane helix) or a region found by protein classification systems (e.g., Pfam (Finn et al., 2016) or
PANTHER (Thomas et al., 2003)) to be conserved among members of the same protein family
(Supplemental Table I and Figure 4).
264
RT-PCR with capillary gel electrophoresis confirmed differential splicing for nine of ten genes
tested
We used a method based on capillary gel electrophoresis of fluorescently tagged PCR products to
assay alternative splicing of ten genes detected as differentially spliced between rice roots and shoots
(Stamm et al., 2012). New rice seedlings were grown under a close-to-identical replication of the RNA-
Seq experiment. Primers were designed to amplify differentially spliced regions, including one primer
that was conjugated to a fluorescent tag. Following PCR amplification of cDNA prepared from the new
rice samples, products were resolved on a capillary-based sequencer and PSI calculated (Table II). In nine
out of ten genes, differential alternative splicing was confirmed. In the one case where differential
alternative splicing was not confirmed, there were very few RNA-Seq read alignments covering the
differentially spliced region, suggesting this was likely a false positive result. The FDR cutoff used to
detect differential splicing in the RNA-Seq data was 0.1, corresponding to 1 in 10 false discoveries, in
line with results from the microcapillary-based analysis.
278
DISCUSSION
In this study we profiled the prevalence of alternative splicing using a well-replicated RNA-Seq data
from 10-day old rice roots and shoots, treated or untreated with exogenous benzyl adenine, a synthetic
cytokinin hormone. We found there was at least one RNA-Seq read supporting each annotated alternative
splicing choice for most of the annotated alternative splicing events, with a distribution skewed toward
supporting a single splicing choice. We identified very few differences in splicing between cytokinin
treated samples and mock, whereas we observed limited but significant differential splicing between roots
and shoots, with the majority of those differences falling within the protein coding region of transcripts.
The observation of very little differential splicing between cytokinin-treated and mock-treated
samples was surprising due to previous evidence in *Arabidopsis* (Palusa et al., 2007). Cytokinin exposure
of *Arabidopsis* seedlings by Palusa et al. identified several SR genes that were differentially spliced when
exposed to cytokinin (Palusa et al., 2007). SR proteins regulate splicing choices in other species as well as
being alternatively spliced themselves (Filichkin et al., 2015). Therefore, we had expected to observe
cytokinin-induced differential splicing of SR genes in rice that would also affect other genes. However,
we detected no rice SR protein differential splicing of SR proteins between cytokinin- and mock-treated
rice samples and only four differentially spliced genes between roots and two in shoots.
The relative lack of differential splicing between cytokinin-treated and mock-treated samples suggests
that cytokinin signaling does not employ alternative splicing as a regulatory mechanism in rice. Cytokinin
signaling involves transfer of phosphate groups between successive elements of a phosphorelay signaling
pathway culminating in phosophorylation-dependent activation of Myb-type transcription factor proteins
called type B ARRs (Raines et al., 2016). Thus, regulation of type B ARR transcription factor activity
occurs at the level of protein, and cytokinin treatment has no or little effect on transcription of type B
ARRs. In addition, most type B transcriptional regulators and other members of the cytokinin signaling
pathway are not highly alternatively spliced. By contrast, a closely related family of similar genes
encoding so-called "pseudo-response regulators" have similar sequence to type B ARRs and are highly
alternatively spliced. Further study is necessary to discern why cytokinin exposure in rice has so few

differences in splicing, whether this is true for all monocots, and how the response to cytokinin compares
to dicots such as *Arabidopsis* and if that indicates differences in cytokinin response in general.
In comparison to the number of differentially spliced genes due to cytokinin exposure, there was a
comparatively large number of differentially spliced genes between shoots and roots. There is a growing
body of evidence that alternative splicing is cell, tissue, and stage specific (Vitulo et al., 2014; Gupta et
al., 2015; Li et al., 2016; Sun et al., 2018). Although the alternatively spliced genes with high minor form
frequency were not enriched for particular Gene Ontology terms, this lack of concentrated function
suggests that alternative splicing has been recruited as a mechanism for diversification of gene function. It
is of note that the majority of differential splicing between roots and shoots fell within protein-coding
sequence, and of those events, nearly all splicing events affected a predicted motif.
In our previous comparison of rice root and shoot tissue, we found that the overall diversity of gene
expression in roots and shoots was similar, but with the most highly expressed genes reflecting tissue-
specific differences (Raines et al., 2016). Sun et al. found that clustering of cucumber tissues by
alternative splicing profiles placed leaf/stem far from root tissue (Sun et al., 2018). It is likely that
alternative splicing plays a role in further delineating differences in expression between these tissues.
320
CONCLUSION
By analyzing the number of reads that supported different splice variants, we identified examples of
differential splicing with confirmation by RT-PCR with capillary gel electrophoresis. There were 90
genes differentially spliced between root and shoot tissues, but only four between cytokinin-treated and
non-treated samples. For most differential splicing events, the protein-coding regions were affected,
strongly suggesting that differential splicing is playing a role in modulating gene function between roots
and shoots.
328
FUNDING AND SUPPORT

This material is based upon work supported by the National Science Foundation through the Postdoctoral
Research Fellowship in Biology under Grant No.1523814 awarded to Nowlan Freese and the Plant
Genome Research Program award number IOS-1238051. NIH award 2R01GM103463-06A1 to AL
supports development of the IGB software.

Table I. Alternative splicing choices producing difference regions evenly divisible by three or with
remainder of 1 or 2. P-value obtained from a binomial test of the null hypothesis that the true probability
of a differentially spliced region having a length divisible by three is 1 in 3 and an alternative hypothesis
that the probability is greater than 1 in 3.

Alternative Splicing		Divisible by 3	Remainder of 1	Remainder of 2	P-value
Location	Event				
Coding region	Annotated as alternatively spliced	3,248	2,466	2,411	3e-36
UTR	Annotated as alternatively spliced	1,152	1,127	1,113	1
Coding region	Minor form is expressed	173	149	153	8e-6
UTR	Minor form is expressed	34	20	13	0.03
Coding region	Differentially spliced	33	24	18	0.03
UTR	Differentially spliced	6	5	11	0.79

340 **Table II.** Differential splicing detected using RNA-Seq and re-tested using capillary gel electrophoresis
341 (CGE). P is the p-value obtained from comparing roots and shoots PSI from CGE.
342

Gene ID	Description	AS type	Avg. RPKM Expression		RNA-Seq PSI (%L)		CGE PSI (%L)		P-value
			Root	Shoot	Root	Shoot	Root	Shoot	
LOC_Os01g25484	ferrodoxin nitrite reductase	RI	300	131	74.2	31.7	29.9	13	5e-05
LOC_Os01g35580	unknown	AS	55.7	34.9	44.0	66.3	49.9	66.9	2.47e-04
LOC_Os01g45274	carbonic anhydrase	ES	171	1,380	96.8	24.3	97.4	15.4	3e-09
LOC_Os01g51290	protein kinase	RI	49.6	49.9	88.4	95.1	13.3	17.2	0.03466
LOC_Os03g05390	citrate transporter	RI	219	174	86.0	96.3	86	95.5	1.2e-04
LOC_Os12g08260	dehydrogenase E1	RI	8.03	30.3	55.7	2.9	4.1	0.87	3e-05
LOC_Os01g61670	ureidoglycolate hydrolase	DS	59	37.8	78.0	31.0	59	37.8	1.6e-09
LOC_Os05g48040	MATE efflux family protein	DS	7.05	6.72	88.1	100.0	89.4	88.27	0.632
LOC_Os02g05830	ribulose bisphosphate carboxylase	RI	4.79	12.3	88.2	10.1	23.9	3.06	8e-04
LOC_Os06g05110	superoxide dismutase	RI	12.5	39.7	38.5	13.6	22.7	6.5	1.3e-07

343
344

Supplemental Table I. Excel spreadsheet file (SuppTable_PSI.xlsx) is provided as a supplement. The
spreadsheet contains output of R script SplicingAnalysis.Rmd run with "coverage_threshold" parameter
set to 10, where coverage_threshold is the number of reads overlapping a differentially spliced region
which supports one or the other splicing choice when comparing mutually exclusive splicing choices. The
R script resides in the project source code repository (https://bitbucket.org/lorainelab/ricealtsplice) in the
"AltSplice" subfolder. Data files produced by the script reside in subfolder "AltSplice/results". The
spreadsheet SuppTable_PSI.xlsx contains worksheets copied from script output files
rootsVshoots_MSU7_10.xlsx, roots_MSU7_10.xlsx, and shoots_MSU7_10.xlsx. The worksheet
comparing roots and shoots alternative splicing (from rootsVshoots_MSU7_10.xlsx) was edited by hand
to include information obtained from manual examination of differentially spliced genes and regions
using Integrated Genome Browser and ProtAnnot. A key describing each column and its meaning is also
included. The first column of each data worksheet contains hyperlinked alternatively spliced region
identifiers. To use the hyperlinks, users should first download and launch Integrated Genome Browser
from http://bioviz.org. Clicking the links sends a message to Integrated Genome Browser instructing it to
zoom and scroll to the alternatively spliced region. Users can then open and view RNA-Seq data sets from
the study by selecting the RNA-Seq folder in the **Available Data Sets** section of the IGB **Data Access**
tab.

362
Figure 1. Alternative splicing annotation. The overlap between an intron in one gene model and an
exon in another gene model defines an alternatively spliced region. Arrows indicate splice sites, named
AS for acceptor site and DS for donor site. Use of sites named AS-L or DS-L causes inclusion of the
differentially spliced region, generating the longer (L) isoform. Similarly, DS-S and AS-S refer to sites
that exclude the differentially spliced region and generate the shorter (S) isoform. (**A**) Alternative donor
sites, in which the U2 snRNP complex forms at alternative locations on the 5' end of introns. (**B**)
Alternative acceptor sites, in which the U1 snRNP complex forms at alternative sites near the 3' end of
alternatively spliced introns. (**C**) Alternatively spliced intron, in which a donor/acceptor site pairing can
either be used or not used, forming a retained intron (RI). (**D**) Alternatively spliced, skipped exon. In
exon skipping, alternative splicing involves four sites, indicated by DS-S/L, AS-L, DS-L, and SD-S/L.
Exon inclusion requires assembly of two spliceosome complexes linking DS-S/L with AS-L and DS-L
with AS-S/L, while exon skipping requires linking DS-S/L and AS-S/L only.
375

All samples, combined

376
Figure 2. Distribution of percent-spliced-in (PSI) for annotated splicing events where each choice
was supported by at least one RNA-Seq alignment. PSI was calculated as $100*L/(S+L)$, where L and
S were the number of reads that supported the splicing choice that included (L) or excluded (S) the
differentially spliced region. Read alignment counts from all twelve libraries were combined to obtain a
global view of alternative splicing occurrence in rice seedlings. The U-shaped character of the distribution
persisted whether lower or higher thresholds of informative reads were used.
383

Figure 3. Scatter plots comparing percent- spliced-in (PSI) between sample types for annotated splicing events. PSI was calculated from RNA-Seq reads obtained from sequencing rice seedling shoots and roots grown hydroponically and subjected to a two-hour treatment with BA, a cytokinin analog, or a mock-treatment (control). PSI is the average of three biological replicates. Only events with at least 15 informative read alignments in all six samples being compared were included. (A) BA-treated rice roots (y axis) compared to mock roots (x axis). (B) BA-treated rice shoots (y axis) compared to mock shoots (x axis). (C) Mock shoots (y axis) compared to mock roots (x axis)

393
394
Figure 4. ProtAnnot and IGB images showing difference in splicing between shoot and root.
ProtAnnot (upper panel) shows coding region exons color-coded by frame, with regions matching
InterPro profiles indicated by green, linked rectangles. Asterisk highlights difference in the PANTHER
InterPro profile PTHR11516 between isoforms 1 and 5 of the LOC_Os12g08260 gene. Integrated
Genome Browser (lower panels) shows a zoomed-in view of RNA-Seq coverage graphs from rice root
(blue) and shoot (green). Y-axis is the number of RNA-Seq aligned sequences with MSU7 gene models in
black below.
402

REFERENCES CITED

Barbosa-Morais NL, Carmo-Fonseca M, Aparício S (2006) Systematic genome-wide annotation of spliceosomal proteins reveals differential gene family expansion. Genome Research **16:** 66-77

Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, Slobodeniuc V, Kutter C, Watt S, Colak R, Kim T, Misquitta-Ali CM, Wilson MD, Kim PM, Odom DT, Frey BJ, Blencowe BJ (2012) The evolutionary landscape of alternative splicing in vertebrate species. Science **338:** 1587-1593

Eddy R, Hahn DT, Moulton T (2012) Optimizing Greenhouse Rice Production: What Is the Best Root Medium? Purdue Methods for Rice Growth

English AC, Patel KS, Loraine AE (2010) Prevalence of alternative splicing choices in Arabidopsis thaliana. BMC Plant Biol **10:** 102

Filichkin S, Priest HD, Megraw M, Mockler TC (2015) Alternative splicing in plants: directing traffic at the crossroads of adaptation and environmental stress. Current Opinion in Plant Biology **24:** 125-135

Filichkin Sergei A, Cumbie Jason S, Dharmawardhana P, Jaiswal P, Chang Jeff H, Palusa Saiprasad G, Reddy ASN, Megraw M, Mockler Todd C (2015) Environmental Stresses Modulate Abundance and Timing of Alternatively Spliced Circadian Transcripts in Arabidopsis. Molecular Plant **8:** 207-227

Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, Chang H-Y, Dosztányi Z, El-Gebali S, Fraser M, Gough J, Haft D, Holliday GL, Huang H, Huang X, Letunic I, Lopez R, Lu S, Marchler-Bauer A, Mi H, Mistry J, Natale DA, Necci M, Nuka G, Orengo CA, Park Y, Pesseat S, Piovesan D, Potter SC, Rawlings ND, Redaschi N, Richardson L, Rivoire C, Sangrador-Vegas A, Sigrist C, Sillitoe I, Smithers B, Squizzato S, Sutton G, Thanki N, Thomas PD, Tosatto Silvio C E, Wu CH, Xenarios I, Yeh L-S, Young S-Y, Mitchell AL (2017) InterPro in 2017—beyond protein family and domain annotations. Nucleic Acids Research **45:** D190-D199

Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A (2016) The Pfam protein families database: towards a more sustainable future. Nucleic Acids Research **44:** D279-D285

Freese NH, Norris DC, Loraine AE (2016) Integrated genome browser: visual analytics platform for genomics. Bioinformatics **32:** 2089-2095

Gulledge AA, Roberts AD, Vora H, Patel K, Loraine AE (2012) Mining Arabidopsis thaliana RNA-seq data with Integrated Genome Browser reveals stress-induced alternative splicing of the putative splicing regulator SR45a. American journal of botany **99:** 219-231

Gupta V, Estrada AD, Blakley I, Reid R, Patel K, Meyer MD, Andersen SU, Brown AF, Lila MA, Loraine AE (2015) RNA-Seq analysis and annotation of a draft blueberry genome assembly identifies candidate genes involved in fruit ripening, biosynthesis of bioactive compounds, and stage-specific alternative splicing. Gigascience **4:** 5

Kalsotra A, Cooper TA (2011) Functional consequences of developmentally regulated alternative splicing. Nature Reviews. Genetics **12:** 715-729

Kalyna M, Barta A (2004) A plethora of plant serine/arginine-rich proteins: redundancy or evolution of novel gene functions? Biochemical Society transactions **32:** 561-564

Katz Y, Wang ET, Airoldi EM, Burge CB (2010) Analysis and design of RNA sequencing experiments for identifying isoform regulation. Nat Methods **7:** 1009-1015

Kawahara Y, de la Bastide M, Hamilton JP, Kanamori H, McCombie WR, Ouyang S, Schwartz DC, Tanaka T, Wu J, Zhou S, Childs KL, Davidson RM, Lin H, Quesada-Ocampo L, Vaillancourt B, Sakai H, Lee SS, Kim J, Numa H, Itoh T, Buell CR, Matsumoto T (2013) Improvement of the Oryza sativa Nipponbare reference genome using next generation sequence and optical map data. Rice **6:** 1-10

Keller M, Hu Y, Mesihovic A, Fragkostefanakis S, Schleiff E, Simm S (2016) Alternative splicing in
tomato pollen in response to heat stress. DNA Research**:** dsw051
Li S, Yamada M, Han X, Ohler U, Benfey PN (2016) High resolution expression map of the
Arabidopsis root reveals alternative splicing and lincRNA regulation. Developmental cell **39:** 508-
522
Loraine AE, McCormick S, Estrada A, Patel K, Qin P (2013) RNA-seq of Arabidopsis pollen
uncovers novel transcription and alternative splicing. Plant Physiol **162:** 1092-1109
Mall T, Eckstein J, Norris D, Vora H, Freese NH, Loraine AE (2016) ProtAnnot: an App for
Integrated Genome Browser to display how alternative splicing and transcription affect proteins.
Bioinformatics **32:** 2499-2501
Mei W, Boatwright L, Feng G, Schnable JC, Barbazuk WB (2017) Evolutionarily Conserved
Alternative Splicing Across Monocots. Genetics **207:** 465-480
Naftelberg S, Schor IE, Ast G, Kornblihtt AR (2015) Regulation of alternative splicing through
coupling with transcription and chromatin structure. Annual review of biochemistry **84:** 165-198
Palusa SG, Ali GS, Reddy ASN (2007) Alternative splicing of pre‐mRNAs of Arabidopsis
serine/arginine‐rich proteins: regulation by hormones and stresses. The Plant Journal **49:** 1091-1107
Plass M, Agirre E, Reyes D, Camara F, Eyras E (2008) Co-evolution of the branch site and SR
proteins in eukaryotes. Trends in Genetics **24:** 590-594
Raines T, Blakley IC, Tsai Y-C, Worthen JM, Franco-Zorrilla JM, Solano R, Schaller GE, Loraine
AE, Kieber JJ (2016) Characterization of the cytokinin-responsive transcriptome in rice. BMC Plant
Biology **16:** 260
Reddy ASN, Marquez Y, Kalyna M, Barta A (2013) Complexity of the Alternative Splicing Landscape
in Plants. The Plant Cell **25:** 3657-3683
Sakai H, Lee SS, Tanaka T, Numa H, Kim J, Kawahara Y, Wakimoto H, Yang CC, Iwamoto M,
Abe T, Yamada Y, Muto A, Inokuchi H, Ikemura T, Matsumoto T, Sasaki T, Itoh T (2013) Rice
Annotation Project Database (RAP-DB): an integrative and interactive database for rice genomics.
Plant Cell Physiol **54:** e6
Salz HK (2011) Sex Determination in Insects: a binary decision based on alternative splicing. Current
opinion in genetics & development **21:** 395-400
Stamm S, Smith C, Lührmann R (2012) Alternative pre-mRNA Splicing: Theory and Protocols. Wiley-
Blackwell
Sun Y, Hou H, Song H, Lin K, Zhang Z, Hu J, Pang E (2018) The comparison of alternative splicing
among the multiple tissues in cucumber. BMC Plant Biology **18:** 5
Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan
A, Narechania A (2003) PANTHER: A Library of Protein Families and Subfamilies Indexed by
Function. Genome Research **13:** 2129-2141
Vaneechoutte D, Estrada AR, Lin YC, Loraine AE, Vandepoele K (2017) Genome-wide
characterization of differential transcript usage in Arabidopsis thaliana. Plant J **92:** 1218-1231
Vitulo N, Forcato C, Carpinelli EC, Telatin A, Campagna D, D'Angelo M, Zimbello R, Corso M,
Vannozzi A, Bonghi C, Lucchin M, Valle G (2014) A deep survey of alternative splicing in grape
reveals changes in the splicing machinery related to tissue, stress condition and genotype. BMC Plant
Biology **14:** 99-99
Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq:
accounting for selection bias. Genome Biol **11:** R14
496